

ON POINTS OF CONVERGENCE LATTICES AND SOBRIETY FOR CONVERGENCE SPACES

FRÉDÉRIC MYNARD

ABSTRACT. We characterize the convergence spaces (X, ξ) such that the space of points of $(\mathbb{P}X, \lim_\xi)$ in the category of convergence lattices is (X, ξ) . On the way, we study variants of sobriety and of the axiom T_D in convergence spaces. New phenomena appear when leaving the realm of topological spaces. We obtain new hindsight into the space of points of a convergence lattice and study a special quotient of it, which, in the case $L = (\mathbb{P}X, \lim_\xi)$ for a topological space (X, ξ) , turns out to be homeomorphic to the sobrification of X .

1. PRELIMINARIES AND INTRODUCTION

1.1. **Convergence Spaces.** Let $\mathbb{P}X$ denote the powerset of X . If $\mathcal{A} \subset \mathbb{P}X$,

$$\mathcal{A}^\uparrow = \{B \subset X : \exists A \in \mathcal{A}, A \subset B\}$$

and

$$\mathcal{A}^\# = \{B \subset X : \forall A \in \mathcal{A}, A \cap B \neq \emptyset\}.$$

We also write $\mathcal{A} \# \mathcal{B}$, and say that \mathcal{A} and \mathcal{B} mesh, if $\mathcal{B} \subset \mathcal{A}^\#$, equivalently, $\mathcal{A} \subset \mathcal{B}^\#$.

A convergence ξ on a set X is a relation between the set $\mathbb{F}\mathbb{P}X$ of (set-theoretic) filters on X and the set X , denoted $x \in \lim_\xi \mathcal{F}$ (or $x \in \lim \mathcal{F}$ if there's no risk of ambiguity) if \mathcal{F} and x are ξ -related, subject to the following two axioms:

(monotone) $\mathcal{F} \subset \mathcal{G} \implies \lim_\xi \mathcal{F} \subset \lim_\xi \mathcal{G}$

(point axiom) $x \in \lim_\xi \{x\}^\uparrow$

for every $x \in X$ and every $\mathcal{F}, \mathcal{G} \in \mathbb{F}\mathbb{P}X$. A convergence is *of finite depth* if additionally

(finite depth) $\lim_\xi(\mathcal{F} \cap \mathcal{G}) = \lim_\xi \mathcal{F} \cap \lim_\xi \mathcal{G}$

for every $\mathcal{F}, \mathcal{G} \in \mathbb{F}\mathbb{P}X$.

Continuity of a map $f : (X, \xi) \rightarrow (Y, \tau)$, in symbols $f \in C(\xi, \tau)$, is simply preservation of limits, that is,

(continuity) $f(\lim_\xi \mathcal{F}) \subset \lim_\tau f[\mathcal{F}]$,

where $f[\mathcal{F}] = \{B \subset Y : f^{-1}(B) \in \mathcal{F}\} \in \mathbb{F}\mathbb{P}Y$ is the image filter of \mathcal{F} under f .

Let **Conv** denote the category of convergence spaces and continuous maps. We denote by $|\cdot| : \mathbf{Conv} \rightarrow \mathbf{Set}$ the forgetful functor, so that $|\xi|$ denotes the underlying set of a convergence ξ and $|f|$ is the underlying function of a morphism. If $|\xi| = |\tau|$,

2000 *Mathematics Subject Classification.* 54A20 .

Key words and phrases. convergence space; sober space; weakly sober space; T_D axiom of separation; space of points of a convergence lattice; sobrification.

we say that ξ is *finer than* τ or that τ is *coarser than* ξ , in symbols, $\xi \geq \tau$, if the identity map of their underlying set belongs to $C(\xi, \tau)$. This order turns the set of convergences on a given set into a complete lattice whose greatest element is the discrete topology, least element is the antidiscrete topology, and whose suprema and infima are given by

$$(1.1) \quad \lim_{\bigvee_{\xi \in \Xi} \xi} \mathcal{F} = \bigcap_{\xi \in \Xi} \lim_{\xi} \mathcal{F} \quad \text{and} \quad \lim_{\bigwedge_{\xi \in \Xi} \xi} \mathcal{F} = \bigcup_{\xi \in \Xi} \lim_{\xi} \mathcal{F}.$$

Conv is a concrete topological category; in particular, for every $f : X \rightarrow |\tau|$, there is the coarsest convergence on X , called *initial convergence* for (f, τ) and denoted $f^{-}\tau$, making f continuous (to τ), and for every $f : |\xi| \rightarrow Y$, there is the finest convergence on Y , called *final convergence* for (f, ξ) and denoted $f\xi$, making f continuous (from ξ). Note that with these notations

$$(1.2) \quad f \in C(\xi, \tau) \iff \xi \geq f^{-}\tau \iff f\xi \geq \tau.$$

Moreover, the initial lift on X of a structured source $(f_i : X \rightarrow |\tau_i|)_{i \in I}$ turns out to be $\bigvee_{i \in I} f_i^{-}\tau_i$ and the final lift on Y of a structured sink $(f_i : |\xi_i| \rightarrow Y)_{i \in I}$ turns out to be $\bigwedge_{i \in I} f_i \xi_i$.

If $f : |\xi| \rightarrow Y$ is surjective, $f\xi$ is also called the *quotient convergence*. If $A \subset |\xi|$, the *subspace convergence* $\xi|_A$ or *induced convergence on A* is $i_A^{-}\xi$ where $i_A : A \rightarrow |\xi|$ is the inclusion map. If Ξ is a set of convergences, then the *product convergence* is

$$\prod_{\xi \in \Xi} \xi = \bigvee_{\xi \in \Xi} p_{\xi}^{-}\xi,$$

where $p_{\xi} : \prod_{\xi \in \Xi} |\xi| \rightarrow |\xi|$ is the projection, that is, $\prod_{\xi \in \Xi} \xi$ is the initial convergence for the family of projections.

Convergences of finite depth form a concretely reflective subcategory of **Conv** with reflector L defined (on objects) by $x \in \lim_{L, \xi} \mathcal{F}$ if there is a finite set \mathbb{D} filters on X each converging to x for ξ such that $\mathcal{F} \geq \bigcap_{\mathcal{D} \in \mathbb{D}} \mathcal{D}$.

Of course, every topology τ can be seen as a convergence given by $x \in \lim_{\tau} \mathcal{F}$ if and only if $\mathcal{F} \geq \mathcal{N}_{\tau}(x)$, where $\mathcal{N}_{\tau}(x)$ denotes the neighborhood filter of x in the topology τ . This turns the category **Top** of topological spaces and continuous maps into a full (concretely reflective) subcategory of (the topological extensional and cartesian-closed category) **Conv**. If (X, ξ) is a convergence space, the *topological modification* $T\xi$ of ξ is the topology on X whose set of closed sets are the ξ -closed subsets of X , that is, the sets $C \subset X$ with

$$(1.3) \quad C \in \mathcal{F} \implies \lim_{\xi} \mathcal{F} \subset C.$$

The ξ -open subsets are complements of ξ -closed sets, that is, subsets O satisfying

$$(1.4) \quad \lim_{\xi} \mathcal{F} \cap O \neq \emptyset \implies O \in \mathcal{F}.$$

$T\xi$ is the finest among topologies coarser than ξ . Note that if \mathcal{O}_{ξ} denotes the set of ξ -open sets and $\mathcal{O}_{\xi}(x) = \mathcal{O}_{\xi} \cap \{x\}^{\uparrow}$, then $\mathcal{N}_{\xi}(x) = (\mathcal{O}_{\xi}(x))^{\uparrow}$.

A subspace $(A, \xi|_A)$ of a convergence space (X, ξ) is *dense* if for every $x \in X$ there is $\mathcal{F} \in \mathbb{F}X$ with $A \in \mathcal{F}$ and $x \in \lim_{\xi} \mathcal{F}$.

1.2. Convergence Lattices. A pointfree generalization was offered in [5] in which the function

$$\lim_{\xi} : \mathbb{F}PX \rightarrow \mathbb{P}X$$

is abstracted away to a monotone function

$$\lim : \mathbb{F}L \rightarrow L$$

from (order-theoretic) filters on a lattice L to L . (point axiom) is not part of the axiomatic in this pointfree version of convergence spaces, though the notion can also be recovered (as so-called *centered* convergence lattices) in an abstract order-theoretic form.

Definition 1. Given a category \mathbf{C} of lattices, a *convergence \mathbf{C} -object* (L, \lim) is a \mathbf{C} -object L together with a monotone map $\lim : \mathbb{F}L \rightarrow L$. The objects of the category \mathbf{C}^{conv} are the convergence \mathbf{C} -objects and the morphisms $\varphi : L \rightarrow L'$ are the \mathbf{C} -morphisms that are *continuous* in the sense that for every $\mathcal{F} \in \mathbb{F}L'$,

$$\text{(ptfree continuity)} \quad \lim_{L'} \mathcal{F} \leq \varphi(\lim_L \varphi^{-1}(\mathcal{F})),$$

where $\varphi^{-1}(\mathcal{F}) = \{\ell \in L : \varphi(\ell) \in \mathcal{F}\}$.

The category \mathbf{Conv} embeds coreflectively into $(\mathbf{C}^{\text{conv}})^{op}$ when \mathbf{C} is the category of frames or of coframes: the *powerset functor* $\mathbb{P} : \mathbf{Conv} \rightarrow (\mathbf{C}^{\text{conv}})^{op}$ sending (X, ξ) to $(\mathbb{P}X, \lim_\xi)$ and $f : (X, \xi) \rightarrow (Y, \tau)$ to $\mathbb{P}f = f^{-1} : \mathbb{P}Y \rightarrow \mathbb{P}X$ (in \mathbf{C}^{conv}) is then right-adjoint to the *point-functor* $\text{pt} : (\mathbf{C}^{\text{conv}})^{op} \rightarrow \mathbf{Conv}$ (the coreflector) where the underlying set of $\text{pt } L$, the set of “points” of L , is the set of \mathbf{C}^{conv} -morphisms from L to $\mathbb{P}(1)$ (which is terminal in $(\mathbf{C}^{\text{conv}})^{op}$ [5, Lemma 2.7]), hence depends on the choice of \mathbf{C} , so we’ll write $\text{pt}_{\mathbf{C}} L$ when different options for the base category \mathbf{C} of lattices are considered.

For a given \mathbf{C} , the convergence structure on $\text{pt } L$ is given by

$$\text{(Conv on pt)} \quad \lim_{\text{pt } L} \mathcal{F} = (\lim_L \mathcal{F}^\circ)^\bullet,$$

where $\ell^\bullet = \{\varphi \in \text{pt } L : \varphi(\ell) = \{1\}\}$ and $\mathcal{F}^\circ = \{\ell \in L : \ell^\bullet \in \mathcal{F}\}$. Finally, if $\varphi \in \mathbf{C}^{\text{conv}}(L, L')$ then $\text{pt } \varphi : \text{pt } L' \rightarrow \text{pt } L$ is defined by $\text{pt}(\varphi)(f) = f \circ \varphi$.

As pointed out in [5, Remark 2.9], a continuous \mathbf{C} -morphism $\varphi : L \rightarrow \mathbb{P}(1)$ can be identified with the filter $\mathcal{F} = \varphi^{-1}(\{1\})$ which satisfies $\lim_L \mathcal{F} \in \mathcal{F}$ by continuity. Hence the set $\text{pt}_{\mathbf{C}} L$ of points of a \mathbf{C}^{conv} -object L can be identified with sets of specific filters as follows ⁽¹⁾:

cat. \mathbf{C} of	$\text{pt}_{\mathbf{C}} L = \{\mathcal{F} \in \mathbb{F}L : \lim_L \mathcal{F} \in \mathcal{F} \text{ and } \dots\}$	ref.
lattices	\mathcal{F} is prime	[5, Remark 2.10]
frames	\mathcal{F} is completely prime	[5, Remark 2.11]
coframes	\mathcal{F} is principal of a \vee -prime element	[5, Remark 2.12]

Lemma 2. [5, Lemma 2.19(6)] *Let L be a convergence lattice and let $\mathcal{U} \in \text{pt } L$. Then $(\{\mathcal{U}\}^\dagger)^\circ = \mathcal{U}$.*

Proof. $\ell \in (\{\mathcal{U}\}^\dagger)^\circ$ if and only if $\ell^\bullet \in \{\mathcal{U}\}^\dagger$, that is, if and only if $\mathcal{U} \in \ell^\bullet$, equivalently, $\ell \in \mathcal{U}$. \square

¹Recall that a proper filter \mathcal{F} on a lattice L is *prime* if $\ell_1 \vee \ell_2 \in \mathcal{F}$ implies that ℓ_1 or ℓ_2 belongs to \mathcal{F} and *completely prime* if for every $A \subset L$, $\bigvee_{\ell \in A} \ell \in \mathcal{F}$ implies that $\ell \in \mathcal{F}$ for some $\ell \in A$. An element $\ell \in L$ is *join-prime* if $\ell \leq \ell_1 \vee \ell_2$ implies $\ell \leq \ell_1$ or $\ell \leq \ell_2$.

It turns out that filters of the form $\{x\}^\uparrow$ are always points of $\mathbb{P}(X)$ in \mathbf{C}^{conv} provided that \mathbf{C} be an admissible category ⁽²⁾ of lattices [5, Remark 2.14]. As a result ⁽³⁾:

Proposition 3. *A convergence space (X, ξ) is always (homeomorphic to) a dense subspace of $\text{pt}((\mathbb{P}X, \lim_\xi))$.*

Proof. The map $h : (X, \xi) \rightarrow \text{pt}((\mathbb{P}X, \lim_\xi))$ defined by $h(x) = \{x\}^\uparrow$ is an embedding, i.e., an homeomorphism onto its image. The map h is clearly one-to-one. To see that it is an embedding, let $L = \text{pt}((\mathbb{P}X, \lim_\xi))$ and recall that $\lim_{\text{pt } L} h[\mathcal{F}] = (\lim_\xi h[\mathcal{F}]^\circ)^\bullet$ where $h[\mathcal{F}]^\circ = \{A \in \mathbb{P}(X) : A^\bullet \in h[\mathcal{F}]\}$ and $A^\bullet = \{\mathcal{G} \in \text{pt } L : A \in \mathcal{G}\}$. Noting that $A^\bullet \in h[\mathcal{F}]$ if and only if $A \in \mathcal{F}$, we see that $h(x) \in \lim_{\text{pt } L} h[\mathcal{F}]$ if and only if $x \in \lim_\xi \mathcal{F}$.

Suppose $\mathcal{G} \in \text{pt } L$. Then $\lim_\xi \mathcal{G} \in \mathcal{G}$ (when \mathcal{G} is seen as an element of $\mathbb{P}X$), that is, $\mathcal{G} \in (\lim_\xi \mathcal{G})^\bullet$. Moreover $\mathcal{G} \subset h[\mathcal{G}]^\circ$ so that $\lim_\xi \mathcal{G} \subset \lim_\xi h[\mathcal{G}]^\circ$ and thus $\mathcal{G} \in (\lim_\xi \mathcal{G})^\bullet \subset (\lim_\xi h[\mathcal{G}]^\circ)^\bullet$, that is, $\mathcal{G} \in \lim_{\text{pt } L} h[\mathcal{G}]$. Hence $h(X)$ is a dense subspace of $\text{pt } L$. To see that $\mathcal{G} \subset h[\mathcal{G}]^\circ$, note that if $G \in \mathcal{G}$ then

$$h(G) = \{\{x\}^\uparrow : x \in G\} \subset G^\bullet$$

so that $G^\bullet \in h[\mathcal{G}]$, that is, $G \in h[\mathcal{G}]^\circ$. □

As filters of the form $\{x\}^\uparrow$ are exactly completely prime filters in $\mathbb{P}(X)$ they are the only points in \mathbf{C}^{conv} if \mathbf{C} is the category of frames [5, Remark 2.16]. Similarly, as join-prime elements of $\mathbb{P}(X)$ are singletons, points of $\mathbb{P}(X)$ in \mathbf{C}^{conv} are exactly singletons if \mathbf{C} is the category of coframes [5, Remark 2.17]. Therefore, in both cases, $\text{pt}((\mathbb{P}X, \lim_\xi))$ is homeomorphic to (X, ξ) .

On the other hand, prime filters of $\mathbb{P}(X)$ are the *ultrafilters* on X . Hence, when \mathbf{C} is the category **Lat** of lattices and (X, ξ) is a convergence space, the set of points of $(\mathbb{P}(X), \lim_\xi)$ in \mathbf{C}^{conv} is

$$(1.5) \quad \text{pt}_{\mathbf{Lat}} \mathbb{P}(X) = \{\mathcal{U} \in \mathbb{U}X : \lim_\xi \mathcal{U} \in \mathcal{U}\},$$

where $\mathbb{U}X$ denotes the set of ultrafilters on X [5, Remark 2.15]. Though it contains the “usual points” identified with their principal ultrafilter $\{x\}^\uparrow$, there may in general be others (See Lemma 6 below). One purpose of this note is to clarify when this does not happen, that is:

Problem 4. Under what condition on a convergence space (X, ξ) are all ultrafilters $\mathcal{U} \in \mathbb{U}X$ with $\lim_\xi \mathcal{U} \in \mathcal{U}$ principal?

First we should point out that the question at hand was fully solved [7, Theorem 2.2 with Addendum I] for the case where (X, ξ) is topological. Though our motivations leading to Problem 4 are distinct from those of [7] and come from [5], we are thus essentially exploring a generalization of [7, Theorem 2.2] from topological spaces to convergence spaces. Such an extension may seem of little relevance, but in light of the question of finding out when $\text{pt}_{\mathbf{Lat}} \mathbb{P}(X)$ are just the “usual points of X ” in the context of convergence lattices, this is a most natural problem and

²A category \mathbf{C} of lattices is *admissible* if $\mathbb{P}(X)$ is a \mathbf{C} -object for every set X and there are two classes \mathcal{I} and \mathcal{J} of index sets such that for all pairs L, L' of \mathbf{C} -objects, the \mathbf{C} -morphisms from L to L' are exactly monotonic maps that preserve all I -indexed infima that exist in L for $I \in \mathcal{I}$ and all J -indexed suprema that exist in L for $J \in \mathcal{J}$. See [5, Def. 2.5]

³Save density, this was already observed in [5, Lemma 2.26]

restricting ourselves to topological spaces is then highly undesirable. Corollary 47 gives a full answer to Problem 4 for a large class of convergence spaces.

It turns out that some interesting subtleties appear in this wider context of convergence spaces. On the way, we study variants of sobriety for convergence spaces and take the opportunity to correct a related misstatement in [5, Remark 2.15].

Though Problem 4 stems from a question on $\text{pt } L$ in the very particular case where $L = (\mathbb{P}X, \lim_\xi)$ for a convergence space (X, ξ) , we study more in depth the space $\text{pt } L$ for a general convergence lattice L . A particular natural quotient $\text{pt}' L$ of $\text{pt } L$ turns out to be of importance. In particular, when $L = (\mathbb{P}X, \lim_\xi)$ for a topological space (X, ξ) , the space $\text{pt}' L$ is homeomorphic to the sobrification of (X, ξ) .

2. IRREDUCIBLE FILTERS, SOBRIETY, AND CONVERGENCE VARIANTS

A filter \mathcal{F} on a convergence space (X, ξ) is *irreducible* in the sense of [7] if

$$\lim_\xi \mathcal{F} \in \mathcal{F}.$$

Hence, elements of $\text{pt}_{\mathbf{C}} \mathbb{P}(X)$ are specific types of irreducible filters. As we have seen, when \mathbf{C} is either the category of frames or that of coframes, points turn out to be exactly principal ultrafilters without the irreducibility condition even playing a role. This is not the case for $\text{pt}_{\mathbf{Lat}} \mathbb{P}(X)$ whose elements are exactly irreducible ultrafilters. Note that if \mathcal{F} is irreducible and $\mathcal{G} \geq \mathcal{F}$ then \mathcal{G} is irreducible too and if $f : (X, \xi) \rightarrow (Y, \tau)$ is continuous and \mathcal{F} is irreducible on X then $f[\mathcal{F}]$ is irreducible on Y .

The notion is related to that of a *compact filter* in the sense of [2, 3, 11, 10]. A filter \mathcal{F} on a convergence space (X, ξ) is *compact* (resp. *compactoid*) if

$$(2.1) \quad \mathcal{H} \# \mathcal{F} \implies \text{adh } \mathcal{H} \# \mathcal{F} \quad (\text{resp. } \text{adh } \mathcal{H} \neq \emptyset),$$

for every $\mathcal{H} \in \mathbb{F}X$.

Lemma 5. *Let (X, ξ) be a convergence space.*

- (1) *If \mathcal{F} is irreducible then \mathcal{F} is compact;*
- (2) *A compact ultrafilter is irreducible.*

Proof. (1). If \mathcal{F} is irreducible and $\mathcal{H} \# \mathcal{F}$ then $\lim \mathcal{F} \subset \text{adh } \mathcal{H}$ so that $\text{adh } \mathcal{H} \in \mathcal{F} \subset \mathcal{F}^\#$.

(2). If $\mathcal{U} \in \mathbb{U}X$ is compact, then taking $\mathcal{H} = \mathcal{U}$ in (2.1) we obtain $\lim \mathcal{U} \in \mathcal{U}^\# = \mathcal{U}$ and thus \mathcal{U} is irreducible. \square

Hence an ultrafilter is irreducible if and only if it is compact. However, there are compact filters that are not irreducible. For instance, a neighborhood filter of the real line with its usual topology is compact but not irreducible.

Recall that a topological space X is called *Noetherian* if every decreasing sequence of closed subsets is eventually constant. It is well-known that X is Noetherian if and only if all of its subsets are compact. In [5, Remark 2.15], it is erroneously stated that a topological space is Noetherian if and only if every compact ultrafilter is principal. On the contrary:

Lemma 6. *A topological space is Noetherian if and only if every filter, equivalently every ultrafilter, is compact. In particular, an infinite Noetherian topological space admits irreducible free ultrafilters.*

Proof. If X is Noetherian, $\mathcal{F} \in \mathbb{F}X$ and \mathcal{H} is a filter $\mathcal{H} \# \mathcal{F}$ then $\text{adh } \mathcal{H} \cap F \neq \emptyset$ for every $F \in \mathcal{F}$ because F is compact. Conversely, if every ultrafilter is compact and $A \subset X$ then $\lim \mathcal{U} \cap A \neq \emptyset$ whenever $A \in \mathcal{U} \in \mathbb{U}X$ because $\text{adh } \mathcal{U} = \lim \mathcal{U} \in \mathcal{U}^\#$. \square

Recall that a closed subset C of a topological space is *irreducible* if whenever $C \subset D \cup F$ where D and F are closed, then either $C \subset D$ or $C \subset F$. A topological space X is *sober* if for every irreducible closed subset C of X , there is a (necessarily unique) point $x \in X$ with $C = \text{cl}\{x\}$, which we call a *generic point for C* . Note that we can equivalently formulate irreducibility of C in the following alternative terms: If O_1, O_2 are open sets intersecting with C then $O_1 \cap O_2$ also intersects with C .

It was shown in [7] that a topological space X is sober if and only if for every irreducible filter \mathcal{F} on X there is a unique point $x \in X$ with $\lim \mathcal{F} = \lim\{x\}^\uparrow$. Following [7], we can take this as a definition of a *sober convergence space* and call *weakly sober* a convergence space in which for every irreducible *ultrafilter* \mathcal{U} there is a unique x with $\lim \mathcal{U} = \lim\{x\}^\uparrow$. The two notions (sober and weakly sober) coincide for topological spaces, but not for convergence spaces. Though [7, Remark 1.9] mentions this, it provides neither a proof nor an example and does not explore the conditions on a convergence space for weak sobriety and sobriety to coincide. We fill this gap here (Example 14, Proposition 8).

In this context, it is useful to point out that the condition

$$(S_0) \quad (x \in \lim \mathcal{F} \text{ and } t \in \lim\{x\}^\uparrow) \implies t \in \lim \mathcal{F}$$

is automatically true in a topological space because in this case $t \in \lim\{x\}^\uparrow$ implies $\mathcal{O}(t) \subset \mathcal{O}(x)$. However condition (S_0) is not generally true in an arbitrary convergence space. This (vacuous for topological spaces) weak diagonal axiom is called S_0 in [14].

Recall (e.g., [4]) that a convergence space is T_0 if points can be distinguished by the convergence structure, that is, if

$$x \neq t \implies \{\mathcal{F} \in \mathbb{F}X : x \in \lim \mathcal{F}\} \neq \{\mathcal{F} \in \mathbb{F}X : t \in \lim \mathcal{F}\},$$

and T_1 if singletons are closed. Of course, a topological space is T_0 (resp. T_1) in the usual sense if and only if it is in the convergence sense.

Note that if a convergence has closed limits (that is, $\lim \mathcal{F}$ is closed for every filter) then it is S_0 , but not conversely. Of course, in a topological space limits are closed. It might be useful to have Figure 2 below in mind when considering the next few results:




FIGURE 2.1. weak diagonality and separation conditions

Lemma 7. (1) *If limit sets of principal ultrafilters are closed (in particular if the convergence is S_0), then*

$$\{x, y\} \subset \lim\{x\}^\uparrow \cap \lim\{y\}^\uparrow \iff \lim\{x\}^\uparrow = \lim\{y\}^\uparrow.$$

(2) *If X is T_0 and S_0 then*

$$(2.2) \quad \{x, y\} \subset \lim\{x\}^\uparrow \cap \lim\{y\}^\uparrow \implies x = y.$$

Proof. It is clear that if $\lim\{x\}^\uparrow = \lim\{y\}^\uparrow$ then in particular $\{x, y\} \subset \lim\{x\}^\uparrow \cap \lim\{y\}^\uparrow$. Conversely, if $\lim\{x\}^\uparrow$ is closed and belongs to $\{y\}^\uparrow$, then $\lim\{y\}^\uparrow \subset \lim\{x\}^\uparrow$ and similarly for the reverse inclusion if $\lim\{y\}^\uparrow$ belongs to $\{x\}^\uparrow$.

For the second part if $x \neq y$, there is a filter \mathcal{F} with $\text{card}(\lim \mathcal{F} \cap \{x, y\}) = 1$ because the convergence is T_0 . Say $x \in \lim \mathcal{F}$. Then $y \notin \lim\{x\}^\uparrow$ for otherwise $y \in \lim \mathcal{F}$ by S_0 , and similarly $x \notin \lim\{y\}^\uparrow$ if $x \in \lim \mathcal{F}$, so that $\{x, y\} \not\subset \lim\{x\}^\uparrow \cap \lim\{y\}^\uparrow$. \square

In view of Lemma 7, in a T_0 topological space a generic point of an irreducible closed set is necessarily unique. However,

$$(2.3) \quad \lim\{x\}^\uparrow = \lim\{y\}^\uparrow \implies x = y$$

requires both S_0 and T_0 for general convergence spaces as shown by Examples 9 and 10 below.


Proposition 8. *A weakly sober convergence space in which limits of irreducible filters are closed (in particular, a topological space) is sober.*

Proof. If X is a weakly sober convergence space and \mathcal{F} is an irreducible filter on X , then an ultrafilter \mathcal{U} finer than \mathcal{F} is also irreducible, so that $\lim \mathcal{U} = \lim\{x_{\mathcal{U}}\}^\uparrow$ for a unique point $x_{\mathcal{U}} \in X$. Moreover, $\lim \mathcal{F} \subset \lim \mathcal{U}$ because $\mathcal{U} \geq \mathcal{F}$ and as $\lim \mathcal{F} \in \mathcal{F} \subset \mathcal{U}$ and $\lim \mathcal{F}$ is closed, $\lim \mathcal{U} \subset \lim \mathcal{F}$. Hence $\lim \mathcal{U} = \lim \mathcal{F}$ for every


$\mathcal{U} \in \mathbb{U}(\mathcal{F})$. Hence all points $x_{\mathcal{U}}$ coincide and this point is the unique generic point for \mathcal{F} . \square

It turns out that all new phenomena in the realm of general convergences (as opposed to topological spaces) can be observed with finite examples. Filters on a finite sets are all principal and a finitely deep convergence on a finite set is entirely determined by its restriction to principal ultrafilters. Hence such convergences can easily be described by diagrams in which $x \rightarrow y$ means that $y \in \lim\{x\}^\uparrow$ and we include such pictures in each relevant example, using this convention and systematically omitting the loops (i.e., we do not depict the automatic convergence relation $x \in \lim\{x\}^\uparrow$).

Example 9 (A T_0 convergence with $x \neq y$ and $\lim\{x\}^\uparrow = \lim\{y\}^\uparrow$). Let $X = \{x, y, z\}$ with the convergence of finite depth given by $\lim\{x\}^\uparrow = \lim\{y\}^\uparrow = \lim\{x, y\}^\uparrow = \{x, y\}$ and $\lim\{z\}^\uparrow = \{z, x\}$. By finite depth, $\lim\{x, y, z\}^\uparrow = \lim\{y, z\}^\uparrow = \lim\{x, z\}^\uparrow = \{x\}$.



Example 10 (A (non T_0) S_0 convergence space with $x \neq y$ and $\lim\{x\}^\uparrow = \lim\{y\}^\uparrow$). Let $X = \{x, y, z\}$ the convergence of finite depth given by $\lim\{x\}^\uparrow = \lim\{y\}^\uparrow = \lim\{x, y\}^\uparrow = \{x, y\}$ and $\lim\{z\}^\uparrow = \{z\}$. By finite depth, $\lim\{x, y, z\}^\uparrow = \lim\{y, z\}^\uparrow = \lim\{x, z\}^\uparrow = \emptyset$.



Note that weakly sober implies (2.3): if there are points $x \neq y$ with $\lim\{x\}^\uparrow = \lim\{y\}^\uparrow$ then $\{x\}^\uparrow$ is an irreducible ultrafilter with two different generic points, so that the space is not weakly sober. Hence the easy fact that finite T_0 topological spaces are sober fails to extend to convergence spaces, because of the uniqueness requirement on generic points, though this requirement is vacuous in topological spaces. Therefore, it makes sense to also consider the variants without this requirement:

A convergence space X is *quasi-sober* (resp. *weakly quasi-sober*) if for every irreducible filter \mathcal{F} (respectively, for every irreducible ultrafilter \mathcal{F}) on X there is a (not necessarily unique) point x of X with $\lim \mathcal{F} = \lim\{x\}^\uparrow$. Example 9 is T_0 and quasi-sober but not weakly sober. Example 10 is S_0 and quasi-sober but not weakly sober. A quasi-sober space does not need to be T_0 for the antidiscrete topology is quasi-sober. Of course, if a convergence space satisfies (2.3) and is quasi weakly sober, it is weakly sober. Lemma 7 states that this is the case in a T_0 space with closed limits of principal ultrafilters. More generally, let us call a convergence space *antisymmetric* if (2.2) holds and *almost antisymmetric*, or *aas* for short, if (2.3) holds. Note that:

- Proposition 11.** (1) *A convergence space is weakly sober if and only if it is weakly quasi-sober and aas.*
 (2) *An antisymmetric space is T_0 .*

(3) In a convergence in which limits of principal ultrafilters are closed, the following are equivalent:

- (a) weakly sober;
- (b) weakly quasi-sober and antisymmetric;
- (c) weakly quasi-sober and aas.

In particular a weakly sober space in which limits of principal ultrafilters are closed is T_0 .


Proof. (1) If the space is weakly sober and $\lim\{x\}^\uparrow = \lim\{y\}^\uparrow$ then $x = y$ by uniqueness of generic points for limits of irreducible ultrafilters, hence the space is aas. Conversely, aas ensure uniqueness of generic points in a weakly quasi-sober space.

(2) Suppose $\{\mathcal{F} \in \mathbb{F}X : x \in \lim \mathcal{F}\} = \{\mathcal{F} \in \mathbb{F}X : y \in \lim \mathcal{F}\}$. In particular $\{x, y\} \subset \lim\{x\}^\uparrow \cap \lim\{y\}^\uparrow$ so that $x = y$ by aas.

For (3), to see (a) \implies (b), suppose $\{x, y\} \subset \lim\{x\}^\uparrow \cap \lim\{y\}^\uparrow$. Then $\lim\{x\}^\uparrow \subset \lim\{y\}^\uparrow$ because $\lim\{y\}^\uparrow \in \{x\}^\uparrow$ and $\lim\{y\}^\uparrow$ is closed, and similarly for the reverse inclusion. (b) \implies (c) is clear and (a) \iff (c) is (1). \square

A topological sober space is always T_0 but things are slightly more complicated for the general case:


Example 12 (A sober convergence that is not T_0 , whose modification of finite depth is not sober). Let $X = \{x, y, s, t\}$ with the convergence given by $\lim\{x\}^\uparrow = \{x, y, s\}$, $\lim\{y\}^\uparrow = \{x, y, t\}$, $\lim\{s\}^\uparrow = \{s\}$ and $\lim\{t\}^\uparrow = \{t\}$, and all other filters do not converge (note that this convergence is not of finite depth).



Then $\lim^{-1}(x) = \lim^{-1}(y) = \{\{x\}^\uparrow, \{y\}^\uparrow\}$ but $\lim\{x\}^\uparrow \neq \lim\{y\}^\uparrow$ and the condition of sobriety is satisfied. Note that the modification of finite depth $L\xi$ would add the irreducible filter $\{x, y\}^\uparrow$ because the $\lim_{L\xi}\{x, y\}^\uparrow = \lim_\xi\{x\}^\uparrow \cap \lim_\xi\{y\}^\uparrow = \{x, y\}$ but this limit has no generic point, hence sobriety would fail.

In other words, the modification of finite depth of a sober convergence may fail to be sober. On the other hand, the example above is easily modified to make it of a finite depth:

Example 13 (A sober convergence of finite depth that is not T_0). Let $X = \{x, y, s, t, z\}$ with the convergence of finite depth given by $\lim\{x\}^\uparrow = \{x, y, z, s\}$, $\lim\{y\}^\uparrow = \{x, y, z, t\}$, $\lim\{z\}^\uparrow = \{x, y, z\}$, $\lim\{s\}^\uparrow = \{s\}$ and $\lim\{t\}^\uparrow = \{t\}$.




This convergence is not T_0 as the same filters converge to x, y and z . By finite depth,

$$\lim\{x, y\}^\uparrow = \lim\{x, z\}^\uparrow = \lim\{y, z\}^\uparrow = \lim\{x, y, z\}^\uparrow = \{x, y, z\},$$

so that these four filters are irreducible and all have the same unique generic point z .


Example 14 (A T_0 weakly sober convergence of finite depth that is not sober). On $X = \{x, y, t, s, w, z\}$ consider the convergence of finite depth given by $\lim\{x\}^\uparrow = \{x, y, s, t\}$, $\lim\{y\}^\uparrow = \{x, y, t, w\}$, $\lim\{t\}^\uparrow = \{t\}$, $\lim\{s\}^\uparrow = \{s\}$, $\lim\{w\}^\uparrow = w$ and $\lim\{z\}^\uparrow = \{z, x\}$.



This is a T_0 weakly sober convergence of finite depth that is not sober, because $\{x, y\}^\uparrow$ is irreducible but does not have a generic point.

Note that if $\xi \geq \tau$ and τ is (weakly) sober then ξ is (weakly) sober. In particular, if $T\xi$ is (weakly) sober, so is ξ but the converse is false:

Example 15 (A T_0 sober convergence of finite depth with a non-sober topological modification). Take on $X = \{x, y, z\}$ the finitely deep convergence given by $\lim\{x\}^\uparrow = \{x, y\}$, $\lim\{y\}^\uparrow = \{x, y, z\}$ and $\lim\{z\}^\uparrow = \{y, z\}$. Then by finite depth, $\lim\{x, y\}^\uparrow = \{x, y\}$, $\lim\{x, z\}^\uparrow = \lim\{x, y, z\}^\uparrow = \{y\}$, and $\lim\{y, z\}^\uparrow = \{y, z\}$.



The irreducible filters are $\{x\}^\uparrow$, $\{y\}^\uparrow$, $\{z\}^\uparrow$, $\{x, y\}^\uparrow$ and $\{y, z\}^\uparrow$, x is the only generic point for $\lim\{x\}^\uparrow$ and for $\lim\{x, y\}^\uparrow$, y is the only generic point for $\lim\{y\}^\uparrow$, z is the only generic point for $\lim\{z\}^\uparrow$ and for $\lim\{y, z\}^\uparrow$. Hence this convergence is sober, but its topological modification is antidiscrete, hence is not even T_0 .

As observed in [7], an arbitrary product of (weakly) sober convergence spaces is (weakly) sober and a subspace of a (weakly) sober convergence does not need to be (weakly) sober, though a closed subspace does.

3. MORE ON THE CONVERGENCE SPACE $\text{pt } L$

Proposition 16. *For every convergence lattice L the convergence space $\text{pt } L$ is weakly quasi-sober.*

Proof. Suppose $\lim_{\text{pt } L} \mathcal{U} \in \mathcal{U}$ for some $\mathcal{U} \in \mathbb{U}(\text{pt } L)$. Then $\mathcal{U}^\circ \in \text{pt } L$ and moreover $\lim_{\text{pt } L} \{\mathcal{U}^\circ\}^\uparrow = \lim_{\text{pt } L} \mathcal{U}$. Indeed, \mathcal{U}° is prime for if $\ell \vee m \in \mathcal{U}^\circ$ then, since elements of $\text{pt } L$ are prime filters, $(\ell \vee m)^\bullet = \ell^\bullet \cup m^\bullet \in \mathcal{U}$ and \mathcal{U} is an ultrafilter so either ℓ^\bullet or m^\bullet belongs to \mathcal{U} , that is, either $\ell \in \mathcal{U}^\circ$ or $m \in \mathcal{U}^\circ$. Moreover, as $\lim_{\text{pt } L} \mathcal{U} = (\lim_{\xi} \mathcal{U}^\circ)^\bullet \in \mathcal{U}$ we conclude that $\lim_{\xi} \mathcal{U}^\circ \in \mathcal{U}^\circ$, by definition of \mathcal{U}° . The fact that $\lim_{\text{pt } L} \{\mathcal{U}^\circ\}^\uparrow = \lim_{\text{pt } L} \mathcal{U}$ follows from Lemma 2. \square

Proposition 17. *Let (L, \lim_L) be a convergence lattice satisfying*

$$(3.1) \quad \lim_L x \wedge \lim_L y \in x \cap y \implies x = y$$

for every $x, y \in \text{pt } L$. Then the convergence space $\text{pt } L$ is aas, hence weakly sober.

Proof. Assume (3.1) and suppose $x, y \in \text{pt } L$ with $\lim_{\text{pt } L} \{x\}^\uparrow = \lim_{\text{pt } L} \{y\}^\uparrow$, that is,

$$(\lim_L(\{x\}^\uparrow)^\circ)^\bullet = (\lim_L(\{y\}^\uparrow)^\circ)^\bullet.$$

In view of Lemma 2, $(\lim_L x)^\bullet = (\lim_L y)^\bullet$ so that $\lim_L x \wedge \lim_L y \in x \cap y$. As a result of (3.1), $x = y$.

In view of Propositions 16 and 11, $\text{pt } L$ is not only quasi weakly sober but also weakly sober. \square

An element ℓ of a convergence lattice (L, \lim) is *closed* (e.g., [12]⁴) if $\lim \mathcal{F} \leq \ell$ whenever $\ell \in \mathcal{F}$ (equivalently whenever $\ell \in \mathcal{F}^\#$ where $\mathcal{F}^\# = \{m \in L : \forall f \in \mathcal{F} \ m \wedge f > \perp\}$) and *open* if $\lim \mathcal{F} \wedge \ell > \perp$ implies $\ell \in \mathcal{F}$ ⁵. This is an abstraction of the case $L = (\mathbb{P}X, \lim_\xi)$ in which closed elements are exactly the closed subsets of (X, ξ) and open elements are exactly the open subsets. Let \mathcal{O}_L and \mathcal{C}_L denote the sublattices of L formed by open elements and closed elements respectively.

Lemma 18. *If $\ell \in L$ is closed then ℓ^\bullet is a closed subset of $\text{pt } L$. If $\ell \in L$ is open, then ℓ^\bullet is an open subset of $\text{pt } L$.*

Proof. Let $x \in \lim_{\text{pt } L} \mathcal{F}$ with $\ell^\bullet \in \mathcal{F}$. We need to show that $x \in \ell^\bullet$, equivalently, that $\ell \in x$. Since $\lim_L \mathcal{F}^\circ \in x$ and $\ell \in \mathcal{F}^\circ$ for ℓ closed, we conclude that $\lim_L \mathcal{F}^\circ \leq \ell$, so that $\ell \in x$.

Let $\mathcal{F} \in \mathbb{F}\mathbb{P} \text{pt } L$ with $x \in \lim_{\text{pt } L} \mathcal{F} \cap \ell^\bullet$, that is, $\ell \in x$ and $\lim_L \mathcal{F}^\circ \in x$ so that $\ell \wedge \lim_L \mathcal{F}^\circ \neq \perp$ and ℓ is open, hence $\ell \in \mathcal{F}^\circ$, that is, $\ell^\bullet \in \mathcal{F}$. \square

Proposition 19. *If (L, \lim_L) is a convergence lattice in which*

- (1) $\lim_L \mathcal{F}$ is closed for every $\mathcal{F} \in \mathbb{F}L$ then limits sets are closed in the convergence space $\text{pt } L$.
- (2) $\lim_L x$ is closed for every $x \in \text{pt } L$ then $\lim_{\text{pt } L} \{x\}^\uparrow$ is closed for every $x \in \text{pt } L$.

Proof. Let $\lim_{\text{pt } L} \mathcal{F} \in \mathcal{G} \in \mathbb{F}\mathbb{P}(\text{pt } L)$. By definition $\lim_{\text{pt } L} \mathcal{F} = (\lim_L \mathcal{F}^\circ)^\bullet$ is closed because $\lim_L \mathcal{F}^\circ$ is closed, by Lemma 18. The second part is the particular case where $\mathcal{F} = \{x\}^\uparrow$ for $x \in \text{pt } L$, with the observation that in this case $\mathcal{F}^\circ = x$ by Lemma 2. \square

Corollary 20. *If (L, \lim_L) is a convergence lattice in which $\lim_L x$ is closed whenever $x \in \text{pt } L$, the following are equivalent:*

- (1) $\text{pt } L$ is weakly sober;
- (2) $\text{pt } L$ is antisymmetric;
- (3) $\text{pt } L$ is aas;
- (4) (3.1) for every $x, y \in \text{pt } L$;
- (5) $\lim_L x = \lim_L y \implies x = y$ for every $x, y \in \text{pt } L$.

⁴called *quasi-closed* in [5].

⁵called *fully open* in [9] and a weaker notion than the notion of open element in [5]

Proof. By Propositions 16 and 19, $\text{pt } L$ is a weakly quasi-sober space in which limits of principal ultrafilters are closed. In view of Proposition 11, (1), (2) and (3) are equivalent. That (4) \implies (5) is clear and (4) \implies (3) is Proposition 17. To see (5) \implies (4) suppose that $\lim_L x \wedge \lim_L y \in x \cap y$. Then $\lim_L y \leq \lim_L x$ because $\lim_L x$ is closed and belongs to y , and similarly for the reverse inequality. Hence by (5), $x = y$.

To see that (1) \implies (5) suppose $\lim_L x = \lim_L y$ so that, in view of Lemma 2, $\lim_L(\{x\}^\uparrow)^\circ = \lim_L(\{y\}^\uparrow)^\circ$ so that $\lim_{\text{pt } L}\{x\}^\uparrow = \lim_{\text{pt } L}\{y\}^\uparrow$. By uniqueness of generic point of irreducible ultrafilters in $\text{pt } L$, $x = y$. \square

In the case $L = (\mathbb{P}X, \lim_\xi)$, this means:

Corollary 21. *If (X, ξ) is a convergence space in which limits of irreducible ultrafilters are closed then (X, ξ) is a dense subspace of $\text{pt}(\mathbb{P}X, \lim_\xi)$ which is weakly sober if and only if*

$$(3.2) \quad \lim_\xi \mathcal{U} = \lim_\xi \mathcal{W} \implies \mathcal{U} = \mathcal{W}$$

for every pair \mathcal{U}, \mathcal{W} of irreducible ultrafilters.

We will see with Remark 38 below that the condition (3.2) turns out to be very restrictive.

An alternative approach to force (3.1) is to consider the equivalence relation \sim on $\text{pt } L$ defined by $x \sim y$ if $\lim_L x = \lim_L y$ and consider the quotient set

$$\text{pt}' L := \text{pt } L / \sim,$$

endowed with the quotient convergence. To be explicit, with $q : \text{pt } L \rightarrow \text{pt}' L$ the canonical surjection, this means

$$\begin{aligned} \lim_{\text{pt}' L} \mathcal{F} &= \bigcup_{q[\mathcal{G}] \leq \mathcal{F}} q(\lim_{\text{pt } L} \mathcal{G}) \\ &= \bigcup_{q[\mathcal{G}] \leq \mathcal{F}} q((\lim_L \mathcal{G}^\circ)^\bullet). \end{aligned}$$

Remark 22. Note that we can identify $\text{pt}' L$ with $\{\lim_L x : x \in \text{pt } L\} \subset L$ and in this interpretation, the canonical surjection q is the restriction $\lim'_L : \text{pt } L \rightarrow \text{pt}' L$ of \lim_L to points, when points of L are interpreted as filters on L . Hence we can consider the order induced by L on $\text{pt}' L$ ⁽⁶⁾ which we also denote \leq . Note that $\text{pt}' L$ is not a sublattice of L in general. In contrast the subset \mathcal{C}_L of closed elements of L is a sublattice of L .

Note that

$$(3.3) \quad \forall c \in \mathcal{C}_L \forall x \in \text{pt } L \quad (c \in x \iff \lim_L x \leq c),$$

because $c \in x \implies \lim_L x \leq c$ follows from c closed, and the converse is true whether c is closed or not, because $\lim_L x \in x$. Hence $x \cap \mathcal{C}_L = (\uparrow \lim_L x) \cap \mathcal{C}_L$ whenever $x \in \text{pt } L$.

⁶namely, if $x, y \in \text{pt}' L$, let $x \sqsubseteq y$ denote $\lim_L t \leq \lim_L s$ where $t \in q^{-1}(x)$ and $s \in q^{-1}(y)$, which is well-defined by definition of \sim , but with the interpretation

$$\text{pt}' L = \{\lim_L x : x \in \text{pt } L\} \subset L$$

There is no need to distinguish \sqsubseteq from the order \leq of L .

Lemma 23. *If (L, \lim_L) is a convergence lattice with closed limits, that is, $\lim_L : \mathbb{F}L \rightarrow \mathcal{C}_L$, then*

$$x \in \lim_{\text{pt } L} \mathcal{F} \iff \lim_L x \leq \lim_L \mathcal{F}^\circ,$$

that is,

$$(3.4) \quad \lim_{\text{pt } L} \mathcal{F} = q^{-1}(\text{pt}' L \cap \downarrow \lim_L \mathcal{F}^\circ).$$

Thus

$$(3.5) \quad \lim_{\text{pt}' L} \mathcal{H} = \bigcup_{q[\mathcal{F}] \leq \mathcal{H}} \text{pt}' L \cap \downarrow \lim_L \mathcal{F}^\circ.$$

In particular, $\lim_{\text{pt}' L} \mathcal{H} = \downarrow \lim_{\text{pt}' L} \mathcal{H}$.

Proof. By definition, $x \in \lim_{\text{pt } L} \mathcal{F}$ if and only if $\lim_L \mathcal{F}^\circ \in x$, which is equivalent to $\lim_L x \leq \lim_L \mathcal{F}^\circ$ by (3.3). The formula (3.4) is a rephrasing from which (3.5) follows because q is onto and thus $qq^{-1} = \text{id}_{\text{pt } L}$. \square

In particular, if $\mathcal{H} = \{\lim_L x\}^\uparrow$ is a principal ultrafilter on $\text{pt}' L$ then:

Corollary 24. *If (L, \lim_L) is a convergence lattice with closed limits, then*

$$(3.6) \quad \lim_{\text{pt}' L} \{\lim_L x\}^\uparrow = \text{pt}' L \cap \downarrow \lim_L x.$$

Thus $\text{pt}' L$ is an aas convergence space.

Proof. Note that $\lim_{\text{pt}' L} \{\lim_L x\}^\uparrow = \bigcup_{q[\mathcal{F}] \leq \{\lim_L x\}^\uparrow} \text{pt}' L \cap \downarrow \lim_L \mathcal{F}^\circ$ by Lemma 23 and that $q[\mathcal{F}] \leq \{\lim_L x\}^\uparrow$ means $(\lim_L x)^\bullet \in \mathcal{F}^\#$ because $\lim_L x \in q[F]$ if there is $y \in F$ with $\lim_L y = \lim_L x$ and thus $y \in (\lim_L x)^\bullet \cap F$. As $\lim_L x$ is a closed element of L , $(\lim_L x)^\bullet$ is a closed subset of $\text{pt } L$ by Lemma 18, so that $\lim_{\text{pt } L} \mathcal{F} = (\lim_L \mathcal{F}^\circ)^\bullet \subset (\lim_L x)^\bullet$. Hence if $\lim_L y \in \text{pt}' L \cap \downarrow \lim_L \mathcal{F}^\circ$ then $y \in (\lim_L \mathcal{F}^\circ)^\bullet \subset (\lim_L x)^\bullet$, that is, $\lim_L x \in y$ and $\lim_L x$ is closed thus $\lim_L y \leq \lim_L x$. Consequently, $\lim_L y \in \text{pt}' L \cap \downarrow \lim_L x$. Conversely, if $\lim_L y \leq \lim_L x$ then $\lim_L x \in y$ and thus $\mathcal{F} = \{y\}^\uparrow$ satisfies $q[\mathcal{F}] \leq \{\lim_L x\}^\uparrow$ and $\lim_L y = \lim_L \mathcal{F}^\circ$ because $\mathcal{F}^\circ = y$ by Lemma 18, which completes the proof of (3.6).

As a result of (3.6), if $\lim_{\text{pt}' L} \{\lim_L x\}^\uparrow = \lim_{\text{pt}' L} \{\lim_L y\}^\uparrow$ then $\lim_L x \leq \lim_L y$ and $\lim_L y \leq \lim_L x$ so that $\lim_L x = \lim_L y$. Hence, $\text{pt}' L$ is aas. \square

Lemma 25. *If U is an open or a closed subset of $\text{pt } L$ then U is saturated, that is, $U = q^{-1}(q(U))$. In particular, if ℓ is either an open or a closed element of L then ℓ^\bullet is saturated.*

Proof. Suppose U is open and $x \in q^{-1}(q(U))$ that is, $\lim_L x \in q(U)$, that is, there is $t \in U$ with $\lim_L t = \lim_L x$. As

$$\lim_{\text{pt } L} \{t\}^\uparrow = (\lim_L(\{t\}^\uparrow))^\bullet = (\lim_L t)^\bullet = (\lim_L x)^\bullet = \lim_{\text{pt } L} \{x\}^\uparrow,$$

we conclude that $\lim_{\text{pt } L} \{x\}^\uparrow \cap U \neq \emptyset$ so that $U \in \{x\}^\uparrow$ because U is open, that is, $x \in U$. If U is closed instead, then $\lim_{\text{pt } L} \{t\}^\uparrow = \lim_{\text{pt } L} \{x\}^\uparrow \subset U$ and we conclude similarly. The case of ℓ^\bullet follows from Lemma 18. \square

Corollary 26. *$q = \lim'_L : \text{pt } L \rightarrow \text{pt}' L$ is an open and closed map (sends open sets to open sets and closed sets to closed sets). Moreover, the initial topology for \lim'_L is the topological modification of $\text{pt } L$:*

$$\text{T}(\text{pt } L) = q^-(\text{T } \text{pt}' L) = q^-(\text{T } q(\text{pt } L)).$$

Lemma 27. *If c is a closed element of (L, \lim_L) then*

$$q^{-1}(\text{pt}' L \setminus \downarrow c) = \text{pt} L \setminus c^\bullet.$$

Proof. $x \in q^{-1}(\text{pt}' L \setminus \downarrow c)$ if and only if $\lim_L x \not\leq c$, which, in view of (3.3), is equivalent to $c \notin x$, that is, to $x \in \text{pt} L \setminus c^\bullet$. \square

Note that if $L = (\mathbb{P}X, \lim_\xi)$ then for every open subset U of $\text{pt} L$ there is an open element ℓ of L with $U = \ell^\bullet$ and similarly for closed sets. Indeed:

Proposition 28. *Let $L = (\mathbb{P}X, \lim_\xi)$. Every subset U of $\text{pt} L$ that is either open or closed satisfies $U = (U \cap X)^\bullet$.*

Proof. Assume that U is open. Let $\mathcal{U} \in (U \cap X)^\bullet$, that is, $(U \cap X) \in \mathcal{U}$. As $\lim_{\text{pt} L} \{\mathcal{U}\}^\uparrow = (\lim_\xi(\{\mathcal{U}\}^\uparrow)^\circ)^\bullet = (\lim_\xi \mathcal{U})^\bullet$ and there is $x \in \lim_\xi \mathcal{U} \cap U \cap X$, we conclude that $\{x\}^\uparrow \in (\lim_\xi \mathcal{U})^\bullet \cap U$. Hence $\lim_{\text{pt} L} \{\mathcal{U}\}^\uparrow \cap U \neq \emptyset$ and U is $\text{pt} L$ -open so that $U \in \{\mathcal{U}\}^\uparrow$, that is, $\mathcal{U} \in U$. Conversely, if $\mathcal{U} \in U$, since $\mathcal{U} \in \lim_{\text{pt} L} h[\mathcal{U}]$ where $h : X \rightarrow \text{pt} L$ is $h(x) = \{x\}^\uparrow$ and U is open, $U \cap X \in h[\mathcal{U}]$, that is, $\mathcal{U} \in (U \cap X)^\bullet$.

Assume now that U is closed. If $\mathcal{U} \in U$ then $\lim_{\text{pt} L} \{\mathcal{U}\}^\uparrow = (\lim_\xi \mathcal{U})^\bullet \subset U$ and thus $\lim_\xi \mathcal{U} \subset U \cap X$ so that $U \cap X \in \mathcal{U}$, that is, $\mathcal{U} \in (U \cap X)^\bullet$. Conversely, if $\mathcal{U} \in (U \cap X)^\bullet$, that is, $U \cap X \in \mathcal{U}$, then $\mathcal{U} \in \lim_{\text{pt} L} h[\mathcal{U}] \subset U$ so that $\mathcal{U} \in U$. \square

Let us say that a convergence lattice (L, \lim_L) has enough closed elements if for every closed subset C of $\text{pt} L$ there is a closed element $c \in L$ with $C = c^\bullet$. We similarly say that (L, \lim_L) has enough open elements if every open subset O of $\text{pt} L$ is of the form u^\bullet for some open element u of L . By Proposition 28, $L = (\mathbb{P}X, \lim_\xi)$ always has enough closed elements and enough open elements.

Theorem 29. *If (L, \lim_L) is a convergence lattice with enough closed elements, then the topological modification of $\text{pt}' L$ is given by the open sets*

$$\{\text{pt} L' \setminus \downarrow c : c \in \mathcal{C}_L\}.$$

Proof. In view of Lemma 27, sets of the form $\text{pt} L' \setminus \downarrow c$ for a closed element $c \in L$ are $\text{pt}' L$ -open, because their preimages under q are $\text{pt} L$ -open by Lemma 27. If now V is $\text{pt}' L$ -open, that is, $q^{-1}(V)$ is $\text{pt} L$ -open, then $\text{pt} L \setminus q^{-1}(V)$ is closed, hence there is a closed element c of L with $q^{-1}(V) = \text{pt} L \setminus c^\bullet$. In view of Lemma 27, $q^{-1}(V) = q^{-1}(\text{pt}' L \setminus \downarrow c)$ and q is surjective, so that $V = \text{pt}' L \setminus \downarrow c$. \square

Remark 30. Recall that the upper topology of a poset (X, \leq) is that in which sets $\{X \setminus \downarrow x : x \in X\}$ form a subbase of open sets (e.g., [6]). Note that when L has closed limits (that is, $\lim_L : \mathbb{F}L \rightarrow \mathcal{C}_L$), then $\text{pt}' L \subset \mathcal{C}_L$ and the topology induced on $\text{pt}' L$ by the upper topology of \mathcal{C}_L is given by the subbase $\{\text{pt} L' \setminus \downarrow c : c \in \mathcal{C}_L\}$, so that, in view of Theorem 29, it coincides with the quotient topology induced by $\lim'_L : \text{pt} L \rightarrow \text{pt}' L$ where $\text{pt} L$ carries the topological modification of its standard convergence.

4. \mathcal{Z} -REGULARITY AND STRUCTURE OF $\text{pt}' L$

If $\mathcal{Z} \subset L$ we say that (L, \lim_L) is \mathcal{Z} -regular if

$$\lim_L \mathcal{F} = \lim_L \uparrow (\mathcal{F} \cap \mathcal{Z})$$

for every $\mathcal{F} \in \mathbb{F}L$. This is a generalization of the notion defined for convergence spaces in [4, Section VII.2], namely, if $\mathcal{Z} \subset \mathbb{P}X$, then (X, ξ) is \mathcal{Z} -regular if $\lim_\xi \mathcal{F} = \lim_\xi (\mathcal{F} \cap \mathcal{Z})^\uparrow$ for every filter $\mathcal{F} \in \mathbb{F}\mathbb{P}X$.

Note that a convergence space (X, ξ) is \mathcal{O}_ξ -regular if and only if it is topological⁽⁷⁾, equivalently, the convergence lattice $L = (\mathbb{P}X, \lim_\xi)$ is \mathcal{O}_L -regular.

Note that:

Proposition 31. *Let $f : (X, \xi) \rightarrow (Y, \tau)$ be a continuous, onto, and open map. If ξ is topological, so is τ .*

Proof. We show τ is \mathcal{O} -regular. To this end, let $y \in \lim_\tau \mathcal{G}$. In view of [4, Exercise XV.1.29], $\mathcal{G} \geq f[\mathcal{N}_\xi(x)]$ for every $x \in f^{-1}(y)$. Because f is open $f[\mathcal{N}_\xi(x)] \subset (\mathcal{G} \cap \mathcal{O}_\tau)^\uparrow$ and $y \in \lim_\tau f[\mathcal{N}_\xi(x)]$ by continuity. Hence, $y \in \lim_\tau (\mathcal{G} \cap \mathcal{O}_\tau)^\uparrow$. \square

Similarly a convergence space (X, ξ) is topologically regular (in the sense of [4, Section VI.4]) if and only if it is \mathcal{C}_ξ -regular, equivalently, the convergence lattice $L = (\mathbb{P}X, \lim_\xi)$ is \mathcal{C}_L -regular.

In [5], a convergence lattice L is called *classical* if it is regular with respect to the set of complemented elements of L (see [5, Lemma 2.29]). In [12] and [9], $**$ -regular convergence frames are considered. They are those convergence frames that are \mathcal{Z} -regular for the set $\mathcal{Z} = \{\ell \in L : \ell = \ell^{**}\}$, where ℓ^* denotes the pseudocomplement of ℓ (note that every element of a frame has a pseudocomplement).

Lemma 32. *Let (L, \lim_L) be a convergence lattice. Then $\text{pt } L$ is a \mathcal{Z} -regular convergence space for $\mathcal{Z} = \{A \subset \text{pt } L : \exists \ell \in L (A = \ell^\bullet)\}$. Moreover $\text{pt}' L$ is $q(\mathcal{Z})$ -regular where $q(\mathcal{Z}) = \{q(A) : A \in \mathcal{Z}\}$.*

Proof. \mathcal{Z} -regularity of $\text{pt } L$ follows from $\lim_{\text{pt } L} \mathcal{F} = (\lim_L \mathcal{F}^\circ)^\bullet$ and $\mathcal{F}^\circ = ((\mathcal{F} \cap \mathcal{Z})^\uparrow)^\circ$. Now,

$$\begin{aligned} \lim_{\text{pt}' L} \mathcal{F} &= \bigcup_{q[\mathcal{G}] \leq \mathcal{F}} q(\lim_{\text{pt } L} \mathcal{G}) \\ &= \bigcup_{q[\mathcal{G}] \leq \mathcal{F}} q(\lim_{\text{pt } L} (\mathcal{G} \cap \mathcal{Z})^\uparrow) \end{aligned}$$

and $q(\mathcal{G} \cap \mathcal{Z}) \subset \mathcal{F} \cap \mathcal{Z} \subset \mathcal{F}$ so that

$$\lim_{\text{pt}' L} \mathcal{F} = \bigcup_{q[\mathcal{G} \cap \mathcal{Z}] \subset \mathcal{F} \cap \mathcal{Z}} q(\lim_{\text{pt } L} (\mathcal{G} \cap \mathcal{Z})^\uparrow) \subset \lim_{\text{pt}' L} (\mathcal{F} \cap \mathcal{Z})^\uparrow.$$

\square

Lemma 33. *If (L, \lim_L) is \mathcal{Z} -regular where ℓ^\bullet is saturated for every $\ell \in \mathcal{Z}$, then*

$$\lim_{\text{pt}' L} \mathcal{U} = q(\lim_{\text{pt } L} q^{-1}[\mathcal{U}]) = q(\lim_{\text{pt } L} \mathcal{W}),$$

for every ultrafilter \mathcal{U} on $\text{pt}' L$ on every $\mathcal{W} \in \mathbb{U}(q^{-1}[\mathcal{U}])$.

Proof. Let L be \mathcal{Z} -regular. Because \mathcal{U} is an ultrafilter, $\lim_{\text{pt}' L} \mathcal{U} = \bigcup_{\mathcal{W} \in \mathbb{U}(q^{-1}[\mathcal{U}])} q((\lim_L \mathcal{W}^\circ)^\bullet)$. Since $\lim_L \mathcal{W}^\circ = \lim_L \uparrow (\mathcal{W}^\circ \cap \mathcal{Z})$ by \mathcal{Z} -regularity, it is enough to show that $\mathcal{W}^\circ \cap \mathcal{Z} = \mathcal{Z} \cap (q^{-1}[\mathcal{U}])^\circ$ whenever $\mathcal{W} \in \mathbb{U}(q^{-1}[\mathcal{U}])$. This follows from the observation that $\ell \in \mathcal{W}^\circ \cap \mathcal{Z}$ if and only if $\ell \in \mathcal{Z}$ and $\ell^\bullet \in \mathcal{W}$, and ℓ^\bullet is saturated and thus $\ell^\bullet \in q^{-1}[q[\mathcal{W}]] = q^{-1}[\mathcal{U}]$. \square

⁷Indeed, it is clear that a topological space is \mathcal{O}_ξ -regular, and conversely, if ξ is \mathcal{O}_ξ -regular, then

$$x \in \lim\{x\}^\uparrow = \lim(\{x\}^\uparrow \cap \mathcal{O}_\xi)^\uparrow = \lim \mathcal{N}_\xi(x),$$

so that ξ is topological.

In view of Lemma 25, Lemma 33 applies in particular if $\mathcal{Z} = \mathcal{O}_L$ or $\mathcal{Z} = \mathcal{C}_L$.

Proposition 34. *If (L, \lim_L) is \mathcal{O}_L -regular and has enough open elements, then $\text{pt } L$ is topological. If $\text{pt } L$ is topological, so is $\text{pt}' L$.*

Proof. Since L has enough open sets,

$$\begin{aligned} \lim_{\text{pt } L} (\mathcal{F} \cap \mathcal{O}_{\text{pt } L})^\uparrow &= \lim_{\text{pt } L} (\mathcal{F} \cap \{u^\bullet : u \in \mathcal{O}_L\})^\uparrow \\ &= (\lim_L ((\mathcal{F} \cap \{u^\bullet : u \in \mathcal{O}_L\})^\uparrow)^\circ)^\bullet \\ &= (\lim_L \uparrow (\mathcal{O}_L \cap \mathcal{F}^\circ))^\bullet \\ &= (\lim_L \mathcal{F}^\circ)^\bullet = \lim_{\text{pt}' L} \mathcal{F}, \end{aligned}$$

because L is \mathcal{O}_L -regular. Hence $\text{pt } L$ is \mathcal{O} -regular, that is, topological. As q is a continuous, onto, and open map by Corollary 26, $\text{pt}' L$ is also topological by Proposition 31. \square

Corollary 35. *If L is \mathcal{O}_L -regular, has enough open sets, and has closed limits then $\text{pt}' L$ is a sober topological space.*

Proof. Let \mathcal{U} be an irreducible ultrafilter on $\text{pt}' L$, that is, $\lim_{\text{pt}' L} \mathcal{U} \in \mathcal{U}$. In view of Lemma 33, $\lim_{\text{pt}' L} \mathcal{U} = q(\lim_{\text{pt } L} \mathcal{W})$ for every $\mathcal{W} \in \mathbb{U}(q^{-1}[\mathcal{U}])$, so that

$$q^{-1}(\lim_{\text{pt}' L} \mathcal{U}) = q^{-1}(q(\lim_{\text{pt } L} \mathcal{W})) = \lim_{\text{pt } L} \mathcal{W} \in q^{-1}[\mathcal{U}] \subset \mathcal{W}$$

because each $\lim_{\text{pt } L} \mathcal{W}$ is closed (because L , hence also $\text{pt } L$, has closed limits by Proposition 19) hence saturated by Lemma 25. In other words, each \mathcal{W} is an irreducible ultrafilter so that $\lim_{\text{pt } L} \mathcal{W} = \lim_{\text{pt } L} \{x\}^\uparrow = (\lim_L x)^\bullet$ for some $x \in \text{pt } L$ because, in view of Proposition 25, $\text{pt } L$ is weakly quasi-sober. Thus $\lim_{\text{pt}' L} \mathcal{U} = q((\lim_L x)^\bullet) = \lim_{\text{pt}' L} \{\lim_L x\}^\uparrow$ by Corollary 24. Hence $\text{pt}' L$ is weakly quasi-sober and aas by Corollary 24, hence weakly sober. In view of Proposition 11 it is sober because it is topological. \square

In view of Lemma 18 and Proposition 28, $(\mathbb{P}X, \lim_\xi)$ always has enough open elements and enough closed elements. Moreover, it has closed limits when ξ has closed limits, in particular if ξ is topological. Hence, Proposition 34 and Corollary 35 apply to the effect that:

Corollary 36. *If (X, ξ) is a topological space and $L = (\mathbb{P}X, \lim_\xi)$ then $\text{pt } L$ is topological and weakly quasi-sober and $\text{pt}' L$ is a topological sober space.*

In fact, in this case $\text{pt}' L$ is exactly the *sobrification* of the topological space X , as we will see in Section 5 below.

5. SOBRIFICATION AND THE SPACE $\text{pt}(\mathbb{P}X, \lim_\xi)$

Recall that a closed subset C of a convergence space is *irreducible* if $C \subset D$ or $C \subset F$ whenever $C \subset D \cup F$ where D and F are closed. Let us call C *c-irreducible* if it satisfies this property even if C is not necessarily closed. Note that C is c-irreducible if and only if for every open sets O and U intersecting C , $O \cap U \cap C \neq \emptyset$. Obviously, C is c-irreducible if and only if $\text{cl } C$ is an irreducible closed set.

Lemma 37. *In a convergence space, limits of irreducible filters are c-irreducible and the closure of every c-irreducible set is the limit of a \mathbb{T} - ξ -irreducible ultrafilter.*

Proof. If $\lim_{\xi} \mathcal{F} \in \mathcal{F}$ then $\lim_{\xi} \mathcal{F}$ is a c-irreducible set. Indeed, if $\lim_{\xi} \mathcal{F} \subset D \cup F$ then $D \cup F \in \mathcal{F} \subset \mathcal{F}^{\#}$ so that either $D \in \mathcal{F}^{\#}$ or $F \in \mathcal{F}^{\#}$. Hence $\lim_{\xi} \mathcal{F} \subset D$ or $\lim_{\xi} \mathcal{F} \subset F$ because D and F are closed.

If C is a c-irreducible set then $\{O \in \mathcal{O}_{\xi} : O \cap C \neq \emptyset\}$ is a filter base because C is c-irreducible. Let \mathcal{U} be an ultrafilter containing this filter base. Note that \mathcal{U} is finer than all $\mathcal{N}_{\xi}(x)$ for $x \in C$, so that $C \subset \lim_{\mathbb{T}\xi} \mathcal{U}$ hence $\text{cl} C \subset \lim_{\mathbb{T}\xi} \mathcal{U}$. Note that \mathcal{U} is $\mathbb{T}\xi$ -irreducible, for if $\lim_{\mathbb{T}\xi} \mathcal{U} \notin \mathcal{U}$ then $X \setminus C \in \mathcal{U}$ which is not the case. Moreover, if $t \notin \text{cl} C$, there is $O = X \setminus \text{cl} C \in \mathcal{N}(t)$ that is disjoint from C hence does not belong to \mathcal{U} . In other words, $\lim_{\mathbb{T}\xi} \mathcal{U} = \text{cl} C$. \square

Remark 38. In particular, in a topological space every irreducible closed set C is the limit of an irreducible ultrafilter—in fact, of many, for every ultrafilter containing the filter base $\{O \in \mathcal{O}_{\xi} : O \cap C \neq \emptyset\}$ satisfies this condition. As a result, if X is a T_0 topological space then (3.2) is equivalent to every irreducible ultrafilters being principal, a property characterized for topological spaces in [7, Theorem 2.2] and more generally for (some) convergence spaces in Corollary 47 below to be equivalent with X being both sober and T_D . To see that (3.2) implies that irreducible ultrafilters are principal, note that if there is a free irreducible ultrafilter on a topological space, then $C = \lim \mathcal{U}$ is an infinite irreducible closed set and the non-maximal filter-base $\{O \in \mathcal{O}_{\xi} : O \cap C \neq \emptyset\}$ admits one, hence many finer free ultrafilters all of which admit C as their limit, so that (3.2) fails. Conversely, if all irreducible ultrafilters are principal, then every irreducible closed set has a generic point, which is necessarily unique in a T_0 topological space, so that (3.2) is satisfied.

Corollary 39. *If \mathcal{F} is an irreducible filter on a convergence space (X, ξ) then*

$$\text{cl}_{\xi}(\lim_{\xi} \mathcal{F}) = \lim_{\mathbb{T}\xi} \mathcal{F}.$$

Proof. We only need to show that $\lim_{\mathbb{T}\xi} \mathcal{F} \subset \text{cl}_{\xi}(\lim_{\xi} \mathcal{F})$ for the reverse inclusion is always true. In view of Lemma 37, $\lim_{\xi} \mathcal{F}$ is c-irreducible and thus, for every ultrafilter \mathcal{U} of the filter-base $\{O \in \mathcal{O}_{\xi} : O \cap \lim_{\xi} \mathcal{F} \neq \emptyset\}$, $\text{cl}_{\xi}(\lim_{\xi} \mathcal{F}) = \lim_{\mathbb{T}\xi} \mathcal{U}$. Note that the condition $O \cap \lim_{\xi} \mathcal{F} \neq \emptyset$ implies that $O \in \mathcal{F}$, so that \mathcal{F} contains this filter-base. \square

Recall (e.g., [8]) that the *sobrification* of a T_0 topological space X is the set ${}^s X$ of irreducible closed subsets of X endowed with the topology $\{{}^s O : O \in \mathcal{O}_{\xi}\}$ where ${}^s O = \{C \in {}^s X : C \cap O \neq \emptyset\}$ ⁸. Because X is T_0 the map $e : X \rightarrow {}^s X$ defined by $e(x) = \text{cl}\{x\}$ is one-to-one. It is a dense embedding because ${}^s O \cap e(X) = e(O)$.

Note that in view of Lemma 37 we can identify elements of $\text{pt}' L$ (equivalence classes of irreducible ultrafilters with the same limit) and elements of ${}^s X$. In fact, since the topology of ${}^s X$ is the topology induced on ${}^s X$ by the upper topology of the space of closed subsets of X ordered by inclusion because

$${}^s O = \{C \in {}^s X : C \cap O \neq \emptyset\} = {}^s X \setminus \downarrow (X \setminus O),$$

we conclude via Theorem 29 that ${}^s X$ is homeomorphic to $\text{pt}' L$ endowed with the quotient topology induced by $q = \lim'_L : \text{pt} L \rightarrow \text{pt}' L$, but this is the structure of $\text{pt}' L$, since $\text{pt}' L$ is topological by Corollary 36.

⁸to see that this defines indeed a topology, note that ${}^s(\bigcup_{O \in \mathcal{B}} O) = \bigcup_{O \in \mathcal{B}} {}^s O$ and that ${}^s(O \cap U) = {}^s O \cap {}^s U$ because an irreducible closed sets intersects $O \cap U$ whenever it intersects open sets O and U .

Theorem 40. *Let (X, ξ) be a topological and let $L = (\mathbb{P}X, \lim_\xi)$. Then $\text{pt}' L$ is the sobrification ${}^s X$ of X , and $e = q \circ h$ is a dense embedding in the diagram below whenever (X, ξ) is T_0 :*

$$\begin{array}{ccc} X & \xrightarrow{h: x \mapsto \{x\}^\uparrow} & \text{pt}(\mathbb{P}X, \lim_\xi) \\ e: x \mapsto \text{cl}\{x\} \downarrow & & \downarrow q \\ {}^s X & \xlongequal{\quad} & \text{pt}'(\mathbb{P}X, \lim_\xi) \end{array}$$

6. THE AXIOM T_D IN CONVERGENCE SPACES

Recall that a topological space X is T_D if for every $x \in X$ there is $U \in \mathcal{O}(x)$ with $U \setminus \{x\}$ open (e.g., [13]).

Lemma 41. *In a topological space X , the following are equivalent:*

- (1) X is T_D ;
- (2) for every $x \in X$ there is $U \in \mathcal{O}(x)$ with

$$\lim\{x\}^\uparrow \cap U = \{x\};$$

- (3) for every $x \in X$ and every filter \mathcal{F} on X that converges to x , there is $A \in \mathcal{F}$ with

$$\lim\{x\}^\uparrow \cap A \subset \{x\};$$

- (4) for every $x \in X$ and every ultrafilter \mathcal{U} on X that converges to x , there is $A \in \mathcal{U}$ with

$$\lim\{x\}^\uparrow \cap A \subset \{x\}.$$

Proof. (1) \implies (2): If X is T_D and $x \in X$, pick $U \in \mathcal{O}(x)$ with $U \setminus \{x\}$ open. Then $\lim\{x\} \cap U = \{x\}$ because $t \in \lim\{x\} \cap (U \setminus \{x\})$ would imply $t \in \{x\}$ which is impossible.

(2) \implies (3): If $x \in \lim \mathcal{F}$, then $\mathcal{F} \geq \mathcal{O}(x)$ and we can take the U from (1) as A .

(3) \implies (4) is obvious.

(4) \implies (1): For every $\mathcal{U} \in \mathbb{U}(\mathcal{N}(x))$, there is $A_{\mathcal{U}} \in \mathcal{U}$ with $\lim\{x\} \cap A_{\mathcal{U}} \subset A_{\mathcal{U}}$. Hence there is a finite subset F of $\mathbb{U}(\mathcal{N}(x))$ with $\bigcup_{\mathcal{U} \in F} A_{\mathcal{U}} \in \mathcal{N}(x)$. Let $U \in \mathcal{O}(x)$ with $U \subset \bigcup_{\mathcal{U} \in F} A_{\mathcal{U}}$. Then $U \setminus \{x\}$ is open. Indeed, if $t \in \lim \mathcal{U} \cap (U \setminus \{x\})$ for some ultrafilter \mathcal{U} then $U \in \mathcal{U}$ because U is open. If $U \setminus \{x\} \notin \mathcal{U}$ then $U^c \cup \{x\} \in \mathcal{U}$, hence $\mathcal{U} = \{x\}^\uparrow$ and (4) ensures a contradiction. \square

Note that (3) and (4) make sense for general convergences and are equivalent ⁽⁹⁾. Thus we call a *convergence space* T_D if it satisfies (3), equivalently (4). As far as I know, this is the first time this axiom is introduced for convergence spaces.

Just as for topological space, T_D lies between T_0 and T_1 : T_1 obviously implies T_D as in this case $\lim\{x\}^\uparrow = \{x\}$ for every x . On the other hand, if X is T_D and

9

Proof. If ξ satisfies (4) and $x \in \lim \mathcal{F}$ then $x \in \lim \mathcal{U}$ for every $\mathcal{U} \in \mathbb{U}(\mathcal{F})$ so that there is $A_{\mathcal{U}} \in \mathcal{U}$ with $\lim\{x\}^\uparrow \cap A_{\mathcal{U}} \subset \{x\}$. Hence there is a finite subset F of $\mathbb{U}(\mathcal{F})$ with $A = \bigcup_{\mathcal{U} \in F} A_{\mathcal{U}} \in \mathcal{F}$. Then

$$\lim\{x\} \cap A = \bigcup_{\mathcal{U} \in F} \lim\{x\} \cap A_{\mathcal{U}} \subset \{x\}.$$

\square

$\{\mathcal{F} \in \mathbb{F}X : x \in \lim \mathcal{F}\} = \{\mathcal{F} \in \mathbb{F}X : t \in \lim \mathcal{F}\}$ then $\{x, t\} \subset \lim\{x\}^\uparrow \cap \lim\{t\}^\uparrow$ so by T_D there is $A \in \{x\}^\uparrow$ with $\lim\{t\}^\uparrow \cap A \subset \{t\}$ which ensures $x = t$ because $x \in \lim\{t\}^\uparrow \cap A$.

Note that in all the examples of Section 2, we have $x \neq y$ with $\{x, y\} \in \lim\{x\}^\uparrow \cap \lim\{y\}^\uparrow$ so that $\{y\}^\uparrow$ is an ultrafilter converging to x and $y \in \lim\{x\}^\uparrow$. Hence $y \in \lim\{x\}^\uparrow \cap A$ for every $A \in \{y\}^\uparrow$ and the space is *not* T_D .

Preservation properties for T_D easily extend from topologies to convergences with this definition:

Proposition 42. *If $\xi \leq \tau$ and ξ is T_D , so is τ ; a subspace of a T_D convergence space is T_D ; a finite product of T_D convergence spaces is T_D .*

Proof. If $x \in \lim_\tau \mathcal{F} \subset \lim_\xi \mathcal{F}$ there is $A \in \mathcal{F}$ with $\lim_\xi \{x\}^\uparrow \cap A \subset \{x\}$ and the conclusion follows from $\lim_\tau \{x\}^\uparrow \subset \lim_\xi \{x\}^\uparrow$. If $A \subset X$ with inclusion map $i : A \rightarrow X$ and $x \in A \cap \lim_{\xi_A} \mathcal{F}$, $x \in \lim_\xi i[\mathcal{F}]$ so there is $F \in i[\mathcal{F}]$ with $\lim_\xi \{x\}^\uparrow \cap F \subset \{x\}$ so that $\lim_{\xi_A} \{x\}^\uparrow \cap (A \cap F) \subset \{x\}$ and (A, ξ_A) is T_D . Suppose $(X_1, \xi_1), \dots, (X_n, \xi_n)$ are T_D convergence spaces and let $(x_1, \dots, x_n) \in \lim_{\prod_{i=1}^n \xi_i} \mathcal{F}$, that is $x_i \in \lim_{\xi_i} p_i[\mathcal{F}]$ for every i , where $p_i : \prod_{i=1}^n X_i \rightarrow X_i$ is the projection. There is $A_i \in p_i[\mathcal{F}]$ with $\lim_{\xi_i} \{x_i\}^\uparrow \cap A_i \subset \{x_i\}$ so that $\prod_{i=1}^n A_i \in \prod_{i=1}^n p_i[\mathcal{F}] \subset \mathcal{F}$ and then $\lim_{\prod_{i=1}^n \xi_i} \{(x_1, \dots, x_n)\}^\uparrow \cap \prod_{i=1}^n A_i \subset \{(x_1, \dots, x_n)\}$. \square

On the other hand, a countable product of topological T_D spaces (e.g., countably many copies of the Sierpiński space) may fail to be T_D (See e.g., [1]).

However:

- Lemma 43.** (1) *A T_D convergence space is antisymmetric.*
 (2) *Moreover, if the space is finite the converse is true, that is, if \rightarrow is antisymmetric then the convergence is T_D .*
 (3) *If $\lim\{x\}^\uparrow$ is closed for every x then \rightarrow is transitive. In particular if the convergence is S_0 then \rightarrow is transitive, and conversely if the space is finite and of finite depth.*

Hence when the convergence is T_D and limits of principal ultrafilters are closed, we have a *specialization order* defined by $x \geq y$ if $y \in \lim\{x\}^\uparrow$.

Proof. 1. Assume there are points $x \neq y$ with $\{x, y\} \subset \lim\{x\}^\uparrow \cap \lim\{y\}^\uparrow$. Then $y \in \lim\{x\}^\uparrow \cap A$ for every $A \in \{y\}^\uparrow$ and the convergence is not T_D .

2. Conversely for a finite space, if the convergence is not T_D there is a (principal because the space is finite) ultrafilter $\{y\}^\uparrow$ converging to x such that $\lim\{x\}^\uparrow \cap A \cap (X \setminus \{x\}) \neq \emptyset$ for every $A \in \{y\}^\uparrow$, in particular for $A = \{y\}$. Hence $y \neq x$ and $y \in \lim\{x\}^\uparrow$ and thus $\{x, y\} \subset \lim\{x\}^\uparrow \cap \lim\{y\}^\uparrow$.

3. That \rightarrow is transitive if $\lim\{x\}^\uparrow$ is closed for every x is clear. For the converse for a finite space, assume $\mathcal{F} = \{F\}^\uparrow$ converges to x and $\{x\}^\uparrow$ converges to y . For every $t \in F$, $t \rightarrow x$ and $x \rightarrow y$, so $t \rightarrow y$ by transitivity. By finite depth, $y \in \bigcap_{t \in F} \lim\{t\}^\uparrow = \lim \bigcap_{t \in F} \{t\}^\uparrow = \lim\{F\}^\uparrow$. Hence, the space is S_0 and thus limits of principal ultrafilters are closed. \square

Note that in view of Lemma 43, a *finite*, finitely deep, weakly sober and non-sober convergence like Example 14 cannot be T_D . This is in fact general:

Proposition 44. *In a T_D weakly sober space every irreducible ultrafilter is principal. If moreover the convergence is of finite depth, then the only irreducible filters are principal ultrafilters and thus the convergence is also sober.*


Proof. If ξ is weakly sober and \mathcal{U} is an irreducible filter, then there is a (unique) x with $\lim \mathcal{U} = \lim\{x\}^\uparrow \in \mathcal{U}$. As ξ is T_D there is $U \in \mathcal{U}$ with $\lim\{x\}^\uparrow \cap U \subset \{x\}$. As a result, $\{x\} \in \mathcal{U}$ and \mathcal{U} is principal. Hence if \mathcal{F} is irreducible, every ultrafilter finer than \mathcal{F} is also irreducible, hence principal. As a result, \mathcal{F} is the principal filter of a finite set F . If the convergence is moreover of finite depth,

$$\lim \mathcal{F} = \lim\{F\}^\uparrow = \bigcap_{x \in F} \lim\{x\}^\uparrow \in \mathcal{F} = \{F\}^\uparrow.$$

By Lemma 43 (1), F is a singleton. The space is then sober. \square

In view of Proposition 42, if $T\xi$ is T_D so is the finer convergence ξ , but the converse is false:

Example 45 (A finitely deep T_D convergence whose topological modification is not T_D). Consider on $X = \{x, y, z\}$ the finitely deep convergence determined by $\lim\{x\}^\uparrow = \{x, y\}$, $\lim\{y\}^\uparrow = \{y, z\}$ and $\lim\{z\}^\uparrow = \{z, x\}$.



This convergence is T_D By Lemma 43, but its topological modification is antidiscrete, hence not even T_0 .

7. PROBLEM 4

Theorem 46. *Let (X, ξ) be a convergence space.*

- (1) *If ξ is weakly sober and T_D then every irreducible filter is principal.*
- (2) *If ξ is S_0 and T_0 the converse is true, that is, ξ is weakly sober and T_D whenever every irreducible ultrafilter is principal.*

Proof. (1) is Proposition 44.

(2) if ξ is T_0 and S_0 and every irreducible ultrafilter is principal then ξ is weakly sober: if $\lim \mathcal{U} \in \mathcal{U}$ there is x with $\mathcal{U} = \{x\}^\uparrow$ and thus $\lim \mathcal{U} = \lim\{x\}^\uparrow$. If $t \neq x$ satisfies $\lim\{t\}^\uparrow = \lim\{x\}^\uparrow$, by T_0 there is $\mathcal{F} \in \mathbb{F}X$ with $\text{card}(\lim \mathcal{F} \cap \{t, x\}) = 1$, which is not compatible with S_0 . Moreover, ξ is T_D . Suppose to the contrary that there is $x \in X$ and $\mathcal{U} \in \mathbb{U}X$ with $x \in \lim \mathcal{U}$ such that $\lim\{x\}^\uparrow \cap U \cap (X \setminus \{x\}) \neq \emptyset$ for every $U \in \mathcal{U}$. In particular, $\lim\{x\}^\uparrow \in \mathcal{U}^\# = \mathcal{U}$ and $\lim\{x\}^\uparrow \subset \lim \mathcal{U}$ because ξ is S_0 . Hence $\lim \mathcal{U} \in \mathcal{U}$ and \mathcal{U} is irreducible, hence principal, that is, $\mathcal{U} = \{t\}^\uparrow$ and $\lim\{x\}^\uparrow \subset \lim\{t\}^\uparrow$. Hence $x \in \lim\{t\}^\uparrow$ and $t \in \lim\{x\}^\uparrow$ by taking $U = \{t\}$ in $\lim\{x\}^\uparrow \cap U \cap (X \setminus \{x\}) \neq \emptyset$, the argument above for the uniqueness of a generic point applies to the effect that $x = t$. But that is not compatible with $\lim\{x\}^\uparrow \cap U \cap (X \setminus \{x\}) \neq \emptyset$ for $U = \{t\} = \{x\}$ and we have a contradiction. \square

Corollary 47. *Let (X, ξ) be a S_0 and T_0 convergence space of finite depth. The following are equivalent:*

- (1) *every irreducible ultrafilter is principal;*
- (2) *every irreducible filter is principal;*
- (3) *ξ is weakly sober and T_D ;*
- (4) *ξ is sober and T_D ;*
- (5) *Every subspace of ξ is sober;*

- (6) If $\theta \geq \xi$ then θ is sober;
(7) $\text{pt}_{\mathbf{Lat}}(\mathbb{P}X, \lim_{\xi})$ is homeomorphic to (X, ξ) .

Proof. Equivalence between points 1 through 4 follows directly from Theorem 46, Proposition 44.

(1) \iff (5): By Proposition 44, sober and weakly sober are equivalent in a T_D convergence of finite depth, hence in all its subspaces. Let $A \subset X$ with inclusion map $i : A \rightarrow X$, and let \mathcal{U} be an ultrafilter on A with $\lim_{\xi_A} \mathcal{U} \in \mathcal{U}$. Hence $i(\lim_{\xi_A} \mathcal{U}) \in i[\mathcal{U}]$ and $i(\lim_{\xi_A} \mathcal{U}) \subset \lim_{\xi} i[\mathcal{U}]$ so that the ultrafilter $i[\mathcal{U}]$ on (X, ξ) is irreducible. By (1), there is $x \in X$ with $i[\mathcal{U}] = \{x\}^{\uparrow}$. As $A \in i[\mathcal{U}]$, $x \in A$ and thus $\lim_{\xi_A} \mathcal{U} = \lim_{\xi_A} \{x\}^{\uparrow}$ and A is weakly sober.

Conversely, if there is a free ultrafilter \mathcal{U} that is irreducible, if $\lim \mathcal{U}$ does not have a generic point, then the space is not weakly sober. If $\lim \mathcal{U}$ has a generic point, it is unique by Lemma 7, say, $\lim \mathcal{U} = \lim \{x_0\}^{\uparrow}$. Let $A = \lim \mathcal{U} \setminus \{x_0\}$ with the induced convergence. Then $A \in \mathcal{U}$ (because $\lim \mathcal{U} \in \mathcal{U}$ and $X \setminus \{x_0\} \in \mathcal{U}$ as \mathcal{U} is free) and $\lim_{\xi_A} \mathcal{U} = A \in \mathcal{U}$ but $\lim_{\xi_A} \mathcal{U}$ has not generic point. Hence, (A, ξ_A) is a non weakly sober subspace.

(1) \iff (6) : If $\mathcal{U} \in \mathbb{U}X$ with $\lim_{\theta} \mathcal{U} \in \mathcal{U}$ then $\lim_{\xi} \mathcal{U} \in \mathcal{U}$ because $\lim_{\theta} \mathcal{U} \subset \lim_{\xi} \mathcal{U}$, so that \mathcal{U} is ξ -irreducible, hence principal. Hence it has a unique generic point, because ξ is S_0 and T_0 , hence almost antisymmetric. Thus θ is sober.

Conversely, suppose there is a free ξ -irreducible ultrafilter \mathcal{U} . Let θ be defined by $x \in \lim_{\theta} \mathcal{F}$ if and only if $\mathcal{F} = \{x\}^{\uparrow}$ whenever $x \notin \lim_{\xi} \mathcal{U}$ and $x \in \lim_{\theta} \mathcal{F}$ if and only if $\mathcal{F} = \mathcal{U}$ or $\mathcal{F} = \{x\}^{\uparrow}$ whenever $x \in \lim_{\xi} \mathcal{U}$. Then $\theta \geq \xi$ and θ is not sober: \mathcal{U} is θ -irreducible because $\lim_{\theta} \mathcal{U} = \lim_{\xi} \mathcal{U} \in \mathcal{U}$ but does not have a generic point because singletons are closed and $\lim_{\theta} \mathcal{U}$ is infinite.

(1) \iff (7): Let $L = (\mathbb{P}X, \lim_{\xi})$. By (1) the map $h : \text{pt}_{\mathbf{Lat}} L \rightarrow (X, \xi)$ defined by $h(\{x\}^{\uparrow}) = x$ is onto and thus is an homeomorphism by Proposition 3.

Conversely, if $\text{pt} L$ is homeomorphic to (X, ξ) but there are free ξ -irreducible ultrafilters then, in view of Proposition 3, $\text{pt} L$ contains an homeomorphic copy of itself as a dense *proper* subset via an homeomorphism $j : \text{pt} L \rightarrow j(\text{pt} L) \subsetneq \text{pt} L$, which is not possible, for $j(\text{pt} L)$ is closed as an homeomorphic image of a closed subset of $\text{pt} L$, hence $\text{pt} L \setminus j(\text{pt} L)$ is open, making it impossible for $j(\text{pt} L)$ to be dense in $\text{pt} L$. \square

Note that in view of Remark 38, when (X, ξ) is topological, we can add (3.2) for every pair \mathcal{U}, \mathcal{W} of irreducible ultrafilters as an additional equivalent condition in Corollary 47.

REFERENCES

- [1] G. Brümmer. Initial quasi-uniformities. *Indag. Math.*, 31:403–409, 1969.
- [2] S. Dolecki. Convergence-theoretic characterizations of compactness. *Topology and its Applications*, 125:393–417, 2002.
- [3] S. Dolecki, G. H. Greco, and A. Lechicki. Compactoid and compact filters. *Pacific J. Math.*, **117**:69–98, 1985.
- [4] S. Dolecki and F. Mynard. *Convergence Foundations of Topology*. World Scientific, 2016.
- [5] J. Goubault-Larrecq and F. Mynard. Convergence without points. *Houston Journal of Mathematics*, 46(1):227–282, 2020.
- [6] Jean Goubault-Larrecq. *Non-Hausdorff topology and domain theory*, volume 22 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2013. [On the cover: Selected topics in point-set topology].
- [7] R.E. Hoffmann. Irreducible filters and sober spaces. *Manuscripta Math*, 22(4):365–380, 1977.

- [8] R.E. Hoffmann. On the sobrification remainder ${}^s x \setminus x$. *Pacific J. Math.*, 83(1):145–156, 1979.
- [9] F. Mynard. More pointfree convergence: on convergence frames. *in preparation*.
- [10] F. Mynard. Products of compact filters and applications to classical product theorems. *Topology and its Applications*, 154(4):953–968, 2007.
- [11] F. Mynard. Relations that preserve compact filters. *Applied Gen. Top.*, 8(2):171–185, 2007.
- [12] F. Mynard. Approach theory and pointfree convergence. *Top. Proc.*, 61(31-47), 2023.
- [13] J. Picado and A. Pultr. *Frames and locales: Topology without points*. Frontiers in Mathematics. Birkhäuser/Springer Basel AG, Basel, 2012.
- [14] M. Schroder. Compactness theorems. In *Categorical Topology*, pages 566–577. Springer-Verlag, 1974.

NJCU, DEPARTMENT OF MATHEMATICS, 2039 KENNEDY BLVD JERSEY CITY, NJ 07305,
USA

Email address: `fmynard@njcu.edu`