Conditional probability tensor decompositions for multivariate categorical response regression

Aaron J. Molstad^{†*} and Xin Zhang^{‡*}
[†]University of Florida and [‡]Florida State University

Abstract

In many modern regression applications, the response consists of multiple categorical random variables whose probability mass is a function of a common set of predictors. In this article, we propose a new method for modeling such a probability mass function in settings where the number of response variables, the number of categories per response, and the dimension of the predictor are large. We introduce a latent variable model which implies a low-rank tensor decomposition of the conditional probability tensor. This model is based on the connection between the conditional independence of responses, or lack thereof, and the rank of their conditional probability tensor. Conveniently, our model can be interpreted in terms of a mixture of regressions and can thus be fit using maximum likelihood. We derive an efficient and scalable penalized expectation maximization algorithm to fit this model and examine its statistical properties. We demonstrate the encouraging performance of our method through both simulation studies and an application to modeling the functional classes of genes. **Keywords:** Dimension reduction; Generalized linear model; Latent variable model; Mixture regression; Tensor decomposition.

1 Introduction

We consider the problem of modeling the conditional distribution of multiple categorical response variables as a function of a p-dimensional vector of predictors, i.e., a multivariate categorical response regression analysis. To make matters concrete, for an integer $M \geq 2$, let $\mathbf{Y} = (Y_1, \dots, Y_M)$ be the multivariate categorical response, where each component Y_m has $c_m \geq 2$ many categories with numerically coded support $Y_m \in [c_m] = \{1, \dots, c_m\}$ for all $m \in [M] = \{1, \dots, M\}$. The regression models and methods developed in this article allow

^{*}The authors contributed equally to this work and are listed in alphabetical order. Corresponding author: Aaron J. Molstad (amolstad@ufl.edu). Research for this paper was supported in part by grants DMS-2053697 (XZ), DMS-2113590 (XZ) and DMS-2113589 (AJM) from the U.S. National Science Foundation.

the predictor $\mathbf{X} \in \mathbb{R}^p$ to be either continuous or discrete (or mixed), and allow the predictors $\mathbf{x} \in \mathbb{R}^p$ to be either random or fixed. The regression problem is essentially the study of the conditional distribution $\mathbf{Y} \mid \mathbf{X}$ whose joint probability mass function consists of

$$P_{j_1...j_M}(\mathbf{x}) = \Pr(Y_1 = j_1, ..., Y_M = j_M \mid \mathbf{X} = \mathbf{x}) \ge 0, \quad j_m \in [c_m], \quad m \in [M].$$
 (1)

From (1), we can define the Mth order tensor $\mathbf{P}(\mathbf{x}) \in \mathbb{R}^{c_1 \times \cdots \times c_M}$ whose (j_1, \dots, j_M) th element is $P_{j_1 \dots j_M}(\mathbf{x})$. This conditional probability tensor $\mathbf{P}(\mathbf{x})$ fully characterizes the conditional distribution of $\mathbf{Y} \mid \mathbf{X} = \mathbf{x}$ and is thus the quantity of interest in our study.

1.1 Overview

To fit (1), there are numerous approaches one could consider. At one extreme, each response could be modeled separately using, say, multinomial logistic regression. This approach is scalable to a large number of responses and high-dimensional predictors (Zhu and Hastie, 2004; Simon et al., 2013; Vincent and Hansen, 2014), but entirely ignores the dependence between response variables. On the other extreme, one could define a univariate categorical response variable based on the set of all possible category combinations and methods designed for a univariate categorical response could be applied. For example, in applications with only binary responses, this would require treating M binary response variables as a univariate categorical variable with 2^{M} categories. In Section 6, we consider a genomic application with M=14 binary response variables, so this approach would lead to an unwieldy $2^M=16384$ categories. This approach allows for arbitrary dependence among responses, but in so doing, treats P(x) as a vector and thus fails to exploit its special tensor structure. Moreover, this approach would require an enormous amount of data for model fitting. If even a single category combination is not observed in the training data, this approach cannot be applied directly. Finally, when the number of categorical responses is even moderately large, model interpretation will be difficult because the number of parameters grows exponentially with the number of responses.

For the problem of modeling the conditional probability tensor function \mathbf{P} , we refer to these two approaches as separate modeling (of each response) and vectorized modeling (of the combined-category response), respectively. The objective of this article is to propose a novel alternative to these two approaches; a method which can model complex dependencies among response variables like vectorized modeling, yet provides fitted models which can be computed and interpreted with the ease and scalability of separate modeling. Our approach exploits the connection between the conditional independence of Y_1, \ldots, Y_M , or lack thereof,

and the rank of the conditional probability tensor $\mathbf{P}(\mathbf{x})$. Later, we prove that separate modeling implicitly assumes $\mathbf{P}(\mathbf{x})$ is rank one, whereas vectorized modeling assumes no explicit upper bound on the rank of $\mathbf{P}(\mathbf{x})$. We may thus characterize models for which $\mathbf{P}(\mathbf{x})$ is low rank as intermediate to these two extremes. Neatly, we later show that both the separate model and vectorized model can be characterized as "edge-cases" of a rank-constrained $\mathbf{P}(\mathbf{x})$ when the rank is fixed at one or the rank is allowed to be as large as $\prod_{m=1}^{M} c_m / \max_{k \in [M]} c_k$, respectively. Section 2.4 contains detailed comparisons of these approaches.

Motivated by this observation, in this article we propose a new method for multivariate categorical response which assumes the low-rankness of $\mathbf{P}(\mathbf{x})$. We show that pursuing a low-rank decomposition of the conditional probability tensor $\mathbf{P}(\mathbf{x})$ provides a natural, intuitive, and scalable way to model the complex dependencies among responses. However, because $\mathbf{P}(\mathbf{x})$ consists of the probabilities from (1), there are intrinsic constraints (nonnegativity and "sum-to-one") not often encountered in standard tensor decomposition problems. It is thus nontrivial to perform a low-rank tensor decomposition on $\mathbf{P}(\mathbf{x})$ even just for a single \mathbf{x} . To handle these difficulties, we propose a new latent variable interpretation of the conditional probability tensor function \mathbf{P} . This interpretation naturally allows \mathbf{P} to be characterized in terms of a mixture of regressions model, and implies a low-rank $\mathbf{P}(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}^p$. Moreover, this new latent variable formulation allows models to be fit using penalized maximum likelihood, which can accommodate large p, large M, and large c_m 's.

Before formally describing our proposed method and model, we first situate our work among existing methods for fitting (1).

1.2 Existing approaches

In the statistical literature on categorical data analysis, existing methods for multivariate categorical response regression are primarily focused on developing parametric links between predictors and multiple categorical responses that allow for model interpretation in terms of marginal probabilities and higher-order associations (Molenberghs and Lesaffre, 1999; Glonek, 1996; Ekholm et al., 2000; McCullagh and Nelder, 1989). For example, one set of link functions correspond to log-linear models, and another to multivariate logistic models (Glonek and McCullagh, 1995). Other approaches propose nonparametric regression functions and allow for some response-specific predictors (Gao et al., 2001). In general, these works adopt the vectorized modeling approach and are often not feasible when $M \geq 3$ and p is large, where the interactions among response variables are difficult to parameterize.

More recently, in the high-dimensional regime, Molstad and Rothman (2021) proposed

a novel penalty to enforce linear restrictions on the regression coefficient tensor under the vectorized model with multinomial link. Their penalty can lead to fitted models that can be interpreted in terms of which predictors affect only the marginal distributions of responses, the log odds ratios, or neither. However, their method does not easily generalize to more than two response variables. Theoretically, the estimation error bound for their method scales exponentially with M (more specifically, scales in $\prod_{m=1}^{M} c_m$) rather than linearly (such as $\sum_{m=1}^{M} c_m$) as does our method.

Of course, others have recognized the need for models and methods specifically designed for multivariate categorical response. One class of methods, based on the notion of binary relevance, comes from the literature on multi-label classification (Tsoumakas and Katakis, 2007) in machine learning. Many binary relevance methods fit separate univariate models, and thus fail to account for dependence in the multivariate response (Dembczyński et al., 2012; Montañes et al., 2014; Zhang et al., 2018). One binary relevance approach that accounts for dependence uses classifier chains (Read et al., 2011; Senge et al., 2013). This approach fits univariate categorical response regressions for Y_1, Y_2, \ldots, Y_M successively. For each univariate fit, the responses from previous fits are used as predictors in subsequent fits. For example, one would fit $Y_1 \mid \mathbf{X}$, then $Y_2 \mid \mathbf{X}, Y_1$, and so on. The fitted models are thus typically interpreted in terms of specific (univariate) conditional distributions for each response, rather than the joint distribution of interest. There is also an extensive literature on unsupervised modeling for multivariate categorical data (Fienberg, 2000; Dunson and Xing, 2009; Bhattacharya and Dunson, 2012), but this is not applicable to regression. Finally, we note that the problem of modeling P is not related to recent work on categorical data analysis focused on handling high-dimensional categorical predictors (e.g., Stokell et al., 2021).

Our model and method is related to – although fundamentally distinct from – existing research on tensor decompositions and regression. We give a very brief and selective survey in the following and refer the interested readers to Kolda and Bader (2009) and Bi et al. (2020). First, the idea of jointly modeling multiple responses using a (regularized) tensor decomposition is conceptually similar to that in tensor response regression (e.g., Li and Zhang, 2017), where the response is typically a continuous-valued tensor. However, our study is fundamentally different due to the discrete nature of the response. For example, continuous-valued tensor decompositions (e.g., Sun et al., 2017) are not applicable since they may not result in a valid probability tensor. Secondly, our regression problem is also distinct from recent studies focused on binary or categorical tensor decompositions (e.g., Wang and Li, 2020). Extensions of these unsupervised learning methods to our context is

nontrivial, especially in settings where the predictor is high-dimensional. Notably, Yang and Dunson (2016) also used the term "conditional probability tensor", though their focus was on estimating conditional probabilities of a categorical Y on multivariate categorical predictor X, which is fundamentally different from (1).

1.3 Contributions and organization

The main contributions of this paper are as follows. First, we propose a new and general method for modeling \mathbf{P} which allows practitioners to consider alternatives to the separate and vectorized modeling approaches. Crucially, unlike existing approaches which do not assume conditional independence, our method is scalable to large p, large M, and large c_m 's, without sacrificing flexibility or interpretability. The scalability allows for a broad range of potential applications, and the interpretability – in terms of both predictors selected and the estimated rank of the conditional probability tensor – allows practitioners to gain novel scientific insights.

Second, by connecting the low rank decomposition of P(x) to a latent variable model, we provide a new inferential and estimable object in the statistical analysis of a (conditional) probability tensor. Although the rank of a probability tensor is neatly connected to statistical independence, the current understanding of probability tensors of this type is mainly on the population level from an applied mathematics perspectives (e.g., Robeva and Seigal, 2019). Some statistical approaches were developed in recent years (Amiridi et al., 2019), but these are not applicable to regression.

Third, in order to accommodate different scenarios, we propose two new group lassotype penalties to achieve highly interpretable global and local variable selection. We devise an efficient algorithm which we prove to produce a sequence of iterates that monotonically increase the penalized observed data log-likelihood. Statistical properties are also established to illustrate how, in an idealized setting, our method scales with respect to the number of responses, the number of categories per response, and the number of predictors.

The rest of the article is organized as follows. Section 2 introduces the population model after establishing the connection between tensor low-rankness and conditional independence. Section 3 proposes a penalized expectation maximization (EM) algorithm for model fitting. In Section 4, we study the statistical properties of our estimator in an idealized setting. Sections 5, 6, and 7, respectively, contain extensive simulation experiments, a real data example, and a short discussion. The Supplementary Materials contain all additional technical details and numerical results.

2 Model

2.1 Tensor notation

For a positive integer M, an M-way tensor (also known as an Mth order tensor) is an array object $\mathbf{A} \in \mathbb{R}^{p_1 \times \cdots \times p_M}$ for positive integers p_1, \ldots, p_M . For example, a vector is a one-way tensor, and a matrix is a two-way tensor. The tensor rank decomposition, also known as the CANDECOMP/PARAFAC decomposition, is a generalization of the singular value decomposition for matrices (Hitchcock, 1927; Carroll and Chang, 1970). A rank-one tensor $\mathbf{A} \in \mathbb{R}^{p_1 \times \cdots \times p_M}$ can be written as the outer product of M vectors: $\mathbf{A} = \mathbf{a}^{(1)} \circ \cdots \circ \mathbf{a}^{(M)}$, which is defined element-wise as $\mathbf{A}_{i_1,\ldots,i_M} = \mathbf{a}^{(1)}_{i_1} \cdots \mathbf{a}^{(M)}_{i_M}$ for all $i_m \in [p_m]$ and $m \in [M]$. In general, a rank-R tensor $(R \geq 2)$ can be written as the sum of R rank-one tensors, each formed as the outer product of vectors $\mathbf{a}_r^{(m)}$, $r \in [R]$, $m \in [M]$: $\mathbf{A} = \sum_{r=1}^R \mathbf{a}_r^{(1)} \circ \cdots \circ \mathbf{a}_r^{(M)}$. A tensor is said to be rank-R if it can be decomposed into R rank-one tensors but not into r rank-one tensors for any r < R.

2.2 Characterizations of the conditional probability tensor

The goal of this paper is to provide a statistical modeling framework for the Mth order tensor $\mathbf{P}(\mathbf{x}) \in \mathbb{R}^{c_1 \times \cdots \times c_M}$, whose (j_1, \ldots, j_M) th element is the conditional probability function $P_{j_1...j_M}(\mathbf{x})$ defined in (1). To that end, we first provide some characterizations of $\mathbf{P}(\mathbf{x})$ and its rank-R decomposition which motivate our latent variable model proposed in Section 2.3.

The probability tensor defined in (1) must have at least rank-one because rank-zero, which corresponds to the tensor of zeros, would not yield a valid probability mass function. Our first theorem establishes the connection between the rank-one conditional probability tensor and the conditional independence of the responses given the predictor.

Theorem 1. If $P(\mathbf{x}) \in \mathbb{R}^{c_1 \times \cdots \times c_M}$ is rank-one, or equivalently, can be decomposed as

$$\mathbf{P}(\mathbf{x}) = \mathbf{p}_1(\mathbf{x}) \circ \cdots \circ \mathbf{p}_M(\mathbf{x}), \tag{2}$$

for some $\mathbf{p}_m(\mathbf{x}) \in \mathbb{R}^{c_m}$, then (2) must be true for $\mathbf{p}_m(\mathbf{x}) = \{\Pr(Y_m = 1 \mid \mathbf{X} = \mathbf{x}), \dots, \Pr(Y_m = c_m \mid \mathbf{X} = \mathbf{x})\}^{\top}$. Consequently, (2) is equivalent to the conditional independence of Y_1, \dots, Y_M given $\mathbf{X} = \mathbf{x}$.

The tensor rank-one decomposition in (2) is not unique. Theorem 1 shows that a rank-one probability tensor can be decomposed into marginal probability vectors without loss of generality. By constructing $\mathbf{p}_m(\mathbf{x})$ from $\Pr(Y_m = j_m \mid \mathbf{X} = \mathbf{x})$ in the above tensor

rank decomposition, we present an identifiable decomposition with naturally non-negative components.

For any nonzero tensor, there always exist a constant $R \geq 1$ such that a rank-R tensor decomposition holds. However, the relationship between the tensor rank and the tensor dimension is not as straightforward as in the matrix case (i.e., the rank of a matrix is no more than its row or column dimension). We next consider generalizing the rank-one structure to an arbitrary rank-R. Motivated by Theorem 1, the next theorem considers the decomposition of $\mathbf{P}(\mathbf{x})$ into a sum of R rank-one probability tensors (e.g., see Figure 1). Similar to (2), the unconstrained vectors $\mathbf{p}_{mr}(\mathbf{x}) \in \mathbb{R}^{c_m}$ in the tensor rank decomposition will be restricted to the set of c_m -dimensional vectors with nonnegative elements, denoted by $\mathbb{R}^{c_m}_+$, without loss of generality.

Theorem 2. Suppose P(x) is the sum of R rank-one probability tensors,

$$\mathbf{P}(\mathbf{x}) = \sum_{r=1}^{R} \delta_r \mathbf{P}_r(\mathbf{x}) = \sum_{r=1}^{R} \delta_r \mathbf{p}_{1r}(\mathbf{x}) \circ \cdots \circ \mathbf{p}_{Mr}(\mathbf{x}), \tag{3}$$

where $R \geq 1$, $\delta_r \in \mathbb{R}$, and $\mathbf{p}_{mr}(\mathbf{x}) \in \mathbb{R}^{c_m}$. There exists $\delta_r > 0$ and $\mathbf{p}_{mr}(\mathbf{x}) \in \mathbb{R}^{c_m}_+$ with $\sum_{r=1}^R \delta_r = 1$ and $\|\mathbf{p}_{mr}(\mathbf{x})\|_1 = 1$ for $(m,r) \in [M] \times [R]$ such that (3) holds. Moreover, given \mathbf{x} , there exists a rank R such that (3) holds and $R \leq \prod_{m=1}^M c_m / \max_{k \in [M]} c_k$.

The general rank-R decomposition in (3) is a key assumption of our modeling approach. Based on Theorem 2, we can henceforth assume (3) holds for some $\delta_r > 0$ (with $\sum_{r=1}^R \delta_r = 1$) and $\mathbf{p}_{mr}(\mathbf{x}) \in \mathbb{R}_+^{c_m}$ (with $\|\mathbf{p}_{mr}(\mathbf{x})\|_1 = 1$), without loss of generality. This provides identifiability of the decomposition and, more importantly, a conditional independence interpretation of the \mathbf{p}_{mr} we discuss momentarily. Theorem 2 extends Theorem 1 of Dunson and Xing (2009) from unconditional probability mass function to conditional. In particular, it establishes the upper bound on the rank R while Dunson and Xing (2009) only showed the existence of such rank. For M = 2, Theorem 2 implies that the singular value decomposition of $\mathbf{P}(\mathbf{x}) = \sum_{r=1}^R \delta_r \mathbf{p}_{1r}(\mathbf{x}) \mathbf{p}_{2r}^{\top}(\mathbf{x})$ holds for rank $R \leq \min(c_1, c_2)$. This well-known fact for the singular value decomposition of a matrix is thus extended to the probability matrix decomposition with nonnegative, unit length (in L_1 -norm) singular vectors $\mathbf{p}_{1r}(\mathbf{x})$'s, $\mathbf{p}_{2r}(\mathbf{x})$'s, and singular values δ_r 's that sum to one.

As mentioned, direct estimation of a rank-R probability tensor in (3) is not practical or even possible. First of all, the tensor is varying with values of $\mathbf{x} \in \mathbb{R}^p$ and has various positivity and sum-to-one constraints. Standard or even nonnegative tensor decomposition methods would therefore fail. Moreover, the tensor rank decomposition itself has several




Figure 1: Diagram of the decomposition of the probability tensor $\mathbf{P}(\mathbf{x})$ (white cube) into the sum of R=3 rank-one tensors. The gray rank-one tensor represents, for example, $\delta_1\mathbf{p}_{11}(\mathbf{x}) \circ \cdots \circ \mathbf{p}_{M1}(\mathbf{x})$ from the decomposition in (3).

limitations such as identifiability, existence of an optimal low-rank solution (De Silva and Lim, 2008), and NP-hard computation (Hillar and Lim, 2013). Additional orthogonality constraints that are often imposed (Sun et al., 2017) may conflict with the validity of our probability mass tensor. Next, we propose a new latent variable model for the conditional probability tensor under which (3) holds for all $\mathbf{x} \in \mathbb{R}^p$.

2.3 Latent variable model

To provide a statistical interpretation of (3), we connect it to the conditional independence result established for the rank-one case. In particular, we exploit that each rank-one tensor in (3) consists of vectors $\mathbf{p}_{mr}(\mathbf{x})$ which, appealing to Theorem 1, can be interpreted as marginal conditional probabilities for the *m*th response. We thus introduce a latent categorical variable $Z \in [R] = \{1, \ldots, R\}$ that is independent of \mathbf{X} such that Y_1, \ldots, Y_M are conditionally independent given \mathbf{X} and Z. Specifically, we take $\delta_r = \Pr(Z = r \mid \mathbf{X} = \mathbf{x}) = \Pr(Z = r)$ and $\mathbf{p}_{mr}(\mathbf{x}) = \Pr(Y_m = j_m \mid \mathbf{X} = \mathbf{x}, Z = r)$ such that (3) is satisfied in Theorem 2. It follows that

$$P_{j_{1}...j_{M}}(\mathbf{x}) = \sum_{r=1}^{R} \Pr(Z = r | \mathbf{X} = \mathbf{x}) \Pr(Y_{1} = j_{1}, ..., Y_{M} = j_{M} | \mathbf{X} = \mathbf{x}, Z = r)$$

$$= \sum_{r=1}^{R} \left\{ \Pr(Z = r) \prod_{m=1}^{M} \Pr(Y_{m} = j_{m} | \mathbf{X} = \mathbf{x}, Z = r) \right\}.$$
(4)

This connection between (3) and (4) naturally leads to our latent variable model, which assumes a population level tensor decomposition (3) for all values of \mathbf{x} by assuming a parametric link for the conditional probability function $\Pr(Y_m = j_m \mid \mathbf{X} = \mathbf{x}, Z = r)$ for $(m,r) \in [M] \times [R]$. With this latent categorical variable $Z \in [R]$, our proposed model is thus




Figure 2: Diagram of the latent variable model interpretation of (3). The highlighted (in gray) part indicates that the probability tensor of $\mathbf{Y} \mid (\mathbf{X}, Z = 1)$ is rank-one and that responses are conditionally independent. The regression parameters $\boldsymbol{\theta}_r$ can characterized as $\boldsymbol{\theta}_r = \{\boldsymbol{\beta}_{mr}\}_{m \in [M]}$, where $\boldsymbol{\beta}_{m1}$ is the regression coefficient matrix for $Y_m \mid \mathbf{X}$ represented by the bottom-most (terminal) gray nodes.

$$Pr(Z = r) = \delta_r, \quad f(\mathbf{Y} \mid \mathbf{X} = \mathbf{x}, Z = r, \boldsymbol{\theta}) = \prod_{m=1}^{M} f_r(Y_m \mid \mathbf{X} = \mathbf{x}, \boldsymbol{\beta}_{mr}), \tag{5}$$

where f and f_r denote generic probability mass functions and $\boldsymbol{\beta}_{mr}$ (to be specified later) is the model parameter for the regression of Y_m on \mathbf{X} given Z = r. The parameters in this model are denoted $\boldsymbol{\theta} = \{\delta_r, \boldsymbol{\theta}_r\}_{r \in [R]} = \{\delta_r, \boldsymbol{\beta}_{1r}, \dots, \boldsymbol{\beta}_{Mr}\}_{r \in [R]}$. Of course, we can write the joint probability mass function of interest as a mixture of regressions without Z as

$$f(\mathbf{Y} \mid \mathbf{X} = \mathbf{x}, \boldsymbol{\theta}) = \sum_{r=1}^{R} \delta_r f_r(\mathbf{Y} \mid \mathbf{X} = \mathbf{x}, \boldsymbol{\theta}_r) = \sum_{r=1}^{R} \left\{ \delta_r \prod_{m=1}^{M} f_r(Y_m \mid \mathbf{X} = \mathbf{x}, \boldsymbol{\beta}_{mr}) \right\}.$$
(6)

The model (5) an intuitive interpretation: there are R latent states indexed by Z, and conditional on the latent state and \mathbf{X} , the response variables are independent. Figure 2 provides a visualization of this interpretation. For each value of the latent variable $Z \in [R]$, we have a distinct sub-model with conditionally independent responses (i.e., separate models). Each branch from Z corresponds to a rank-one probability tensor (e.g., the gray rank-one tensor from Figure 1). The vectors $\mathbf{p}_{mr}(\mathbf{x})$ in (3) naturally consist of probabilities $\Pr(Y_m = 1 \mid Z = r, \mathbf{X} = \mathbf{x}), \ldots, \Pr(Y_m = c_m \mid Z = r, \mathbf{X} = \mathbf{x})$, the terminal nodes of the diagram in Figure 2.

Notably, maximum likelihood estimates of $\boldsymbol{\theta}$, and consequently, of \mathbf{P} , can be obtained even when all category combinations (as described in our discussion of vectorized modeling) are not observed in the training data. This is because each set of parameters $\boldsymbol{\theta}_r$ corresponds to marginal conditional probabilities for each response.

For the remainder, we assume a multinomial logistic regression model for each $Y_m \mid (\mathbf{X} = \mathbf{x}, Z = r)$. The regression coefficients $\boldsymbol{\beta}_{mr} \in \mathbb{R}^{p \times c_m}$ thus characterize the c_m possible outcomes of a multinomial random variable Y_m (based on a single trial), whose mass function

 $f_r(Y_m \mid \mathbf{X} = \mathbf{x}, \boldsymbol{\beta}_{mr})$ consists of the probabilities

$$\Pr(Y_m = j \mid \mathbf{X} = \mathbf{x}, \boldsymbol{\beta}_{mr}) = \frac{\exp(\boldsymbol{\beta}_{mrj}^{\top} \mathbf{x})}{\sum_{k=1}^{c_m} \exp(\boldsymbol{\beta}_{mrk}^{\top} \mathbf{x})}, \quad j \in [c_m],$$
 (7)

where $\boldsymbol{\beta}_{mrj} \in \mathbb{R}^p$ is the *j*th column of $\boldsymbol{\beta}_{mr} \in \mathbb{R}^{p \times c_m}$. Under (7), the $\boldsymbol{\beta}_{mr}$ are not identifiable. This can be resolved by imposing the condition $\sum_{j=1}^{c_m} \boldsymbol{\beta}_{mrj} = 0$, a "sum-to-zero" constraint. Under this constraint, the model formulation and parameterization lead to identifiable parameters and conditional distributions. Later, we explain that when using our proposed penalties, this constraint is enforced automatically.

2.4 Comparison with alternative approaches

In this section, we compare our model assumptions (6) and (7) to the two other approaches for fitting M categorical responses $\mathbf{Y} = (Y_1, \dots, Y_M)$ on the predictor $\mathbf{X} \in \mathbb{R}^p$. For the sake of comparison, we use multinomial logistic links for all approaches.

The first, and most naive, direct approach is separate modeling. This model assumes that

$$\Pr(Y_m = j \mid \mathbf{X} = \mathbf{x}) = \frac{\exp(\boldsymbol{\eta}_{mj}^\top \mathbf{x})}{\sum_{k=1}^{c_m} \exp(\boldsymbol{\eta}_{mk}^\top \mathbf{x})}, \quad j \in [c_m], \ m \in [M],$$
(8)

where $\eta_{mj} \in \mathbb{R}^p$ for $j \in [c_m]$ and $m \in [M]$. This model is equivalent to our model with R = 1. If (8) is true, then our model with R > 1 becomes over-parameterized but can still provide consistent estimates of η_{mj} 's (with some asymptotic efficiency loss). If (6) and (7) are true for R > 1, then separate fitting based on (8) will not only lose the interrelationship between responses, but will be incorrect for the marginal probabilities $\Pr(Y_m \mid \mathbf{X} = \mathbf{x}), m \in [M]$. This may be somewhat surprising, but can be seen from the latent variable representation. That is,

$$\Pr(Y_m = j \mid \mathbf{X} = \mathbf{x}) = \sum_{r=1}^R \delta_r \Pr(Y_m = j \mid \mathbf{X} = \mathbf{x}, Z = r) = \sum_{r=1}^R \delta_r \frac{\exp(\boldsymbol{\beta}_{mrj}^\top \mathbf{x})}{\sum_{k=1}^{c_m} \exp(\boldsymbol{\beta}_{mrk}^\top \mathbf{x})},$$

which can not be rewritten as proportional to $\exp(\boldsymbol{\eta}_{mj}^{\top}\mathbf{x})$. Intuitively, the latent variable Z introduces heterogeneity, and hence nonlinearity, in the conditional probability function $\Pr(Y_m = j \mid \mathbf{X} = \mathbf{x})$. As a result, separate model fitting is insufficient even when there is only one response variable. This is analogous to fitting a linear model to a mixture of regressions with heterogeneous sub-populations.

The second direct approach is vectorized modeling. As mentioned, this approach trans-

forms **Y** into a univariate categorical response Y_* with $c_* = \prod_{m=1}^{M} c_m$ categories. The corresponding multinomial logistic regression model assumes

$$\Pr(\mathbf{Y}_* = j \mid \mathbf{X} = \mathbf{x}) = \frac{\exp(\mathbf{\gamma}_j^{\top} \mathbf{x})}{\sum_{k=1}^{c_*} \exp(\mathbf{\gamma}_k^{\top} \mathbf{x})}, \quad j \in [c_*],$$
(9)

where $\gamma_j \in \mathbb{R}^p$ for $j \in [c_*]$. For example, Molstad and Rothman (2021) assume (9) and impose linear restrictions on the matrix of γ_j 's. Similar to the separate fitting approach, if (6) and (7) are true for R > 1, then this joint fitting based on (9) will also be incorrect even for marginal probabilities $\Pr(Y_m \mid \mathbf{X} = \mathbf{x}), m \in [M]$. On the other hand, if (9) is correct, then the rank R in our model may be as large as c_* . That is, we have c_* rank-one tensors that each consists of one element of the probability tensor P(x). The number of free parameters is $p(c_*-1) = p(\prod_m^M c_m - 1)$ for (9) and $(R-1) + pR \sum_m (c_m - 1)$ for our latent variable model (6) and (7). For example, consider the scenario where $c = c_1 = \cdots = c_M$. Then the number of free parameters becomes $p(c^M-1)$ for the vectorized model (9) and (R-1)+pR(c-1)Mfor the rank R version of our model. As the number of responses M increases, the complexity of (9) increases exponentially the order of $O(pc^M)$ while our model's complexity increases linearly in the order of O(pcMR). As the number of categories for each response c increases, our model's complexity still increases linearly in c but the complexity of (9) increases more rapidly as c^M as $M \geq 2$. To gain further intuition, when c = M = 4 and p = 100 (as in our simulation studies), the joint model has 25500 free parameters and our model has 2401 when R = 2 or 3602 when R = 3.

A practical advantage of our approach is that its estimation algorithm is much more scalable to large c, M, and p than estimators based on the vectorized model (9). When M is large, methods that fit (9) become infeasible partly due to the enormous number of parameters. We discuss our computational approach in the subsequent section.

Finally, our new latent variable approach is related to the existing latent variable regression model of Bandeen-Roche et al. (1997), who assume Y is independent of \mathbf{X} given the latent variable Z, but model Z given \mathbf{X} with a generalized linear model. Compared to the model of Bandeen-Roche et al. (1997), (5) is arguably more flexible due to the larger number of regression parameters.

3 Penalized maximum likelihood estimation

Let the observed data be $\{(\mathbf{Y}_i, \mathbf{x}_i)\}_{i=1}^n$ where $\mathbf{Y}_i = (Y_{1i}, \dots, Y_{Mi})$ for $i \in [n] = \{1, \dots, n\}$. Recall that $\boldsymbol{\theta}$ denotes all of the unknown parameters $\{(\delta_r, \boldsymbol{\beta}_{1r}, \dots, \boldsymbol{\beta}_{Mr})\}_{r \in [R]} \in \mathcal{D}^R$ where $\mathcal{D} = (0, 1) \times \mathbb{R}^{p \times c_1} \times \dots \times \mathbb{R}^{p \times c_M}$ with $\sum_{r=1}^R \delta_r = 1$. The conditional log-likelihood of $\mathbf{Y} \mid \mathbf{X}$ evaluated at $\boldsymbol{\theta}$ is

$$\sum_{i=1}^{n} \log \left[\sum_{r=1}^{R} \delta_r \left\{ \prod_{m=1}^{M} f_r(Y_{mi} \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\beta}_{mr}) \right\} \right]. \tag{10}$$

In this section, we first describe the standard EM algorithm for maximizing (10) over \mathcal{D}^R . The standard EM algorithm, which iterates between the expectation (E) step and the maximization (M) step, is only applicable in the classical low-dimensional setting. To address settings with high-dimensional predictors, we later discuss how to maximize a penalized version of (10). In particular, we devise a computational algorithm that replaces the penalized M-step with an approximation guaranteed to monotonically increase the objective function.

3.1 The EM algorithm and its parallel M-step

The standard EM algorithm will deal with the complete-data log-likelihood; that is, the log-likelihood of $(\mathbf{Y}, Z) \mid \mathbf{X}$, treating Z as if it were observable. Let $Z_{ir} = \mathbf{1}(Z_i = r)$. Recalling that Z is independent of \mathbf{X} with $\Pr(Z = r \mid \mathbf{X}) = \Pr(Z = r) = \delta_r$, the log-likelihood of $(\mathbf{Y}, Z) \mid \mathbf{X}$ evaluated at $\boldsymbol{\theta}$ is thus

$$\mathcal{L}(\boldsymbol{\theta}) = \sum_{i=1}^{n} \sum_{r=1}^{R} Z_{ir} \log \left\{ f_r(\mathbf{Y}_i \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\theta}_r) \right\} + \sum_{i=1}^{n} \sum_{r=1}^{R} Z_{ir} \log(\delta_r),$$
(11)

where $f_r(\mathbf{Y}_i \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\theta}_r) = \prod_{m=1}^M f_r(Y_{mi} \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\beta}_{mr})$ by definition. Each iteration of the EM algorithm, indexed by $t = 0, 1, 2, \ldots$, consists of two steps. In the E-step, we compute the \mathcal{Q} -function at tth iterate $\boldsymbol{\theta}^{(t)}$, $\mathcal{Q}(\boldsymbol{\theta} \mid \boldsymbol{\theta}^{(t)}) = \mathrm{E}\left[\mathcal{L}(\boldsymbol{\theta}) \mid \{(\mathbf{Y}_i, \mathbf{x}_i)\}_{i=1}^n, \boldsymbol{\theta}^{(t)}\right]$. To do so, we first compute the conditional estimate of $\pi_{ir} = \mathrm{E}(Z_{ir})$, the probability the $Z_i = r$, given the observed data and $\boldsymbol{\theta}^{(t)}$,

$$\pi_{ir}^{(t)} = \frac{\delta_r^{(t)} f_r(\mathbf{Y}_i \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\theta}_r^{(t)})}{\sum_{s=1}^R \delta_s^{(t)} f_s(\mathbf{Y}_i \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\theta}_r^{(t)})}.$$
(12)

Then we can express the Q-function as

$$\mathcal{Q}(\boldsymbol{\theta} \mid \boldsymbol{\theta}^{(t)}) = \sum_{i=1}^{n} \sum_{r=1}^{R} \left[\pi_{ir}^{(t)} \sum_{m=1}^{M} \log\{f_r(Y_{mi} \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\beta}_{mr})\} + \pi_{ir}^{(t)} \log(\delta_r) \right]. \tag{13}$$

In the M-step, we compute $\boldsymbol{\theta}^{(t+1)}$, which we define as the maximizer of (13) with respect to $\boldsymbol{\theta}$. One can verify that $\delta_r^{(t+1)} = n^{-1} \sum_{i=1}^n \pi_{ir}^{(t)}$, so the main challenge is maximizing the \mathcal{Q} -function with respect to the regression coefficients $\boldsymbol{\beta}_{mr} \in \mathbb{R}^{p \times c_m}$ for $(m, r) \in [M] \times [R]$.

From the first term in the Q-function, one can see that the maximization with respect to the β_{mr} is separable across each (m, r) combination. Therefore, for $(m, r) \in [M] \times [R]$ in an embarassingly parallel fashion, we need only compute

$$\boldsymbol{\beta}_{mr}^{(t+1)} = \underset{\boldsymbol{\beta}_{mr} \in \mathbb{R}^{p \times c_m}}{\operatorname{argmax}} \ell_{mr}(\boldsymbol{\beta}_{mr} \mid \boldsymbol{\theta}^{(t)}),$$

$$\ell_{mr}(\boldsymbol{\beta}_{mr} \mid \boldsymbol{\theta}^{(t)}) = \sum_{i=1}^{n} \pi_{ir}^{(t)} \log\{f_r(Y_{mi} \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\beta}_{mr})\}.$$
(14)

The solution to the above optimization problem is obtained by fitting a weighted multinomial logistic regression model of Y_m on \mathbf{X} . This could be done using a modified version of the standard computational approaches, e.g., a quasi-Newton method. This special structure naturally lends itself to settings with large M and large R.

The update (14) reinforces the generality of our latent variable model. We could replace the assumption of multinomial logistic link in (7) with a different assumption on $f_r(Y_{mi} | \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\beta}_{mr})$. The only necessary modification of the estimation procedure is in (14).

3.2 Penalties on the tensor parameter

To address the p > n case, we propose to maximize a penalized version of (10). Imposing penalties on each β_{mr} separately is possible, but may lead to fitted models which are difficult to interpret. Moreover, by imposing penalties across both mixture and response components, efficiency can be greatly improved. To achieve this, first organize all $\beta_{mr} \in \mathbb{R}^{p \times c_m}$ into a tensor parameter $\mathbf{B} \in \mathbb{R}^{p \times R \times C}$, where $C = \sum_{m=1}^{M} c_m$ and define the mth mode-2 slice of \mathbf{B} as $\mathbf{B}_{[:,m,:]} = (\beta_{m1}, \ldots, \beta_{mR}) \in \mathbb{R}^{p \times C}$ for $m \in [M]$. We propose to estimate the parameters of \mathbf{P} using

$$\underset{\boldsymbol{\theta} \in \mathcal{D}^R}{\operatorname{argmax}} \left\{ \mathcal{F}(\boldsymbol{\theta}) - \mathcal{P}_{\lambda}(\mathbf{B}) \right\}, \tag{15}$$




Figure 3: Visualizing the global and local variable selections and the corresponding sparsity patterns in the tensor regression parameter **B**. The two plots in middle respectively show the mode-1 slices and the mode-3 fibers to be selected. Our blockwise proximal gradient descent algorithm updates one row of $\mathbf{B}_{(1)}$ at a time for the penalty \mathcal{G}_{λ} , and updates one column of $\mathbf{B}_{(3)}$ at a time for the penalty \mathcal{H}_{λ} .

where \mathcal{F} is the observed data conditional log-likelihood in (10) and \mathcal{P}_{λ} is a sparsity-inducing penalty by the tuning parameter $\lambda > 0$. To compute (15), we need only modify the M-step of the EM algorithm from Section 3.1 to be replaced with the following joint optimization problem

$$\underset{\mathbf{B} \in \mathbb{R}^{p \times R \times C}}{\operatorname{argmax}} \mathcal{M}_{\lambda}(\mathbf{B} \mid \boldsymbol{\theta}^{(t)}), \quad \mathcal{M}_{\lambda}(\mathbf{B} \mid \boldsymbol{\theta}^{(t)}) = \sum_{m=1}^{M} \sum_{r=1}^{R} \ell_{mr}(\boldsymbol{\beta}_{mr} \mid \boldsymbol{\theta}^{(t)}) - \mathcal{P}_{\lambda}(\mathbf{B}), \quad (16)$$

where $\ell_{mr}(\cdot \mid \boldsymbol{\theta}^{(t)})$ is defined in (14) and \mathcal{P}_{λ} is a sparsity-inducing penalty with tuning parameter $\lambda > 0$. If $\lambda = 0$, (16) would reduce to (14). However, when $\lambda > 0$, the penalized objective function $\mathcal{M}_{\lambda}(\mathbf{B} \mid \boldsymbol{\theta}^{(t)})$ may not be separable across responses and mixture components depending on the choice of penalty \mathcal{P}_{λ} : we propose two such penalties which correspond to distinct types of variable selection.

First, we consider global variable selection. We say that the jth predictor is irrelevant if a change in the jth component of \mathbf{x} does not change $\mathbf{P}(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}^p$. Under our model assumption on \mathbf{P} , for the jth variable to be irrelevant it must be that $\mathbf{B}_{[j,:,:]}$

 $(d_{j11} \cdot \mathbf{1}_{c_1}^{\mathsf{T}}, d_{j12} \cdot \mathbf{1}_{c_1}^{\mathsf{T}}, \dots, d_{jMR} \cdot \mathbf{1}_{c_M}^{\mathsf{T}})^{\mathsf{T}} \in \mathbb{R}^{MC}$ for constants $d_{jmr} \in \mathbb{R}$, $(m, r) \in [M] \times [R]$. Recall that we have over-parameterized **B** in the sense that each $\boldsymbol{\beta}_{mr} \in \mathbb{R}^{p \times c_m}$ has only $(c_m - 1)$ identifiable columns (see the "sum-to-zero" constraint under equation (7)). This means that we may replace the d_{jmr} 's with 0 without loss of generality and ensure parameter identifiability. This equivalence between predictor irrelevance and sparsity in **B** is discussed more in the theoretical analysis (Section 4). For such global variable selection, we propose the following penalty term as \mathcal{P}_{λ} in (16),

$$\mathcal{G}_{\lambda}(\mathbf{B}) = \lambda \sum_{j=1}^{p} \|\mathbf{B}_{[j,:,:]}\|_{F} = \lambda \sum_{j=1}^{p} \sqrt{\sum_{r=1}^{R} \sum_{m=1}^{M} \|[\boldsymbol{\beta}_{mr}]_{j,:}\|_{2}^{2}},$$
(17)

where $\lambda > 0$ is a user-specified tuning parameter and $[\boldsymbol{\beta}_{mr}]_{j,:}$ is the jth row of $\boldsymbol{\beta}_{mr}$. The penalty \mathcal{G}_{λ} is nondifferentiable when for some $j \in [p] = \{1, \ldots, p\}$, $[\boldsymbol{\beta}_{mr}]_{j,k} = 0$ for all $k \in [c_m]$ and $(m,r) \in [M] \times [R]$. This penalty thus links the $\boldsymbol{\beta}_{mr}$ across both latent states and response variables. For large values of λ , this penalty will encourage estimates of \mathbf{P} such that many predictors are estimated to be irrelevant by encouraging zeros across the same rows of all RM coefficient matrices.

Although \mathcal{G}_{λ} can achieve a highly interpretable global form of variable selection, an alternative penalty, \mathcal{H}_{λ} , allows for variable selection specific to each latent state (i.e., local variable selection). Specifically, we also propose the penalty

$$\mathcal{H}_{\lambda}(\boldsymbol{\theta}) = \lambda \sum_{j=1}^{p} \sum_{r=1}^{R} \|\mathbf{B}_{[j,r,:]}\|_{2} = \lambda \sum_{j=1}^{p} \sum_{r=1}^{R} \sqrt{\sum_{m=1}^{M} \|[\boldsymbol{\beta}_{mr}]_{j,:}\|_{2}^{2}}.$$
 (18)

In contrast to \mathcal{G}_{λ} , the penalty \mathcal{H}_{λ} assumes that for a particular value of the latent variable Z, a possibly unique set of predictors are important. This penalty allows practitioners to characterize the categories of latent variable Z in terms of the predictors selected as relevant or not.

Figure 3 provides a visualization of the global and local variable selection in terms of the sparsity of **B**. The global penalty acts on entire mode-1 slices of the tensor parameter **B**. As shown in the plot, the mode-1 matricization $\mathbf{B}_{(1)} \in \mathbb{R}^{p \times RC}$ transforms each mode-1 slice into a row vector. The penalty $\mathcal{G}_{\lambda}(\mathbf{B})$ is thus the group lasso penalty on rows of $\mathbf{B}_{(1)}$. On the other hand, the local penalty is targeting on a more refined sparsity pattern that is shown as mode-3 fibers of **B**. Analogously, the mode-3 matricization $\mathbf{B}_{(3)} \in \mathbb{R}^{C \times pR}$ aligns the fibers across $r \in [R]$. The penalty $\mathcal{H}_{\lambda}(\mathbf{B})$ is thus the group lasso penalty on columns of $\mathbf{B}_{(3)}$.

For concreteness, we focus on computing (15) using the penalty \mathcal{G}_{λ} ; only trivial modifications of our algorithm are needed to accommodate \mathcal{H}_{λ} .

3.3 Penalized EM algorithm

In order to describe our approach, we express the objective function as $\mathcal{M}_{\lambda}(\mathbf{B} \mid \boldsymbol{\theta}^{(t)}) = \mathcal{M}_{\lambda}(\mathbf{B}_{1}, \mathbf{B}_{2}, \dots, \mathbf{B}_{p} \mid \boldsymbol{\theta}^{(t)})$ where $\mathbf{B}_{j} = \text{vec}(\mathbf{B}_{[j,:,:]}) \in \mathbb{R}^{RC}$ denotes the jth row of $\mathbf{B}_{(1)}$. Similarly, we let $\mathcal{M}_{\lambda,j}(\cdot \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)}) : \mathbb{R}^{RC} \to \mathbb{R}$ denote \mathcal{M}_{λ} as a function of its jth argument alone with $\mathbf{B}_{1}, \dots, \mathbf{B}_{j-1}$ fixed at $\mathbf{B}_{1}^{(t+1)}, \dots, \mathbf{B}_{j-1}^{(t+1)}$ and $\mathbf{B}_{j+1}, \dots, \mathbf{B}_{p}$ at $\mathbf{B}_{j+1}^{(t)}, \dots, \mathbf{B}_{p}^{(t)}$. By construction, $\mathcal{M}_{0,j}(\cdot \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)})$ is simply the unpenalized version of $\mathcal{M}_{\lambda,j}(\cdot \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)})$.

Our approach is to update each \mathbf{B}_{j} by maximizing a penalized quadratic minorizer (Lange, 2016) of $\mathcal{M}_{0,j}(\cdot \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)})$ constructed at $\mathbf{B}_{j}^{(t)}$ with all other \mathbf{B}_{k} for $k \neq j$ fixed at their current iterates. Our minorizing objective function for $\mathcal{M}_{\lambda,j}(\mathbf{b} \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)})$, where $\mathbf{b} \in \mathbb{R}^{RC}$, is defined as

$$\widetilde{\mathcal{M}}_{\lambda,j}(\mathbf{b} \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)}) = \mathcal{M}_{0,j}(\mathbf{B}_{j}^{(t)} \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)}) - (2\tau_{j})^{-1} \|\mathbf{b} - \mathbf{B}_{j}^{(t)}\|_{2}^{2} - \lambda \|\mathbf{b}\|_{2} + \operatorname{tr} \{ \nabla \mathcal{M}_{0,j}(\mathbf{B}_{j}^{(t)} \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)})^{\top} (\mathbf{b} - \mathbf{B}_{j}^{(t)}) \},$$
(19)

for which, with a sufficiently small step size $\tau_j > 0$, we have the following result.

Theorem 3. Let $\tilde{\mathbf{x}}_j = (\mathbf{x}_{1j}, \dots, \mathbf{x}_{nj})^{\top} \in \mathbb{R}^n$ where \mathbf{x}_{ij} is the jth component \mathbf{x}_i . For any fixed $0 < \tau_j \le \{RM \|\tilde{\mathbf{x}}_j\|_2 (\max_{k \in [M]} \sqrt{c_k})\}^{-1}$, we have $\widetilde{\mathcal{M}}_{\lambda,j}(\mathbf{b} \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)}) \ge \mathcal{M}_{\lambda,j}(\mathbf{B}_j^{(t)} \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)})$ for all $\mathbf{b} \in \mathbb{R}^{RC}$. Thus, if we define

$$\mathbf{B}_{j}^{(t+1)} = \operatorname*{argmax}_{\mathbf{b} \in \mathbb{R}^{RC}} \widetilde{\mathcal{M}}_{\lambda,j}(\mathbf{b} \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)}),$$

then we are guaranteed that $\mathcal{M}_{\lambda,j}(\mathbf{B}_{j}^{(t+1)} \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)}) \geq \mathcal{M}_{\lambda,j}(\mathbf{B}_{j}^{(t)} \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)})$ by the minorize-maximize principle (Lange, 2016). In addition, the maximizer of $\widetilde{\mathcal{M}}_{\lambda,j}(\cdot \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)})$ has the closed form

$$\mathbf{B}_{j}^{(t+1)} = \max\left(1 - \frac{\lambda \tau_{j}}{\|\mathbf{u}^{(t)}\|_{2}}, 0\right) \mathbf{u}^{(t)}, \tag{20}$$

where $\mathbf{u}^{(t)} = \mathbf{B}_j^{(t)} + \tau_j \nabla \mathcal{M}_{0,j}(\mathbf{B}_j^{(t)} \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)}).$

The theoretical range for τ_j is not used in our implementation. Instead, we select τ_j using an Armijo-type backtracking line search which allows us to consider larger step sizes τ_j while

maintaining the descent property described in Theorem 3. To obtain the complete (t+1)th iterate of \mathbf{B} , we compute (20) for $j \in [p]$ in a random order. This randomized approach is similar to the random permutation cyclical coordinate descent algorithm of Wright and Lee (2020). After all p rows of \mathbf{B} are updated once, we proceed to the next E-step, letting $\{\delta_r^{(t+1)}, \beta_{r1}^{(t+1)}, \dots, \beta_{rM}^{(t+1)}\}_{r \in [R]}$. It is important to note that by cycling through the rows of \mathbf{B} only once, $\mathbf{B}^{(t+1)}$ is not the argument maximizing $\mathcal{M}_{\lambda}(\cdot \mid \boldsymbol{\theta}^{(t)})$. If we repeatedly cycled through all p rows, the iterates would eventually converge to the global maximizer of $\mathcal{M}_{\lambda}(\cdot \mid \boldsymbol{\theta}^{(t)})$. However, we found this approximation scheme was more computationally efficient than solving the M-step exactly at each iteration, and we can easily verify that it ensures ascent of the objective function.

Lemma 1. As long as each τ_j is chosen according to Theorem 3 (or by backtracking line search), the objective function from (15) evaluated at $\boldsymbol{\theta}^{(t+1)}$ is guaranteed to be no less than the objective function from (15) evaluated at $\boldsymbol{\theta}^{(t)}$. That is, the sequence of iterates $\{\boldsymbol{\theta}^{(t)}\}_{t=1}^{\infty}$ generated by Algorithm 1 monotonically increase the value of the objective function from (15).

Lemma 1 relies on the fact that our algorithm is an instance of the expectation conditional-maximization algorithm (Meng and Rubin, 1993). We summarize the entire algorithm in Algorithm 1. To accommodate \mathcal{H}_{λ} rather than \mathcal{G}_{λ} , we would need only replace $\|\mathbf{b}\|_2$ in (20) with $\sum_{r=1}^{R} \|\mathbf{b}_r\|_2$ where $\mathbf{b} = (\mathbf{b}_1^{\top}, \dots, \mathbf{b}_R^{\top})^{\top}$ with each $\mathbf{b}_j \in \mathbb{R}^C$. We would then apply an updating equation like (20) based on the subvectors of $\mathbf{u}^{(t)}$ corresponding to each \mathbf{b}_r .

3.4 Implementation details

In mixture modeling in low-dimensional settings, it is common to consider multiple values at which to initialize EM algorithms. However, we find such schemes are not necessary in our application. To fit our model with the penalties described in the previous section, one must select a tuning parameter value λ . Thus, one generally solves (15) for a grid of candidate tuning parameter values. To avoid convergence to bad local maxima, we initialize the algorithm for solving (15) at $\delta_r = 1/R$ for $r \in [R]$ and at different random values for the β_{mr} for each candidate λ . This way, since many values of λ lead to very similar models if initialized identically, we effectively try multiple random starting values while computing the solution path. In both our simulation studies and real data analyses, we did not observe bad local maxima to be an issue for exactly this reason.

Algorithm 1 Algorithm for maximizing the penalized observed data log-likelihood (15)

- 1. Initialize $\boldsymbol{\theta}^{(0)} = \{\delta_r^{(0)}, \boldsymbol{\beta}_{1r}^{(0)}, \dots, \boldsymbol{\beta}_{Mr}^{(0)}\}_{r \in [R]}$ where $\delta_r^{(0)} > 0$ is the probability that Z = r (i.e., $\sum_{r=1}^R \delta_r^{(0)} = 1$) and $\boldsymbol{\beta}_{mr}^{(0)} \in \mathbb{R}^{p \times c_m}$ such that for $(r, m) \in [R] \times [M]$.
- 2. Repeat the following steps for t = 0, 1, ..., until convergence.
 - (a) E-step: For each $r \in [R]$ and $i \in [n]$, compute the conditional probability that $Z_i = r$,

$$\pi_{ir}^{(t)} = \frac{\delta_r^{(t)} f_r(\mathbf{Y}_i \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\theta}_r^{(t)})}{\sum_{s=1}^R \delta_s^{(t)} f_s(\mathbf{Y}_i \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\theta}_r^{(t)})},$$

where $f_r(\mathbf{Y}_i \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\theta}_r) = \prod_{m=1}^M f_r(Y_{mi} \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\beta}_{mr})$ with each $f_r(Y_{mi} \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\beta}_{mr})$ being the multinomial probability mass function described in (7).

- (b) M-step: Perform (i) and (ii).
 - (i) For each $j \in [p]$ in a random order, compute

$$\mathbf{B}_{j}^{(t+1)} = \max\left(1 - \frac{\lambda \tau_{j}}{\|\mathbf{u}^{(t)}\|_{2}}, 0\right) \mathbf{u}^{(t)}$$

$$(21)$$

where
$$\mathbf{u}^{(t)} = \mathbf{B}_j^{(t)} + \tau_j \nabla \mathcal{M}_{0,j}(\mathbf{B}_j^{(t)} \mid \mathbf{B}_{-j}^{(t+1,t)}, \boldsymbol{\theta}^{(t)}).$$

- (ii) For each $r \in [R]$, compute $\delta_r^{(t+1)} = n^{-1} \sum_{i=1}^n \pi_{ir}^{(t)}$.
- 3. Output $\widehat{\boldsymbol{\theta}} = \boldsymbol{\theta}^{(t+1)}$ at convergence.

Regarding the choice of R, we found that cross-validation may not be necessary. In both our simulation studies and real data example, we found that overspecifying R often led to no worse performance than did selecting R by cross-validation. This can be partly explained by the fact that when using our penalties, the penalized EM algorithm automatically forces some δ_r estimates to be close to zero (e.g., less than 10^{-8}) when λ is sufficiently large. Consequently, computing the solution path for our estimator explores both varying levels of sparsity in the regression coefficients and implicitly, various values of R. See Section 5.3 for further details.

Regarding convergence, we terminate Algorithm 1 when the change in objective function value (15) is less than 10^{-8} for consecutive iterations. Code to implement our method and reproduce the results from Section 5 is available for download at [URL omitted for anonymity].

4 Statistical analysis of penalized M-step

To better understand the performance of our method, we study the statistical error involved in the penalized M-step of Algorithm 1 from Section 3.3. In the finite-sample analysis of the maximizer of $\mathcal{M}_{\lambda}(\mathbf{B} \mid \boldsymbol{\theta}^{(t)})$, it is very challenging to establish uniform concentration inequalities about the stochastic objective function $\mathcal{M}_{\lambda}(\mathbf{B} \mid \boldsymbol{\theta}^{(t)})$ and its gradient $\nabla \mathcal{M}_{\lambda}(\mathbf{B} \mid \boldsymbol{\theta}^{(t)})$, which depend on the estimates $\boldsymbol{\theta}^{(t)}$. For example, a theoretical study of the EM algorithm may require sample splitting; given a total of n samples and T iterations, the sample-splitting EM algorithm would use T subsets of size n/T to break the dependence of $\mathbf{B}^{(t+1)} = \operatorname{argmax}_{\mathbf{B}} \mathcal{M}_{\lambda}(\mathbf{B} \mid \boldsymbol{\theta}^{(t)})$ on $\boldsymbol{\theta}^{(t)}$. See Balakrishnan et al. (2017) and Zhang et al. (2020) for examples. Because of this challenge, we leave the finite-sample statistical analysis of our penalized EM algorithm as future research. Instead, we study an idealized estimator $\widehat{\mathbf{B}}^{\dagger}$ from a modified version of $\mathcal{M}_{\lambda}(\mathbf{B} \mid \boldsymbol{\theta}^{(t)})$ which replaces the $\pi_{ir}^{(t)}$ with the "oracle" information Z_{ir} . That is, by analyzing the maximizer of the penalized conditional log-likelihood of $(\mathbf{Y}, Z) \mid \mathbf{X}$ in (11), we derive bounds that are meant to illustrate how p, c_1, \ldots, c_M , and the sparsity of the $\boldsymbol{\beta}_{mr}$ affect estimation of the regression parameter \mathbf{B} in an idealized scenario.

Throughout this section, let $|\mathcal{A}|$ denote the cardinality of a set \mathcal{A} . Similarly, let $\|\mathbf{A}\|_{1,2} = \sum_k \|\mathbf{A}_{k,:}\|_2$ be the norm which sums the Euclidean norms of the rows of its matrix-valued argument. To simplify matters, we treat $\mathbf{X} = (\mathbf{x}_1, \cdots \mathbf{x}_n)^{\top} \in \mathbb{R}^{n \times p}$ as nonrandom and standardized such that $\sum_{i=1}^n \mathbf{X}_{i,j}^2 = n$ for $j \in [p]$. We focus on the penalized estimator using the global variable selection penalty $\mathcal{G}_{\lambda}(\mathbf{B})$. To avoid cumbersome tensor notation and operators, we re-define $\mathbf{B} = \mathbf{B}_{(1)} \in \mathbb{R}^{p \times RC}$ as the matrix parameter in this section.

The estimator we study, $\widehat{\mathbf{B}}^{\dagger}$, is defined formally as

$$\underset{\mathbf{B} \in \mathbb{R}^{p \times RC}}{\operatorname{argmin}} \left\{ -\frac{1}{n} \sum_{i=1}^{n} \sum_{r=1}^{R} \left[Z_{ir} \sum_{m=1}^{M} \log \left\{ f_r(Y_{mi} \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\beta}_{mr}) \right\} \right] + \mathcal{G}_{\lambda}(\mathbf{B}) \right\}. \tag{22}$$

where $Z_{ir} = \mathbf{1}(Z_i = r)$ for $(i, r) \in [n] \times [R]$. The above estimator hence does not depend on $\boldsymbol{\theta}^{(t)}$. We will treat R as a fixed and allow M, c_1, \ldots, c_m, p , and n to tend to infinity. Our objective is to establish an error bound on $\widehat{\mathbf{B}}^{\dagger} - \mathbf{B}^{\dagger}$ where we define $\mathbf{B}^{\dagger} = \operatorname{argmin}_{\mathbf{B} \in \pi^*} \mathcal{G}_{\lambda}(\mathbf{B})$ with π^* denoting the set of all \mathbf{B} which lead to the true probabilities $\mathbf{P}(\mathbf{x})$ for all $\mathbf{x} \in \mathbb{R}^p$. In the Supplementary Material, we show that \mathbf{B}^{\dagger} is uniquely defined, does not depend on λ , and that for each irrelevant predictor, the corresponding row $\mathbf{B}_{j,:}^{\dagger} \in \mathbb{R}^{RC}$ is zero. Thus, define $\mathcal{S} = \{j : \mathbf{B}_{j,:}^{\dagger} \neq 0, j \in [p]\}$ and $\mathcal{S}^c := \{j : \mathbf{B}_{j,:}^{\dagger} = 0, j \in [p]\}$ as the set of relevant and

irrelevant predictors, respectively.

Before we describe the needed conditions and assumptions, a few comments are in order. First, the estimator in (22) is distinct from the estimator which estimates each β_{mr} or even $(\beta_{1r}, \ldots, \beta_{Mr})$ separately across $r \in [R]$. The penalty we use, \mathcal{G}_{λ} , necessarily ties the estimators together so that even in the case that the Z_i are known, all components are estimated jointly. Second, the Z_i 's are random so our statistical analysis must consider the joint distribution of the Z_i and the responses. Finally, we mention that the estimator in (22) is also relevant to the emerging literature on regression with heterogeneous sub-populations. In that context, Z_i is the indicator that a subject belongs to a particular sub-population (hence observable), and then our estimator from (22) is adjusting for Z in order to harmonize the data (Fortin et al., 2017) while our penalty \mathcal{G}_{λ} can identify homogeneous structures across sub-population (Tang and Song, 2016).

Our first assumption is a new version of the restricted eigenvalue condition which applies to data generated from a mixture of regressions model. This assumption will depend on $n_{\min} = \min_{r \in [R]} \sum_{i=1}^{n} Z_{ir}$, the minimum number of samples observed across the R latent states. Let $\Delta = (\tilde{\Delta}_{11}, \ldots, \tilde{\Delta}_{M1}, \tilde{\Delta}_{12}, \ldots, \tilde{\Delta}_{MR}) \in \mathbb{R}^{p \times RC}$ with $\tilde{\Delta}_{mr} \in \mathbb{R}^{p \times c_m}$ for $(r, m) \in [R] \times [M]$, and let $\{\sum_{i=1}^{n} \Psi_{mr}^{\dagger}(\mathbf{x}_i) \otimes \mathbf{x}_i \mathbf{x}_i^{\top}\}$ be the Hessian of $f_r(Y_{mi} \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\beta}_{mr}^{\dagger})$ with respect to the vectorization of its argument at $\boldsymbol{\beta}_{mr}^{\dagger}$. The exact form of the positive semidefinite matrix $\Psi_{mr}^{\dagger}(\mathbf{x}_i) \in \mathbb{R}^{c_m \times c_m}$ is given in the Supplementary Material. In addition, let $\mathcal{N}(n_{\min}) = \{\mathcal{N}_1, \ldots, \mathcal{N}_R\}$ denote the set of R-element partitions of [n] where $\min_{r \in [R]} |\mathcal{N}_r| \geq n_{\min}$.

A1. (Restricted eigenvalue condition) There exists a constant k > 0 such that

$$0 < k \le \kappa(\mathcal{S}, n_{\min}) = \inf_{(\Delta, \mathcal{A}) \in \mathbb{C}(\mathcal{S}) \times \mathcal{N}(n_{\min})} \sum_{m=1}^{M} \sum_{r=1}^{R} \frac{\tilde{\Delta}_{mr}^{\top} \left[\sum_{i \in \mathcal{A}_r} \{ \Psi_{mr}^{\dagger}(\mathbf{x}_i) \otimes \mathbf{x}_i \mathbf{x}_i^{\top} \} \right] \tilde{\Delta}_{mr}}{n \|\Delta\|_F^2},$$

where
$$\mathbb{C}(S) = \{ \Delta \in \mathbb{R}^{p \times R \sum_{m=1}^{M} c_m} : \Delta \neq 0, \|\Delta_{S^c,:}\|_{1,2} \leq 3 \|\Delta_{S,:}\|_{1,2} \}.$$

Explicitly, the quantity $\kappa(\mathcal{S}, n_{\min})$ is a function of n_{\min} and \mathcal{S} . The minimum sample size n_{\min} determines the space over which the infimum is taken with respect to \mathcal{A} as it defines $\mathcal{N}(n_{\min})$. If Δ were fixed, taking the infimum with respect to \mathcal{A} would correspond to finding the worst possible – in the sense of minimizing $\kappa(\mathcal{S}, n_{\min})$ – partition of the subjects into the R latent states with each state having at least n_{\min} subjects. The set \mathcal{S} , in contrast, defines the set $\mathbb{C}(\mathcal{S})$. Implicitly, $\kappa(\mathcal{S}, n_{\min})$ is a function of R, M, and the δ_r 's. In fact, to use this quantity in our proofs, we need the following additional assumption.

A2. $(\delta_r \text{ bounds})$ There exists a constant $v \in (0, 1/2)$ such that $v \leq \delta_r \leq 1 - v$ for

$$r \in [R]$$
.

Lastly, we also make an assumption about the data generating process.

A3. (Data generating process) The data $\{(Z_i, \mathbf{Y}_i)\}_{i=1}^n$ are independent with $\Pr(Z_i = r) = \delta_r$, and $f(\mathbf{Y}_i \mid \mathbf{X}_i = \mathbf{x}_i, Z_i = r, \boldsymbol{\theta}) = \prod_{m=1}^M f_r(Y_{mi} \mid \mathbf{X}_i = \mathbf{x}_i, \boldsymbol{\beta}_{mr}^{\dagger})$ where f_r is as defined in (7) for $r \in [R]$.

We are ready to state our main result, which will depend on a sample size condition, C1.

Theorem 4. Let $k_1 > 2$ and $k_2 > 2$ be fixed constants. Suppose $\mathbf{A1}$ - $\mathbf{A3}$ hold. If $\lambda = \{\sum_{m=1}^{M} c_m/n\}^{1/2} + \{4Mk_2 \log p/n\}^{1/2}$, then for n sufficiently large such that condition $\mathbf{C1}$ holds,

$$\|\hat{\mathbf{B}}^{\dagger} - \mathbf{B}^{\dagger}\|_{F} \leq \frac{3k_{1}}{\kappa(\mathcal{S}, n_{\min})} \left(\sqrt{\frac{|\mathcal{S}| \sum_{m=1}^{M} c_{m}}{4n}} + \sqrt{\frac{k_{2}|\mathcal{S}|M \log p}{n}} \right)$$

with probability at least $1 - p^{1-k_2/2} - \sum_{r=1}^{R} \exp[-2n\{\delta_r - (n_{\min} - 1)/n\}^2]$.

The condition on the sample size (C1) needed for the result of Theorem 4 is as follows.

C1. (Sample size) Let $\phi_n = \max_{i \in [n]} \|\mathbf{x}_i\|_2 k_1 \lambda \sqrt{54|\mathcal{S}|} / \{2\kappa(\mathcal{S}, n_{\min})\}$ where λ is as prescribed in Theorem 4. The sample size n is sufficiently large with respect to $p, c_1, \ldots, c_M, M, k_1, k_2$, and $|\mathcal{S}|$ such that $e^{-\phi_n} + \phi_n - 1 - \phi_n^2 / k_1 > 0$.

Our error bound illustrates how our idealized M-step scales with respect to the number of response variables and number of categories per response. It is especially instructive to compare the error bound in Theorem 4 to that from Molstad and Rothman (2021). Recall that Molstad and Rothman (2021) propose an estimator of the regression coefficient tensor under the vectorized model. Unsurprisingly, their error bound scales in $\prod_{m=1}^{M} c_m$ rather than $\sum_{m=1}^{M} c_m$. This can be explained by the fact that the regression coefficient tensor of interest under the vectorized model has $p \prod_{m=1}^{M} c_m$ elements whereas under our model assumptions, there are only $Rp \sum_{m=1}^{M} c_m$ unknown regression coefficients. These results may suggest that when M is reasonably large, the method of Molstad and Rothman (2021) will perform poorly whereas our method may perform well as long as (5) with small R serves as a reasonable approximation to \mathbf{P} .

5 Simulation studies

5.1 Data generating models and competing methods

In this section, we compare our method for estimating conditional probability tensors to numerous alternative approaches. We consider data generating models with R, the training sample size, the mixture probabilities, and the magnitude of entries in the $\boldsymbol{\beta}_{mr}$'s varying. For a given R, we generate n independent realizations of the predictor $\mathbf{X} \sim N_{100}(0, \boldsymbol{\Sigma})$ where $\boldsymbol{\Sigma}_{j,k} = 0.5^{|j-k|}$ for $(j,k) \in [p] \times [p]$ and generate Y_1, \ldots, Y_4 from

$$\Pr(Y_1 = j_1, \dots, Y_4 = j_4 \mid \mathbf{X} = \mathbf{x}) = \sum_{r=1}^R \delta_r \left\{ \prod_{m=1}^4 \Pr(Y_m = j_m \mid \mathbf{X} = \mathbf{x}, Z = r) \right\}$$

where $\Pr(Y_m = j_m \mid \mathbf{X} = \mathbf{x}, Z = r) = \exp(\mathbf{x}^{\top} \boldsymbol{\beta}_{mrj_m})/\{\sum_{k=1}^4 \exp(\mathbf{x}^{\top} \boldsymbol{\beta}_{mrk})\}$ for $m \in [4], j_m \in [4]$, and $r \in [R]$. We set M = 4 and $c_1 = \cdots = c_4 = 4$ throughout. When R = 2, we set $\delta_2 = 1 - \delta_1$, whereas when R = 3, we set $\delta_2 = 2/3 - \delta_1$ and $\delta_3 = 1/3$. In both cases, we consider various values of δ_1 . In each setting we consider, we randomly select five predictors to affect the conditional probability tensor. Each of the five corresponding rows of the $\boldsymbol{\beta}_{mr}$'s has entries which are drawn independently from $N(0, \sigma_{\beta}^2)$. To select tuning parameters, we generate a validation set of size 200 from the same data generating model. To quantify performance of the various estimators, we also generate a testing set of size 1000.

We consider multiple versions of our method with candidate $R \in [4]$ (denoted Mix-1 through Mix-4) and \mathcal{P}_{λ} taken to be \mathcal{G}_{λ} . We compare these variations of our method to two methods which implicitly assume independence: fitting separate multinomial logistic regression models to each response with (a) group-lasso penalty on the rows of each of the regression coefficient matrices (Sep-Group) and (b) the L_1 -norm penalty applied the regression coefficient matrices (Sep-L1). Sep-Group is equivalent to our method with R = 1 and $\mathcal{G}_{\lambda}(\mathbf{B})$ replaced with $\sum_{m=1}^{M} \lambda_m \sum_{j=1}^{p} ||[\mathcal{G}_{m1}]_{j,:}||_2$. While it may seem natural to want to compare to the vectorized modeling approach, which treats all possible category combinations as a 4⁴-dimensional multinomial, this is not possible since when even with n = 450 (the largest training sample size we consider), the probability that all category combinations are observed is nearly zero in every scenario.

For all considered methods, tuning parameters are chosen to minimize the negative log-likelihood evaluated on the validation set. For the methods which fit separate models, tuning parameters are chosen for each response separately. We evaluate the perfor-

mance by calculating the square-root average Kullback-Leibler divergence on the testing set $\{n_{\text{test}}^{-1} \sum_{i=1}^{n_{\text{test}}} \sum_{j_1=1}^{c_1} \cdots \sum_{j_M=1}^{c_M} \hat{P}_{j_1...j_M}(\mathbf{x}_i) \log(\hat{P}_{j_1...j_M}(\mathbf{x}_i)/P_{j_1...j_M}(\mathbf{x}_i))\}^{1/2}$, where $\mathbf{P}(\mathbf{x}_i)$ is the true probability mass function of $\mathbf{Y} \mid \mathbf{X} = \mathbf{x}_i$ and $\hat{\mathbf{P}}(\mathbf{x}_i)$ is the estimate based on a particular fitted model.

5.2 Results

In Figure 4, we display the square-root average Kullback-Leibler divergences on the testing set for each of the six methods with R=2. Focusing first on the top row, where coefficients tend to have smaller magnitude, we see that when n=75, there is only minor differences between methods. When n=150 or n=300, however, we notice that Mix-2, Mix-3, and Mix-4 all tend to outperform the other competitors. This is not surprising given that each of these methods contains the model with the true number of components as a special case. Another interesting aspect of these results is that as δ_1 approaches 0.5, the differences in performance become more apparent. This too is not surprising because when $\delta_1=0.1$, Mix-1 could serve as a reasonable approximation to **P**. Results are effectively the same when $\sigma_{\beta}=2$, except that differences between the methods are more pronounced even in the case that n=75 or n=150.

We display analogous results for the data generating model with R=3 in Figure 5. Here, we see results similar to those when R=2. Specifically, when n=75 and coefficient magnitudes are (relatively) small, there is little distinction between the methods. As n=150, we see a more clear seperation between the versions of our method with different numbers of mixture components. Recall that when R=3, $(\delta_1,\delta_2,\delta_3)=(\delta_1,2/3-\delta_1,1/3)$ so that when $\delta_1=1/3$ (rounded to 0.33 in the figure), all three components are equally weighted, whereas when $\delta_1=0.58$, one could expect that **P** can be well approximated by R=2 components. Indeed, in both top and bottom rows of Figure 5, we see the difference between Mix-2 and Mix-3 is more pronounced when $\delta_1=0.33$ relative to when $\delta_1\geq 0.5$. Just as in the case with R=2, the magnitude of the regression coefficients β_{mr} appears to have an effect as well. Specifically, when $\sigma_{\beta}=2$, the difference between estimators is much greater than under similar settings with $\sigma_{\beta}=1$.

In Section S2 of the Supplementary Material, we present results for the same set of competitors under identical data generating models, but using Hellinger distance as a performance metric. In brief, relative performances are very similar to those based on KL divergence.




Figure 4: The square-root average Kullback-Leibler divergence evaluated on the testing set for the six considered methods. Results displayed are from 100 independent replications with (top row) $(\sigma_{\beta}, R) = (1, 2)$ and (bottom row) $(\sigma_{\beta}, R) = (2, 2)$.

5.3 Effect of overspecifying R

In the simulation results displayed in Figures 4 and 5, we see that overspecifying the number of mixture components seems to have little effect on estimation. In Figure 5, we see that Mix-4 tended to perform slightly better than Mix-3, which has the correctly specified number of components. To explore whether more extreme overspecification has a similar effect, we repeated the simulations displayed in the bottom panel of Figure 5 and added three additional versions of our method with $R \in \{5, 10, 15\}$, which we call Mix-5, Mix-10, and Mix-15, respectively. We display results in Figure S3 of the Supplementary Material. Based on the results, it seems that extreme overspecification of R may even improve estimation accuracy when n is small: Mix-15 seemed to perform as well or better than all other methods in each scenario. However, it is important to note that for the particular tuning parameters chosen for, say, Mix-15, there were often only two nonzero estimates of the δ_r 's. This is one particularly appealing feature of the solution path for fitting (15): for large values of λ , even when R is large, the solution will have one $\delta_r = 1$ and $\delta_{r'} = 0$ for $r' \neq r$. As λ




Figure 5: The square-root average Kullback-Leibler divergence evaluated on the testing set for the six considered methods. Results displayed are from 100 independent replications with (top row) $(\sigma_{\beta}, R) = (1, 3)$ and (bottom row) $(\sigma_{\beta}, R) = (2, 3)$.

decreases, eventually a second δ_r becomes nonzero so that the fitted model effectively has R=2. This continues with each additional δ_r becoming nonzero as $\lambda \to 0$. We illustrate this phenomenon in Figure 6 of Section 6.

6 Modeling functional classes of genes

In this section, we apply our method to the problem of modeling a gene's functional classes based on both the gene's expression and phylogenetic profile. The dataset we analyzed, which was collected on yeast, was originally studied in Elisseeff and Weston (2001) and can be downloaded from https://www.uco.es/kdis/mllresources/. The predictors consist of p = 103 components, which are the collection of both the gene's expression and phylogenetic profile. In these data, there are M = 14 functional classes (including metabolism, energy, protein synthesis, etc.). Each of the n = 2417 genes can be characterized as belonging to multiple functional classes. For example, a gene may affect both metabolism and protein

	R = 1	R=2	R = 3	R = 4	R = 5
Joint error rate		82.23			
	,	(0.084)	,	,	` /
Deviance		16883			
	(13.64)	(28.46)	(20.40)	(31.88)	(36.53)

Table 1: Results based on five hundred replications: (top row) Joint error rates, i.e., proportion of testing subjects for which all fourteen classes were properly predicted; (bottom row) average deviance. Standard errors are provided in parentheses.




Figure 6: Results based on five hundred replications: (left) test set deviance for five versions of the mixture model with $R \in \{1, 2, 3, 4, 5\}$, (middle) ordered estimates of the δ_r for the version of our method with R = 5, and (right) number of predictors selected for each component mixture (in descending order according to the $\hat{\delta}_r$).

synthesis. Thus, it is natural to treat each functional class assignment as a binary response so that we have $c_1 = \cdots = c_{14} = 2$.

Because of the relatively large number of response variables, we used the version of our method with the penalty \mathcal{H}_{λ} described in (18). This penalty allows different components of the predictor to be relevant for different values of the latent variable Z. In the case that R = 1, this is equivalent to the penalty \mathcal{G}_{λ} , but differs for R > 1.

To compare our method's performance across different choices of R, for 500 independent replications we split the data into training, validation, and testing sets of size 1500, 500, and 417, respectively. For $R \in \{1, ..., 5\}$ separately, we fit the model to the training set and selected the tuning parameter λ by minimizing the negative log-likelihood evaluated on the validation set. Then, we compute the deviance and misclassification accuracy on the testing set for the selected model. In Table 1, we display the average joint error rate and the average testing set deviance for our method with $R \in [5]$. Evidently, R = 5 performed best in terms of both classification accuracy and deviance. We also observe that R = 4 and R = 5 are not

significantly different in either performance metric. This type of finding coheres with the simulation study results which suggest that performance is not hurt when R is over-specified.

We display the test set deviance for $R \in \{1, ..., 5\}$ in the leftmost panel of Figure 6. Here, we see there appears to be little difference between R = 4 and R = 5 in terms of testing set deviance. Notably, the version of our method assuming independent responses (R = 1) performs much worse than the versions with R > 1; adding even the second mixture component (R = 2) decreased testing set deviance by nearly 15%. In the two rightmost plots in Figure 6, we display the estimated δ_r and the number of predictors selected in the corresponding β_{mr} 's. We see that in general, even when R = 5, the selected model effectively has R = 4. We also notice that the mixture component with the highest probability often had 75 or more predictors included in the model, whereas those mixture components with smaller probabilities tended to have fewer.

7 Discussion

In this article, we propose a general modeling strategy for the regression of a multivariate categorical response on a high-dimensional predictor based on the population-level tensor rank decomposition of the conditional probability tensor function. Our approach exploits the connection between the tensor rank decomposition and the independence of the response variables conditioning on the predictor and a latent discrete variable Z. Numerically, our method is shown to perform well with a large number of response variables and large number of categories per response – a setting where many existing competitors fail or cannot be applied. Our results also suggest that our method is insensitive to over-specification of the number of mixtures R.

Based on our theoretical analysis of an idealized and simplified estimator, we conjecture that the convergence rate of the penalized EM algorithm, under suitable assumptions (see, for example, Balakrishnan et al., 2017; Cai et al., 2019), is $\sqrt{|\mathcal{S}|M\log p/n}$ when treating R and c_m 's as constants. We leave this as a future theoretical study. Promising future research directions also include (i) initialization of the algorithm and (ii) alternative convex penalties on the tensor \mathbf{B} . Regarding (i), methods such as Sedghi et al. (2016) are applicable to our setting and could potentially improve over the random initialization that we are currently using. For (ii), recent advances on tensor nuclear norm penalties (e.g., Raskutti et al., 2019) neatly fit in our optimization framework and can take advantage of the tensor construction of \mathbf{B} .

References

- Amiridi, M., Kargas, N., and Sidiropoulos, N. D. (2019). Statistical learning using hierarchical modeling of probability tensors. In 2019 IEEE Data Science Workshop (DSW), pages 290–294. IEEE.
- Balakrishnan, S., Wainwright, M. J., and Yu, B. (2017). Statistical guarantees for the EMalgorithm: From population to sample-based analysis. *The Annals of Statistics*, 45(1):77–120.
- Bandeen-Roche, K., Miglioretti, D. L., Zeger, S. L., and Rathouz, P. J. (1997). Latent variable regression for multiple discrete outcomes. *Journal of the American Statistical Association*, 92(440):1375–1386.
- Bhattacharya, A. and Dunson, D. B. (2012). Simplex factor models for multivariate unordered categorical data. *Journal of the American Statistical Association*, 107(497):362–377.
- Bi, X., Tang, X., Yuan, Y., Zhang, Y., and Qu, A. (2020). Tensors in statistics. *Annual Review of Statistics and Its Application*, 8.
- Cai, T. T., Ma, J., and Zhang, L. (2019). Chime: Clustering of high-dimensional gaussian mixtures with em algorithm and its optimality. *The Annals of Statistics*, 47(3):1234–1267.
- Carroll, J. D. and Chang, J.-J. (1970). Analysis of individual differences in multidimensional scaling via an n-way generalization of "eckart-young" decomposition. *Psychometrika*, 35(3):283–319.
- De Silva, V. and Lim, L.-H. (2008). Tensor rank and the ill-posedness of the best low-rank approximation problem. SIAM Journal on Matrix Analysis and Applications, 30(3):1084–1127.
- Dembczyński, K., Waegeman, W., Cheng, W., and Hüllermeier, E. (2012). On label dependence and loss minimization in multi-label classification. *Machine Learning*, 88(1-2):5–45.
- Dunson, D. B. and Xing, C. (2009). Nonparametric bayes modeling of multivariate categorical data. *Journal of the American Statistical Association*, 104(487):1042–1051.
- Ekholm, A., McDonald, J. W., and Smith, P. W. (2000). Association models for a multi-variate binary response. *Biometrics*, 56(3):712–718.

- Elisseeff, A. and Weston, J. (2001). A kernel method for multi-labelled classification. Advances in neural information processing systems, 14:681–687.
- Fienberg, S. E. (2000). Contingency tables and log-linear models: Basic results and new developments. *Journal of the American Statistical Association*, 95(450):643–647.
- Fortin, J.-P., Parker, D., Tunç, B., Watanabe, T., Elliott, M. A., Ruparel, K., Roalf, D. R., Satterthwaite, T. D., Gur, R. C., and Gur, R. E. (2017). Harmonization of multi-site diffusion tensor imaging data. *Neuroimage*, 161:149–170.
- Gao, F., Wahba, G., Klein, R., and Klein, B. (2001). Smoothing spline anova for multivariate bernoulli observations with application to ophthalmology data. *Journal of the American Statistical Association*, 96(453):127–160.
- Glonek, G. F. (1996). A class of regression models for multivariate categorical responses. Biometrika, 83(1):15–28.
- Glonek, G. F. and McCullagh, P. (1995). Multivariate logistic models. *Journal of the Royal Statistical Society: Series B (Methodological)*, 57(3):533–546.
- Hillar, C. J. and Lim, L.-H. (2013). Most tensor problems are NP-hard. *Journal of the ACM*, 60(6):1–39.
- Hitchcock, F. L. (1927). The expression of a tensor or a polyadic as a sum of products. Journal of Mathematics and Physics, 6(1-4):164–189.
- Kolda, T. G. and Bader, B. W. (2009). Tensor decompositions and applications. SIAM Review, 51(3):455–500.
- Lange, K. (2016). MM optimization algorithms. SIAM.
- Li, L. and Zhang, X. (2017). Parsimonious tensor response regression. *Journal of the American Statistical Association*, 112(519):1131–1146.
- McCullagh, P. and Nelder, J. A. (1989). Generalized Linear Models, volume 37. CRC Press.
- Meng, X.-L. and Rubin, D. B. (1993). Maximum likelihood estimation via the ecm algorithm: A general framework. *Biometrika*, 80(2):267–278.
- Molenberghs, G. and Lesaffre, E. (1999). Marginal modelling of multivariate categorical data. *Statistics in Medicine*, 18(17-18):2237–2255.

- Molstad, A. J. and Rothman, A. J. (2021). A likelihood-based approach for multivariate categorical response regression in high dimensions. *Journal of the American Statistical Association*.
- Montañes, E., Senge, R., Barranquero, J., Quevedo, J. R., del Coz, J. J., and Hüllermeier, E. (2014). Dependent binary relevance models for multi-label classification. *Pattern Recognition*, 47(3):1494–1508.
- Raskutti, G., Yuan, M., and Chen, H. (2019). Convex regularization for high-dimensional multiresponse tensor regression. *The Annals of Statistics*, 47(3):1554–1584.
- Read, J., Pfahringer, B., Holmes, G., and Frank, E. (2011). Classifier chains for multi-label classification. *Machine Learning*, 85(3):333.
- Robeva, E. and Seigal, A. (2019). Duality of graphical models and tensor networks. *Information and Inference: A Journal of the IMA*, 8(2):273–288.
- Sedghi, H., Janzamin, M., and Anandkumar, A. (2016). Provable tensor methods for learning mixtures of generalized linear models. In *Artificial Intelligence and Statistics*, pages 1223–1231.
- Senge, R., Coz Velasco, J. J. d., and Hüllermeier, E. (2013). Rectifying classifier chains for multi-label classification. *Space*, 2 (8).
- Simon, N., Friedman, J., and Hastie, T. (2013). A blockwise descent algorithm for group-penalized multiresponse and multinomial regression. arXiv preprint arXiv:1311.6529.
- Stokell, B. G., Shah, R. D., and Tibshirani, R. J. (2021). Modelling high-dimensional categorical data using nonconvex fusion penalties. *Journal of the Royal Statistical Society:* Series B (Statistical Methodology), 83(3):579–611.
- Sun, W. W., Lu, J., Liu, H., and Cheng, G. (2017). Provable sparse tensor decomposition. Journal of the Royal Statistical Society: Series B (Statistical Methodology), 79(3):899–916.
- Tang, L. and Song, P. X. (2016). Fused lasso approach in regression coefficients clustering: learning parameter heterogeneity in data integration. *The Journal of Machine Learning Research*, 17(1):3915–3937.
- Tsoumakas, G. and Katakis, I. (2007). Multi-label classification: An overview. *International Journal of Data Warehousing and Mining (IJDWM)*, 3(3):1–13.

- Vincent, M. and Hansen, N. R. (2014). Sparse group lasso and high dimensional multinomial classification. *Computational Statistics and Data Analysis*, 71:771–786.
- Wang, M. and Li, L. (2020). Learning from binary multiway data: Probabilistic tensor decomposition and its statistical optimality. *The Journal of Machine Learning Research*, 21:154–1.
- Wright, S. and Lee, C.-p. (2020). Analyzing random permutations for cyclic coordinate descent. *Mathematics of Computation*, 89(325):2217–2248.
- Yang, Y. and Dunson, D. B. (2016). Bayesian conditional tensor factorizations for high-dimensional classification. *Journal of the American Statistical Association*, 111(514):656–669.
- Zhang, L., Ma, R., Cai, T. T., and Li, H. (2020). Estimation, confidence intervals, and large-scale hypotheses testing for high-dimensional mixed linear regression. arXiv preprint arXiv:2011.03598.
- Zhang, M.-L., Li, Y.-K., Liu, X.-Y., and Geng, X. (2018). Binary relevance for multi-label learning: an overview. Frontiers of Computer Science, 12(2):191–202.
- Zhu, J. and Hastie, T. (2004). Classification of gene microarrays by penalized logistic regression. *Biostatistics*, 5(3):427–443.