

POINTED HOPF ALGEBRAS OVER NON-ABELIAN GROUPS WITH NON-SIMPLE STANDARD BRAIDINGS

IVÁN ANGIOÑO, SIMON LENTNER, GUILLERMO SANMARCO

ABSTRACT. We construct finite-dimensional Hopf algebras whose coradical is the group algebra of a central extension of an abelian group. They fall into families associated to a semisimple Lie algebra together with a Dynkin diagram automorphism. We show conversely that every finite-dimensional pointed Hopf algebra over a non-abelian group with non-simple infinitesimal braiding of rank at least 4 is of this form.

We follow the steps of the Lifting Method by Andruskiewitsch–Schneider. Our starting point is the classification of finite-dimensional Nichols algebras over non-abelian groups by Heckenberger–Vendramin, which consist of low rank exceptions and large rank families. We prove that the large rank families are cocycle twists of Nichols algebras constructed by the second author as foldings of Nichols algebras of Cartan type over abelian groups by outer automorphisms. This enables us to give uniform Lie-theoretic descriptions of the large rank families, prove generation in degree one and construct liftings. We also show that every lifting is a cocycle deformation of the corresponding coradically graded Hopf algebra using an explicit presentation by generators and relations of the Nichols algebra.

On the level of tensor categories, we construct families of graded extensions of the representation category of a quantum group by a group of diagram automorphism.

CONTENTS

1. Introduction	2
1.1. Background	2
1.2. Goals and content	5
2. Preliminaries	6
2.1. The Nichols algebra of a braided vector space	6
2.2. Nichols algebras over non-abelian groups	8
2.3. Heckenberger–Vendramin classification	10
2.4. On the structure of the group G	12
3. Foldings of Nichols algebras and trivializing the action of the center	13
3.1. Categorical action on Yetter-Drinfeld modules	13
3.2. Folding construction	15

2020 *Mathematics Subject Classification.* 16T05, 17B37.

The work of I. A. was partially supported by CONICET and Secyt (UNC). The authors were partially supported by the Alexander von Humboldt Foundation through the Research Group Linkage Programme.

3.3. Folding data for trivial action of κ	16
3.4. Trivializing the action of κ via a twist	18
4. Generation in degree one	21
5. Generators and relations for Nichols algebras	21
5.1. Types α_2 and α_3	22
5.2. A kind of distinguished pre-Nichols algebra	27
5.3. The subalgebra of coinvariants	30
5.4. A presentation of the Nichols algebra	34
5.5. Rigidity of Nichols algebras	35
6. Liftings of Nichols algebras	37
6.1. Liftings of type α_2	38
6.2. The general case	40
6.3. Foldings of liftings	43
7. Future directions	46
Appendix A. Proof of Theorem 3.18	47
A.1. Group Cohomology Tools	47
A.2. Non-abelian groups and 2-cocycles	50
A.3. Proof of Proposition 3.15	52
Acknowledgements	55
References	56

1. INTRODUCTION

1.1. **Background.** Groups and Lie algebras have in common that they both admit a tensor product of representations and a dual representation; in other words, their categories of representations are tensor categories. More generally, the category of representations of a Hopf algebra is also a tensor category. Prominent examples of Hopf algebras are the quantum groups $U_q(\mathfrak{g})$ by Drinfeld-Jimbo [Dr, J], which are deformations of the enveloping algebra of a semisimple Lie algebra \mathfrak{g} by a formal parameter q , and the small quantum groups $u_q(\mathfrak{g})$ by Lusztig [Lu2], which are finite-dimensional non-semisimple quotients of $U_q(\mathfrak{g})$ for q a root of unity. One of the initial motivations for quantum groups was their relation to the monodromy of certain differential equations in conformal field theory [FBZ] and invariants of knots and 3-manifolds [Tur]. Small quantum groups, their representation categories and their semisimplification are related to Lie algebras in positive characteristic [Lu1, AJS] and affine Lie algebras [Fu], and they have again applications to topology [KL] and conformal field theory [FGST, L3].

While the classification of finite-dimensional semisimple Hopf algebras is still a very hard problem, we may ask for the classification of Hopf algebras H with a *given* maximal cosemisimple part H_0 , called the coradical. For example, if $H_0 = \mathbb{k}G$ is a group ring, then H is called a pointed Hopf algebra. Andruskiewitsch and Schneider proposed a program called the *Lifting method* to classify finite-dimensional Hopf algebras whose coradical is a subalgebra, that we sketch now. Every Hopf algebra comes with a coalgebra filtration

$H_0 \subseteq H_1 \subseteq \dots$ called the coradical filtration, where H_0 is the coradical. If we assume that H_0 is a subalgebra, then the associated graded coalgebra $\text{gr } H$ is a Hopf algebra; we pay special attention to the subspace $\bar{H}_1 = H_1/H_0$ and the subalgebra $\mathcal{B}(\bar{H}_1)$ generated by \bar{H}_1 , which is the Nichols algebra discussed below. As an example, for the (infinite-dimensional) quantum group $H = U_q(\mathfrak{g})$ the coradical H_0 is the group algebra spanned by the root lattice, the space \bar{H}_1 is spanned by the simple root vectors E_i, F_i , the Nichols algebra is the tensor product of the quantum Borel algebras $U_q(\mathfrak{g})^\pm$, and in the graded algebra $\text{gr } H$ the relation $[E_i, F_i] = 0$ holds in contrast to the nontrivial relation $[E_i, F_i] = K_i - K_i^{-1}$ in H . Then the classification proceeds in three steps: First, one classifies all finite-dimensional Nichols algebras over a fixed coradical H_0 . Second, one checks if $H_0 \oplus \bar{H}_1$ generate the entire algebra $\text{gr } H$ which is known as the generation in degree one problem. Third, one determines all possible Hopf algebras H associated to each Nichols algebra in the first step, the so-called liftings. Using these ideas, Andruskiewitsch and Schneider obtained the whole classification of finite-dimensional Hopf algebras over \mathbb{C} with coradical H_0 a group algebra of an abelian group of order not divisible by 2, 3, 5, 7 [AS3]. The Nichols algebras appearing in this case are the Borel parts of small quantum groups, generation in degree one always holds and the possible liftings are described by deforming relations such as $[E_i, F_i] = 0$ and $E_i^\ell = 0$.

The Nichols algebra $\mathcal{B}(V)$ of a braided vector space V , or more generally of an object V in a braided tensor category, is a quotient of the tensor algebra of V , which satisfies several universal properties. In particular, it is the unique Hopf algebra (in the braided tensor category) which is generated by V and contains V as the space of primitive elements. It is a difficult problem to determine the structure of the Nichols algebra of a given braided vector space, even to determine whether it is finite-dimensional. In view of the Lifting Method for pointed Hopf algebras, we need to consider Yetter-Drinfeld modules V over a group algebra $H_0 = \mathbb{k}G$. If G is a finite abelian group, then the braiding is of diagonal type. For braidings of diagonal type, Heckenberger classified the complex finite-dimensional Nichols algebras [H2]. Besides the Nichols algebras $u_q(\mathfrak{g})^+$ coming from small quantum groups, he found Nichols algebras associated to Lie superalgebras, and several exceptional Nichols algebras involving roots of unity of order divisible by 2, 3, 5, 7; many of the later are now known to be related to Lie superalgebras in positive characteristic [AA]. A main structural insight [H1, HeY, HS3] is the existence of generalized root systems and Weyl groupoids. Roughly speaking, the generalization comes from the fact that different Weyl chambers might be in fact different, since reflections may change the braiding and even the Cartan matrix - an effect that already appears for contragredient Lie superalgebras. Nevertheless, finite Weyl groupoids can be classified [CH] and show again a pattern of Lie algebra and Lie superalgebra series, plus low rank exceptions. The new Nichols algebras lead to new pointed Hopf algebras over abelian groups of order divisible by 2, 3, 5, 7, which include the small quantum groups $u_q(\mathfrak{g})$ associated to Lie superalgebras \mathfrak{g} . Using a presentation by generators and relations of these Nichols algebras of diagonal type, Angiono proved generation in degree one [An2]. The final answer about the liftings was established a few years later [AnG] and involves the proof that every lifting is a cocycle deformation of the associated graded Hopf algebra in the sense of [DoT, Ma].

We now discuss Nichols algebras $\mathcal{B}(V)$ of Yetter-Drinfeld modules V over a non-abelian group G . In this case, irreducible Yetter-Drinfeld modules V are parametrized by a conjugacy class g^G in G and a representation ρ of the centralizer G^g of g . A general Yetter-Drinfeld module V is the direct sum of such irreducible modules, their number is called the rank of V and they correspond to the simple roots of a generalized root system. The study of Nichols algebras over non-abelian groups starts with [MiS], where G is a Coxeter group and g is a reflection. They treat two main examples: The symmetric group \mathbb{S}_n has a single conjugacy class of reflections or transpositions with $\binom{n}{2}$ elements, accordingly V is a Yetter-Drinfeld module of rank 1 (i.e. irreducible) with dimension $\binom{n}{2}$. As it turns out, the associated Nichols algebras for $n = 3, 4, 5$ have dimension 12, 576, 8294400 and were considered by Fomin and Kirillov [FK] in a very different context; for $n \geq 6$ they are conjecturally infinite-dimensional. On the other hand, the dihedral group \mathbb{D}_4 has two conjugacy classes of reflections, each with two elements. Accordingly, $V = V_1 \oplus V_2$ has dimension $2 + 2$ and rank 2. As it turns out, the generalized root system is of type A_2 , indicating roughly that there is a space of braided commutators V_{12} , associated to the third conjugacy class with two elements, and all higher commutators vanish. Since the Nichols algebras of the irreducible modules V_1, V_2, V_{12} have each dimension 4, the Nichols algebra $\mathcal{B}(V)$ has dimension 4^3 .

The study now naturally branches into two directions: Nichols algebras of rank 1, meaning of irreducible Yetter-Drinfeld modules, and Nichols algebras of rank > 1 composed of the former via root system theory. In rank 1, more finite-dimensional examples of Nichols algebras were discovered in [Gr, AG, HLV], and later the research concentrated on successfully ruling out finite-dimensional Nichols algebras over most simple groups, see [ACG] and the references there.

In rank > 1 , Lentner has developed the folding method for central extensions of Hopf algebras [L1, L2]. When applied to Nichols algebras, it takes a Nichols algebra $\mathcal{B}(V)$ over a group Γ with an additional (twisted) symmetry group Z , and converts it to a Nichols algebra $\mathcal{B}(\tilde{V})$ over the central extension of Γ . In particular, starting with Γ an abelian group, $\mathcal{B}(V) = u_q(\mathfrak{g})^+$ the Borel part of a small quantum group for $q^2 = -1$ and Z generated by a diagram automorphism of \mathfrak{g} , the folding method produces a Nichols algebra over central extensions G of the abelian group Γ . There are cases where \mathfrak{g} is simple, namely ${}^2A_{2n+1}$ and 2E_6 , and cases where \mathfrak{g} consists of two copies of the same simply-laced Lie algebra interchanged by $Z = \mathbb{Z}_2$, which we denote ${}^2A_n^2$ and ${}^2D_n^2$ and ${}^2E_n^2$, $n = 6, 7, 8$. The root system attached to $\mathcal{B}(V)$ is the folded root system considered in Lie theory, with Z -orbits of roots becoming the new roots and G -conjugacy classes. For example, the smallest case ${}^2A_2^2$ reproduces the Nichols algebra over \mathbb{D}_4 discussed above, and the 6 roots of A_2^2 become 3 roots of A_2 , each attached to a conjugacy class with two elements. In view of the exotic examples above, let us mention that these Nichols algebras are close to the quantum group case: By construction, their braiding can still be diagonalized, the group G is close to abelian, and the Weyl groupoid is again a Weyl group: they are *standard*.

Around the same time, Heckenberger, Schneider and Vendramin started a systematic classification of finite-dimensional Nichols algebras over non-abelian groups over arbitrary fields using the root system theory, starting in rank 2 in [HS1] and culminating in a full classification in rank ≥ 2 [HV] in 2014. Their surprising observation was that the existence

of a finite root system severely restricts the possible groups G , so that only very few of the (not yet fully classified) Nichols algebras of rank 1 can appear in rank ≥ 2 . They found (in characteristic zero) three new exceptional Nichols algebras in rank 2 and 3, three infinite series $\alpha_n, \delta_n, \gamma_n$, and exceptional ones ϵ_n, ϕ_4 with standard root systems A_n, D_n, C_n , respectively E_n, F_4 , whose conjugacy classes have two elements for short roots and one element for long roots. Note that these families precisely match the folding above.

1.2. Goals and content. The purpose of this paper is to construct all pointed Hopf algebras whose group of group-like elements is a non-abelian group G and the Nichols algebra is of standard type $\alpha_n, \delta_n, \gamma_n, \epsilon_n$ or ϕ_4 . Our first main result is that these series are all twists of the folded Nichols algebras by 2-cocycles of the group G . This opens the possibility for a uniform treatment of these Nichols algebras. Our second main result is that all Hopf algebras with coradical and a infinitesimal braiding as above are already generated in degree one. Since the folding method also applies to Hopf algebras, we construct a family of liftings inspired by those of diagonal type, and then fold accordingly. Our last main result is that these are already all liftings. To achieve this, we give an explicit presentation of the Nichols algebras by generators and relations, which also allows us to prove that these Nichols algebras are rigid.

We now discuss the content of this paper in more detail.

In §2 we review the preliminaries and the classification result by Heckenberger–Vendramin.

Next we recall the folding construction in §3. Because we wish to work also with Hopf algebras, we give an exposition that emphasizes the origins of the folding method in this setting, as in [L1]: For a given Hopf algebra H and a group of biGalois objects, the direct sum is again a Hopf algebra. Now we specialize to the case where the Hopf algebra is a smash product $H = \mathcal{B}(V) \# \mathbb{k}\Gamma$ and the biGalois objects are based on a 2-cocycle σ on Γ and a twisted symmetry of V . Then, the result of the folding is again a smash product of the centrally extended group with the folded Nichols algebra. Our first main result is Theorem 3.18, which states that the infinite families of Nichols algebras are twists of foldings over central extensions $Z \rightarrow G \rightarrow \Gamma$. By construction, Z acts trivially on the folded Nichols algebras. We thus have to go through the classification of Heckenberger and Vendramin, assume a particular action of Z and prove that the resulting groups G possess sufficient cohomology to twist this action to zero. For large rank, the action of Z turns out to be always trivial, but for small rank there are different possibilities depending on the structure of the group. For example, ${}^2A_2^2$ can be realized over central extensions such as \mathbb{D}_4 and \mathbb{Q}_8 , in the former case we find nontrivial Z -action and sufficient cohomology, in the latter case we only find trivial Z -action. Since these computations are lengthy and involve tools from group cohomology, they are confined to the Appendix A.

In §4 we state our second main result, Theorem 4.1, which gives a positive answer to the generation in degree one problem.

The main result of §5 is Theorem 5.7, where the Nichols algebra is presented by generators and relations, and a PBW basis is obtained. Using the diagonal setting as inspiration, we give G -homogeneous generators and relations, which is necessary to deform the Hopf algebra structure. We also show that our Nichols algebras are rigid in the sense of [AnKM].

In §6 we construct a big family of liftings using Hopf-Galois objects as in [AAG, AnG, AnS]. Our last main result is Theorem 6.3, where we show that this family exhaust all liftings, and that all liftings are cocycle deformation of the associated graded Hopf algebra.

In §7 we list some open questions and future directions of research.

2. PRELIMINARIES

Conventions. We denote $\mathbb{N} = \{1, 2, 3, \dots\}$ and $\mathbb{N}_0 = \{0\} \cup \mathbb{N}$. Given $k < \theta$ in \mathbb{N}_0 we put $\mathbb{I}_{k,\theta} = \{n \in \mathbb{N}_0 : k \leq n \leq \theta\}$ and $\mathbb{I}_\theta = \mathbb{I}_{1,\theta}$. When θ is clear from the context we just write $\mathbb{I} = \mathbb{I}_\theta$. The canonical basis of \mathbb{Z}^θ is denoted by $(\alpha_i)_{i \in \mathbb{I}_\theta}$.

We work over an algebraically closed field \mathbb{k} of characteristic zero and use \mathbb{k}^\times to denote the group of non-zero elements. If $N \in \mathbb{N}$, we use \mathbb{G}_N to denote the subgroup of N -th roots of unity; the subset of those with order N is \mathbb{G}'_N .

Given a group G and an element g , we use g^G and G^g to denote the conjugacy class and the centralizer of g , respectively. By \widehat{G} we mean the group of characters, and $\mathbb{k}G$ stands for the group algebra. If K is another group, then a pairing (also called a bicharacter) is a map $P : G \times K \rightarrow \mathbb{k}^\times$ such that for all $g, g' \in G, k, k' \in K$:

$$P(gg', k) = P(g, k)P(g', k), \quad P(g, kk') = P(g, k)P(g, k').$$

A skew-polynomial algebra in variables z_1, \dots, z_k is a quotient of the free algebra in these variables by an ideal generated by $z_i z_j - t_{ij} z_j z_i$, $1 \leq i, j \leq k$, for some $t_{ij} \in \mathbb{k}^\times$.

We denote Hopf algebras by tuples $(H, \mu, \Delta, \mathcal{S})$ where μ is the multiplication, Δ the comultiplication and \mathcal{S} the antipode, which we always assume bijective. The subspace of primitive elements is $\mathcal{P}(H)$. The group of group-like elements is $G(H)$. If $\delta : V \rightarrow H \otimes V$ is a left H -comodule we write $\delta(v) = v_{-1} \otimes v_0$; for $g \in G(H)$ we put $V_g := \{v \in V : \delta(v) = g \otimes v\}$. We refer to [Mo] for any unexplained terminology on Hopf algebras and to [A+, §2] for preliminaries on Hopf-Galois objects and cocycle deformations.

2.1. The Nichols algebra of a braided vector space. A *braided vector space* is a pair (V, c) where V is a vector space and $c \in GL(V \otimes V)$ satisfies

$$(c \otimes \text{id})(\text{id} \otimes c)(c \otimes \text{id}) = (\text{id} \otimes c)(c \otimes \text{id})(\text{id} \otimes c).$$

By declaring the elements of V to be primitive, the tensor algebra $T(V)$ becomes an \mathbb{N}_0 -graded *braided* Hopf algebra. There is a largest coideal $\mathcal{J}(V)$ among those that trivially intersect $\mathbb{k} \oplus V$; it happens to be graded so we denote $\mathcal{J}(V) = \bigoplus_{n \geq 2} \mathcal{J}^n(V)$. The Nichols algebra of (V, c) is defined as the quotient $\mathcal{B}(V) = T(V)/\mathcal{J}(V)$. This is again an \mathbb{N}_0 -graded braided Hopf algebra, which is strictly graded as a coalgebra and generated by V as an algebra, see [HS3, §7]. Any intermediate quotient $\mathcal{B} = T(V)/\mathcal{J}$ by an \mathbb{N}_0 -homogeneous Hopf ideal \mathcal{J} is called a *pre-Nichols algebra* of V .

The *braided commutator* of $T(V)$ is defined by

$$[-, -]_c = \text{mult}(\text{id} - c) : T(V) \otimes T(V) \rightarrow T(V).$$

If $u \in V$ and $v \in T(V)$ we denote $(\text{ad}_c u)v = [u, v]_c$. In §2.2.1 will define $\text{ad}_c u$ for arbitrary $u \in T(V)$. For a fixed basis $(x_i)_{i \in I}$ of V and $k \geq 2$ we set

$$(2.1) \quad x_{i_1 \dots i_k} := (\text{ad}_c x_{i_1}) \cdots (\text{ad}_c x_{i_{k-1}}) x_{i_k}, \quad i_j \in \mathbb{I}.$$

Example 2.1. Given $\mathfrak{q} = (q_{ij})_{i,j \in \mathbb{I}}$ a matrix of elements of \mathbb{k}^\times , there is a braided vector space $(V, c^{\mathfrak{q}})$ where V has basis $(\mathbf{x}_i)_{i \in \mathbb{I}}$ and $c^{\mathfrak{q}}$ is given by

$$(2.2) \quad c^{\mathfrak{q}}(\mathbf{x}_i \otimes \mathbf{x}_j) = q_{ij} \mathbf{x}_j \otimes \mathbf{x}_i, \quad i, j \in \mathbb{I}.$$

A braided vector space is called *of diagonal type* [AS2] if (2.2) holds in some basis of V for some $\mathfrak{q} = (q_{ij})_{i,j \in \mathbb{I}}$. In this case, we denote the Nichols algebra of (V, c) by $\mathcal{B}_{\mathfrak{q}}$, which is now $\mathbb{Z}^{\mathbb{I}}$ -graded; we refer to \mathfrak{q} as the *braiding matrix*. The Dynkin diagram of \mathfrak{q} is a graph with \mathbb{I} as the set of vertices, each vertex i labelled with q_{ii} . There is an edge between $i \neq j$ if and only if $\tilde{q}_{ij} := q_{ij}q_{ji} \neq 1$; such an edge is labelled with this scalar.

We say that \mathfrak{q} is *of Cartan type* [AS1] if there is a Cartan matrix $\mathbf{a} = (a_{ij})$ such that

$$q_{ij}q_{ji} = q_{ii}^{a_{ij}}, \quad \text{for all } i, j \in \mathbb{I}.$$

If some q_{ii} is not a root of unity, then the integers a_{ij} are uniquely determined. Otherwise we impose $-\text{ord } q_{ii} < a_{ij} \leq 0$ for all $j \neq i$. In this case we say that \mathfrak{q} is of Cartan type \mathbf{a} .

Although braided vector spaces of Cartan type seem quite *simple*, the structure of the corresponding Nichols algebra is related either with quantized enveloping algebras (when the entries of \mathfrak{q} are not roots of unity), or with Frobenius-Lusztig kernels.

The following example of Cartan type will be particularly relevant in later sections.

Example 2.2. Fix a finite Cartan matrix \mathbf{a} with simply-laced Dynkin diagram. Assume that $\mathfrak{q} = (q_{ij})$ satisfies the following conditions:

$$(2.3) \quad q_{ii} = -1; \quad q_{ij}q_{ji} = \begin{cases} -1, & a_{ij} = -1, \\ 1, & a_{ij} = 0; \end{cases} \quad i \neq j \in \mathbb{I}.$$

Then \mathfrak{q} is of Cartan type \mathbf{a} .

Let $\beta_1 < \beta_2 < \cdots < \beta_M$ be a convex order on the set of positive roots Δ_+ of \mathbf{a} . In [An2] we can find a *root vector* $\mathbf{x}_\beta \in \mathcal{B}_{\mathfrak{q}}$ for each $\beta \in \Delta_+$, of $\mathbb{Z}^{\mathbb{I}}$ -degree β , obtained recursively as braided commutator of root vectors with smaller degree.

In this case, the Nichols algebra $\mathcal{B}_{\mathfrak{q}}$ is presented by generators $(\mathbf{x}_i)_{i \in \mathbb{I}}$ and relations

$$(2.4) \quad \mathbf{x}_\alpha^2 = 0, \quad \alpha \in \Delta_+;$$

$$(2.5) \quad [\mathbf{x}_{ijk}, \mathbf{x}_j]_c = 0, \quad a_{ij} = a_{jk} = -1;$$

$$(2.6) \quad \mathbf{x}_{ij} = 0, \quad a_{ij} = 0.$$

Moreover, by [An2, AA], a basis for $\mathcal{B}_{\mathfrak{q}}$ is given by the set

$$(2.7) \quad \{\mathbf{x}_{\beta_1}^{n_1} \mathbf{x}_{\beta_2}^{n_2} \cdots \mathbf{x}_{\beta_M}^{n_M} \mid n_i \in \{0, 1\}\}.$$

Remark 2.3. Let \mathfrak{q} as above. In §5 we will need the following constructions, due to [An4].

(I) The *distinguished pre-Nichols algebra* $\widetilde{\mathcal{B}}_{\mathfrak{q}}$ is the quotient of $T(V)$ by (2.5), (2.6) and

$$(2.8) \quad \mathbf{x}_{ij} = 0, \quad a_{ij} = -1.$$

The set $\{\mathbf{x}_{\beta_1}^{n_1} \mathbf{x}_{\beta_2}^{n_2} \dots \mathbf{x}_{\beta_M}^{n_M} \mid n_i \in \mathbb{N}_0\}$ is a basis of $\widetilde{\mathcal{B}}_{\mathfrak{q}}$.

(II) Let $\pi : \widetilde{\mathcal{B}}_{\mathfrak{q}} \rightarrow \mathcal{B}_{\mathfrak{q}}$ be the canonical projection. By [An4], the subalgebra of coinvariants $\mathcal{Z}_{\mathfrak{q}} := \widetilde{\mathcal{B}}_{\mathfrak{q}}^{\text{co}\pi}$ is a q -polynomial algebra with generators \mathbf{x}_{β}^2 , $\beta \in \Delta_+$.

(III) Let $\widehat{\mathcal{B}}_{\mathfrak{q}}$ be the quotient of $T(V)$ by (2.5), (2.6) and \mathbf{x}_i^2 , $i \in \mathbb{I}$. Then $\widehat{\mathcal{B}}_{\mathfrak{q}}$ is a pre-Nichols algebra, which coincides with the quotient of $\widetilde{\mathcal{B}}_{\mathfrak{q}}$ by \mathbf{x}_i^2 , $i \in \mathbb{I}$. The set

$$\{\mathbf{x}_{\beta_1}^{n_1} \mathbf{x}_{\beta_2}^{n_2} \dots \mathbf{x}_{\beta_M}^{n_M} \mid n_i \in \{0, 1\} \text{ if } \beta_i \text{ is simple, } n_i \in \mathbb{N}_0 \text{ otherwise}\}$$

is a basis of $\widehat{\mathcal{B}}_{\mathfrak{q}}$, so its Hilbert series is

$$\mathcal{H}_{\widehat{\mathcal{B}}_{\mathfrak{q}}} = \left(\prod_{\beta \in \Delta_+ - \{\alpha_i\}} \frac{1}{1 - t^{\beta}} \right) \left(\prod_{i \in \mathbb{I}} 1 + t_i \right).$$

2.2. Nichols algebras over non-abelian groups. The goal of this subsection is to introduce the notion of Weyl groupoids. These play a fundamental role in the classification achieved in [HV]. We refer to the book [HS3] for details and unexplained terminology.

2.2.1. Yetter-Drinfeld modules over groups. For a group G , the category of Yetter-Drinfeld modules ${}_{\mathbb{k}G}^{\mathbb{k}G}\mathcal{YD}$ consist of G -graded vector spaces $V = \bigoplus_{g \in G} V_g$ endowed with a G -action such that $h \cdot V_g \subset V_{hgh^{-1}}$ for all $h, g \in G$. This is a braided tensor category where the braiding $c_{V,W} : V \otimes W \rightarrow W \otimes V$ is given by $c(v \otimes w) = g \cdot w \otimes v$ for $v \in V_g$ and $w \in W$. We recall that a G -grading on a vector space $V = \bigoplus_{g \in G} V_g$ is equivalent to a $\mathbb{k}G$ -comodule structure $\delta : V \rightarrow V \otimes \mathbb{k}G$, declaring $\delta(v) = g \otimes v$ if and only if $v \in V_g$. The *support* of V is

$$\text{supp } V = \{g \in G \mid V_g \neq 0\}.$$

Let $(R, \mu, \Delta, \mathcal{S})$ be a Hopf algebra in ${}_{\mathbb{k}G}^{\mathbb{k}G}\mathcal{YD}$. The braided commutator defined above satisfies the following identity: If $u \in R_g$ and $v \in R_h$ for some $g, h \in G$, then for any $w \in R$,

$$(2.9) \quad [[u, v]_c, w] = [u, [v, w]_c] - (g \cdot v)[u, w]_c + [u, h \cdot w]_c v.$$

R admits a *braided* adjoint representation $\text{ad}_c : R \rightarrow \text{End}(R)$ given by

$$(\text{ad}_c u)v = \mu(\mu \otimes \mathcal{S})(\text{id} \otimes c)(\Delta \otimes \text{id})(u \otimes v), \quad u, v \in R.$$

When u is primitive, $\text{ad}_c u$ and $[u, -]_c$ coincide. Notice also that

$$(2.10) \quad g \cdot ((\text{ad}_c u)v) = (\text{ad}_c g \cdot u)(g \cdot v), \quad g \in G, u \in \mathcal{P}(R), v \in R.$$

2.2.2. The Nichols algebra of a Yetter-Drinfeld module. Each $V \in {}_{\mathbb{k}G}^{\mathbb{k}G}\mathcal{YD}$ is a braided vector space, so it has a Nichols algebra $\mathcal{B}(V)$ as discussed in §2.1. In this setting, $T(V)$ and $\mathcal{B}(V) = T(V)/\mathcal{J}(V)$ turn out to be \mathbb{N}_0 -graded Hopf algebras in ${}_{\mathbb{k}G}^{\mathbb{k}G}\mathcal{YD}$.

2.2.3. *Skew derivations.* There is a criterion to decide if any given element of $T(V)$ belongs to $\mathcal{J}(V)$. Fix a basis x_1, \dots, x_r of V with x_i of degree h_i . For each i , we define a skew derivation $\partial_{x_i} \in \text{End}(T(V))$ recursively in $V^{\otimes n}$, $n \geq 0$. For $n = 0$, put $\partial_{x_i}(1) = 0$; in $V^{\otimes 1}$ put $\partial_{x_i}(x_j) = \delta_{i,j}$ and in general define

$$\partial_{x_i}(xy) = x\partial_{x_i}(y) + \partial_{x_i}(x)h_i \cdot y, \quad x, y \in T(V).$$

Then, for any $n \geq 2$, we have $x \in \mathcal{J}^n(V)$ if and only if $\partial_{x_i}(x) \in \mathcal{J}^{n-1}(V)$ for all i . The compositions of these derivations with the braided adjoint action satisfy

$$(2.11) \quad \partial_{x_i}((\text{ad}_c u)v) = u\partial_{x_i}(v) - \partial_{x_i}(g \cdot v)h_i \cdot u, \quad u \in V_g, v \in T(V), i \in \mathbb{I}_r.$$

2.2.4. *Simple Yetter-Drinfeld modules.* Fix $g \in G$ and (V, ρ) an irreducible representation of the centralizer G^g . We consider the induced G -module $\mathbb{k}G \otimes_{G^g} V$ endowed with the G -grading determined by declaring the degree of $x \otimes v$ to be gxg^{-1} for all $x \in G$ and $v \in V$. This is a simple Yetter-Drinfeld module over G , which is denoted by $M(g^G, \rho)$ since it depends on the conjugacy class g^G rather than the element g itself. Moreover, all the simple objects of ${}^{\mathbb{k}G}\mathcal{YD}$ arise in this way, and if G is finite the category ${}^{\mathbb{k}G}\mathcal{YD}$ is semisimple. There is a concrete description of $M(g^G, \rho)$ in [A, Example 24], which we will use several times to perform computations in ${}^{\mathbb{k}G}\mathcal{YD}$.

Example 2.4. Let G be a finite group. Assume that $g \in G$ is such that $g^G = \{g, g\kappa\}$ for some $\kappa \neq e \in G$. Then there exists $g_0 \in G$ such that $g_0g = \kappa gg_0$, $G^g = G^{g\kappa} = G^{g^{-1}}$ is a subgroup of index two since $G = G^g \cup g_0G^g$, and $\kappa \in Z(G)$, $\kappa^2 = e$.

Let χ be a one-dimensional representation of G^g , that is $V = \mathbb{k}$ and $\chi \in \widehat{G^g}$, and $M = M(g^G, \chi)$. Then $\dim M = 2$: we may fix a basis $\{x, y\}$ such that $y = g_0 \cdot x$ and the coaction is given by $\rho(x) = g \otimes x$, $\rho(y) = g\kappa \otimes y$. As $hg_0, g_0^{-1}h \in G^g$ for all $h \notin G^g$, the action satisfies

$$h \cdot x = \begin{cases} \chi(h)x, & h \in G^g, \\ \chi(g_0^{-1}h)y, & h \notin G^g; \end{cases} \quad h \cdot y = \begin{cases} \chi(g_0^{-1}hg_0)y, & h \in G^g, \\ \chi(hg_0)x, & h \notin G^g; \end{cases} \quad k \in G.$$

Thus M is of diagonal type with braiding matrix $\begin{bmatrix} \chi(g) & \chi(g\kappa) \\ \chi(g\kappa) & \chi(g) \end{bmatrix}$; from this we deduce that $\dim \mathcal{B}(M) < \infty$ if and only if $\chi(g) \in \{-1\} \cup \mathbb{G}'_3$.

2.2.5. *Weyl groupoid.* Next we recall the definition of the Weyl groupoid of a non-simple element $M \in {}^{\mathbb{k}G}\mathcal{YD}$ such that $\dim \mathcal{B}(M) < \infty$. Let $M = \bigoplus_{i \in \mathbb{I}} M_i$, where each summand M_i is simple: $M_i = M(g_i^G, \rho_i) \in {}^{\mathbb{k}G}\mathcal{YD}$ for some $g_i \in G$, ρ_i an irreducible representation of G^{g_i} .

For each $i \neq j \in \mathbb{I}$, set $(\text{ad } M_i)^0 M_j := M_j$, and for $n \in \mathbb{N}$,

$$(\text{ad } M_i)^n M_j := \{(\text{ad}_c m_1) \cdots (\text{ad}_c m_n)m \mid m_\ell \in M_i, m \in M_j\} \subset \mathcal{B}(M).$$

The *generalized Cartan matrix* of M is $C^M = (c_{ij}^M) \in \mathbb{Z}^{\mathbb{I} \times \mathbb{I}}$, where

$$(2.12) \quad c_{ii}^M = 2, \quad c_{ij}^M = -\min\{n \in \mathbb{N}_0 \mid (\text{ad } M_i)^{n+1} M_j = 0\}, \quad j \neq i.$$

The i -reflection of M is $\rho_i M = \sum_{j \in \mathbb{I}} \underline{M}_j$, where

$$\underline{M}_j = \begin{cases} (\text{ad } M_i)^{-c_{ij}^M} M_j, & j \neq i; \\ M_i^*, & j = i. \end{cases}$$

Each $\underline{M}_j \in \frac{\mathbb{k}^G}{\mathbb{k}G} \mathcal{YD}$ is simple and $\dim \mathcal{B}(M) = \dim \mathcal{B}(\rho_i M)$.

These reflections generate the *Weyl groupoid* of M , see [HS3].

2.3. Heckenberger–Vendramin classification. Let $M = \bigoplus_{i \in \mathbb{I}_\theta} M_i$ be a finite dimensional Yetter-Drinfeld module over a non-abelian group G , where each M_i is simple. Here θ is called the *rank* of M .

In [HV, Theorem 2.5] the authors classify Yetter-Drinfeld modules as above of rank at least 2, such that the associated Nichols algebra is finite-dimensional. To be precise, one needs to assume that the support of M generates G , and impose a mild non-degeneracy condition on the braiding between different summands of M . Up to a few exceptions in ranks two and three, the classification consists on families α_θ , γ_θ and δ_θ of arbitrary rank, and types ϵ_θ , $\theta = 6, 7, 8, \phi_4$, which resemble the classification of finite-dimensional Lie algebras. These Yetter-Drinfeld modules are invariant under the Weyl groupoid action; we say that they are *standard*, adopting the terminology used for diagonal type, see [AA]. In this paper we study this class; next we give an explicit description of each module.

2.3.1. Types α_θ , δ_θ and ϵ_θ . Fix a simply-laced indecomposable Cartan matrix $\mathbf{a} = (a_{ij}) \in \mathbb{Z}^{\theta \times \theta}$ of finite type; i.e. of type A_θ , $\theta \geq 2$, D_θ , $\theta \geq 4$, or E_θ for $\theta \in \mathbb{I}_{6,8}$. Let Δ_+ be the set of positive roots. Following [HV, Lemma 6.2] we describe a Yetter-Drinfeld module $M = \bigoplus_{i \in \mathbb{I}} M_i$ over a non-abelian group G with simply-laced skeleton and Cartan graph of standard type \mathbf{a} . Assume there exist

- $\kappa \in Z(G)$ such that $\kappa \neq 1$, $\kappa^2 = 1$,
- $g_i \in G$ with $g_i^G = \{g_i, \kappa g_i\}$ for all $i \in \mathbb{I}_\theta$,
- $\chi_i \in \widehat{G^{g_i}}$ such that $\chi_i(g_i) = -1$ for all $i \in \mathbb{I}_\theta$,

satisfying the following:

$$(2.13) \quad g_i g_j = \kappa g_j g_i, \quad \chi_i(\kappa g_j^2) \chi_j(\kappa g_i^2) = 1, \quad a_{ij} = -1;$$

$$(2.14) \quad g_i g_j = g_j g_i, \quad \chi_i(g_j) \chi_j(g_i) = 1, \quad a_{ij} = 0;$$

$$(2.15) \quad \chi_i(\kappa) \chi_j(\kappa) = 1, \quad a_{ij} = 0;$$

For $i \in \mathbb{I}_\theta$ let $M_i = M(g_i^G, \chi_i) \in \frac{\mathbb{k}^G}{\mathbb{k}G} \mathcal{YD}$, which has a basis $\{x_i, x_{\bar{i}}\}$ with coaction $x_i \mapsto g_i \otimes x_i$, $x_{\bar{i}} \mapsto \kappa g_i \otimes x_{\bar{i}}$. As in [A, Example 37] the braidings c_{M_i, M_j} for $i, j \in \mathbb{I}_\theta$ are determined by

$$(2.16) \quad \begin{bmatrix} c(x_i \otimes x_i) & c(x_i \otimes x_{\bar{i}}) \\ c(x_{\bar{i}} \otimes x_i) & c(x_{\bar{i}} \otimes x_{\bar{i}}) \end{bmatrix} = \begin{bmatrix} -x_i \otimes x_i & -\chi_i(\kappa) x_{\bar{i}} \otimes x_i \\ -\chi_i(\kappa) x_i \otimes x_{\bar{i}} & -x_{\bar{i}} \otimes x_{\bar{i}} \end{bmatrix};$$

$$(2.17) \quad \begin{bmatrix} c(x_i \otimes x_j) & c(x_i \otimes x_{\bar{j}}) \\ c(x_{\bar{i}} \otimes x_j) & c(x_{\bar{i}} \otimes x_{\bar{j}}) \end{bmatrix} = \begin{bmatrix} x_{\bar{j}} \otimes x_i & \chi_j(g_i^2) x_j \otimes x_i \\ \chi_j(\kappa) x_{\bar{j}} \otimes x_{\bar{i}} & \chi_j(\kappa g_i^2) x_j \otimes x_{\bar{i}} \end{bmatrix}, \quad a_{ij} = -1;$$

$$(2.18) \quad \begin{bmatrix} c(x_i \otimes x_j) & c(x_i \otimes x_{\bar{j}}) \\ c(x_{\bar{i}} \otimes x_j) & c(x_{\bar{i}} \otimes x_{\bar{j}}) \end{bmatrix} = \begin{bmatrix} \chi_j(g_i) x_j \otimes x_i & \chi_j(\kappa g_i) x_{\bar{j}} \otimes x_i \\ \chi_j(\kappa g_i) x_j \otimes x_{\bar{i}} & \chi_j(g_i) x_{\bar{j}} \otimes x_{\bar{i}} \end{bmatrix}, \quad a_{ij} = 0.$$

The Cartan matrix of $M = \bigoplus_{i \in \mathbb{I}} M_i$ is \mathbf{a} , and we have $\dim \mathcal{B}(M) = 2^{2|\Delta|}$.

Remark 2.5. • The braiding on M_i is of diagonal type and $c_{M_i, M_i}^2 = \text{id}$.

- If $a_{ij} = 0$, then $M_i \oplus M_j$ is of diagonal type and $c_{M_i, M_j} c_{M_j, M_i} = \text{id}_{M_j \otimes M_i}$.
- If $a_{ij} = -1$, then $M_i \oplus M_j$ is of diagonal type if and only if $\chi_i(\kappa) = \chi_j(\kappa) = 1$, but here $c_{M_i, M_j} c_{M_j, M_i} \neq \text{id}_{M_j \otimes M_i}$.

2.3.2. *Type γ_θ , $\theta \geq 3$.* Following [HV, Lemma 7.6] we describe, for each rank $\theta \geq 3$, a Yetter-Drinfeld module M of type γ_θ over a non-abelian group G . Assume there exist

- $\kappa \in Z(G)$ such that $\kappa \neq 1$, $\kappa^2 = 1$,
- $g_1, \dots, g_\theta \in G$ with $g_i^G = \{g_i, \kappa g_i\}$ for $i \in \mathbb{I}_{\theta-1}$ and $g_\theta^G = \{g_\theta\}$,
- $\chi_i \in \widehat{G^{g_i}}$ such that $\chi_i(g_i) = -1$,

satisfying (2.13), (2.14) and (2.15) for $i, j \in \mathbb{I}_{\theta-1}$, and

$$(2.19) \quad g_i g_\theta = g_\theta g_i, \quad \chi_i(g_\theta) \chi_\theta(g_i) = 1, \quad i < \theta - 1;$$

$$(2.20) \quad \chi_{\theta-1}(g_\theta) \chi_\theta(g_{\theta-1}) = -1.$$

Let $M_i = M(g_i^G, \chi_i) \in \frac{\mathbb{k}G}{\mathbb{k}G} \mathcal{YD}$. Then $M = \bigoplus_{i \in \mathbb{I}} M_i$ is of type γ_θ and $\dim \mathcal{B}(M) = 2^{2\theta^2 - \theta}$. Notice that $M_1 \oplus \dots \oplus M_{\theta-1}$ is of type $\alpha_{\theta-1}$ so, by §2.3.1, we have a basis $\{x_i, x_{\bar{i}}\}$ of M_i such that for $i, j \in \mathbb{I}_{\theta-1}$ the braiding c_{M_i, M_j} is determined by (2.16), (2.17) and (2.18). On the other hand $M_\theta = \mathbb{k}\{x_\theta\}$ is one-dimensional concentrated in degree $g_\theta \in Z(G)$. The braidings c_{M_θ, M_θ} , c_{M_i, M_θ} , c_{M_θ, M_i} , $i \in \mathbb{I}_{\theta-1}$, are determined by

$$(2.21) \quad c(x_\theta \otimes x_\theta) = -x_\theta \otimes x_\theta;$$

$$(2.22) \quad c(x_i \otimes x_\theta) = \chi_\theta(g_i) x_\theta \otimes x_i, \quad c(x_{\bar{i}} \otimes x_\theta) = \chi_\theta(\kappa g_i) x_\theta \otimes x_{\bar{i}};$$

$$(2.23) \quad c(x_\theta \otimes x_i) = \chi_i(g_\theta) x_i \otimes x_\theta, \quad c(x_\theta \otimes x_{\bar{i}}) = \chi_i(g_\theta) x_{\bar{i}} \otimes x_\theta.$$

2.3.3. *Type ϕ_4 .* Following [HV, Lemma 9.2] we describe a Yetter-Drinfeld module M over a non-abelian group G with Cartan matrix of type F_4 . Assume there exist

- $\kappa \in Z(G)$ such that $\kappa \neq 1$, $\kappa^2 = 1$,
- $g_1, \dots, g_4 \in G$ with $g_i^G = \{g_i, \kappa g_i\}$ for $i = 1, 2$ and $g_i^G = \{g_i\}$ for $i = 3, 4$,
- $\chi_i \in \widehat{G^{g_i}}$ such that $\chi_i(g_i) = -1$

satisfying the following:

$$(2.24) \quad \chi_1(g_3) \chi_3(g_1) = \chi_1(g_4) \chi_4(g_1) = \chi_2(g_4) \chi_4(g_2) = 1;$$

$$(2.25) \quad \chi_3(g_4) \chi_4(g_3) = \chi_2(g_3) \chi_3(g_2) = -1;$$

$$(2.26) \quad g_1 g_2 = \kappa g_2 g_1, \quad \chi_1(\kappa g_2^2) \chi_2(\kappa g_1^2) = 1.$$

Let $M_i = M(g_i^G, \chi_i)$. The Cartan matrix of $M = \bigoplus_{i \in \mathbb{I}} M_i$ is of type F_4 and $\dim \mathcal{B}(M) = 2^{36}$. The braidings c_{M_i, M_j} are given as in types α_θ , γ_θ , depending on i, j .

2.4. On the structure of the group G . Fix an abelian group Γ and $x, y \in \mathbb{Z}_2 \times \mathbb{Z}_2$ such that $\mathbb{Z}_2 \times \mathbb{Z}_2 = \langle x, y \rangle$. For every $u, v \in \Gamma$ and every $\kappa \in \Gamma$ such that $\kappa^2 = e$, there exists a 2-cocycle $\beta \in H^2(\mathbb{Z}_2 \times \mathbb{Z}_2, \Gamma)$ such that

$$\beta(x, x) = u, \quad \beta(x, y) = \kappa, \quad \beta(y, x) = e, \quad \beta(y, y) = v.$$

We denote by $\Gamma_{u,v,\kappa}$ the associated central extension of Γ by $\mathbb{Z}_2 \times \mathbb{Z}_2$:

$$(2.27) \quad \Gamma \hookrightarrow \Gamma_{u,v,\kappa} \twoheadrightarrow \mathbb{Z}_2 \times \mathbb{Z}_2 = \langle x, y \rangle,$$

where $[x, y] = \kappa$, $x^2 = u$, $y^2 = v$.

Next we describe some general features of a group G realizing the braidings described in §2.3.1, 2.3.2, 2.3.3. Let $M = \bigoplus_{i \in \mathbb{I}} M_i$ be of type α_θ , γ_θ , δ_θ , ϵ_θ , or ϕ_4 . Here $M_i = M(g_i^G, \chi_i) \in {}_{\mathbb{k}G}^{\mathbb{k}G} \mathcal{YD}$, hence $\text{supp } M = \bigcup_{i \in \mathbb{I}} g_i^G$. Note that

- There exists $\kappa \in Z(G)$ such that $\kappa^2 = e$ and $g_i^G = \{g_i, \kappa g_i\}$ for all i such that $g_i \notin Z(G)$.
- For i with $g_i^G = \{g_i, \kappa g_i\}$, there exists $j \neq i$ such that $g_i g_j = \kappa g_j g_i$; hence $g_j^G = \{g_j, \kappa g_j\}$.

The relevance of the central extensions constructed above is explained next.

Lemma 2.6. *Let $i \neq j \in \mathbb{I}$ be such that $g_i g_j = \kappa g_j g_i$, and let $N = G^{g_i} \cap G^{g_j}$. Then*

- The subgroup N is normal, and $G/N \simeq \mathbb{Z}_2 \times \mathbb{Z}_2$.*
- If $G = \langle \text{supp } M \rangle$, then N is an abelian subgroup, generated by the elements*

$$\begin{aligned} g_k, & \quad \text{for all } k \in \mathbb{I} \text{ such that } g_k g_i = g_i g_k, g_k g_j = g_j g_k, \\ g_k g_i, & \quad \text{for all } k \in \mathbb{I} \text{ such that } g_k g_i = g_i g_k, g_k g_j = \kappa g_j g_k, \text{ and} \\ g_k g_j, & \quad \text{for all } k \in \mathbb{I} \text{ such that } g_k g_j = g_j g_k, g_k g_i = \kappa g_i g_k. \end{aligned}$$

- If N is abelian, then $G \simeq N_{g_i^2, g_j^2, \kappa}$.*

Proof. For (a), we note first that $[G : G^{g_i}] = [G : G^{g_j}] = 2$, so both G^{g_i} and G^{g_j} are normal subgroups, and $[G : N_{ij}] = 4$ since $G = G_i G_j$. Let $g \in G$:

- If $g g_i g^{-1} = g_i$, $g g_j g^{-1} = g_j$, then $gN = N$.
- If $g g_i g^{-1} = g_i$, $g g_j g^{-1} = \kappa g_j$, then $gN = g_i N$.
- If $g g_i g^{-1} = \kappa g_i$, $g g_j g^{-1} = g_j$, then $gN = g_j N$.
- If $g g_i g^{-1} = \kappa g_i$, $g g_j g^{-1} = \kappa g_j$, then $gN = g_i g_j N$.

Hence $G/N = \{N, g_i N, g_j N, g_i g_j N\}$. Since $g_i^2, g_j^2 \in N$, we get $G/N \simeq \mathbb{Z}_2 \times \mathbb{Z}_2$.

(b) and (c) are straightforward. \square

A more detailed description of these groups is postponed to the Appendix.

2.4.1. *The parity vector.* An important invariant of $M = \bigoplus_{i \in \mathbb{I}} M_i \in {}_{\mathbb{k}G}^{\mathbb{k}G}\mathcal{YD}$ is

$$(2.28) \quad \mathbf{P} := (\chi_i(\kappa))_{1 \leq i \leq \theta} \in \{\pm 1\}^\theta.$$

If $\mathbf{P} = (1, \dots, 1)$, then the G -action on M factors to one of $G/\langle \kappa \rangle$. We show next that this happens in most of the cases, in which case M is a braided vector space of diagonal type.

Lemma 2.7. *Let M be either of type α_θ , $\theta \geq 4$, γ_θ , $\theta \geq 5$, δ_θ , $\theta \geq 5$ or ϵ_θ , $\theta = 6, 7, 8$. Then $\mathbf{P} = (1, \dots, 1)$. In other words, κ acts trivially on M .*

Proof. We consider first the case α_4 . By (2.14), $g_1, g_2 \in G^{g_4}$, and by (2.13), $\kappa = g_1 g_2 g_1^{-1} g_2^{-1}$; hence $\chi_4(\kappa) = 1$. By (2.15), $\chi_i(\kappa) = 1$ for all $i \in \mathbb{I}_4$.

Now the proof for α_θ , $\theta \geq 5$, δ_θ , $\theta \geq 5$ or ϵ_θ , $\theta = 6, 7, 8$ follows from taking into account all the submodules of type α_4 . The same fact says that, for type γ_θ , $\chi_i(\kappa) = 1$ for all $i \in \mathbb{I}_{\theta-1}$. Next we use that $g_{\theta-2}, g_{\theta-1} \in G^{g_\theta}$ and $\kappa = g_{\theta-1} g_{\theta-2} g_{\theta-1}^{-1} g_{\theta-2}^{-1}$ to deduce that $\chi_\theta(\kappa) = 1$. \square

3. FOLDINGS OF NICHOLS ALGEBRAS AND TRIVIALIZING THE ACTION OF THE CENTER

Motivated by Lemma 2.7, we pay special attention to Yetter-Drinfeld modules where κ acts trivially. We will show that these examples are related to diagonal braidings (of Cartan type) via the folding construction for Nichols algebras, developed by the second author in [L1, L2]. Then we show that for the other cases the action of the central element can be trivialized via a twist. First, we introduce basic notions needed for the folding construction.

3.1. Categorical action on Yetter-Drinfeld modules. Given a group G and a 2-cocycle $\sigma \in Z^2(G, \mathbb{k}^\times)$, we get a pairing $b_\sigma: G \times G \rightarrow \mathbb{k}^\times$ given by $b_\sigma(g, h) = \sigma(hgh^{-1}, h)\sigma^{-1}(h, g)$. Each 2-cocycle σ yields a tensor functor $F_\sigma: {}_{\mathbb{k}G}^{\mathbb{k}G}\mathcal{YD} \rightarrow {}_{\mathbb{k}G}^{\mathbb{k}G}\mathcal{YD}$ as follows:

- for an object M , let $F_\sigma(M)$ denote the same $\mathbb{k}G$ -comodule;
- the $\mathbb{k}G$ -module structure on $F_\sigma(M)$ is given by

$$g \cdot_\sigma m = b_\sigma(g, m_{-1})g \cdot m_0, \quad g \in G, m \in M;$$

- on Hom spaces F_σ is the identity, thus F_σ is \mathbb{k} -linear, faithful and exact;
- the monoidal structure $J_\sigma: F_\sigma(M) \otimes F_\sigma(N) \rightarrow F_\sigma(M \otimes N)$ is defined by

$$J_\sigma(m \otimes n) = \sigma(m_{-1}, n_{-1})m_0 \otimes n_0, \quad m \in M, n \in N.$$

We note that J_σ satisfy the hexagon axiom thanks to the cocycle condition on σ .

Given $V, W \in {}_{\mathbb{k}G}^{\mathbb{k}G}\mathcal{YD}$, one can see that $F_\sigma \otimes F_\sigma$ commutes with $c_{V, W}$ if and only if $\sigma(ghg^{-1}, g)\sigma(hgh^{-1}, h) = \sigma(g, h)\sigma(h, g)$ for all $g \in \text{supp } V$, $h \in \text{supp } W$. Hence, F_σ is braided if and only if that equality holds for all $g, h \in G$, which certainly happens if G is abelian. Anyhow, these functors patch together to an action by tensor autoequivalences of the group $Z^2(G, \mathbb{k}^\times)$ on ${}_{\mathbb{k}G}^{\mathbb{k}G}\mathcal{YD}$:

- the trivial cocycle acts as the identity, and
- $F_\eta F_\sigma = F_{\eta\sigma}$ for all $\eta, \sigma \in Z^2(G, \mathbb{k}^\times)$.

Given a Hopf algebra \mathcal{B} in $\mathbb{k}^G\mathcal{YD}$ we set, by abuse of notation,

$$(3.1) \quad \sigma : \mathcal{B}\#\mathbb{k}G \otimes \mathcal{B}\#\mathbb{k}G \rightarrow \mathbb{k}, \quad \sigma(x\#g \otimes y\#h) = \epsilon(x)\epsilon(y)\sigma(g, h).$$

Lemma 3.1. (i) *The map σ is a Hopf 2-cocycle for $\mathcal{B}\#\mathbb{k}G$.*

(ii) *For all $M \in \mathbb{k}^G\mathcal{YD}$, we have $(\mathcal{B}(M)\#\mathbb{k}G)_\sigma \simeq \mathcal{B}(F_\sigma(M))\#\mathbb{k}G$.*

Proof. (i) is clear. For (ii), apply [A+, 4.14 (a) & (b)]. \square

Here is the first notion towards the folding construction.

Definition 3.2. A *folding datum* is a triple (σ, M, \mathbf{u}) where $\sigma \in Z^2(\Gamma, \mathbb{k}^\times)$, $M \in \mathbb{k}^\Gamma\mathcal{YD}$ and $\mathbf{u} : F_\sigma(M) \rightarrow M$ is an isomorphism in $\mathbb{k}^\Gamma\mathcal{YD}$.¹

We will mainly deal with folding data coming from the following source.

Example 3.3. Fix $g_i \in \Gamma$, $\chi_i \in \widehat{\Gamma}$, $\sigma \in Z^2(\Gamma, \mathbb{k}^\times)$. Let $f : \mathbb{I} \rightarrow \mathbb{I}$ be a permutation such that

$$g_{f(i)} = g_i \quad \text{for all } i \in \mathbb{I}.$$

Consider $M = \bigoplus_{i \in \mathbb{I}} M(g_i, \chi_i) \in \mathbb{k}^\Gamma\mathcal{YD}$, and let $0 \neq x_i \in M(g_i, \chi_i)$. Note that M is of diagonal type with braiding matrix $\mathbf{q} = (q_{ij})_{i,j \in \mathbb{I}}$, $q_{ij} = \chi_j(g_i)$. Then the linear isomorphism $\mathbf{u} : F_\sigma(M) \rightarrow M$, $x_i \mapsto x_{f(i)}$ is in $\mathbb{k}^\Gamma\mathcal{YD}$ if and only if

$$\chi_{f(j)} = b_\sigma(-, g_j)\chi_j \quad \text{for all } j \in \mathbb{I}.$$

In this case we have that

$$q_{i,f(j)} = b_\sigma(g_i, g_j)q_{ij} \quad \text{for all } i, j \in \mathbb{I},$$

and f induces an automorphism of the Dynkin diagram of \mathfrak{q} because $b_\sigma(g_i, g_i) = 1$ and $b_\sigma(g_j, g_i)b_\sigma(g_i, g_j) = 1$ for all $i \neq j \in \mathbb{I}$.

Fixed M , the folding data form a group with unit $(1, M, \text{id})$ and product

$$(\sigma, M, \mathbf{u}) * (\sigma', M, \mathbf{u}') = (\sigma\sigma', M, \mathbf{u} \circ F_\sigma(\mathbf{u}')).$$

The next results are extracted from [L1, Part I].

Remark 3.4. (a) Let (σ, M, \mathbf{u}) be a folding datum, $H = \mathcal{B}(M)\#\mathbb{k}\Gamma$. By Lemma 3.1, \mathbf{u} induces a Hopf algebra isomorphism $\mathbf{u} : H_\sigma \rightarrow H$.

(b) The map $(1 \otimes \mathbf{u})\Delta_{H_\sigma} : H_\sigma \rightarrow H_\sigma \times H$ makes H_σ a right H -Galois object. Moreover ${}_\sigma H$ is an (H, H) -bi-Galois object.

(c) Given two folding data (σ, M, \mathbf{u}) , $(\sigma', M, \mathbf{u}')$, the map

$$(\text{id} \otimes \mathbf{u}')\Delta_{H_{\sigma\sigma'}} : H_{\sigma\sigma'} \rightarrow H_\sigma \square H_{\sigma'}$$

is an isomorphism of bi-Galois objects.

¹In other words, the pair (M, \mathbf{u}) is an object in the category $(\mathbb{k}^\Gamma\mathcal{YD})^\sigma$ of σ -equivariant objects.

- (d) The map in (c) determines a group homomorphism from the group of folding data over $M \in \frac{\mathbb{k}\Gamma}{\mathbb{k}\Gamma}\mathcal{YD}$ to the group of bi-Galois objects of $H = \mathcal{B}(M)\#\mathbb{k}\Gamma$.

3.2. Folding construction. Let $1 \rightarrow \Sigma \rightarrow G \rightarrow \Gamma \rightarrow 1$ be a central extension of a finite abelian group Γ by a finite abelian group Σ . Fix a set-theoretic section $s : \Gamma \rightarrow G$, and let

$$\tau \in Z^2(\Gamma, \Sigma), \quad \tau(g, h) = s(g)s(h)s(gh)^{-1}, \quad g, h \in \Gamma.$$

For each $t \in \widehat{\Sigma}$, we denote by $\sigma_t \in Z^2(\Gamma, \mathbb{k}^\times)$ the 2-cocycle $\sigma_t = t \circ \tau$. The assignment $t \mapsto \sigma_t$ is a group homomorphism $\widehat{\Sigma} \rightarrow Z^2(\Gamma, \mathbb{k}^\times)$. Now we fix

- a Yetter-Drinfeld module M over $\mathbb{k}\Gamma$,
- isomorphisms $\mathbf{u}_t : F_{\sigma_t}(M) \rightarrow M$, $t \in \widehat{\Sigma}$, in $\frac{\mathbb{k}\Gamma}{\mathbb{k}\Gamma}\mathcal{YD}$ such that the map $t \mapsto (\sigma_t, M, \mathbf{u}_t)$ is a group homomorphism from $\widehat{\Sigma}$ to the group of folding data for M . In particular, we have $\mathbf{u}_0 = \text{id} : {}_{\sigma_0}H = H \rightarrow H$.

Remark 3.5. M becomes a $\widehat{\Sigma}$ -module, where $t \in \widehat{\Sigma}$ acts by the automorphism \mathbf{u}_t . As Σ is finite abelian, the $\widehat{\Sigma}$ -action diagonalizes and M decomposes as a direct sum of Σ -eigenspaces:

$$M = \bigoplus_{p \in \Sigma} M^p, \quad M^p = \{m \in M : \mathbf{u}_t(m) = t(p)m \text{ for all } t \in \widehat{\Sigma}\}.$$

Theorem 3.6. [L2, Theorem 3.6] *Let Γ , G and M as above. The following structure define a $\mathbb{k}G$ -Yetter-Drinfeld module \widetilde{M} :*

- as a vector space, $\widetilde{M} = M$,
- the G -action is obtained by pulling back the Γ -action (hence Σ acts trivially),
- the G -grading is given by $\widetilde{M}_g = M_g^{gs(g)^{-1}} := M_g \cap M^{gs(g)^{-1}}$, for each $g \in G$.

Also, as a braided vector space, $\widetilde{M} = M$. □

Next we introduce the folding construction, which produces a Nichols algebra over G starting from folding data on a Nichols algebra over Γ . The procedure gives a central extension of Hopf algebras and is related with the *Fourier transform* developed in [AG].

Theorem 3.7. *Let $H := \mathcal{B}(M)\#\mathbb{k}\Gamma$. There exists a Hopf algebra structure on $\widetilde{H} := \bigoplus_{t \in \widehat{\Sigma}} H_{\sigma_t}$ given by*

$$\Delta|_{\sigma_t H} = \bigoplus_{t't''=t} (\text{id} \otimes \mathbf{u}_{t''}) \Delta_H : H_{\sigma_t} \rightarrow \bigoplus_{t't''=t} H_{\sigma_{t'}} \otimes H_{\sigma_{t''}}; \quad \epsilon|_H = \epsilon_H, \quad \epsilon|_{H_{\sigma_t}} = 0, t \neq 0.$$

Moreover, $\widetilde{H} \cong \mathcal{B}(\widetilde{M})\#\mathbb{k}G$ as Hopf algebras.

Remark 3.8. By the results above, the group of folding data induces a homomorphism of 2-groups $\widehat{\Sigma} \rightarrow \text{BiGal}(H)$. This in turn defines a homomorphism of 2-groups $\widehat{\Sigma} \rightarrow \text{BrPic}(\text{Rep}(H))$ by [Mm] and thus defines by [EGNO] a Σ -extension of the tensor category $\text{Rep}(H)$. This tensor category coincides with $\text{Rep}(\widetilde{H})$.

A way to see this fact comes from the equivariantization process applied to Hopf algebras since the folding data gives a functor from (the category defined from) Σ to the Drinfeld double of $\text{Rep}(H)$. Reciprocally, $\text{Rep}(H)$ is the de-equivariantization of $\text{Rep}(\tilde{H})$ associated to a central extension of Hopf algebras [AnGP], see also [L1, Theorem 3.6].

3.3. Folding data for trivial action of κ . Next we realize most of the examples of types $\alpha_\theta, \gamma_\theta, \delta_\theta, \epsilon_\theta$ and ϕ_4 as foldings of braided vector spaces of diagonal type. In all cases we can proceed as in Example 3.3 with $\Sigma = \mathbb{Z}_2 = \{e, \kappa\}$.

Example 3.9. Fix a finite Cartan matrix $\mathbf{a} = (a_{ij})_{i,j \in \mathbb{I}}$ with simply-laced Dynkin diagram. Assume that Γ is a finite abelian group generated by $g_i, i \in \mathbb{I}$, which admits a 2-cocycle

$$(3.2) \quad \tau \in Z^2(\Gamma, \Sigma) \text{ such that} \quad \tau(g_i, g_j) = \begin{cases} \kappa^{a_{ij}}, & i < j \in \mathbb{I}; \\ e & i \geq j \in \mathbb{I}. \end{cases}$$

Let G be the extension of Γ by Σ associated to τ . Thus G is generated by g_i and κ ; in G we have $g_i g_j = \kappa^{a_{ij}} g_j g_i$ for $i \neq j \in \mathbb{I}$. Assume further that we have $\chi_i \in \widehat{\Gamma}, i \in \mathbb{I}$, satisfying

$$\chi_i(g_i) = -1, \quad \chi_i(g_j) \chi_j(g_i) = (-1)^{a_{ij}}, \quad i \neq j \in \mathbb{I}.$$

Then $V = \bigoplus_{i \in \mathbb{I}} \mathbb{k}^{X_i}$ is of Cartan type \mathbf{a} with $q = -1$, as in Example 2.2.

Here, $\widehat{\Sigma} = \{e, \mathbf{t}\}$, with $\mathbf{t}(\kappa) = -1$. Set $\sigma := \mathbf{t} \circ \tau$, and

$$g_{i+\theta} = g_i \quad \chi_{i+\theta} := b_\sigma(-, g_i) \chi_i \in \widehat{\Gamma}, \quad i \in \mathbb{I}, \quad M := \bigoplus_{i \in \mathbb{I}_{2\theta}} \mathbb{k}^{X_i}.$$

Then $M = V \oplus F_\sigma(V)$, and is of Cartan type with Cartan matrix $\tilde{\mathbf{a}} := \begin{pmatrix} \mathbf{a} & 0 \\ 0 & \mathbf{a} \end{pmatrix}$.

Set also $f : \mathbb{I}_{2\theta} \rightarrow \mathbb{I}_{2\theta}, i \mapsto i + \theta$ modulo 2θ . Then $\mathbf{u} : F_\sigma(M) \rightarrow M$ as in Example 3.3 is a folding datum, and the map from $\widehat{\Sigma}$ to the group of folding data such that $\mathbf{t} \mapsto (\sigma, M, \mathbf{u})$ is a group homomorphism. Following [L2], if \mathbf{a} is of type $X_\theta \in \{A_\theta | \theta \geq 2\} \cup \{D_\theta | \theta \geq 4\} \cup \{E_\theta | \theta = 6, 7, 8\}$, we use ${}^2X_\theta^2$ to denote the corresponding folding of $X_\theta \times X_\theta$ by f, \mathbf{u} as above.

Example 3.10. Fix a finite abelian group Γ generated by $g_i, i \in \mathbb{I}_4$, which admits

$$(3.3) \quad \tau \in Z^2(\Gamma, \Sigma) \text{ such that} \quad \tau(g_i, g_j) = \kappa^{\delta_{i3}\delta_{j4}}, \quad i, j \in \mathbb{I}_4.$$

Let G be the extension of Γ by Σ associated to τ . Now G is generated by g_i and κ ; the relations $g_3 g_4 = \kappa g_4 g_3$, and $g_i g_j = g_j g_i$ for $\{i, j\} \neq \{3, 4\}$ hold in G .

Assume further that $\chi_i \in \widehat{\Gamma}, i \in \mathbb{I}$, satisfy

$$\chi_i(g_i) = -1, \quad i \in \mathbb{I}_4; \quad \chi_i(g_j) \chi_j(g_i) = (-1)^{\delta_{i+1,j}}, \quad i < j \in \mathbb{I}_4.$$

Again, fix $\mathbf{t} \in \widehat{\Sigma}$ such that $\mathbf{t}(\kappa) = -1$. Set $\sigma := \mathbf{t} \circ \tau$, and

$$g_{i+2} = g_i \quad \chi_{i+2} := b_\sigma(-, g_i) \chi_i \in \widehat{\Gamma}, \quad i \in \{3, 4\}, \quad M := \bigoplus_{i \in \mathbb{I}_6} \mathbb{k}^{X_i}.$$

Then M is of Cartan type E_6 . Let $f : \mathbb{I}_6 \rightarrow \mathbb{I}_6$ be the bijection that exchanges $3 \leftrightarrow 5$ and $4 \leftrightarrow 6$. Then $\mathbf{u} : F_\sigma(M) \rightarrow M$ as in Example 3.3 is a folding datum, and the map $\mathbf{t} \mapsto (\sigma, M, \mathbf{u})$ is a group homomorphism from $\widehat{\Sigma}$ to the group of folding data. Following [L2], 2E_6 denotes a folding as above.

Example 3.11. Fix a finite abelian group Γ generated by $g_i, i \in \mathbb{I}$, which admits a 2-cocycle

$$(3.4) \quad \tau \in Z^2(\Gamma, \Sigma) \text{ such that} \quad \tau(g_i, g_j) = \begin{cases} \kappa, & j = i + 1 < \theta, \\ e, & \text{otherwise.} \end{cases}$$

Let G be the extension of Γ by Σ associated to τ . Thus G is generated by g_i and κ ; in G , we have the relations $g_i g_{i+1} = \kappa g_{i+1} g_i$ if $i < \theta - 1$, and $g_i g_j = g_j g_i$ otherwise.

Assume further that $\chi_i \in \widehat{\Gamma}$, $i \in \mathbb{I}$, satisfy

$$\chi_i(g_i) = -1, \quad i \in \mathbb{I}; \quad \chi_i(g_j) \chi_j(g_i) = (-1)^{\delta_{i+1,j}}, \quad i < j \in \mathbb{I}.$$

Again, fix $\mathbf{t} \in \widehat{\Sigma}$ such that $\mathbf{t}(\kappa) = -1$. Set $\sigma := \mathbf{t} \circ \tau$, and

$$g_{2\theta-i} = g_i \quad \chi_{2\theta-i} := b_\sigma(-, g_i) \chi_i \in \widehat{\Gamma}, \quad i \in \mathbb{I}_{\theta-1}, \quad M := \bigoplus_{i \in \mathbb{I}_{2\theta-1}} \mathbb{k}^{\chi_i}.$$

Then M is of Cartan type $A_{2\theta-1}$. Let $f : \mathbb{I}_{2\theta-1} \rightarrow \mathbb{I}_{2\theta-1}$, $f(i) = 2\theta - i$. Then $\mathbf{u} : F_\sigma(M) \rightarrow M$ as in Example 3.3 is a folding datum, and the map $\mathbf{t} \mapsto (\sigma, M, \mathbf{u})$ is a group homomorphism from $\widehat{\Sigma}$ to the group of folding data. Following [L2], ${}^2A_{2n-1}$ denotes the folding above.

Remark 3.12. In the three examples above, we can take $\Gamma = \mathbb{Z}_2^{\mathbb{I}}$, see [L2, §5].

Theorem 3.13. *Let M as in §2.3.1, §2.3.2 or §2.3.3. Assume that κ acts trivially on M .*

- (a) *If M is of type $\alpha_\theta, \delta_\theta$ or ϵ_θ , then $\mathcal{B}(M)$ is a folded Nichols algebra as in Example 3.9.*
- (b) *If M is of type ϕ_4 , then $\mathcal{B}(M)$ is a folded Nichols algebra as in Example 3.10.*
- (c) *If M is of type γ_θ , then $\mathcal{B}(M)$ is a folded Nichols algebra as in Example 3.11.*

Proof. Since κ acts trivially, this follows by [L2, Theorems 5.6, 5.7, 5.8]. \square

Remark 3.14. Fix a braided vector space M of type either $\alpha_\theta, \theta \geq 4, \delta_\theta, \theta \geq 5$, or $\epsilon_\theta, \theta = 6, 7, 8$ (type γ_θ was already considered in general), with Cartan matrix $\mathbf{a} = (a_{ij})_{i,j \in \mathbb{I}_\theta}$. By Lemma 2.7, $\chi_i(\kappa) = 1$ for all $i \in \mathbb{I}$, so M is of diagonal type. We exhibit a basis in which the braiding is of diagonal type, and we give the braiding matrix.

- (I) Set $q_{ii} = -1$ for all $i \in \mathbb{I}_\theta$, and $q_{ij} = \chi_j(g_i)$ if $a_{ij} = 0$.
- (II) Let $1 \leq i < j \leq \theta$ be such that $a_{ij} = -1$. Let $q_{ij} \in \mathbb{k}^\times$ be such that $q_{ij}^2 = \chi_j(g_i^2)$, and set $q_{ji} := -q_{ij}^{-1}$. By Step 4 of Proposition 5.2, if $k > i$ also satisfies that $a_{ik} = -1$ we may choose $q_{ik} = q_{ij}$.

(III) We also set $\mathfrak{q} = (q_{ij})_{i,j \in \mathbb{I}_{2\theta}}$, where

$$q_{ij} = \begin{cases} -1, & i \leq \theta < j \text{ or } j \leq \theta < i; \\ -q_{i-\theta, j-\theta}, & i, j > \theta, a_{i-\theta, j-\theta} = -1; \\ q_{i-\theta, j-\theta}, & i, j > \theta, a_{i-\theta, j-\theta} = 0; \\ -1, & i = j > \theta. \end{cases}$$

Given $i \in \mathbb{I}_\theta$, there is $j \neq i$ in \mathbb{I}_θ such that $a_{ij} \neq 0$. If possible, take $j > i$ such that $a_{ij} \neq 0$; otherwise take $j < i$ with $a_{ij} \neq 0$. By (II) above, we can define

$$\mathbf{x}_i := x_i + q_{ij}x_{\bar{j}}, \quad \mathbf{x}_{\bar{i}} := x_i - q_{ij}x_{\bar{j}}, \quad i \in \mathbb{I}.$$

Using (2.16), (2.17) and (2.18) we verify that

$$c(\mathbf{x}_i \otimes \mathbf{x}_j) = \begin{cases} -1, & i = j; \\ q_{ij}, & i \neq j. \end{cases} \quad c(\mathbf{x}_{\bar{i}} \otimes \mathbf{x}_{\bar{j}}) = \begin{cases} -1, & i = j \\ -q_{ij}, & a_{ij} = -1; \\ q_{ij}, & a_{ij} = 0. \end{cases}$$

$$c(\mathbf{x}_i \otimes \mathbf{x}_{\bar{j}}) = \begin{cases} -1, & i = j \\ -q_{ij}, & a_{ij} = -1; \\ q_{ij}, & a_{ij} = 0. \end{cases} \quad c(\mathbf{x}_{\bar{i}} \otimes \mathbf{x}_j) = \begin{cases} -1, & i = j \\ q_{ij}, & a_{ij} = -1; \\ q_{ij}, & a_{ij} = 0. \end{cases}$$

so the braiding matrix of M is \mathfrak{q} , and M is, respectively, of type $A_\theta \times A_\theta$, $D_\theta \times D_\theta$ or $E_\theta \times E_\theta$, both copies with parameter $q = -1$.

3.4. Trivializing the action of κ via a twist. Retain the notation introduced in §3.3. Thus G is a non-abelian group and $M = \bigoplus_{i \in \mathbb{I}} M_i \in \mathbb{k}_G^G \mathcal{YD}$, where $M_i = M(g_i^G, \chi_i)$.

For the cases not covered by Lemma 2.7, we will show the existence of a 2-cocycle σ such that κ acts trivially on $F_\sigma(M)$. Recall the parity vector $\mathbf{P} = (\chi_1(\kappa), \dots, \chi_\theta(\kappa))$ from (2.28).

Let $\sigma \in H^2(G, \mathbb{k}^\times)$. Following §3.1, the *twisted Yetter-Drinfeld module* associated to σ is $F_\sigma(M) = M^\sigma = \bigoplus_{i \in \mathbb{I}} M(g_i, \chi_i^\sigma)$, where $\chi_i^\sigma \in \widehat{G}^{g_i}$ is given by the following formula:

$$\chi_i^\sigma(h) = \sigma(hg_ih^{-1}, h)\sigma^{-1}(h, g_i)\chi_i(h), \quad h \in G.$$

Since κ is central in G , we have $\chi_i^\sigma(\kappa) = \sigma(g_i, \kappa)\sigma^{-1}(\kappa, g_i)\chi_i(\kappa)$.

Proposition 3.15. *Let G be a non-abelian group, $M \in \mathbb{k}_G^G \mathcal{YD}$ of type $\alpha_2, \alpha_3, \delta_4, \gamma_3, \gamma_4$ or ϕ_4 such that $\text{supp } M$ generates G . There exists $\sigma \in H^2(G, \mathbb{k}^\times)$ with $\chi_i^\sigma(\kappa) = 1$ for all i .*

Let us outline the strategy that will be used in the Appendix to prove this statement.

- (i) We go through the cases and list the possible $\mathbf{P} = (\chi_i(\kappa))_i \in \{\pm 1\}^n$.
- (ii) It is sufficient to consider one \mathbf{P} representing each Weyl groupoid orbit. The i -th reflection of M is

$$\rho_i M = \bigoplus_{j \in \mathbb{I}} M(g_j g_i^{-c_{ij}}, \chi_j \chi_i^{-c_{ij}}),$$

and the parity vector of $\rho_i M$ is $P' = (\chi_j(\kappa)\chi_i(\kappa)^{-c_{ij}})_{j \in \mathbb{I}}$.

- (iii) Next we introduce an auxiliary *minimal* group G^{\min} . Namely $G^{\min} \subseteq \text{End } M$ is generated by (the action of) g_i . The definition of G^{\min} depends only on the scalars $\chi_i(g_j), \chi_i(\kappa)$. It is enough to prove Proposition 3.15 for this group, since the asserted 2-cocycle σ on G^{\min} can be pulled back to G .
- (iv) In the next steps we show case-wise that there exists $\sigma \in H^2(G^{\min}, \mathbb{k}^\times)$ such that

$$\sigma(g_i, \kappa)\sigma(\kappa, g_i)^{-1} = \chi_i(\kappa), \quad \text{for all } i \in \mathbb{I}.$$
- (v) For type α_2 there are two Weyl groupoid orbits for $P = (\chi_1(\kappa), \chi_2(\kappa))$, namely $\{(1, 1)\}$ and $\{(1, -1), (-1, -1), (-1, 1)\}$, the first one corresponding to P trivial. For $(-1, 1)$ we get three different types of groups according to the order of $\chi_2(g_1^2)$: we find the desired cocycle using semidirect product decompositions.
- (vi) The cases α_3 and γ_3 are treated using spectral sequences arguments for a central extension. Necessary information about the structure of the group (minimal orders of central elements for example) enters conveniently via the existence of a 1-dimensional representation, constructed from the structure of M .
- (vii) The remaining cases are treated using two simultaneous extensions.

We postpone the proof until §A.3 since we need technical results on group cohomology.

By assumption G is a central extension of an abelian group Γ by $\langle \kappa \rangle \simeq \mathbb{Z}_2$, say:

$$1 \longrightarrow \mathbb{Z}_2 \longrightarrow G \longrightarrow \Gamma \longrightarrow 1.$$

To illustrate the proof of Proposition 3.15, we give an example where such central extensions are related to symplectic forms on \mathbb{Z}_2^θ .

Example 3.16. Let $\Gamma = \mathbb{Z}_2^\theta$ with generators $g_i, i \in \mathbb{I}$. The commutator in G defines a symplectic form on Γ such that the radical \mathbb{Z}_2^r is the image of the center $Z(G)$; in particular g_i, g_j commute iff the symplectic form on them is zero; hence the size of the conjugacy class of g_i is 1 or 2 depending if g_i is in the radical or not. This symplectic form of type (θ, r) is uniquely determined.

Central extensions $\mathbb{Z}_2 \rightarrow G \rightarrow \Gamma$ are classified by quadratic forms with fixed symplectic form. For type α_2 we have $(\theta, r) = (2, 0)$ and there are two types of central extensions of order 8 with this commutator structure, namely the extraspecial groups 2_-^{2+1} (the quaternion group) and 2_+^{2+1} (the dihedral group). The group 2_-^{2+1} has defining relations $g_1^2 = g_2^2 = \kappa$ and trivial second cohomology $H^2(2_-^{2+1}, \mathbb{k}^\times)$, while the group 2_+^{2+1} has defining relations $g_1^2 = e, g_2^2 = \kappa$ (depending on a choice of generators) and non-trivial second cohomology $H^2(2_+^{2+1}, \mathbb{k}^\times) = \{1, \sigma\}$. Our proof works for the group 2_-^{2+1} because the relations (2.13) defining M imply

$$\chi_1(\kappa) = \chi_1(g_1^2) = 1, \quad \chi_2(\kappa) = \chi_2(g_2^2) = 1.$$

For 2_+^{2+1} , both $P = (1, 1)$ and $P = (-1, 1)$ are possible and we have a 2-cocycle σ with

$$\sigma(g_1, \kappa)\sigma^{-1}(\kappa, g_1) = -1, \quad \sigma(g_2, \kappa)\sigma^{-1}(\kappa, g_2) = 1.$$

The other choices of generators for 2_+^{2+1} work similarly and also follow from the first choice by using Weyl groupoid reflections.

The cases $\alpha_3, \delta_4, \gamma_3, \gamma_4, \phi_4$ present similar behavior because $\frac{1}{2}(\theta - r) = 1$ in all of them. For each case there is an underlying extraspecial group 2_{\pm}^{2+1} .

Remark 3.17. The proof becomes more involved for an arbitrary abelian group Γ because there are many central extensions by \mathbb{Z}_2 , parametrized by the powers of the generators, and the existence of a non-trivial group cohomology is very sensitive to these choices.

We are ready to state the main result of this section, which states that each Nichols algebras of type $\alpha_\theta, \gamma_\theta, \delta_\theta, \epsilon_\theta$, or ϕ_4 is a twist of the corresponding Nichols algebra of diagonal type as in §3.3.

Theorem 3.18. *Let G be a finite non-abelian group, $M \in \mathbb{k}_G^G \mathcal{YD}$ of type either $\alpha_\theta, \gamma_\theta, \delta_\theta, \epsilon_\theta$ or ϕ_4 whose support generates G . Then there exists $\sigma \in H^2(G, \mathbb{k})$ such that $F_\sigma(M)$ is of diagonal type.*

Proof. If M is of type $\alpha_\theta, \theta \geq 4, \gamma_\theta, \theta \geq 5, \delta_\theta, \theta \geq 5$ or $\epsilon_\theta, \theta = 6, 7, 8$, then κ acts trivially by Lemma 2.7, so M is naturally a Yetter-Drinfeld module over $\Gamma = G/\langle \kappa \rangle$, a finite abelian group, thus M is itself of diagonal type. For the other cases, we apply Proposition 3.15. \square

Let G and $M = \bigoplus_{i \in \mathbb{I}} M_i \in \mathbb{k}_G^G \mathcal{YD}$ as above. Consider

$$(3.5) \quad \ell := \begin{cases} 2 & \text{for type } \phi_4, \\ \theta - 1 & \text{for type } \gamma_\theta, \\ \theta & \text{otherwise.} \end{cases}$$

Keeping the notation used for §2.3.1, 2.3.2, 2.3.3, we have

- If $i \leq \ell$, then $\dim M_i = 2, g_i^G = \{g_i, \kappa g_i\}$, where $\kappa \in Z(G)$ satisfies $\kappa^2 = 1$. We fix a basis $x_i, x_{\bar{i}}$ of M_i as above.
- If $i > \ell$, then $\dim M_i = 1, g_i^G = \{g_i\}$. For a basis, we fix any nonzero element x_i in M_i .

Remark 3.19. Let \mathbf{G} be the group generated by $g_i, i \in \mathbb{I}$ and κ with relations

$$g_i \kappa = \kappa g_i, \quad \kappa^2 = e, \quad g_i g_j = \begin{cases} \kappa g_j g_i, & i \neq j \leq \ell, a_{ij} = -1, \\ g_j g_i & \text{otherwise,} \end{cases}$$

where $\mathbf{a} = (a_{ij})$ denotes the Cartan matrix of M . Then \mathbf{G} is a central extension of $\mathbb{Z}^{\mathbb{I}}$ by \mathbb{Z}_2 , and the braided vector space M has a realization over \mathbf{G} . Moreover, the subgroup of G generated by $\text{supp } M$ is a quotient of \mathbf{G} .

The $\mathbb{Z}^{\mathbb{I}}$ -grading on $T(M)$ and its homogeneous quotients (in particular, $\mathcal{B}(M)$) is given by the induced coaction of $\mathbb{k}\mathbb{Z}^{\mathbb{I}} \simeq \mathbb{k}\mathbf{G}/\langle \kappa \rangle$.

4. GENERATION IN DEGREE ONE

Using Theorem 3.18 and generation-in-degree-one for the diagonal setting [An2], we get

Theorem 4.1. *Let H be a finite-dimensional pointed Hopf algebra with infinitesimal braiding M of type $\alpha_\theta, \gamma_\theta, \delta_\theta, \epsilon_\theta$ or ϕ_4 . Then*

$$\text{gr } H \simeq \mathcal{B}(M) \# \mathbb{k}G(H).$$

In other words, H is generated by skew-primitive and group-like elements.

Proof. Let R be the diagram of H , so $\text{gr } H \simeq R \# \mathbb{k}G(H)$. Put also $\mathcal{B} := R^*$. Notice that $W := M^*$ is of the same type as M ; we fix $g_i, i \in \mathbb{I}, \kappa$ as in §2.2. Let G be the subgroup of $G(H)$ generated by $g_i, i \in \mathbb{I}$. Then $\mathcal{B} \in {}^{\mathbb{k}G} \mathcal{YD}$ is a finite-dimensional pre-Nichols algebra of W ; i.e. $\mathcal{B} = \bigoplus_{n \geq 0} \mathcal{B}^n$, where $\mathcal{B}^n = R(n)^*$, is a graded Hopf algebra such that $\mathcal{B}^0 = \mathbb{k}1$ and is generated as an algebra by $W = \mathcal{B}^1$.

For $\sigma \in H^2(G, \mathbb{k}^\times)$ as in Theorem 3.18, set $\sigma : \mathcal{B} \# \mathbb{k}G \otimes \mathcal{B} \# \mathbb{k}G \rightarrow \mathbb{k}$ as in (3.1), and let $\mathbb{H} := (\mathcal{B} \# \mathbb{k}G)_\sigma, \mathbb{H}^n := \mathcal{B}^n \# \mathbb{k}G$. Then $\mathbb{H} = \bigoplus_{n \geq 0} \mathbb{H}^n$ is a graded coalgebra, because twisting by σ leaves the coalgebra structure unchanged. Thus \mathbb{H} is pointed with coradical $\mathbb{H}^0 \simeq \mathbb{k}G$ by [Mo, 5.3.4]. As σ is trivial in degree > 0 , \mathbb{H} is a graded Hopf algebra: By [A+, 4.14 (a)], $\mathbb{H} \simeq \mathcal{B}' \# \mathbb{k}G$, where $\mathcal{B}' \in {}^{\mathbb{k}G} \mathcal{YD}$ is a pre-Nichols algebra of $F_\sigma(W)$.

As $\kappa \in Z(G)$ and it acts trivially on W^σ , we have that $\kappa \in Z(H)$. Set $Q := \mathbb{H}/\mathbb{H}(\kappa - 1), \Gamma = G/\langle \kappa \rangle$. The G -actions on $F_\sigma(W)$ and on \mathcal{B}' induce respective Γ -actions on them. Also, $F_\sigma(W)$ and \mathcal{B}' become $\mathbb{k}\Gamma$ -comodules via $\pi : G \twoheadrightarrow \Gamma$. Moreover, with these structures both \mathcal{B}' and $F_\sigma(W)$ are ${}^{\mathbb{k}\Gamma} \mathcal{YD}$, and $\mathcal{B}' \in {}^{\mathbb{k}\Gamma} \mathcal{YD}$ is a pre-Nichols algebra of $F_\sigma(W)$. We identify $\mathbb{H} \simeq \mathcal{B}' \# \mathbb{k}G$ and consider the map $\Phi : \mathbb{H} \rightarrow \mathcal{B}' \# \mathbb{k}\Gamma, \Phi(x \# g) = x \# \pi(g)$. Then Φ is a surjective Hopf algebra map such that $\Phi(\kappa - 1) = 0$, hence Φ induces a surjective Hopf algebra map $\phi : Q \rightarrow \mathcal{B}' \# \mathbb{k}\Gamma$. As \mathbb{H} is a central extension of Q by $\mathbb{k}\mathbb{Z}_2$ (since $\kappa^2 = 1, \kappa \neq 1$),

$$\dim Q = \frac{1}{2} \dim \mathbb{H} = \frac{1}{2} \dim \mathcal{B}'|G| = \dim \mathcal{B}'|\Gamma|,$$

so ϕ is an isomorphism.

Now Γ is an abelian group since for each $i \neq j$ either $g_i g_j = g_j g_i$ or $g_i g_j = \kappa g_j g_i$ in G , and Γ is generated by the images of the g_i 's (see §2.2). Hence [An2] implies that $\mathcal{B}' = \mathcal{B}(F_\sigma(W))$, so $\mathcal{B} = \mathcal{B}(W)$ by [A+, 4.14 (b)]. Dualizing, $R = \mathcal{B}(M)$, as desired. \square

5. GENERATORS AND RELATIONS FOR NICHOLS ALGEBRAS

In this section we exhibit a presentation by generators and relations for the Nichols algebras of the Yetter-Drinfeld modules $M = \bigoplus_{i \in \mathbb{I}} M_i \in {}^{\mathbb{k}G} \mathcal{YD}$ as in §2.3.1, 2.3.2, 2.3.3, which are standard, i.e. all reflection $\rho_i M$ are of the same type as M . In particular, the root system Δ^M is a classical one. We may assume that G is the group in Remark 3.19.

5.1. **Types α_2 and α_3 .** We give a presentation of Nichols algebras of types α_2 and α_3 . They will be a key step toward the presentation for the general case, since all relations which are not powers of root vectors are supported on smaller submodules of these types.

5.1.1. *Type α_2 .* Let G be a group, $e \neq \kappa$, $g_1, g_2 \in G$ such that

$$g_1 g_2 = \kappa g_2 g_1, \quad \kappa^2 = e, \quad g_i^G = \{g_i, \kappa g_i\}, \quad i = 1, 2.$$

Following [HS1], the subgroup of G generated by g_1 and g_2 is a quotient of

$$(5.1) \quad G_2 = \langle g_1, g_2, \kappa \mid \kappa^2 = 1, \kappa g_1 = g_1 \kappa, \kappa g_2 = g_2 \kappa, g_2 g_1 = \kappa g_1 g_2 \rangle.$$

Assume there are $\chi_i \in \widehat{G^{g_i}}$ such that $\chi_i(g_i) = -1$ and $\chi_1(\kappa g_2^2) \chi_2(\kappa g_1^2) = 1$. For $i \in \mathbb{I}_2$ set $M_i = M(g_i^G, \chi_i) \in \mathbb{k}_G^G \mathcal{YD}$, thus $M = M_1 \oplus M_2$ is of type α_2 by [HS1, Theorem 4.6].

Proposition 5.1. *Let $M = M_1 \oplus M_2 \in \mathbb{k}_G^G \mathcal{YD}$ of type α_2 as above. Then $\mathcal{B}(M)$ is presented by generators $x_1, x_{\overline{1}}, x_2, x_{\overline{2}}$ and relations*

$$(5.2) \quad x_i^2 = x_{\overline{i}}^2 = 0, \quad (\text{ad}_c x_i) x_{\overline{i}} = 0, \quad i \in \mathbb{I}.$$

$$(5.3) \quad x_{1\overline{2}} = -\chi_2(g_1^2) x_{\overline{1}\overline{2}}, \quad x_{\overline{1}\overline{2}} = -\chi_1(\kappa) x_{12};$$

$$(5.4) \quad x_{12}^2 = 0, \quad [x_{12}, x_{\overline{1}\overline{2}}]_c = 0.$$

The following set is a PBW basis of $\mathcal{B}(M)$:

$$(5.5) \quad \{x_2^a x_{\overline{2}}^b x_{12}^c x_{\overline{1}\overline{2}}^d x_1^e x_{\overline{1}}^f : a, b, c, d, e, f \in \{0, 1\}\}.$$

Proof. We proceed in several steps.

Step 1. By (2.16) the Nichols algebra of M_i for $i = 1, 2$ is a quantum linear space, and (5.6) implies that relations (5.2) hold in $\mathcal{B}(M)$.

Step 2. The inclusion $M_{12} := (\text{ad}_c M_1) M_2 \hookrightarrow \mathcal{B}(M)$ extends to a \mathbb{Z}^2 -graded algebra inclusion $\mathcal{B}(M_{12}) \hookrightarrow \mathcal{B}(M)$ and the multiplication

$$(5.6) \quad \mathcal{B}(M_2) \otimes \mathcal{B}(M_{12}) \otimes \mathcal{B}(M_1) \longrightarrow \mathcal{B}(M)$$

is an isomorphism of \mathbb{Z}^2 -graded objects in $\mathbb{k}_G^G \mathcal{YD}$, where M_1 sits in degree α_1 , M_2 in degree α_2 , and M_{12} in degree $\alpha_1 + \alpha_2$.

This follows by [HS1, Theorem 4.6]. In order to find defining relations for $\mathcal{B}(M_{12}) \subset \mathcal{B}(M)$, we need a more explicit description of the structure of M_{12} .

Step 3. (a) The set $\{x_{12}, x_{\overline{1}\overline{2}}\}$ is a basis of M_{12} , and the braiding in this basis is of diagonal type with matrix

$$(5.7) \quad \begin{pmatrix} -1 & -\chi_1(\kappa) \chi_2(\kappa) \\ -\chi_1(\kappa) \chi_2(\kappa) & -1 \end{pmatrix}.$$

(b) Relations (5.3) and (5.4) hold in $\mathcal{B}(M)$.

Proof. Note that $\partial_{x_2}(x_{12}) = x_1$ and $\partial_{x_2}(x_{\overline{12}}) = x_{\overline{1}}$, thus x_{12} and $x_{\overline{12}}$ are linearly independent in $\mathcal{B}(M)$. Relations (5.3) are verified using the skew-derivations of $T(M_1 \oplus M_2)$. The braiding of M_{12} is obtained from a straight-forward computation in ${}_{\mathbb{k}G}^{\mathbb{k}G}\mathcal{YD}$. Now (5.4) follows from (5.6) and (5.7). \square

Note that $\mathcal{B}(M_{12})$ is presented by the relations (5.4) and $(x_{\overline{12}})^2 = 0$, which has not been included above since it can be deduced from the previous ones, as we show in Step 4.

With (5.6) in mind, the next step towards exhibiting a presentation of $\mathcal{B}(M)$ should be to find braided commutations between $\mathcal{B}(M_i)$ and $\mathcal{B}(M_{12})$ for $i = 1, 2$. Such relations are known to exist, since $(\text{ad } M_i)^2 M_j = 0$ for $i \neq j$ by [HS1, Lemma 4.2]. However, we show next that these can be deduced from some of the already established relations.

Step 4. Let A denote the quotient of $T(M_1 \oplus M_2)$ by the ideal generated by (5.2) and (5.3). In A the following relations hold

$$(5.8) \quad (\text{ad}_c x_1)x_{12} = 0, \quad (\text{ad}_c x_{\overline{1}})x_{12} = 0, \quad (\text{ad}_c x_1)x_{\overline{12}} = 0, \quad (\text{ad}_c x_{\overline{1}})x_{\overline{12}} = 0;$$

$$(5.9) \quad [x_{12}, x_2]_c = 0, \quad [x_{12}, x_{\overline{2}}]_c = 0, \quad [x_{\overline{12}}, x_2]_c = 0, \quad [x_{\overline{12}}, x_{\overline{2}}]_c = 0;$$

$$(5.10) \quad x_{\overline{12}}^2 = 0.$$

Proof. First, $(\text{ad}_c x_1)x_{12} = (\text{ad}_c x_1^2)x_2 = 0$ since $x_1^2 = 0$. Analogously $(\text{ad}_c x_{\overline{1}})x_{\overline{12}} = 0$ follows from $x_{\overline{1}}^2 = 0$. Next, using that x_{12} is a scalar multiple of $x_{\overline{12}}$, we get that $(\text{ad}_c x_{\overline{1}})x_{12} = 0$ follows from $x_{\overline{1}}^2 = 0$. Since $x_{\overline{12}}$ is a scalar multiple of $x_{\overline{12}}$ we get $(\text{ad}_c x_1)x_{\overline{12}} = 0$ from $x_1^2 = 0$. Thus (5.8) hold.

For (5.9), unpacking the definitions we see that $[x_{12}, x_2]_c = [x_{\overline{12}}, x_{\overline{2}}]_c = 0$ follow from $x_2^2 = x_{\overline{2}}^2 = 0$. Now $[x_{12}, x_{\overline{2}}]_c = 0$ follows using the previous argument, since x_{12} is a scalar multiple of $x_{\overline{12}}$. The remaining relation holds similarly.

Finally, we show that (5.10) follows from (5.8), (5.9) and (5.3). In fact,

$$(5.11) \quad \begin{aligned} (x_{12})^2 &= (x_1 x_2 - x_{\overline{2}} x_1)(\text{ad}_c x_1)(x_2) \\ &= -(\chi_2(\kappa g_1^2))^{-1} x_1 (\text{ad}_c x_1)(x_2) x_{\overline{2}} - \chi_2(g_1^2) x_{\overline{2}} (\text{ad}_c x_{\overline{1}})(x_2) x_1 \\ &= -\chi_2(\kappa)(\text{ad}_c x_{\overline{1}})(x_2) x_1 x_{\overline{2}} + \chi_2(\kappa g_1^2)(\text{ad}_c x_{\overline{1}})(x_2) x_2 x_1 \\ &= -\chi_2(\kappa)(\text{ad}_c x_{\overline{1}})(x_2)(\text{ad}_c x_1)(x_{\overline{2}}) = \chi_2(\kappa g_1^2)(x_{\overline{12}})^2, \end{aligned}$$

as claimed. \square

We are ready to give a presentation of $\mathcal{B}(M_1 \oplus M_2)$. Let R denote the quotient of $T(M_1 \oplus M_2)$ by the ideal generated by (5.2), (5.3) and (5.4). We already know from (5.6) and Step 3 that the canonical projection $T(M_1 \oplus M_2) \rightarrow \mathcal{B}(M)$ factors to a surjective

algebra map $R \rightarrow \mathcal{B}(M)$. We show that this map is injective. Since $\dim \mathcal{B}(M) = 2^6$, it is enough to verify that the set (5.5) linearly generates R . Let J denote the subspace spanned by (5.5) in R . Since J contains 1, it is enough to show that J is a left ideal, which reduces to verify that $x_i I \subset I$ and $x_{\bar{i}} I \subset I$ for $i = 1, 2$. Clearly $x_2 J \subset J$; as $x_{\bar{2}} x_2$ is a scalar multiple of $x_2 x_{\bar{2}}$, it is equally clear that $x_{\bar{2}} J \subset J$. So we need to verify that $x_1 J \subset J$ and $x_{\bar{1}} J \subset J$, which follow since the (5.8) and (5.9) hold in R by Step 4. \square

5.1.2. *Type α_3 .* Let G denote a non-abelian group and let $M = M_1 \oplus M_2 \oplus M_3$ in ${}_{\mathbb{k}G}\mathcal{YD}$ of type α_3 . By [HV, Lemma 5.2] for $i \in \mathbb{I}_2$, the subgroup $\langle \kappa, g_i, g_{i+1} \rangle \subset G$ is a quotient of G_2 , see (5.1). Next we describe $\mathcal{B}(M)$.

Proposition 5.2. *For M of type α_3 , the Nichols algebra $\mathcal{B}(M)$ is presented by generators $x_i, x_{\bar{i}}, i \in \mathbb{I}_3$ and relations*

$$(5.12) \quad x_i^2 = x_{\bar{i}}^2 = 0, \quad (\text{ad}_c x_i) x_{\bar{i}} = 0, \quad i \in \mathbb{I};$$

$$(5.13) \quad x_{i\bar{j}} = -\chi_j(g_i^2) x_{\bar{i}j}, \quad x_{\bar{i}j} = -\chi_i(\kappa) x_{ij}, \quad i < j, a_{ij} = -1;$$

$$(5.14) \quad x_{ij}^2 = 0, \quad [x_{ij}, x_{i\bar{j}}]_c = 0, \quad i < j, a_{ij} = -1;$$

$$(5.15) \quad x_{13} = x_{\bar{1}\bar{3}} = 0, \quad x_{\bar{1}3} = x_{1\bar{3}} = 0;$$

$$(5.16) \quad x_{123}^2 = 0, \quad [x_{123}, x_{\bar{1}\bar{2}\bar{3}}]_c = 0;$$

$$(5.17) \quad (\text{ad}_c x_2) x_{123} = 0, \quad (\text{ad}_c x_2) x_{\bar{1}\bar{2}\bar{3}} = 0.$$

A PBW basis of $\mathcal{B}(M)$ is given by

$$(5.18) \quad \{x_3^{a_3} x_{\bar{3}}^{b_3} x_{23}^{a_{23}} x_{\bar{2}\bar{3}}^{b_{23}} x_2^{a_2} x_{\bar{2}}^{b_2} x_{123}^{a_{123}} x_{\bar{1}\bar{2}\bar{3}}^{b_{123}} x_{12}^{a_{12}} x_{\bar{1}\bar{2}}^{b_{12}} x_1^{a_1} x_{\bar{1}}^{b_1} : a_\beta, b_\beta \in \{0, 1\}\}.$$

Proof. Again, we proceed in several steps.

Step 1. The multiplication map is an isomorphism of \mathbb{Z}^3 -graded objects in ${}_{\mathbb{k}G}\mathcal{YD}$:

$$\mathcal{B}(M_3) \otimes \mathcal{B}(M_{23}) \otimes \mathcal{B}(M_2) \otimes \mathcal{B}(M_{123}) \otimes \mathcal{B}(M_{12}) \otimes \mathcal{B}(M_1) \simeq \mathcal{B}(M)$$

This follows by [HS1, Theorem 2.6]. Next we give some relations that hold in $\mathcal{B}(M)$.

Step 2. The relations (5.12), (5.13) and (5.15) hold in $\mathcal{B}(M)$.

We know that $M_{13} = 0$, so (5.15) hold, while (5.12) and (5.13) follow since $M_1 \oplus M_2$ and $M_2 \oplus M_3$ are of type α_2 . \square

Following the treatment in §5.1.1, we describe $\mathcal{B}(M_{123}) \subset \mathcal{B}(M)$. As a first step in this direction, we seek for a basis of M_{123} .

Step 3. Let A denote the quotient of $T(M_1 \oplus M_2 \oplus M_3)$ by the ideal generated by (5.15) and (5.13). In A the following relations hold

$$(5.19) \quad x_{1\bar{2}\bar{3}} = -\chi_2(g_1^2) x_{\bar{1}\bar{2}\bar{3}}, \quad x_{\bar{1}\bar{2}3} = -\chi_1(\kappa) x_{123};$$

Proof. We only verify the first one. We compute

$$\begin{aligned} x_{1\bar{2}3} &= [x_1, [x_{\bar{2}}, x_3]_c]_c = [[x_1, x_{\bar{2}}]_c, x_3]_c + (g_1 \cdot x_{\bar{2}})[x_1, x_3]_c - \chi_3(\kappa)[x_1, x_{\bar{3}}]x_{\bar{2}} \\ &= [[x_1, x_{\bar{2}}]_c, x_3]_c = -\chi_2(g_1^2)[[x_{\bar{1}}, x_2]_c, x_3]_c = -\chi_2(g_1^2)[x_{\bar{1}}, [x_2, x_3]_c]_c, \end{aligned}$$

where the second equality follows by (2.9), the third from (5.15), the fourth from (5.13), and the fifth one by (2.9) and (5.15). \square

Surprisingly, there are further restrictions on the character χ_2 :

Step 4. If $M_1 \oplus M_2 \oplus M_3$ is of type α_3 , then $\chi_2(g_1^2) = \chi_2(g_3^2)$.

Proof. We compute the action of g_2 on $x_{123} \in \mathcal{B}(M)$ following two different approaches. Applying (2.10) first, followed by (5.13) we get

$$\begin{aligned} g_2 \cdot x_{123} &= -(\text{ad}_c x_{\bar{1}})(\text{ad}_c x_2)x_{\bar{3}} = \chi_3(g_2^2)x_{\bar{1}\bar{2}3} \\ &= \chi_3(g_2^2)(x_{\bar{1}}((\text{ad}_c x_{\bar{2}})x_3) - \chi_2(\kappa g_1^2)\chi_3(\kappa g_1)((\text{ad}_c x_2)x_3)x_{\bar{1}}) \end{aligned}$$

On the other hand, if we first unpack the definition of $\text{ad}_c x_1$ and then let g_2 act, we get

$$\begin{aligned} g_2 \cdot x_{123} &= g_2 \cdot (x_1((\text{ad}_c x_2)x_3) - \chi_3(g_1)((\text{ad}_c x_{\bar{2}})x_3)x_1) \\ &= -x_{\bar{1}}((\text{ad}_c x_2)x_{\bar{3}}) + \chi_2(\kappa)\chi_3(g_1)((\text{ad}_c x_{\bar{2}})x_{\bar{3}})x_{\bar{1}} \\ &= \chi_3(g_2^2)x_{\bar{1}}((\text{ad}_c x_{\bar{2}})x_3) - \chi_3(g_1)((\text{ad}_c x_2)x_3)x_{\bar{1}} \\ &= \chi_3(g_2^2)(x_{\bar{1}}((\text{ad}_c x_{\bar{2}})x_3) - \chi_2(\kappa g_3^2)\chi_3(\kappa g_1)((\text{ad}_c x_2)x_3)x_{\bar{1}}) \end{aligned}$$

where the second equality follows by (2.10), the third one from (5.13) and the last one from $\chi_2(\kappa g_3^2)\chi_3(\kappa g_2^2) = 1$.

These two equations give $(\chi_2(g_1^2) - \chi_2(g_3^2))((\text{ad}_c x_2)x_3)x_{\bar{1}} = 0$, and the claim follows because $\partial_{x_{\bar{1}}}((\text{ad}_c x_2)x_3)x_{\bar{1}} = (\text{ad}_c x_2)x_3 \neq 0$. \square

The next result is analogue to Step 3 in Proposition 5.1, and its proof follows from similar arguments, so we only give a sketch.

Step 5. (a) A basis of M_{123} is $\{x_{123}, x_{\bar{1}\bar{2}3}\}$, where the braid is diagonal with matrix

$$(5.20) \quad \begin{pmatrix} -1 & -\chi_1(\kappa)\chi_2(\kappa)\chi_3(\kappa) \\ -\chi_1(\kappa)\chi_2(\kappa)\chi_3(\kappa) & -1 \end{pmatrix}.$$

(b) Relations (5.16) hold in $\mathcal{B}(M)$.

Proof. (a) The set $\{x_{123}, x_{\bar{1}\bar{2}3}\}$ linearly spans M_{123} by Step 3, and it is linearly independent since so are $\partial_{x_3}(x_{123}) = x_{12}$ and $\partial_{x_3}(x_{\bar{1}\bar{2}3}) = x_{\bar{1}\bar{2}}$. The braiding is computed using (2.10) and Step 4. Now (b) follows from (a) and Step 1. \square

The Nichols algebra of M_{123} is presented by the relations (5.16) and $x_{123}^2 = 0$; we will show that this last relation can be deduced from others.

Braided commutations between M_i and M_{123} for $i = 1, 3$ can now be deduced:

Step 6. Let A denote the quotient of $T(M_1 \oplus M_2 \oplus M_3)$ by the ideal generated by (5.2), (5.15) and (5.13). In A the following relations hold

$$(5.21) \quad (\text{ad}_c x_1)x_{123} = 0, \quad (\text{ad}_c x_{\overline{1}})x_{123} = 0, \quad (\text{ad}_c x_1)x_{\overline{123}} = 0, \quad (\text{ad}_c x_{\overline{1}})x_{\overline{123}} = 0;$$

$$(5.22) \quad [x_{123}, x_3]_c = 0, \quad [x_{123}, x_{\overline{3}}]_c = 0, \quad [x_{\overline{123}}, x_3]_c = 0, \quad [x_{\overline{123}}, x_{\overline{3}}]_c = 0;$$

$$(5.23) \quad (\text{ad}_c x_{\overline{2}})x_{123} = -\chi_1(g_2^{-2})(\text{ad}_c x_2)x_{\overline{123}}, \quad (\text{ad}_c x_{\overline{2}})x_{\overline{123}} = -\chi_1(g_2^{-2})(\text{ad}_c x_2)x_{123}.$$

Proof. The relations (5.21) can be verified using the argument in the proof of Step 4 of Proposition 5.1. For the first relation in (5.22), use (2.9) to get

$$[x_{123}, x_3]_c = [[x_1, x_{23}]_c, x_3]_c = [x_1, [x_{23}, x_3]_c]_c - x_{\overline{2}}[x_1, x_3]_c + [x_1, x_{\overline{3}}]_c x_2 = 0.$$

The three remaining relations follow similarly. For (5.23), use (2.9), (5.13) and (5.8) to get

$$\begin{aligned} (\text{ad}_c x_{\overline{2}})x_{123} &= [x_{\overline{2}1}, x_{23}] + (g_2 \kappa \cdot x_1)[x_{\overline{2}}, x_{23}]_c - \chi_3(g_1)[x_{\overline{2}}, x_{\overline{2}3}]_c x_1 \\ &= -\chi_1(g_2^{-2})[x_{\overline{2}1}, x_{23}] = -\chi_1(g_2^{-2})(\text{ad}_c x_2)x_{\overline{123}}. \end{aligned}$$

and the other relation follows analogously. \square

We employ skew derivations to verify braided commutations between M_2 and M_{123} .

Step 7. Relations (5.17) hold in $\mathcal{B}(M)$.

Proof. We focus on the first relation. One can see directly from (2.10) that

$$\partial_{x_i}(M_{123}) = \partial_{x_{\overline{i}}}(M_{123}) = 0, \quad \partial_{x_3}(x_{123}) = x_{12}, \quad \partial_{x_{\overline{3}}}(x_{\overline{123}}) = x_{\overline{12}}.$$

Now, using (2.10) and also Step 4 we get

$$\partial_{x_{\overline{3}}}(x_{123}) = -\chi_2(\kappa)x_{1\overline{2}}, \quad \partial_{x_3}(x_{\overline{123}}) = -\chi_3(g_2^2)x_{\overline{12}}.$$

For $i \in \mathbb{I}_2$, both ∂_{x_i} and $\partial_{x_{\overline{i}}}$ annihilate the first relation, and we compute

$$\partial_{x_3}((\text{ad}_c x_2)x_{123}) = x_2 x_{12} - \chi_3(g_2^2)x_{\overline{12}}x_{\overline{2}} = x_2 x_{12} + \chi_3(\kappa g_2^2)x_{12}x_{\overline{2}} = \chi_3(\kappa g_2^2)[x_{12}, x_{\overline{2}}]_c = 0,$$

where the third equality follows by (5.13). Similarly, $\partial_{x_{\overline{3}}}((\text{ad}_c x_2)x_{123}) = \chi_2(\kappa)[x_{\overline{12}}, x_{\overline{2}}]_c = 0$. The other relations follow analogously. \square

Next we deduce that braided brackets between $\mathcal{B}(M_{23})$ and $\mathcal{B}(M_{12})$ can be rewritten in terms of intermediate factors of the decomposition given in Step 1.

Step 8. We have $[\mathcal{B}(M_{12}), \mathcal{B}(M_{23})]_c \subset \mathcal{B}(M_2) \otimes \mathcal{B}(M_{123})$ in $\mathcal{B}(M)$.

Proof. This follows from $x_2^2 = 0$, $x_{\bar{2}}^2 = 0$, (5.13) and (5.17). As an illustration:

$$[x_{12}, x_{23}]_c = (\text{ad}_c x_1)(\text{ad}_c x_2^2)x_3 - x_{\bar{2}}x_{123} - x_{12\bar{3}}x_2,$$

which belongs to $\mathbb{k}x_{\bar{2}}x_{123} + \mathbb{k}x_2x_{\bar{1}23}$. \square

Step 9. Let A denote the quotient of $T(M_1 \oplus M_2 \oplus M_3)$ by the ideal generated by (5.13), (5.19), (5.21), (5.22), and (5.17). Then in A we have

$$(5.24) \quad x_{123}^2 = \chi_2(g_1^4)\chi_3(g_2^2)x_{\bar{1}23}^2$$

Proof. Use (5.21), (5.22), and (5.17) several times to get explicit braided commutations between x_{123} and each of its monomials:

$$\begin{aligned} x_{123}^2 &= (x_1x_2x_3 - x_1x_{\bar{3}}x_2 - \chi_3(g_1)x_{\bar{2}}x_3x_1 + \chi_3(\kappa g_1)x_{\bar{3}}x_{\bar{2}}x_1)x_{123} \\ &= \chi_2(g_1^2)x_{\bar{1}23}(x_1(x_2x_{\bar{3}} - \chi_3(g_2^2)x_3x_2) - \chi_3(\kappa g_1)(x_{\bar{2}}x_{\bar{3}} - \chi_3(\kappa g_2^2)x_3x_{\bar{2}})x_1) \\ &= \chi_2(g_1^4)\chi_3(g_2^2)x_{\bar{1}23}^2, \end{aligned}$$

where the last equality follows from (5.13) and (5.19). \square

As in the proof of Proposition 5.1, by Steps 2, 5 and 7, there exists an algebra surjection from \mathcal{B} , the algebra presented by relations in Proposition 5.2, onto $\mathcal{B}(M)$. Now we use Steps 1, 3, 6, 8, 9 together with the fact that $\dim \mathcal{B}(M) = 2^{12}$ to conclude that $\mathcal{B} = \mathcal{B}(M)$. \square

5.2. A kind of distinguished pre-Nichols algebra. For Nichols algebras of diagonal type, a presentation by generators and relations was achieved by the first author in [An2]. A fundamental role is played by an intermediate quotient known as the distinguished pre-Nichols algebra. Inspired by that construction, we define a pre-Nichols algebra $\widehat{\mathcal{B}}(M)$ for each one of the braidings M described in §2.3.1, 2.3.2, 2.3.3. More precisely, our construction resembles the algebra $\widehat{\mathcal{B}}_{\mathfrak{q}}$ introduced in Remark 2.3.

This pre-Nichols algebra will also play a key role in §6, where we describe the liftings.

Given $M = \bigoplus_{i \in \mathbb{I}} M_i$ as in §2.3.1, 2.3.2, 2.3.3, let σ as in Theorem 3.18, and denote by \mathfrak{q} the braiding matrix of $W := F_\sigma(M)$ (see §3.1). Recall the index ℓ defined in (3.5). Consider

$$\bar{\cdot} : \mathbb{I}_\ell \rightarrow \mathbb{I}_{\theta+\ell}, \quad \bar{i} := \begin{cases} i + \theta, & \text{for types } \alpha_\theta, \delta_\theta, \epsilon_\theta, \\ \theta + \ell - i + 1, & \text{otherwise,} \end{cases}$$

so $\mathbb{I}_{\theta+\ell} = \mathbb{I}_\theta \cup \{\bar{i} : i \in \mathbb{I}_\ell\}$. We fix a basis \mathbf{x}_i , $i \in \mathbb{I}_{\theta+\ell}$, such that

- \mathbf{x}_i , $\mathbf{x}_{\bar{i}}$ is a basis of $F_\sigma(M_i)$ for each $i \in \mathbb{I}_\ell$;
- $\mathbf{x}_i = x_i$ if $\ell < i \leq \theta$;
- the braiding in this basis is given by \mathfrak{q} ; i.e. $c(\mathbf{x}_i \otimes \mathbf{x}_j) = q_{ij}\mathbf{x}_j \otimes \mathbf{x}_i$.

Let $\mathbf{A} = (\mathbf{a}_{ij})_{i,j \in \mathbb{I}_{\theta+\ell}}$ be the Cartan matrix of \mathfrak{q} .

Remark 5.3. Let $\Xi : \mathbb{Z}^{\theta+\ell} \rightarrow \mathbb{Z}^\theta$ be the group morphism such that

$$\Xi(\alpha_i) = \alpha_i, \quad i \leq \theta; \quad \Xi(\alpha_i) = \alpha_j, \quad i > \theta, i = \bar{j}.$$

This map *identifies* the two elements of the basis above corresponding to each $F_\sigma(M_i)$ when $\dim M_i = 2$. Thus, if \mathcal{B} is an \mathbb{N}_0^θ -graded pre-Nichols algebra such that $F_\sigma(\mathcal{B})$ is $\mathbb{N}_0^{\theta+\ell}$ -graded (for the usual grading as pre-Nichols algebra of diagonal type), then

$$\dim \mathcal{B}_\beta = \sum_{\gamma \in \Xi^{-1}(\beta)} \dim F_\sigma(\mathcal{B})_\gamma, \quad \beta \in \mathbb{N}_0^\theta.$$

Thus the Hilbert series $\mathcal{H}_{\mathcal{B}}$ is the image of $\mathcal{H}_{F_\sigma(\mathcal{B})}$ under Ξ .

Let $\widehat{\mathcal{B}}(M)$ be the algebra generated $x_i, i \in \mathbb{I}, x_{\bar{j}}, j \in \mathbb{I}_\ell$, subject to the relations

$$(5.25) \quad (\text{ad}_c x_i)x_{\bar{i}}, \quad x_i^2, \quad x_{\bar{i}}^2, \quad i \in \mathbb{I}_\ell;$$

$$(5.26) \quad x_{\bar{i}}^2, \quad i > \ell;$$

$$(5.27) \quad x_{\bar{i}\bar{j}} + \chi_j(g_i^2)x_{\bar{i}\bar{j}}, \quad x_{\bar{i}\bar{j}} + \chi_i(\kappa)x_{ij}, \quad i < j \leq \ell, a_{ij} = -1;$$

$$(5.28) \quad x_{ij}, \quad x_{\bar{i}\bar{j}}, \quad x_{\bar{i}\bar{j}}, \quad x_{\bar{i}\bar{j}}, \quad i < j \leq \ell, a_{ij} = 0;$$

$$(5.29) \quad x_{ij}, \quad x_{\bar{i}\bar{j}}, \quad i \leq \ell < j, a_{ij} = 0;$$

$$(5.30) \quad (\text{ad}_c x_j)x_{ij\bar{i}} = 0, \quad a_{ij} = -2.$$

$$(5.31) \quad (\text{ad}_c x_j)x_{ijk}, \quad (\text{ad}_c x_j)x_{\bar{i}\bar{j}\bar{k}}, \quad i < j < k, a_{ji} = a_{jk} = -1.$$

Proposition 5.4. $\widehat{\mathcal{B}}(M)$ is a graded pre-Nichols algebra with Hilbert series

$$(5.32) \quad \mathcal{H}_{\widehat{\mathcal{B}}(M)} = \prod_{\substack{\beta \in \Delta_+^M - \{\alpha_i\} \\ \dim M_\beta = 2}} \left(\frac{1}{1-t^\beta} \right)^2 \prod_{j \in \mathbb{I}_\ell} (1+t_j)^2 \prod_{\substack{\beta \in \Delta_+^M - \{\alpha_i\} \\ \dim M_\beta = 1}} \left(\frac{1}{1-t^\beta} \right) \prod_{j > \ell} (1+t_j).$$

Proof. First, $\widehat{\mathcal{B}}(M)$ is a quotient of $T(M)$ by an homogeneous ideal, so it is a graded algebra. Also, (5.25)–(5.29) are primitive elements of $T(M)$, while those in (5.31) are primitive modulo the previous relations, see the proofs of Propositions 5.1 and 5.2. Thus $\widehat{\mathcal{B}}(M)$ is a graded pre-Nichols algebra.

We fix $G = \mathbf{G}$ defined in Remark 3.19, and M has a canonical G -Yetter-Drinfeld module structure. Now $\widehat{\mathcal{B}} := F_\sigma(\widehat{\mathcal{B}}(M)) \in \mathbb{k}_G^G \mathcal{YD}$ is a pre-Nichols algebra of W such that

$$\left(\widehat{\mathcal{B}}(M) \# \mathbb{k}G \right)_\sigma \simeq \widehat{\mathcal{B}} \# \mathbb{k}G.$$

We claim that $\widehat{\mathcal{B}} = \widehat{\mathcal{B}}_q$. If so, the statement on the Hilbert series follows from Remarks 2.3 and 5.3. First, note that (2.8) holds in $\widehat{\mathcal{B}}$ for all $i \in \mathbb{I}_{\theta+\ell}$ by (5.25) and (5.26).

To verify (2.6), let $i < j$ be such that $\mathbf{a}_{ij} = 0$. We consider 5 cases.

- ◇ $i \leq \ell, j = \bar{i}$. The space of primitive elements of $T(M)$ of degree $2\alpha_i \in \mathbb{N}_0^\theta$ is three-dimensional, spanned by $x_i^2, x_{\bar{i}}^2$ and $x_{i\bar{i}}$. Thus $\widehat{\mathcal{B}}(M)$ has no primitive elements of degree $2\alpha_i$. On the other hand, the spaces of primitive elements of $T(W)$ of degree $\alpha_i + \alpha_j \in \mathbb{N}_0^{\theta+\ell}$ is one-dimensional spanned by x_{ij} , and those of degrees $2\alpha_i, 2\alpha_j$ are also one-dimensional, spanned by \mathbf{x}_i^2 and \mathbf{x}_j^2 respectively. As the space of homogeneous primitive elements of $\widehat{\mathcal{B}}$ coincides with that of $\widehat{\mathcal{B}}(M)$, we have that $\mathbf{x}_{ij} = 0$ in $\widehat{\mathcal{B}}$.
- ◇ $i \leq \ell, \theta < j, \mathbf{a}_{ik} = 0$, where $j = \bar{k}$. The space of primitive elements of $T(M)$ of degree $\alpha_i + \alpha_k \in \mathbb{N}_0^\theta$ has dimension 4, spanned by $x_{ik}, x_{i\bar{k}}, x_{\bar{i}k}, x_{\bar{i}\bar{k}}$, so the space of primitive elements of $\widehat{\mathcal{B}}(M)$ of the same degree is 0. On the other hand, the space of primitive elements of $T(W)$ of degree $\alpha_i + \alpha_k$ is spanned by $\mathbf{x}_{ik}, \mathbf{x}_{ij}, \mathbf{x}_{k\bar{i}}, \mathbf{x}_{j\bar{i}}$. As the space of homogeneous primitive elements of $\widehat{\mathcal{B}}$ coincides with that of $\widehat{\mathcal{B}}(M)$, we have that $\mathbf{x}_{ij} = 0 = \mathbf{x}_{j\bar{i}}$ in $\widehat{\mathcal{B}}$. This also shows that $\mathbf{x}_{ik} = \mathbf{x}_{k\bar{i}} = 0$.
- ◇ $i \leq \ell, \theta < j, \mathbf{a}_{ik} = -1$, where $j = \bar{k}$. The space of primitive elements of $T(M)$ of degree $\alpha_i + \alpha_k \in \mathbb{N}_0^\theta$ is two-dimensional, spanned by (5.27), so $\widehat{\mathcal{B}}(M)$ has no primitive elements of this degree. On the other hand, the spaces of primitive elements of $T(W)$ of degrees $\alpha_i + \alpha_j$ and $\alpha_{\bar{i}} + \alpha_k$ are one-dimensional, spanned by \mathbf{x}_{ij} and $\mathbf{x}_{k\bar{i}}$ respectively, and those of degrees $\alpha_i + \alpha_k$ and $\alpha_j + \alpha_{\bar{i}}$ are 0 since $a_{ik} = a_{\bar{i}j} = -1$. Hence $\mathbf{x}_{ij} = \mathbf{x}_{k\bar{i}} = 0$ in $\widehat{\mathcal{B}}$.
- ◇ $i \leq \ell < j \leq \theta$. Here $a_{ij} = 0$, and the space of primitive elements of $T(M)$ of degree $\alpha_i + \alpha_j \in \mathbb{N}_0^\theta$ is two-dimensional, spanned by x_{ij} and $x_{i\bar{j}}$, so the space of primitive elements of $\widehat{\mathcal{B}}(M)$ of this degree is 0. On the other hand, the spaces of primitive elements of $T(W)$ of degrees $\alpha_i + \alpha_j$ and $\alpha_{\bar{i}} + \alpha_j$ are one-dimensional, spanned by \mathbf{x}_{ij} and $\mathbf{x}_{j\bar{i}}$, respectively. Hence $\mathbf{x}_{ij} = \mathbf{x}_{j\bar{i}} = 0$ in $\widehat{\mathcal{B}}$.
- ◇ $i, j \leq \ell$. Here $a_{ij} = 0$, respectively $a_{i\bar{j}} = 0$, and this case is the second one.

Now we check (2.5) in $T(W)$, modulo (2.4) and (2.6). We have 3 cases:

- ♡ $j \leq \ell$. The space of primitive elements of $T(M)$, modulo (5.25), (5.26), (5.27), (5.28) and (5.29), of degree $\alpha_i + 2\alpha_j + \alpha_k$ has dimension ≤ 2 and is spanned by (5.31): we can use skew-derivations as in type α_3 , see the proof of Proposition 5.2. Thus the space of primitive elements of $\widehat{\mathcal{B}}(M)$ of this degree is 0. On the other hand, the space of primitive elements of $T(W)$, modulo (2.4) and (2.6), of each degree in $\Xi^{-1}(\alpha_i + 2\alpha_j + \alpha_k)$ is either one-dimensional or 0. Indeed, the non-zero cases are spanned by
 - $[\mathbf{x}_{ijk}, \mathbf{x}_j]_c, [\mathbf{x}_{i\bar{j}k}, \mathbf{x}_{\bar{j}}]_c$ if $j < \ell$;
 - $[\mathbf{x}_{\theta-2\theta-1\theta}, \mathbf{x}_{\theta-1}]_c, [\mathbf{x}_{\theta+2\theta+1\theta}, \mathbf{x}_{\theta+1}]_c$ if $j = \theta - 1$ in type γ_θ ;
 - $[\mathbf{x}_{123}, \mathbf{x}_2]_c, [\mathbf{x}_{653}, \mathbf{x}_5]_c$ if $j = 2$ in type ϕ_4 .

As the space of homogeneous primitive elements of $\widehat{\mathcal{B}}$ coincides with that of $\widehat{\mathcal{B}}(M)$, we deduce that (2.5) hold in $\widehat{\mathcal{B}}$.

♡ $j > \ell$. There are three possibilities: $i = \theta - 1, j = \theta, k = \bar{i}$ in types γ_θ or ϕ_4 , and $i \in \{2, 5\}, j = 3, k = 4$ in type ϕ_4 . The proof is analogous, using (5.30) for the first case, and (5.31) for the last one.

From the analysis above there exists a surjective Hopf algebra map $\widehat{\mathcal{B}} \rightarrow \widehat{\mathcal{B}}_q$. In a similar way, checking spaces of homogeneous primitive elements of appropriate degree, each defining relation of $\widehat{\mathcal{B}}(M)$ annihilates in $F_\sigma^{-1}(\widehat{\mathcal{B}}_q) = F_{\sigma^{-1}}(\widehat{\mathcal{B}}_q)$, so there exists a surjective Hopf algebra map $F_\sigma^{-1}(\widehat{\mathcal{B}}_q) \rightarrow \widehat{\mathcal{B}}(M)$. As F_σ preserves the \mathbb{N}_0 -graduation of the pre-Nichols algebras, both surjective maps are indeed isomorphisms. Hence, $\widehat{\mathcal{B}} = \widehat{\mathcal{B}}_q$ as we claimed. \square

5.3. The subalgebra of coinvariants. Let $\pi : \widehat{\mathcal{B}}(M) \twoheadrightarrow \mathcal{B}(M)$ be the canonical projection, $\mathcal{Z}(M) := \widehat{\mathcal{B}}(M)^{\text{co}\pi}$ the subalgebra of coinvariants. The next step towards the presentation of $\mathcal{B}(M)$ is to describe $\mathcal{Z}(M)$. To uncover the structure of this subalgebra, we will use a cocycle σ as in Theorem 3.18 to translate the problem to the diagonal setting, where the situation is better understood. In particular, we compute the Hilbert series of $\mathcal{Z}(M)$. Since we know that of $\widehat{\mathcal{B}}(M)$, we will thus obtain the Hilbert series of $\mathcal{B}(M)$.

To do so, and also to compute a PBW basis of $\mathcal{B}(M)$ later on, we fix a reduced expression of the element w_0 of maximal length (or equivalently, a convex order on Δ_+) for each type. Using this reduced expression, [HS3] defines a submodule $M_\beta \in \frac{\mathbb{k}G}{\mathbb{k}G} \mathcal{YD}$, $\beta \in \Delta_+$. We exhibit a basis $\{x_\beta\}$ or $\{x_\beta, x_{\bar{\beta}}\}$ of the submodule M_β , depending on its dimension.

α_θ : The set of positive roots is $\Delta_+ = \{\alpha_{ij} : i \leq j \in \mathbb{I}\}$, and

$$\alpha_1 < \alpha_{12} < \alpha_2 < \cdots < \alpha_{\theta-1} < \alpha_{1\theta} < \alpha_{2\theta} < \cdots < \alpha_\theta$$

is a convex order on Δ_+ . By [HS3] the modules M_β can be defined as

$$(5.33) \quad M_{\alpha_{ij}} = (\text{ad}_c M_i) \cdots (\text{ad}_c M_{j-1}) M_j.$$

A basis of $M_{\alpha_{ij}}$ is given by

$$(5.34) \quad x_{\alpha_{ij}} = x_{i i+1 \cdots j}, \quad x_{\overline{\alpha_{ij}}} = x_{\bar{i} i+1 \cdots j}.$$

δ_θ : The positive roots are $\Delta_+ = \{\alpha_{ij} : i \leq j \in \mathbb{I}\} - \{\alpha_{\theta-1\theta}\} \cup \{\alpha_{i\theta-2} + \alpha_\theta : i \in \mathbb{I}_{\theta-2}\} \cup \{\alpha_{i\theta-2} + \alpha_{j\theta} : i < j \in \mathbb{I}_{\theta-2}\}$, and a convex order on Δ_+ is given by

$$\begin{aligned} \alpha_1 &< \alpha_{12} < \alpha_2 < \cdots < \alpha_{\theta-2} < \alpha_{1\theta-1} < \cdots < \alpha_{\theta-1} < \cdots < \alpha_{\theta-1} \\ &< \alpha_{1\theta} + \alpha_{2\theta-2} < \cdots < \alpha_{1\theta} + \alpha_{\theta-2} < \alpha_{2\theta} + \alpha_{3\theta-2} < \cdots < \alpha_{2\theta} + \alpha_{\theta-2} \\ &< \cdots < \alpha_{\theta-3\theta} + \alpha_{\theta-2} < \alpha_{1\theta} < \cdots < \alpha_{\theta-2\theta} < \alpha_\theta. \end{aligned}$$

The Yetter-Drinfeld modules $M_{\alpha_{ij}}, j \neq \theta$, are defined as for type α_θ . For $j = \theta$ let

$$M_{\alpha_{i\theta}} = (\text{ad}_c M_i) \cdots (\text{ad}_c M_{\theta-3})(\text{ad}_c M_{\theta-1})(\text{ad}_c M_{\theta-2}) M_\theta.$$

For the other roots we have

$$\begin{aligned} M_{\alpha_{i\theta-2} + \alpha_\theta} &= (\text{ad}_c M_i) \cdots (\text{ad}_c M_{\theta-2}) M_\theta, \\ M_{\alpha_{i\theta} + \alpha_{j\theta-2}} &= [M_{\alpha_{i\theta-2} + \alpha_\theta}, M_{\alpha_{j\theta-1}}]_c. \end{aligned}$$

A basis of M_β for either $\beta = \alpha_{ij}$ or $\beta = \alpha_{i\theta-2} + \alpha_\theta$ is given as in (5.34). For $\beta = \alpha_{i\theta} + \alpha_{j\theta-2}$, a basis of M_β is

$$(5.35) \quad x_\beta = [x_{i\dots\theta-2\theta}, x_{j\dots\theta-1}]_c, \quad x_{\bar{\beta}} = [x_{\bar{i}i+1\dots\theta-2\theta}, x_{j\dots\theta-1}]_c.$$

ϵ_θ : Here one can fix a convex order as in [An1, §5]. For braidings of diagonal type, a PBW basis is obtained recursively on the height of the roots, starting with $\mathbf{x}_{\alpha_i} = \mathbf{x}_i$ for simple roots, and later $\mathbf{x}_\beta = [\mathbf{x}_{\beta_1}, \mathbf{x}_{\beta_2}]_c$ for some pair (β_1, β_2) such that $\beta_1 + \beta_2 = \beta$, see [An3, Corollary 3.17]. For each non simple root $\beta \in \Delta_+$ we have, accordingly,

$$M_\beta = [M_{\beta_1}, M_{\beta_2}]_c, \quad x_\beta = [x_{\beta_1}, x_{\beta_2}]_c, \quad x_{\bar{\beta}} = [x_{\bar{\beta}_1}, x_{\beta_2}]_c.$$

γ_θ : Now $\Delta_+ = \{\alpha_{ij} : i \leq j \in \mathbb{I}\} \cup \{\alpha_{i\theta} + \alpha_{k\theta-1} : i \leq j \in \mathbb{I}_{\theta-1}\}$, and

$$\begin{aligned} \alpha_1 &< \alpha_{12} < \alpha_2 < \dots < \alpha_{\theta-1} < \alpha_{1\theta} < \alpha_{2\theta} < \dots < \\ \alpha_{1\theta} + \alpha_{1\theta-1} &< \alpha_{1\theta} + \alpha_{2\theta-1} < \dots < \alpha_{1\theta} + \alpha_{\theta-1} < \dots < \\ \alpha_{2\theta} + \alpha_{2\theta-1} &< \dots < \alpha_{2\theta} + \alpha_{\theta-1} < \dots < 2\alpha_{\theta-1} + \alpha_\theta < \alpha_{\theta-1} + \alpha_\theta < \alpha_\theta. \end{aligned}$$

is a convex order associated to the following reduced expression of w_0 :

$$s_1(s_2s_1)(s_3s_2s_1) \cdots (s_{\theta-1} \cdots s_1)(s_\theta s_{\theta-1} \cdots s_1)(s_\theta \cdots s_2) \cdots s_\theta.$$

The Yetter-Drinfeld modules $M_{\alpha_{ij}}$, $\alpha_{ij} \neq \alpha_\theta$, are as in (5.33), and (5.34) is a basis as well. Now, M_{α_θ} is one-dimensional, spanned by x_θ , and for the other roots we check that

$$(5.36) \quad M_{\alpha_{i\theta} + \alpha_{k\theta-1}} = [M_{\alpha_{i\theta}}, M_{\alpha_{k\theta-1}}]_c.$$

A basis of $M_{\alpha_{i\theta} + \alpha_{k\theta-1}}$ is given by

$$(5.37) \quad x_\beta = [x_{i\dots\theta}, x_{k\dots\theta-1}]_c, \quad x_{\bar{\beta}} = [x_{\bar{i}i+1\dots\theta}, x_{k\dots\theta-1}]_c.$$

ϕ_4 : The element w_0 of maximal length has a reduced expression

$$(5.38) \quad s_1s_2s_3s_4s_2s_3s_1s_2s_3s_4s_1s_2s_3s_2s_1s_2s_3s_4s_2s_3s_2s_3s_4s_3,$$

which induces the following convex order on the set of positive roots:

$$\begin{array}{cccccccc} 1, & 12, & 1^22^23, & 1^22^234, & 123, & 1^22^23^24, & 12^23, & 1^22^33^24, \\ 1^22^43^24, & 1^22^43^34, & 1234, & 12^234, & 1^22^43^34^2, & 12^23^24, & 12^33^24, & 2, \\ 2^23, & 2^234, & 23, & 2^23^24, & 234, & 3, & 34, & 4. \end{array}$$

We denote by β_i the i -th root according with this order.

Next we give, for each non-simple root $\beta = 1^a2^b3^c4^d$ such that $d \neq 0$, a basis for each Yetter-Drinfeld submodule M_β (if $d = 0$, then we choose a basis as for γ_θ):

$$\begin{array}{ll} M_{1^22^234} : \{ [x_{12}, x_{\bar{1}234}]_c \}, & M_{1^22^23^24} : \{ [x_{123}, x_{\bar{1}234}]_c \}, \\ M_{1^22^23^324} : \{ [x_{1223}, x_{1234}]_c, [x_{\bar{1}223}, x_{1234}]_c \}, & M_{1^22^43^24} : \{ [x_{1223}, x_{12234}]_c \}, \\ M_{1^22^43^34} : \{ [x_{1223}, x_{122324}]_c \}, & M_{1234} : \{ x_{1234}, x_{\bar{1}234} \}, \\ M_{12^234} : \{ [x_{1234}, x_2]_c, [x_{\bar{1}234}, x_2]_c \}, & M_{1^22^43^34^2} : \{ [x_{12^234}, x_{12^23^24}]_c \}, \end{array}$$

$$\begin{aligned}
M_{12^23^24} &: \{[x_{1234}, x_{23}]_c, [x_{\overline{1}234}, x_{23}]_c\}, & M_{2^234} &: \{x_{\overline{2}234}\}, \\
M_{12^33^24} &: \{[x_{12^23^24}, x_2]_c, [x_{\overline{1}2^23^24}, x_2]_c\}, & M_{2^23^24} &: \{[x_{\overline{2}3}, x_{234}]_c\}, \\
M_{234} &: \{x_{234}, x_{\overline{2}34}\}, & M_{34} &: \{x_{34}\}.
\end{aligned}$$

where x_β is the first vector fixed for M_β , while for $\dim M_\beta = 2$, we denote by $x_{\overline{\beta}}$ the second vectors in the order fixed above.

Remark 5.5. Let $\beta = 1^{a_1}2^{a_2} \dots \theta^{a_\theta} \in \Delta_+^M$, $g_\beta := g_1^{a_1}g_2^{a_2} \dots g_\theta^{a_\theta} \in G$.

- (1) If $\dim M_\beta = 1$, then x_β has G -degree g_β .
- (2) If $\dim M_\beta = 2$, then x_β has G -degree g_β and $x_{\overline{\beta}}$ has G -degree $g_\beta\kappa$.

We omit the details of the proof that either $\{x_\beta\}$ or $\{x_\beta, x_{\overline{\beta}}\}$ is a basis of M_β . The first step is to check that M_β is spanned by $\{x_\beta\}$, respectively $\{x_\beta, x_{\overline{\beta}}\}$, using the defining relations of $\widehat{\mathcal{B}}(M)$; this can be done recursively on the convex (total) order. If $\dim M_\beta = 2$ we see that $x_\beta, x_{\overline{\beta}}$ have different G -degree by Remark 5.5, so they are linearly independent.

Proposition 5.6. *The subalgebra $\mathcal{Z}(M) = \widehat{\mathcal{B}}(M)^{\text{co}\pi}$ of coinvariants under the canonical projection $\pi: \widehat{\mathcal{B}}(M) \rightarrow \mathcal{B}(M)$ is a Hopf subalgebra of $\widehat{\mathcal{B}}(M)$. It is a skew-polynomial algebra in variables*

$$(5.39) \quad x_\beta^2, \quad \beta \in \Delta_+^M - \{\alpha_i\},$$

$$(5.40) \quad [x_\beta, x_{\overline{\beta}}]_c, \quad \beta \in \Delta_+^M - \{\alpha_i\} \text{ such that } \dim M_\beta = 2.$$

Proof. We proceed in several steps. In what follows, $\beta \in \Delta_+$ is not simple.

Step 1. For each $\beta \in \Delta_+$ such that $\dim M_\beta = 1$, we have $x_\beta^2 \in M_\beta^2 \cap \mathcal{Z}(M)$.

For each $\beta \in \Delta_+$ such that $\dim M_\beta = 2$, we have $x_\beta^2, x_{\overline{\beta}}^2, [x_\beta, x_{\overline{\beta}}]_c \in \mathcal{Z}(M)$.

The subalgebra spanned by M_β is isomorphic to the Nichols algebra $\mathcal{B}(M_\beta)$ by [HS2]. Assume first that $\dim M_\beta = 1$. The braiding of M_β satisfies that $c(x_\beta \otimes x_\beta) = -x_\beta \otimes x_\beta$, so $x_\beta^2 = 0$ in $\mathcal{B}(M)$. Thus $x_\beta^2 \in \ker \pi \cap M_\beta^2$, and applying F_σ we get

$$F_\sigma(x_\beta^2) \in F_\sigma(\ker \pi \cap M_\beta^2) = \ker \pi \cap F_\sigma(M_\beta)^2 = \mathcal{Z}(W).$$

Hence $x_\beta^2 \in \mathcal{Z}(M)$, since F_σ leaves the coalgebra structure unchanged.

Now, if $\dim M_\beta = 2$, then the braiding of M_β satisfies

$$c(x_\beta \otimes x_\beta) = -x_\beta \otimes x_\beta, \quad c(x_{\overline{\beta}} \otimes x_{\overline{\beta}}) = -x_{\overline{\beta}} \otimes x_{\overline{\beta}}, \quad c^2(x_\beta \otimes x_{\overline{\beta}}) = x_\beta \otimes x_{\overline{\beta}}.$$

By a similar argument, $x_\beta^2, x_{\overline{\beta}}^2, [x_\beta, x_{\overline{\beta}}]_c \in \mathcal{Z}(M)$. □

Step 2. For each $\beta \in \Delta_+$ such that $\dim M_\beta = 1$, $\{x_\beta\}$ is a basis of $F_\sigma(M_\beta)$.

For each $\beta \in \Delta_+$ such that $\dim M_\beta = 2$, $\{x_\beta, x_{\overline{\beta}}\}$ is a basis of $F_\sigma(M_\beta)$.

The statement certainly holds for simple roots, so we fix a non-simple root β . For types $\alpha_\theta, \delta_\theta, \epsilon_\theta$, we always have $\dim M_\beta = 2$ and $M_\beta = [M_{\beta_1}, M_{\beta_2}]_c$ for some $\beta_1, \beta_2 \in \Delta_+$. Notice that $(\text{ad}_c x_i)x_{\bar{j}} = 0$ for all $i, j \in \mathbb{I}$ since i, \bar{j} belong to different connected components of the Dynkin diagram of type $X_\theta \times X_\theta$. Hence $[\mathbf{x}_\gamma, \mathbf{x}_{\bar{\delta}}]_c$ for all $\gamma, \delta \in \Delta_+$, and arguing recursively,

$$\begin{aligned} F_\sigma(M_\beta) &= F_\sigma([M_{\beta_1}, M_{\beta_2}]_c) = [F_\sigma(M_{\beta_1}), F_\sigma(M_{\beta_2})]_c \\ &= \left[\mathbb{k}\mathbf{x}_{\beta_1} + \mathbb{k}\mathbf{x}_{\bar{\beta}_1}, \mathbb{k}\mathbf{x}_{\beta_2} + \mathbb{k}\mathbf{x}_{\bar{\beta}_2} \right]_c = \mathbb{k} \left[\mathbf{x}_{\beta_1}, \mathbf{x}_{\beta_2} \right]_c + \mathbb{k} \left[\mathbf{x}_{\bar{\beta}_1}, \mathbf{x}_{\bar{\beta}_2} \right]_c = \mathbb{k}\mathbf{x}_\beta + \mathbb{k}\mathbf{x}_{\bar{\beta}}. \end{aligned}$$

For multiply-laced types, the proof follows similarly, case-by-case. \square

Step 3. There exist \mathbb{N}_0^θ -homogeneous elements

- $y_\beta \in M_\beta^2$ of G -degree g_β^2 when $\dim M_\beta = 1$,
- $y_\beta, y_{\bar{\beta}} \in M_\beta^2$ of G -degree g_β^2 , respectively $g_{\bar{\beta}}^2\kappa$, when $\dim M_\beta = 2$,

that q -commute with every G -homogeneous element of $\hat{\mathcal{B}}$. Moreover, $\mathcal{Z}(M)$ is a skew-polynomial algebra in these variables.

Assume that $\dim M_\beta = 2$. Note that, for all $i \in \mathbb{I}$, the elements $\mathbf{x}_i^2, \mathbf{x}_{\bar{i}}^2$ are linear combinations of elements of G -degree g_i^2 and $g_i^2\kappa$ with non-trivial components on each degree. Hence, the elements $\mathbf{x}_\beta^2, \mathbf{x}_{\bar{\beta}}^2$ are written as linear combinations of elements of G -degree g_β^2 and $g_\beta^2\kappa$ with non-trivial components on each degree, since they are obtained applying Lusztig's isomorphisms to appropriate $\mathbf{x}_i^2, \mathbf{x}_{\bar{i}}^2$. Thus there exist y_β and $y_{\bar{\beta}}$ of G -degrees g_β^2 and $g_{\bar{\beta}}^2\kappa$ respectively, that span the same as \mathbf{x}_β^2 and $\mathbf{x}_{\bar{\beta}}^2$. When $\dim M_\beta = 1$ we may choose $y_\beta = \mathbf{x}_\beta^2$.

As $\mathcal{Z}(W)$ is a skew-polynomial algebra in variables $x_\beta^2, x_{\bar{\beta}}^2$ and each element $x_\beta^2, x_{\bar{\beta}}^2$ is skew-central, the same holds with respect to $y_\beta, y_{\bar{\beta}}$.

Let $\beta \in \Delta_+$ be such that $\dim M_\beta = 2$. We set

$$y_\beta := F_\sigma^{-1}(y_\beta) \in M_\beta^2, \quad y_{\bar{\beta}} := F_\sigma^{-1}(y_{\bar{\beta}}) \in M_{\bar{\beta}}^2.$$

Then $y_\beta, y_{\bar{\beta}} \in \mathcal{Z}(M)$ since F_σ preserves the coalgebra structure. Note that

$$F_\sigma(x_i y_\beta) = \sigma(g_i, g_\beta^2) F_\sigma(x_i) y_\beta, \quad F_\sigma(y_\beta x_i) = \sigma(g_\beta^2, g_i) y_\beta F_\sigma(x_i),$$

and these two elements differ up to a non-zero scalar for all $i \in \mathbb{I}$, thus y_β is skew-central. The same happens for $y_{\bar{\beta}}$, and for y_β when $\dim M_\beta = 1$. In particular, the image under F_σ of a multiplication of various y_β 's, $y_{\bar{\beta}}$'s is the multiplication to the corresponding y_β 's, $y_{\bar{\beta}}$'s up to a non-zero scalar, and the Step follows. \square

Step 4. For each $\beta \in \Delta_+$ such that $\dim M_\beta = 1$, we have $\mathbb{k}y_\beta = \mathbb{k}x_\beta^2$.

For each $\beta \in \Delta_+$ such that $\dim M_\beta = 2$, we have $\mathbb{k}y_\beta = \mathbb{k}x_\beta^2, \mathbb{k}y_{\bar{\beta}} = \mathbb{k}[x_\beta, x_{\bar{\beta}}]_c$.

Assume first that $\dim M_\beta = 1$. Then $\dim M_\beta^2 = 1$ in $\widehat{\mathcal{B}}(M)$, and the claim follows since both y_β and x_β^2 are generators of M_β^2 .

Now assume that $\dim M_\beta = 2$. In this case $\dim M_\beta^2 = 3$ in $\widehat{\mathcal{B}}(M)$ since $\dim F_\sigma(M_\beta^2) = 3$ in $\widehat{\mathcal{B}}(W)$. On the other hand, $\dim M_\beta^2 = 1$ in $\mathcal{B}(V)$: it is generated by $x_{\overline{\beta}}x_\beta$ since $x_\beta^2 = x_{\overline{\beta}}^2 = [x_\beta, x_{\overline{\beta}}]_c = 0$. Notice that $\dim F_\sigma(M_\beta)^2 \cap \mathcal{Z}(W) = 2$, and $F_\sigma(M_\beta)^2 \cap \mathcal{Z}(W)$ contains elements with non-trivial components in degrees g_β^2 and $g_{\overline{\beta}}^2\kappa$. Hence $\dim M_\beta^2 \cap \mathcal{Z}(M) = 2$, with one-dimensional homogeneous components of degrees g_β^2 and $g_{\overline{\beta}}^2\kappa$. Thus $\mathbb{k}y_\beta = \mathbb{k}x_\beta^2 = \mathbb{k}x_{\overline{\beta}}^2$ and $\mathbb{k}y_{\overline{\beta}} = \mathbb{k}[x_\beta, x_{\overline{\beta}}]_c$, as claimed. \square

Hence Step 3 shows that $\mathcal{Z}(M)$ is a skew-polynomial algebra, and Step 4 assures that we can choose generating variables as stated. \square

5.4. A presentation of the Nichols algebra. Here we put together the results obtained in §5.2, 5.3 to get a presentation, a PBW basis and the Hilbert series for the Nichols algebra.

Theorem 5.7. (i) *A set of PBW generators for $\mathcal{B}(M)$ is given by*

$$(5.41) \quad x_\beta, \beta \in \Delta_+; \quad x_{\overline{\beta}} \text{ when } \dim M_\beta = 2.$$

The height of $x_\beta, x_{\overline{\beta}}$ is 2 for all $\beta \in \Delta_+$.

(ii) *The Nichols algebra $\mathcal{B}(M)$ is presented by generators $x_i, i \in \mathbb{I}, x_{\overline{j}}, j \in \mathbb{I}_\ell$, and relations (5.25), (5.26), (5.27), (5.28), (5.29), (5.30), (5.31), (5.39) and (5.40).*

Proof. (i) By [HS1, Theorem 2.6], the multiplication map

$$\bigotimes_{\beta \in \Delta_+^M} \mathcal{B}(M_\beta) \rightarrow \mathcal{B}(M)$$

is an isomorphism of \mathbb{Z}^θ -graded objects in ${}_{\mathbb{k}G}^{\mathbb{k}G}\mathcal{YD}$. If $\dim M_\beta = 1$, then M_β has braiding $-id$ and $1, x_\beta$ is a basis of $\mathcal{B}(M_\beta)$. If $\dim M_\beta = 2$, then M_β has braiding as in (2.16): that is, $\mathcal{B}(M_\beta)$ is a quantum plane with basis $1, x_\beta, x_{\overline{\beta}}, x_\beta x_{\overline{\beta}}$, and the claim follows.

(ii) By Proposition 5.4, relations (5.25), (5.26), (5.27), (5.28), (5.29), (5.30) and (5.31) hold in $\mathcal{B}(M)$. Also, (5.39) and (5.40) hold in $\mathcal{B}(M)$ since the subalgebra generated by M_β is isomorphic to $\mathcal{B}(M_\beta)$ as an algebra. Therefore, if \mathcal{B} denotes the quotient of $T(M)$ by all these relations, then there exists a canonical projection $\mathcal{B} \twoheadrightarrow \mathcal{B}(M)$ of graded Hopf algebras. Moreover,

$$\mathcal{B} = \widehat{\mathcal{B}}(V) / \langle \mathcal{Z}(V)^+ \rangle,$$

so $\mathcal{H}_{\widehat{\mathcal{B}}(V)} = \mathcal{H}_{\mathcal{Z}(V)} \mathcal{H}_{\mathcal{B}}$ by [ACS, Lemma 2.4]. By Propositions 5.6 and 5.4,

$$\mathcal{H}_{\mathcal{B}} = \left(\prod_{\beta \in \Delta_+^V: \dim M_\beta=2} (1+t^\beta)^2 \right) \left(\prod_{\beta \in \Delta_+^V: \dim M_\beta=1} (1+t^\beta) \right) = \mathcal{H}_{\mathcal{B}(V)},$$

and we deduce that $\mathcal{B} = \mathcal{B}(M)$. \square

Remark 5.8. The order of the elements in the PBW basis in Theorem 5.7 (i) is given by the expression of the element w_0 of maximal length fixed below. For example, for type ϕ_4 we have the following PBW basis:

$$(5.42) \quad \begin{aligned} & x_4^{a_1} x_{34}^{a_2} x_{234}^{a_3} x_{234}^{a_4} x_{2^2 3^2 4}^{a_5} x_{2^3}^{a_6} x_{2^3}^{a_7} x_{2^2 3^4}^{a_8} x_{2^2 3}^{a_9} x_{2^2}^{a_{10}} x_2^{a_{11}} x_{12^3 3^2 4}^{a_{12}} x_{12^3 3^2 4}^{a_{13}} x_{12^2 3^2 4}^{a_{14}} \\ & x_{12^2 3^2 4}^{a_{15}} x_{1^2 2^4 3^3 4^2}^{a_{16}} x_{12^2 3^4}^{a_{17}} x_{12^2 3^4}^{a_{18}} x_{12^3 4}^{a_{19}} x_{12^3 4}^{a_{20}} x_{3^4}^{a_{21}} x_{1^2 2^4 3^3 4}^{a_{22}} x_{1^2 2^4 3^2 4}^{a_{23}} x_{1^2 2^3 3^2 4}^{a_{24}} \\ & x_{1^2 2^3 3^2 4}^{a_{25}} x_{12^2 3}^{a_{26}} x_{12^2 3}^{a_{27}} x_{1^2 2^2 3^2 4}^{a_{28}} x_{12^3}^{a_{29}} x_{12^3}^{a_{30}} x_{1^2 2^2 3^4}^{a_{31}} x_{1^2 2^2 3}^{a_{32}} x_{12}^{a_{33}} x_{12}^{a_{34}} x_1^{a_{35}} x_1^{a_{36}}, \quad a_i \in \mathbb{I}_{0,1}. \end{aligned}$$

5.5. Rigidity of Nichols algebras. We briefly discuss rigidity for Nichols algebras of types $\alpha_\theta, \gamma_\theta, \delta_\theta, \epsilon_\theta$ and ϕ_4 , inspired by [AnKM] where rigidity for finite-dimensional Nichols algebras over abelian groups is studied. This will come in handy in the next section, where we study the liftings of these Nichols algebras.

Let $\mathcal{R}_M \subset T(M)$ be the set of defining relations as in Theorem 5.7. We start by describing the Yetter-Drinfeld structure of $\mathbb{k}\mathcal{R}_M$. Recall the index ℓ introduced in (3.5).

Remark 5.9. (1) The G -degree of $x_{i\bar{i}}$ is $g_i^2 \kappa$. Using Example 2.4, it is easy to see that

$$(5.43) \quad \bar{\chi}_i : G \rightarrow \mathbb{k}^\times, \quad \bar{\chi}_i(h) := \begin{cases} \chi_i(hg_j^{-1}hg_j) & h \in G^{g_i}, \\ \chi_i(\kappa)\chi_i(h^2) & h \notin G^{g_i}, \end{cases}$$

is a character, and the G -action on $x_{i\bar{i}}$ is given by $\bar{\chi}_i$.

(2) The G -degree of x_i^2 and $x_{\bar{i}}^2$ is g_i^2 . The G -action when $i \leq \ell$ is

$$g \cdot x_i^2 = \begin{cases} \chi_i^2(g)x_i^2, & g \in G^{g_i}, \\ \chi_i^2(g_j^{-1}g)x_i^2, & g \notin G^{g_i}; \end{cases} \quad g \cdot x_{\bar{i}}^2 = \begin{cases} \chi_i^2(g)x_{\bar{i}}^2, & g \in G^{g_i}, \\ \chi_i^2(gg_j)x_{\bar{i}}^2, & g \notin G^{g_i}. \end{cases}$$

If $i > \ell$, then the action on x_i^2 is given by χ_i^2 .

(3) Let $i < j \leq \ell$ be such that $a_{ij} = -1$, and set

$$r_1 := x_{i\bar{j}} + \chi_j(g_i^2)x_{i\bar{j}}, \quad r_2 := x_{\bar{i}\bar{j}} + \chi_i(\kappa)x_{i\bar{j}}.$$

The G -degrees of r_1 and r_2 are $g_i g_j \kappa$ and $g_i g_j$. The G -action is given by

$h \in$	$h \cdot r_1$	$h \cdot r_2$
$G^{g_i} \cap G^{g_j}$	$\chi_i(h)\chi_j(h)r_1$	$\chi_i(h)\chi_j(h)r_2$
$G^{g_i} - G^{g_j}$	$\chi_i(\kappa h)\chi_j(g_i h)r_2$	$\chi_i(\kappa h)\chi_j(g_i^{-1}h)r_2$
$G^{g_j} - G^{g_i}$	$\chi_i(g_j^{-1}h)\chi_j(\kappa h)r_2$	$\chi_i(hg_j)\chi_j(\kappa h)r_1$
$G - (G^{g_i} \cup G^{g_j})$	$\chi_i(hg_j)\chi_j(g_i h)r_1$	$\chi_i(hg_j^{-1})\chi_j(g_i^{-1}h)r_2$

(4) Let $i < j \leq \ell$ be such that $a_{ij} = 0$. Both x_{ij} and $x_{\bar{i}\bar{j}}$ have G -degree $g_i g_j$, while the G -degree of $x_{i\bar{j}}$ and $x_{\bar{i}j}$ is $g_i g_j \kappa$. For the action, as G^{g_i} and G^{g_j} are both subgroups of

index 2, there are two possibilities. If $G^{g_i} \neq G^{g_j^2}$, then we choose $g_a \in G^{g_i} - G^{g_j}$ and $g_b \in G^{g_j} - G^{g_i}$. By Example 2.4, the G -action is given by:

$h \in$	$h \cdot x_{ij}$	$h \cdot x_{i\bar{j}}$	$h \cdot x_{i\bar{j}}$	$h \cdot x_{\bar{i}j}$
$G^{g_i} \cap G^{g_j}$	$\chi_i(h)\chi_j(h)x_{ij}$	$\chi_i(g_b^{-1}hg_b)\chi_j(g_a^{-1}hg_a)x_{i\bar{j}}$	$\chi_i(h)\chi_j(g_a^{-1}hg_a)x_{i\bar{j}}$	$\chi_i(g_b^{-1}hg_b)\chi_j(h)x_{\bar{i}j}$
$G^{g_i} - G^{g_j}$	$\chi_i(h)\chi_j(g_a^{-1}h)x_{i\bar{j}}$	$\chi_i(g_b^{-1}hg_b)\chi_j(hg_a)x_{\bar{i}j}$	$\chi_i(h)\chi_j(hg_a)x_{ij}$	$\chi_i(g_b^{-1}hg_b)\chi_j(g_a^{-1}h)x_{\bar{i}j}$
$G^{g_j} - G^{g_i}$	$\chi_i(g_b^{-1}h)\chi_j(h)x_{i\bar{j}}$	$\chi_i(hg_b)\chi_j(g_a^{-1}hg_a)x_{i\bar{j}}$	$\chi_i(g_b^{-1}h)\chi_j(g_a^{-1}hg_a)x_{i\bar{j}}$	$\chi_i(hg_b)\chi_j(h)x_{ij}$
$G - (G^{g_i} \cup G^{g_j})$	$\chi_i(g_b^{-1}h)\chi_j(g_a^{-1}h)x_{i\bar{j}}$	$\chi_i(hg_b)\chi_j(hg_a)x_{ij}$	$\chi_i(g_b^{-1}h)\chi_j(hg_a)x_{i\bar{j}}$	$\chi_i(hg_b)\chi_j(g_a^{-1}h)x_{\bar{i}j}$

Hence $x_{ij}, x_{i\bar{j}}, x_{i\bar{j}}, x_{\bar{i}j}$ span a 4-dimensional irreducible Yetter-Drinfeld submodule.

In the case $G^{g_i} = G^{g_j}$ we choose $g_a = g_b \notin G^{g_i}$, and the action is described by the second and last rows of the previous table. There are two Yetter-Drinfeld submodules, spanned by $\{x_{ij}, x_{i\bar{j}}\}$, and by $\{x_{i\bar{j}}, x_{\bar{i}j}\}$.

- (5) Let $i \leq \ell < j$ be such that $a_{ij} = 0$. The G -degree of x_{ij} is $g_i g_j$, while the G -degree of $x_{i\bar{j}}$ is $g_i g_j \kappa$. For the action, notice that $g_j \in Z(G)$: if we pick $\mathbf{g} \notin G^{g_i}$, then

$h \in$	$h \cdot x_{ij}$	$h \cdot x_{i\bar{j}}$
G^{g_i}	$\chi_i(h)\chi_j(h)x_{ij}$	$\chi_i(\mathbf{g}^{-1}h\mathbf{g})\chi_j(h)x_{i\bar{j}}$
$G - G^{g_i}$	$\chi_i(\mathbf{g}^{-1}h)\chi_j(h)x_{i\bar{j}}$	$\chi_i(h\mathbf{g})\chi_j(h)x_{ij}$

- (6) For i, j such that $a_{ij} = -2$, the G -degree of $(\text{ad}_c x_j)x_{i\bar{j}}$ is $g_i^2 g_j^2 \kappa$, where G acts via $\bar{\chi}_1 \chi_j^2$.

- (7) Let $i < j < k$ be such that $a_{ji} = a_{jk} = -1$. Consider

$$\mathbf{r}_1 := (\text{ad}_c x_j)x_{ijk}, \quad \mathbf{r}_2 := (\text{ad}_c x_j)x_{\bar{i}jk},$$

which have G -degrees $g_i g_j^2 g_k \kappa$ and $g_i g_j^2 g_k$ respectively. The G -action is given by

$h \in$	$\bar{\chi}_j(h)^{-1}h \cdot \mathbf{r}_1$	$\bar{\chi}_j(h)^{-1}h \cdot \mathbf{r}_2$
$G^{g_i} \cap G^{g_k}$	$\chi_i(h)\chi_k(h)\mathbf{r}_1$	$\chi_i(g_j^{-1}hg_j)\chi_k(h)\mathbf{r}_2$
$G^{g_i} - G^{g_k}$	$\chi_i(h)\chi_k(g_j^{-1}h)\mathbf{r}_2$	$\chi_i(g_j^{-1}hg_j)\chi_k(g_j^{-1}h)\mathbf{r}_1$
$G^{g_k} - G^{g_i}$	$\chi_i(g_j^{-1}h)\chi_k(h)\mathbf{r}_2$	$\chi_i(hg_j)\chi_k(h)\mathbf{r}_1$
$G - (G^{g_i} \cup G^{g_k})$	$\chi_i(g_j^{-1}h)\chi_k(g_j^{-1}h)\mathbf{r}_1$	$\chi_i(hg_j)\chi_k(g_j^{-1}h)\mathbf{r}_2$

- (8) Let $\beta \in \Delta_+^M - \{\alpha_i\}$. The G -degrees of x_β^2 and $[x_\beta, x_{\bar{\beta}}]_c$ are g_β^2 and $g_\beta^2 \kappa$, respectively, where g_β is as in Remark 5.5. We define accordingly $\bar{\chi}_\beta := \bar{\chi}_1^{\alpha_1} \cdots \bar{\chi}_\theta^{\alpha_\theta}$, and G acts on x_β^2 and $[x_\beta, x_{\bar{\beta}}]_c$ via $\bar{\chi}_\beta$.

Remark 5.10. Let $i, j \in \mathbb{I}$ such that $G^{g_i} \neq G^{g_j}$ and $g_i, g_j \notin Z(G)$. Then

$$G^{g_i g_j} = (G^{g_i} \cap G^{g_j}) \cup (G - (G^{g_i} \cup G^{g_j}))$$

²This occurs, for example, in type α_θ whenever $j - i \geq 3$. We may choose $g_a = g_{j-1} \neq g_b = g_{i+1}$.

and the following rule defines a character

$$(5.44) \quad \chi_{ij} : G^{g_i g_j} \rightarrow \mathbb{k}^\times, \quad \chi_{ij}(h) := \begin{cases} \chi_i(h) \chi_j(h), & h \in G^{g_i} \cap G^{g_j}, \\ \chi_i(h g_j) \chi_j(h g_i), & h \notin G^{g_i} \cup G^{g_j}. \end{cases}$$

Theorem 5.11. *Let $M \in \mathbb{k}_G^G \mathcal{YD}$ of type $\alpha_\theta, \gamma_\theta, \delta_\theta, \epsilon_\theta$ or ϕ_4 . Then*

$$\mathrm{Hom}_{\mathbb{k}_G^G}(\mathbb{k} \mathcal{R}_M, M) = 0.$$

Proof. Let $\mathbf{r} \in \mathcal{R}_M$ be G -homogeneous of degree $\mathbf{g} \in G$. By direct computation, $\mathbf{g} \cdot \mathbf{r} = \mathbf{r}$. On the other hand, M has a basis $\{x_i | i \in \mathbb{I}\} \cup \{x_{\bar{j}} | j \in \mathbb{J}\}$, where $\mathbb{J} = \{i \in \mathbb{I} | \dim M_i = 2\}$. Here x_i has degree g_i , while $x_{\bar{j}}$ has degree $g_j \kappa$, and

$$g_i \cdot x_i = -x_i, \quad i \in \mathbb{I}; \quad g_j \kappa \cdot x_{\bar{j}} = -x_{\bar{j}}, \quad j \in \mathbb{J}.$$

Hence the claim follows. \square

Recall that a graded braided bialgebra is *rigid* if it has no non-trivial graded deformations. See [AnKM, §2] and the references therein for details. Next we address rigidity for $\mathcal{B}(M)$.

Theorem 5.12. *Let $M \in \mathbb{k}_G^G \mathcal{YD}$ be of type either $\alpha_\theta, \gamma_\theta, \delta_\theta, \epsilon_\theta$ or ϕ_4 . Then $\mathcal{B}(M)$ is rigid.*

Proof. The category $\mathbb{k}_G^G \mathcal{YD}$ is semisimple and $\mathrm{Hom}_{\mathbb{k}_G^G}(\mathbb{k} \mathcal{R}_M, M) = 0$ by Theorem 5.11. Hence [AnKM, Theorem 5.3] applies for $\mathcal{B}(M)$. \square

Remark 5.13. The previous notion of rigidity is related to another one introduced in [Me] coming from the action of an appropriate algebraic group on the Nichols algebra (viewed as a braided Hopf algebra). In fact, the notion of rigidity in loc. cit. is equivalent to generation in degree one, which holds by Theorem 4.1. This gives a different proof of Theorem 5.12, independent of Theorem 5.11. Anyway, we need Theorem 5.11 to compute liftings.

6. LIFTINGS OF NICHOLS ALGEBRAS

We describe all *liftings* for Nichols algebras of types $\alpha_\theta, \gamma_\theta, \delta_\theta, \epsilon_\theta$ and ϕ_4 . Even when the braided vector space is of diagonal type (that is, when κ acts trivially) we cannot invoke [AnG] since the Yetter-Drinfeld realizations considered here are not *principal*. Nevertheless we will perform an adaptation of the strategy developed in [A+, AnG].

We study the lowest rank type α_2 first, with a double purpose. On the one hand, we will only show all the details in this case, with explicit formulas for the defining relations. On the other hand, it will be the starting point to prove the general case, in which we will conclude that all liftings are cocycle deformations of the associated graded Hopf algebras.

Recall that a lifting of M over G is a finite-dimensional Hopf algebra H with coradical $\mathbb{k}G$ and infinitesimal braiding M . Hence $\mathrm{gr} H \simeq \mathcal{B}(M) \# \mathbb{k}G$ by Theorem 4.1.

The family of liftings of M over G will be indexed by a set $\mathcal{R}_M \subseteq \mathbb{k}^K$ of *deformation parameters*, where K is the number of suitable chosen Yetter-Drinfeld submodules of the subspace spanned by a minimal set \mathcal{G} of generators for the ideal defining $\mathcal{B}(M)$.

For each $\lambda \in \mathcal{R}_M$ and $i \in \mathbb{I}_K$, we define $\lambda^{(i)}, \lambda^{(-i)} \in \mathbb{k}^K$ by

$$(6.1) \quad (\lambda^{(i)})_j := \begin{cases} 0 & j \neq i, \\ \lambda_i & j = i; \end{cases} \quad (\lambda^{(-i)})_j := \begin{cases} \lambda_i & j \neq i, \\ 0 & j = i; \end{cases} \quad j \in \mathbb{I}_K.$$

The aforementioned strategy starts by choosing a *good* stratification $\mathcal{G} = \mathcal{G}_0 \sqcup \mathcal{G}_1 \sqcup \cdots \sqcup \mathcal{G}_l$, meaning that the vector space spanned by \mathcal{G}_k is a Yetter-Drinfeld submodule of $\mathcal{B}(M)$ and the elements of \mathcal{G}_k are primitive in the braided Hopf algebra $\mathcal{B}_k := T(M)/\langle \cup_{j=0}^{k-1} \mathcal{G}_j \rangle$, $k \in \mathbb{I}_{l+1}$, with one possible exception: we do not require primitiveness for the last step.

6.1. Liftings of type α_2 . Let $M \in \frac{\mathbb{k}G}{\mathbb{k}G} \mathcal{YD}$ of type α_2 . Let \mathcal{R}_M be the set of tuples $\lambda = (\mu_1, \mu_2, \lambda_1, \lambda_2, \lambda_{12}, \mu_{12}, \mu'_{12}) \in \mathbb{k}^7$ that satisfy the constraints

$$(6.2) \quad \begin{aligned} \mu_i &= 0 \text{ if either } \chi_i^2 \neq \varepsilon \text{ or } g_i^2 = 1, \quad i \in \mathbb{I}_2, \\ \lambda_i &= 0 \text{ if either } \bar{\chi}_i \neq \varepsilon \text{ or } g_i^2 = \kappa, \quad i \in \mathbb{I}_2, \\ \lambda_{12} &= 0 \text{ if } \chi_{12} \neq \varepsilon, \\ \mu_{12} &= 0 \text{ if either } \bar{\chi}_1 \bar{\chi}_2 \neq \varepsilon \text{ or } (g_1 g_2)^2 = 1. \\ \mu'_{12} &= 0 \text{ if either } \bar{\chi}_1 \bar{\chi}_2 \neq \varepsilon \text{ or } (g_1 g_2)^2 = \kappa. \end{aligned}$$

The definition of $\bar{\chi}_i$ and χ_{ij} was given in (5.43) and (5.44). This subsection is devoted to prove the following:

Theorem 6.1. *Let $M \in \frac{\mathbb{k}G}{\mathbb{k}G} \mathcal{YD}$ be of type α_2 . For each $\lambda \in \mathcal{R}_M$, let $\mathcal{L}(\lambda)$ be the quotient of $T(M) \# \mathbb{k}G$ by the following set of relations:*

$$\begin{aligned} z_i^2 - \mu_i(1 - g_i^2), & \quad z_{\bar{i}} - \lambda_i(1 - g_i^2 \kappa), \\ z_{\bar{1}2} + \chi_1(\kappa)z_{12} - \lambda_{12}(1 - g_1 g_2), & \\ z_{\bar{1}2}^2 + \lambda_1 \mu_2(1 - g_1^2 \kappa)g_2^2 - \chi_1(\kappa)\mu_1 \lambda_2(1 - g_1^2)g_2^2 \kappa - \mu_{12}(1 - g_1^2 g_2^2 \kappa), & \\ [z_{12}, z_{\bar{1}2}]_c + 2(1 + \chi_1 \chi_2(\kappa))\mu_1 \mu_2(1 - g_1^2)g_2^2 - \lambda_1 \lambda_2(\kappa - g_1^2)g_2^2 - \mu'_{12}(1 - g_1^2 g_2^2), & \end{aligned}$$

where we changed the labels $(x_i, x_{\bar{i}})_{i \in \mathbb{I}_2}$ of the generators of $T(M)$ to $(z_i, z_{\bar{i}})_{i \in \mathbb{I}_2}$. Then:

- (a) $\mathcal{L}(\lambda) \simeq L(\mathcal{A}(\lambda), \mathcal{B}(M) \# \mathbb{k}G)$.
- (b) $\mathcal{L}(\lambda)$ is a lifting of M over $\mathbb{k}G$.
- (c) $\mathcal{L}(\lambda)$ is a cocycle deformation of $\mathcal{B}(M) \# \mathbb{k}G$.

Conversely, if L is lifting of M over $\mathbb{k}G$, then there exist $\lambda \in \mathcal{R}_M$ such that $L \simeq \mathcal{L}(\lambda)$.

Fix a Yetter-Drinfeld module M over G of type α_2 . As $x_i^2, x_{\bar{i}}^2, x_{\bar{i}}, x_{\bar{1}2} + \chi_2(g_1^2)x_{\bar{1}2}$ and $x_{\bar{1}2} + \chi_1(\kappa)x_{12}$ are primitive in $T(M)$ and

$$\Delta(x_{\bar{1}2}^2) = x_{\bar{1}2}^2 \otimes 1 + x_{\bar{1}\bar{1}}g_2^2 \otimes x_2^2 - \chi_1(\kappa)x_{\bar{1}}^2 g_2^2 \kappa \otimes x_{\bar{2}\bar{2}} + g_1^2 g_2^2 \kappa \otimes x_{\bar{1}2}^2,$$

$$\begin{aligned} \Delta([x_{12}, x_{\overline{12}}]_c) &= [x_{12}, x_{\overline{12}}]_c \otimes 1 - \chi_1^{-1}(g_2^2)x_{1\overline{1}}g_2^2\kappa \otimes x_{2\overline{2}} + \chi_1(g_2^2)x_1^2g_2^2 \otimes x_2^2 \\ &\quad + \chi_1(g_2^2)x_{\overline{1}}^2g_2^2 \otimes x_2^2 + \chi_1\chi_2(\kappa)x_{\overline{1}}^2g_2^2 \otimes x_2^2 + x_1^2g_2^2 \otimes x_2^2 + g_1^2g_2^2 \otimes [x_{12}, x_{\overline{12}}]_c, \end{aligned}$$

we may choose the following stratification:

$$\mathcal{G}_0 = \{x_i^2, x_{\overline{i}}^2, x_{i\overline{i}}\}, \quad \mathcal{G}_1 = \{x_{1\overline{2}} + \chi_2(g_1^2)x_{\overline{12}}, x_{\overline{12}} + \chi_1(\kappa)x_{12}\}, \quad \mathcal{G}_2 = \{x_{12}^2, [x_{12}, x_{\overline{12}}]_c\}.$$

The Yetter-Drinfeld structure for each stratum is given in Remark 5.9.

Let $\mathcal{H}_k := \mathcal{B}_k \# \mathbb{k}G$. Next we introduce a family of cleft objects of \mathcal{H}_k parametrized by the set \mathcal{R}_M . Given $\lambda \in \mathcal{R}_M$, define $\mathcal{E}_0(\lambda) = \mathcal{B}_0 = T(M)$, $\mathcal{E}_1(\lambda) = \mathcal{B}_1$, but we change the labels of the generators to $(y_i, y_{\overline{i}})_{i \in \mathbb{I}_2}$ in order to differentiate with generators $(x_i, x_{\overline{i}})_{i \in \mathbb{I}_2}$ of the pre-Nichols algebras \mathcal{B}_k . Let

$$\begin{aligned} \mathcal{E}_1(\lambda) &:= \mathcal{E}_0(\lambda) / \langle y_i^2 - \mu_i, y_{\overline{i}}^2 - \mu_i, y_{i\overline{i}} - \lambda_i : i \in \mathbb{I}_2 \rangle, \\ \mathcal{E}_2(\lambda) &:= \mathcal{E}_1(\lambda) / \langle y_{1\overline{2}} + \chi_2(g_1^2)y_{\overline{12}} - \lambda_{12}, y_{\overline{12}} + \chi_1(\kappa)y_{12} - \lambda_{12} \rangle, \\ \mathcal{E}_3(\lambda) &:= \mathcal{E}_2(\lambda) / \langle y_{12}^2 - \mu_{12}, [y_{12}, y_{\overline{12}}]_c - \mu'_{12} \rangle. \end{aligned}$$

Each $\mathcal{E}_i(\lambda)$ is a $\mathbb{k}G$ -module algebra since the ideal is stable under the G -action by (6.2). Thus we may introduce $\mathcal{A}_i(\lambda) := \mathcal{E}_i(\lambda) \# \mathbb{k}G$.

Lemma 6.2. *Let $k \in \mathbb{I}_3$. Then $\mathcal{E}_k(\lambda) \neq 0$ and each $\mathcal{A}_k(\lambda)$ is an \mathcal{H}_k -cleft object. There exists an \mathcal{H}_k -colinear section $\gamma_k: \mathcal{H}_k \rightarrow \mathcal{A}_k$ that restricts to an algebra map $(\gamma_k)|_{\mathbb{k}G} \in \text{Alg}(\mathbb{k}G, \mathcal{A}_k)$.*

Proof. Fix $\lambda \in \mathcal{R}_M$; to simplify the notation we suppress λ and put $\mathcal{E}_k = \mathcal{E}_k(\lambda)$, $\mathcal{A}_k = \mathcal{A}_k(\lambda)$. We prove the claim recursively on k .

For $k = 1$, we notice that $\mathcal{E}_1 \neq 0$ (and a fortiori $\mathcal{A}_1 \neq 0$) by [A+, Lemma 5.16]. Notice that $g_j(y_i^2 - \mu_i)g_j^{-1} = y_i^2 - \lambda_i$ if $i \neq j$, so in \mathcal{A}_1 we have

$$\langle y_i^2 - \mu_i, y_{\overline{i}}^2 - \mu_i : i = 1, 2 \rangle = \langle y_i^2 - \mu_i : i = 1, 2 \rangle.$$

We may refine the stratification and proceed in four steps, quotient out first by x_1^2 , then by x_2^2 , now by $x_{1\overline{1}}$ and finally by $x_{2\overline{2}}$. At each step we consider the subalgebra Y' generated by the relation r in the corresponding pre-Nichols algebra, note that Y' is isomorphic to a polynomial ring in one variable since $r \in \mathcal{P}(T(M))_g - 0$, and for this g we have $g \cdot r = r$. Consider $Y = \mathcal{S}(Y')$. As Y is a polynomial algebra generated by rg^{-1} , there exists an algebra map $\phi: Y \rightarrow \mathcal{A}$ such that $\phi(rg^{-1}) = rg^{-1} - \lambda g^{-1}$, $\lambda \in \mathbb{k}$, which is \mathcal{H} -colinear. Applying repeatedly [Gu, Theorem 8] as in [A+, Proposition 5.19], \mathcal{A}_1 is a \mathcal{H}_1 -cleft object and the existence of the desired section γ_1 follows by [A+, Proposition 5.8].

For $k = 2$, it is enough to show that $\mathcal{A}_2 \neq 0$. Indeed, in that case [Gu, Theorem 8] assures that \mathcal{A}_2 is an \mathcal{H}_2 -cleft object. Now [A+, Proposition 5.8] provides a section γ_2 such that $(\gamma_2)|_{\mathbb{k}G} \in \text{Alg}(\mathbb{k}G, \mathcal{A}_2)$. As in [AnG, Lemma 3.4], non-vanishing of \mathcal{A}_2 would follow from

$$\mathcal{E}_2(\lambda^{(5)}) = \mathcal{B}_1 / \langle y_{1\overline{2}} + \chi_2(g_1^2)y_{\overline{12}} - \lambda_{12}, y_{\overline{12}} + \chi_1(\kappa)y_{12} - \lambda_{12} \rangle \neq 0.$$

Indeed, if $\varpi_1: \mathcal{A}_1(\boldsymbol{\lambda}^{(5)}) = \mathcal{H}_1 \rightarrow \mathcal{E}_2(\boldsymbol{\lambda}^{(5)}) \# \mathbb{k}G$ is the canonical projection, then the composition of the algebra map $(\rho_1)_{|\mathcal{E}_1}: \mathcal{E}_1 \rightarrow \mathcal{A}_1 \otimes \mathcal{H}_1$ with $\text{id} \otimes \varpi_1$ factors through $\mathcal{E}_2 = \mathcal{E}_2(\boldsymbol{\lambda})$.

To check that $\mathcal{E}_2(\boldsymbol{\lambda}^{(5)}) \neq 0$ we use that $(\mathcal{H}_1)_\sigma$ is the bosonization of a pre-Nichols algebra of diagonal type by G , and that the $(1, g_1 g_2 \kappa)$ - and $(1, g_1 g_2)$ -primitive elements $y_{1\bar{2}} + \chi_2(g_1^2) y_{\bar{1}2}$ and $y_{\bar{1}2} + \chi_1(\kappa) y_{12}$ span the same subspace as $\mathbf{x}_{1\bar{2}}$ and $\mathbf{x}_{\bar{1}2}$, see the proof of Proposition 5.4. The quotient $(\mathcal{H}_1)_\sigma / \langle \mathbf{x}_{1\bar{2}} - \lambda_{12}, \mathbf{x}_{\bar{1}2} - \lambda_{12} \rangle$ is not zero by [AAG], and

$$(\mathcal{H}_1)_\sigma / \langle \mathbf{x}_{1\bar{2}} - \lambda_{12}, \mathbf{x}_{\bar{1}2} - \lambda_{12} \rangle \simeq F_\sigma \left(\mathcal{A}_2(\boldsymbol{\lambda}^{(5)}) \right),$$

which implies $\mathcal{A}_2(\boldsymbol{\lambda}^{(5)}) \neq 0$. Notice that

$$g_1 (y_{1\bar{2}} + \chi_2(g_1^2) y_{\bar{1}2} - \lambda_{12}) g_1^{-1} = -\chi_1(\kappa) \chi_2(g_1^2) (y_{\bar{1}2} + \chi_1(\kappa) y_{12} - \lambda_{12}),$$

so in \mathcal{A}_2 ,

$$\langle y_{1\bar{2}} + \chi_2(g_1^2) y_{\bar{1}2} - \lambda_{12}, y_{\bar{1}2} + \chi_1(\kappa) y_{12} - \lambda_{12} \rangle = \langle y_{\bar{1}2} + \chi_1(\kappa) y_{12} - \lambda_{12} \rangle.$$

Finally, as $\mathcal{H}_2 = \widehat{\mathcal{B}}(M) \# \mathbb{k}G$, $\mathcal{H}_3 = \mathcal{B}(M) \# \mathbb{k}G$, we have $\mathcal{H}_2^{\text{co}\pi_2} = \mathcal{Z}(M)$, a skew-polynomial algebra in variables $x_{1\bar{2}}^2, [y_{12}, y_{\bar{1}2}]_c$ by Proposition 5.6. Hence [Gu, Theorem 4] applies and \mathcal{A}_3 is \mathcal{H}_3 -cleft. The claim about γ_3 follows from [A+, Proposition 5.8]. \square

Proof of Theorem 6.1. Now follows by the same procedure as in [AnS, Theorem 5.6], using Theorem 5.11. Indeed, if we define $\mathcal{L}_0(\boldsymbol{\lambda}) = \mathcal{H}_0$,

$$\begin{aligned} \mathcal{L}_1(\boldsymbol{\lambda}) &= \mathcal{L}_0(\boldsymbol{\lambda}) / \langle z_i^2 - \mu_i(1 - g_i^2), z_{i\bar{i}} - \lambda_i(1 - g_i^2 \kappa) \rangle, \\ \mathcal{L}_2(\boldsymbol{\lambda}) &= \mathcal{L}_1(\boldsymbol{\lambda}) / \langle z_{\bar{1}2} + \chi_1(\kappa) z_{12} - \lambda_{12}(1 - g_1 g_2) \rangle, \end{aligned}$$

and $\mathcal{L}_3(\boldsymbol{\lambda}) = \mathcal{L}(\boldsymbol{\lambda})$, we can prove recursively that $\mathcal{L}_i(\boldsymbol{\lambda}) \simeq L(\mathcal{A}_i(\boldsymbol{\lambda}), \mathcal{H}_i)$. \square

6.2. The general case. Let $M \in \frac{\mathbb{k}G}{\mathbb{k}G} \mathcal{YD}$ of type $\alpha_\theta, \gamma_\theta, \delta_\theta, \epsilon_\theta$ or ϕ_4 . The definition of $\bar{\chi}_i$ and χ_{ij} was given in (5.43) and (5.44). The set \mathcal{R}_M of deformation parameters contain

tuples λ satisfying the following constraints:

$$\begin{aligned}
(6.3) \quad & \mu_i = 0 \text{ if either } \chi_i^2 \neq \varepsilon \text{ or } g_i^2 = 1, i \in \mathbb{I}_\theta; \\
& \lambda_i = 0 \text{ if either } \bar{\chi}_i \neq \varepsilon \text{ or } g_i^2 = \kappa, i \in \mathbb{I}_\theta; \\
& \lambda_{ij} = 0 \text{ if } i < j \leq \ell, a_{ij} = -1, \chi_{ij} \neq \varepsilon; \\
& \lambda_{ij} = 0 \text{ if } i < j \leq \ell, a_{ij} = 0, \chi_{ij} \neq \varepsilon; \\
& \lambda'_{ij} = 0 \text{ if } i < j \leq \ell, a_{ij} = 0, \chi_{ij} \neq \varepsilon; \\
& \lambda_{ij} = \lambda'_{ij} \text{ if } i < j \leq \ell, a_{ij} = 0, G^{g_i} \neq G^{g_j}; \\
& \lambda_{ij} = 0 \text{ if } i \leq \ell < j, a_{ij} = 0, \chi_{ij} \neq \varepsilon; \\
& \lambda_{ijk} = 0 \text{ if } i < j < k, a_{ji} = a_{jk} = -1, \bar{\chi}_j \chi_{ik} \neq \varepsilon; \\
& \lambda_{ji} = 0 \text{ if } i < j, a_{ij} = -2, \bar{\chi}_i \chi_j^2 \neq \varepsilon; \\
& \mu_\beta = 0 \text{ if either } \bar{\chi}_\beta \neq \varepsilon \text{ or } g_\beta^2 = 1 (\beta \in \Delta_+^V - \{\alpha_i\}); \\
& \mu'_\beta = 0 \text{ if either } \bar{\chi}_\beta \neq \varepsilon \text{ or } g_\beta^2 = \kappa (\beta \in \Delta_+^V - \{\alpha_i\}, \dim M_\beta = 2).
\end{aligned}$$

In this subsection we prove our last main result:

Theorem 6.3. *Let $M \in \mathbb{k}^G \mathcal{YD}$ of type $\alpha_\theta, \gamma_\theta, \delta_\theta, \epsilon_\theta$ or ϕ_4 . For each $\lambda \in \mathcal{R}_V$, see (6.3), let $\mathcal{L}(\lambda)$ be the quotient of $T(M) \# \mathbb{k}G$, where we change the labels of the generators to $(z_i)_{i \in \mathbb{I}_{\theta+\ell}}$ by the following relations*

$$\begin{aligned}
& z_i^2 - \mu_i(1 - g_i^2), \quad i \in \mathbb{I}_\theta; \\
& z_{\bar{i}} - \lambda_i(1 - g_i^2 \kappa), \quad i \in \mathbb{I}_\ell; \\
& z_{\bar{i}\bar{j}} + \chi_i(\kappa)z_{ij} - \lambda_{ij}(1 - g_i g_j), \quad i < j \leq \ell, a_{ij} = -1; \\
& z_{ij} - \lambda_{ij}(1 - g_i g_j), z_{\bar{i}\bar{j}} - \lambda'_{ij}(1 - g_i g_j \kappa), \quad i < j \leq \ell, a_{ij} = 0; \\
& z_{ij} - \lambda_{ij}(1 - g_i g_j), \quad i \leq \ell < j, a_{ij} = 0; \\
& (\text{ad}_c z_j)z_{\bar{i}\bar{j}} - (2)_{\chi_j(g_i^2 \kappa)} \mu_j \lambda_i (1 - g_j^2) g_i^2 + \mu_i \lambda_j (1 - g_j^2 \kappa) g_i^2 - \lambda_{ji} (1 - g_i^2 g_j^2), \quad a_{ij} = -2; \\
& (\text{ad}_c z_j)z_{ijk} - (2)_{\chi_j(g_i g_k \kappa)} \mu_j \lambda'_{ik} (1 - g_j^2) g_i g_k \kappa + \lambda_{ik} \lambda_j (1 - g_j^2 \kappa) g_i g_k - \lambda_{ijk} (1 - g_i g_j^2 g_k \kappa), \\
& \quad i < j < k, a_{ji} = a_{jk} = -1; \\
& z_\beta^2 - \mathbf{z}_\beta - \mu_\beta(1 - g_\beta^2), \quad \beta \in \Delta_+^M - \{\alpha_i\}; \\
& \left[z_\beta, z_{\bar{\beta}} \right]_c - \mathbf{z}_\beta - \mu'_\beta(1 - g_\beta^2 \kappa), \quad \beta \in \Delta_+^M - \{\alpha_i\}, \dim M_\beta = 2,
\end{aligned}$$

where $\mathbf{z}_\beta, \mathbf{z}'_\beta \in T(M) \# \mathbb{k}G$ are defined recursively on $\beta \in \Delta_+^M - \{\alpha_i\}$ such that $z_\beta^2 - \mathbf{z}_\beta$ is $(g_\beta^2, 1)$ -primitive and $\left[z_\beta, z_{\bar{\beta}} \right]_c - \mathbf{z}'_\beta$ is $(g_\beta^2 \kappa, 1)$ -primitive in the quotient of $T(M) \# \mathbb{k}G$ by the previous relations. Then:

- (a) $\mathcal{L}(\lambda) \simeq L(\mathcal{A}(\lambda), \mathcal{B}(M) \# \mathbb{k}G)$.
- (b) $\mathcal{L}(\lambda)$ is a lifting of M over $\mathbb{k}G$.

(c) $\mathcal{L}(\boldsymbol{\lambda})$ is a cocycle deformation of $\mathcal{B}(M)\#\mathbb{k}G$.

Conversely, if L is lifting of M over $\mathbb{k}G$, then there exist $\boldsymbol{\lambda} \in \mathcal{R}_M$ such that $L \simeq \mathcal{L}_5(\boldsymbol{\lambda})$.

Let $M \in {}_{\mathbb{k}G}^{\mathbb{k}G}\mathcal{YD}$ of type $\alpha_\theta, \gamma_\theta, \delta_\theta, \epsilon_\theta$ or ϕ_4 . We choose first a stratification $\mathcal{G} = \sqcup_{i=0}^4 \mathcal{G}_i$ on the set of defining relations found in Theorem 5.7

$$\begin{aligned} \mathcal{G}_0 &= \{x_i^2, x_{\bar{i}}^2, x_{i\bar{i}} | i \in \mathbb{I}_\ell\} \cup \{x_i^2 | i > \ell\}; \\ \mathcal{G}_1 &= \{x_{i\bar{j}} + \chi_j(g_i^2)x_{i\bar{j}}, x_{i\bar{j}} + \chi_i(\kappa)x_{ij} | i < j \leq \ell, a_{ij} = -1\}; \\ \mathcal{G}_2 &= \{x_{ij}, x_{i\bar{j}}, x_{\bar{i}j}, x_{i\bar{j}} | i < j \leq \ell, a_{ij} = 0\} \cup \{x_{ij}, x_{i\bar{j}} | i \leq \ell < j, a_{ij} = 0\}; \\ \mathcal{G}_3 &= \{r_{ijk} := (\text{ad}_c x_j)x_{ijk}, \bar{r}_{ijk} := (\text{ad}_c x_j)x_{i\bar{j}k} | i < j < k, a_{ji} = a_{jk} = -1\} \\ &\quad \cup \{(\text{ad}_c x_j)x_{i\bar{j}i} | a_{ij} = -2\}; \\ \mathcal{G}_4 &= \{x_\beta^2 | \beta \in \Delta_+^V - \{\alpha_i\}\} \cup \{[x_\beta, x_{\bar{\beta}}]_c | \beta \in \Delta_+^V - \{\alpha_i\}, \dim M_\beta = 2\}. \end{aligned}$$

This stratification is good since

$$\begin{aligned} \Delta(r_{ijk}) &= r_{ijk} \otimes 1 - \chi_j(g_i g_k \kappa) x_j^2 g_i g_k \kappa \otimes x_{i\bar{k}} - \chi_j(g_i^2 g_k^2) x_j^2 g_i g_k \kappa \otimes x_{i\bar{k}} \\ &\quad + x_j x_{\bar{j}} g_i g_k \otimes x_{ik} + \chi_j(g_i g_k \kappa) x_{\bar{j}} x_j g_i g_k \otimes x_{i\bar{k}} + g_i g_j^2 g_k \kappa \otimes r_{ijk}, \\ \Delta(\bar{r}_{ijk}) &= \bar{r}_{ijk} \otimes 1 - \chi_j(g_i g_k \kappa) x_j^2 g_i g_k \otimes x_{i\bar{k}} - \chi_j(g_i^2 g_k^2) x_j^2 g_i g_k \otimes x_{ik} \\ &\quad + x_j x_{\bar{j}} g_i g_k \kappa \otimes x_{i\bar{k}} + \chi_j(g_i g_k \kappa) x_{\bar{j}} x_j g_i g_k \kappa \otimes x_{i\bar{k}} + g_i g_j^2 g_k \otimes \bar{r}_{ijk}. \end{aligned}$$

Set $\mathcal{H}_k := \mathcal{B}_k \#\mathbb{k}G$. The Yetter-Drinfeld structure of each stratum is given in Remark 5.9.

Let $\boldsymbol{\lambda} \in \mathcal{R}_M$. Define $\mathcal{E}_0(\boldsymbol{\lambda}) = \mathcal{B}_0 = T(M)$, but we change the labels of the generators to $(y_i, y_{\bar{i}})_{i \in \mathbb{I}}$ to differentiate from the generators $(x_i, x_{\bar{i}})_{i \in \mathbb{I}}$ of the pre-Nichols algebras \mathcal{B}_k . Let

$$\begin{aligned} \mathcal{E}_1(\boldsymbol{\lambda}) &:= \mathcal{E}_0(\boldsymbol{\lambda}) / \langle y_i^2 - \mu_i, y_{\bar{i}}^2 - \mu_i, y_{i\bar{i}} - \lambda_i : i \in \mathbb{I}_\theta \rangle, \\ \mathcal{E}_2(\boldsymbol{\lambda}) &:= \mathcal{E}_1(\boldsymbol{\lambda}) / \langle y_{i\bar{j}} + \chi_j(g_i^2)y_{i\bar{j}} - \lambda_{ij}, y_{i\bar{j}} + \chi_i(\kappa)y_{ij} - \lambda_{ij} | i < j \leq \ell, a_{ij} = -1 \rangle, \\ \mathcal{E}_3(\boldsymbol{\lambda}) &:= \mathcal{E}_2(\boldsymbol{\lambda}) / \left\langle \begin{array}{l} y_{ij} - \lambda_{ij}, y_{i\bar{j}} - \lambda'_{ij}, y_{\bar{i}j} - \lambda'_{ij}, y_{i\bar{j}} - \lambda_{ij}, i < j \leq \ell, a_{ij} = 0; \\ y_{ij} - \lambda_{ij}, y_{i\bar{j}} - \lambda_{ij}, i \leq \ell < j, a_{ij} = 0 \end{array} \right\rangle, \\ \mathcal{E}_4(\boldsymbol{\lambda}) &:= \mathcal{E}_3(\boldsymbol{\lambda}) / \left\langle \begin{array}{l} (\text{ad}_c y_j)y_{ijk} - \lambda_{ijk}, (\text{ad}_c y_j)y_{i\bar{j}k} - \lambda_{ijk}, a_{ji} = a_{jk} = -1, \\ (\text{ad}_c y_j)y_{i\bar{j}i} - \lambda_{ji}, a_{ij} = -2 \end{array} \right\rangle, \\ \mathcal{E}_5(\boldsymbol{\lambda}) &:= \mathcal{E}_4(\boldsymbol{\lambda}) / \left\langle \begin{array}{l} y_\beta^2 - \mu_\beta, \beta \in \Delta_+^V - \{\alpha_i\}, \\ [y_\beta, y_{\bar{\beta}}]_c - \mu'_\beta, \beta \in \Delta_+^V - \{\alpha_i\}, \dim M_\beta = 2 \end{array} \right\rangle. \end{aligned}$$

Each $\mathcal{E}_i(\boldsymbol{\lambda})$ is a $\mathbb{k}G$ -module algebra since each defining ideal above is stable under the G -action by (6.3). Thus we may introduce $\mathcal{A}_i(\boldsymbol{\lambda}) := \mathcal{E}_i(\boldsymbol{\lambda})\#\mathbb{k}G$.

Lemma 6.4. *Let $k \in \mathbb{I}_5$. Then $\mathcal{E}_k(\boldsymbol{\lambda}) \neq 0$ and each $\mathcal{A}_k(\boldsymbol{\lambda})$ is an \mathcal{H}_k -cleft object. There exists an \mathcal{H}_k -colinear section $\gamma_k: \mathcal{H}_k \rightarrow \mathcal{A}_k$ that restricts to an algebra map $(\gamma_k)|_{\mathbb{k}G} \in \text{Alg}(\mathbb{k}G, \mathcal{A}_k)$.*

Proof. Fix $\boldsymbol{\lambda} \in \mathcal{R}_M$; again, we simplify the notation and write $\mathcal{E}_k = \mathcal{E}_k(\boldsymbol{\lambda})$, $\mathcal{A}_k = \mathcal{A}_k(\boldsymbol{\lambda})$. The proof is analogous to that of Lemma 6.2, recursively on k .

When $k < 5$, the key step is to prove that $\mathcal{E}_k \neq 0$, which implies that $\mathcal{A}_k \neq 0$: if so, then [Gu, Theorem 8] applies again to conclude that \mathcal{A}_k is \mathcal{H}_k -cleft, hence there exists a section γ_k as in the statement by [A+, Proposition 5.8].

To show that $\mathcal{E}_k \neq 0$, it is enough to verify non vanishing when we *deform just one submodule of relations*; that is, to consider the case $\boldsymbol{\lambda} = \boldsymbol{\lambda}^{(i)}$ for each i and then proceed as in [AG, Lemma 3.4]. Indeed, if $\varpi_k : \mathcal{A}_k(\boldsymbol{\lambda}^{(i)}) = \mathcal{H}_{k-1} \rightarrow \mathcal{E}_k(\boldsymbol{\lambda}^{(i)}) \# \mathbb{k}G$ is the canonical projection, then the composition of the algebra map $(\rho_{k-1})|_{\mathcal{E}_{k-1}} : \mathcal{E}_{k-1} \rightarrow \mathcal{A}_{k-1} \otimes \mathcal{H}_{k-1}$ with $(\text{id} \otimes \varpi_k)$ factors through $\mathcal{E}_k = \mathcal{E}_k(\boldsymbol{\lambda})$.

To verify that $\mathcal{E}_k(\boldsymbol{\lambda}^{(i)}) \neq 0$ when the submodule of relations to be deformed is neither $\{(\text{ad}_c x_j)x_{ijk}, (\text{ad}_c x_j)x_{i\bar{j}k}\}$, where $a_{ji} = a_{jk} = -1$, nor $\{(\text{ad}_c x_j)x_{ij\bar{i}}\}$ with $a_{ij} = -2$ we may use Lemma 6.2. For these two exceptions, we adapt the argument given in Lemma 6.2 for relations $y_{1\bar{2}} + \chi_2(g_1^2)y_{\bar{1}2}$ and $y_{\bar{1}2} + \chi_1(\kappa)y_{12}$, then use the cocycle σ to reduce to deformations of Nichols algebras of diagonal type, so the result follows by [AAG, Proposition 4.2].

Finally, for $k = 5$ we have that $\mathcal{H}_4^{\text{co}\pi^4} = \mathcal{Z}(M)$ is a skew-polynomial algebra in variables $x_\beta^2, [x_\beta, x_{\bar{\beta}}]_c$ by Proposition 5.6, thus [Gu, Theorem 4] assures that \mathcal{A}_5 is \mathcal{H}_5 -cleft. The section γ_5 can be chosen so that $(\gamma_k)|_{\mathbb{k}G} \in \text{Alg}(\mathbb{k}G, \mathcal{A}_k)$ by [A+, Proposition 5.8]. \square

We are ready to prove the main theorem of this section.

Proof of Theorem 6.3. We proceed as in [AnS, Theorem 5.6], using correspondingly Theorem 5.11 and Lemma 6.4. Indeed, starting with $\mathcal{L}_0(\boldsymbol{\lambda}) = \mathcal{H}_0$, we define successive quotients $\mathcal{L}_i(\boldsymbol{\lambda})$, $i \in \mathbb{I}_5$, where \mathcal{L}_4 is the quotient by all the relations of $\mathcal{L}(\boldsymbol{\lambda})$ except the last two sets (parametrized by $\beta \in \Delta_+$). Each $\mathcal{L}_k(\boldsymbol{\lambda})$, $k < 5$, is a Hopf algebra since one if obtained from the previous one by recursively quotient by skew-primitive elements. Working as in [AAG, Theorem 1.6], $\mathcal{L}_4(\boldsymbol{\lambda}) \simeq L(\mathcal{A}_4(\boldsymbol{\lambda}), \mathcal{H}_4)$, and there exist $\mathbf{z}_\beta \in \mathcal{L}_4(\boldsymbol{\lambda})$ and $\mathbf{z}'_\beta \in \mathcal{L}_4(\boldsymbol{\lambda})$ as stated below; moreover $\mathcal{L}(\boldsymbol{\lambda}) \simeq L(\mathcal{A}_5(\boldsymbol{\lambda}), \mathcal{H}_5)$. The proof that these are all the liftings follows exactly as in [AnS, Theorem 5.6] \square

6.3. Foldings of liftings. The folding construction in [L1, Part 1] was formulated in the following general setting: Let H be a Hopf algebra and H_σ , $\sigma \in \hat{\Sigma}$ a group of biGalois objects with coherent choice of isomorphisms $\iota_{\sigma,\tau} : H_{\sigma\tau} \cong H_\sigma \square H_\tau$. By [L1, Theorem 1.6] the direct sum of algebras

$$\tilde{H} := \bigoplus_{\sigma \in \hat{\Sigma}} H_\sigma$$

can be endowed with the structure of a Hopf algebra with coproduct $\bigoplus_{\sigma,\tau} \iota_{\sigma,\tau}$.

Conversely by [L1, Theorem 3.6], any Hopf algebra \tilde{H} with Σ a central subgroup is a folding of $H = \tilde{H}/\Sigma^+\tilde{H}$ by Σ . The biGalois objects are quotients of \tilde{H} associated to a central character on Σ . The folding data in Section 3 was formulated specifically for the situation $H = \mathcal{B}(M) \# \mathbb{k}\Gamma$ and for biGalois objects arising from 2-cocycles σ on the group

Γ , trivially extended to H , and twisted Yetter-Drinfeld isomorphisms $\mathbf{u} : \mathcal{B}(M)_\sigma \rightarrow \mathcal{B}(M)$, extended by the identity on G to H . In Theorem 3.6 we have stated the folding solely in terms of σ, \mathbf{u} , while in Theorem 3.7 we have stated the folding with these specific choices of biGalois objects as above.

We now discuss the following alternative systematic way to understand the liftings of folded Nichols algebras, which we constructed in the previous section: Let H' be a lifting of $H = \mathcal{B}(M) \# \mathbb{k}\Gamma$ for a diagonal Nichols algebra, which are classified in [AAG, AnG]. Let again $(H'_\sigma)_{\sigma \in \hat{\Sigma}}$ be a group of biGalois objects over the lifting, then we have a folding \tilde{H}' , whose graded algebra is the folding $\tilde{H} = \mathcal{B}(\tilde{M}) \# \mathbb{k}G$ of H . One source for such biGalois objects could be again folding data (σ, \mathbf{u}) where in addition \cong is compatible with the lifting H' , more precisely, leaves a lifting cocycle invariant. But there are also other possibilities, namely the 2-cocycle σ over Γ could be nontrivially extended to $\mathcal{B}(M) \# \mathbb{k}\Gamma$, which would cause a folding of H that is a lifting of \tilde{H} with values in the new center.

Conversely we obtain in this way all liftings of \tilde{H} where Σ is central, acting trivially on \tilde{M} . We have already shown for each Nichols algebra in Theorem 3.18, that this trivial action can always be achieved by a Doi twist; however, it is not a-priori clear that these Doi twists carry over to the lifting. We will now use this tool to analyze the smallest example:

Example 6.5 (Case ${}^2A_2^2$). We consider the Nichols algebra of type α_2 defined in Section 2.3.1 over a group G generated by g_1, g_2 with $g_1g_2 = \kappa g_2g_1$ and κ central of order two, which is a central extension of the abelian group Γ generated by \bar{g}_1, \bar{g}_2 . We computed all its liftings in Section 6.1. We can conveniently take the group action from Remark 5.9 and the braiding from Section 2.3.1 to replace z_{ij} again by the braided commutator. For example, the first three relations depending on the parameters $\mu_i, \lambda_i, \lambda_{12}$ read

$$\begin{aligned} z_i^2 &= \mu_i(1 - g_i^2), & z_i z_{\bar{i}} + \chi_i(\kappa) z_{\bar{i}} z_i &= \lambda_i(1 - g_i^2 \kappa), \\ z_{\bar{1}} z_{\bar{2}} - \chi_2(\kappa g_1^2) z_2 z_{\bar{1}} + \chi_1(\kappa) z_1 z_2 - \chi_1(\kappa) z_{\bar{2}} z_1 &= \lambda_{12}(1 - g_1 g_2). \end{aligned}$$

Note that acting with a group element on a relation may produce more relations, as we explained in the proof of Lemma 6.2 for the cleft objects. For example, acting with g_j on the first relation produces the relation $z_{\bar{i}}^2 = \mu_i(1 - g_i^2)$.

The associated Nichols algebra and its liftings are foldings iff κ is a central element in the Hopf algebra, i.e. $\chi_1(\kappa) = \chi_2(\kappa) = 1$ (which we saw it is true up to Doi twist). In this case we saw in Remark 3.14 that the braiding diagonalizes in the basis

$$\mathbf{x}_i = z_i + q_{ij} z_{\bar{i}}, \quad \mathbf{x}_{\bar{i}} = z_i - q_{ij} z_{\bar{i}}.$$

The elements $\mathbf{x}_i, \mathbf{x}_{\bar{i}}$ are not G -homogeneous, but Γ -homogeneous with degrees $\bar{g}_i, i \in \mathbb{I}_2$. On the other hand they are G -eigenvectors with g_i, g_j acting on \mathbf{x}_i with eigenvalues $-1, -q_{ji}$

and on $\mathbf{x}_{\bar{i}}$ with eigenvalues $-1, q_{ji}$. The diagonal braiding matrix is of type $A_2 \times A_2$

$$\begin{pmatrix} -1 & -q_{12} & -1 & q_{12} \\ -q_{21} & -1 & q_{21} & -1 \\ -1 & -q_{12} & -1 & q_{12} \\ -q_{21} & -1 & q_{21} & -1 \end{pmatrix}$$

and a twisted symmetry switching the two copies. In the folding construction $z_i, z_{\bar{i}}$ arise as eigenvalues of this symmetry.

We now rewrite the relations in this basis, starting with those involving just one orbit $\mathbf{x}_i, \mathbf{x}_{\bar{i}}$, which is a diagonal Nichols algebra of type $A_1 \times A_1$:

$$\begin{aligned} \frac{1}{4}(\mathbf{x}_i + \mathbf{x}_{\bar{i}})^2 &= \mu_i(1 - g_i^2), & \frac{q_{ij}^{-2}}{4}(\mathbf{x}_i - \mathbf{x}_{\bar{i}})^2 &= \mu_i(1 - g_i^2), \\ \frac{q_{ij}^{-1}}{4}((\mathbf{x}_i + \mathbf{x}_{\bar{i}})(\mathbf{x}_i - \mathbf{x}_{\bar{i}}) + (\mathbf{x}_i - \mathbf{x}_{\bar{i}})(\mathbf{x}_i + \mathbf{x}_{\bar{i}})) &= \lambda_i(1 - g_i^2 \kappa). \end{aligned}$$

These relations rewrite to

$$\begin{aligned} \mathbf{x}_i^2 &= (1 + q_{ij}^2)\mu_i(1 - g_i^2) + q_{ij}\lambda_i(1 - g_i^2\kappa), & \mathbf{x}_i\mathbf{x}_{\bar{i}} + \mathbf{x}_{\bar{i}}\mathbf{x}_i &= 2(1 - q_{ij}^2)\mu_i(1 - g_i^2) = 0, \\ \mathbf{x}_{\bar{i}}^2 &= (1 + q_{ij}^2)\mu_i(1 - g_i^2) - q_{ij}\lambda_i(1 - g_i^2\kappa), & \frac{1}{2}(\mathbf{x}_i^2 - \mathbf{x}_{\bar{i}}^2) &= q_{ij}\lambda_i(1 - g_i^2\kappa), \end{aligned}$$

where we have to take into account that Section 6.1 states that $\mu_i \neq 0$, respectively $\lambda_i \neq 0$, only if $q_{ji}^2 = 1$, so the anti-commutator vanishes.

This is consistent with the possible liftings of diagonal $A_1 \times A_1$:

- The anti-commutator relation admits a non-trivial lifting if $\chi_i\chi_{\bar{i}} = \epsilon$, but in our case $\chi_i(g_j)\chi_{\bar{i}}(g_j) = -1$.
- The truncation relations admit non-trivial liftings if $\chi_i^2 = \epsilon$, which is the case if and only if $q_{ji}^2 = 1$. If the respective lifting parameters are equal, then this lifting datum is compatible with a folding using the group 2-cocycle. This produces the symmetric lifting depending on μ_i .
- On the other hand the antisymmetric lifting depending on λ_i requires a lifting cocycle that is non-trivially extended to the Nichols algebra. The corresponding non-trivial biGalois object is determined by plugging the non-trivial central character $\kappa \mapsto -1$.

We now turn to the relation involving λ_{12} :

$$\begin{aligned} \lambda_{12}(1 - g_1g_2) &= \frac{q_{12}^{-1}}{2}(\mathbf{x}_1 - \mathbf{x}_{\bar{1}})\frac{q_{21}^{-1}}{2}(\mathbf{x}_2 - \mathbf{x}_{\bar{2}}) - q_{12}^2\frac{1}{2}(\mathbf{x}_2 + \mathbf{x}_{\bar{2}})\frac{q_{12}^{-1}}{2}(\mathbf{x}_1 - \mathbf{x}_{\bar{1}}) \\ &\quad + \frac{1}{2}(\mathbf{x}_1 + \mathbf{x}_{\bar{1}})\frac{1}{2}(\mathbf{x}_2 + \mathbf{x}_{\bar{2}}) - \frac{q_{21}^{-1}}{2}(\mathbf{x}_2 - \mathbf{x}_{\bar{2}})\frac{1}{2}(\mathbf{x}_1 + \mathbf{x}_{\bar{1}}) \\ &= \frac{1}{2}(\mathbf{x}_1\mathbf{x}_{\bar{2}} - q_{12}\mathbf{x}_{\bar{2}}\mathbf{x}_1) + \frac{1}{2}(\mathbf{x}_{\bar{1}}\mathbf{x}_2 + q_{12}\mathbf{x}_2\mathbf{x}_{\bar{1}})\lambda_{12}(1 - g_1g_2\kappa) \\ &= -\frac{q_{12}}{2}(\mathbf{x}_1\mathbf{x}_{\bar{2}} - q_{12}\mathbf{x}_{\bar{2}}\mathbf{x}_1) + \frac{q_{12}}{2}(\mathbf{x}_{\bar{1}}\mathbf{x}_2 + q_{12}\mathbf{x}_2\mathbf{x}_{\bar{1}}). \end{aligned}$$

Section 6.1 with χ_{12} in (5.44) applied to g_i^2 and g_1g_2 states that $\lambda_{12} \neq 0$ only if $1 = q_{ii}^2q_{ij}^2$ and $1 = q_{11}q_{21}^2 \cdot q_{22}q_{12}^2$, which is again equivalent to $q_{ij}^2 = q_{ji}^2 = 1$. Possibly reversing 1, 2 we may assume we are in the case $q_{12} = 1, q_{21} = -1$, then adding and subtracting the previous relations returns:

$$\mathbf{x}_{\overline{1}}\mathbf{x}_2 + \mathbf{x}_2\mathbf{x}_{\overline{1}} = 2\lambda_{12} \left(1 - g_1g_2 \frac{\kappa+1}{2}\right), \quad \mathbf{x}_1\mathbf{x}_{\overline{2}} - \mathbf{x}_{\overline{2}}\mathbf{x}_1 = 2\lambda_{12} \left(1 - g_1g_2 \frac{\kappa-1}{2}\right).$$

On the other hand, the diagonal Nichols algebra $A_2 \times A_2$ has such liftings of

- $\mathbf{x}_{\overline{1}}\mathbf{x}_2 - q_{ji}\mathbf{x}_2\mathbf{x}_{\overline{1}}$ if $\chi_{\overline{1}}\chi_2 = \epsilon$, which is the case for $q_{12} = 1, q_{21} = -1$.
- $\mathbf{x}_1\mathbf{x}_{\overline{2}} + q_{ij}\mathbf{x}_{\overline{2}}\mathbf{x}_1$ if $\chi_1\chi_{\overline{2}} = \epsilon$, which is the case for $q_{12} = -1, q_{21} = 1$.

Altogether, there is no \mathbf{u} -symmetric lifting of this type, and the solution we find starts with a lifting H' for one of these relation, again visible at the central character $\kappa \mapsto 1$, and the other of these relations appears in the biGalois object which is nontrivially extended from the group 2-cocycle.

We refrain from discussing the last two relations in a similar manner.

7. FUTURE DIRECTIONS

We conclude by some outlook questions that naturally arise from our work.

Question 1. Is there a modified folding construction that produces the remaining Nichols algebras in Heckenberger–Vendramin classification?

Question 2. Several folded Nichols algebras in [L2] do not appear in [HV] because their support is too small. More precisely, these are the cases 2D_n and 3D_4 and ${}^2A_1^2$ familiar from Lie theory, as well as unfamiliar cases 2A_2 at a third root of unity and several cases involving other diagonal Nichols algebras. We expect that our methods can be applied in these cases.

Question 3. Which modular tensor categories can be constructed from the new pointed Hopf algebras described here?

From the categorical perspective, there is a rather unique Σ -graded extensions of tensor categories with a Σ -crossed braiding [ENOM]. Since the operations of Σ -graded extension and taking Hopf algebra representations commute, this extension could be computed by taking a Σ -symmetric Nichols algebra over an abelian group, which can then be folded to Nichols algebra over the known Σ -extension of the abelian group. To get a braiding, this would require a non-trivial associator (an effect familiar for quantum group of even order root of unity), and for $\Sigma = \mathbb{Z}_2$ conjecturally involve a Tambara–Yamigami category.

Question 4. The Logarithmic Kazhdan–Lusztig Correspondence, see e.g. [FGST, L3] conjectures the existence of a vertex algebra, realized as subalgebra of a free field algebra, whose

tensor category of representations is equivalent to representations of a small quantum group. The folding construction and the previous problem suggests an extension of this construction, where the free field algebra is replaced by an orbifold model.

APPENDIX A. PROOF OF THEOREM 3.18

Here we complete the proof of Theorem 3.18, which states that Nichols algebras of types $\alpha_\theta, \gamma_\theta, \delta_\theta, \epsilon_\theta$ and ϕ_4 become of diagonal type when an appropriate twist is performed. The cases that remain unsolved are $\alpha_2, \alpha_3, \gamma_3, \gamma_4, \delta_4$ and ϕ_4 , which will be dealt with in Proposition 3.15. The other cases are diagonal by Lemma 2.7.³

A.1. Group Cohomology Tools. We start by collecting some useful group extensions and group cohomology statements for later use, following [L1, Chapter 7]. Recall the definition of $\Gamma_{u,v,\kappa}$ in (2.27).

Definition A.1. Let Γ be an abelian group, \mathfrak{t} a generator of \mathbb{Z} . For each $w \in \Gamma, r \in \mathbb{N}$ we consider the abelian group

$$\Gamma(\sqrt[r]{w}) := \Gamma \times \mathbb{Z} / \langle (w, \mathfrak{t}^{-r}) \rangle.$$

We shall identify $g \in \Gamma, \mathfrak{t}^k, k \in \mathbb{N}$, with their images $\overline{(g, e)}, \overline{(e, \mathfrak{t}^k)}$ in $\Gamma(\sqrt[r]{w})$. The defining relation becomes $w = \mathfrak{t}^r$. We think of $\Gamma(\sqrt[r]{w})$ as the set $\{g\mathfrak{t}^k | g \in \Gamma, 0 \leq k < r\}$ with product

$$g\mathfrak{t}^j \cdot h\mathfrak{t}^k = \begin{cases} gh\mathfrak{t}^{j+k}, & \text{if } j+k < r, \\ ghw\mathfrak{t}^{j+k-r}, & \text{if } j+k \geq r, \end{cases} \quad g, h \in \Gamma, 0 \leq j, k < r.$$

Remark A.2. Fix Γ an abelian group, $u, v, w, \kappa \in \Gamma$, where $\kappa^2 = e, r \in \mathbb{N}$.

(i) The inclusion $\Gamma \hookrightarrow \Gamma(\sqrt[r]{w})$ extends to an injective map

$$\Gamma_{u,v,\kappa} \hookrightarrow \Gamma(\sqrt[r]{w})_{u,v,\kappa}.$$

(ii) Fix also $z \in \Gamma, s \in \mathbb{N}$. There is a canonical isomorphism $\Gamma(\sqrt[r]{w})(\sqrt[s]{z}) \cong \Gamma(\sqrt[s]{z})(\sqrt[r]{w})$, which in turn induces an isomorphism

$$\Gamma(\sqrt[r]{w})(\sqrt[s]{z})_{u,v,\kappa} \cong \Gamma(\sqrt[s]{z})(\sqrt[r]{w})_{u,v,\kappa}.$$

Proposition A.3. Let Γ be an abelian group, $u, v, w, \kappa \in \Gamma$, where $\kappa^2 = e, r \in \mathbb{N}$. Given $\sigma \in H^2(\Gamma_{u,v,\kappa}, \mathbb{k}^\times)$, consider $f := f_\sigma: \Gamma \rightarrow \mathbb{k}^\times$ given by

$$f(g) = \sigma(g, w) / \sigma(w, g), \quad g \in \Gamma.$$

³The corresponding folded Nichols algebras are those with symplectic root system [L2, §4] of dimension θ and radical dimension $r = \theta - 2$.

Then σ lifts to a 2-cocycle $\tilde{\sigma} \in H^2(\Gamma(\sqrt[r]{w})_{u,v,\kappa}, \mathbb{k}^\times)$ if and only if there exists $\mathbf{f} \in \widehat{\Gamma_{u,v,\kappa}}$ such that $\mathbf{f}^r = f$, $\mathbf{f}(w) = 1$. In this case,

$$(A.1) \quad \tilde{\sigma}(g, \mathbf{t})\tilde{\sigma}^{-1}(\mathbf{t}, g) = \mathbf{f}(g), \quad \text{for all } g \in \Gamma.$$

If $w = 1$, then any $\sigma \in H^2(\Gamma_{u,v,\kappa}, \mathbb{k}^\times)$ lifts to $\tilde{\sigma} \in H^2(\Gamma(\sqrt[r]{w})_{u,v,\kappa}, \mathbb{k}^\times)$.

Proof. Set $\underline{\Gamma} := \Gamma_{u,v,\kappa}$, $G = \Gamma(\sqrt[r]{w})_{u,v,\kappa}$. We can write G as the following central extension:

$$1 \rightarrow \langle w\mathbf{t}^{-r} \rangle \rightarrow \underline{\Gamma} \times \langle \mathbf{t} \rangle \rightarrow G \rightarrow 1.$$

As \mathbb{k}^\times is divisible, the map $H^1(\langle \mathbf{t} \rangle, \mathbb{k}^\times) \rightarrow H^1(\langle w\mathbf{t}^{-r} \rangle, \mathbb{k}^\times)$ is surjective and $H^2(\langle \mathbf{t} \rangle, \mathbb{k}^\times) = H^2(\langle w\mathbf{t}^{-r} \rangle, \mathbb{k}^\times) = 1$. Hence the exact sequence in [IM, §1] associated to the central extension below is

$$\begin{aligned} 1 \rightarrow H^1(G, \mathbb{k}^\times) \rightarrow H^1(\underline{\Gamma} \times \langle \mathbf{t} \rangle, \mathbb{k}^\times) \rightarrow H^1(\langle w\mathbf{t}^{-r} \rangle, \mathbb{k}^\times) \rightarrow \\ \rightarrow H^2(G, \mathbb{k}^\times) \rightarrow H^2(\underline{\Gamma} \times \langle \mathbf{t} \rangle, \mathbb{k}^\times) \rightarrow \text{Pair}(\underline{\Gamma} \times \langle \mathbf{t} \rangle, \langle w\mathbf{t}^{-r} \rangle). \end{aligned}$$

Using the results above, the Künneth formula for the cohomologies of the direct product and decomposing the pairings we get

$$1 \longrightarrow H^2(G, \mathbb{k}^\times) \longrightarrow H^2(\underline{\Gamma}, \mathbb{k}^\times) \times \text{Pair}(\underline{\Gamma}, \langle \mathbf{t} \rangle) \xrightarrow{\Phi} \text{Pair}(\underline{\Gamma}, \langle w\mathbf{t}^{-r} \rangle) \times \text{Pair}(\langle \mathbf{t} \rangle, \langle w\mathbf{t}^{-r} \rangle),$$

where Φ is defined as follows:

- for $\sigma \in H^2(\underline{\Gamma}, \mathbb{k}^\times)$, we have $\Phi(\sigma) = (B_\sigma, 1)$, where $1 \in \text{Pair}(\langle \mathbf{t} \rangle, \langle w\mathbf{t}^{-r} \rangle)$ is the trivial pairing, and $B_\sigma \in \text{Pair}(\underline{\Gamma}, \langle w\mathbf{t}^{-r} \rangle)$ is given by

$$B_\sigma(g, w\mathbf{t}^{-r}) = \sigma(g, w)\sigma^{-1}(w, g), \quad g \in \underline{\Gamma}.$$

- for $P \in \text{Pair}(\underline{\Gamma}, \langle \mathbf{t} \rangle)$, we have $\Phi(P) = (F'_P, F''_P)$, where

$$F'_P(g, w\mathbf{t}^{-r}) = P(g, \mathbf{t})^{-r}, \quad g \in \underline{\Gamma}; \quad F''_P(\mathbf{t}, w\mathbf{t}^{-r}) = P(w, \mathbf{t}).$$

Hence $(\sigma, P) \in \ker \Phi$ if and only if $\sigma(g, w)\sigma^{-1}(w, g) = P(g, \mathbf{t})^r$ for all $g \in \underline{\Gamma}$ and $P(w, \mathbf{t}) = 1$. If σ lifts to a 2-cocycle $\tilde{\sigma} \in H^2(\Gamma(\sqrt[r]{w})_{u,v,\kappa}, \mathbb{k}^\times)$, then set \mathbf{f} as in (A.1). Reciprocally, if there exists such \mathbf{f} , we define $P \in \text{Pair}(\underline{\Gamma}, \langle \mathbf{t} \rangle)$, by $P(g, \mathbf{t}) = \mathbf{f}(g)$, and get $(\sigma, P) \in \ker \Phi$. By exactness of the sequence, $\ker \Phi$ is the image of the injective map $H^2(G, \mathbb{k}^\times) \longrightarrow H^2(\underline{\Gamma}, \mathbb{k}^\times) \times \text{Pair}(\underline{\Gamma}, \langle \mathbf{t} \rangle)$; thus σ lifts to a 2-cocycle $\tilde{\sigma} \in H^2(\Gamma(\sqrt[r]{w})_{u,v,\kappa}, \mathbb{k}^\times)$. Moreover, P describes the values on the additional generator. The last statement is clear. \square

Corollary A.4. *Let Γ be an abelian group, $u, v, w, z, \kappa \in \Gamma$, where $\kappa^2 = e$, $r, s \in \mathbb{N}$. Then $\sigma \in H^2(\Gamma_{u,v,\kappa}, \mathbb{k}^\times)$ lifts to $\tilde{\sigma} \in H^2(\Gamma(\sqrt[r]{w})(\sqrt[s]{z})_{u,v,\kappa}, \mathbb{k}^\times)$ if and only if there exists $\mathbf{f}, \mathbf{g} \in \widehat{\Gamma_{u,v,\kappa}}$ such that $\mathbf{f}^r = f = \mathbf{g}^s$, $\mathbf{f}(w) = 1 = \mathbf{g}(w)$.*

Proof. Use the isomorphism in Remark A.2 and Proposition A.3. \square

Next we assume that Γ splits as $\Gamma = \Lambda \oplus \Omega$, where $u \in \Lambda$, $v, \kappa \in \Omega$. Recall the extension $\Gamma \hookrightarrow \Gamma_{u,v,\kappa} \twoheadrightarrow \mathbb{Z}_2 \times \mathbb{Z}_2 = \langle x, y \rangle$ from (2.27). Consider

- $\underline{\Lambda}$ the subgroup of $\Gamma_{u,v,\kappa}$ generated by Λ and x ,
- $\underline{\Omega}$ the subgroup of $\Gamma_{u,v,\kappa}$ generated by Ω and y .

Hence $\Gamma_{u,v,\kappa} \simeq \underline{\Lambda} \times \underline{\Omega}$, where x acts on $\underline{\Omega}$ by

$$x \cdot y = \kappa y, \quad x \cdot h = h, \quad h \in \Omega.$$

Let $\tilde{H}^2(\Gamma_{u,v,\kappa}, \mathbb{k}^\times)$ denote the kernel of the restriction map

$$H^2(\Gamma_{u,v,\kappa}, \mathbb{k}^\times) \rightarrow H^2(\underline{\Lambda}, \mathbb{k}^\times).$$

By [T, Theorem 2 (I)], we have that

$$H^2(\Gamma_{u,v,\kappa}, \mathbb{k}^\times) \simeq H^2(\underline{\Lambda}, \mathbb{k}^\times) \oplus \tilde{H}^2(\Gamma_{u,v,\kappa}, \mathbb{k}^\times).$$

By [T, Theorem 2 (II)], there exists an exact sequence

$$(A.2) \quad 0 \longrightarrow H^1(\underline{\Lambda}, \widehat{\underline{\Omega}}) \longrightarrow \tilde{H}^2(\Gamma_{u,v,\kappa}, \mathbb{k}^\times) \xrightarrow{res} H^2(\underline{\Omega}, \mathbb{k}^\times)^{\underline{\Lambda}}.$$

The image of the first map is the subspace of $H^2(\Gamma_{u,v,\kappa}, \mathbb{k}^\times)$ of 2-cocycles which are cohomologically trivial on $\underline{\Lambda}$ and $\underline{\Omega}$. Next we will characterize $H^1(\underline{\Lambda}, \widehat{\underline{\Omega}})$ and describe the shape of 2-cocycles coming from this group.

Proposition A.5. *Let \mathcal{T} denote the set of triples $(P, \chi, \psi) \in \text{Pair}(\Lambda, \Omega) \times \widehat{\Lambda} \times \widehat{\Omega}$ such that $\psi(v)\psi(\kappa) = \chi(u)$; $P(g, \kappa) = 1$, $\chi(g)^2 = P(g, v)$, $\psi(h)^2 = P(u, h)$, for all $g \in \Lambda, h \in \Omega$.*

(a) *The map $H^1(\underline{\Lambda}, \widehat{\underline{\Omega}}) \rightarrow \mathcal{T}$, $\phi \mapsto (P_\phi, \chi_\phi, \phi(x)|_\Omega)$, where*

$$(A.3) \quad P_\phi(g, h) = \phi(g)(h), \quad \chi_\phi(g) = \phi(g)(y), \quad g \in \Lambda, h \in \Omega,$$

is bijective.

(b) *The image $\sigma \in \tilde{H}^2(\Gamma_{u,v,\kappa}, \mathbb{k}^\times) \subseteq H^2(\Gamma_{u,v,\kappa}, \mathbb{k}^\times)$ of a triple $(P, \chi, \psi) \in \mathcal{T}$ (viewed as an element of $H^1(\underline{\Lambda}, \widehat{\underline{\Omega}})$) under the map ∂ in (A.2) satisfies*

$$\frac{\sigma(g, h)}{\sigma(h, g)} = P(g, h), \quad \frac{\sigma(g, y)}{\sigma(y, g)} = \chi(g), \quad \frac{\sigma(x, h)}{\sigma(h, x)} = \psi(h),$$

for all $g \in \Lambda, h \in \Omega$. In particular we have that

$$\frac{\sigma(g, \kappa)}{\sigma(\kappa, g)} = 1 \text{ for all } g \in \Gamma, \quad \frac{\sigma(\kappa, y)}{\sigma(y, \kappa)} = \chi(\kappa), \quad \frac{\sigma(x, \kappa)}{\sigma(\kappa, x)} = \psi(\kappa).$$

Proof. (a) As Λ acts trivially on $\widehat{\underline{\Omega}}$, each crossed morphism $\phi \in H^1(\underline{\Lambda}, \widehat{\underline{\Omega}})$ restricts on Λ to a homomorphism, and hence to a pairing $\Lambda \times \underline{\Omega} \rightarrow \mathbb{k}^\times$, which we think as a pair $(P, \chi) \in \text{Pair}(\Lambda, \Omega) \times \widehat{\Omega}$ as in (A.3) such that $\chi(g)^2 = P(g, v)$ for all $g \in \Lambda$ (because $y^2 = v$). We set $\chi_\phi := \phi(x)|_\Omega : \Omega \rightarrow \mathbb{k}$. As x acts trivially on Ω , χ is a group homomorphism. Hence we have an injective map

$$H^1(\underline{\Lambda}, \widehat{\underline{\Omega}}) \rightarrow \{(P, \chi, \psi) \in \text{Pair}(\Lambda, \Omega) \times \widehat{\Lambda} \times \widehat{\Omega} \mid \chi(g)^2 = P(g, v), \text{ for all } g \in \Lambda\}.$$

If $\xi := \phi(x)(y) \in \mathbb{k}$, then $\xi^2 = \phi(x)(v) = \chi(v)$ and

$$(A.4) \quad \phi(gx^i)(hy^j) = P(g, h)\psi(g)^j\chi(h)^i\xi^{ij}, \quad g \in \Lambda, h \in \Omega, i, j \in \{0, 1\}.$$

Reciprocally, given a triple (P, χ, ψ) as above, set $\phi : \underline{\Lambda} \rightarrow \widehat{\underline{\Omega}}$ as in A.4. Then ϕ is a crossed homomorphism if and only if for all $g \in \Lambda, h \in \Omega$,

$$\begin{aligned} \phi(gx)(hy) &= \phi(xg)(hy) = \phi(x)(hy)(x \cdot \phi(g)(hy)) = P(g, \kappa h)\psi(g)\chi(h)\xi, \\ \phi(u)(h) &= \phi(x^2)(h) = \phi(x)(h)(x \cdot \phi(x)(h)) = \chi(h)^2, \\ \phi(u)(hy) &= \phi(x^2)(hy) = \phi(x)(hy)(x \cdot \phi(x)(hy)) = \chi(h)^2\xi^2\chi(\kappa). \end{aligned}$$

This means $P(g, \kappa) = 1$ for all $g \in \Lambda$, $\chi(h)^2 = P(u, h)$ and $P(u, h)\psi(u) = P(u, h)\chi(v)\chi(\kappa)$ for all $h \in \Omega$.

(b) This follows by explicit computation of the coboundary map ∂ , see for example the proof of [T, Theorem 2 (II)]. \square

The next result will allow us to reduce the question about the existence of a 2-cocycle just for groups of order a power of 2.

Proposition A.6. *Let $\Gamma = \Gamma_2 \times \Gamma_{\text{odd}}$, where $|\Gamma_2| = 2^n$ for some $n \in \mathbb{N}$, and $|\Gamma_{\text{odd}}|$ is odd. Let $u = u_2\mathbf{u}$, $v = v_2\mathbf{v}$, with $u_2, v_2 \in \Gamma_2$, $\mathbf{u}, \mathbf{v} \in \Gamma_{\text{odd}}$. Then*

$$\Gamma_{u, v, \kappa} \simeq (\Gamma_2)_{u_2, v_2, \kappa} \times \Gamma_{\text{odd}}, \quad H^2(\Gamma_{u, v, \kappa}, \mathbb{k}^\times) \simeq H^2((\Gamma_2)_{u_2, v_2}, \mathbb{k}^\times) \times H^2(\Gamma_{\text{odd}}, \mathbb{k}^\times).$$

Proof. Let $m, n \in \mathbb{N}_0$ be such that $|\mathbf{u}| = 2m + 1$, $|\mathbf{v}| = 2n + 1$. We write \tilde{x}, \tilde{y} for the extra generators of $(\Gamma_2)_{u_2, v_2, \kappa}$ and keep x, y for those in $\Gamma_{u, v, \kappa}$. Then

$$\begin{aligned} \Gamma_{u, v, \kappa} &\rightarrow (\Gamma_2)_{u_2, v_2, \kappa} \times \Gamma_{\text{odd}}, \\ g_2\mathbf{g}x^i y^j &\mapsto g_2\mathbf{u}^{i(m-1)}\mathbf{v}^{j(n-1)}\tilde{x}^i\tilde{y}^j\mathbf{g}, \quad g_2 \in \Gamma_2, \mathbf{g} \in \Gamma_{\text{odd}}, i, j \in \{0, 1\}, \end{aligned}$$

is a group isomorphism. The isomorphism between cohomology groups follows by Künneth's formula. \square

A.2. Non-abelian groups and 2-cocycles. Next we discuss how to apply Propositions A.3 and A.5 to the main classes of groups $\Gamma_{u, v, \kappa}$ that will appear in the proof below in order to obtain the desired 2-cocycles.

A.2.1. If $\Gamma = \mathbb{Z}_2 = \langle \mathbf{k} \rangle$ then $\Gamma_{1, 1, \mathbf{k}}, \Gamma_{1, \mathbf{k}, \mathbf{k}}, \Gamma_{\mathbf{k}, 1, \mathbf{k}}$ are isomorphic to the dihedral group of order 8.⁴

a) We apply Proposition A.5 to $\Gamma_{1, \mathbf{k}, \mathbf{k}}$ with $\Lambda = 1, \Omega = \Gamma, P = 1, \chi = 1, \psi(\mathbf{k}) = -1$; then $H^2(\Gamma_{1, \mathbf{k}, \mathbf{k}}, \mathbb{k}^\times) \simeq \mathbb{Z}_2 = \langle \sigma \rangle$, where

$$\sigma(x, \mathbf{k})\sigma^{-1}(\mathbf{k}, x) = -1, \quad \sigma(y, \mathbf{k})\sigma^{-1}(\mathbf{k}, y) = 1.$$

⁴On the other hand, $\Gamma_{\mathbf{k}, \mathbf{k}, \mathbf{k}}$ is the quaternion group, which has trivial cohomology.

- b) We apply Proposition A.3 to $\Gamma_{1,\mathbf{k},\mathbf{k}}$, where $f(x) = -1$, $f(y) = 1$. If either $w = 1$ or $2 \nmid r$, then σ can be lifted to $\Gamma(\sqrt[r]{w})$; but when $w = \mathbf{k}$, $2 \mid r$, the lift does not exist. As smallest example, $\Gamma(\sqrt[2]{\mathbf{k}})$ is the almost extraspecial group 2^{3+1} , which has cohomology \mathbb{Z}_2^2 : all 2-cocycles are lifts of the trivial 2-cocycle on $\Gamma_{1,\mathbf{k},\mathbf{k}}$ with $f = 1$ and $\mathbf{f} = \pm 1$.
- c) More generally, if there exists a surjective map $\pi : \Gamma \rightarrow \mathbb{Z}_2$ such that $\pi(u) = 1$, $\pi(v) = \mathbf{k} = \pi(\kappa)$, then the pullback of σ from $(\mathbb{Z}_2)_{1,\mathbf{k},\mathbf{k}}$ to $\Gamma_{u,v,\kappa}$ can be lifted to $\Gamma(\sqrt[r]{w})$ either when $\pi(w) = 1$ or $2 \nmid r$. For $\pi(w) = \kappa$, $2 \mid r$, it can also be lifted if there is a character $\mathbf{f} : \Gamma \rightarrow \mathbb{k}^\times$ such that

$$\mathbf{f}(u) \in \mathbb{G}_r, \quad 2 \nmid r / \text{ord}(\mathbf{f}(u)), \quad \mathbf{f}(v) \in \mathbb{G}'_{r/2}, \quad \mathbf{f}(\kappa) = 1.$$

A.2.2. Let $\Gamma = \mathbb{Z}_2 \times \mathbb{Z}_2 = \langle \mathbf{u} \rangle \times \langle \mathbf{k} \rangle$. Reordering generators, the non-trivial possibilities for $\Gamma_{u,v,\kappa}$ are $\Gamma_{\mathbf{u},1,\mathbf{k}}$ and $\Gamma_{\mathbf{u},\mathbf{k},\mathbf{k}}$, which are groups of order 16 with Gap Id 3,4 and Hall-Senior number $\#_{16}9, \#_{16}10$, see [L1, Chapter 7].

- a) We apply Proposition A.5 to $\Gamma_{\mathbf{u},1,\mathbf{k}}$ with $\Lambda = \langle \mathbf{u} \rangle$, $\Omega = \langle \mathbf{k} \rangle$, $P \equiv 1$, $\psi(\mathbf{u}) = \chi(\mathbf{k}) = -1$: we get a 2-cocycle σ such that

$$\frac{\sigma(x, \mathbf{k})}{\sigma(\mathbf{k}, x)} = -1, \quad \frac{\sigma(x, \mathbf{u})}{\sigma(\mathbf{u}, x)} = 1, \quad \frac{\sigma(y, \mathbf{k})}{\sigma(\mathbf{k}, y)} = 1, \quad \frac{\sigma(y, \mathbf{u})}{\sigma(\mathbf{u}, y)} = -1;$$

for the non-trivial choice $P(\mathbf{u}, \mathbf{k}) = -1$ there is no suitable ψ .

For the decomposition $\Lambda = 1$, $\Omega = \Gamma$, we have the non-trivial choice $P = 1$, $\psi = 1$, $\chi(\mathbf{u}) = \chi(\mathbf{k}) = -1$. We obtain a 2-cocycle σ' such that

$$\frac{\sigma'(x, \mathbf{k})}{\sigma'(\mathbf{k}, x)} = -1, \quad \frac{\sigma'(x, \mathbf{u})}{\sigma'(\mathbf{u}, x)} = -1, \quad \frac{\sigma'(y, \mathbf{k})}{\sigma'(\mathbf{k}, y)} = 1, \quad \frac{\sigma'(y, \mathbf{u})}{\sigma'(\mathbf{u}, y)} = 1.$$

Accordingly it is known that $H^2(\#_{16}9, \mathbb{k}^\times) = \mathbb{Z}_2 \times \mathbb{Z}_2$.

- b) We apply Proposition A.3 to $\Gamma_{\mathbf{u},1,\mathbf{k}}$ and the 2-cocycle σ constructed above, where $\mathbf{w} = \mathbf{k}^s \mathbf{u}^t$, $f(x) = (-1)^s$, $f(y) = (-1)^t$, $s, t \in \{0, 1\}$.
- c) Assume that there exists a surjective map $\pi : \Gamma \rightarrow \mathbb{Z}_2 \times \mathbb{Z}_2$ such that $\pi(u) = \mathbf{u}$, $\pi(v) = 1$, $\pi(\kappa) = \mathbf{k}$. The pullback of σ from $(\mathbb{Z}_2 \times \mathbb{Z}_2)_{\mathbf{u},1,\mathbf{k}}$ to $\Gamma_{u,v,\kappa}$ can be lifted to $\Gamma(\sqrt[r]{w})$ if either $\pi(w) = 1$ or $2 \nmid r$. If $\pi(w) = \kappa$, $2 \mid r$, then the 2-cocycle can be lifted to $\Gamma_{u,v,\kappa}$ if there exists $\mathbf{f} \in \widehat{\Gamma}$ such that $\mathbf{f}(\kappa) = 1$ and

$$\mathbf{f}(u) \text{ (resp. } \mathbf{f}(v)) = \xi \in \begin{cases} \mathbb{G}_r, 2 \nmid r / \text{ord}(\xi), & \text{if } s = 1 \text{ (resp. } t = 1), \\ \mathbb{G}'_{r/2}, & \text{if } s = 0 \text{ (resp. } t = 0). \end{cases}$$

A.2.3. Fix $t \in \mathbb{N}$. Set $k = 4t$, $\Gamma = \mathbb{Z}_k \times \mathbb{Z}_k$ with generators \mathbf{u} , \mathbf{v} , and consider $\mathbf{k} := \mathbf{v}^{2t}$. The group $\Gamma_{\mathbf{u},\mathbf{v},\mathbf{k}}$ has order $4k^2$, center Γ , and is presented by generators x, y and relations $x^{2k} = y^{2k} = 1$, $[x, y] = y^k$, where $\mathbf{u} = x^2$, $\mathbf{v} = y^2$, $\mathbf{k} = y^k$.

Moreover $\langle y \rangle \simeq \mathbb{Z}_{2k}$ is a normal subgroup, $\langle x \rangle \simeq \mathbb{Z}_{2k}$ and $\Gamma_{\mathbf{u},\mathbf{v},\mathbf{k}} \simeq \mathbb{Z}_{2k} \times \mathbb{Z}_{2k}$ with action $x \cdot y = y^{k+1}$ (notice that $x^2 \cdot y = y$).

- a) We apply Proposition A.5 to $\Gamma_{\mathbf{u},\mathbf{v},\mathbf{k}}$ with $\Lambda = \langle \mathbf{u} \rangle$, $\Omega = \langle \mathbf{v} \rangle$. Fix $\xi \in \mathbb{G}_k$.

- The pairings $P : \Lambda \times \Omega$ such that $P(\mathbf{u}, \mathbf{k}) = 1$ are given by

$$P(\mathbf{u}, \mathbf{v}) = \xi^{2i} \quad \text{for some } i \in \mathbb{I}_{2t}.$$

- $\chi \in \widehat{\Lambda}$ satisfies $\chi(\mathbf{u})^2 = P(\mathbf{u}, \mathbf{v})$ if and only if $\chi(\mathbf{u}) = p_1 \xi^i$ for some $p_1 \in \{\pm 1\}$. Analogously, $\psi \in \widehat{\Omega}$ is given by $\psi(\mathbf{v}) = p_2 \xi^i$, $p_2 \in \{\pm 1\}$.
- As $\chi(\mathbf{k}) = (-1)^i$, the condition $\psi(\mathbf{v})\psi(\mathbf{k}) = \chi(\mathbf{u})$ always holds when i is even, and for i odd there are two choices since we need $p_1 p_2 = -1$.

Altogether, when i is even we obtain $4t$ different 2-cocycles $\sigma_{ip_1 p_2}$ with $\chi(\mathbf{k}) = 1$, and for i odd we have $2t$ different 2-cocycles $\sigma_{ip_1 p_2}$ with $\chi(\mathbf{k}) = -1$. Set $\sigma := \sigma_{1+-}$.

- b) We apply Proposition A.3 to $\Gamma_{\mathbf{u}, \mathbf{v}, \mathbf{k}}$, σ as constructed above and $\mathbf{w} = \mathbf{u}^s \mathbf{v}^t$: here, $f(x) = \psi(\mathbf{v}^t) = (-\xi)^t$ and $f(y) = \chi(\mathbf{u}^s) = \xi^s$.
- c) More generally, fix a surjective map $\pi : \Gamma \rightarrow \mathbb{Z}_k \times \mathbb{Z}_k$, $u, v, \kappa \in \Gamma$ such that $\pi(u) = \mathbf{u}$, $\pi(v) = \mathbf{v}$, $\pi(\kappa) = \mathbf{k}$, $w = u^s v^t$, $s, t \in \mathbb{I}_k$, and $r \in \mathbb{N}$. The pullback of σ from $(\mathbb{Z}_k \times \mathbb{Z}_k)_{\mathbf{u}, \mathbf{v}, \mathbf{k}}$ to $\Gamma_{u, v, \kappa}$ can be lifted to $\Gamma(\sqrt[r]{w})$ if there exists $\mathbf{f} \in \widehat{\Gamma}$ such that $\mathbf{f}(x)^r = (-\xi)^t$, $\mathbf{f}(y)^r = \xi^s$ and $\mathbf{f}(\kappa) = 1$.

A.3. Proof of Proposition 3.15. We proceed case-by-case. We use the representations $M(g, \chi)$ coming from the Yetter-Drinfeld structure for each group in order to get a triple as in Proposition A.5, which in turn gives a 2-cocycle, then use Proposition A.3 when we need to extend the group accordingly. For each $M(g, \chi)$ we choose a basis given by centralizer coset representatives and give the corresponding matrices: When the matrix is a multiple of the identity we just write the corresponding scalar.

A.3.1. Type α_2 . Here $M = M(g_1, \chi_1) \oplus M(g_2, \chi_2)$, see §5.1.1. The four possibilities for the parity vector $\mathbf{P} = (\chi_1(\kappa), \chi_2(\kappa))$ fall into two orbits under the Weyl groupoid action, namely $(1, 1)$ and $\{(-1, 1), (-1, -1), (-1, 1)\}$. As the first one corresponds to trivial action of κ , we just need to study $(-1, 1)$. Set $\mathbf{a} := \chi_2(g_1^2) = -\chi_1(g_2^2)^{-1}$, $k = \text{ord } \mathbf{a}$. We compute $M(g_1, \chi_1)$, $M(g_2, \chi_2)$, $M(g_1 g_2, \chi_1 \chi_2)$.

	κ	g_1	g_2	$g_1^2 = u$	$g_2^2 = v$
$M(g_1, \chi_1)$	-1	$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$	$\begin{pmatrix} 0 & -\mathbf{a}^{-1} \\ 1 & 0 \end{pmatrix}$	1	$-\mathbf{a}^{-1}$
$M(g_2, \chi_2)$	1	$\begin{pmatrix} 0 & \mathbf{a} \\ 1 & 0 \end{pmatrix}$	$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$	\mathbf{a}	1
$M(g_1 g_2, \chi_1 \chi_2)$	-1	$\begin{pmatrix} 0 & \mathbf{a} \\ 1 & 0 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 \\ -\mathbf{a}^{-1} & 0 \end{pmatrix}$	\mathbf{a}	$-\mathbf{a}^{-1}$

From the actions above we can read off the group G^{\min} explicitly:

$$G^{\min} = \left\langle g_1, g_2, \kappa \mid [g_1, g_2] = \kappa, [g_1, \kappa] = [g_2, \kappa] = (g_1^2 \kappa)^k = (g_2^2)^k = \kappa^2 = 1, \right. \\ \left. \text{if } 2 \mid k \text{ we add } (g_1^2 \kappa)^{k/2} = \kappa \right\rangle.$$

By Proposition A.6 we reduce to 2-groups, so we have three cases:

- $k = 1$, that is $\mathbf{a} = 1$. Then $x^2 = 1, y^2 = \kappa$ and the group G^{\min} is the dihedral group of order 8, see §A.2.1 a).

- $k = 2$, that is $\mathbf{a} = -1$. Here $x^2 = u$ has order 2, $y^2 = v = 1$, $(xy)^2 = \kappa u$. Then G^{\min} is the group $\#_{16}9$ and such 2-cocycle exists, see §A.2.2 a).
- $k = 2^n$, $n \geq 2$. In this case $x^{2^n} = y^{2^n} = 1$ and $y^{2^{n-1}} = \kappa$. Then $|G^{\min}| = 2^{2+2n}$, the center is $\mathbb{Z}_{2^n} \times \mathbb{Z}_{2^n}$, and moreover $G^{\min} \simeq \mathbb{Z}_{2^{n+1}} \rtimes \mathbb{Z}_{2^{n+1}}$ as in A.2.3 a), so there exists such 2-cocycle.

A.3.2. *Type α_3 .* Here $M = M(g_1, \chi_1) \oplus M(g_2, \chi_2) \oplus M(g_3, \chi_3)$, see §5.1.2. As $\chi_1(\kappa) = \chi_3(\kappa)$, there are four choices of $\mathbf{P} = (\chi_i(\kappa))_{i \in \mathbb{I}_3}$, which fall into two orbits under the Weyl groupoid action, namely

$$\{(1, 1, 1)\} \quad \text{and} \quad \{(-1, 1, -1), (-1, -1, -1), (1, -1, 1)\}.$$

Thus we just need to study $\mathbf{P} = (-1, 1, -1)$. We compute the representations $M(g_i, \chi_i)$, which contains the previous case A_2 . Set

$$\mathbf{a} := \chi_2(g_1^2) = -\chi_1(g_2^2)^{-1}, \quad \mathbf{b} := \chi_3(g_1) = \chi_1(g_3)^{-1}, \quad \mathbf{c} = \chi_2(g_1 g_3^{-1}).$$

Hence the action of G on M is given by:

	κ	g_1	g_2	g_3
$M(g_1, \chi_1)$	-1	$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} 0 & -\mathbf{a}^{-1} \\ 1 & 0 \end{pmatrix}$	$\begin{pmatrix} \mathbf{b}^{-1} & 0 \\ 0 & -\mathbf{b}^{-1} \end{pmatrix}$
$M(g_2, \chi_2)$	1	$\begin{pmatrix} 0 & \mathbf{a} \\ 1 & 0 \end{pmatrix}$	$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$	$\begin{pmatrix} 0 & \mathbf{c}\mathbf{a} \\ \mathbf{c} & 0 \end{pmatrix}$
$M(g_3, \chi_3)$	-1	$\begin{pmatrix} \mathbf{b} & 0 \\ 0 & -\mathbf{b} \end{pmatrix}$	$\begin{pmatrix} 0 & -(\mathbf{c}^2\mathbf{a})^{-1} \\ 1 & 0 \end{pmatrix}$	$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$

Using that $g_1 g_2 = \kappa g_2 g_1$, $g_3 g_2 = \kappa g_2 g_3$, $g_1 g_3 = g_3 g_1$, κ is central and $\kappa^2 = 1$, cf. §5.1.2, the subgroup $\tilde{\Gamma}$ generated by⁵

$$u := g_1^2, \quad v := g_2^2, \quad \mathbf{t} := g_3 g_1^{-1} \kappa, \quad \kappa,$$

is contained in $Z(G^{\min})$, and $G^{\min}/\tilde{\Gamma}$ has four elements: $\tilde{\Gamma}$, $g_1 \tilde{\Gamma}$, $g_2 \tilde{\Gamma}$, $g_1 g_2 \tilde{\Gamma}$. From here we check that $G^{\min} \simeq \tilde{\Gamma}_{u,v,\kappa}$, where $x \mapsto g_1$, $y \mapsto g_2$.

Let Γ be the subgroup generated by u , v , κ . Set $N_1 = \text{lcm}(\text{ord } \mathbf{a}, \text{ord } \mathbf{b}^2)$, $N_2 = \text{lcm}(\text{ord } \mathbf{a}, \text{ord } \mathbf{c}^2)$, $N_3 = \text{lcm}(\text{ord } \mathbf{b}, \text{ord } \mathbf{c})$. As $\mathbf{t}^{N_3} = \text{id}$, we can define

$$\begin{aligned} r &:= \min\{s \in \mathbb{I}_{N_3} \mid \mathbf{t}^s \in \Gamma\} \\ &= \min\{s \in \mathbb{I}_{N_3} \mid \exists m \in \mathbb{I}_{N_1}, n \in \mathbb{I}_{N_2} : \mathbf{c}^{2n} = \mathbf{b}^{2m+2r}, \mathbf{a}^n \mathbf{b}^r = 1, \mathbf{a}^m = \mathbf{c}^r\}. \end{aligned}$$

We have that $\tilde{\Gamma} \simeq \Gamma(\sqrt[r]{w})$ for $w = \mathbf{t}^r$. The action of u , v , \mathbf{t} on M_1 , M_2 , M_3 is given, respectively, by the following scalars:

$$(1, \mathbf{a}, \mathbf{b}^{-2}), \quad (\mathbf{a}^{-1}, 1, \mathbf{a}^{-1} \mathbf{c}^{-2}), \quad (\mathbf{b}^{-1}, \mathbf{c}, \mathbf{b}^{-1}).$$

We also set $k_1 = \text{ord } \mathbf{a}$, $k_2 = \text{ord } \mathbf{b}$, $k_3 = \text{ord } \mathbf{c}$. Then

$$\Gamma \simeq \langle \kappa, u, v \mid \kappa^2 = 1, u^{N_1} = 1, v^{N_2} = 1, \kappa = (u^i v)^{k_1/2} \text{ if } 2 \mid k_1, \mathbf{b}^{2i} = \mathbf{c}^{-k_1} \rangle.$$

⁵We set the generators of $\tilde{\Gamma}$ according with generators for the symplectic root system $n = 3$, $r = 1$ in [L2, Thm. 4.5].

As $k_1|N_1, N_2$, there exists a surjective map $\Gamma \twoheadrightarrow \Gamma'$, where Γ' is an abelian group as in §A.3.1, so there exists a 2-cocycle σ' for Γ' such that $\frac{\sigma'(g_2, \kappa)}{\sigma'(\kappa, g_2)} = 1$, $\frac{\sigma'(g_1, \kappa)}{\sigma'(\kappa, g_1)} = -1 = \frac{\sigma'(g_3, \kappa)}{\sigma'(\kappa, g_3)}$. Let σ be the pullback of σ' on Γ : we look for a lift on $G^{\min} \simeq \Gamma(\sqrt[r]{w})_{u, v, \kappa}$, so we look for a character \mathbf{f} as in Proposition A.3.

By Proposition A.6 it is enough to solve the case in which the three k_i are powers of 2. We split in three cases as in §A.3.1:

- $k = 1$, that is $\mathbf{a} = 1$. Either $w = 1$ or else $k_2 > k_3$, $w = \kappa$ and $r = k_2/2 > 1$. In the second case we construct $\mathbf{f} \in \widehat{\Gamma}$ in $M(g_1, \chi_1)^* \otimes M(g_3, \chi_1)$: that is, $\mathbf{f}(u) = \mathbf{b}^2 \in \mathbb{G}'_r$, $\mathbf{f}(v) = \mathbf{c}^{-2} \in \mathbb{G}'_{r/2}$, $\mathbf{f}(\kappa) = 1$. Then we apply §A.2.1 c).
- $k = 2$, that is $\mathbf{a} = -1$. If $k_2 > k_3$, then $r = k_2/2$, $w = \kappa$; if $k_2 < k_3$, then $r = k_3/2$, $w = u$; otherwise $k_2 = k_3$, $r = k_2/2$ and $w = u\kappa$. In any case we construct $\mathbf{f} \in \widehat{\Gamma}$ in $M(g_1, \chi_1)^* \otimes M(g_3, \chi_1)$ as in §A.2.2 c), and there exists such a lift.
- $k = 2^n$, $n \geq 2$. Here $r = \max(k_2/k_1, k_3/k_1, 1)$, and we construct $\mathbf{f} \in \widehat{\Gamma}$, again in $M(g_1, \chi_1)^* \otimes M(g_3, \chi_1)$:

$$\mathbf{f}(\kappa) = 1, \quad \mathbf{f}(u) = \mathbf{b}^2, \quad \mathbf{f}(v) = \mathbf{c}^2.$$

This \mathbf{f} satisfies the conditions in §A.2.3 c), and there exists such a lift.

A.3.3. *Type δ_4 .* As $\chi_1(\kappa) = \chi_3(\kappa) = \chi_4(\kappa)$, the choices of $\mathbf{P} = (\chi_i(\kappa))_{i \in \mathbb{I}_4}$ fall into two orbits under the Weyl groupoid action, namely

$$\{(1, 1, 1, 1)\} \quad \text{and} \quad \{(-1, 1, -1, -1), (-1, -1, -1, -1), (1, -1, 1, 1)\},$$

where the second entry denotes the center node in the Dynkin diagram. Now we study $\mathbf{P} = (-1, 1, -1, -1)$. We fix the central elements ⁶ $z = g_3 g_1^{-1}$, $z' = g_4 g_1^{-1}$. Then this case can be achieved by combining the previous result of extending α_2 to α_3 by z and by z' , see Corollary A.4.

A.3.4. *Type γ_3 .* Here $M = M(g_1, \chi_1) \oplus M(g_2, \chi_2) \oplus M(g_3, \chi_3)$, with $g_3 \in Z(G)$, $\chi_3 \in \widehat{G}$. As $\kappa = [g_1, g_2]$, we have that $\chi_3(\kappa) = 1$. The possible $\mathbf{P} = (\chi_i(\kappa))_{i \in \mathbb{I}_3}$ fall into two orbits under the Weyl groupoid action, namely $\{(1, 1, 1)\}$ and $\{(-1, 1, 1), (-1, -1, 1), (1, -1, 1)\}$. Fix $\mathbf{P} = (-1, 1, 1)$ and set

$$\mathbf{a} := \chi_2(g_1^2) = -\chi_1(g_2^{-2}), \quad \mathbf{b} := \chi_3(g_1) = \chi_1(g_3^{-1}), \quad \mathbf{c} := \chi_3(g_2) = -\chi_2(g_3^{-1}).$$

Now we compute the representations $M(g_i, \chi_i)$:

	κ	g_1	g_2	g_3
$M(g_1, \chi_1)$	-1	$\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$	$\begin{pmatrix} 0 & -\mathbf{a}^{-1} \\ 1 & 0 \end{pmatrix}$	\mathbf{b}^{-1}
$M(g_2, \chi_2)$	+1	$\begin{pmatrix} 0 & \mathbf{a} \\ 1 & 0 \end{pmatrix}$	$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$	$-\mathbf{c}^{-1}$
$M(g_3, \chi_3)$	+1	\mathbf{b}	\mathbf{c}	-1

⁶According with generators for the symplectic root system $n = 4$, $r = 2$ in [L2].

If $u = g_1^2$, $v = g_2^2$, $\mathfrak{t} = g_3$, then $G^{\min} \simeq \Gamma(\sqrt[r]{w})_{u,v,\kappa}$ for $\Gamma = \langle u, v, \kappa \rangle$ and appropriate $w \in \Gamma$, $r \in \mathbb{N}$. Moreover G^{\min} is isomorphic to the one in §A.3.2, hence there exists a 2-cocycle as in Proposition 3.15.

A.3.5. *Type γ_4 .* Here we have central elements $z = g_3 g_1^{-1}$, $z' = g_4$, and this case is solved by combining the previous result of extending A_2 to A_3 by z and by z' , see Corollary A.4.

A.3.6. *Type ϕ_4 .* Here $M = M(g_1, \chi_1) \oplus M(g_2, \chi_2) \oplus M(g_3, \chi_3) \oplus M(g_4, \chi_4)$, with $g_3, g_4 \in Z(G)$, $\chi_3, \chi_4 \in \widehat{G}$, and $\chi_3(\kappa) = \chi_4(\kappa) = 1$. The possible $\mathbf{P} = (\chi_i(\kappa))_{i \in \mathbb{I}_4}$ fall into two orbits under the Weyl groupoid action: $\{(1, 1, 1, 1)\}$ and $\{(-1, 1, 1, 1), (-1, -1, 1, 1), (1, -1, 1, 1)\}$. Fix $\mathbf{P} = (-1, 1, 1, 1)$ and set

$$\begin{aligned} \mathbf{a} &:= \chi_2(g_1^2) = -\chi_1(g_2^{-2}), & \mathbf{b} &:= \chi_3(g_1) = \chi_1(g_3^{-1}), & \mathbf{c} &:= \chi_3(g_2) = -\chi_2(g_3^{-1}), \\ \mathbf{b}' &:= \chi_4(g_1) = \chi_1(g_4^{-1}), & \mathbf{c}' &:= \chi_4(g_2) = \chi_2(g_4^{-1}), & \mathbf{d} &:= \chi_4(g_3) = -\chi_3(g_2^{-1}). \end{aligned}$$

Now we compute the representations $M_i := M(g_i, \chi_i)$:

	κ	g_1	g_2	g_3	g_4
$M(g_1, \chi_1)$	-1	$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$	$\begin{pmatrix} 0 & -\mathbf{a}^{-1} \\ 1 & 0 \end{pmatrix}$	\mathbf{b}^{-1}	\mathbf{b}'^{-1}
$M(g_2, \chi_2)$	1	$\begin{pmatrix} 0 & \mathbf{a} \\ 1 & 0 \end{pmatrix}$	$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$	$-\mathbf{c}^{-1}$	\mathbf{c}'^{-1}
$M(g_3, \chi_3)$	1	\mathbf{b}	\mathbf{c}	-1	$-\mathbf{d}^{-1}$
$M(g_4, \chi_4)$	1	\mathbf{b}'	\mathbf{c}'	\mathbf{d}	-1

Set $u = g_1^2$, $v = g_2^2$. We will construct a 2-cocycle σ on G^{\min} by using appropriate 2-cocycles from the previous cases:

- Set $G_{12} = \langle g_1, g_2 \rangle$, $\Gamma = \langle \kappa, u, v \rangle$; let $\widetilde{G}_{12}, \Gamma_{12} \subset \text{End}(M_1 \oplus M_2)$ be the subgroups obtained by restriction. Then Γ, Γ_{12} are central subgroups, $G_{12} = \Gamma_{u,v,\kappa}$, $\widetilde{G}_{12} = (\Gamma_{12})_{u,v,\kappa}$, with canonical projections $G_{12} \twoheadrightarrow \widetilde{G}_{12}$, $\Gamma \twoheadrightarrow \Gamma_{12}$, and $\widetilde{G}_{12}, \Gamma_{12}$ are as in §A.3.1. Hence there exists a 2-cocycle as we need: the pullback σ_{12} on G_{12} satisfies $\frac{\sigma_{12}(g_1, \kappa)}{\sigma_{12}(\kappa, g_1)} = -1$, $\frac{\sigma_{12}(g_2, \kappa)}{\sigma_{12}(\kappa, g_2)} = 1$.
- For $j = 3, 4$ we set $G_{12j} = \langle g_1, g_2, g_j \rangle$, $\Gamma_{12j} = \langle \kappa, u, v, g_j \rangle$. Then Γ_{12j} is a central subgroup of the form $\Gamma_{12j} \simeq \Gamma(\sqrt[r_j]{w_j})$ for appropriate $r_j \in \mathbb{N}$, $w_j \in \Gamma$: the proof for $j = 3$ is the same as in §A.3.4 since $M_1 \oplus M_2 \oplus M_3$ is of type C_3 , and for $j = 4$ we have the same structure (luckily). Using the same argument as in §A.3.4 we check the existence of 2-cocycle σ_{12j} on G_{12j} such that $\frac{\sigma_{12j}(g_1, \kappa)}{\sigma_{12j}(\kappa, g_1)} = -1$, $\frac{\sigma_{12j}(g_2, \kappa)}{\sigma_{12j}(\kappa, g_2)} = \frac{\sigma_{12j}(g_j, \kappa)}{\sigma_{12j}(\kappa, g_j)} = 1$.
- Finally, $G^{\min} \simeq \Gamma(\sqrt[r_3]{w_3})(\sqrt[r_4]{w_4})_{u,v,\kappa} \simeq \Gamma(\sqrt[r_4]{w_4})\Gamma(\sqrt[r_3]{w_3})_{u,v,\kappa}$. The existence of a 2-cocycle σ on G^{\min} such that

$$\frac{\sigma(g_1, \kappa)}{\sigma(\kappa, g_1)} = -1, \quad \frac{\sigma(g_2, \kappa)}{\sigma(\kappa, g_2)} = \frac{\sigma(g_3, \kappa)}{\sigma(\kappa, g_3)} = \frac{\sigma(g_4, \kappa)}{\sigma(\kappa, g_4)} = 1$$

follows from the 2-cocycles σ_{12j} on G_{12j} , $j = 3, 4$ and Corollary A.4.

This concludes the proof of Proposition 3.15. \square

Acknowledgements. The authors thank Istvan Heckenberger for many useful discussions, suggestions and hospitality. In particular, for the discussion of Lemma 2.7.

REFERENCES

- [A] N. Andruskiewitsch. *An Introduction to Nichols Algebras*. In Quantization, Geometry and Noncommutative Structures in Mathematics and Physics. A. Cardona, P. Morales, H. Ocampo, S. Paycha, A. Reyes, eds., pp. 135–195, Springer (2017).
- [AA] N. Andruskiewitsch, I. Angiono. *On finite-dimensional Nichols algebras of diagonal type*. Bull. Math. Sci. **7** (2017), 353–573.
- [A+] N. Andruskiewitsch, I. Angiono, A. García Iglesias, A. Masuoka, C. Vay. *Lifting via cocycle deformation*. J. Pure Appl. Alg. **218** (2014), 684–703.
- [AAG] N. Andruskiewitsch, I. Angiono, A. García Iglesias. *Liftings of Nichols algebras of diagonal type I. Cartan type A*. Int. Math. Res. Not. IMRN **2017** (9) (2017), 2793–2884.
- [ACG] N. Andruskiewitsch, G. Carnovale, G. García. *Finite-dimensional pointed Hopf algebras over finite simple groups of Lie type V. Mixed classes in Chevalley and Steinberg groups*. Manuscripta Math., to appear.
- [AG] N. Andruskiewitsch, M. Graña, *From racks to pointed Hopf algebras*. Adv. Math. **178** (2003), 177–243.
- [AS1] N. Andruskiewitsch, H.-J. Schneider. *Finite quantum groups and Cartan matrices*, Adv. Math. **154** (2000), 1–45.
- [AS2] ———. *Pointed Hopf algebras*. In Recent developments in Hopf algebras Theory, MSRI Publ. **43** (2002), 1–68, Cambridge Univ. Pr.
- [AS3] ———. *On the classification of finite-dimensional pointed Hopf algebras*, Ann. of Math. **171** (2010), 375–417.
- [An1] I. Angiono, *On Nichols algebras with standard braiding*, Algebra Number Theory **1** (2009), 35–106.
- [An2] ———. *On Nichols algebras of diagonal type*. J. Reine Angew. Math. **683** (2013), 189–251.
- [An3] ———. *A presentation by generators and relations of Nichols algebras of diagonal type and convex orders on root systems*. J. Eur. Math. Soc. **17** (2015), 2643–2671.
- [An4] ———. *Distinguished pre-Nichols algebras*. Transf. Groups **21** (2016), 1–33.
- [ACS] I. Angiono, E. Campagnolo and G. Sanmarco. *Finite GK-dimensional pre-Nichols algebras of super and standard type*, arXiv:2009.04863.
- [AnG] I. Angiono and A. García Iglesias. *Liftings of Nichols algebras of diagonal type II. All liftings are cocycle deformations*. Selecta Math. **25** (2019), Paper No. 5, 95 pp.
- [AnGP] I. Angiono, C. Galindo, M. Pereira. *De-equivariantization of Hopf algebras*. Algebr. Represent. Theory **17** (2014), 161–180.
- [AnKM] I. Angiono, M. Kochetov, M. Mastnak. *On rigidity of Nichols algebras*. J. Pure Appl. Algebra **219** (2015), 5539–5559.
- [AnS] I. Angiono, G. Sanmarco. *Pointed Hopf algebras over non-abelian groups with decomposable braidings. I*. J. Algebra **549** (2020), 78–111.
- [AJS] H.H. Andersen, J.C. Jantzen, W. Soergel, *Representations of quantum groups at a p th root of unity and of semisimple groups in characteristic p : independence of p* . Astérisque **220** (1994), 321.
- [CH] M. Cuntz, I. Heckenberger, *Finite Weyl groupoids of rank three*. Trans. Amer. Math. Soc. **364** (2012), 1369–1393.
- [DoT] Y. Doi, M. Takeuchi, *Multiplication alteration by two-cocycles - the quantum version*, Comm. Algebra **22** (1994), 5715–5732.

- [Dr] V. Drinfeld, *Quantum groups*, Proc. Int. Congr. Math., Berkeley 1986, Vol. **1** (1987), 798–820.
- [EGNO] P. Etingof, S. Gelaki, D. Nikshych, V. Ostrik, *Tensor categories*. Mathematical Surveys and Monographs **205**, AMS, Providence, RI, 2015, xvi+343pp.
- [ENOM] P. Etingof, D. Nikshych, V. Ostrik, E. Meir, *Fusion categories and homotopy theory*. Quantum Topol. **1** (2010), 209–273.
- [FBZ] E. Frenkel, D. Ben-Zvi, *Vertex Algebras and Algebraic Curves*, Mathematical Surveys and Monographs 88, American Mathematical Society (2004).
- [FGST] B. Feigin, A. Gainutdinov, A. Semikhatov, I. Tipunin, *Kazhdan-Lusztig correspondence for the representation category of the triplet W -algebra in logarithmic CFT*, Theor. Math. Phys. **148** (2006).
- [FK] S. Fomin, K. N. Kirillov, *Quadratic algebras, Dunkl elements, and Schubert calculus*. Progr. Math. **172** (1999), 146–182.
- [Fu] J. Fuchs, *Affine Lie algebras and quantum groups*, Cambridge Monographs Math. Phys. (1994).
- [Gr] M. Graña, *On Nichols algebras of low dimension*, Contemp. Math. **267** (2000).
- [Gu] R. Günther, *Crossed products for pointed Hopf algebras*. Comm. Algebra, **27** (1999), 4389–4410.
- [H1] I. Heckenberger. *The Weyl groupoid of a Nichols algebra of diagonal type*. Invent. Math. **164** (2006), 175–188 .
- [H2] ——— *Classification of arithmetic root systems*. Adv. Math. **220** (2009), 59–124 .
- [HLV] I. Heckenberger, A. Lochmann, L. Vendramin, *Braided racks, Hurwitz actions and Nicholsalgebras with many cubic relations*, Transform. Groups **17** (2012), 157–194.
- [HS1] I. Heckenberger, H.-J. Schneider. *Nichols algebras over groups with finite root system of rank two I*. J. Algebra **324** (2010), 3090–3114.
- [HS2] I. Heckenberger, H.-J. Schneider. *Right coideal subalgebras of Nichols algebras and the Duflo order on the Weyl groupoid*. Isr. J. Math. **197** (2013), 139–187.
- [HS3] ——— *Hopf algebras and root systems*. Mathematical Surveys and Monographs **247**. Providence, RI: American Mathematical Society (AMS) (ISBN 978-1-4704-5232-2/hbk; 978-1-4704-5680-1/ebook). xix, 582 p. (2020).
- [HV] I. Heckenberger, L. Vendramin. *A classification of Nichols algebras of semi-simple Yetter-Drinfeld modules over non-abelian groups*. J. Eur. Math. Soc. **19** (2017), 299–356.
- [HeY] I. Heckenberger, H. Yamane, *A generalization of Coxeter groups, root systems, and Matsumoto’s theorem*, Math. Z. **259** (2008), 255–276.
- [IM] N. Iwahori, H. Matsumoto, *Several remarks on projective representations of finite groups*. J. Faculty of Science Univ. Tokyo **10** (1964), 129–146.
- [J] M. Jimbo, *A q -difference analogue of $U(\mathfrak{g})$ and the Yang Baxter equation*, Lett. Math. Phys. **10** (1985), 63–69.
- [KL] T. Kerler, V.V. Lyubashenko, *Non-Semisimple Topological Quantum Field Theories for 3-Manifolds with Corners*, Springer Lect. Notes Math., vol. 1765, Springer-Verlag, New York (2001).
- [L1] S. Lentner, *Orbifoldizing Hopf and Nichols Algebras*. Dissertation, LMU München (2012), <https://edoc.ub.uni-muenchen.de/15363/>.
- [L2] S. Lentner, *New large-rank Nichols algebras over non-abelian groups with commutator subgroup \mathbb{Z}_2* , J. Algebra **419** (2014), 1–33.
- [L3] S.D. Lentner, *Quantum groups and Nichols algebras acting on conformal field theories*, Adv. Math., to appear.

- [Lu1] G. Lusztig, *Modular representations and quantum groups*. Contemp. Math. **82** 59–77, Amer. Math. Soc., Providence, RI, 1989.
- [Lu2] G. Lusztig, *Introduction to quantum groups*. Birkhäuser (1993).
- [Ma] A. Masuoka, *Abelian and non-abelian second cohomologies of quantized enveloping algebras*, J. Algebra **320** (2008), 1–47.
- [Me] E. Meir, *Geometric perspective on Nichols algebras*. J. Algebra **601** (2022), 390–422.
- [MiS] A. Milinski, H.-J. Schneider, *Pointed Indecomposable Hopf Algebras over Coxeter Groups*. Contemp. Math. **267** (2000), 215–236.
- [Mm] M. Mombelli, *The Brauer-Picard group of the representation category of finite supergroup algebras*. Rev. Unión Mat. Argent. **55** (2014), 83–117.
- [Mo] S. Montgomery. *Hopf algebras and their actions on rings*, CMBS **82**, Amer. Math. Soc. (1993).
- [T] K. Tahara, *On the Second Cohomology Groups of Semidirect Products*, Math. Z. **129** (1972), 365–379.
- [Tur] V. Turaev, *Quantum invariants of knots and 3-manifolds*, de Gruyter Studies in Mathematics 18 Walter de Gruyter & Co (1994).

FACULTAD DE MATEMÁTICA, ASTRONOMÍA, FÍSICA Y COMPUTACIÓN, UNIVERSIDAD NACIONAL DE CÓRDOBA. CIEM – CONICET. MEDINA ALLENDE S/N (5000) CIUDAD UNIV., CÓRDOBA, ARGENTINA

Email address: `angiono@famaf.unc.edu.ar`

ALGEBRA AND NUMBER THEORY. UNIVERSITY OF HAMBURG, BUNDESSTRASSE 55, 20146 HAMBURG

Email address: `simon.lentner@uni-hamburg.de`

DEPARTMENT OF MATHEMATICS, IOWA STATE UNIVERSITY, AMES, IA 50011, USA

Email address: `sanmarco@iastate.edu`