

Classical and quantum bicosmology with noncommutativity

Nahomi Kan*

National Institute of Technology, Gifu College, Motosu-shi, Gifu 501-0495, Japan

Takuma Aoyama[†] and Kiyoshi Shiraishi[‡]

*Graduate School of Sciences and Technology for Innovation,
Yamaguchi University, Yamaguchi-shi, Yamaguchi 753-8512, Japan*

(Dated: June 23, 2022)

Recently, Falomir, Gamboa, Méndez, Gondolo and Maldonado proposed a bicosmology scenario for solving some cosmological problems. Their trick is to introduce noncommutativity into the momentum space of the two scale factors. In the present paper, we revisit their model and firstly we consider exact classical solutions in a model with constant noncommutativity between dynamical variables as well as between canonical momenta. Further, we propose a hypothesis that the noncommutativity appears if the scale factors are small, and illustrate the behavior of the classical solution in that case with momentum-space noncommutativity. Finally, we write down the Wheeler–DeWitt equation in that case and investigate the behavior of the solution.

PACS numbers: 04.20.Fy, 04.20.Jb, 04.60.Kz, 45.20.Jj, 98.80.Qc, 98.80.Jk.

* kan@gifu-nct.ac.jp

† b014vbv@yamaguchi-u.ac.jp

‡ shiraish@yamaguchi-u.ac.jp

I. INTRODUCTION

The bicosmology scenario has been proposed by Falomir, Gamboa, Méndez, Gondolo and Maldonado [1–4]. In their scenario, two scale factors are introduced to represent two causally separated regions. From a different point of view, it can be regarded as a kind of cosmological models in bigravity theory (for example, see Refs. [5–7]). However, instead of the mass term by mixing of two scale factors in bigravity, mixing is guaranteed to be due to the noncommutativity in the Hamiltonian formulation of the classical bicosmology.

Noncommutative cosmology has been studied by many authors [8–27], not only classically but also quantum mechanically. Thus, we intend to study various aspects of classical and quantum bicosmology.

In the present paper, we revisit the model of bicosmology and examine classical and quantum cosmological solutions. Although the main analyses in later sections are illustrated when canonical momenta are noncommutative, we also consider noncommutativity in dynamical variables in the minisuperspace at first.

Next, we observe the case where the parameter representing the noncommutativity is not constant. The most natural idea is that noncommutativity is almost ineffective on cosmological evolution when the universe is large, but is highly effective at the very early stage of the universe, when the universe is small. We also write down the Wheeler–DeWitt (WDW) equation in that case and investigate the behavior of the solution.

In Sec. II, we describe the setting of our model of bicosmology and obtain classical solutions for a commutative case. In Sec. III, we study classical bicosmology with noncommutativity in the phase space. In Sec. IV, we define the localized noncommutativity in momentum space and investigate classical solutions in the system. The quantum bicosmology by the WDW equation in the system is studied in Sec. V. Finally, conclusion and discussion are given in the last section.

II. (COMMUTATIVE) BICOSMOLOGY MODEL: WARM-UP

We consider the following action consisting only of the Einstein–Hilbert term and the cosmological term:

$$S \propto \int d^D x \sqrt{-g} [R - 2\Lambda], \quad (2.1)$$

where R is the Ricci scalar derived from the metric $g_{\mu\nu}$ ($\mu, \nu = 0, 1, \dots, D-1$), and g is the determinant of $g_{\mu\nu}$. We also assumed that the cosmological constant Λ is a positive constant. As a first step in an analysis, it is great importance to investigate cases where such a toy model has exact solutions. It will be also useful for consideration on numerical analyses of elaborate models with further variations in future.

First, in this section, let us review a commutative version of bicosmology and use it for comparison with the noncommutative case later.

We take an ansatz for the metric tensor as

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = -N^2 dt^2 + a^2(t) d\mathbf{x}^2, \quad (2.2)$$

where $t = x^0$, $d\mathbf{x}^2 \equiv \sum_{\mu=i}^{D-1} (dx^i)^2$, and N stands for the lapse function. Then, one can derive the following Lagrangian for the scale factor from the action (2.1):

$$L = -\frac{1}{2N} a^{\frac{D-3}{2}} \dot{a}^2 - \frac{N}{(D-1)(D-2)} \Lambda a^{D-1}, \quad (2.3)$$

where the dot indicates the derivative with respect to time t and total derivatives have been dropped. Now, we switch the dynamical variable a to x , which is defined as

$$x \equiv \frac{2}{D-1} a^{\frac{D-1}{2}}. \quad (2.4)$$

Then, the Lagrangian takes the form

$$L = -\frac{1}{2N} \dot{x}^2 - \frac{N}{2} \gamma^2 x^2. \quad (2.5)$$

where

$$\gamma^2 = \frac{D-1}{2(D-2)} \Lambda. \quad (2.6)$$

Note that this Lagrangian is just that of an inverted harmonic oscillator.

In the bicosmology scenario [1–4], the second scale factor b , which represents the causally disconnected region, is introduced. Here, we adopt the case that the Lagrangian for the second scale factor b is just a copy of that for the first scale factor a , though the authors of Refs. [1–4] considered generally different values for cosmological constants. Therefore, the total Lagrangian takes the form of the two-dimensional inverted harmonic oscillator:

$$L = -\frac{1}{2N} (\dot{x}^2 + \dot{y}^2) - \frac{N}{2} \gamma^2 (x^2 + y^2), \quad (2.7)$$

where

$$y = \frac{2}{D-1} b^{\frac{D-1}{2}}. \quad (2.8)$$

To construct the Hamiltonian, we have only to follow the standard way. The conjugate momenta are found to be

$$\pi_x \equiv \frac{\partial L}{\partial \dot{x}} = -\frac{\dot{x}}{N}, \quad \pi_y \equiv \frac{\partial L}{\partial \dot{y}} = -\frac{\dot{y}}{N}, \quad (2.9)$$

and the Hamiltonian is finally obtained as

$$H = N \left[-\frac{1}{2}(\pi_x^2 + \pi_y^2) + \frac{1}{2}\gamma^2(x^2 + y^2) \right]. \quad (2.10)$$

The Hamiltonian constraint is assigned from $\frac{\partial H}{\partial N} = 0$. We thus set $N = 1$ hereafter.

The Poisson brackets are placed as usual:

$$\{x, \pi_x\} = \{y, \pi_y\} = 1, \quad (2.11)$$

and others are zero. Then, the usual Hamilton's equations are obtained as

$$\begin{aligned} \dot{x} = \{x, H\} &= \frac{\partial H}{\partial \pi_x} = -\pi_x, & \dot{y} = \{y, H\} &= \frac{\partial H}{\partial \pi_y} = -\pi_y, \\ \dot{\pi}_x = \{\pi_x, H\} &= -\frac{\partial H}{\partial x} = -\gamma^2 x, & \dot{\pi}_y = \{\pi_y, H\} &= -\frac{\partial H}{\partial y} = -\gamma^2 y, \end{aligned} \quad (2.12)$$

and subsequently, we can derive the equations of motion as

$$\ddot{x} - \gamma^2 x = 0, \quad \ddot{y} - \gamma^2 y = 0. \quad (2.13)$$

The solution of the equations is

$$x(t) = C_1 \cos \varphi_0 \exp[\gamma t] + C_2 \sin \varphi_0 \exp[-\gamma t], \quad (2.14)$$

$$y(t) = C_1 \sin \varphi_0 \exp[\gamma t] - C_2 \cos \varphi_0 \exp[-\gamma t], \quad (2.15)$$

where the arrangement of the constant coefficients C_1 , C_2 , and the phase angle φ_0 have been chosen as the Hamiltonian constraint $H = 0$ holds.

Now, a warm-up exercise is over. In the next section, we consider classical noncommutative bicosmology.

III. CLASSICAL NONCOMMUTATIVE BICOSMOLOGY

The authors of bicosmology [1–4] introduced noncommutativity into the momentum space of the system. We first consider noncommutative dynamical variables as well.

We presume that the corresponding noncommutative system is defined by replacement¹

$$x \rightarrow X, \quad \pi_x \rightarrow \Pi_X, \quad y \rightarrow Y, \quad \pi_y \rightarrow \Pi_Y, \quad (3.1)$$

and the Hamiltonian takes the form

$$H_{NC} = -\frac{1}{2}(\Pi_X^2 + \Pi_Y^2) + \frac{1}{2}\gamma^2(X^2 + Y^2). \quad (3.2)$$

Here, we assume the noncommutative phase space in the minisuperspace, which is realized by the deformed Poisson brackets

$$\{X, \Pi_X\} = \{Y, \Pi_Y\} = 1, \quad \{X, Y\} = \Theta, \quad \{\Pi_X, \Pi_Y\} = B, \quad (3.3)$$

and others are zero. We assume the parameters which represent noncommutativities, Θ and B , are constant for this time.

Now, the canonical equations can be obtained as

$$\begin{aligned} \dot{X} &= \{X, H_{NC}\} = -\Pi_X + \gamma^2\Theta Y, & \dot{Y} &= \{Y, H_{NC}\} = -\Pi_Y - \gamma^2\Theta X, \\ \dot{\Pi}_X &= \{\Pi_X, H_{NC}\} = -\gamma^2 X - B\Pi_Y, & \dot{\Pi}_Y &= \{\Pi_Y, H_{NC}\} = -\gamma^2 Y + B\Pi_X, \end{aligned} \quad (3.4)$$

and consequently, the equations of motion for X and Y can be revealed as

$$\ddot{X} - (\gamma^2\Theta - B)\dot{Y} - \gamma^2(1 - B\Theta)X = 0, \quad \ddot{Y} + (\gamma^2\Theta - B)\dot{X} - \gamma^2(1 - B\Theta)Y = 0. \quad (3.5)$$

These equations can be easily solved, especially if we considered a combination

$$\ddot{Z} + i(\gamma^2\Theta - B)\dot{Z} - \gamma^2(1 - B\Theta)Z = 0, \quad (3.6)$$

where $Z \equiv X + iY$.

The general solutions which satisfy the Hamiltonian constraint $H_{NC} = 0$ are exhibited as follows.

¹ Note that the usage of letters may be slightly different from the work so far in bicosmology [1–4].

- if $|B + \gamma^2\Theta| < 2\gamma$

$$X(t) = C_1 \exp\left[\frac{\sqrt{4\gamma^2 - (\gamma^2\Theta + B)^2}}{2}t\right] \cos\left[\frac{B - \gamma^2\Theta}{2}t + \varphi_1\right] \\ + C_2 \exp\left[-\frac{\sqrt{4\gamma^2 - (\gamma^2\Theta + B)^2}}{2}t\right] \sin\left[\frac{B - \gamma^2\Theta}{2}t + \varphi_2\right], \quad (3.7)$$

$$Y(t) = C_1 \exp\left[\frac{\sqrt{4\gamma^2 - (\gamma^2\Theta + B)^2}}{2}t\right] \sin\left[\frac{B - \gamma^2\Theta}{2}t + \varphi_1\right] \\ - C_2 \exp\left[-\frac{\sqrt{4\gamma^2 - (\gamma^2\Theta + B)^2}}{2}t\right] \cos\left[\frac{B - \gamma^2\Theta}{2}t + \varphi_2\right], \quad (3.8)$$

where C_1 , C_2 , φ_1 , and φ_2 are constants. The two phase angles should satisfy

$$\sin(\varphi_2 - \varphi_1) = \frac{B + \gamma^2\Theta}{2\gamma}, \quad (3.9)$$

for the Hamiltonian constraint.

- if $|B + \gamma^2\Theta| = 2\gamma$

$$X(t) = C \cos\left[\frac{B - \gamma^2\Theta}{2}t + \varphi_1\right] \\ + C(B + \gamma^2\Theta) \sin(\varphi_1 - \varphi_2)t \cos\left[\frac{B - \gamma^2\Theta}{2}t + \varphi_2\right], \quad (3.10)$$

$$Y(t) = C \sin\left[\frac{B - \gamma^2\Theta}{2}t + \varphi_1\right] \\ + C(B + \gamma^2\Theta) \sin(\varphi_1 - \varphi_2)t \sin\left[\frac{B - \gamma^2\Theta}{2}t + \varphi_2\right], \quad (3.11)$$

where C , φ_1 , and φ_2 are constants.

- if $|B + \gamma^2\Theta| > 2\gamma$

$$X(t) = Ce^{-\eta} \cos\left[\frac{B - \gamma^2\Theta + \sqrt{(B + \gamma^2\Theta)^2 - 4\gamma^2}}{2}t + \varphi_1\right] \\ + Ce^{\eta} \sin\left[\frac{B - \gamma^2\Theta - \sqrt{(B + \gamma^2\Theta)^2 - 4\gamma^2}}{2}t + \varphi_2\right], \quad (3.12)$$

$$Y(t) = Ce^{-\eta} \sin\left[\frac{B - \gamma^2\Theta + \sqrt{(B + \gamma^2\Theta)^2 - 4\gamma^2}}{2}t + \varphi_1\right] \\ - Ce^{\eta} \cos\left[\frac{B - \gamma^2\Theta - \sqrt{(B + \gamma^2\Theta)^2 - 4\gamma^2}}{2}t + \varphi_2\right], \quad (3.13)$$

where C , φ_1 , and φ_2 are constants. The constant η is determined by

$$\cosh(2\eta) = \frac{|B + \gamma^2\Theta|}{2\gamma}. \quad (3.14)$$

All of these noncommutative classical solutions are always accompanied by trigonometric functions, and each scale factor becomes zero at a certain time in the past or the future. The

case with $B = \gamma^2\Theta$ is interesting, since there is no slow oscillatory behavior, but we naturally think that B and $\gamma^2\Theta$ do not take so large values at present, and they are expected to be much less than H_0 (the Hubble constant), so they do not affect the present cosmic expansion. Therefore, we have not to think the relation $B \sim \gamma^2\Theta$ so seriously. It is interesting, however, that the parameters of the noncommutativity may have been large only in the early stage of the small universe. This possibility will be investigated in the next section.

By the way, $X^2 + Y^2$ in each case is found to be

$$X^2 + Y^2 = C_1^2 \exp\left[\sqrt{4\gamma^2 - (\gamma^2\Theta + B)^2}t\right] + C_2^2 \exp\left[-\sqrt{4\gamma^2 - (\gamma^2\Theta + B)^2}t\right] + 2C_1C_2 \frac{B + \gamma^2\Theta}{2\gamma}, \quad (|B + \gamma^2\Theta| < 2\gamma) \quad (3.15)$$

$$X^2 + Y^2 = C^2 \left[1 + 2(B + \gamma^2\Theta) \sin(\varphi_1 - \varphi_2) \cos(\varphi_1 - \varphi_2)t + (B + \gamma^2\Theta)^2 \sin^2(\varphi_1 - \varphi_2)t^2\right], \quad (|B + \gamma^2\Theta| = 2\gamma) \quad (3.16)$$

$$X^2 + Y^2 = 2C^2 \left[\frac{|B + \gamma^2\Theta|}{2\gamma} - \sin[\sqrt{(B + \gamma^2\Theta)^2 - 4\gamma^2}t]\right]. \quad (|B + \gamma^2\Theta| > 2\gamma) \quad (3.17)$$

It turns out that $X^2 + Y^2$ does not oscillate slowly, except for a rapid oscillation in the case of $|B + \gamma^2\Theta| > 2\gamma$, so it can at least be a measure of the average size of the two scale factors, although its physical meaning needs to be discussed in later sections.

In the rest of this section, we write down the formulation of noncommutative system using undeformed canonical variables. Similar formulation is already known as a necessary technique for quantum cosmology, i.e., obtaining the WDW equation [8–27].

To this end, we set the variables as follows:

$$X = \frac{1}{\sqrt{1 + \frac{\beta\theta}{4}}}\left(x - \frac{\theta}{2}\pi_y\right), \quad Y = \frac{1}{\sqrt{1 + \frac{\beta\theta}{4}}}\left(y + \frac{\theta}{2}\pi_x\right), \\ \Pi_X = \frac{1}{\sqrt{1 + \frac{\beta\theta}{4}}}\left(\pi_x + \frac{\beta}{2}y\right), \quad \Pi_Y = \frac{1}{\sqrt{1 + \frac{\beta\theta}{4}}}\left(\pi_y - \frac{\beta}{2}x\right), \quad (3.18)$$

where x , π_x , y , and π_y obey the usual Poisson brackets (2.11). If the constant parameters θ and β are related to Θ and B as

$$\Theta = \frac{\theta}{1 + \frac{\beta\theta}{4}}, \quad B = \frac{\beta}{1 + \frac{\beta\theta}{4}}, \quad (3.19)$$

one can find that the brackets for X , Π_X , Y , Π_Y defined by (3.18) realize the noncommutative Poisson brackets (3.3). Then, the Hamiltonian H_{NC} is expressed as

$$H_{NC} = -\frac{1 - \frac{\gamma^2\theta^2}{4}}{2\left(1 + \frac{\beta\theta}{4}\right)}(\pi_x^2 + \pi_y^2) + \frac{\beta - \gamma^2\theta}{2\left(1 + \frac{\beta\theta}{4}\right)}(x\pi_y - y\pi_x) + \frac{\gamma^2 - \frac{\beta^2}{4}}{2\left(1 + \frac{\beta\theta}{4}\right)}(x^2 + y^2). \quad (3.20)$$

The equivalence of the equations motion can be checked easily. Since one can see

$$\begin{aligned}
\dot{x} &= \frac{\partial H_{NC}}{\partial \pi_x} = -\frac{1 - \frac{\gamma^2 \theta^2}{4}}{1 + \frac{\beta \theta}{4}} \pi_x - \frac{\beta - \gamma^2 \theta}{2(1 + \frac{\beta \theta}{4})} y, \\
\dot{y} &= \frac{\partial H_{NC}}{\partial \pi_y} = -\frac{1 - \frac{\gamma^2 \theta^2}{4}}{1 + \frac{\beta \theta}{4}} \pi_y + \frac{\beta - \gamma^2 \theta}{2(1 + \frac{\beta \theta}{4})} x, \\
\dot{\pi}_x &= -\frac{\partial H_{NC}}{\partial x} = -\frac{\gamma^2 - \frac{\beta^2}{4}}{1 + \frac{\beta \theta}{4}} x - \frac{\beta - \gamma^2 \theta}{2(1 + \frac{\beta \theta}{4})} \pi_y, \\
\dot{\pi}_y &= -\frac{\partial H_{NC}}{\partial y} = -\frac{\gamma^2 - \frac{\beta^2}{4}}{1 + \frac{\beta \theta}{4}} y + \frac{\beta - \gamma^2 \theta}{2(1 + \frac{\beta \theta}{4})} \pi_x,
\end{aligned} \tag{3.21}$$

one can finally find the identical equations with (3.4),

$$\begin{aligned}
\dot{X} &= \frac{1}{\sqrt{1 + \frac{\beta \theta}{4}}} \left(\dot{x} - \frac{\theta}{2} \dot{\pi}_y \right) = -\Pi_X + \gamma^2 \Theta Y, \\
\dot{Y} &= \frac{1}{\sqrt{1 + \frac{\beta \theta}{4}}} \left(\dot{y} + \frac{\theta}{2} \dot{\pi}_x \right) = -\Pi_Y - \gamma^2 \Theta X, \\
\dot{\Pi}_X &= \frac{1}{\sqrt{1 + \frac{\beta \theta}{4}}} \left(\dot{\pi}_x + \frac{\beta}{2} \dot{y} \right) = -\gamma^2 X - B \Pi_Y, \\
\dot{\Pi}_Y &= \frac{1}{\sqrt{1 + \frac{\beta \theta}{4}}} \left(\dot{\pi}_y - \frac{\beta}{2} \dot{x} \right) = -\gamma^2 Y + B \Pi_X.
\end{aligned} \tag{3.22}$$

The check is completed in this way, as mentioned above.

IV. LOCALIZED NONCOMMUTATIVITY IN MOMENTA: CLASSICAL SYSTEM

If the parameter $\theta = 0$, (3.18) is simplified as

$$X = x, \quad Y = y, \quad \Pi_X = \pi_x + \frac{\beta}{2} y, \quad \Pi_Y = \pi_y - \frac{\beta}{2} x. \tag{4.1}$$

This is just the case that the authors of Refs. [1–4] considered. This model with noncommutativity in canonical momenta has several advantages over the simplicity. One of the advantages is the configuration variables X and Y are trivially expressed by x and y . Especially, in the study of general noncommutative quantum cosmology, we have to bring out the Wigner function [28–30] to analyze the solution of WDW equation if $\{X, Y\} \neq 0$ (see Ref. [25] for example). Even in the classical model of cosmology, the analysis becomes at

least tedious in general cases. The model including generic noncommutativity in the phase space of the minisuperspace is left for future work.

Another advantage of considering only nontrivial Poisson bracket $\{\Pi_X, \Pi_Y\} \neq 0$ is that there can be an analogy with a magnetic flux [1–4]. In Ref. [2], the authors explicitly pointed out the quantum Hall effect as the analogous system.²

Here, in the present paper, we propose localized noncommutativity in the minisuperspace. As mentioned already, it can be supposed that the noncommutativity is ineffective in the present era, but the great effect might exist in the early universe when the scale factors are very small.

Now, we assume that the parameter β in (4.1) is a function of $r = \sqrt{x^2 + y^2}$. Accordingly, the Poisson bracket of Π_X and Π_Y becomes

$$\{\Pi_X, \Pi_Y\} = \beta(r) + \frac{r\beta'(r)}{2}, \quad (4.2)$$

where the prime denotes the derivative with respect to r . Thus, the relation between β and B is slightly modified if there are position dependence.

The Hamiltonian (3.2) now takes the form

$$H_{NC} = -\frac{1}{2}(\pi_x^2 + \pi_y^2) + \frac{1}{2}\beta(r)(x\pi_y - y\pi_x) + \frac{1}{2}\left(\gamma^2 - \frac{\beta^2(r)}{4}\right)(x^2 + y^2). \quad (4.3)$$

The canonical equations

$$\begin{aligned} \dot{x} &= \frac{\partial H_{NC}}{\partial \pi_x} = -\pi_x - \frac{\beta(r)}{2}y, & \dot{y} &= \frac{\partial H_{NC}}{\partial \pi_y} = -\pi_y + \frac{\beta(r)}{2}x, \\ \dot{\pi}_x &= -\frac{\partial H_{NC}}{\partial x} = -\left(\gamma^2 - \frac{\beta(r)(\beta(r) + r\beta'(r))}{4}\right)x - \frac{\beta(r)}{2}\pi_y - \frac{\beta'(r)x}{2r}(x\pi_y - y\pi_x), \\ \dot{\pi}_y &= -\frac{\partial H_{NC}}{\partial y} = -\left(\gamma^2 - \frac{\beta(r)(\beta(r) + r\beta'(r))}{4}\right)y + \frac{\beta(r)}{2}\pi_x - \frac{\beta'(r)y}{2r}(x\pi_y - y\pi_x), \end{aligned} \quad (4.4)$$

leads to the following equations of motion:

$$\ddot{x} + \beta\dot{y} + \frac{\beta'}{2r}(x\dot{x} + y\dot{y})y - \left(\gamma^2 - \frac{r\beta\beta'}{4}\right)x + \frac{\beta'}{2r}\left[x\dot{y} - y\dot{x} - \frac{\beta}{2}(x^2 + y^2)\right]x = 0, \quad (4.5)$$

$$\ddot{y} - \beta\dot{x} - \frac{\beta'}{2r}(x\dot{x} + y\dot{y})x - \left(\gamma^2 - \frac{r\beta\beta'}{4}\right)y + \frac{\beta'}{2r}\left[x\dot{y} - y\dot{x} - \frac{\beta}{2}(x^2 + y^2)\right]y = 0. \quad (4.6)$$

By use of the polar coordinates

$$X = x = r \cos \phi, \quad Y = y = r \sin \phi, \quad (4.7)$$

² The analogy is pointed out also by the recent paper [31].

the equations of motion reduce to

$$\ddot{r} - r \left(\dot{\phi} - \frac{\beta}{2} \right)^2 - \left(\gamma^2 - \frac{\beta(\beta + r\beta')}{4} \right) r + \frac{\beta'}{2} r^2 \left(\dot{\phi} - \frac{\beta}{2} \right) = 0, \quad (4.8)$$

$$\frac{d}{dt} \left[r^2 \left(\dot{\phi} - \frac{\beta}{2} \right) \right] = 0. \quad (4.9)$$

On the other hand, the Hamiltonian constraint reads

$$\begin{aligned} H_{NC} &= -\frac{1}{2}\dot{r}^2 - \frac{1}{2}r^2\dot{\phi}^2 + \frac{1}{2}\gamma^2 r^2 \\ &= -\frac{1}{2}\dot{r}^2 - \frac{1}{2}r^2 \left(\dot{\phi} - \frac{\beta}{2} \right)^2 - \frac{\beta}{2}r^2 \left(\dot{\phi} - \frac{\beta}{2} \right) + \frac{1}{2} \left(\gamma^2 - \frac{\beta^2}{4} \right) r^2 = 0. \end{aligned} \quad (4.10)$$

The equation (4.9) can be integrated and if we set

$$L = r^2 \left(\dot{\phi} - \frac{\beta(r)}{2} \right) = \text{constant}, \quad (4.11)$$

the constraint equation becomes

$$\dot{r}^2 + \frac{L^2}{r^2} + L\beta(r) - \left(\gamma^2 - \frac{\beta^2(r)}{4} \right) r^2 = 0, \quad (4.12)$$

or, equivalently

$$\dot{r}^2 = \gamma^2 r^2 - \left(\frac{L}{r} + \frac{r\beta(r)}{2} \right)^2. \quad (4.13)$$

Now, we find that the equations of motion can be integrated and have very simple forms even if β is not constant as long as β is an arbitrary function of r .

We would like to show a concrete example of the solution. For this purpose, we adopt the following localized form of $\beta(r)$:

$$\beta(r) = \frac{\beta_0}{1 + e^{\mu(r-r_c)}}, \quad (4.14)$$

where β_0 , μ , and r_c are positive constants. This typical form shows that $\beta \sim \beta_0$ for $r < r_c$ and $\beta \sim 0$ for $r > r_c$, i.e., β is finite in the vicinity of the origin $X = Y = 0$. This form mimics a solenoid with radius r_c . In Fig. 1, we try to compare the noncommutative case with the commutative case. For the noncommutative case, we choose $\beta_0 = 5$, $\mu = 10$, and $r_c = 1$. The evolution of $X = x$ and $Y = y$ can be clearly seen if the solutions of (4.9) and (4.13) are plotted on an (X, Y) plane as shown in Fig. 1. The arrows in the figures indicate the normalized “velocity” vector

$$\frac{1}{\sqrt{\dot{X}^2 + \dot{Y}^2}} (\dot{X}, \dot{Y}), \quad (4.15)$$




FIG. 1. The normalized velocity vectors which indicate the flow of evolution of X and Y (see the text). (Top row) The commutative case ($\beta = 0$). The figures are plotted for $L = -2, -1, 0, 1, 2$, from left to right. (Bottom row) The case with the localized noncommutativity. For the choice of parameters, see the text. The figures are plotted for $L = -2, -1, 0, 1, 2$, from left to right.

at each point. In Fig. 1, the gray curves (quarter circles) indicate that the right-hand side of (4.13) becomes zero.

Roughly speaking, non zero β contributes to the “angular velocity” $\dot{\phi}$. Thus, the effective potential

$$V_{eff}(r) = -\frac{1}{2}\gamma^2 r^2 + \frac{1}{2} \left(\frac{L}{r} + \frac{r\beta(r)}{2} \right)^2. \quad (4.16)$$

has a positive barrier for a finite β even if $L = 0$. For a negative L and positive β , there appears a region where the solution is approximately described by (3.12) and (3.13) (with $\Theta = 0$ and $B = \beta_0$ in this time) because of the appearance of the small region of $V_{eff} < 0$ (in the region of $r < r_c$). In any case, a sufficiently large β_0 can forbid the small value of X and Y simultaneously; In short, $r = \sqrt{X^2 + Y^2} < \approx r_c$ is the forbidden region.

This means that the scale factors of bicosmology at the beginning of the classical universe is small but finite, in the sense of “average”, $r = \sqrt{X^2 + Y^2} \approx r_c$. The discussion on the averaging will be shown later.

V. LOCALIZED NONCOMMUTATIVITY IN MOMENTA: QUANTUM SYSTEM

The same system treated in the previous section is adopted also in this section, and we consider quantum cosmology [32–36] here. The WDW equation is obtained by the following replacement in the Hamiltonian constraint:

$$\pi_x \rightarrow -i \frac{\partial}{\partial x} \quad \pi_y \rightarrow -i \frac{\partial}{\partial y}. \quad (5.1)$$

Incidentally, one can confirm the commutators

$$[X, \Pi_X] = [Y, \Pi_Y] = i, \quad [\Pi_X, \Pi_Y] = i \left(\beta(r) + \frac{r}{2} \beta'(r) \right), \quad (5.2)$$

by using the set of operators

$$X = x, \quad Y = y, \quad \Pi_X = -i \frac{\partial}{\partial x} + \frac{\beta(r)}{2} y, \quad \Pi_Y = -i \frac{\partial}{\partial y} - \frac{\beta(r)}{2} x. \quad (5.3)$$

The WDW equation in our model can be written as, after the coordinate transformation (4.7),

$$\left[\frac{1}{r} \frac{\partial}{\partial r} r \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial \phi^2} - i \beta(r) \frac{\partial}{\partial \phi} + \left(\gamma^2 - \frac{\beta^2(r)}{4} \right) r^2 \right] \Psi(r, \theta) = 0, \quad (5.4)$$

where Ψ is the so-called wave function of the universe. Note that the metric of the minisuperspace is Euclidean (not Lorentzian).

The fundamental solution for this WDW equation takes the form

$$\psi_L(r) e^{-iL\phi}, \quad (5.5)$$

and then, the function ψ_L obeys the equation

$$\left[\frac{1}{r} \frac{\partial}{\partial r} r \frac{\partial}{\partial r} - \frac{L^2}{r^2} - L\beta(r) + \left(\gamma^2 - \frac{\beta^2(r)}{4} \right) r^2 \right] \psi_L(r) = 0. \quad (5.6)$$

The general solution then takes the form $\Psi = \sum_{L=-\infty}^{\infty} A_L \psi_L(r) e^{-iL\phi}$, where A_L denotes the amplitude.

For a constant β , the solution for $\psi_L(r)$ which is finite at $r = 0$ is expressed by using the confluent hypergeometric function as [37, 38]

$$\psi_L(r) = r^{|L|} e^{-i \frac{\sqrt{\gamma^2 - \frac{\beta^2}{4}}}{2} r^2} {}_1F_1 \left(\frac{|L| + 1}{2} + \frac{\beta L}{4i \sqrt{\gamma^2 - \frac{\beta^2}{4}}}; |L| + 1; i \sqrt{\gamma^2 - \frac{\beta^2}{4}} r^2 \right) \quad (\gamma^2 > \beta^2/4), \quad (5.7)$$

$$\psi_L(r) = r^{|L|} e^{-\sqrt{\frac{\beta^2}{4} - \gamma^2} r^2} {}_1F_1\left(\frac{|L|+1}{2} + \frac{\beta L}{4\sqrt{\frac{\beta^2}{4} - \gamma^2}}; |L|+1; \sqrt{\frac{\beta^2}{4} - \gamma^2} r^2\right) \quad (\gamma^2 < \beta^2/4). \quad (5.8)$$

Especially, for the commutative case $\beta \equiv 0$, $\psi_L(r) \propto J_{|L|/2}(\gamma r^2/2)$, where $J_\nu(z)$ is the Bessel function.

For a relatively small value of β and a finite L , the first peak of the wave function ψ_L is pushed out to larger r due to the ‘‘centrifugal force’’.




FIG. 2. The numerical solutions $\psi_0(r)$ ($L = 0$) are plotted for $\beta_0 = 0, 5, 10, 15$, from left to right. There is no scale on the vertical axis because an overall coefficient of the fundamental solution is arbitrary.

In the case with the localized noncommutativity, proposed in the previous section, the situation is the same. The most interesting case is when $L = 0$ with the localized noncommutativity. In Fig. 2, we plot $\psi_0(r)$ for various values of β_0 , where $\beta(r)$ is the same as (4.14), and $r_c = 1$, $\mu = 10$, and $\gamma = 1$. This behavior can be understood by considering the effective potential V_{eff} (4.16). As seen from Fig. 3, the localized noncommutativity creates the potential barrier and the first peak of the wave function is located outside $r = r_c$; It can be said that the wave function represents the tunneling phenomenon. The possibility that our universe start with a small but finite scale (in the present case, $r \approx r_c$) seems to be very interesting; The scenario is very akin to the model of early quantum cosmology, in which the potential barrier is supplied by the curvature of space [32].

Now, we confront the question: why should we focus on the $L = 0$ mode? One answer is the setting of $L = 0$ seems most simple and probable.³ Another answer is that we might consider averaging the (‘‘angular’’) variables, such that⁴

$$\frac{1}{2\pi} \int_0^{2\pi} \Psi(r, \theta) d\theta \propto \psi_0(r). \quad (5.9)$$

³ However, this is not so scientific but rather philosophical argument that exists universally in study of cosmology.

⁴ An additional advantage of the averaging is that we do not have to worry about the boundary conditions at $x = 0$ and at $y = 0$.




FIG. 3. A schematic view of the effective potential $V_{eff}(r)$ for $L = 0$. The gray curve indicates $\beta(r)$.

Looking back, even in the classical noncommutative solutions, averaging gave interesting results. Further discussions are given in the last section.

VI. CONCLUSION AND DISCUSSION

In this work, we investigated classical and quantum solutions for a toy model of bicosmology. We considered a symmetric model of two scale factors. Then, we derived the classical equations of the model when noncommutativities exist, both between the canonical momenta and between the dynamical variables of the minisuperspace. Further, we studied the system with the localized noncommutativity in momenta and classical and quantum cosmological solutions are investigated. In this system, the tunneling wave function can be found and it may provide a novel mechanism of the creation of the universe.

We must comment on the aforementioned averaging of variables. According to the original idea advocated in Refs. [1–4], two scale factors represent the causally disconnected regions. Naturally, it can be supposed that such two regions are extracted from a lot of typical regions in the universe. Thus, considering the averaging is not so novel procedure. As the average shown here of the scale factors, however, a strange fractional power appears (such as $x \propto a^{\frac{D-1}{2}}$), so further consideration is needed for expressing the proper averaging in terms of classical metrics. As for the wave function of the universe, the averaging is a speculative but interesting conjecture, since it explains the creation of a finite size universe.

An important problem is finding the mechanism to determine the critical value of r_c (as well as the finite β_0); This is unfortunately a difficult problem with few clues at the present

time.

On the other hand, an immediate issue is the introduction of the curvature of space and matter fields into the model. We point out that the introduction of phantom (scalar) fields [39–43] to the model allows the signature of the minisuperspace metric unchanged, so that one may select the well-defined boundary condition of the wave function at the origin of the minisuperspace, though there are various difficulties associated with the phantom field.

In addition, it is worth further studying the case with general noncommutativity configuration variables (such as Sec. III) and other possibility such as $[X, \Pi_X] = if(\Pi_X, \Pi_Y)$ (see Refs. [37, 38], for example), by classical and quantum approaches.

ACKNOWLEDGMENTS

We would like to thank Taiga Hasegawa for useful communication.

-
- [1] H. Falomir, J. Gamboa, F. Méndez and P. Gondolo, “Inflation without inflation: A model for dark energy”, *Phys. Rev.* **D96** (2017) 083534. DOI: 10.1103/PhysRevD.96.083534 arXiv:1707.04670 [gr-qc].
 - [2] H. Falomir, J. Gamboa, P. Gondolo and F. Méndez, “Magnetic seed and cosmology as quantum hall effect”, *Phys. Lett.* **B785** (2018) 399. DOI: 10.1016/j.physletb.2018.08.055 arXiv:1801.07575 [hep-th].
 - [3] H. Falomir, J. Gamboa and F. Méndez, “A noncommutative model of cosmology with two metrics”, *Symmetry* **2020** (2020) 435. DOI: 10.3390/sym12030435 arXiv:1807.08359 [hep-th].
 - [4] C. Maldonado and F. Méndez, “Bimetric universe with matter”, *Phys. Rev.* **D103** (2021) 123505. DOI: 10.1103/PhysRevD.103.123505 arXiv:2101.11235 [gr-qc].
 - [5] K. Aoki and K. Maeda, “dark matter in ghost-free bigravity theory”, *Phys. Rev.* **D90** (2014) 124089. DOI: 10.1103/PhysRevD.90.124089 arXiv:1409.0202 [gr-qc].
 - [6] A. Schmidt-May and M. von Strauss, “Recent developments in bimetric theory”, *J. Phys.* **A49** (2016) 183001. DOI: 10.1088/1751-8113/49/18/183001 arXiv:1512.00021 [hep-th].
 - [7] F. Darabi and M. Mousavi, “Classical and quantum cosmology of minimal massive bigravity”, *Phys. Lett.* **B761** (2016) 269. DOI: 10.1016/j.physletb.2016.08.031 arXiv:1512.03333 [gr-qc].

- [8] H. García-Compeán, O. Obregón and C. Ramírez, “Noncommutative quantum cosmology”, *Phys. Rev. Lett.* **88** (2002) 161301. DOI: 10.1103/PhysRevLett.88.161301 hep-th/0107250.
- [9] G. D. Barbosa and N. Pinto-Neto, “Noncommutative geometry and cosmology”, *Phys. Rev.* **D70** (2004) 103512. DOI: 10.1103/PhysRevD.70.103512 hep-th/0407111.
- [10] L. O. Pimentel and C. Mora, “Noncommutative quantum cosmology”, *Gen. Rel. Grav.* **37** (2005) 817. DOI: 10.1007/s10714-005-0066-3 gr-qc/0408100.
- [11] L. O. Pimentel and O. Obregón, “Non commuting quantum cosmology with scalar matter”, *Gen. Rel. Grav.* **38** (2006) 553. DOI: 10.1007/s10714-006-0246-9
- [12] M. Aguero, J. A. S. Aguilar, C. Ortiz, M. Sabido and J. Socorro, “Noncommutative Bianchi type II quantum cosmology”, *Int. J. Theor. Phys.* **46** (2007) 2928. DOI: 10.1007/s10773-007-9405-3
- [13] W. Guzmán, M. Sabido and J. Socorro, “Noncommutativity and scalar field cosmology”, *Phys. Rev.* **D76** (2007) 087302. DOI: 10.1103/PhysRevD.76.087302 arXiv:0712.1520 [gr-qc].
- [14] W. Guzmán, M. Sabido and J. Socorro, “Towards an inflationary scenario in noncommutative quantum cosmology”, *Rev. Mex. Fís.* **53** suppl. 4 (2007) 94.
- [15] W. Guzmán, M. Sabido and J. Socorro, “Towards noncommutative supersymmetric quantum cosmology”, *AIP Conf. Proc.* **1318** (2010) 209. DOI: 10.1063/1.3531633 arXiv:0812.4999 [hep-th].
- [16] E. Mena, O. Obregón and M. Sabido, “On the WKB approximation of noncommutative quantum cosmology”, *Rev. Mex. Fís.* **53** suppl. 4 (2007) 118.
- [17] E. Mena, O. Obregón and M. Sabido, “WKB-type approximation to noncommutative quantum cosmology”, *Int. J. Mod. Phys.* **D18** (2009) 95. DOI: 10.1142/S0218271809014376 gr-qc/0701097.
- [18] C. Ortiz, E. Mena, M. Sabido and J. Socorro, “(Non)commutative isotropization in Bianchi I with barotropic perfect fluid and Λ cosmological”, *Int. J. Theor. Phys.* **47** (2008) 1240. DOI: 10.1007/s10773-007-9557-1 gr-qc/0703101.
- [19] J. Socorro, L. O. Pimentel, C. Ortiz and M. Aguero, “Scalar field in the Bianchi I: Noncommutative classical and quantum cosmology”, *Int. J. Theor. Phys.* **48** (2009) 3567. DOI: 10.1007/s10773-009-0164-1 arXiv:0910.2449 [gr-qc].
- [20] C. Bastos, O. Bertolami, N. C. Dias and J. N. Prata, “Phase-space noncommutative quantum cosmology”, *Phys. Rev.* **D78** (2008) 023516. DOI: 10.1103/PhysRevD.78.023516

- arXiv:0712.4122 [gr-qc].
- [21] C. Bastos, O. Bertolami, N. C. Dias and J. N. Prata, “Noncommutative quantum cosmology”, *J. Phys. Conf. Ser.* **174** (2009) 012053. DOI: 10.1088/1742-6596/174/1/012053 arXiv:0812.3488 [gr-qc].
- [22] C. Bastos, O. Bertolami, N. Dias and J. N. Prata, “Noncommutative quantum mechanics and quantum cosmology”, *Int. J. Mod. Phys. A* **24** (2009) 2741. DOI: 10.1142/S0217751X09046138 arXiv:0904.0400 [hep-th].
- [23] M. Maceda, A. Macías and L. O. Pimentel, “Homogeneous noncommutative quantum cosmology”, *Phys. Rev. D* **78** (2008) 064041 DOI: 10.1103/PhysRevD.78.064041.
- [24] O. Obregón and I. Quiros, “Can noncommutative effects account for the present speed up of the cosmic expansion?”, *Phys. Rev. D* **84** (2011) 044005. DOI: 10.1103/PhysRevD.84.044005 arXiv:1011.3896 [gr-qc].
- [25] N. Kan, M. Kuniyasu, K. Shiraishi and K. Takimoto, “Accelerating cosmologies in an integrable model with noncommutative minisuperspace variables”, *J. Phys. Comm.* **4** (2020) 075010. DOI: 10.1088/2399-6528/aba1d3 arXiv:1903.07895 [gr-qc].
- [26] S. Jalalzadeh, M. Rashki and S. abarghouei Nejad, “Classical universe arising from quantum cosmology”, *Phys. Dark Universe* **30** (2020) 100741. DOI: 10.1016/j.dark.2020.100741 arXiv:2001.00556 [gr-qc].
- [27] H. García-Compeán, O. Obregón and C. Ramírez, “Topics in supersymmetric and noncommutative quantum cosmology”, *Universe* **7** (2021) 434. DOI: 10.3390/universe7110434
- [28] E. Wigner, “On the quantum correction for thermodynamic equilibrium”, *Phys. Rev.* **40** (1932) 749. DOI: 10.1103/PhysRev.40.749.
- [29] W. B. Case, “Wigner functions and Weyl transforms for pedestrians”, *Am. J. Phys.* **76** (2008) 937. DOI: 10.1119/1.2957889.
- [30] J. Weinbub and D. K. Ferry, “Recent advances in Wigner function approaches”, *App. Phys. Rev.* **5** (2018) 041104. DOI: 10.1063/1.5046663.
- [31] E. Guendelman and D. Singleton, “Momentum gauge fields and non-commutative space-time”, arXiv:2206.02638 [quant-ph].
- [32] J. B. Hartle and S. W. Hawking, “Wave function of the universe”, *Phys. Rev. D* **28** (1983) 2960. DOI: 10.1103/PhysRevD.28.2960

- [33] S. W. Hawking, “The quantum state of the universe”, Nucl. Phys. **B239** (1984) 257. DOI: 10.1016/0550-3213(84)90093-2.
- [34] J. J. Halliwell, “Introductory lectures on quantum cosmology”, in Proceedings of the Jerusalem Winter School on Quantum Cosmology and Baby Universe (edited by T. Piran, World Scientific, Singapore, 1991), DOI: 10.1142/9789814503501_0003. arXiv:0909.2566 [gr-qc].
- [35] C. Kiefer, *Quantum Gravity*, Int. Ser. Monogr. Phys. Vol. **155** (Clarendon Press, Oxford, 2012). DOI: 10.1093/acprof:oso/9780199585205.001.0001
- [36] C. Kiefer, “Conceptual problems on quantum gravity and quantum cosmology”, ISRN Mathematical Physics **2013** (2013) 509316. DOI: 10.1155/2013/509316. arXiv:1401.3578 [gr-qc].
- [37] A. Boumali and Z. Selama, “Two-dimensional Klein–Gordon oscillator in the presence of a minimal length”, Phys. Part. Nuclei Lett. **15** (2018) 473. DOI: 10.1134/S1547477118050047 arXiv:1706.08593 [quant-ph].
- [38] Z. Selama and A. Boumali, “Two-dimensional boson oscillator under a magnetic field in the presence of a minimal length in the non-commutative space”, Rev. Mex. Fis. **67** (2021) 226. DOI: 10.31349/RevMexFis.67.226
- [39] R. R. Caldwell, “A phantom menace? cosmological consequences of a dark energy component with super-negative equation of state”, Phys. Lett. **B545** (2002) 23. DOI: 10.1016/S0370-2693(02)02589-3 astro-ph/9908168.
- [40] R. R. Caldwell, M. Kamionkowski and N. N. Weinberg, “Phantom energy: Dark energy with $w < -1$ causes a cosmic doomsday”, Phys. Rev. Lett. **91** (2003) 071301. DOI: 10.1103/PhysRevLett.91.071301 astro-ph/0302506.
- [41] M. P. Dabrowski, T. Stachowiak and M. Szydlowski, “Phantom cosmologies”, Phys. Rev. **D68** (2003) 103519. DOI: 10.1103/PhysRevD.68.103519 hep-th/0307128.
- [42] M. P. Dabrowski, C. Kiefer and B. Sandhöfer, “Quantum phantom cosmology”, Phys. Rev. **D74** (2006) 044022. DOI: 10.1103/PhysRevD.74.044022 hep-th/0605229.
- [43] D. E. Kaplan and R. Sundrum, “A symmetry for the cosmological constant”, JHEP **0607** (2006) 042. DOI: 10.1088/1126-6708/2006/07/042 hep-th/0505265.