

# Some useful operators on differential forms in Galilean and Carrollian spacetimes

Marián Fecko\*

Department of Theoretical Physics

Faculty of Mathematics, Physics and Informatics

Comenius University in Bratislava

*Differential forms in Minkowski spacetime is a folklore subject. In Galilean and Carrollian spacetimes it does not seem to be quite so. This might be caused by absence of Hodge star operator. There are, however, useful analogs of Hodge star operator also in the two spacetimes, namely intertwining operators between corresponding representations on forms. Their use could make differential forms as interesting tool for writing down Galilean and Carrollian invariant equations as forms in Minkowski spacetime are known to be for writing down (as well as manipulating) Poincaré invariant equations.*

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Forms in Galilean, Carrollian and Minkowski spacetime</b>	<b>5</b>
2.1	Parametrization . . . . .	5
2.2	Generators of rotations . . . . .	6
2.3	Generators of boosts . . . . .	8
2.4	List of all generators $\rho_p$ . . . . .	9
<b>3</b>	<b>Intertwining operators <math>a_{qp}</math></b>	<b>10</b>
3.1	What intertwining operators we are interested in . . . . .	10
3.2	General structure of all $a_{qp}$ (no invariance, yet) . . . . .	12
3.3	How $a_{qp}$ are restricted by rotation invariance alone . . . . .	12
3.4	Restrictions on $a_{qp}$ added by boosts . . . . .	13

---

\*e-mail: [Marian.Fecko@fmph.uniba.sk](mailto:Marian.Fecko@fmph.uniba.sk)

<b>4</b>	<b>Results - non-trivial operators <math>a_{qp}</math></b>	<b>13</b>
4.1	Which operators $a_{qp}$ are trivial . . . . .	13
4.2	Non-trivial operators $a_{qp}$ in Lorentzian spacetime . . . . .	14
4.3	Non-trivial operators $a_{qp}$ in Galilean spacetime . . . . .	15
4.4	Non-trivial operators $a_{qp}$ in Carrollian spacetime . . . . .	17
4.5	A note on the degree $\pm 1$ operators . . . . .	18
4.6	Invariant operators and invariant subspaces . . . . .	19
<b>5</b>	<b>Summary and conclusions</b>	<b>21</b>
<b>A</b>	<b>Appendices</b>	<b>22</b>
A.1	Details of solving equations for $a_{qp}$ . . . . .	22
A.1.1	Operators $a_{00}, a_{40}, a_{04}$ and $a_{44}$ . . . . .	22
A.1.2	Operators $a_{02}, a_{20}, a_{42}$ and $a_{24}$ . . . . .	23
A.1.3	Operators $a_{01}, a_{03}, a_{41}$ and $a_{43}$ . . . . .	23
A.1.4	Operators $a_{10}, a_{30}, a_{14}$ and $a_{34}$ . . . . .	24
A.1.5	Operators $a_{11}, a_{33}, a_{13}$ and $a_{31}$ . . . . .	24
A.1.6	Operators $a_{12}$ and $a_{32}$ . . . . .	26
A.1.7	Operators $a_{21}$ and $a_{23}$ . . . . .	27
A.1.8	Operator $a_{22}$ . . . . .	28
A.2	Galilean and Carrollian Hodge star operators . . . . .	28

# 1 Introduction

Differential forms in spacetime provide a well-established tool used in theoretical physics. For example, the fact that

$$d * F = -J \quad dF = 0 \tag{1}$$

represent Maxwell equations and its consistency condition

$$dJ = 0 \tag{2}$$

expresses local conservation of electric charge, is a folklor knowledge.

There are many reasons why forms are so effective.

One of their attractive features is that expressing laws of physics in terms of forms is often the most succinct way how to achieve *manifest* invariance of the resulting equations. And this is because forms themselves, as well as important *operators* on forms, behave nicely w.r.t. relevant diffeomorphisms.

For equations (1) and (2), say, let

$$f : x \mapsto x' = \Lambda x + a \quad \Lambda^T \eta \Lambda = \eta \tag{3}$$

be a Poincaré transformation, regarded as a transformation between two *inertial* observers in *Minkowski* spacetime. Then well-known properties of the two operators present in the equations,  $d$  and  $*$ ,

$$f^*d = df^* \quad (4)$$

$$f^* * = * f^* \quad (5)$$

result in statements that

- if  $F$  is a solution of (1) generated by  $J$ ,
- then  $f^*F$  is a solution of (1) generated by  $f^*J$

and, in addition,

- if  $dJ = 0$  (i.e. the first observer sees local conservation of electric charge),
- then  $d(f^*J) = 0$  (i.e. the second observer sees the same phenomenon).

So, both observers use the same equations (1) and (2), and the vocabulary between their descriptions of the same phenomenon is just making of *pull-back* of corresponding forms (solutions of (1) and (2)).

Notice that crucial element of this striking simplicity is the (simplest possible) behavior

- (4) of the *exterior derivative*  $d$  and
- (5) of the *Hodge star* operator  $*$ .

Now, there is an important difference, in this respect, between  $d$  and  $*$ .

While  $d$  needs *no* additional structure on a smooth manifold  $M$  and (4) is true for *any* diffeomorphism  $f : M \rightarrow M$ , the Hodge star  $* \equiv *_{g,o}$  (see e.g. [1] - [7])

- is only defined on a (pseudo)Riemannian oriented manifold  $(M, g, o)$

$$* : \Omega^p \rightarrow \Omega^{n-p} \quad \Omega^p \equiv \Omega^p(M, g, o) \quad (6)$$

- behaves in a more complicated way w.r.t. a *general* diffeomorphism, namely

$$f^*(*_{g,o}\alpha) = *_{f^*g, f(o)}(f^*\alpha) \quad (7)$$

or, in short,

$$f^* *_{g,o} = *_{f^*g, f(o)} f^* \quad (8)$$

(it is said to be *natural* w.r.t. diffeomorphisms of  $M$ ).

Then, clearly, for (orientation preserving) *isometries* of  $(M, g)$ , the Hodge star operator becomes *invariant*

$$f^*g = g \quad \Rightarrow \quad f^* *_{g,o} = *_{g,o} f^* \quad (9)$$

In particular, if  $g \equiv \eta$  corresponds to *Minkowski spacetime*  $(M, g) = (\mathbb{R}^4, \eta)$ , the corresponding Hodge star  $* = *_{\eta}$  is invariant w.r.t. *Poincaré* transformations.

So, the important property (5) is based, in our case, on

- the existence of *metric tensor* on the (Minkowski) spacetime,

- such that transformations between inertial observers are its *isometries*.

A lot of research is recently done on *Galilean* and *Carrollian* spacetimes, see e.g. [8] - [14]; also long lists of references in recent PhD theses [15] and [16].) There, by definition, Galilean or Carrollian transformations, respectively, play formally role of transformations connecting various inertial observers. What about repeating the same story there?

According to what was written above, we have no problem with  $d$  (it works perfectly anywhere, including the two “new” spacetimes). Concerning the Hodge star operator, one should

- take a metric tensor on Galilean (or Carrollian) spacetime
- such that Galilean (or Carrollian) transformations become its isometries

and then invariance of the corresponding Hodge star is guaranteed by (9).

There is, however, a serious problem with this natural idea: It is well known that

- *neither* Galilean invariant metric tensor on Galilean spacetime
- *nor* Carrollian invariant metric tensor on Carrollian spacetime

do exist. So, the routine procedure of construction of  $*$  *fails*, there.

However, it is not excluded, apriori, that still some useful *analogs* of the star operator exist in the two spacetimes. In order to contemplate on them, we can

- either try to suitably modify the standard *formula* for the Hodge operator (this is done in [17])
- or try to identify which particular *property* of the Hodge star is *the* most important for us and then try to find, if possible, *all* operators with the property (this is the idea to be examined in the present paper).

From the point of view of striving for writing down (and manipulating with) *invariant* physics equations, *invariance* property of the operator is to be on the top of our interest. So operators with property (5) w.r.t.

- Galilean transformations  $f$  in Galilean spacetime and
- Carrollian transformations  $f$  in Carrollian spacetime

are to be of interest for us.

Now, we concentrate on *differential forms* (on particular spacetime  $M$ ). Then, in any particular space  $\Omega^p(M)$  a *representation*  $\rho_p$  of the symmetry group (the group of Poincaré, Galilean or Carrollian transformations) is induced (via pull-back) by  $f$ . In this language the invariance property (5) of  $*$  w.r.t. Poincaré transformations, as an example, may be expressed in terms of the commutative diagram

$$\begin{array}{ccc}
 \Omega^p & \xrightarrow{*} & \Omega^{n-p} \\
 \rho_p \downarrow & & \downarrow \rho_{n-p} \\
 \Omega^p & \xrightarrow[*]{} & \Omega^{n-p}
 \end{array}
 \quad \text{i.e.} \quad
 \rho_{n-p} \circ * = * \circ \rho_p
 \quad (10)$$

This says, in representation theory parlance, that  $*$  may be regarded as *intertwining operator* between representations  $\rho_p$  and  $\rho_{n-p}$ .

Then it looks as a promising idea, on the route to find *useful analogs* of the Hodge star for Galilean and Carrollian spacetimes, to search for all <sup>1</sup> *intertwining operators* on forms, acting between *general* representations  $\rho_p$  and  $\rho_q$ :

$$\begin{array}{ccc}
 \Omega^p & \xrightarrow{a_{qp}} & \Omega^q \\
 \rho_p \downarrow & & \downarrow \rho_q \\
 \Omega^p & \xrightarrow{a_{qp}} & \Omega^q
 \end{array}
 \quad \text{i.e.} \quad
 \rho_q \circ a_{qp} = a_{qp} \circ \rho_p
 \quad (11)$$

Any such operator may be then combined with the exterior derivative  $d$  in order to write down Galilean (or Carrollian) *explicitly invariant* differential equation.

In what follows we will see that such operators do exist and that their list depends heavily on the choice of particular spacetime (Galilean, Carrollian, Minkowski).

In particular, in Minkowski spacetime, the *only* (non-trivial) intertwining operator turns out to be the Hodge star.

- In the Galilean and Carrollian spacetimes, on the contrary, there are operators
- acting similarly to (Minkowski) Hodge star (already known from [17])
- with no similarity to (Minkowski) Hodge star.

So, in this sense, the structure of intertwining operators on forms is *richer* in the Galilean and Carrollian spacetimes than it is in the Minkowski one. (See also discussion in Section 4.6.)

## 2 Forms in Galilean, Carrollian and Minkowski spacetime

### 2.1 Parametrization

In order to write down explicitly equations (11), we need to determine explicit form of the matrices  $\rho_p$  for

- all three spacetimes (Galilean, Carrollian, Minkowski)
- all degrees  $p = 0, 1, 2, 3, 4$  of differential forms.

Let us start with a suitable coordinates and parametrization of forms.

In all three cases we have (global) coordinates  $(t, x^i) = (t, x, y, z)$  adapted to particular spacetime structures. In these coordinates, we can use, in *all three* cases, parametrization of differential forms well-known from the Minkowski spacetime (see e.g. Section 16.1 in [7]). That is, any  $p$ -form  $\alpha$ ,  $p = 0, 1, 2, 3, 4$ , may be first

---

<sup>1</sup>We only mean all “non-differential”, purely linear algebraic, as it is the case for the Hodge star.

uniquely decomposed as

$$\alpha = dt \wedge \hat{s} + \hat{r} \quad (12)$$

where the two hatted forms,  $(p-1)$ -form  $\hat{s}$  and  $p$ -form  $\hat{r}$ , respectively, are *spatial*, meaning that they do not contain (“temporal”) differential  $dt$  in its coordinate expression. Being spatial,  $\hat{s}$  and  $\hat{r}$  may be, depending on degree, expressed as

$$f \quad \mathbf{a} \cdot d\mathbf{r} \quad \mathbf{a} \cdot d\mathbf{S} \quad f dV \quad (13)$$

where

$$\mathbf{a} \cdot d\mathbf{r} = a_x dx + a_y dy + a_z dz \quad (14)$$

$$\mathbf{a} \cdot d\mathbf{S} = a_x dS_x + a_y dS_y + a_z dS_z \quad (15)$$

$$\equiv a_x dy \wedge dz + a_y dz \wedge dx + a_z dx \wedge dy \quad (16)$$

$$dV = dx \wedge dy \wedge dz \quad (17)$$

So, the general decomposition formula (12) may be further specified, for the five relevant degrees of differential forms, as follows:

$$\Omega^0 : \alpha = f \quad (18)$$

$$\Omega^1 : \alpha = f dt + \mathbf{a} \cdot d\mathbf{r} \quad (19)$$

$$\Omega^2 : \alpha = dt \wedge \mathbf{a} \cdot d\mathbf{r} + \mathbf{b} \cdot d\mathbf{S} \quad (20)$$

$$\Omega^3 : \alpha = dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV \quad (21)$$

$$\Omega^4 : \alpha = f dt \wedge dV \quad (22)$$

(Notice that the time coordinate  $t$  itself may be present in *components* of the spatial parts, for example  $a_x = a_x(t, x, y, z)$  in general.)

Alternatively, we can uniquely identify forms from (18) - (22) just with columns

$$f \quad \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} \quad \begin{pmatrix} \mathbf{a} \\ \mathbf{b} \end{pmatrix} \quad \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} \quad f \quad (23)$$

Now we go on to study action of corresponding spacetime symmetry groups on forms within this parametrization.

## 2.2 Generators of rotations

In above mentioned coordinates, what is common for all three spacetimes is action of

- time and space translation
- spatial (3D) rotations

$$t' = t + t_0 \quad (24)$$

$$\mathbf{r}' = R\mathbf{r} + \mathbf{r}_0 \quad (25)$$

where  $R$  denotes  $3 \times 3$  rotation matrix.

What is different, is action of boosts (see below, Section 2.3).

Concerning scrambling of components of forms, time and space *translations* alone do *nothing* (since  $dt_0$  and  $d\mathbf{r}_0$  vanish), so we can forget about them and restrict to rotations and boosts.

Moreover, we can restrict to *infinitesimal* rotations and boosts since even if  $\rho_p$  in the commutative diagram (11) were meant as operators representing corresponding symmetry *Lie group*, the same intertwining operators enter the “infinitesimal version” of the commutative diagram, where  $\rho_p$  already denote operators of the *derived* representation of the corresponding symmetry *Lie algebra*.

For infinitesimal rotations about  $j$ -th axis, formulas (24) - (25) reduce to

$$\begin{pmatrix} t' \\ \mathbf{r}' \end{pmatrix} = \begin{pmatrix} t \\ \mathbf{r} \end{pmatrix} + \epsilon \begin{pmatrix} 0 & 0 \\ 0 & l_j \end{pmatrix} \begin{pmatrix} t \\ \mathbf{r} \end{pmatrix} \quad (26)$$

where

$$(l_j)_{ik} = \epsilon_{ijk} \quad (27)$$

Formula (26) induces action on coordinate basis of forms

$$dt' = dt \quad (28)$$

$$d\mathbf{r}' = (\mathbb{1} + \epsilon l_j) d\mathbf{r} \quad (29)$$

$$d\mathbf{S}' = (\mathbb{1} + \epsilon l_j) d\mathbf{S} \quad (30)$$

$$dV' = dV \quad (31)$$

From requirement

$$f' = f \quad (32)$$

$$f' dt' + \mathbf{a}' \cdot d\mathbf{r}' = f dt + \mathbf{a} \cdot d\mathbf{r} \quad (33)$$

$$dt' \wedge \mathbf{a}' \cdot d\mathbf{r}' + \mathbf{b}' \cdot d\mathbf{S}' = dt \wedge \mathbf{a} \cdot d\mathbf{r} + \mathbf{b} \cdot d\mathbf{S} \quad (34)$$

$$dt' \wedge \mathbf{a}' \cdot d\mathbf{S}' + f' dV' = dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV \quad (35)$$

$$f' dt' \wedge dV' = f dt \wedge dV \quad (36)$$

we can compute, for each degree, primed components in terms of unprimed and get

$$\Omega^0 \quad f' = f \quad (37)$$

$$\Omega^1 \quad \begin{pmatrix} f' \\ \mathbf{a}' \end{pmatrix} = \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} + \epsilon \begin{pmatrix} 0 & 0 \\ 0 & l_j \end{pmatrix} \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} \quad (38)$$

$$\Omega^2 \quad \begin{pmatrix} \mathbf{a}' \\ \mathbf{b}' \end{pmatrix} = \begin{pmatrix} \mathbf{a} \\ \mathbf{b} \end{pmatrix} + \epsilon \begin{pmatrix} l_j & 0 \\ 0 & l_j \end{pmatrix} \begin{pmatrix} \mathbf{a} \\ \mathbf{b} \end{pmatrix} \quad (39)$$

$$\Omega^3 \quad \begin{pmatrix} f' \\ \mathbf{a}' \end{pmatrix} = \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} + \epsilon \begin{pmatrix} 0 & 0 \\ 0 & l_j \end{pmatrix} \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} \quad (40)$$

$$\Omega^4 \quad f' = f \quad (41)$$

All the formulas may be written (for rotation about  $j$ -th axis) as multiplication of the correspondent component column by matrix

$$\mathbb{1} + \epsilon S_j \quad (42)$$

where

$$\Omega^0 \quad S_j = 0 \quad (43)$$

$$\Omega^1 \quad S_j = \begin{pmatrix} 0 & 0 \\ 0 & l_j \end{pmatrix} \quad (44)$$

$$\Omega^2 \quad S_j = \begin{pmatrix} l_j & 0 \\ 0 & l_j \end{pmatrix} \quad (45)$$

$$\Omega^3 \quad S_j = \begin{pmatrix} 0 & 0 \\ 0 & l_j \end{pmatrix} \quad (46)$$

$$\Omega^4 \quad S_j = 0 \quad (47)$$

## 2.3 Generators of boosts

Let us display all three kinds of infinitesimal boosts along  $\mathbf{n}$ :

$$\text{Lorentzian} \quad t' = t + \epsilon \mathbf{n} \cdot \mathbf{r} \quad (48)$$

$$\mathbf{r}' = \mathbf{r} + \epsilon \mathbf{n} t \quad (49)$$

$$\text{Galilean} \quad t' = t \quad (50)$$

$$\mathbf{r}' = \mathbf{r} + \epsilon \mathbf{n} t \quad (51)$$

$$\text{Carrollian} \quad t' = t + \epsilon \mathbf{n} \cdot \mathbf{r} \quad (52)$$

$$\mathbf{r}' = \mathbf{r} \quad (53)$$

(As for the Carrollian boosts, see (9) in [22] or II.10 in [8].) The following formulas conveniently cover all three cases at once:

$$t' = t + \epsilon \lambda_1 \mathbf{n} \cdot \mathbf{r} \quad (54)$$

$$\mathbf{r}' = \mathbf{r} + \epsilon \lambda_2 \mathbf{n} t \quad (55)$$

The three special cases clearly correspond to values

$$\text{Lorentzian} \quad (\lambda_1, \lambda_2) = (1, 1) \quad (56)$$

$$\text{Galilean} \quad (\lambda_1, \lambda_2) = (0, 1) \quad (57)$$

$$\text{Carrollian} \quad (\lambda_1, \lambda_2) = (1, 0) \quad (58)$$

Induced action on coordinate basis of forms comes out to be

$$\Omega^1 \quad dt' = dt + \epsilon \lambda_1 \mathbf{n} \cdot d\mathbf{r} \quad (59)$$

$$d\mathbf{r}' = d\mathbf{r} + \epsilon \lambda_2 \mathbf{n} dt \quad (60)$$

$$\Omega^2 \quad dt' \wedge d\mathbf{r}' = dt \wedge d\mathbf{r} - \epsilon \lambda_1 (\mathbf{n} \cdot \mathbf{l}) d\mathbf{S} \quad (61)$$

$$d\mathbf{S}' = d\mathbf{S} + \epsilon \lambda_2 dt \wedge (\mathbf{n} \cdot \mathbf{l}) d\mathbf{r} \quad (62)$$

$$\Omega^3 \quad dt' \wedge d\mathbf{S}' = dt \wedge d\mathbf{S} + \epsilon \lambda_1 \mathbf{n} dV \quad (63)$$

$$dV' = dV + \epsilon \lambda_2 dt \wedge (\mathbf{n} \cdot d\mathbf{S}) \quad (64)$$

$$\Omega^4 \quad dt' \wedge dV' = dt \wedge dV \quad (65)$$

Writing down again (32) - (36) we get boost (along  $x_j$ ) analogs of (37) - (41)

$$\Omega^0 \quad f' = f \quad (66)$$

$$\Omega^1 \quad \begin{pmatrix} f' \\ \mathbf{a}' \end{pmatrix} = \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} + \epsilon \begin{pmatrix} 0 & \lambda_2 e_j^T \\ \lambda_1 e_j & 0 \end{pmatrix} \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} \quad (67)$$

$$\Omega^2 \quad \begin{pmatrix} \mathbf{a}' \\ \mathbf{b}' \end{pmatrix} = \begin{pmatrix} \mathbf{a} \\ \mathbf{b} \end{pmatrix} + \epsilon \begin{pmatrix} 0 & -\lambda_2 l_j \\ \lambda_1 l_j & 0 \end{pmatrix} \begin{pmatrix} \mathbf{a} \\ \mathbf{b} \end{pmatrix} \quad (68)$$

$$\Omega^3 \quad \begin{pmatrix} f' \\ \mathbf{a}' \end{pmatrix} = \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} + \epsilon \begin{pmatrix} 0 & \lambda_1 e_j^T \\ \lambda_2 e_j & 0 \end{pmatrix} \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} \quad (69)$$

$$\Omega^4 \quad f' = f \quad (70)$$

(where  $e_j$  denotes unit vector along  $x_j$ ). They can be written via multiplication by

$$\mathbb{1} + \epsilon N_j \quad (71)$$

where

$$\Omega^0 \quad N_j = 0 \quad (72)$$

$$\Omega^1 \quad N_j = \begin{pmatrix} 0 & \lambda_2 e_j^T \\ \lambda_1 e_j & 0 \end{pmatrix} \quad (73)$$

$$\Omega^2 \quad N_j = \begin{pmatrix} 0 & -\lambda_2 l_j \\ \lambda_1 l_j & 0 \end{pmatrix} \quad (74)$$

$$\Omega^3 \quad N_j = \begin{pmatrix} 0 & \lambda_1 e_j^T \\ \lambda_2 e_j & 0 \end{pmatrix} \quad (75)$$

$$\Omega^4 \quad N_j = 0 \quad (76)$$

## 2.4 List of all generators $\rho_p$

If we denote, in general,

$$\rho_p = S_j \text{ and } N_j \text{ on components of } p\text{-forms} \quad (77)$$

then the full list of generators on (suitably parametrized components of)  $p$ -forms explicitly reads

$$\rho_0 \quad S_j = 0 \quad N_j = 0 \quad (78)$$

$$\rho_1 \quad S_j = \begin{pmatrix} 0 & 0 \\ 0 & l_j \end{pmatrix} \quad N_j = \begin{pmatrix} 0 & \lambda_2 e_j^T \\ \lambda_1 e_j & 0 \end{pmatrix} \quad (79)$$

$$\rho_2 \quad S_j = \begin{pmatrix} l_j & 0 \\ 0 & l_j \end{pmatrix} \quad N_j = \begin{pmatrix} 0 & -\lambda_2 l_j \\ \lambda_1 l_j & 0 \end{pmatrix} \quad (80)$$

$$\rho_3 \quad S_j = \begin{pmatrix} 0 & 0 \\ 0 & l_j \end{pmatrix} \quad N_j = \begin{pmatrix} 0 & \lambda_1 e_j^T \\ \lambda_2 e_j & 0 \end{pmatrix} \quad (81)$$

$$\rho_4 \quad S_j = 0 \quad N_j = 0 \quad (82)$$

Observe a simple rule, which can save some computations in the future:

- Interchanging  $(\lambda_1, \lambda_2) \leftrightarrow (\lambda_2, \lambda_1)$  leads to interchanging  $\rho_1 \leftrightarrow \rho_3$

or, as a formula,

$$\rho_3(\lambda_1, \lambda_2) = \rho_1(\lambda_2, \lambda_1) \quad (83)$$

### 3 Intertwining operators $a_{qp}$

#### 3.1 What intertwining operators we are interested in

On Minkowski, Galilean or Carrollian spacetime, respectively, let

$$a_{qp} : \Omega^p \rightarrow \Omega^q \quad (84)$$

be a linear operator realized as a *matrix* on *components* of the forms. (Notice the order of indices convention - mapping from  $p$  to  $q$  is denoted as  $a_{qp}$  rather than vice versa.)

Recall (see (23)) that the components of  $p$ -forms are parametrized, for  $p = 0, \dots, 4$ , either by single objects or by pairs

$$f \quad \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} \quad \begin{pmatrix} \mathbf{a} \\ \mathbf{b} \end{pmatrix} \quad \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} \quad f \quad (85)$$

Therefore, the matrices  $a_{qp}$  are necessarily *block* matrices of (block) dimensions  $1 \times 1$ ,  $1 \times 2$ ,  $2 \times 1$  or  $2 \times 2$ . As an example, matrices  $a_{12}$  and  $a_{02}$  are of the form

$$a_{12} = \begin{pmatrix} \mathbf{c}^T & \mathbf{d}^T \\ C & D \end{pmatrix} \quad a_{02} = (\mathbf{c}^T \quad \mathbf{d}^T) \quad (86)$$

for  $\mathbf{c}, \mathbf{d}$  (column) vectors (i.e.  $\mathbf{c}^T, \mathbf{d}^T$  row vectors) and  $C, D$   $3 \times 3$  matrices.

Within the spaces  $\Omega^p$  and  $\Omega^q$ , the corresponding (Lorentzian, Galilean or Carrollian) transformations induce (as generators) matrices  $\rho_p$  and  $\rho_q$ , respectively,

acting on components of forms and explicitly given, for both rotations and boosts, by (78) - (82).

Now we want  $a_{qp}$  to be *intertwining operator* for the two representations. So we require that standard commutative diagram holds:

$$\begin{array}{ccc} \Omega^p & \xrightarrow{a_{qp}} & \Omega^q \\ \rho_p \downarrow & & \downarrow \rho_q \\ \Omega^p & \xrightarrow{a_{qp}} & \Omega^q \end{array} \quad \text{i.e.} \quad \rho_q \circ a_{qp} = a_{qp} \circ \rho_p \quad (87)$$

Thus, we are to solve equation (87) for each  $q, p = 0, \dots, 4$  (and any  $j$  hidden in  $\rho_p$  and  $\rho_q$ ).

Realize that, as is always the case for intertwining operators, solutions constitute a *vector space*, so that we are actually to find *a basis* of the space of solutions.

As an example, for  $a_{12}$  and  $a_{02}$ , we are to solve

$$\rho_1 \circ a_{12} = a_{12} \circ \rho_2 \quad \rho_0 \circ a_{02} = a_{02} \circ \rho_2 \quad (88)$$

for

- matrices  $a_{12}$  and  $a_{02}$  (i.e. for unknown  $\mathbf{c}$ ,  $\mathbf{d}$ ,  $C$  and  $D$  in (86))
- $\rho_0$ ,  $\rho_1$  and  $\rho_2$  displayed in (78), (79) and (80)

Performing this for *rotation* generators  $\rho_0$ ,  $\rho_1$  and  $\rho_2$  *alone* (i.e. only for  $S_j$  operators from (78), (79) and (80)) we get

$$a_{12} = \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ k_1 \mathbb{1} & k_2 \mathbb{1} \end{pmatrix} \quad a_{02} = 0 \equiv (\mathbf{0}^T \quad \mathbf{0}^T) \quad (89)$$

So,  $a_{02}$  is completely fixed by rotations alone (it *vanishes* - there is *no non-vanishing* intertwining  $a_{02}$  *irrespective* of the spacetime under consideration) and in  $a_{12}$  still two free constants survive.

Repeating the same procedure with  $a_{12}$  (already with ansatz (89)) for *boost* generators  $\rho_1$  and  $\rho_2$  we get the following restrictions on  $k_1, k_2$  (see the details in Appendix A.1):

$$k_1 \lambda_2 = k_2 \lambda_2 = k_2 \lambda_1 = 0 \quad (90)$$

(notice that they already depend on  $\lambda_1, \lambda_2$ , i.e. on whether we speak of Lorentzian, Galilean or Carrollian case). This should be analyzed for all three cases (56) - (58) separately and the result is that the *only non-vanishing* operator is (constant multiple of)

$$a_{12} = \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ \mathbb{1} & \mathbb{0} \end{pmatrix} \quad \text{for } \textit{Carrollian} \text{ spacetime} \quad (91)$$

When translated back into the language of complete forms (rather than just their component column), it produces the following (intertwining) mapping  $\Omega^2 \rightarrow \Omega^1$ :

$$a_{12} : \quad dt \wedge \mathbf{a} \cdot d\mathbf{r} + \mathbf{b} \cdot d\mathbf{S} \mapsto \mathbf{a} \cdot d\mathbf{r} \quad \text{for } \textit{Carrollian} \text{ spacetime} \quad (92)$$

In what follows we perform similar calculation for *all*  $a_{qp}$  (and *all three* spacetimes).

### 3.2 General structure of all $a_{qp}$ (no invariance, yet)

Apriori, just from the parametrization structure (85) of spaces  $\Omega^q$  and  $\Omega^p$  (with no intertwining properties, yet) the operators  $a_{qp}$

$$a_{00} \quad a_{01} \quad a_{02} \quad a_{03} \quad a_{04} \quad (93)$$

$$a_{10} \quad a_{11} \quad a_{12} \quad a_{13} \quad a_{14} \quad (94)$$

$$a_{20} \quad a_{21} \quad a_{22} \quad a_{23} \quad a_{24} \quad (95)$$

$$a_{30} \quad a_{31} \quad a_{32} \quad a_{33} \quad a_{34} \quad (96)$$

$$a_{40} \quad a_{41} \quad a_{42} \quad a_{43} \quad a_{44} \quad (97)$$

should be parametrized as follows (we already encountered  $a_{12}$  and  $a_{02}$  in (86)):

$$k \quad (k \ \mathbf{c}^T) \quad (\mathbf{c}^T \ \mathbf{d}^T) \quad (k \ \mathbf{c}^T) \quad k \quad (98)$$

$$\begin{pmatrix} k \\ \mathbf{c} \end{pmatrix} \quad \begin{pmatrix} k \ \mathbf{c}^T \\ \mathbf{d} \ C \end{pmatrix} \quad \begin{pmatrix} \mathbf{c}^T \ \mathbf{d}^T \\ C \ D \end{pmatrix} \quad \begin{pmatrix} k \ \mathbf{c}^T \\ \mathbf{d} \ C \end{pmatrix} \quad \begin{pmatrix} k \\ \mathbf{c} \end{pmatrix} \quad (99)$$

$$\begin{pmatrix} \mathbf{c} \\ \mathbf{d} \end{pmatrix} \quad \begin{pmatrix} \mathbf{c} \ C \\ \mathbf{d} \ D \end{pmatrix} \quad \begin{pmatrix} C \ D \\ E \ F \end{pmatrix} \quad \begin{pmatrix} \mathbf{c} \ C \\ \mathbf{d} \ D \end{pmatrix} \quad \begin{pmatrix} \mathbf{c} \\ \mathbf{d} \end{pmatrix} \quad (100)$$

$$\begin{pmatrix} k \\ \mathbf{c} \end{pmatrix} \quad \begin{pmatrix} k \ \mathbf{c}^T \\ \mathbf{d} \ C \end{pmatrix} \quad \begin{pmatrix} \mathbf{c}^T \ \mathbf{d}^T \\ C \ D \end{pmatrix} \quad \begin{pmatrix} k \ \mathbf{c}^T \\ \mathbf{d} \ C \end{pmatrix} \quad \begin{pmatrix} k \\ \mathbf{c} \end{pmatrix} \quad (101)$$

$$k \quad (k \ \mathbf{c}^T) \quad (\mathbf{c}^T \ \mathbf{d}^T) \quad (k \ \mathbf{c}^T) \quad k \quad (102)$$

Notice the symmetry of the structure of the matrices: The pattern (type of matrix) happens to be symmetric with respect to both

- reflection across the *central horizontal* axis and
- reflection across the *central vertical* axis.

This is just a simple consequence of the symmetry of parametrizations (85) w.r.t. left-right reflection over the center. So there are altogether 9 *types* of matrices  $a_{qp}$ .

### 3.3 How $a_{qp}$ are restricted by rotation invariance alone

After applying rotational invariance alone (i.e. property (87) with just rotation generators  $S_j$ ), the matrices (98) - (102) simplify a lot. In short the simplification may be described as application of the following rules: All

- scalars remain intact (all scalars are rotation invariant)
- vectors become zero (no rotation invariant vectors except for zero)
- matrices become multiples of unity (no other rotation invariant matrices)

What we get is the following, much simpler, parametrization:

$$k \quad (k \ \mathbf{0}^T) \quad (\mathbf{0}^T \ \mathbf{0}^T) \quad (k \ \mathbf{0}^T) \quad k \quad (103)$$

$$\begin{pmatrix} k \\ \mathbf{0} \end{pmatrix} \quad \begin{pmatrix} k_1 & \mathbf{0}^T \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} \quad \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ k_1 \mathbb{1} & k_2 \mathbb{1} \end{pmatrix} \quad \begin{pmatrix} k_1 & \mathbf{0}^T \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} \quad \begin{pmatrix} k \\ \mathbf{0} \end{pmatrix} \quad (104)$$

$$\begin{pmatrix} \mathbf{0} \\ \mathbf{0} \end{pmatrix} \quad \begin{pmatrix} \mathbf{0} & k_1 \mathbb{1} \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} \quad \begin{pmatrix} k_1 \mathbb{1} & k_2 \mathbb{1} \\ k_3 \mathbb{1} & k_4 \mathbb{1} \end{pmatrix} \quad \begin{pmatrix} \mathbf{0} & k_1 \mathbb{1} \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} \quad \begin{pmatrix} \mathbf{0} \\ \mathbf{0} \end{pmatrix} \quad (105)$$

$$\begin{pmatrix} k \\ \mathbf{0} \end{pmatrix} \quad \begin{pmatrix} k_1 & \mathbf{0}^T \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} \quad \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ k_1 \mathbb{1} & k_2 \mathbb{1} \end{pmatrix} \quad \begin{pmatrix} k_1 & \mathbf{0}^T \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} \quad \begin{pmatrix} k \\ \mathbf{0} \end{pmatrix} \quad (106)$$

$$k \quad (k \ \mathbf{0}^T) \quad (\mathbf{0}^T \ \mathbf{0}^T) \quad (k \ \mathbf{0}^T) \quad k \quad (107)$$

### 3.4 Restrictions on $a_{qp}$ added by boosts

There are further restrictions on operators  $a_{qp}$  displayed in (103) - (107) added by boosts. Since boost generators already depend on  $(\lambda_1, \lambda_2)$ , these restrictions depend on the choice of particular spacetime. So we get, in general, different set of operators  $a_{qp}$  for different spacetimes.

The detailed computations are presented in Appendix A.1.

Here we give the results (and comments).

## 4 Results - non-trivial operators $a_{qp}$

### 4.1 Which operators $a_{qp}$ are trivial

It is evident from structure of (87) that

- *zero* operator (for any pair of spaces,  $\Omega^q \rightarrow \Omega^p$ ) as well as
- *unit* operator within any space ( $\hat{1} : \Omega^p \rightarrow \Omega^p$ )

are solutions, i.e. intertwining operators.

Unit operators form part of those found by our computation; so

- $a_{00}$ ,  $a_{11}$ ,  $a_{22}^{(1)}$ ,  $a_{33}$  and  $a_{44}$  from (201), (228) and (259)

are in a sense trivial and they may be omitted from our further discussion. (And they clearly work in *any* spacetime.)

In what follows we summarize all *non-trivial* intertwining operators found in our systematic computations. They depend, as we have seen, on particular spacetime.

## 4.2 Non-trivial operators $a_{qp}$ in Lorentzian spacetime

In Lorentzian (Minkowski) spacetime we have found the following non-trivial operators:

$$a_{40} = 1 \quad (108)$$

$$a_{31} = \begin{pmatrix} 1 & \mathbf{0}^T \\ \mathbf{0} & \mathbb{1} \end{pmatrix} \quad (109)$$

$$a_{22} = \begin{pmatrix} 0 & \mathbb{1} \\ -\mathbb{1} & 0 \end{pmatrix} \quad (110)$$

$$a_{13} = \begin{pmatrix} 1 & \mathbf{0}^T \\ \mathbf{0} & \mathbb{1} \end{pmatrix} \quad (111)$$

$$a_{04} = 1 \quad (112)$$

(see (201), (229), (232) and (260); we use brief notation  $a_{22}$  for the longer one  $a_{22}^{(2)}$ , ignoring thus the existence of the (trivial) *unit* operator  $a_{22}^{(1)}$ ).

From the matrices displayed above we can read-off how the five operators actually act on corresponding differential forms:

$$a_{40} : \quad f \mapsto f dt \wedge dV \quad (113)$$

$$a_{31} : \quad f dt + \mathbf{a} \cdot d\mathbf{r} \mapsto dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV \quad (114)$$

$$a_{22} : \quad dt \wedge \mathbf{a} \cdot d\mathbf{r} + \mathbf{b} \cdot d\mathbf{S} \mapsto dt \wedge \mathbf{b} \cdot d\mathbf{r} - \mathbf{a} \cdot d\mathbf{S} \quad (115)$$

$$a_{13} : \quad dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV \mapsto f dt + \mathbf{a} \cdot d\mathbf{r} \quad (116)$$

$$a_{04} : \quad f dt \wedge dV \mapsto f \quad (117)$$

Now if we compare this with the action of standard (Minkowskian) *Hodge star* operator  $* \equiv *_{\text{Min}}$  (see e.g. Sec.16.1 in [7]) on forms

$$*f = f dt \wedge dV \quad (118)$$

$$*(f dt + \mathbf{a} \cdot d\mathbf{r}) = dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV \quad (119)$$

$$*(dt \wedge \mathbf{a} \cdot d\mathbf{r} + \mathbf{b} \cdot d\mathbf{S}) = dt \wedge \mathbf{b} \cdot d\mathbf{r} - \mathbf{a} \cdot d\mathbf{S} \quad (120)$$

$$*(dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV) = f dt + \mathbf{a} \cdot d\mathbf{r} \quad (121)$$

$$*(f dt \wedge dV) = -f \quad (122)$$

we see that

- we merely “rediscovered” *standard* Hodge star in Minkowski spacetime
- *nothing more* than the Hodge star is intertwining on forms, there.

Invariance of the Hodge star is well known. So what we *actually* learned about the Minkowski case is that the Hodge star is in fact *the only* relevant operator with the property, there. (And this is fairly plausible result; otherwise we certainly met the other operators in important physics equations.)

As we will see in the next two paragraphs, the situation in Galilean as well as Carrollian spacetimes turns out to be *refreshingly different*.

### 4.3 Non-trivial operators $a_{qp}$ in Galilean spacetime

In *Galilean* spacetime we have found quite a lot of non-trivial operators.

First, the following five *analogs* of Lorentzian ones:

$$a_{40} = 1 \quad (123)$$

$$a_{31} = \begin{pmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & \mathbb{1} \end{pmatrix} \quad (124)$$

$$a_{22} = \begin{pmatrix} 0 & \mathbb{1} \\ 0 & 0 \end{pmatrix} \quad (125)$$

$$a_{13} = \begin{pmatrix} 1 & \mathbf{0}^T \\ \mathbf{0} & 0 \end{pmatrix} \quad (126)$$

$$a_{04} = 1 \quad (127)$$

(see (201), (230), (234) and (261)); we use brief notation  $a_{22}$  for the longer one  $a_{22}^{(2)}$ , ignoring thus the existence of the (trivial) *unit* operator  $a_{22}^{(1)}$ ). From the matrices displayed above we can read-off how the five operators actually act on corresponding differential forms:

$$a_{40} : \quad f \mapsto f dt \wedge dV \quad (128)$$

$$a_{31} : \quad f dt + \mathbf{a} \cdot d\mathbf{r} \mapsto dt \wedge \mathbf{a} \cdot d\mathbf{S} \quad (129)$$

$$a_{22} : \quad dt \wedge \mathbf{a} \cdot d\mathbf{r} + \mathbf{b} \cdot d\mathbf{S} \mapsto dt \wedge \mathbf{b} \cdot d\mathbf{r} \quad (130)$$

$$a_{13} : \quad dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV \mapsto f dt \quad (131)$$

$$a_{04} : \quad f dt \wedge dV \mapsto f \quad (132)$$

In *Lorentzian* case, see (113) - (117), we realized that *this type* of operators was nothing but the *Hodge star* acting on various degrees of forms, see (118) - (122). Here this is, in a sense, a similar story. One can introduce “Galilean Hodge star” in Galilean spacetime (see Appendix A.2 and more details, including motivation, in [17]) and it turns out that (128) - (132) just reproduce the formulas valid for the Galilean Hodge star. (Actually there are as many as two possibilities for the Galilean Hodge described in [17] and the last operator,  $a_{04}$ , reproduces one of them (the other gives zero.) So we can say again, like we did at the end of Section 4.2, that in (128) - (132) we “merely rediscovered” *Galilean* Hodge star in Galilean spacetime.

Now there are, however, in addition to (123) - (127), operators with *no analogy* to Lorentzian situation. They act as follows:

$$\Omega^0 \xrightarrow{a_{10}} \Omega^1 \xrightarrow{a_{21}} \Omega^2 \xrightarrow{a_{32}} \Omega^3 \xrightarrow{a_{43}} \Omega^4 \quad (133)$$

Here

$$a_{10} = \begin{pmatrix} 1 \\ \mathbf{0} \end{pmatrix} \quad (134)$$

$$a_{21} = \begin{pmatrix} \mathbf{0} & \mathbb{1} \\ \mathbf{0} & 0 \end{pmatrix} \quad (135)$$

$$a_{32} = \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ 0 & \mathbb{1} \end{pmatrix} \quad (136)$$

$$a_{43} = \begin{pmatrix} 1 & \mathbf{0}^T \end{pmatrix} \quad (137)$$

(see (217), (251), (243) and (210)). So their actual action on corresponding differential forms reads

$$a_{10} : \quad f \mapsto f dt \quad (138)$$

$$a_{21} : \quad f dt + \mathbf{a} \cdot d\mathbf{r} \mapsto dt \wedge \mathbf{a} \cdot d\mathbf{r} \quad (139)$$

$$a_{32} : \quad dt \wedge \mathbf{a} \cdot d\mathbf{r} + \mathbf{b} \cdot d\mathbf{S} \mapsto dt \wedge \mathbf{b} \cdot d\mathbf{S} \quad (140)$$

$$a_{43} : \quad dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV \mapsto f dt \wedge dV \quad (141)$$

We see that the rule how these operators work is nothing but application of *exterior product*

$$(\dots) \mapsto dt \wedge (\dots) \quad (142)$$

on the form under consideration. This is clearly

- an operator of degree +1 and it is also
- *Galilean-invariant* operation on forms,

since

$$dt' = dt \quad (143)$$

for Galilean boosts (59). (Speak nothing of rotations and translations.)

Finally, it should be noted that there were still two more operators found, namely

$$a_{03} = \begin{pmatrix} 1 & \mathbf{0}^T \end{pmatrix} \quad (144)$$

$$a_{14} = \begin{pmatrix} 1 \\ \mathbf{0} \end{pmatrix} \quad (145)$$

(see (210) and (217)), i.e. with action on forms

$$a_{03} : \quad dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV \mapsto f \quad (146)$$

$$a_{14} : \quad f dt \wedge dV \mapsto f dt \quad (147)$$

Both of them are, however, just *compositions* of operators already described previously, namely

$$a_{03} = a_{04} \circ a_{43} \quad (148)$$

$$a_{14} = a_{10} \circ a_{04} \quad (149)$$

## 4.4 Non-trivial operators $a_{qp}$ in Carrollian spacetime

In *Carrollian* spacetime we have found quite a lot of non-trivial operators, too.

First, the following five *analogs* of Lorentzian ones:

$$a_{40} = 1 \quad (150)$$

$$a_{31} = \begin{pmatrix} 1 & \mathbf{0}^T \\ \mathbf{0} & 0 \end{pmatrix} \quad (151)$$

$$a_{22} = \begin{pmatrix} 0 & 0 \\ \mathbb{1} & 0 \end{pmatrix} \quad (152)$$

$$a_{13} = \begin{pmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & \mathbb{1} \end{pmatrix} \quad (153)$$

$$a_{04} = 1 \quad (154)$$

(see (201), (231), (233) and (262); we use brief notation  $a_{22}$  for the longer one  $a_{22}^{(2)}$ , ignoring thus the existence of the (trivial) *unit* operator  $a_{22}^{(1)}$ ). From the matrices displayed above we can read-off how the five operators actually act on corresponding differential forms:

$$a_{40} : \quad f \mapsto f dt \wedge dV \quad (155)$$

$$a_{31} : \quad f dt + \mathbf{a} \cdot d\mathbf{r} \mapsto f dV \quad (156)$$

$$a_{22} : \quad dt \wedge \mathbf{a} \cdot d\mathbf{r} + \mathbf{b} \cdot d\mathbf{S} \mapsto \mathbf{a} \cdot d\mathbf{S} \quad (157)$$

$$a_{13} : \quad dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV \mapsto \mathbf{a} \cdot d\mathbf{r} \quad (158)$$

$$a_{04} : \quad f dt \wedge dV \mapsto f \quad (159)$$

Here the story is similar to the Galilean case. One can introduce “Carrollian Hodge star” in Carrollian spacetime (see Appendix A.2 and more details, including motivation, in [17]) and it turns out that (155) - (159) just reproduce the formulas valid for the Carrollian Hodge star.

So we can say again, like we did at the end of Sections 4.2 and 4.3, that in (155) - (159) we “merely rediscovered” *Carrollian* Hodge star in Carrollian spacetime.

In addition, again, there are operators with *no analogy* to Lorentzian situation (albeit with a *strong* analogy to the *Galilean* one). They act as follows:

$$\Omega^0 \xleftarrow{a_{01}} \Omega^1 \xleftarrow{a_{12}} \Omega^2 \xleftarrow{a_{23}} \Omega^3 \xleftarrow{a_{34}} \Omega^4 \quad (160)$$

Here

$$a_{01} = \begin{pmatrix} 1 & \mathbf{0}^T \end{pmatrix} \quad (161)$$

$$a_{12} = \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ \mathbb{1} & 0 \end{pmatrix} \quad (162)$$

$$a_{23} = \begin{pmatrix} \mathbf{0} & 0 \\ \mathbf{0} & \mathbb{1} \end{pmatrix} \quad (163)$$

$$a_{34} = \begin{pmatrix} 1 \\ \mathbf{0} \end{pmatrix} \quad (164)$$

(see (208), (241), (253) and (219)) so that their actual action on corresponding differential forms reads

$$a_{01} : \quad f dt + \mathbf{a} \cdot d\mathbf{r} \mapsto f \quad (165)$$

$$a_{12} : \quad dt \wedge \mathbf{a} \cdot d\mathbf{r} + \mathbf{b} \cdot d\mathbf{S} \mapsto \mathbf{a} \cdot d\mathbf{r} \quad (166)$$

$$a_{23} : \quad dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV \mapsto \mathbf{a} \cdot d\mathbf{S} \quad (167)$$

$$a_{34} : \quad f dt \wedge dV \mapsto f dV \quad (168)$$

We can see that the rule how these four operators work is nothing but application of *interior product*

$$(\dots) \mapsto i_{\partial_t}(\dots) \quad (169)$$

on the form under consideration. It is clearly

- an operator of degree  $-1$  and
- it is also *Carrollian-invariant* operation on forms

since

$$\partial_{t'} = \partial_t \quad (170)$$

for Carrollian boost (52) and (53). (Speak nothing of rotations and translations.)

Finally, it should be noted that there were still two more operators found, namely

$$a_{30} = \begin{pmatrix} 1 \\ \mathbf{0} \end{pmatrix} \quad (171)$$

$$a_{41} = \begin{pmatrix} 1 & \mathbf{0}^T \end{pmatrix} \quad (172)$$

(see (208) and (219)), i.e. with action on forms

$$a_{30} : \quad f \mapsto f dV \quad (173)$$

$$a_{41} : \quad f dt + \mathbf{a} \cdot d\mathbf{r} \mapsto f dt \wedge dV \quad (174)$$

Both of them are, however, just *compositions* of operators already described previously, namely

$$a_{30} = a_{34} \circ a_{40} \quad (175)$$

$$a_{41} = a_{40} \circ a_{01} \quad (176)$$

## 4.5 A note on the degree $\pm 1$ operators

In Galilean and Carrollian spacetimes, invariant tensor fields are folklor knowledge.

In Galilean case, say, two of them are, within our notation (in adapted coordinates), tensors

$$\delta^{ij} \partial_i \otimes \partial_j \in \mathcal{T}_0^2 \quad dt \in \mathcal{T}_1^0 \quad (177)$$

In Carrollian case we have similarly

$$\delta_{ij}dx^i \otimes dx^j \in \mathcal{T}_2^0 \quad \partial_t \in \mathcal{T}_0^1 \quad (178)$$

For example, for the Galilean case, we can already see them, to credit just some fathers-founders, in (differently looking) component presentations as

$$g^{\alpha\beta}t_\beta = 0 \quad \text{in 1963 paper [18],} \quad (179)$$

$$h^{ab}t_b = 0 \quad \text{in 1966 paper [19],} \quad (180)$$

$$\gamma^{\alpha\beta}\psi_\beta = 0 \quad \text{in 1972 paper [20],} \quad (181)$$

(they all describe the same equation).

On the other side, among the most recent occurrence, we can mention notation and terminology

$$\text{Galilean :} \quad \gamma \quad \text{spatial cometric} \quad (182)$$

$$\tau \quad \text{clock one-form} \quad (183)$$

$$\text{Carrollian :} \quad h \quad \text{spatial metric} \quad (184)$$

$$\xi \quad \text{carrollian vector field} \quad (185)$$

with

$$\gamma(\tau, \cdot) = 0 \quad h(\xi, \cdot) = 0 \quad (186)$$

from 2020 in [21].

Now a simple observation is that, just being aware of tensor fields  $\tau$  and  $\xi$  from (183) and (185), it should be clear from the very beginning that two *invariant* degree  $\pm 1$  *operators* are available,

$$\tau \wedge (\dots) \text{ on } \Omega(\text{Galilei}) \quad i_\xi(\dots) \text{ on } \Omega(\text{Carroll}) \quad (187)$$

Notice that both may be regarded as *differentials* in corresponding de Rham complex.

Also notice that they have *no counterpart* for forms on *Minkowski* spacetime since there is neither invariant one-form nor invariant vector field, there. We identify these interpretations of our “degree  $\pm 1$  operators” in (142) and (169).

## 4.6 Invariant operators and invariant subspaces

In Galilean and Carrollian spacetimes, some features of resulting invariant operators differ from those in Minkowski spacetime. Namely

- analogs of Hodge star happen to be degenerate
- there are operators connecting spaces which were not connected before.

There is a simple *formal* reason for these features could appear. Namely, the representations of Galilean and Carrollian groups on components of forms possess *nontrivial invariant* subspaces.

Due to Schur's lemma, any intertwining operator between two *irreducible* representations is either zero or isomorphism. So, for irreducible representations, *non-zero* intertwining operators are necessarily isomorphisms, and therefore they operate between spaces of equal dimensions.

Now representations of Lorentz group on the space of components of  $p$ -forms are *irreducible*. Indeed, we see from explicit formulas (namely from (23) and (66) - (70)) that the space is

- either 1-dimensional (0-forms and 4-forms)
- or spanned by *pairs*  $((f, \mathbf{a})$  or  $(\mathbf{a}, \mathbf{b}))$
- the two elements within the pair are *independent* for rotations
- the two elements within the pair *need one another* for boosts
- so for rotations *and boosts* they *need one another*
- so the *whole* representation space is needed
- so the representation is irreducible.

Dimensionality of spaces of components of  $p$ -forms for  $p = 0, \dots, 4$  is

$$1 \quad 4 \quad 6 \quad 4 \quad 1 \tag{188}$$

so, just because of these *dimensions*, non-trivial intertwining operators are only possible between spaces  $\Omega^0 \leftrightarrow \Omega^4$ ,  $\Omega^1 \leftrightarrow \Omega^3$  and within each space,  $\Omega^p \rightarrow \Omega^p$ . This is exactly used by the (Minkowski) Hodge star (the last possibility for  $p = 2$ ). And, as an example, non-zero operator  $\Omega^2 \rightarrow \Omega^3$  *cannot* be intertwining.

For Galilean and Carrollian spacetimes, the situation is different.

There *are*, contrary to Minkowski case, *nontrivial invariant* sub-spaces not only for rotations but also for Galilean (as well as Carrollian) *boosts*.

Namely formulas for infinitesimal rotations and Galilean/Carrollian boosts, based on generators listed in Section 2.4, readily show that, in our language of components for  $p$ -forms (85),

$$f \quad \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} \quad \begin{pmatrix} \mathbf{a} \\ \mathbf{b} \end{pmatrix} \quad \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} \quad f \tag{189}$$

there are invariant subspaces spanned on elements

$$\text{Galilean case} \quad f \quad \begin{pmatrix} f \\ \mathbf{0} \end{pmatrix} \quad \begin{pmatrix} \mathbf{a} \\ \mathbf{0} \end{pmatrix} \quad \begin{pmatrix} 0 \\ \mathbf{a} \end{pmatrix} \quad f \tag{190}$$

$$\text{Carrollian case} \quad f \quad \begin{pmatrix} 0 \\ \mathbf{a} \end{pmatrix} \quad \begin{pmatrix} \mathbf{0} \\ \mathbf{b} \end{pmatrix} \quad \begin{pmatrix} f \\ \mathbf{0} \end{pmatrix} \quad f \tag{191}$$

So, we have *non-trivial* invariant subspaces in  $\Omega^1$ ,  $\Omega^2$  and  $\Omega^3$  and, therefore, representations  $\rho_1$ ,  $\rho_2$  and  $\rho_3$  are not irreducible.

Then, nothing (so simple as in the Lorentzian case) excludes, e.g., a *non-zero intertwining* operator  $\Omega^2 \rightarrow \Omega^3$ . And indeed, we have found, in Galilean case, the operator

$$a_{32} : \begin{pmatrix} \mathbf{a} \\ \mathbf{b} \end{pmatrix} \mapsto \begin{pmatrix} 0 \\ \mathbf{b} \end{pmatrix} \tag{192}$$

(see (136)). Notice that

$$\text{Ker } a_{32} \leftrightarrow \begin{pmatrix} \mathbf{a} \\ \mathbf{0} \end{pmatrix} \quad \text{and} \quad \text{Im } a_{32} \leftrightarrow \begin{pmatrix} 0 \\ \mathbf{a} \end{pmatrix} \quad (193)$$

Kernel and image spaces of *any* intertwining operator are known to be invariant subspaces, and here we see that they coincide with particular invariant subspaces displayed in (190).

Similarly, the new Galilean Hodge star (acting on 1-forms)

$$a_{31} : \begin{pmatrix} f \\ \mathbf{a} \end{pmatrix} \mapsto \begin{pmatrix} 0 \\ \mathbf{a} \end{pmatrix} \quad (194)$$

has no reason to be isomorphism. And it indeed seized its chance:

$$\text{Ker } a_{31} \leftrightarrow \begin{pmatrix} f \\ \mathbf{0} \end{pmatrix} \quad \text{and} \quad \text{Im } a_{31} \leftrightarrow \begin{pmatrix} 0 \\ \mathbf{a} \end{pmatrix} \quad (195)$$

Again, they coincide with particular invariant subspaces displayed in (190).

## 5 Summary and conclusions

In expressing Poincaré invariant equations in terms of differential forms on Minkowski spacetime, Hodge star operator plays a key role. One of decisive reasons for that is its Poincaré invariance, i.e. its commutation with (pull-back of) any Poincaré transformation.

Now the pull-back induces, for each degree  $p$  of differential forms, a (finite-dimensional) representation  $\rho_p$  of the Poincaré group and the commutation of the Hodge operator may be treated as *intertwining* property of the latter.

In Galilean and Carrollian (rather than Minkowski) spacetimes, there is no (standard) Hodge star operator, since there is no (invariant) spacetime metric available.

One can, however, find useful *analogs* of the Hodge star.

In [17] this is done via proper modification of the standard formula (making use of suitable invariant tensors).

Here we pursue the idea of intertwining operators in detail. In particular we try to find *all* (algebraic) intertwining operators  $a_{qp}$  from  $p$ -forms to  $q$ -forms, i.e.

$$a_{qp} : \Omega^p \rightarrow \Omega^q \quad \rho_q \circ a_{qp} = a_{qp} \circ \rho_p \quad (196)$$

see (11).

What we found may be summarized as follows:

1. In *Minkowski* spacetime, the Hodge star is *the only* intertwining operator.
2. In *Galilean* spacetime, there is
  - Galilean Hodge star from [17] plus

- degree +1 operator  $\alpha \mapsto dt \wedge \alpha$ .

3. In *Carrollian* spacetime, there is

- Carrollian Hodge star from [17] plus

- degree  $-1$  operator  $\alpha \mapsto i_{\partial_t} \wedge \alpha$ .

So these (algebraic) operators, when combined with exterior derivative  $d$ , which is *known* to be intertwining (differential) operator as well, lead to systems of (Poincaré, Galilean, Carrollian) *invariant equations*. As an example, equations

$$d * F = -J \quad dF = 0 \quad (197)$$

(discussed in Section 1) correspond to Poincaré/Galilean/Carrollian electrodynamics depending upon *which* Hodge star is used; see more in [17]).

## A Appendices

### A.1 Details of solving equations for $a_{qp}$

Here we provide details of solving equations (87) for  $a_{qp}$ .

As mentioned in Section 3.3, the first step is taking *rotations* generators  $S_j$  alone. This leads to simplification (103) - (107). Here we repeat, starting already with these simplified expressions, the same step with *boosts* generators  $N_j$  alone. Since they already depend on  $(\lambda_1, \lambda_2)$ , results become dependent on particular spacetime.

Due to the symmetry of the pattern mentioned in Section 3.2 we can typically compute several operators at once (perform a single computation valid for a particular group of operators, namely for the “orbit” of the symmetry of the pattern).

All *nonzero* results (including trivial *unit* operators) are highlighted as being boxed.

#### A.1.1 Operators $a_{00}$ , $a_{40}$ , $a_{04}$ and $a_{44}$

For  $a_{00}$ ,  $a_{40}$ ,  $a_{04}$  and  $a_{44}$ , the equations read

$$\rho_0 \circ a_{00} = a_{00} \circ \rho_0 \quad \rho_0 \circ a_{04} = a_{00} \circ \rho_4 \quad (198)$$

$$\rho_4 \circ a_{40} = a_{40} \circ \rho_0 \quad \rho_4 \circ a_{44} = a_{44} \circ \rho_4 \quad (199)$$

However, since

$$\rho_0 = \rho_4 = 0 \quad (200)$$

they are completely void and therefore, in all of them,  $k$  is *arbitrary*. So, we can write explicit results (for the *bases* of corresponding 1-dimensional spaces of intertwining operators)

$$\boxed{a_{00} = a_{04} = a_{40} = a_{44} = 1} \quad \boxed{\text{Lorentz, Galilei, Carroll}} \quad (201)$$

### A.1.2 Operators $a_{02}$ , $a_{20}$ , $a_{42}$ and $a_{24}$

This is even easier. *All* of them *vanish* just because of rotation invariance alone (they are parametrized by two vectors and there is no rotation invariant vector, see (103)) - (107)):

$$a_{02} = a_{42} = (\mathbf{0}^T \quad \mathbf{0}^T) \quad a_{20} = a_{24} = \begin{pmatrix} \mathbf{0} \\ \mathbf{0} \end{pmatrix} \quad \text{Lorentz, Galilei, Carroll} \quad (202)$$

### A.1.3 Operators $a_{01}$ , $a_{03}$ , $a_{41}$ and $a_{43}$

For  $a_{01}$ ,  $a_{03}$ ,  $a_{41}$  and  $a_{43}$ , the equations read

$$\rho_0 \circ a_{01} = a_{01} \circ \rho_1 \quad \rho_0 \circ a_{03} = a_{03} \circ \rho_3 \quad (203)$$

$$\rho_4 \circ a_{41} = a_{41} \circ \rho_1 \quad \rho_4 \circ a_{43} = a_{43} \circ \rho_3 \quad (204)$$

Because of (200) they simplify to

$$0 = a_{01} \circ \rho_1 \quad 0 = a_{03} \circ \rho_3 \quad (205)$$

$$0 = a_{41} \circ \rho_1 \quad 0 = a_{43} \circ \rho_3 \quad (206)$$

From this we see that

$$- a_{01} = a_{41} \text{ and } a_{03} = a_{43}$$

$$- a_{03} \text{ may be produced from } a_{01} \text{ by swapping } \lambda_1 \leftrightarrow \lambda_2 \text{ (see (83))}$$

Therefore we only need to really compute (say)  $a_{01}(\lambda_1, \lambda_2)$ . The left equation in (205) explicitly reads

$$(k \quad \mathbf{0}^T) \begin{pmatrix} 0 & \lambda_2 e_j^T \\ \lambda_1 e_j & 0 \end{pmatrix} = (0 \quad \mathbf{0}^T) \quad \text{i.e.} \quad k\lambda_2 = 0 \quad (207)$$

So, for

$$- \text{Carrollian case } (\lambda_2 = 0) \text{ } k \text{ is arbitrary (} a_{01} \text{ does not vanish)}$$

$$- \text{Lorentzian and Galilean case } (\lambda_2 \neq 0) \text{ } k \text{ is zero (} a_{01} \text{ does vanish).}$$

Just swapping  $(\lambda_1, \lambda_2) \mapsto (\lambda_2, \lambda_1)$  we get that for

$$- \text{Galilean case } (\lambda_1 = 0) \text{ } k \text{ is arbitrary (} a_{03} \text{ does not vanish)}$$

$$- \text{Lorentzian and Carrollian case } (\lambda_2 \neq 0) \text{ } k \text{ is zero (} a_{30} \text{ does vanish).}$$

We have

$$\boxed{a_{01} = a_{41}} = (1 \quad \mathbf{0}^T) \quad \boxed{\text{Carroll}} \quad (208)$$

$$= (0 \quad \mathbf{0}^T) \quad \text{Lorentz, Galilei} \quad (209)$$

$$\boxed{a_{03} = a_{43}} = (1 \quad \mathbf{0}^T) \quad \boxed{\text{Galilei}} \quad (210)$$

$$= (0 \quad \mathbf{0}^T) \quad \text{Lorentz, Carroll} \quad (211)$$

#### A.1.4 Operators $a_{10}$ , $a_{30}$ , $a_{14}$ and $a_{34}$

For  $a_{10}$ ,  $a_{30}$ ,  $a_{14}$  and  $a_{34}$ , the equations read

$$\rho_1 \circ a_{10} = a_{10} \circ \rho_0 \qquad \rho_3 \circ a_{30} = a_{30} \circ \rho_0 \qquad (212)$$

$$\rho_1 \circ a_{14} = a_{14} \circ \rho_4 \qquad \rho_3 \circ a_{34} = a_{34} \circ \rho_4 \qquad (213)$$

Because of (200) they simplify to

$$\rho_1 \circ a_{10} = 0 \qquad \rho_3 \circ a_{30} = 0 \qquad (214)$$

$$\rho_1 \circ a_{14} = 0 \qquad \rho_3 \circ a_{34} = 0 \qquad (215)$$

From this we see that

-  $a_{10} = a_{14}$  and  $a_{30} = a_{34}$

-  $a_{30}$  may be produced from  $a_{10}$  by swapping  $\lambda_1 \leftrightarrow \lambda_2$  (see (83))

Therefore we only need to really compute (say)  $a_{10}(\lambda_1, \lambda_2)$ . The left equation in (214) explicitly reads

$$\begin{pmatrix} 0 & \lambda_2 e_j^T \\ \lambda_1 e_j & 0 \end{pmatrix} \begin{pmatrix} k \\ \mathbf{0} \end{pmatrix} = \begin{pmatrix} 0 \\ \mathbf{0} \end{pmatrix} \qquad \text{i.e.} \qquad k\lambda_1 = 0 \qquad (216)$$

So, for

- Galilean case ( $\lambda_1 = 0$ )  $k$  is arbitrary ( $a_{10}$  *does not* vanish)

- Lorentzian and Carrollian case ( $\lambda_1 \neq 0$ )  $k$  is zero ( $a_{10}$  *does* vanish).

Just swapping  $(\lambda_1, \lambda_2) \mapsto (\lambda_2, \lambda_1)$  we get that for

- Carrollian case ( $\lambda_2 = 0$ )  $k$  is arbitrary ( $a_{30}$  *does not* vanish)

- Lorentzian and Galilean case ( $\lambda_2 \neq 0$ )  $k$  is zero ( $a_{30}$  *does* vanish).

We have

$$\boxed{a_{10} = a_{14}} = \begin{pmatrix} 1 \\ \mathbf{0} \end{pmatrix} \qquad \boxed{\text{Galilei}} \qquad (217)$$

$$= \begin{pmatrix} 0 \\ \mathbf{0} \end{pmatrix} \qquad \text{Lorentz, Carroll} \qquad (218)$$

$$\boxed{a_{30} = a_{34}} = \begin{pmatrix} 1 \\ \mathbf{0} \end{pmatrix} \qquad \boxed{\text{Carroll}} \qquad (219)$$

$$= \begin{pmatrix} 0 \\ \mathbf{0} \end{pmatrix} \qquad \text{Lorentz, Galilei} \qquad (220)$$

#### A.1.5 Operators $a_{11}$ , $a_{33}$ , $a_{13}$ and $a_{31}$

For  $a_{11}$ ,  $a_{33}$ ,  $a_{13}$  and  $a_{31}$ , the equations read

$$\rho_1 \circ a_{11} = a_{11} \circ \rho_1 \qquad \rho_1 \circ a_{13} = a_{13} \circ \rho_3 \qquad (221)$$

$$\rho_3 \circ a_{33} = a_{33} \circ \rho_3 \qquad \rho_3 \circ a_{31} = a_{31} \circ \rho_1 \qquad (222)$$

From this we see that

- we only need to really compute  $a_{11}(\lambda_1, \lambda_2)$  and  $a_{13}(\lambda_1, \lambda_2)$
- $a_{33}$  may be produced from  $a_{11}$  by swapping  $\lambda_1 \leftrightarrow \lambda_2$  (see (83))
- $a_{31}$  may be produced from  $a_{13}$  by swapping  $\lambda_1 \leftrightarrow \lambda_2$  (see (83))

The equation for  $a_{11}$  (left in (221)) explicitly reads

$$\begin{pmatrix} 0 & \lambda_2 e_j^T \\ \lambda_1 e_j & 0 \end{pmatrix} \begin{pmatrix} k_1 & \mathbf{0}^T \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} = \begin{pmatrix} k_1 & \mathbf{0}^T \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} \begin{pmatrix} 0 & \lambda_2 e_j^T \\ \lambda_1 e_j & 0 \end{pmatrix} \quad (223)$$

i.e.

$$k_2 \lambda_2 = k_1 \lambda_2 \quad k_2 \lambda_1 = k_1 \lambda_1 \quad (224)$$

Solution is

$$k_1 = k_2 \quad \text{for all three spacetimes} \quad (225)$$

And since the system (224) remains intact for swapping  $\lambda_1 \leftrightarrow \lambda_2$ , we get  $a_{11} = a_{33}$ .

The equation for  $a_{13}$  (right in (221)) explicitly reads

$$\begin{pmatrix} 0 & \lambda_2 e_j^T \\ \lambda_1 e_j & 0 \end{pmatrix} \begin{pmatrix} k_1 & \mathbf{0}^T \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} = \begin{pmatrix} k_1 & \mathbf{0}^T \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} \begin{pmatrix} 0 & \lambda_1 e_j^T \\ \lambda_2 e_j & 0 \end{pmatrix} \quad (226)$$

i.e.

$$k_2 \lambda_2 = k_1 \lambda_1 \quad (227)$$

Here we have for  $a_{13}$

- $\lambda_1 = \lambda_2 = 1$  (Lorentz)  $\Rightarrow (k_1 = k_2 = \text{arbitrary})$
- $\lambda_1 = 0$  (Galilei)  $\Rightarrow k_2 = 0$  ( $k_1$  arbitrary)
- $\lambda_2 = 0$  (Carroll)  $\Rightarrow k_1 = 0$  ( $k_2$  arbitrary)

Swapping  $\lambda_1 \leftrightarrow \lambda_2$  in (227) produces swapping  $k_1 \leftrightarrow k_2$  for solutions. So it produces as  $a_{31}$  swapping Galilei  $\leftrightarrow$  Carroll for the above conclusion for  $a_{13}$ .

Altogether we come to conclusion:

$$\boxed{a_{11} = a_{33}} = \begin{pmatrix} 1 & \mathbf{0}^T \\ \mathbf{0} & \mathbb{1} \end{pmatrix} \quad \boxed{\text{Lorentz, Galilei, Carroll}} \quad (228)$$

$$\boxed{a_{13}} = \begin{pmatrix} 1 & \mathbf{0}^T \\ \mathbf{0} & \mathbb{1} \end{pmatrix} \quad \boxed{\text{Lorentz}} \quad (229)$$

$$= \begin{pmatrix} 1 & \mathbf{0}^T \\ \mathbf{0} & 0 \end{pmatrix} \quad \boxed{\text{Galilei}} \quad (230)$$

$$= \begin{pmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & \mathbb{1} \end{pmatrix} \quad \boxed{\text{Carroll}} \quad (231)$$

$$\boxed{a_{31}} = \begin{pmatrix} 1 & \mathbf{0}^T \\ \mathbf{0} & \mathbb{1} \end{pmatrix} \quad \boxed{\text{Lorentz}} \quad (232)$$

$$= \begin{pmatrix} 1 & \mathbf{0}^T \\ \mathbf{0} & 0 \end{pmatrix} \quad \boxed{\text{Carroll}} \quad (233)$$

$$= \begin{pmatrix} 0 & \mathbf{0}^T \\ \mathbf{0} & \mathbb{1} \end{pmatrix} \quad \boxed{\text{Galilei}} \quad (234)$$

### A.1.6 Operators $a_{12}$ and $a_{32}$

For  $a_{12}$  and  $a_{32}$ , the equations read

$$\rho_1 \circ a_{12} = a_{12} \circ \rho_2 \quad (235)$$

$$\rho_3 \circ a_{32} = a_{32} \circ \rho_2 \quad (236)$$

Now the swapping argument is more tricky since also  $\rho_2$  changes, so it is safer to solve both equations apart.

The equation for  $a_{12}$  explicitly reads

$$\begin{pmatrix} 0 & \lambda_2 e_j^T \\ \lambda_1 e_j & 0 \end{pmatrix} \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ k_1 \mathbb{1} & k_2 \mathbb{1} \end{pmatrix} = \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ k_1 \mathbb{1} & k_2 \mathbb{1} \end{pmatrix} \begin{pmatrix} 0 & -\lambda_2 l_j \\ \lambda_1 l_j & 0 \end{pmatrix} \quad (237)$$

i.e.

$$k_1 \lambda_2 = k_2 \lambda_2 = k_2 \lambda_1 = 0 \quad (238)$$

This results in

- $k_2 = 0$  in all three cases
- $k_1 = 1$  for  $\lambda_2 = 0$  (Carroll), otherwise (Lorentz, Galilei)  $k_1 = 0$

For  $a_{32}$ , the equation explicitly reads

$$\begin{pmatrix} 0 & \lambda_1 e_j^T \\ \lambda_2 e_j & 0 \end{pmatrix} \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ k_1 \mathbb{1} & k_2 \mathbb{1} \end{pmatrix} = \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ k_1 \mathbb{1} & k_2 \mathbb{1} \end{pmatrix} \begin{pmatrix} 0 & -\lambda_2 l_j \\ \lambda_1 l_j & 0 \end{pmatrix} \quad (239)$$

i.e.

$$k_1 \lambda_1 = k_2 \lambda_1 = k_1 \lambda_2 = 0 \quad (240)$$

This results in

- $k_1 = 0$  in all three cases
- $k_2 = 1$  for  $\lambda_1 = 0$  (Galilei), otherwise (Lorentz, Carroll)  $k_1 = 0$

So, we can conclude, that

$$\boxed{a_{12}} = \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ \mathbb{1} & 0 \end{pmatrix} \quad \boxed{\text{Carroll}} \quad (241)$$

$$= \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ 0 & 0 \end{pmatrix} \quad \text{Lorentz, Galilei} \quad (242)$$

$$\boxed{a_{32}} = \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ 0 & \mathbb{1} \end{pmatrix} \quad \boxed{\text{Galilei}} \quad (243)$$

$$= \begin{pmatrix} \mathbf{0}^T & \mathbf{0}^T \\ 0 & 0 \end{pmatrix} \quad \text{Lorentz, Carroll} \quad (244)$$

### A.1.7 Operators $a_{21}$ and $a_{23}$

For  $a_{21}$  and  $a_{23}$ , the equations read

$$\rho_2 \circ a_{21} = a_{21} \circ \rho_1 \quad (245)$$

$$\rho_2 \circ a_{23} = a_{23} \circ \rho_3 \quad (246)$$

Again the swapping argument is more tricky since also  $\rho_2$  changes, so it is safer to solve both equations apart.

The equation for  $a_{21}$  explicitly reads

$$\begin{pmatrix} 0 & -\lambda_2 l_j \\ \lambda_1 l_j & 0 \end{pmatrix} \begin{pmatrix} \mathbf{0} & k_1 \mathbb{1} \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} = \begin{pmatrix} \mathbf{0} & k_1 \mathbb{1} \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} \begin{pmatrix} 0 & \lambda_2 e_j^T \\ \lambda_1 e_j & 0 \end{pmatrix} \quad (247)$$

i.e.

$$k_1 \lambda_1 = k_2 \lambda_2 = k_2 \lambda_1 = 0 \quad (248)$$

This results in

- $k_2 = 0$  in all three cases
- $k_1 = 1$  for  $\lambda_1 = 0$  (Galilei), otherwise (Lorentz, Carroll)  $k_1 = 0$

For  $a_{23}$ , the equation explicitly reads

$$\begin{pmatrix} 0 & -\lambda_2 l_j \\ \lambda_1 l_j & 0 \end{pmatrix} \begin{pmatrix} \mathbf{0} & k_1 \mathbb{1} \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} = \begin{pmatrix} \mathbf{0} & k_1 \mathbb{1} \\ \mathbf{0} & k_2 \mathbb{1} \end{pmatrix} \begin{pmatrix} 0 & \lambda_1 e_j^T \\ \lambda_2 e_j & 0 \end{pmatrix} \quad (249)$$

i.e.

$$k_1 \lambda_1 = k_2 \lambda_2 = k_1 \lambda_2 = 0 \quad (250)$$

This results in

- $k_1 = 0$  in all three cases
- $k_2 = 1$  for  $\lambda_2 = 0$  (Carroll), otherwise (Lorentz, Galilei)  $k_2 = 0$

So, we can conclude, that

$$\boxed{a_{21}} = \begin{pmatrix} \mathbf{0} & \mathbb{1} \\ \mathbf{0} & 0 \end{pmatrix} \quad \boxed{\text{Galilei}} \quad (251)$$

$$= \begin{pmatrix} \mathbf{0} & 0 \\ \mathbf{0} & 0 \end{pmatrix} \quad \text{Lorentz, Carroll} \quad (252)$$

$$\boxed{a_{23}} = \begin{pmatrix} \mathbf{0} & 0 \\ \mathbf{0} & \mathbb{1} \end{pmatrix} \quad \boxed{\text{Carroll}} \quad (253)$$

$$= \begin{pmatrix} \mathbf{0} & 0 \\ \mathbf{0} & 0 \end{pmatrix} \quad \text{Lorentz, Galilei} \quad (254)$$

### A.1.8 Operator $a_{22}$

For  $a_{22}$ , the equation reads

$$\rho_2 \circ a_{22} = a_{22} \circ \rho_2 \quad (255)$$

Explicitly:

$$\begin{pmatrix} 0 & -\lambda_2 l_j \\ \lambda_1 l_j & 0 \end{pmatrix} \begin{pmatrix} k_1 \mathbb{1} & k_2 \mathbb{1} \\ k_3 \mathbb{1} & k_4 \mathbb{1} \end{pmatrix} = \begin{pmatrix} k_1 \mathbb{1} & k_2 \mathbb{1} \\ k_3 \mathbb{1} & k_4 \mathbb{1} \end{pmatrix} \begin{pmatrix} 0 & -\lambda_2 l_j \\ \lambda_1 l_j & 0 \end{pmatrix} \quad (256)$$

i.e.

$$k_1 \lambda_2 = k_4 \lambda_2 \quad k_1 \lambda_1 = k_4 \lambda_1 \quad k_2 \lambda_1 = -k_3 \lambda_2 \quad (257)$$

This results in

- $k_1 = k_4$  in all three cases (from the first two equations)
- $k_2 = -k_3$  (Lorentz),  $k_3 = 0$  (Galilei),  $k_2 = 0$  (Carroll)

So, we can conclude, that

$$a_{22} = q_1 a_{22}^{(1)} + q_2 a_{22}^{(2)} \quad (258)$$

where

$$\boxed{a_{22}^{(1)}} = \begin{pmatrix} \mathbb{1} & 0 \\ 0 & \mathbb{1} \end{pmatrix} \quad \boxed{\text{Lorentz, Galilei, Carroll}} \quad (259)$$

$$\boxed{a_{22}^{(2)}} = \begin{pmatrix} 0 & \mathbb{1} \\ -\mathbb{1} & 0 \end{pmatrix} \quad \boxed{\text{Lorentz}} \quad (260)$$

$$= \begin{pmatrix} 0 & \mathbb{1} \\ 0 & 0 \end{pmatrix} \quad \boxed{\text{Galilei}} \quad (261)$$

$$= \begin{pmatrix} 0 & 0 \\ \mathbb{1} & 0 \end{pmatrix} \quad \boxed{\text{Carroll}} \quad (262)$$

## A.2 Galilean and Carrollian Hodge star operators

In [17], Galilean and Carrollian invariant ‘‘Hodge star’’ operators were proposed via appropriate modification of the standard Hodge operator formula. In order to compare the results obtained in this way with our intertwining operators discussed in this paper, we list the results from [17], here. Just for the sake of completeness we start with the ‘‘standard’’ (Minkowski) Hodge star (see Section 16.1 in [7]).

*Poincaré* invariant Hodge star:

$$*f = f dt \wedge dV \quad (263)$$

$$*(f dt + \mathbf{a} \cdot d\mathbf{r}) = dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV \quad (264)$$

$$*(dt \wedge \mathbf{a} \cdot d\mathbf{r} + \mathbf{b} \cdot d\mathbf{S}) = dt \wedge \mathbf{b} \cdot d\mathbf{r} - \mathbf{a} \cdot d\mathbf{S} \quad (265)$$

$$*(dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV) = f dt + \mathbf{a} \cdot d\mathbf{r} \quad (266)$$

$$*(f dt \wedge dV) = -f \quad (267)$$

*Galilean* invariant Hodge star:

$$*f = f dt \wedge dV \quad (268)$$

$$*(f dt + \mathbf{a} \cdot d\mathbf{r}) = dt \wedge \mathbf{a} \cdot d\mathbf{S} \quad (269)$$

$$*(dt \wedge \mathbf{a} \cdot d\mathbf{r} + \mathbf{b} \cdot d\mathbf{S}) = dt \wedge \mathbf{b} \cdot d\mathbf{r} \quad (270)$$

$$*(dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV) = f dt \quad (271)$$

$$*(f dt \wedge dV) = -f \quad (272)$$

*Carrollian* invariant Hodge star:

$$*f = f dt \wedge dV \quad (273)$$

$$*(f dt + \mathbf{a} \cdot d\mathbf{r}) = f dV \quad (274)$$

$$*(dt \wedge \mathbf{a} \cdot d\mathbf{r} + \mathbf{b} \cdot d\mathbf{S}) = -\mathbf{a} \cdot d\mathbf{S} \quad (275)$$

$$*(dt \wedge \mathbf{a} \cdot d\mathbf{S} + f dV) = \mathbf{a} \cdot d\mathbf{r} \quad (276)$$

$$*(f dt \wedge dV) = -f \quad (277)$$

In fact, for both new cases, Galilean as well as Carrollian, there are as many as *two* formulas for each degree of forms and some of them lead to zero operator. What is displayed here is always the *nonzero* choice.

## Acknowledgement

The author acknowledges support from grant VEGA 1/0703/20.

## References

- [1] P.Bamberg, S.Sternberg: *A Course in Mathematics for Students of Physics*, Cambridge University Press, Cambridge, 1990
- [2] B.F.Schutz: *Geometrical Methods of Mathematical Physics*, Cambridge University Press, Cambridge, 1982
- [3] M.Crampin, F.A.E.Pirani: *Applicable Differential Geometry*, Cambridge University Press, Cambridge, 1986
- [4] M.Göckeler, T.Schücker: *Differential Geometry, Gauge Theories and Gravity*, Cambridge University Press, Cambridge, 1987
- [5] T.Frankel: *The Geometry of Physics, An Introduction* Cambridge University Press 1997 (1-st ed), 2004 (2-nd ed), 2012 (3-rd ed)
- [6] A.Trautman: *Differential Geometry for Physicists*, Napoli, Bibliopolis 1984

- [7] M.Fecko: *Differential Geometry and Lie Groups for Physicists*,  
Cambridge University Press, Cambridge, 2006 (paperback 2011)
- [8] Ch.Duval, G.W.Gibbons, P.A.Horvathy, P.M.Zhang:  
Carroll versus Newton and Galilei: two dual non-Einsteinian concepts of time,  
*Classical and Quantum Gravity*, Volume 31, Number 8 (2014)  
arXiv:1402.0657 [gr-qc]
- [9] D.Van den Bleeken, C.Yunus:  
Newton-Cartan, Galileo-Maxwell and Kaluza-Klein,  
*Classical and Quantum Gravity*, 33, 13 (2016)  
arXiv:1512.03799v2 [math-ph]
- [10] J.Figueroa-O’Farrill, R.Grassie, S.Prohazka: Geometry and BMS Lie Algebras  
of Spatially Isotropic Homogeneous Spacetimes,  
*Journal of High Energy Physics* 119 (2019); arXiv:2009.01948 [hep-th]
- [11] L.Ciambelli, R.G.Leigh, Ch.Marteau and P.M.Petropoulos:  
Carroll structures, null geometry, and conformal isometries,  
*Physical Review D* 100, 046010 (2019); arXiv:1905.02221 [hep-th]
- [12] K.Morand:  
Embedding Galilean and Carrollian geometries I. Gravitational waves,  
*Journal of Mathematical Physics* 61, 082502 (2020)  
arXiv:1811.12681v2 [hep-th]
- [13] K.Banerjee, R.Basu, A.Mehra, A.Mohan, A.Sharma:  
Interacting Conformal Carrollian Theories: Cues from Electrodynamics,  
*Physical Review D* 103, 105001 (2021); arXiv:2008.02829v4 [hep-th]
- [14] M.Henneaux, P.Salgado-Rebolledo:  
Carroll contractions of Lorentz-invariant theories,  
*Journal of High Energy Physics* 180 (2021); arXiv:2109.06708 [hep-th]
- [15] R.Grassie: Beyond Lorentzian Symmetry,  
PhD Thesis, University of Edinburgh (2021)  
arXiv:2107.09495v1 [hep-th]
- [16] D.Hansen: Beyond Lorentzian Physics,  
PhD Thesis, ETH Zurich (2021)  
<https://www.research-collection.ethz.ch/handle/20.500.11850/488630>
- [17] M. Fecko: Galilean and Carrollian invariant Hodge star operators,  
arXiv:2206.09788 [math-ph] (2022)
- [18] A.Trautman: Sur la théorie newtonienne de la gravitation,  
*C.R.Acad.Sci. Paris*, t.257, p.617-720 (1963)

- [19] A.Trautman: Comparison of Newtonian and Relativistic Theories of Space-Time, pp. 413–425 in: *Perspectives in Geometry and Relativity*, Essays in honor of V.Hlavatý, ed. by B. Hoffmann, Indiana Univ. Press, Bloomington, 1966.
- [20] H.P.Künzle: Galilei and Lorentz structures on spacetime: Comparison of the corresponding geometry and physics, Ann. Inst. H. Poincaré. Phys. Théor. 17, 4, 337-362. (1972)
- [21] J.Figueroa-O’Farrill:  
On the intrinsic torsion of spacetime structures,  
Report number EMPG-20-14; arXiv:2009.01948 [hep-th] (2020)
- [22] J.-M.Lévy Leblond: Une nouvelle limite non-relativiste du groupe de Poincaré, Annales de l’ I.H.P. Physique théorique vol. 3, no. 1, pp. 1-12, (1965)