

Frugal Splitting Operators: Representation, Minimal Lifting and Convergence

Martin Morin *

Sebastian Banert*

Pontus Giselsson*

Abstract

We consider frugal splitting operators for finite sum monotone inclusion problems, i.e., splitting operators that use exactly one direct or resolvent evaluation of each operator of the sum. A novel representation of these operators in terms of what we call a generalized primal-dual resolvent is presented. This representation reveals a number of new results regarding lifting numbers, existence of solution maps, and parallelizability of the forward/backward evaluations. We show that the minimal lifting is $n - 1 - f$ where n is the number of monotone operators and f is the number of direct evaluations in the splitting. Furthermore, we show that this lifting number is only achievable as long as the first and last evaluation are resolvent evaluations. In the case of frugal resolvent splitting operators, these results recovers the results of Ryu and Malitsky–Tam. The representation also enables a unified convergence analysis and we present a generally applicable theorem for the convergence and Fejér monotonicity of fixed point iterations of frugal splitting operators with cocoercive direct evaluations. We conclude by constructing a new convergent and parallelizable frugal splitting operator with minimal lifting.

1 Introduction

It is well known that a zero of a maximally monotone operator can be found by performing a fixed point iteration of the resolvent of the operator [1]. However, the resolvent of an operator is not always easily computable, even in the case when the operator is a finite sum of maximally monotone operators for which each resolvent is easily computable. This has led to the development of splitting methods that use each term separately to form convergent fixed point iterations that find a zero of the sum of the operators. In this work, we will consider a general class of splitting operators for finite sums of maximally monotone operators which we call *frugal splitting operators*. Informally, the class contains all operators whose fixed points encode the zeros of the sum of the monotone operators and can be computed with exactly one evaluation of each operator, either directly or via a resolvent. Apart from the operator evaluations, only predetermined linear combinations of the input and operator evaluations are allowed. The class covers the classic Douglas–Rachford [2] and forward-backward operators [3, 4] along with many others, for instance [5–16].

We provide an equivalence between frugal splitting operators and a class of operators we call *generalized primal-dual resolvents*. The generalized primal-dual resolvents are inspired by the works of [17–20] that also made similar generalizations of the resolvent. These works are powerful modeling tools able to capture many different algorithms but our resolvent generalization is the first to provably fully cover the class of frugal splitting operators. This novel representation is a key difference to the related works of [5, 6] which examined frugal resolvent splitting operators, i.e., frugal

*Department of Automatic Control, Lund University, (martin.morin@control.lth.se, sebastian.banert@control.lth.se, pontus.giselsson@control.lth.se)

splitting operators that only use resolvents and no direct evaluations. These works focused on the *lifting number* of frugal splitting operators and, while we also provide minimal lifting results, our representation allows us to easily design new splitting operators and analyze the convergence of their fixed point iterations in a general setting. It also allows us to relax one assumption of [5, 6] regarding the existence of a *solution map*. The existence of a solution map for any frugal splitting operator is evident directly from the representation, and so are properties like step-sizes and parallelizability of the individual resolvent/direct evaluations.

The general idea behind lifting is to trade computational complexity for storage complexity by creating an easier to solve problem in some higher dimensional space. In the context of frugal splitting operators, this means that, while the monotone inclusion problem lies in some Hilbert space \mathcal{H} , the splitting operator maps to and from \mathcal{H}^d for some non-zero natural number d which we call the lifting number. For example, the Douglas–Rachford and forward-backward splitting operators have lifting number one while the splitting operators of the primal-dual method of Chambolle–Pock [7] have lifting number two².

In the three operator case, Ryu [5] showed that the smallest possible lifting number for a frugal resolvent splitting operator is two. Malitsky and Tam [6] later expanded this result to sums of n maximally monotone operators and found that the minimal lifting is $n - 1$. In this paper, we show via a simple rank constraint on a real matrix in the generalized primal-dual resolvent that this lifting number can be reduced to $n - 1 - f$ where f is the number of operators that are evaluated directly and not via resolvents. We also show that the minimal lifting number is dependent on the order of the direct and resolvent evaluations. In particular, if the first or last operator is evaluated directly, the minimal lifting under these assumptions is $n - f$ instead of $n - 1 - f$. An example of this can be found in three operator splitting of Davis and Yin [8] which places its only direct evaluation second which allows them to achieve a lifting number of one. This is to be compared to the primal-dual methods of Vũ and Condat [9, 10] which perform a direct evaluation first and hence require the higher lifting number of two in the three operator case.

We provide sufficient conditions for the convergence of fixed point iteration of a frugal splitting operator with cocoercive direct evaluations and show that the generated sequence is Fejér monotone w.r.t. to the fixed points of the splitting. There are a number of different general or unified approaches for analyzing algorithm classes [17–25]. Many of these approaches can be applied to a frugal resolvent splitting as well but our analysis has the advantage that it is performed directly on the generalized primal-dual resolvent representation. Both the conditions for convergence and the conditions for a generalized primal-dual resolvent to be a frugal resolvent splitting are then constraints on the same set of matrices, which simplifies the design of new frugal resolvent splittings. As an example, we construct a new frugal splitting operator with resolvent and direct evaluations that is convergent and parallelizable. This is in contrast to the minimal lifting method of [6] that only uses backward evaluations and requires sequential evaluation of the resolvents.

1.1 Outline

In Section 2, we introduce some preliminary notation and results together with the main monotone inclusion problem. We properly define and discuss our definition of a frugal splitting operator in Section 3. Our representation theorem can be found in Section 4. Section 5 contains some general remarks on the representation: the relationship between fixed points of the splitting and solutions to the monotone inclusion is proven, how resolvent step-sizes and the dependency between forward/backward evaluation is encoded in the representation is demonstrated, and how a representation can be derived is illustrated via an example. The minimal lifting results in terms of a rank bound on a structured matrix can be found in Section 6. This result recovers the results of Malitsky and Tam [6] and show that the minimal lifting number depends on whether the first or last operator

²The Chambolle–Pock method consider monotone inclusion problem that allow for composition with some linear operators. We will therefore implicitly assume that the linear operators are identity when discussing Chambolle–Pock or other similar methods for monotone inclusion with composition.

evaluation is a forward or backward evaluation. Convergence under cocoercive forward evaluations is proved in Section 7. In Section 7.1 we apply the convergence theorem and reestablish the convergence criterion of the three operator splitting of Davis and Yin [8] as well as provide conditions for the convergence of a fixed point iteration of forward-backward with Nesterov-like momentum. The last part of the paper, Section 8, contains the construction of a new frugal splitting operator with minimal lifting and parallelizable forward/backward evaluations. The paper is ended with a short conclusion in Section 9. In the accompanying supplement many more examples of representations of frugal splitting operators and application of our convergence theorem can be found.

2 Preliminaries

Let $\mathbb{N} = \{0, 1, \dots\}$ be the set of natural numbers and $\mathbb{N}_+ = \{1, 2, \dots\}$ be the set of non-zero natural numbers. The cardinality of a set A is denoted by $|A|$. A subset B of A is denoted by $B \subseteq A$ while a strict subset is denoted by $B \subset A$.

Let \mathbb{R} be the set of real numbers. We refer to the range and kernel of a matrix $A \in \mathbb{R}^{n \times m}$ as $\text{ran } A$ and $\text{ker } A$ respectively. These are linear subspaces of \mathbb{R}^n and \mathbb{R}^m respectively and their orthogonal complement with respect to the standard Euclidean inner product are denoted by $(\text{ran } A)^\perp$ and $(\text{ker } A)^\perp$ respectively, which also are linear subspaces.

Let \mathcal{H} be a real Hilbert space with $\dim \mathcal{H} > 0$. The inner product and norm on \mathcal{H} are given by $\langle \cdot, \cdot \rangle$ and $\|\cdot\|$ respectively. With $H: \mathcal{H} \rightarrow \mathcal{H}$ being a bounded linear operator, we let $\langle \cdot, \cdot \rangle_H = \langle H(\cdot), \cdot \rangle$ and $\|\cdot\|_H = \langle H(\cdot), \cdot \rangle$. If H is self-adjoint and strongly positive then $\langle \cdot, \cdot \rangle_H$ is a inner product and $\|\cdot\|_H$ is a norm.

An operator $A: \mathcal{H} \rightarrow 2^{\mathcal{H}}$ is *monotone* if

$$\langle u - v, x - y \rangle \geq 0$$

for all $x, y \in \mathcal{H}$ and $u \in Ax$ and $v \in Ay$. An operator $A: \mathcal{H} \rightarrow 2^{\mathcal{H}}$ is μ -*strongly monotone* where $\mu > 0$ if

$$\langle u - v, x - y \rangle \geq \mu \|x - y\|^2$$

for all $x, y \in \mathcal{H}$ and $u \in Ax$ and $v \in Ay$. A monotone operator is *maximal* if its graph is not contained in the graph of any other monotone operator. An operator $A: \mathcal{H} \rightarrow 2^{\mathcal{H}}$ is said to have *full domain* if $Ax \neq \emptyset$ for all $x \in \mathcal{H}$. An operator $A: \mathcal{H} \rightarrow 2^{\mathcal{H}}$ is *single valued* if Ax is a singleton for all $x \in \mathcal{H}$; we will make no distinction between single valued operators $A: \mathcal{H} \rightarrow 2^{\mathcal{H}}$ and ordinary mappings $A: \mathcal{H} \rightarrow \mathcal{H}$. An operator $A: \mathcal{H} \rightarrow \mathcal{H}$ is β -*cocoercive* if

$$\langle Ax - Ay, x - y \rangle \geq \beta \|Ax - Ay\|^2$$

for all $x, y \in \mathcal{H}$. The inverse of a β -cocoercive operator is β -strongly monotone. The *resolvent* of a maximally monotone operator is $J_A = (\text{Id} + A)^{-1}$.

Let U, V and W be sets and define the operator $A: U \times W \rightarrow V$. With the notation $A_w u = v$ we mean $A(u, w) = v$ and we define $A_w = A(\cdot, w)$. Since A_w is an operator from U to V , instead of writing $A: U \times W \rightarrow V$ we will say that $A_{(\cdot)}: U \rightarrow V$ is *parameterized* by W .

Definition 2.1 (Matrix as Operator). *Let $B \in \mathbb{R}^{n \times m}$ and $z = (z_1, \dots, z_m) \in \mathcal{H}^m$. We define Bz as³*

$$Bz = \begin{bmatrix} B_{11}z_1 + \dots + B_{1m}z_m \\ \vdots \\ B_{n1}z_1 + \dots + B_{nm}z_m \end{bmatrix}.$$

Lemma 2.1. *Let $A \in \mathbb{R}^{n \times m}$ and $B \in \mathbb{R}^{n \times d}$. If $\text{ran } A \subseteq \text{ran } B$, there exists a unique $S \in \mathbb{R}^{d \times m}$ such that $A = BS$ and $\text{ran } S \subseteq (\text{ker } B)^\perp$. If $\text{ran } A = \text{ran } B$, such an S satisfies $\text{ran } S = (\text{ker } B)^\perp$.*

³This operator could more accurately be represented by the matrix $B \otimes \text{Id}$ where \otimes is the tensor product.

Proof. Since $\text{ran } A \subseteq \text{ran } B$, the columns of A lie in the span of the columns of B . This means that there exists a matrix $S \in \mathbb{R}^{d \times m}$ such that $A = BS$.

Let S be a matrix such that $A = BS$. If $\text{ran } S \supset (\ker B)^\perp$, we can always construct a different S' such that $\text{ran } S' \subseteq (\ker B)^\perp$. Let Π_\perp be the orthogonal projection onto $(\ker B)^\perp$ in the standard Euclidean inner product, then $B = B\Pi_\perp$ and hence $A = BS = B\Pi_\perp S = BS'$ where we define $S' = \Pi_\perp S$. It is clear that $\text{ran } S' \subseteq (\ker B)^\perp$.

To show uniqueness, let S and S' be such that $A = BS = BS'$ and $\text{ran } S \subseteq (\ker B)^\perp$ and $\text{ran } S' \subseteq (\ker B)^\perp$ and let $x \in \mathbb{R}^m$ be such that $Sx \neq S'x$. Since $Sx - S'x \notin \ker B$ we have

$$0 \neq B(Sx - S'x) = BSx - BS'x = Ax - Ax = 0$$

which clearly is a contradiction and Sx and $S'x$ must then be equal for all $x \in \mathbb{R}^m$ and hence is $S = S'$.

To show the last statement, assume $\text{ran } A = \text{ran } B$ and let S be the unique matrix that satisfies $A = BS$ and $\text{ran } S \subseteq (\ker B)^\perp$. Assume $\text{ran } S \subset (\ker B)^\perp$, then $\text{rank } S < \dim(\ker B)^\perp = \text{rank } B$ and $\text{rank } A \leq \min(\text{rank } B, \text{rank } S) < \text{rank } B$. However, this contradicts $\text{ran } A = \text{ran } B$ and hence $\text{ran } S = (\ker B)^\perp$. \square

Lemma 2.2. *Let $A \in \mathbb{R}^{m \times n}$ and $B \in \mathbb{R}^{d \times n}$. If $\ker A \supseteq \ker B$, there exists a unique $S \in \mathbb{R}^{m \times d}$ such that $A = SB$ and $\ker S \supseteq (\text{ran } B)^\perp$. If $\ker A = \ker B$ such an S satisfies $\ker S = (\text{ran } B)^\perp$.*

Proof. This is simply the dual to Lemma 2.1. If $\ker A \supseteq \ker B$ then $\text{ran } A^T \subseteq \text{ran } B^T$ and Lemma 2.1 then implies the existence of a unique $S^T \in \mathbb{R}^{d \times m}$ with $\text{ran } S^T \subseteq (\ker B^T)^\perp$ such that $A^T = B^T S^T$ or equivalently $A = SB$. Since $\text{ran } S^T = (\ker S)^\perp$ and $(\ker B^T)^\perp = \text{ran } B$ we have $(\ker S)^\perp \subseteq \text{ran } B$ or equivalently $\ker S \supseteq (\text{ran } B)^\perp$. If $\ker A = \ker B$ then $\text{ran } A^T = \text{ran } B^T$ and Lemma 2.1 then yields $\ker S = (\text{ran } B)^\perp$. \square

2.1 Problem

This paper concerns the problem of finding a zero of a finite sum of operators,

$$\text{find } x \in \mathcal{H} \text{ such that } 0 \in \sum_{i=1}^n A_i x \tag{1}$$

where $A_i: \mathcal{H} \rightarrow 2^{\mathcal{H}}$ is maximally monotone for all $i \in \{1, \dots, n\}$ and some of the operators are single valued, i.e., $A_i: \mathcal{H} \rightarrow \mathcal{H}$ for $i \in F$ for some $F \subseteq \{1, \dots, n\}$. Note, F is allowed to be empty.

The aim is to examine a class of iterative methods for solving this problem and in order to simplify the notation we make the following definition.

Definition 2.2 (Operator Tuples). *With $F \subseteq \{1, \dots, n\}$,*

$$\mathcal{A}_n^F = \{(A_1, \dots, A_n) \in \mathcal{M}(\mathcal{H})^n \mid A_i \text{ is single valued for all } i \in F\}$$

where $\mathcal{M}(\mathcal{H})$ is the set of all maximally monotone operators from \mathcal{H} to $2^{\mathcal{H}}$. Furthermore, we define $\mathcal{A}_n = \mathcal{A}_n^\emptyset$ and we note that $\mathcal{A}_n^F \subseteq \mathcal{A}_n$ for all $F \subseteq \{1, \dots, n\}$.

With each tuple $A = (A_1, \dots, A_n) \in \mathcal{A}_n^F$ we can associate a monotone inclusion problem of the form (1). This allows us to consider all solution methods of (1) to be parameterized by \mathcal{A}_n^F , i.e., every solution method takes a tuple $A \in \mathcal{A}_n^F$ as input—alongside any other algorithm parameters—and generates a solution to the monotone inclusion problem associated with A . We will give the exact definition of the considered class of solution methods in Section 3. The remainder of this section will be used for further preparation and introduction of some key operators.

Instead of solving (1) directly, we will instead work with a family of primal-dual problems. Let $p \in \{1, \dots, n\}$, the primal-dual problem is then

$$\text{find } (y_1, \dots, y_n) \in \mathcal{H}^n \text{ such that } \begin{cases} 0 \in A_i^{-1}y_i - y_p \text{ for all } i \in \{1, \dots, n\} \setminus \{p\}, \\ 0 \in A_p y_p + \sum_{j \in \{1, \dots, n\} \setminus \{p\}} y_j. \end{cases} \quad (2)$$

We call p the primal index since the corresponding variable, y_p , solves the primal-problem (1). The equivalence between (2) and (1) is straightforward to show and holds in the sense that if $y_p \in \mathcal{H}$ is a solution to (1) then there exists $y_i \in \mathcal{H}$ for all $i \in \{1, \dots, n\} \setminus \{p\}$ such that (y_1, \dots, y_n) solves (2). Conversely, if $(y_1, \dots, y_n) \in \mathcal{H}^n$ is a solution to (2) then y_p solves (1).

This primal-dual problem is in fact also a monotone inclusion problem and we can identify a primal-dual operator $\Phi_{A,p}: \mathcal{H}^n \rightarrow 2^{\mathcal{H}^n}$ parameterized by \mathcal{A}_n^F and primal index $p \in \{1, \dots, n\}$ such that (2) can be written as

$$\text{find } y \in \mathcal{H}^n \text{ such that } 0 \in \Phi_{A,p}y := \Delta_{A,p}y + \Gamma_p y \quad (3)$$

where $\Delta_{A,p}: \mathcal{H}^n \rightarrow 2^{\mathcal{H}^n}$ and $\Gamma_p \in \mathbb{R}^{n \times n}$ are defined as

$$(\Delta_{A,p}(y_1, \dots, y_n))_i = \begin{cases} A_i^{-1}y_i & \text{if } i \neq p, \\ A_i y_i & \text{if } i = p \end{cases} \quad \text{and} \quad \Gamma_p = R_p - R_p^T \quad (4)$$

where $(\Delta_{A,p}y)_i$ denotes the i th block of $\Delta_{A,p}y \in \mathcal{H}^n$ for all $i \in \{1, \dots, n\}$ and $R_p \in \mathbb{R}^{n \times n}$ is the matrix with ones on the p th row and zeros in all other positions. All of these operators are maximally monotone: Γ_p as an operator on $\mathcal{H} \rightarrow \mathcal{H}$ is bounded linear and skew-adjoint, $\Delta_{A,p}$ is separable w.r.t. the components with each component operator being maximally monotone, and $\Delta_{A,p}$ is the sum of two maximally monotone operators with one having full domain. The primal-dual operator $\Phi_{A,p}$ will feature extensively in the rest of the paper. In fact, we will show that the considered class of operators always can be written as taking a resolvent-like step of the primal-dual operator $\Phi_{A,p}$ for some choice of primal index p .

For illustration, consider the case when $p = n$, then

$$\Delta_{A,n}(y_1, \dots, y_n) = \begin{bmatrix} A_1^{-1}y_1 \\ \vdots \\ A_{n-1}^{-1}y_{n-1} \\ A_n y_n \end{bmatrix} \quad \text{and} \quad \Gamma_n = \begin{bmatrix} 0 & \cdots & 0 & -1 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & 0 & -1 \\ 1 & \cdots & 1 & 0 \end{bmatrix},$$

while if $p = 1$, we get

$$\Delta_{A,n}(y_1, \dots, y_n) = \begin{bmatrix} A_1 y_1 \\ A_2^{-1}y_2 \\ \vdots \\ A_n^{-1}y_n \end{bmatrix} \quad \text{and} \quad \Gamma_n = \begin{bmatrix} 0 & 1 & \cdots & 1 \\ -1 & 0 & \cdots & 0 \\ \vdots & \cdots & \vdots & \vdots \\ -1 & 0 & \cdots & 0 \end{bmatrix}.$$

3 Frugal Splitting Operators

A common way of solving (1) associated with some $A \in \mathcal{A}_n^F$ is with a fixed point iteration. These are methods that, given some initial iterate $z_0 \in \mathcal{H}^d$ and operator $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ parameterized by \mathcal{A}_n^F , iteratively perform

$$z_{k+1} = T_A z_k \quad (5)$$

for $k \in \mathbb{N}$. The operator $T_{(\cdot)}$ is such that $\{z_k\}_{k \in \mathbb{N}}$ converges to a fixed point and that a solution to (1) can be recovered from each fixed point of T_A . However, it is hard to make any interesting statements in this abstract setting without making further assumptions on $T_{(\cdot)}$. For instance, we could choose

$T_A: \mathcal{H} \rightarrow \mathcal{H}$ as the projection onto the solution set of (1). The fixed point iteration would then converge in one iteration but the computation of $T_A z$ might not even be feasible. Therefore, the standard approach when considering these kinds of fixed point iterations is to restrict $T_{(\cdot)}$ to have a known structure and relatively cheap computational cost. With this motivation in mind, we will in this paper consider the following class of operators.

Definition 3.1 (Frugal Splitting Operator). *Let $d \in \mathbb{N}_+$ and $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ be parameterized by \mathcal{A}_n^F . We say that $T_{(\cdot)}$ is a frugal splitting operator over \mathcal{A}_n^F if, for all $A = (A_1, \dots, A_n) \in \mathcal{A}_n^F$,*

- (i) $\text{fix } T_A \neq \emptyset \iff \text{zer } \sum_{i=1}^n A_i \neq \emptyset$,
- (ii) $T_A = \tau_{n, A_n} \circ \tau_{n-1, A_{n-1}} \circ \dots \circ \tau_{1, A_1}$ where $\tau_{k, A_k}: \mathcal{H}^{d_k} \rightarrow \mathcal{H}^{d_{k+1}}$ for all $k \in \{1, \dots, n\}$ with $d_1 = d_{n+1} = d$. For each $k \in \{1, \dots, n\}$, there exist $a_{ij}, b_i, c_j \in \mathbb{R}$ and $\gamma > 0$ for $i \in \{1, \dots, d_{k+1}\}$ and $j \in \{1, \dots, d_k\}$ such that

$$(\tau_{k, A_k}(z_1, \dots, z_{d_k}))_i = \sum_{j=1}^{d_k} a_{ij} z_j + b_i \begin{cases} J_{\gamma A_k}(\sum_{j=1}^{d_k} c_j z_j) & \text{if } k \notin F \\ A_k(\sum_{j=1}^{d_k} c_j z_j) & \text{if } k \in F \end{cases}$$

for all $(z_1, \dots, z_{d_k}) \in \mathcal{H}^{d_k}$ and $i \in \{1, \dots, d_{k+1}\}$ where $(\tau_{k, A_k}(z_1, \dots, z_{d_k}))_i$ denotes the i th block of $\tau_{k, A_k}(z_1, \dots, z_{d_k}) \in \mathcal{H}^{d_{k+1}}$

We call a frugal splitting operator over \mathcal{A}_n a resolvent splitting operator.

In line with [5], we call the class frugal since the computational requirement Definition 3.1(ii) states that each operator in A is evaluated exactly once, either directly or via a resolvent. We will also refer to a direct evaluation as a forward evaluation while a resolvent evaluation will be referred to as a backward step or backward evaluation. Apart from forward and backward evaluations, only predetermined vector additions and scalar multiplications of the input and results of the operator evaluations are allowed. This ensures that the evaluation cost of a frugal splitting operator is a known quantity that is mainly determined by the cost of the operator and resolvent evaluations. We also see that (ii) specifies evaluation order, i.e., it specifies that the operators A_i in the tuple A must be used in the order they appear in A when evaluating T_A . However, since the operators of the original monotone inclusion problem can be arbitrary rearranged, this entails no loss of generality.

This definition is functionally the same as the definitions of Ryu and Malitsky–Tam [6, 25] but with the addition of forward evaluations being allowed. It also covers many classic operators like the operators used in the forward-backward method [3, 4] and Douglas–Rachford method [2], but also operators of more recent methods such that the three operator splitting of Davis and Yin [8] and primal-dual methods in the style of Chambolle and Pock [7] among many others. However, it does not allow for multiple evaluations of an operator and can therefore not cover the forward-backward-forward of Tseng [26] although it does cover the forward-reflected-backward method of Malitsky and Tam [11]. Since it also does not allow for second order information or online search criteria it does not cover Newton’s method, backtracking gradient descent or other similar methods.

Both Ryu [25] and Malitsky–Tam [6] assumed the existence of a computationally tractable *solution map* $S_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}$ that maps fixed points of a splitting operator to solutions of (1). One of our results, Proposition 5.1, shows that this is unnecessary since (i) and (ii) together imply the existence of such a solution map. Also, note that Definition 3.1 only considers the solution encoding and the computational requirements of a splitting and does not make any assumptions regarding the convergence of a fixed point iteration of a splitting. In fact, both [6, 25] provided examples of resolvent splitting operators whose fixed point iteration failed to converge in general. We will therefore analyze fixed point iterations of frugal splitting operators separately in Section 7, there we provide sufficient convergence conditions that are applicable to any frugal splitting operator.

4 Representation of Frugal Splitting Operators

Although Definition 3.1 fully defines all operators we aim to consider, we do not find it conducive for analysis. In this section, we will therefore develop an equivalent representation of the class of

frugal splitting operators that will be used in the subsequent analysis. In Section 5, we will remark on some of the properties we find appealing of this particular representation.

The representation will be in the form of what we call a generalized primal-dual resolvent.

Definition 4.1 (Generalized Primal-Dual Resolvent). *Let $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ be parameterized by \mathcal{A}_n^F . We say that $T_{(\cdot)}$ is a generalized primal-dual resolvent if there exist $p \in \{1, \dots, n\}$, $M \in \mathbb{R}^{n \times n}$, $N \in \mathbb{R}^{n \times d}$, $U \in \mathbb{R}^{d \times d}$ and $V \in \mathbb{R}^{d \times n}$ such that for each $A \in \mathcal{A}_n^F$ and each $z \in \text{dom } T_A$ there exists $y \in \mathcal{H}^n$ such that*

$$\begin{aligned} y &= (M + \Phi_{A,p})^{-1} N z, \\ T_A z &= z - U z + V y, \end{aligned} \tag{6}$$

where $\Phi_{A,p}$ is defined in (3).

We call the tuple (p, M, N, U, V) a representation of $T_{(\cdot)}$.

When $M = N = \gamma^{-1} I$ and $U = V = \theta I$ where $I \in \mathbb{R}^{n \times n}$ is the identity matrix, we see that a generalized primal-dual resolvent becomes an ordinary relaxed resolvent operator of the primal-dual operator $\Phi_{A,p}$,

$$(1 - \theta) \text{Id} + \theta(\gamma^{-1} \text{Id} + \Phi_{A,p})^{-1} \circ \gamma^{-1} \text{Id}$$

where $\theta \in (0, 2]$ and $\gamma > 0$. While the ordinary relaxed resolvent operator is always well defined, the operator $(M + \Phi_{A,p})^{-1} \circ N$ used in (6) is not necessarily single valued or computationally tractable for an arbitrary choice of M , N and tuples $A \in \mathcal{A}_n^F$. However, for the purposes of this paper, it is not necessary to find general conditions for when a generalized primal-dual resolvent is well defined. The objective is to parameterize frugal splitting operators and, since they are single valued and computable by Definition 3.1, if we find conditions for (p, M, N, U, V) to be a representation of frugal splitting operator over \mathcal{A}_n^F it will also guarantee the single valuedness and computational tractability of $(M + \Phi_{A,p})^{-1} \circ N$ for all $A \in \mathcal{A}_n^F$.

All generalized primal-dual resolvents are not frugal splitting operators and further restrictions on the parameters p , M , N , U and V in Definition 4.1 are required. Of these parameters, most of the structural requirements will be on the kernel M and we make the following definition of the necessary kernel structure.

Definition 4.2 (p -Kernel over \mathcal{A}_n^F). *We call a matrix $M \in \mathbb{R}^{n \times n}$ a p -kernel over \mathcal{A}_n^F if $p \notin F$ and $M + \Gamma_p$ is lower triangular with $M_{i,i} \geq 0$ for $i \in \{1, \dots, n\}$ and $M_{i,i} = 0$ if and only if $i \in F$ where $M_{i,i}$ denotes the i th element on the diagonal of M . The matrix Γ_p is defined in (4).*

With this definition in place we can now state our main representation theorem.

Theorem 4.1. *Let $F \subset \{1, \dots, n\}$ and $p \in \{1, \dots, n\} \setminus F$. An operator $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ parameterized by \mathcal{A}_n^F is a frugal splitting operator over \mathcal{A}_n^F if and only if it is a generalized primal-dual resolvent with representation (p, M, N, U, V) where*

- (i) M is a p -kernel over \mathcal{A}_n^F ,
- (ii) $\ker \begin{bmatrix} N & -M \end{bmatrix} \supseteq \ker \begin{bmatrix} U & -V \end{bmatrix}$,
- (iii) $\text{ran } U \supseteq \text{ran } V$,

and $M \in \mathbb{R}^{n \times n}$, $N \in \mathbb{R}^{n \times d}$, $U \in \mathbb{R}^{d \times d}$ and $V \in \mathbb{R}^{d \times n}$.

Proof. Follows directly from Lemmas 4.1 to 4.3 in Section 4.1. □

Note that we require F to be a strict subset of $\{1, \dots, n\}$ and hence there always exists $p \in \{1, \dots, n\} \setminus F$.

Theorem 4.1 gives conditions on all four matrices of a representation (p, M, N, U, V) . However, it turns out that these conditions make the kernel M uniquely defined, given N , U and V . This leads to the following corollary which will be useful both in the examination of minimal lifting in Section 6 and in the convergence analysis in Section 7.

Corollary 4.1. *Let $F \subset \{1, \dots, n\}$ and $p \in \{1, \dots, n\} \setminus F$. A generalized primal-dual resolvent with representation (p, SUP, SU, U, UP) where $U \in \mathbb{R}^{d \times d}$, $S \in \mathbb{R}^{n \times d}$ and $P \in \mathbb{R}^{d \times n}$ is a frugal splitting operator if*

- (i) SUP is a p -kernel over \mathcal{A}_n^F ,
- (ii) $\text{ran } P \subseteq (\ker U)^\perp$ and $\ker S \supseteq (\text{ran } U)^\perp$.

Furthermore, for any frugal splitting operator $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ over \mathcal{A}_n^F with representation (p, M, N, U, V) , there exist matrices $S \in \mathbb{R}^{n \times d}$ and $P \in \mathbb{R}^{d \times n}$ such that $M = SUP$, $N = SU$ and $V = UP$ and the conditions (i) and (ii) are satisfied.

Proof. It is straightforward to verify that a representation (p, SUP, SU, U, UP) that satisfies the conditions of the theorem satisfies the conditions of Theorem 4.1 and hence is a representation of a frugal splitting operator over \mathcal{A}_n^F .

Assume $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ is a frugal splitting operator over \mathcal{A}_n^F , Theorem 4.1 ensures the existence of a representation (p, M, N, U, V) . The theorem further states that $\text{ran } U \supseteq \text{ran } V$ and Lemma 2.1 then proves the unique existence of $P \in \mathbb{R}^{d \times n}$ with $\text{ran } P \subseteq (\ker U)^\perp$ such that

$$V = UP.$$

Theorem 4.1 also states $\ker \begin{bmatrix} U & -V \end{bmatrix} \subseteq \ker \begin{bmatrix} N & -M \end{bmatrix}$ and Lemma 2.2 yields the unique existence of $S \in \mathbb{R}^{n \times d}$ with $\ker S \supseteq (\text{ran} \begin{bmatrix} U & -V \end{bmatrix})^\perp$ such that

$$N = SU \quad \text{and} \quad M = SV = SUP.$$

However, since $\text{ran } U \supseteq \text{ran } V$ we have $\text{ran} \begin{bmatrix} U & -V \end{bmatrix} = \text{ran } U$, this concludes the proof. \square

4.1 Representation Lemmas

The following lemmas together prove Theorem 4.1. They show the existence of a generalized primal-dual resolvent representation of a frugal splitting operator, the properties the matrices of such a representation must have, and the converse statement that any generalized primal-dual resolvent with such properties is a frugal splitting operator.

Lemma 4.1. *Let $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ be a frugal splitting operator over \mathcal{A}_n^F where $F \subset \{1, \dots, n\}$, then $T_{(\cdot)}$ is a generalized primal-dual resolvent. For each $p \in \{1, \dots, n\} \setminus F$, there exists a representation (p, M, N, U, V) of $T_{(\cdot)}$ where M is a p -kernel over \mathcal{A}_n^F .*

Proof. Let $p \in \{1, \dots, n\} \setminus F$, and let $A \in \mathcal{A}_n^F$ and $z \in \text{dom } T_A$ be arbitrary and consider the evaluation of $T_A z$.

From Definition 3.1(ii) we know that $T_A = \tau_{n, A_n} \circ \dots \circ \tau_{1, A_1}$ where $\tau_{i, A_i}: \mathcal{H}^{d_i} \rightarrow \mathcal{H}^{d_{i+1}}$ and $d_1 = d_{n+1} = d$. If we introduce variables for the intermediate results, i.e., $z_1 = z$ and $z_{i+1} = \tau_{i, A_i} z_i$ for all $i \in \{1, \dots, n\}$, we can write $T_A z = z_{n+1}$. Furthermore, from the definition of τ_{i, A_i} in Definition 3.1(ii) we can conclude that for all $i \in \{1, \dots, n\}$ there exist matrices $B_i \in \mathbb{R}^{d_{i+1} \times d_i}$, $C_i \in \mathbb{R}^{d_{i+1} \times 1}$ and $D_i \in \mathbb{R}^{1 \times d_i}$ such that

$$z_{i+1} = \tau_{i, A_i} z_i = \begin{cases} B_i z_i + C_i J_{\gamma_i A_i} D_i z_i & \text{if } i \notin F, \\ B_i z_i + C_i A_i D_i z_i & \text{if } i \in F. \end{cases}$$

Now, for all $i \in \{1, \dots, n\} \setminus (F \cup \{p\})$, we apply the Moreau identity, $J_{\gamma_i A_i} = \text{Id} - \gamma_i J_{\gamma_i^{-1} A_i^{-1}} \circ \gamma_i^{-1} \text{Id}$, and rewrite $\tau_{i, A_i} z_i$ as

$$\begin{aligned} z_{i+1} &= B_i z_i + C_i J_{\gamma_i A_i} D_i z_i \\ &= (B_i + C_i D_i) z_i - (\gamma_i C_i) J_{\gamma_i^{-1} A_i^{-1}} (\gamma_i^{-1} D_i) z_i \\ &= \widehat{B}_i z_i + \widehat{C}_i (l_{i, i} \text{Id} + \widehat{A}_i)^{-1} \widehat{D}_i z_i \end{aligned}$$

where $\widehat{B}_i = B_i + C_i D_i$, $\widehat{C}_i = -\gamma_i C_i$, $\widehat{D}_i = D_i$, $l_{i,i} = \gamma_i$ and $\widehat{A}_i = A_i^{-1}$. We can write $\tau_{i,A_i} z_i$ in a similar form for all $i \in F$ as well,

$$\begin{aligned} z_{i+1} &= B_i z_i + C_i A_i D_i z_i \\ &= \widehat{B}_i z_i + \widehat{C}_i (l_{i,i} \text{Id} + \widehat{A}_i)^{-1} \widehat{D}_i z_i \end{aligned}$$

where $\widehat{B}_i = B_i$, $\widehat{C}_i = C_i$, $\widehat{D}_i = D_i$, $l_{i,i} = 0$ and $\widehat{A}_i = A_i^{-1}$. Finally, since $p \notin F$ we can write $\tau_{p,A_p} z_p$ as

$$\begin{aligned} z_{p+1} &= B_p z_p + C_p J_{\gamma_p A_p} D_p z_p \\ &= B_p z_p + C_p (\gamma_p^{-1} \text{Id} + A_p)^{-1} (\gamma_p^{-1} D_p) z_p \\ &= \widehat{B}_p z_p + \widehat{C}_p (l_{p,p} \text{Id} + \widehat{A}_p)^{-1} \widehat{D}_p z_p \end{aligned}$$

where $\widehat{B}_p = B_p$, $\widehat{C}_p = C_p$, $\widehat{D}_p = \gamma_p^{-1} D_p$, $l_p = \gamma_p^{-1}$ and $\widehat{A}_p = A_p$. With these notations in place, we define $y_i \in \mathcal{H}$ as

$$y_i = (l_{i,i} \text{Id} + \widehat{A}_i)^{-1} \widehat{D}_i z_i, \quad \forall i \in \{1, \dots, n\}$$

which gives

$$z_{i+1} = \widehat{B}_i z_i + \widehat{C}_i y_i, \quad \forall i \in \{1, \dots, n\}$$

and for clarity we note that $\widehat{B}_i \in \mathbb{R}^{d_{i+1} \times d_i}$, $\widehat{C}_i \in \mathbb{R}^{d_{i+1} \times 1}$ and $\widehat{D}_i \in \mathbb{R}^{1 \times d_i}$ for all $i \in \{1, \dots, n\}$. Unrolling this iteration gives

$$z_i = \widehat{B}_{(i-1)\dots 1} z + \sum_{j=1}^{i-1} \widehat{B}_{(i-1)\dots(j+1)} \widehat{C}_j y_j, \quad \forall i \in \{1, \dots, n\}$$

where $\widehat{B}_{a\dots b} = \widehat{B}_a \widehat{B}_{a-1} \dots \widehat{B}_b$ and $\widehat{B}_{a\dots b} = I$ if $a < b$ where $I \in \mathbb{R}^{d_b \times d_b}$ is the identity matrix. The expression for y_i can then be rewritten as

$$l_{i,i} y_i + \widehat{A}_i y_i \ni \widehat{D}_i \widehat{B}_{(i-1)\dots 1} z + \sum_{j=1}^{i-1} \widehat{D}_i \widehat{B}_{(i-1)\dots(j+1)} \widehat{C}_j y_j, \quad \forall i \in \{1, \dots, n\}.$$

If we define $N_i = \widehat{D}_i \widehat{B}_{(i-1)\dots 1} \in \mathbb{R}^{1 \times d}$, $l_{i,j} = -\widehat{D}_i \widehat{B}_{(i-1)\dots(j+1)} \widehat{C}_j \in \mathbb{R}$ for $j < i$ and rearrange this expression we get

$$\sum_{j=1}^i l_{i,j} y_j + \widehat{A}_i y_i \ni N_i z, \quad \forall i \in \{1, \dots, n\}.$$

If we define $y = (y_1, \dots, y_n) \in \mathcal{H}^n$, the lower triangular matrix $L \in \mathbb{R}^{n \times n}$ with elements $L_{i,j} = l_{i,j}$ for all $j \leq i$ and the matrix $N \in \mathbb{R}^{n \times d}$ whose i th row is given by N_i , this can be written as $Ly + \Delta_{A,p} y \ni Nz$ or equivalently

$$y = (L + \Delta_{A,p})^{-1} Nz = (M + \Phi_{A,p})^{-1} Nz$$

where $M = L - \Gamma_p$. Note, this inverse $(M + \Phi_{A,p})^{-1}$ is single valued by the fact that $y_i = (l_{i,i} \text{Id} + \widehat{A}_i)^{-1} \widehat{D}_i z_i$, and $(l_{i,i} \text{Id} + \widehat{A}_i)^{-1}$ is single valued by construction. Finally, if we define the matrices $U = I - \widehat{B}_{n\dots 1} \in \mathbb{R}^{d \times d}$ and $V \in \mathbb{R}^{d \times n}$ where the i th column of V is given by \widehat{C}_i we can write the expression of z_{n+1} as

$$T_A z = z_{z+1} = z - Uz + Vy$$

which shows that (p, M, N, U, V) is a representation of $T_{(\cdot)}$. Since $M + \Gamma_p = L$ is lower triangular with diagonal elements $M_{i,i} = l_{i,i}$ and $l_{i,i} = 0$ if and only if $i \in F$ and otherwise $l_{i,i} > 0$, we can from Definition 4.2 we conclude that M is a p -kernel over \mathcal{A}_n^F . \square

Lemma 4.2. *Let (p, M, N, U, V) be a representation of a frugal splitting operator over \mathcal{A}_n^F with $p \notin F$, then*

- (i) $\ker \begin{bmatrix} N & -M \end{bmatrix} \supseteq \ker \begin{bmatrix} U & -V \end{bmatrix}$,
- (ii) $\text{ran } U \supseteq \text{ran } V$.

Proof. Let $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ be a frugal splitting operator over \mathcal{A}_n^F with representation (p, M, N, U, V) , i.e., for each $z \in \text{dom } T_{(\cdot)}$ there exists $y \in \mathcal{H}^n$ such that

$$\begin{cases} y = (M + \Phi_{(\cdot),p})^{-1}Nz, \\ T_{(\cdot)}z = z - Uz + Vy \end{cases} \quad \text{or equivalently} \quad \begin{cases} Nz - My \in \Phi_{(\cdot),p}y, \\ Uz - Vy = z - T_{(\cdot)}z. \end{cases} \quad (7)$$

We show the necessity of (i) by contradiction, i.e., we show that if $Uz - Vy = 0$ but $Nz - My \neq 0$ for some $z \in \mathbb{R}^d$ and $y \in \mathbb{R}^n$ then $T_{(\cdot)}$ is not a frugal splitting operator. In particular we will show that if $Uz - Vy = 0$ while $Nz - My \neq 0$, we can always construct some operator tuple $A \in \mathcal{A}_n^F$ such that $\text{fix } T_A \neq \emptyset$ but $\text{zer } \sum_{i=1}^n A_i = \emptyset$. This contradicts Definition 3.1(i) and hence $T_{(\cdot)}$ cannot be a frugal splitting operator.

Before we start constructing this counterexample, we note that if there exist points $z \in \mathbb{R}^d$ and $y \in \mathbb{R}^n$ such that $Uz - Vy = 0$ but $Nz - My \neq 0$ there exist points $z \in \mathcal{H}^d$ and $y \in \mathcal{H}^n$ such that $Uz - Vy = 0$ and $Nz - My \neq 0$. Now, let $z \in \mathcal{H}^d$ and $y = (y_1, \dots, y_n) \in \mathcal{H}^n$ be such points and define $(a_1, \dots, a_n) = Nz - My$. By assumption we have $(a_1, \dots, a_n) \neq 0$ and we can further split this into two cases, either $a_p \neq 0$ or otherwise $a_p = 0$ but then $a_i \neq 0$ for some $i \neq p$. We will show that both of these cases result in a contradiction to $T_{(\cdot)}$ being a frugal splitting operator.

First, assume $a_p \neq 0$ and define $A = (A_1, \dots, A_n) \in \mathcal{A}_n^F$ where the operators are defined as

$$A_i x = \begin{cases} a_p - \sum_{j \in \{1, \dots, n\} \setminus \{p\}} y_j & \text{if } i = p, \\ y_i & \text{otherwise} \end{cases}$$

for all $x \in \mathcal{H}$. The primal-dual operator $\Phi_{A,p}$ of this tuple evaluated at the y we specified above is

$$(\Phi_{A,p}y)_i = \begin{cases} a_p & \text{if } i = p, \\ \mathcal{H} & \text{otherwise} \end{cases}$$

where $(\Phi_{A,p}y)_i$ is the i th block of $\Phi_{A,p}y \in \mathcal{H}^n$. We can conclude that $Nz - My \in \Phi_{A,p}y$ and (7) then gives that $T_A z = z - Uz + Vy = z$ since $Uz - Vy = 0$ by assumption and we can conclude that $z \in \text{fix } T_A$. However, $\sum_{i=1}^n A_i x = a_p \neq 0$ for all $x \in \mathcal{H}$ and hence $\emptyset = \text{zer } \sum_{i=1}^n A_i$ which contradicts Definition 3.1(i).

Now, assume $a_p = 0$ but $a_i \neq 0$ for some $i \in \{1, \dots, n\} \setminus \{p\}$. Let $B = (B_1, \dots, B_n) \in \mathcal{A}_n^F$ where for all $x \in \mathcal{H}$ the operators are defined as

$$\begin{aligned} (\forall i \in F \text{ s.t. } a_i = 0) \quad & B_i x = y_i, \\ (\forall i \in F \text{ s.t. } a_i \neq 0) \quad & B_i x = \begin{cases} y_i + a_i(1 - \frac{\|a_i\|^2}{\langle a_i, x - y_p \rangle}) & \text{if } \langle a_i, x - y_p \rangle > 0 \\ \emptyset & \text{otherwise} \end{cases}, \\ (\forall i \in \{1, \dots, n\} \setminus F) \quad & B_i x = \begin{cases} \mathcal{H} & \text{if } x = y_p + a_i \\ \emptyset & \text{otherwise} \end{cases}. \end{aligned}$$

Evaluating the primal-dual operator $\Phi_{B,p}$ at y and remembering that $a_p = 0$ yields

$$(\Phi_{B,p}y)_i = \begin{cases} \mathcal{H} & \text{if } i = p, \\ \mathcal{H} & \text{if } i \in F \text{ and } a_i = 0, \\ a_i & \text{otherwise} \end{cases}$$

and hence $Nz - My \in \Phi_{B,p}y$ and $T_B z = z - Uz + Vy = z$ which implies $\emptyset \neq \text{zer } \sum_{B_i \in B} B_i$. Since $T_{(\cdot)}$ is a frugal splitting operator this implies that $\text{zer } \sum_{i=1}^n B_i \neq \emptyset$. Noting that $p \notin F$

we can see that $\text{dom } B_p = \{y_p + a_p\} = \{y_p\}$ since $a_p = 0$ by assumption and therefore must $\text{zer } \sum_{i=1}^n B_i = \{y_p\}$. However, let i be such that $a_i \neq 0$, it is then clear for the definition of B_i that $B_i y_p = \emptyset$ which contradicts $\sum_{i=1}^n B_i y_p = 0$. This concludes the necessity of (i).

For the necessity of (ii), let $y = (y_1, \dots, y_n) \in \mathcal{H}^n$ be arbitrary and define $C = (C_1, \dots, C_n) \in \mathcal{A}_n^F$ where the operators are defined as

$$C_i x = \begin{cases} x - y_p + y_i & \text{if } i \neq p, \\ x - \sum_{j=1}^n y_j & \text{if } i = p \end{cases}$$

for all $x \in \mathcal{H}$. From adding these operators, we see that $\{y_p\} = \text{zer } \sum_{i=1}^n C_i$ and $\{y\} = \text{zer } \Phi_{C,p}$. Since $T_{(\cdot)}$ is a frugal splitting operator we must then have $\emptyset \neq \text{fix } T_C$. Let $z \in \text{fix } T_C$, (7) then implies the existence $y' \in \mathcal{H}^d$ such that $Uz - Vy' = 0$ and $Nz - My' \in \Phi_{C,p} y'$. But, (i) implies that $Nz - My' = 0$ and hence must $y' \in \text{zer } \Phi_{C,p} = \{y\}$, i.e., $y' = y$ and $Uz = Vy$. Since the choice of y is arbitrary this implies that for all $y \in \mathcal{H}^n$ there exists $z \in \mathcal{H}^d$ such that $Uz = Vy$ which then implies the existence of $z \in \mathbb{R}^d$ for each $y \in \mathbb{R}^n$ such that $Uz = Vy$, i.e., $\text{ran } U \supseteq \text{ran } V$. This concludes the proof. \square

Lemma 4.3. *Let (p, M, N, U, V) be the representation of a generalized primal-dual resolvent where*

- (i) M is a p -kernel over \mathcal{A}_n^F ,
- (ii) $\ker \begin{bmatrix} N & -M \end{bmatrix} \supseteq \ker \begin{bmatrix} U & -V \end{bmatrix}$,
- (iii) $\text{ran } U \supseteq \text{ran } V$,

then the generalized primal-dual resolvent is a frugal splitting operator over \mathcal{A}_n^F .

Proof. Let $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ be a generalized primal-dual resolvent with representation (p, M, N, U, V) . Let $A \in \mathcal{A}_n^F$, for each $z \in \text{dom } T_A$ there exists $y \in \mathcal{H}^n$ such that

$$\begin{cases} y = (M + \Phi_{A,p})^{-1} Nz, \\ T_A z = z - Uz + Vy \end{cases} \quad \text{or equivalently} \quad \begin{cases} Nz - My \in \Phi_{A,p} y, \\ Uz - Vy = z - T_A z. \end{cases} \quad (8)$$

Item (i) implies that Definition 3.1(ii) holds for this operator. To show this we simply use the arguments of the proof of Lemma 4.1 in reverse.

If $z \in \text{fix } T_A$, then $Uz - Vy = 0$ and (ii) implies $0 = Nz - My \in \Phi_{A,p} y$ and hence $y \in \text{zer } \Phi_{A,p} \neq \emptyset$. This proves the right implication of Definition 3.1(i).

For the left implication of Definition 3.1(i), let $y^* \in \text{zer } \Phi_{A,p}$. Then (iii) implies that there exists $z \in \mathcal{H}^d$ such that $Uz - Vy^* = 0$ and (ii) then implies $Nz - My^* = 0$. Since $y^* \in \text{zer } \Phi_{A,p}$ we then have $0 = Nz - My^* \in \Phi_{A,p} y^*$ and z and y^* then satisfies (8) which proves that $z \in \text{fix } T_A \neq \emptyset$. \square

5 Remarks on the Frugal Splitting Operator Representation

The fact that the primal-dual operator appears directly in the representation and one only needs to consider simple range and structure constraints of a handful of matrices makes the representation easy to work with, both when analyzing and constructing splittings, see for instance Sections 6 to 8. However, the representation also reveals the relationship between fixed points of a splitting and solutions to (1) and in simple ways encodes properties like parallelizability and the resolvent step-sizes. In the following sections we discuss and illustrate these things as well as demonstrate an approach for deriving a representation of a frugal splitting operator.

5.1 Solution Map and Fixed Points

The following two propositions reveal the relationship between fixed points of the splitting operator and solutions to the monotone inclusion problem for any frugal splitting operator. They also

clarify why there is no need to assume the existence of a solution map in Definition 3.1 since there always exists a map that maps fixed points of a frugal splitting operator to solutions of the monotone inclusion problem (1). Furthermore, we see that this solution map is always evaluated within the evaluation of a frugal splitting operator itself.

Proposition 5.1. *Let $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ be a frugal splitting operator over \mathcal{A}_n^F with representation (p, M, N, U, V) where $p \notin F$. If $z \in \text{fix } T_A$ for $A \in \mathcal{A}_n^F$, then there exists $(y_1, \dots, y_n) \in \mathcal{H}^n$ such that*

$$(y_1, \dots, y_n) = (M + \Phi_{A,p})^{-1}Nz \in \text{zer } \Phi_{A,p} \quad \text{and} \quad y_p \in \text{zer } \sum_{i=1}^n A_i.$$

Proof. Let $A \in \mathcal{A}_n^F$ and $z \in \text{fix } T_A$, Definition 4.1 then gives the existence of $y = (y_1, \dots, y_n) \in \mathcal{H}^n$ such that

$$\begin{cases} y = (M + \Phi_{A,p})^{-1}Nz, \\ z = z - Uz + Vy \end{cases} \quad \text{or equivalently} \quad \begin{cases} Nz - My \in \Phi_{A,p}y, \\ Uz - Vy = 0 \end{cases}$$

Since $T_{(\cdot)}$ is a frugal splitting operator and Theorem 4.1(ii) necessarily hold for frugal splitting operators, $Uz - Vy = 0$ implies $Nz - My = 0$ and hence that $y \in \text{zer } \Phi_{A,p}$. That $y_p \in \text{zer } \sum_{i=1}^n A_i$ follows from the equivalence between the primal-dual problem (3) and the primal problem (1). \square

Proposition 5.2. *Let $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ be a frugal splitting operator over \mathcal{A}_n^F and let (p, SUP, SU, U, UP) be a representation of T_A that satisfies Corollary 4.1. With $A \in \mathcal{A}_n^F$, the set of fixed points satisfies*

$$\text{fix } T_A \supseteq P \text{zer } \Phi_{A,p}$$

where $P \text{zer } \Phi_{A,p} = \{Py \mid y \in \text{zer } \Phi_{A,p}\}$. Equality in the inclusion holds if and only if U has full rank.

Proof. Let $y \in \text{zer } \Phi_{A,p}$, then $0 = SUPy - SUPy \in \Phi_{A,p}y$ or equivalently $y = (SUP + \Phi_{A,p})^{-1}SUPy$. By letting $z = Py$ in the definition of the generalized primal-dual resolvent we conclude that

$$\begin{aligned} y &= (SUP + \Phi_{A,p})^{-1}SUPy, \\ T_A Py &= Py - U(Py - Py) = Py, \end{aligned}$$

which implies that $Py \in \text{fix } T_A$. Since $y \in \text{zer } \Phi_{A,p}$ was arbitrary we have $\text{fix } T_A \supseteq P \text{zer } \Phi_{A,p}$.

Assume U does not have full rank. Then, there exists $z \in \mathcal{H}^d$ such that $Uz = 0$ and $z \neq 0$. Since $\text{ran } P \subseteq (\ker U)^\perp$ by Corollary 4.1, there exists no $y' \in \mathcal{H}^n$ such that $z = Py'$ and hence $Py + z \notin P \text{zer } \Phi_{A,p}$. However, we have

$$\begin{aligned} y &= (SUP + \Phi_{A,p})^{-1}SUPy = (SUP + \Phi_{A,p})^{-1}SU(Py + z), \\ T_A(Py + z) &= Py + z - U(Py + z - Py) = Py + z, \end{aligned}$$

and hence $Py + z \in \text{fix } T_A$. The equality $\text{fix } T_A = P \text{zer } \Phi_{A,p}$ can therefore not hold if U does not have full rank.

Let $z \in \text{fix } T_A$ and assume U has full rank, i.e., U is invertible. It then exists $y \in \mathcal{H}^n$ such that

$$\begin{aligned} SUz - SUPy &\in \Phi_{A,p}y, \\ U(z - Py) &= z - T_A z = 0, \end{aligned}$$

but this implies that $z = Py$ and that $0 = SU(z - Py) \in \Phi_{A,p}y$ and hence that $y \in \text{zer } \Phi_{A,p}$ and $z \in P \text{zer } \Phi_{A,p}$. Since $z \in \text{fix } T_A$ was arbitrary we have $\text{fix } T_A \subseteq P \text{zer } \Phi_{A,p}$ and the opposite inclusion from before then gives equality. \square

5.2 Evaluation Order and Parallelizability

The order of the evaluations of the different operators is specified in a frugal splitting operator, Definition 3.1. However, the definition only states that it should be possible to compute the frugal splitting operator using this order, it does not exclude the possibility of computing the frugal splitting operator with some other evaluation order or with some of the evaluations being performed in parallel. Especially the ability of performing forward and/or backward evaluations in parallel is of particular interest since it can allow for distributed or multithreaded implementations. Fittingly, how one operator evaluation depends on previous evaluations is directly encoded in the kernel. It is therefore straightforward to both identify and construct parallelizable kernels, something we will use in Section 8 where we construct a new parallelizable frugal splitting operator with minimal lifting.

If we define $L = M + \Gamma_p$ for a representation (p, M, N, U, V) of a frugal splitting operator over \mathcal{A}_n^F we can write the inverse of the generalized primal-dual resolvent as

$$(M + \Phi_{A,p})^{-1} = (L + \Delta_{A,p})^{-1}.$$

Since M is a p -kernel, we know from Definition 4.2 that L is lower triangular matrix which means that $(y_1, \dots, y_n) = (L + \Delta_{A,p})^{-1}(z_1, \dots, z_n)$ can be computed with back-substitution,

$$y_i = (L_{i,i} \text{Id} + A_i^{-1})^{-1} (z_i - \sum_{j=1}^{i-1} L_{i,j} y_j) \quad (9)$$

for $i \neq p$ while if $i = p$ then A_i^{-1} is simply replaced by A_i . It is clear that the strict lower triangular part of L , $L_{i,j}$ for $j < i$, determines the dependency on the results of previous forward or backward evaluations⁴. Hence, if for $i \in \mathbb{N}$ and some $j < i$ the element $L_{i,j} = 0$, then the i th evaluation does not directly depend on the result of the j th. If the i th evaluation does not depend on any other evaluation that depends on the j th evaluation, then the i th and j th evaluation can be performed in parallel.

For example, if we take the 4-kernel over $\mathcal{A}_4^{\{2\}}$ as

$$M = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 1 \\ -1 & 0 & -1 & 1 \end{bmatrix} \quad \text{which yields} \quad L = M + \Gamma_5 = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 1 \end{bmatrix}$$

we see that, in any frugal splitting operator with this kernel, the second evaluation directly depends on the first; the third depends directly on only the second but depends indirectly on the first; the fourth is independent of the third, depends directly on the second and depends indirectly on the first. The second must therefore be performed after the first while the last two must be performed after the second but can be performed in parallel with each other.

5.3 Step-Sizes

It is possible to identify the resolvent step-sizes directly from the kernel. The inverse in (9) for $i \neq p$ and $L_{i,i} > 0$ can be written as

$$(L_{i,i} \text{Id} + A_i^{-1})^{-1} = L_{i,i}^{-1} (\text{Id} - J_{L_{i,i} A_i}).$$

From this we see that the diagonal elements of L act as step-sizes in the resolvents of the frugal splitting operator and hence so do the diagonal elements of M since $L = M + \Gamma_p$ and Γ_p is skew-symmetric. Similarly we can conclude that $L_{p,p}^{-1}$ or equivalently $M_{p,p}^{-1}$ is the step-size used in the

⁴The strictly lower triangular matrix $\tilde{L} \in \mathbb{R}^{n \times n}$ with elements $\tilde{L}_{i,j} = 1$ if $L_{i,j} \neq 0$ and $\tilde{L}_{i,j} = 0$ if $L_{i,j} = 0$ for $j < i$ is the transpose of adjacency matrix of the directed dependency graph of the operator evaluations, i.e., the graph of n nodes where there is an edge from the i th to j th node if the result of forward or backward evaluation of A_i is used in the argument of the forward or backward evaluation of A_j .

resolvent corresponding to the primal index. From Definition 4.2 we know that a forward evaluation is performed on the i th operator if and only if $L_{i,i} = M_{i,i} = 0$. Since they are zero, the diagonal elements of the kernel corresponding to forward evaluations are therefore not step-sizes. In fact, what would count as a step-size for a forward evaluation for a general frugal splitting operator is ill-defined. The same evaluation could be used in several different forward steps and we can therefore not point to a single scalar step-size. Examples of all these cases can be seen in the kernel presented in Section 5.4.

5.4 Example of a Representation

The process of finding representations of frugal splitting operators is quite straightforward. Roughly speaking it is as follows: select a primal index; apply Moreau's identity to all backward steps that do not correspond to the primal index; explicitly define the results of the forward evaluations and backward steps; rearrange and identify the primal-dual operator and the generalized primal-dual resolvent representation. We demonstrate the process on the three operator splitting of Davis–Yin [8] which encodes the zeros of $A_1 + A_2 + A_3$ where $A = (A_1, A_2, A_3) \in \mathcal{A}_3^{\{2\}}$ as the fixed points of the operator

$$T_A = J_{\gamma A_3} \circ (2J_{\gamma A_1} - \text{Id} - \gamma A_2 \circ J_{\gamma A_1}) + \text{Id} - J_{\gamma A_1}.$$

In order for fixed point iterations of T_A to always be convergent it is further required that A_2 is cocoercive and that the step-size γ is sufficiently small. However, we leave the convergence analysis of frugal splitting operators to Section 7.

To find a representation, we first choose primal index p in the representation (p, M, N, U, V) . Since it is required that $p \notin \{2\}$ we have the choice of either $p = 1$ or $p = 3$. We choose $p = 3$. Applying Moreau's identity to the resolvents of A_i for all $i \neq p$ (in this case only $J_{\gamma A_1}$) and defining the result of each forward and backward-step yields

$$\begin{aligned} y_1 &= (\text{Id} + \gamma^{-1} A_1^{-1})^{-1}(\gamma^{-1} z), \\ y_2 &= A_2(z - \gamma y_1), \\ y_3 &= (\text{Id} + \gamma A_3)^{-1}(2(z - \gamma y_1) - z - \gamma y_2), \\ T_A z &= y_3 + z - (z - \gamma y_1). \end{aligned}$$

Rearranging the first three lines such that we only have z on the left and y_1, y_2 and y_3 and unscaled operators on the right yields

$$\begin{aligned} z &\in A_1^{-1} y_1 + \gamma y_1, \\ z &\in A_2^{-1} y_2 + \gamma y_1, \\ \gamma^{-1} z &\in A_3 y_3 + 2y_1 + y_2 + \gamma^{-1} y_3, \\ T_A z &= \gamma y_1 + y_3. \end{aligned}$$

If we define $y = (y_1, y_2, y_3)$, we see that the first three lines can be written as

$$\begin{bmatrix} 1 \\ 1 \\ \gamma^{-1} \end{bmatrix} z \in \Delta_{A,3} y + \begin{bmatrix} \gamma & 0 & 0 \\ \gamma & 0 & 0 \\ 2 & 1 & \gamma^{-1} \end{bmatrix} y = \Phi_{A,3} y + \begin{bmatrix} \gamma & 0 & 1 \\ \gamma & 0 & 1 \\ 1 & 0 & \gamma^{-1} \end{bmatrix} y$$

and T_A can then be written as

$$\begin{aligned} y &= \left(\begin{bmatrix} \gamma & 0 & 1 \\ \gamma & 0 & 1 \\ 1 & 0 & \gamma^{-1} \end{bmatrix} + \Phi_{A,3} \right)^{-1} \begin{bmatrix} 1 \\ 1 \\ \gamma^{-1} \end{bmatrix} z, \\ T_A z &= z - [1] z + [\gamma \ 0 \ 1] y. \end{aligned}$$

From this we can easily identify the matrices M, N, U and V in the representation $(3, M, N, U, V)$ by comparing to Definition 4.1.

6 Minimal Lifting

Definition 6.1 (Lifting). *The lifting number, or lifting, of a frugal splitting operator $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ over \mathcal{A}_n^F is the number $d \in \mathbb{N}_+$.*

The lifting number represents how much memory, proportional to the problem variable in (1), is needed to store data between iterations in a fixed point iteration of the splitting operator. For instance, if we are trying to find a zero of operators from \mathbb{R}^N to $2^{(\mathbb{R}^N)}$ with a frugal splitting operator with lifting number 3, i.e., $T_{(\cdot)}: \mathbb{R}^{3N} \rightarrow \mathbb{R}^{3N}$, we need to be able to store a variable in \mathbb{R}^{3N} between iterations⁵. For the reason, we are interested in finding lower bounds for the lifting number and frugal splitting operators that attain these bounds.

Definition 6.2 (Minimal Lifting). *A frugal splitting operator $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ over \mathcal{A}_n^F has minimal lifting if $d \leq d'$ for all frugal splitting operators $T'_{(\cdot)}: \mathcal{H}^{d'} \rightarrow \mathcal{H}^{d'}$ over \mathcal{A}_n^F . Furthermore, we say that d is the minimal lifting over \mathcal{A}_n^F .*

The equivalent representation of a frugal splitting operator in Theorem 4.1 is very useful when it comes to examining this minimal lifting. In fact, a lower bound on the lifting number is directly given by Corollary 4.1.

Corollary 6.1. *Let (p, M, N, U, V) be a representation of a frugal splitting operator $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ over \mathcal{A}_n^F , then*

- (i) $d \geq \text{rank } U \geq \text{rank } N \geq \text{rank } M$,
- (ii) $d \geq \text{rank } U \geq \text{rank } V \geq \text{rank } M$.

Proof. Corollary 4.1 states that there exist matrices $S \in \mathbb{R}^{n \times d}$ and $P \in \mathbb{R}^{d \times n}$ such that $M = SUP$, $N = SU$ and $V = UP$. The results then follow directly from the fact that $U \in \mathbb{R}^{d \times d}$ and that a product of matrices cannot have greater rank than any of its factors. \square

Since the kernel rank bounds the lifting number, it is of great interest how small the rank of any valid kernel can be made, which leads us to the following definition.

Definition 6.3 (Minimal Kernel Rank). *A matrix $M \in \mathbb{R}^{n \times n}$ is a minimal p -kernel over \mathcal{A}_n^F if it is a p -kernel over \mathcal{A}_n^F and $\text{rank } M \leq \text{rank } M'$ for all p -kernels M' over \mathcal{A}_n^F . The minimal p -kernel rank over \mathcal{A}_n^F is $\text{rank } M$ where M is a minimal p -kernel M over \mathcal{A}_n^F .*

The existence of a minimal p -kernel over \mathcal{A}_n^F follows from the fact that $\text{rank } M \in \mathbb{N}$ for all real matrices and a p -kernel over \mathcal{A}_n^F exists for all $n \geq 2$, $F \in \{1, \dots, n\}$ and $p \in \{1, \dots, n\} \setminus F$. We can use the minimal p -kernel rank over \mathcal{A}_n^F and Corollary 6.1 to provide a lower bound on the minimal lifting number, but, this is not enough to establish that that lower bound actually can be attained. For that we need to show that it is always possible to construct a frugal splitting operator over \mathcal{A}_n^F from a p -kernel over \mathcal{A}_n^F .

Proposition 6.1. *Let $F \subset \{1, \dots, n\}$ and $p \in \{1, \dots, n\} \setminus F$. If M is a p -kernel over \mathcal{A}_n^F with $\text{rank } M = d$ and matrices $K \in \mathbb{R}^{n \times d}$ and $H \in \mathbb{R}^{d \times n}$ satisfy $\text{ran } K = (\ker M)^\perp$ and $\ker H = (\text{ran } M)^\perp$, then (p, M, MK, HMK, HM) is a representation of a frugal splitting operator over \mathcal{A}_n^F with lifting number d . Furthermore, such matrices K and H exist for all p -kernels over \mathcal{A}_n^F .*

Proof. We first show that (p, M, MK, HMK, HM) satisfies the conditions in Theorem 4.1.

Theorem 4.1(i) is directly given by the assumption on M . Now, let $x \in \mathbb{R}^d$ and $y \in \mathbb{R}^n$ be such that $HMKx = HMy$. Since $\ker H \cap \text{ran } M = \{0\}$, this implies $MKx = My$ which proves

⁵It is possible for the internal operations needed to evaluate the splitting operator itself to require additional memory. However, since this is highly problem and implementation dependent, we do not consider this engineering problem.

Theorem 4.1(ii). For Theorem 4.1(iii), let $y \in \mathbb{R}^n$ be arbitrary and let $y^\parallel \in \ker M$ and $y^\perp \in (\ker M)^\perp$ be such that $y = y^\parallel + y^\perp$. Since $\text{ran } K = (\ker M)^\perp$, there exists $x \in \mathbb{R}^d$ such that $y^\perp = Kx$ and hence $MKx = My^\perp = M(y^\parallel + y^\perp) = My$, multiplying both sides with H from the left finally gives Theorem 4.1(iii).

Finally, let M be an arbitrary p -kernel over \mathcal{A}_n^F . Let $\text{rank } M = d$, then the kernel M has d linearly independent columns and d linearly independent rows. Define $K \in \mathbb{R}^{n \times d}$ such that the columns of K are d linearly independent rows of M and define $H \in \mathbb{R}^{d \times n}$ such that the rows of H are d linearly independent columns of M , then $\text{ran } K = (\ker M)^\perp$ and $\ker H = (\text{ran } M)^\perp$. This concludes the proof. \square

With these results in hand, our main minimal lifting result can be stated and proved.

Theorem 6.1. *The minimal lifting number of a frugal splitting operator over \mathcal{A}_n^F is equal to the minimal p -kernel rank over \mathcal{A}_n^F for arbitrary $p \in \{1, \dots, n\} \setminus F$.*

Proof. Let $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ be a frugal splitting operator over \mathcal{A}_n^F with minimal lifting and let $p \in \{1, \dots, n\} \setminus F$. Theorem 4.1 states that $T_{(\cdot)}$ has a representation (p, M, N, U, V) and Corollary 6.1 then directly gives that the lifting d is greater or equal to the minimal p -kernel rank over \mathcal{A}_n^F . Proposition 6.1 proves that d is equal to the minimal p -kernel rank over \mathcal{A}_n^F since otherwise we could construct a frugal splitting operator with smaller lifting. Since the choice of $p \in \{1, \dots, n\} \setminus F$ was arbitrary and d is independent of p , the minimal p -kernel rank over \mathcal{A}_n^F is the same for all $p \in \{1, \dots, n\} \setminus F$. \square

This theorem reduces the problem of finding the minimal lifting over \mathcal{A}_n^F to finding the minimal p -kernel rank over \mathcal{A}_n^F . Furthermore, since Proposition 6.1 proves the existence results necessary for Theorem 6.1 by construction, it provides a clear way of constructing frugal splitting operators with lifting equal to the kernel rank. Hence, even the construction of frugal splitting operators with minimal lifting can be seen as a problem of finding p -kernels over \mathcal{A}_n^F with minimal rank. We will give an example of this in Section 8 and end this section by making Theorem 6.1 concrete by finding the minimal kernel rank over \mathcal{A}_n^F and the corresponding minimal lifting results.

Corollary 6.2. *Let $n \geq 2$ and $F \subset \{1, \dots, n\}$. The minimal lifting over \mathcal{A}_n^F is $n - |F|$ if $1 \in F$ or $n \in F$, otherwise it is $n - 1 - |F|$, where $|F|$ is the cardinality of F .*

Proof. Let M be a p -kernel over \mathcal{A}_n^F . It must have the structure

$$M = \left[\begin{array}{c|c|c} L_1 & \mathbf{1} & \mathbf{0} \\ * & l_p & -\mathbf{1} \\ * & * & L_2 \end{array} \right] \in \mathbb{R}^{n \times n}$$

where $L_1 \in \mathbb{R}^{(p-1) \times (p-1)}$ and $L_2 \in \mathbb{R}^{(n-p) \times (n-p)}$ are lower triangular matrices and $l_p > 0$ is a real number. The symbols $*$, $\mathbf{1}$ and $\mathbf{0}$ respectively denotes an arbitrary real matrix, a matrix of all ones, and a matrix of all zeros, all of appropriate sizes for their position. Since reordering columns and rows does not change the rank of a matrix there exist matrices

$$M_c = \left[\begin{array}{c|c|c} L_1 & \mathbf{0} & \mathbf{1} \\ * & L_2 & * \\ * & -\mathbf{1} & l_p \end{array} \right] \in \mathbb{R}^{n \times n} \quad \text{and} \quad M_r = \left[\begin{array}{c|c|c} l_p & * & -\mathbf{1} \\ \mathbf{1} & L_1 & \mathbf{0} \\ * & * & L_2 \end{array} \right] \in \mathbb{R}^{n \times n}.$$

such that $\text{rank } M = \text{rank } M_c = \text{rank } M_r$.

Consider the first $n - 1$ columns of M_c . Since L_1 and L_2 are lower triangular, we see that the number of linearly independent columns is greater or equal to the number of non-zero diagonal elements of L_1 and L_2 . Furthermore, since the diagonal elements of L_1 and L_2 are diagonal elements of M , $M_{i,i} = 0$ if and only if $i \in F$ and $p \notin F$, there are at least $n - 1 - |F|$ linearly independent columns

among the first $n - 1$ columns of M_c , hence, $\text{rank } M_c \geq n - 1 - |F|$. This bound can be attained by selecting the first column of M_c parallel to the last and putting zeros in all other positions that are allowed to be zero. If $1 \in F$ then $M_{1,1} = (L_1)_{1,1} = 0$ and the last column of M_c is not in the span of the first $n - 1$ columns and hence $\text{rank } M_c \geq n - |F|$. This bound is attained by putting zeros in all positions of M_c that are allowed to be zero. By considering rows of M_r instead of columns of M_c we can analogously conclude that $\text{rank } M_r \geq n - 1 - |F|$ and if $n \in F$ (and hence $M_{n,n} = (L_2)_{n-p,n-p} = 0$), then $\text{rank } M_r \geq n - |F|$. These bounds are also similarly attained.

Since $\text{rank } M = \text{rank } M_c = \text{rank } M_r$, we clearly have that $\text{rank } M \geq n - 1 - |F|$ and if $1 \in F$ or $n \in F$ then $\text{rank } M \geq n - |F|$. Choices of M that attain these bounds can be constructed by reordering the rows and columns of the choices of M_c and M_r that attain their respective bounds. The minimal kernel rank over \mathcal{A}_n^F is then clearly $n - |F|$ if $1 \in F$ or $n \in F$, otherwise it is $n - 1 - |F|$. The lifting result then follows from Theorem 6.1. \square

In the case of frugal resolvent splitting operators, i.e., $F = \emptyset$, this is the same lower bound as the one found by Malitsky and Tam [6]. When considering frugal splitting operators with forward evaluations, i.e., $F \neq \emptyset$, this corollary uncovers an interesting phenomenon where the minimal lifting depends on the evaluation order. A lower lifting number is possible as long as the first or last evaluation is not a forward evaluation. This makes it clear why the three operator splitting of Davis and Yin, see Section 5.4, achieves a lifting number of one while the corresponding methods of Vũ and Condat [9, 10] require a lifting of two. Davis and Yin perform the forward evaluation between two resolvent evaluations, while Vũ and Condat performs the forward evaluation first.

7 Convergence

We will now consider the convergence of fixed point iterations $z_{k+1} = T_A z_k$ for frugal splitting operators $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ over \mathcal{A}_n^F and $A \in \mathcal{A}_n^F$. From Corollary 4.1 we know that these iterations can, for each $p \in \{1, \dots, n\} \setminus F$, be written as

$$\begin{cases} y_k = (SUP + \Phi_{A,p})^{-1} S U z_k, \\ z_{k+1} = z_k - U(z_k - P y_k) \end{cases} \quad \text{or equivalently} \quad \begin{cases} S U(z_k - P y_k) \in \Phi_{A,p} y_k, \\ U(z_k - P y_k) = z_k - z_{k+1} \end{cases} \quad (10)$$

for some matrices $S \in \mathbb{R}^{n \times d}$, $P \in \mathbb{R}^{d \times n}$ and $U \in \mathbb{R}^{d \times d}$. In certain cases, this iteration can be analyzed with existing theory. For instance, if S is symmetric positive definite, $P = I$ and $U = \theta I$ for some $\theta \in (0, 1]$ this is an averaged fixed point iteration of a resolvent in the metric given by θS . However, for general frugal splitting operators, M is only rarely symmetric, especially when considering frugal splitting operators with minimal lifting. Only in the cases when $n = 1$ and $n = 2$ can a minimal p -kernel be symmetric⁶, see the proofs of Corollary 6.2. For this reason we will derive sufficient convergence conditions without any symmetry requirements on the kernel under the following assumption.

Assumption 7.1. Let $A = (A_1, \dots, A_n) \in \mathcal{A}_n^F$ be such that the following hold

- (i) $\text{zer } \Phi_{A,p} \neq \emptyset$ for all $p \in \{1, \dots, n\}$,
- (ii) A_i is β_i -cocoercive for all $i \in F$.

For this A , define two diagonal matrices $B \in \mathbb{R}^{n \times n}$ and $B^\dagger \in \mathbb{R}^{n \times n}$ where $B_{i,i} = \beta_i$ and $B_{i,i}^\dagger = \beta_i^{-1}$ for all $i \in F$. All other elements of the matrices are zero.

Cocoercivity of the forward evaluated operators is a standard setting for proving convergence of forward-backward like methods. However, there exist frugal splitting operators whose fixed point iterations converge under only Lipschitz assumptions on the operator used for forward evaluations.

⁶These symmetric minimal kernels correspond to the kernels used in the proximal point method and Douglas–Rachford splitting, respectively.

An example of this is the forward-reflected-backward splitting [11], but it should be noted that it does not have minimal lifting.

Since inverses of cocoercive operators are strongly monotone, primal-dual operators for tuples that satisfy Assumption 7.1 have the following strong-monotonicity-like property.

Lemma 7.1. *Let $F \subset \{1, \dots, n\}$, $p \in \{1, \dots, n\} \setminus F$ and let $A \in \mathcal{A}_n^F$ satisfy Assumption 7.1, then*

$$\langle u - y, x - y \rangle \geq \|x - y\|_B^2$$

for all $x, y \in \mathcal{H}^n$ and for all $u \in \Phi_{A,p}x$, $v \in \Phi_{A,p}y$.

Proof. With some slight abuse of notation for set valued operators, we first note that for all $x, y \in \mathcal{H}^n$ we have

$$\begin{aligned} & \langle \Phi_{A,p}x - \Phi_{A,p}y, x - y \rangle \\ &= \langle \Delta_{A,p}x - \Delta_{A,p}y, x - y \rangle + \langle \Gamma_p x - \Gamma_p y, x - y \rangle \\ &= \langle \Delta_{A,p}x - \Delta_{A,p}y, x - y \rangle \\ &= \langle A_p x_p - A_p y_p, x_p - y_p \rangle + \sum_{i \in \{1, \dots, n\} \setminus \{p\}} \langle A_i^{-1} x_i - A_i^{-1} y_i, x_i - y_i \rangle \end{aligned}$$

where the fact that Γ_p is skew-adjoint was used. For all $i \in \{1, \dots, n\} \setminus F$, the operator A_i is monotone and so is A_i^{-1} , while for all $i \in F$, A_i is β_i -cocoercive and hence A_i^{-1} is β_i -strongly monotone. Noting that $p \notin F$ by assumption and using these properties yields

$$\langle \Phi_{A,p}x - \Phi_{A,p}y, x - y \rangle \geq \sum_{f \in F} \beta_f \|x_f - y_f\|^2 = \|x - y\|_B^2. \quad \square$$

This property is used to derive our main convergence theorem.

Theorem 7.1. *Let $F \subset \{1, \dots, n\}$ and let $A \in \mathcal{A}_n^F$ satisfy Assumption 7.1. Let $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$ be a frugal splitting operator over \mathcal{A}_n^F and let (p, SUP, SU, U, UP) be a representation of $T_{(\cdot)}$ according to Corollary 4.1. Let the sequences $\{z_k\}_{k \in \mathbb{N}}$ and $\{y_k\}_{k \in \mathbb{N}}$ be generated by (10) for this representation and some $z_0 \in \mathcal{H}^d$.*

Consider the following conditions on a symmetric matrix $Q \in \mathbb{R}^{d \times d}$,

- (a) $(I - I_F)(P^T Q - S)U = 0$,
- (b) $Q \succ 0$ and $W \succ 0$,

where $W = QU + (QU)^T - U^T QU - \frac{1}{2}(P^T QU - SU)^T B^\dagger (P^T QU - SU)$, $I \in \mathbb{R}^{n \times n}$ is the identity matrix, and $I_F \in \mathbb{R}^{n \times n}$ is the diagonal matrix with $(I_F)_{i,i} = 1$ if $i \in F$, otherwise $(I_F)_{i,i} = 0$. If (a) holds, then

$$\|z_{k+1} - Py\|_Q^2 \leq \|z_k - Py\|_Q^2 - \|z_k - Py_k\|_W^2$$

for all $k \in \mathbb{N}$ and all $y \in \text{zer } \Phi_{A,p}$. If (b) also holds, then

- (i) $z_k - Py_k \rightarrow 0$,
- (ii) $\Phi_{A,p}y_k \ni SU(z_k - Py_k) \rightarrow 0$,
- (iii) $y_k \rightarrow y^*$,
- (iv) $z_k \rightarrow Py^* \in \text{fix } T_A$,

for some $y^* \in \text{zer } \Phi_{A,p}$ as $k \rightarrow \infty$.

Proof. Let $Q \in \mathbb{R}^{d \times d}$ be a symmetric matrix and let $k \in \mathbb{N}$. For arbitrary $y \in \text{zer } \Phi_{A,p}$, we have

$$\begin{aligned} \|z_{k+1} - Py\|_Q^2 &= \|z_k - Py - U(z_k - Py_k)\|_Q^2 \\ &= \|z_k - Py\|_Q^2 + \|z_k - Py_k\|_{U^T QU}^2 - 2\langle QU(z_k - Py_k), z_k - Py \rangle. \end{aligned}$$

Insert $Py_k - Py_k$ on the right hand side of the inner product to get

$$\begin{aligned}\|z_{k+1} - Py\|_Q^2 &= \|z_k - Py\|_Q^2 + \|z_k - Py_k\|_{U^TQU}^2 - 2\langle QU(z_k - Py_k), z_k - Py_k \rangle \\ &\quad - 2\langle QU(z_k - Py_k), Py_k - Py \rangle \\ &= \|z_k - Py\|_Q^2 - \|z_k - Py_k\|_{QU+(QU)^T-U^TQU}^2 \\ &\quad - 2\langle P^TQU(z_k - Py_k), y_k - y \rangle.\end{aligned}$$

From (10) we know that $SU(z_k - Py_k) \in \Phi_{A,p}y_k$ and Lemma 7.1 then gives that

$$0 \leq \langle SU(z_k - Py_k), y_k - y \rangle - \|y_k - y\|_B^2.$$

Add two times this inequality to the previous equality to get

$$\begin{aligned}\|z_{k+1} - Py\|_Q^2 &\leq \|z_k - Py\|_Q^2 - \|z_k - Py_k\|_{QU+(QU)^T-U^TQU}^2 - 2\|y_k - y\|_B^2 \\ &\quad - 2\langle P^TQU(z_k - Py_k), y_k - y \rangle + 2\langle SU(z_k - Py_k), y_k - y \rangle \\ &= \|z_k - Py\|_Q^2 - \|z_k - Py_k\|_{QU+(QU)^T-U^TQU}^2 - 2\|y_k - y\|_B^2 \\ &\quad - 2\langle (P^TQU - SU)(z_k - Py_k), y_k - y \rangle.\end{aligned}$$

Assume that condition (a) is satisfied, then $P^TQU - SU = I_F(P^TQU - SU)$ and since $I_F = (\frac{1}{2}B^\dagger)^{1/2}(2B)^{1/2}$, we have

$$\begin{aligned}\|z_{k+1} - Py\|_Q^2 &\leq \|z_k - Py\|_Q^2 - \|z_k - Py_k\|_{QU+(QU)^T-U^TQU}^2 - 2\|y_k - y\|_B^2 \\ &\quad - 2\langle (2^{-1}B^\dagger)^{1/2}(P^TQU - SU)(z_k - Py_k), (2B)^{1/2}(y_k - y) \rangle.\end{aligned}$$

Using Young's inequality then finally results in

$$\begin{aligned}\|z_{k+1} - Py\|_Q^2 &\leq \|z_k - Py\|_Q^2 - \|z_k - Py_k\|_{QU+(QU)^T-U^TQU}^2 \\ &\quad + \|(2^{-1}B^\dagger)^{1/2}(P^TQU - SU)(z_k - Py_k)\|^2 \\ &= \|z_k - Py\|_Q^2 - \|z_k - Py_k\|_W^2\end{aligned}\tag{11}$$

where $W = QU + (QU)^T - U^TQU - \frac{1}{2}(P^TQU - SU)^T B^\dagger (P^TQU - SU)$, which proves the first statement.

From here on we assume that (b) also holds, which implies that both $\|\cdot\|_Q$ and $\|\cdot\|_W$ indeed are norms on \mathcal{H}^d . The k used in (11) is arbitrary and the inequality holds for all $k \in \mathbb{N}$. Adding the inequalities for $k = 0, 1, \dots$ implies

$$\|z_k - Py_k\|_W^2 \rightarrow 0,$$

which implies that (i) holds. Statement (ii) follows directly from (i) and (10). Inequality (11) further implies

$$\|z_{k+1} - Py\|_Q^2 \leq \|z_k - Py\|_Q^2$$

for all $k \in \mathbb{N}$ and all $y \in \text{zer } \Phi_{A,p}$ which in turn implies the boundedness of $\{z_k\}_{k \in \mathbb{N}}$ and the convergence of $\{\|z_k - Py\|_Q^2\}_{k \in \mathbb{N}}$ for all $y \in \text{zer } \Phi_{A,p}$. These two facts along with (i) and (ii) will be used to prove (iii). Statement (iv) then follows directly from the weak continuity of P and (iii).

For the proof of (iii), we will first show that $\{y_k\}_{k \in \mathbb{N}}$ is bounded and hence has weak sequential cluster points. Then we will show that these cluster points are in $\text{zer } \Phi_{A,p}$ and lastly we will show that there is at most one cluster point. The convergence $y_k \rightharpoonup y^*$ for some $y^* \in \text{zer } \Phi_{A,p}$ then follows from [27, Lemma 2.46].

Since SUP is a p -kernel over \mathcal{A}_n^F , the operator $(SUP + \Phi_{A,p})^{-1} \circ SU$ can be evaluated using a finite number of vector additions, scalar multiplications, resolvents and evaluations of the cocoercive operators A_i with $i \in F$. All of these operations are Lipschitz continuous and hence must $(SUP +$

$\Phi_{A,p})^{-1} \circ SU$ be Lipschitz continuous, let us say with constant L . Furthermore, with $y \in \text{zer } \Phi_{A,p}$ we have $y = (SUP + \Phi_{A,p})^{-1} SUPy$ and

$$\|y_k - y\| = \|(SUP + \Phi_{A,p})^{-1} SUz_k - (SUP + \Phi_{A,p})^{-1} SUPy\| \leq L\|z_k - Py\|.$$

As noted above, the sequence $\{z_k\}_{k \in \mathbb{N}}$ is bounded and hence is $\{y_k\}_{k \in \mathbb{N}}$ also bounded and has convergent subsequences [27, Lemma 2.45]. From (ii), the maximal monotonicity of $\Phi_{A,p}$, see Section 2.1, and the weak-strong continuity of maximally monotone operators [27, Proposition 20.37] we can conclude that any weak sequential cluster point of $\{y_k\}_{k \in \mathbb{N}}$ lies in $\text{zer } \Phi_{A,p}$.

What remains to be shown is that $\{y_k\}_{k \in \mathbb{N}}$ possesses at most one weak sequential cluster point. Let $\{y_{k_i}\}_{i \in \mathbb{N}}$ and $\{y_{k_j}\}_{j \in \mathbb{N}}$ be sub-sequences such that $y_{k_i} \rightharpoonup a \in \text{zer } \Phi_{A,p}$ and $y_{k_j} \rightharpoonup b \in \text{zer } \Phi_{A,p}$. Since P as an operator on \mathcal{H}^n is linear and hence weakly continuous, (i) implies that

$$z_{k_i} \rightharpoonup Pa \quad \text{and} \quad z_{k_j} \rightharpoonup Pb.$$

Furthermore, $\{\|z_k - Pa\|_Q^2\}_{k \in \mathbb{N}}$ and $\{\|z_k - Pb\|_Q^2\}_{k \in \mathbb{N}}$ both converge and hence

$$\langle z_k, Pa - Pb \rangle_Q = \frac{1}{2}\|z_k - Pa\|_Q^2 - \frac{1}{2}\|z_k - Pb\|_Q^2 - \frac{1}{2}\|Pa\|_Q^2 + \frac{1}{2}\|Pb\|_Q^2 \rightarrow \nu$$

for some $\nu \in \mathbb{R}$. In particular, it means that

$$\langle z_{k_i}, Pa - Pb \rangle_Q \rightarrow \langle Pa, Pa - Pb \rangle_Q = \nu = \langle Pb, Pa - Pb \rangle_Q \leftarrow \langle z_{k_j}, Pa - Pb \rangle_Q.$$

This implies

$$0 = \langle Pa - Pb, Pa - Pb \rangle_Q = \|Pa - Pb\|_Q^2$$

and hence $Pa = Pb$. Since $y \in \text{zer } \Phi_{A,p}$ implies $y = (SUP + \Phi_{A,p})^{-1} SUPy$ we have

$$a = (SUP + \Phi_{A,p})^{-1} SUPa = (SUP + \Phi_{A,p})^{-1} SUPb = b.$$

This proves that there is at most one weak sequential cluster point. Hence, $\{y_k\}_{k \in \mathbb{N}}$ converge weakly to some $y^* \in \text{zer } \Phi_{A,p}$, i.e., (iii) holds. \square

A few remarks on this theorem are in order. The sequence generated by a fixed point iteration of a frugal splitting operator that satisfies Theorem 7.1 is Fejér monotone, see [27, Definition 5.1] with respect to $P \text{zer } \Phi_{A,p}$ in the Hilbert space given by $\langle \cdot, \cdot \rangle_Q$. This is true even for methods without minimal lifting such as momentum methods, see Section 7.1. We will in particular in show that forward-backward splitting with Nesterov-like momentum satisfies Theorem 7.1 and hence is Fejér monotone. Furthermore, we see that condition (a) becomes stricter when the set F on which we make forward evaluations becomes smaller. This puts stronger restrictions on the structure of the frugal splitting operator. For instance, if we assume that S , U and P are square and invertible and $F = \emptyset$, condition (a) states that $Q = (P^T)^{-1}S$ and hence $(P^T)^{-1}S$ must be symmetric. In other cases, we have more freedom in choosing $Q \succ 0$ such that $W \succ 0$ and (b) of Theorem 7.1 is satisfied. It should also be noted that, although $W \succ 0$ is a quadratic expression in Q , it can be transformed to an equivalent positive definite condition that is linear in Q by using a Schur complement.

From condition (b) we conclude that $W \succ 0$ implies that $U + U^T \succ 0$ and hence must U have full rank d . This is convenient since, given a frugal splitting operator with representation (p, M, N, U, V) where U is invertible, it is easy to find a representation of the form used in Theorem 7.1, i.e., (p, SUP, SU, U, UP) . In fact, this factorization can be expressed only in terms of the matrices N , U and V ,

$$S = NU^{-1} \quad \text{and} \quad P = U^{-1}V,$$

which implies that $M = NU^{-1}V$ must hold for such a frugal splitting operator.

There are frugal splitting operators where U is rank deficient but their convergence are of no interest. This is because they are either guaranteed to not converge in general or they can be reduced to a frugal splitting operator whose representation have a full rank U without loosing any information.

To see this, let (p, SUP, SU, U, UP) be a representation satisfying Corollary 4.1 of a frugal splitting operator $T_{(\cdot)}: \mathcal{H}^d \rightarrow \mathcal{H}^d$. Let $\Pi \in \mathbb{R}^{d \times d}$ be the projection matrix onto $\text{ran } U^T$ —the projection is here in the standard Euclidean \mathbb{R}^d space. The projection matrix on the kernel of U is then $\bar{\Pi} = I - \Pi$. Consider the fixed point iteration $z_{k+1} = T_A z_k$ for some $A \in \mathcal{A}_n^F$ and $z_0 \in \mathcal{H}^d$. If we define $z_0^\parallel = \Pi z_0$ and $z_0^\perp = \bar{\Pi} z_0$ this fixed point iteration can be written as $z_k = z_k^\parallel + z_k^\perp$ where

$$\begin{aligned} y_k &= (SUP + \Phi_{A,p})^{-1} SU z_k^\parallel, \\ z_{k+1}^\parallel &= z_k^\parallel - \Pi U(z_k^\parallel - P y_k), \\ z_{k+1}^\perp &= z_k^\perp - \bar{\Pi} U(z_k^\parallel - P y_k), \end{aligned}$$

for all $k \in \mathbb{N}$. Without going into details, if $\text{ran } U = \text{ran } U^T$ then $z_k^\perp = z_0^\perp$ for all $k \in \mathbb{N}$ and only the $\{z_k^\parallel\}_{k \in \mathbb{N}}$ sequence is of interest. Furthermore, the $\{z_k^\parallel\}_{k \in \mathbb{N}}$ sequence can be recovered from a fixed point iteration of a frugal splitting operator with lifting equal to the rank of U . If instead $\text{ran } U \neq \text{ran } U^T$, it is always possible to find an operator tuple $A \in \mathcal{A}_n^F$ and initial point $z_0 \in \mathcal{H}^d$ such that $z_k^\parallel = z_0$, $z_k^\perp = -kc$ and $z_k = z_0 - kc$ for all $k \in \mathbb{N}$ and some $c \in \mathcal{H}^d \setminus \{0\}$. Hence, $\{z_k\}_{k \in \mathbb{N}}$ will always diverge for this choice of A and z_0 .

7.1 Applications of Theorem 7.1

Theorem 7.1 recovers many well-known convergence results, for instance forward-backward splitting [3, 4], Douglas–Rachford splitting [2], the Chambolle–Pock method [7] and the minimal lifting methods of Ryu [5] and Malitsky–Tam [6]. We will not present all these results here and settle for presenting the results for the three operator splitting of Davis and Yin [8]. We will also present convergence conditions for the fixed point iteration of the forward-backward operator with Nesterov-like momentum [28, 29]. To save on space, we will not derive any of the representations and simply state the primal index p and the matrices U, S and P of a representation (p, SUP, SU, U, UP) , see Corollary 4.1. Similarly, for the convergence results we just state the matrices Q and W of Theorem 7.1. Detailed derivations of all the listed examples and more can be found in the supplement.

Three Operator Splitting of Davis–Yin The three operator splitting of Davis and Yin has already been presented in Section 5.4 but we restate it here,

$$x_{k+1} = x_k - J_{\gamma A_1} x_k + J_{\gamma A_3} \circ (2 J_{\gamma A_1} - \text{Id} - \gamma A_2 \circ J_{\gamma A_1}) x_k$$

where $x_0 \in \mathcal{H}$, $\gamma > 0$ and $A \in \mathcal{A}_3^{\{2\}}$. The representation derived in that section can be factored into a representation of the form $(3, SUP, SU, U, UP)$ where

$$U = [1], \quad S = \begin{bmatrix} 1 \\ 1 \\ \gamma^{-1} \end{bmatrix} \quad \text{and} \quad P = [\gamma \quad 0 \quad 1].$$

To prove convergence, we choose $Q = [\gamma^{-1}]$ in Theorem 7.1 and, assuming A satisfies Assumption 7.1, this results in $0 = (I - I_F)(P^T Q - S)U$ and $W = [\gamma^{-1} - \frac{1}{2\beta_2}]$. If $\gamma < 2\beta_2$, then $x_k \rightarrow P y^*$ for $y^* \in \text{zer } \Phi_{A,3}$ and

$$J_{\gamma A_3} \circ (2 J_{\gamma A_1} - \text{Id} - \gamma A_2 \circ J_{\gamma A_1}) x_k = J_{\gamma A_1} x_k + x_{k+1} - x_k \rightarrow x^* \in \text{zer } A_1 + A_2 + A_3$$

and hence also $J_{\gamma A_1} x_k \rightarrow x^*$. This is the same convergence condition as the one presented in [8].

Forward-Backward with Nesterov Momentum A fixed point iteration of a forward-backward operator with Nesterov-like momentum [28] can be written as

$$x_{k+1} = J_{\gamma A_2}(x_k + \theta(x_k - x_{k-1})) - \gamma A_1(x_k + \theta(x_k - x_{k-1}))$$

for $\lambda > 0$, $\theta \in \mathbb{R}$, $x_0, x_{-1} \in \mathcal{H}$ and $A \in \mathcal{A}_2^{\{1\}}$. In the proximal-gradient setting, this is also the same update that is used in the FISTA method [29]. Nesterov momentum gained popularity due to it achieving optimal convergence rates in the smooth optimization setting. However, these faster convergence rates require a momentum parameter θ that varies between iterations, something the fixed point iterations considered in this paper will not allow.

We wish to remove the dependency on previous iterations and hence introduce an extra iterate as

$$\begin{aligned} x_{k+1} &= J_{\gamma A_2}(x_k + \theta y_k - \gamma A_1(x_k + \theta y_k)), \\ y_{k+1} &= x_{k+1} - x_k \end{aligned}$$

where $x_0, y_0 \in \mathcal{H}$. This is a fixed point iteration of the frugal splitting operator given by the representation $(2, SUP, SU, U, UP)$ where

$$U = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}, \quad S = \begin{bmatrix} 1 - \theta & \theta \\ \gamma^{-1}(1 - \theta) & \gamma^{-1}\theta \end{bmatrix} \quad \text{and} \quad P = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}.$$

The lifting number is two and is hence not minimal even though the kernel SUP has rank one and is minimal. In fact, the kernel is the same as for the ordinary forward-backward splitting. Theorem 7.1 with

$$Q = \gamma^{-1} \begin{bmatrix} 1 - \theta & \theta \\ \theta & a \end{bmatrix} \quad \text{and} \quad W = \gamma^{-1} \begin{bmatrix} 1 - \theta - \hat{\gamma} - a & \theta(1 - \hat{\gamma}) \\ \theta(1 - \hat{\gamma}) & a - \theta^2 \hat{\gamma} \end{bmatrix}$$

where $a > 0$ and $\hat{\gamma} = \frac{\gamma}{2\beta_1} > 0$ yields the convergence of $x_k \rightarrow x^* \in \text{zer } A_1 + A_2$ and $y_k \rightarrow 0$ if A satisfies Assumption 7.1 and both $Q \succ 0$ and $W \succ 0$, i.e.,

$$0 < 1 - \theta - \frac{\theta^2}{a}, \quad 0 < a - \theta^2 \hat{\gamma} \quad \text{and} \quad 0 < 1 - \theta - \hat{\gamma} - a - \frac{\theta^2(1 - \hat{\gamma})^2}{a - \theta^2 \hat{\gamma}}.$$

If we restrict these results to $\theta > 0$, these conditions hold with $a = \theta$ if

$$0 < \frac{\gamma}{2\beta_1} < \frac{1 - 3\theta}{(1 + \theta)(1 - 2\theta)} \quad \text{and} \quad 0 < \theta < \frac{1}{3}.$$

It is easily verified that this condition also implies the existence of an $a > 0$ that makes $Q \succ 0$ and $W \succ 0$ even in the case when $\theta = 0$. In fact, it reduces to the well known result $0 < \gamma < 2\beta_1$ for ordinary forward-backward splitting. As mentioned before, if these conditions hold then Theorem 7.1 gives Fejér monotonicity of $\{(x_k, y_k)\}_{k \in \mathbb{N}}$ w.r.t. $P \text{zer } \Phi_{A,p} = \text{zer}(A_1 + A_2) \times \{0\}$ in the norm $\|\cdot\|_Q$.

8 A New Frugal Splitting Operator With Minimal Lifting

We will now derive a new frugal splitting operator with minimal lifting. The approach we will take is the same as the one outlined in Proposition 6.1, i.e., we will select a primal index p , a kernel M and matrices H and K and form a representation as (p, M, MK, HMK, HM) . As long as the matrices satisfy Proposition 6.1, the resulting generalized primal-dual resolvent is a frugal splitting operator with lifting equal to the rank of the kernel. Therefore, if we choose a kernel with minimal rank, the resulting frugal splitting operator will have minimal lifting.

As noted in Section 6, the minimal lifting number of a frugal splitting operator over \mathcal{A}_n^F depends on F . In particular, from Corollary 6.2 we see that it is not only a question regarding whether F is empty or not; the minimal lifting number depends on the actual order of the forward and backward evaluations. For this reason, we choose to construct a frugal splitting operator over \mathcal{A}_n^F where F is such that $1 \notin F$ and $n \notin F$. Corollary 6.2 then guarantees that the minimal lifting in this setting is $n - 1 - |F|$ instead of potentially being $n - |F|$. To further simplify the setup we assume that all the single valued evaluations come one after another as $F = \{n - f, \dots, n - 1\}$ where $f = |F|$ is the number of single valued operators.

Theorem 8.1. Let $n, f \in \mathbb{N}$ be such that $n \geq 2$ and $f \leq n-2$. Let $A = (A_1, \dots, A_n) \in \mathcal{A}_n^F$ where $F = \{n-f, \dots, n-1\}$ if $f > 0$ and $F = \emptyset$ if $f = 0$. Let $z_{i,0} \in \mathcal{H}$ for all $i \in \{1, \dots, n-1-f\}$ and

$$\begin{aligned} x_{1,k} &= J_{\lambda A_1} z_{1,k}, \\ x_{i,k} &= J_{\theta^{-1} \lambda A_i} (x_{1,k} + \theta^{-1} z_{i,k}) && \text{for all } i \in \{2, \dots, n-1-f\}, \\ x_{i,k} &= \lambda A_i x_{1,k} && \text{for all } i \in \{n-f, \dots, n-1\}, \\ \bar{x}_k &= \sum_{j=n-f}^{n-1} x_{j,k} + \sum_{j=2}^{n-1-f} z_{i,k} + \theta(x_{1,k} - x_{j,k}), \\ x_{n,k} &= J_{\lambda A_n} (2x_{1,k} - z_{1,k} - \bar{x}_k), \\ z_{i,k+1} &= z_{i,k} - \theta(x_{i,k} - x_{n,k}) && \text{for all } i \in \{1, \dots, n-1-f\} \end{aligned}$$

for all $k \in \mathbb{N}$ where $\gamma > 0$ and $\theta > 0$. If A satisfies Assumption 7.1 and

$$\frac{\lambda}{2} \sum_{i=n-f}^{n-1} \beta_i^{-1} < 2 - \theta(n-1-f)$$

then $x_{n,k} \rightarrow x^* \in \text{zer} \sum_{i=1}^n A_i$.

Proof. Let $T_{(\cdot)}: \mathcal{H}^{n-1-f} \rightarrow \mathcal{H}^{n-1-f}$ be the generalized primal-dual resolvent with representation (n, M, MK, HMK, HM) with the matrices

$$\begin{aligned} M &= \left[\begin{array}{c|c|c|c} 1 & & & 1 \\ \hline \mathbf{1} & \frac{1}{\theta} I & & \mathbf{1} \\ \hline \mathbf{1} & & & \mathbf{1} \\ \hline 1 & & & 1 \end{array} \right] \in \mathbb{R}^{n \times n}, \quad K = \left[\begin{array}{c|c} \frac{1}{2} & \\ \hline & I \\ \hline & \\ \hline \frac{1}{2} & \end{array} \right] \in \mathbb{R}^{n \times (n-1-f)} \\ \text{and } H &= \theta \left[\begin{array}{c|c|c|c} \frac{1}{2+f} & & & \frac{1}{2+f} \\ \hline & I & & \\ \hline & & \frac{1}{2+f} \mathbf{1}^T & \\ \hline & & & \frac{1}{2+f} \end{array} \right] \in \mathbb{R}^{(n-1-f) \times n}. \end{aligned}$$

where $\theta > 0$, $I \in \mathbb{R}^{(n-2-f) \times (n-2-f)}$ is the identity matrix, $\mathbf{1}$ are column vectors of ones with appropriate sizes and empty blocks denote zero matrices. Note, the third rows and columns vanish completely when $f = 0$ while the second rows and columns vanish when $f = n-2$. The matrix M is an n -kernel over \mathcal{A}_n^F with rank $n-1-f$, which is minimal, see Corollary 6.2. Furthermore, these matrices satisfy $\text{ran } K = \text{ran } M^T = (\ker M)^\perp$ and $\ker H = (\text{ran } H^T)^\perp = (\text{ran } M)^\perp$ and hence is $T_{(\cdot)}$ a frugal splitting operator by Proposition 6.1. For later use we note that $T_{(\cdot)}$ also can be represented as (n, SUP, SU, U, UP) with $U = HMK$, $S = MK(HMK)^{-1}$, $P = (HMK)^{-1}HM$.

We first consider the case when $\lambda = 1$, the general case when $\lambda > 0$ will be proved below. In this case, if we define $z_k = (z_{1,k}, \dots, z_{n-1-f,k})$, the update of z_{k+1} can be written as

$$z_{k+1} = T_A z_k.$$

It can also be verified that $(y_{1,k}, \dots, y_{n,k}) = (M + \Phi_{A,n})^{-1} MK z_k$ satisfies

$$\begin{aligned} y_{1,k} &= z_{1,k} - x_{1,k}, \\ y_{i,k} &= z_{1,k} + \theta(x_{1,k} - x_{i,k}) && \text{for all } i \in \{2, \dots, n-1-f\}, \\ y_{i,k} &= x_{i,k} && \text{for all } i \in \{n-f, \dots, n\}, \end{aligned}$$

for all $k \in \mathbb{N}$. Furthermore, if we choose

$$Q = \theta^{-1} \left[\begin{array}{c|c} 1 & \\ \hline & I \end{array} \right] \in \mathbb{R}^{(n-1-f) \times (n-1-f)}.$$

then $I_F(P^T Q - S)U = (P^T Q - S)U$ where I_F is defined in Theorem 7.1 and hence does condition (a) of Theorem 7.1 hold. Condition (b) of Theorem 7.1 with this Q reads as

$$Q \succ 0 \quad \text{and} \quad W = \left[\frac{2 - \theta(n-1-f) - \frac{1}{2} \sum_{i=n-f}^{n-1} \beta_i^{-1}}{\frac{1}{\theta} I} \right] \succ 0$$

which holds if and only if $\theta > 0$ and $\frac{1}{2} \sum_{i=n-f}^{n-1} \beta_i^{-1} < 2 - \theta(n-1-f)$. In this case, Theorem 7.1 gives the convergence of

$$(y_{1,k}, \dots, y_{n,k}) \rightarrow (y_1^*, \dots, y_n^*) \in \text{zer } \Phi_{A,n}$$

and $x_{n,k} = y_{n,k} \rightarrow y_n^* \in \text{zer } \sum_{i=1}^n A_i$.

Finally, consider the general case with arbitrary $\lambda > 0$. We then see that the update of the theorem can be written as

$$z_{k+1} = T_{\lambda A} z_k$$

for all $k \in \mathbb{N}$ where $\lambda A = (\lambda A_1, \dots, \lambda A_n)$. Since A_i is β_i -cocoercive for all $i \in F$, λA_i will be $\beta_i \lambda^{-1}$ -cocoercive for all $i \in F$ and hence will λA satisfy Assumption 7.1. Applying Theorem 7.1 to this fixed point iteration with the scaled operator tuple then gives the convergence of

$$(y_{1,k}, \dots, y_{n,k}) \rightarrow (y_1^*, \dots, y_n^*) \in \text{zer } \Phi_{\lambda A, n}$$

and $x_{n,k} = y_{n,k} \rightarrow y_n^* \in \text{zer } \sum_{i=1}^n \lambda A_i$ if $\theta > 0$ and $\frac{\lambda}{2} \sum_{i=n-f}^{n-1} \beta_i^{-1} < 2 - \theta(n-1-f)$. This proves the theorem. \square

Note that the relaxation factor θ also occurs as a step-size scaling on some of the resolvents. This was necessary in order to ensure convergence. We also see that, in order to find a step-size that ensures convergence, $\theta < \frac{2}{n-1-f}$ is needed, i.e., the relaxation parameter must decrease with the number of operators. However, this is counteracted by the fact that the step-size of most of the resolvents are scaled with θ^{-1} and hence will also increase with the number of operators.

There are no step-size restrictions when no forward evaluations are used, $f = 0$. Furthermore, we see that the step-size bound only depends on the sum of the inverse cocoercivity constants. This is natural since all forward steps are evaluated at the same point and the results are simply added together. The update can therefore equivalently be seen as evaluating $\hat{A} = \sum_{i=n-f}^{n-1} A_i$ instead of each operator individually and \hat{A} is a $\sum_{i=n-f}^{n-1} \beta_i^{-1}$ -cocoercive operator.

8.1 Relation to Other Methods With Minimal Lifting

When $n = 3$, $f = 1$ and $\theta = 1$, this method reduces to the three operator splitting of Davis and Yin, see Section 5.4. When $n = 3$ and $f = 0$, the method is closely related to the three operator resolvent splitting operator of Ryu [5]. That method uses a frugal splitting operator that is calculated as $(\hat{z}_1, \hat{z}_2) = T_{(A_1, A_2, A_3)}(z_1, z_2)$ where

$$\begin{aligned} x_1 &= J_{\lambda A_1}(z_1), \\ x_2 &= J_{\lambda A_2}(z_2 + x_1), \\ x_3 &= J_{\lambda A_3}(-z_1 - z_2 + x_1 + x_2), \\ \hat{z}_1 &= z_1 + \theta(x_3 - x_1), \\ \hat{z}_2 &= z_2 + \theta(x_3 - x_2) \end{aligned}$$

and we see that, if we consider the unrelaxed case $\theta = 1$, then this update is the same as our update in Theorem 8.1. For other choices of θ , our proposed method differs in that the step-size in the computation of x_2 is scaled with θ^{-1} . As noted by Malitsky and Tam [6], a straightforward

extension of Ryu’s method to four operators fails to converge for all $\theta > 0$ in some cases and we found this step-size scaling to be the key that allowed us to establish convergence for $n > 3$.

In the case when $f = 0$ and $n > 2$ is arbitrary, the only other method with minimal lifting known to us is the method of Malitsky and Tam [6]. It uses a frugal splitting operator where $(\hat{z}_1, \dots, \hat{z}_{n-1}) = T_{(A_1, \dots, A_n)}(z_1, \dots, z_{n-1})$ is calculated as

$$\begin{aligned} x_1 &= J_{\gamma A_1}(z_1), \\ x_i &= J_{\gamma A_i}(z_i - z_{i-1} + x_{i-1}) && \text{for all } i \in \{2, \dots, n-1\}, \\ x_n &= J_{\gamma A_n}(-z_{n-1} + x_1 + x_{n-1}), \\ \hat{z}_i &= z_i + \theta(x_{i+1} - x_i) && \text{for all } i \in \{1, \dots, n-1\}. \end{aligned}$$

One feature of our method is that it can calculate $x_{i,k}$ in parallel for all $i \in \{2, \dots, n-1\}$, which cannot be done in the Malitsky–Tam method since each resolvent depends on the previous one. However, the Malitsky–Tam method converges for all $\theta \in (0, 1)$ while our method requires that $\theta \in (0, \frac{2}{n-1})$, i.e., our method requires a smaller relaxation parameter for larger n . As mentioned before though, our method actually increases the step-sizes for the resolvents of A_2, \dots, A_{n-1} with n , which might offset the decreasing relaxation. Generalized primal-dual resolvent representations of both Ryu’s and Malitsky–Tam’s minimal lifting methods can be found in the supplement. There we also re-establish their convergence conditions using Theorem 7.1.

9 Conclusion

We have presented an explicit parameterization of all frugal splitting operators. The parameterization is in terms of what we call generalized primal-dual resolvents, and we have provided necessary and sufficient conditions for a generalized primal-dual resolvent being a frugal splitting operator. This allows for a unified analysis and both minimal lifting and convergence results that are applicable to all frugal splitting operators. The minimal lifting results of Ryu and Malitsky–Tam were expanded beyond resolvent splitting operators to general frugal splitting operators with forward evaluations and we showed that the lifting number depends on the order of forward and backward evaluations. We further presented a new convergent frugal splitting operator with minimal lifting that allows for most of the forward and/or backward steps to be computed in parallel. This is in contrast to the method of Malitsky–Tam that requires all resolvents to be computed sequentially. In the triple-backward case the method is the same as the minimal lifting method of Ryu if neither method uses relaxation. The slight difference in how relaxation is introduced is crucial to extend Ryu’s method to an arbitrary number of operators. In the double-backward-single-forward case the method reduces to three operator splitting by Davis and Yin.

References

- [1] R. T. Rockafellar, “Monotone Operators and the Proximal Point Algorithm,” *SIAM Journal on Control and Optimization*, vol. 14, no. 5, pp. 877–898, 1976.
- [2] P. L. Lions and B. Mercier, “Splitting Algorithms for the Sum of Two Nonlinear Operators,” *SIAM Journal on Numerical Analysis*, vol. 16, no. 6, pp. 964–979, 1979.
- [3] A. A. Goldstein, “Convex Programming in Hilbert Space,” *Bulletin of the American Mathematical Society*, vol. 70, no. 5, pp. 709–711, 1964.
- [4] E. S. Levitin and B. T. Polyak, “Constrained Minimization Methods,” *USSR Computational mathematics and mathematical physics*, vol. 6, no. 5, pp. 1–50, 1966.
- [5] E. K. Ryu, “Uniqueness of DRS as the 2 Operator Resolvent-Splitting and Impossibility of 3 Operator Resolvent-Splitting,” *Mathematical Programming*, vol. 182, no. 1, pp. 233–273, 2020.

- [6] Y. Malitsky and M. K. Tam, “Resolvent Splitting for Sums of Monotone Operators with Minimal Lifting,” *arXiv:2108.02897v1 [math]*, 2021. [Online]. Available: <http://arxiv.org/abs/2108.02897v1>
- [7] A. Chambolle and T. Pock, “A First-Order Primal-Dual Algorithm for Convex Problems with Applications to Imaging,” *Journal of Mathematical Imaging and Vision*, vol. 40, no. 1, pp. 120–145, 2011.
- [8] D. Davis and W. Yin, “A Three-Operator Splitting Scheme and its Optimization Applications,” *Set-Valued and Variational Analysis*, vol. 25, no. 4, pp. 829–858, 2017.
- [9] B. C. Vũ, “A splitting algorithm for dual monotone inclusions involving cocoercive operators,” *Advances in Computational Mathematics*, vol. 38, no. 3, pp. 667–681, 2013.
- [10] L. Condat, “A Primal–Dual Splitting Method for Convex Optimization Involving Lipschitzian, Proxiable and Linear Composite Terms,” *Journal of Optimization Theory and Applications*, vol. 158, no. 2, pp. 460–479, 2013.
- [11] Y. Malitsky and M. K. Tam, “A Forward-Backward Splitting Method for Monotone Inclusions Without Cocoercivity,” *SIAM Journal on Optimization*, vol. 30, no. 2, pp. 1451–1472, 2020.
- [12] F. Alvarez and H. Attouch, “An Inertial Proximal Method for Maximal Monotone Operators via Discretization of a Nonlinear Oscillator with Damping,” *Set-Valued Analysis*, vol. 9, no. 1, pp. 3–11, 2001.
- [13] A. Moudafi and M. Oliny, “Convergence of a Splitting Inertial Proximal Method for Monotone Operators,” *Journal of Computational and Applied Mathematics*, vol. 155, no. 2, pp. 447–454, 2003.
- [14] E. K. Ryu and B. C. Vũ, “Finding the Forward-Douglas–Rachford-Forward Method,” *Journal of Optimization Theory and Applications*, vol. 184, no. 3, pp. 858–876, 2020.
- [15] R. I. Boş, E. R. Csetnek, and C. Hendrich, “Inertial Douglas–Rachford splitting for monotone inclusion problems,” *Applied Mathematics and Computation*, vol. 256, pp. 472–487, 2015.
- [16] H. Raguét, J. Fadili, and G. Peyré, “A Generalized Forward-Backward Splitting,” *SIAM Journal on Imaging Sciences*, vol. 6, no. 3, pp. 1199–1226, 2013.
- [17] P. Latafat and P. Patrinos, “Asymmetric Forward–Backward–Adjoint Splitting for Solving Monotone Inclusions Involving Three Operators,” *Computational Optimization and Applications*, vol. 68, no. 1, pp. 57–93, 2017.
- [18] P. Giselsson, “Nonlinear Forward-Backward Splitting with Projection Correction,” *SIAM Journal on Optimization*, pp. 2199–2226, 2021.
- [19] M. Morin, S. Banert, and P. Giselsson, “Nonlinear Forward-Backward Splitting with Momentum Correction,” *arXiv:2112.00481v2 [math]*, 2022. [Online]. Available: <http://arxiv.org/abs/2112.00481v2>
- [20] M. N. Bui and P. L. Combettes, “Warped Proximal Iterations for Monotone Inclusions,” *Journal of Mathematical Analysis and Applications*, vol. 491, no. 1, p. 124315, 2020.
- [21] L. Lessard, B. Recht, and A. Packard, “Analysis and Design of Optimization Algorithms via Integral Quadratic Constraints,” *SIAM Journal on Optimization*, vol. 26, no. 1, pp. 57–95, 2016.
- [22] Y. Drori and M. Teboulle, “Performance of First-Order Methods for Smooth Convex Minimization: A Novel Approach,” *Mathematical Programming*, vol. 145, no. 1, pp. 451–482, 2014.
- [23] A. B. Taylor, J. M. Hendrickx, and F. Glineur, “Exact Worst-Case Performance of First-Order Methods for Composite Convex Optimization,” *SIAM Journal on Optimization*, vol. 27, no. 3, pp. 1283–1313, 2017.

- [24] ———, “Smooth Strongly Convex Interpolation and Exact Worst-Case Performance of First-Order Methods,” *Mathematical Programming*, vol. 161, no. 1, pp. 307–345, 2017.
- [25] E. K. Ryu, A. B. Taylor, C. Bergeling, and P. Giselsson, “Operator Splitting Performance Estimation: Tight Contraction Factors and Optimal Parameter Selection,” *SIAM Journal on Optimization*, vol. 30, no. 3, pp. 2251–2271, 2020.
- [26] P. Tseng, “A Modified Forward-Backward Splitting Method for Maximal Monotone Mappings,” *SIAM Journal on Control and Optimization*, vol. 38, no. 2, pp. 431–446, 2000.
- [27] H. H. Bauschke and P. L. Combettes, *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*, 2nd ed., ser. CMS Books in Mathematics. Springer International Publishing, 2017.
- [28] Y. Nesterov, “A Method For Solving A Convex Programming Problem With Rate of Convergence $O(1/k^2)$,” *Soviet Math. Doklady*, vol. v.269, no. No.3, pp. 543–547, 1983.
- [29] A. Beck and M. Teboulle, “A Fast Iterative Shrinkage-Thresholding Algorithm for Linear Inverse Problems,” *SIAM Journal on Imaging Sciences*, vol. 2, no. 1, pp. 183–202, 2009.
- [30] B. T. Polyak, “Some Methods of Speeding up the Convergence of Iteration Methods,” *USSR Computational Mathematics and Mathematical Physics*, vol. 4, no. 5, pp. 1–17, 1964.
- [31] L. M. Briceño-Arias, “Resolvent splitting with minimal lifting for composite monotone inclusions,” *arXiv:2111.09757v2 [math]*, 2021. [Online]. Available: <http://arxiv.org/abs/2111.09757v2>
- [32] P. L. Combettes and J. Eckstein, “Asynchronous Block-Iterative Primal-Dual Decomposition Methods for Monotone Inclusions,” *Mathematical Programming*, vol. 168, no. 1, pp. 645–672, 2018.

Supplement

A Derivation of Representations and Convergence Conditions

In this section we will verify the representations and convergence conditions for the frugal splitting operators stated in the paper. For posterity we will also derive representations and convergence conditions for a number of frugal splitting operators not previously mentioned.

We will keep a consistent notation in each of the examples presented. For a representation of a frugal splitting operator $T_{(\cdot)}$ over \mathcal{A}_n^F , the goal is to find matrices M, N, U, V such that

$$\begin{aligned} y &= (M + \Phi_{(\cdot),p})^{-1}Nz, \\ T_{(\cdot)}z &= z - Uz - Vy \end{aligned}$$

for some $p \in \{1, \dots, n\}$ and that Theorem 4.1 is satisfied. Such a representation (p, M, N, U, V) can always be factorized in terms of matrices S and P such that

$$M = SUP, \quad N = SU \quad \text{and} \quad V = UP,$$

and where $\text{ran } P \subseteq (\ker U)^\perp$ and $\ker S \supseteq (\text{ran } U)^\perp$, see Corollary 4.1. The convergence conditions of Theorem 7.1 for a fixed point iterations of T_A for some $A \in \mathcal{A}_n^F$ that satisfies Assumption 7.1 is stated in terms of this factorization and can be written as

$$\begin{aligned} Q &\succ 0, \\ (I - I_F)(P^T Q - S)U &= 0, \\ W &= QU + (QU)^T - U^T QU - \frac{1}{2}U^T(P^T Q - S)^T B^\dagger(P^T Q - S)U \succ 0 \end{aligned}$$

where Q is a symmetric matrix that needs to be found for each frugal splitting operator. When constructing new frugal splitting operators, we find it more convenient to work with a factorization of the representation (p, M, N, U, V) in terms of matrices H and K such that

$$N = MK, \quad V = HM \quad \text{and} \quad U = HMK,$$

where $\text{ran } K = (\ker M)^\perp$ and $\ker H = (\text{ran } M)^\perp$, see Proposition 6.1. This allows us to first design a kernel and then easily find N, U and V that results in a first order splitting. Furthermore, some of the frugal splitting operators will be presented without step-size. A variable step-size can be added to these method simply by scaling the operator tuple in the same way as for our new frugal splitting operator in Theorem 8.1.

A.1 Forward-Backward

The forward-backward operator [3, 4] is

$$T_{(A_1, A_2)} = J_{\gamma A_2} \circ (\text{Id} - \gamma A_1)$$

where $\gamma > 0$ and $(A_1, A_2) \in \mathcal{A}_2^{\{1\}}$. To derive a representation we choose primal index $p = 2$. The next step would be to apply the Moreau identity to all backward steps with index $i \neq p$ but since there are no such backward steps we can directly define the results of each forward and backward evaluation

$$\begin{aligned} y_1 &= A_1 z, \\ y_2 &= (\text{Id} + \gamma A_2)^{-1}(z - \gamma y_1), \\ T_A z &= y_2. \end{aligned}$$

e rewrite the first two lines such that the input is in the left and the result of all forward and backward steps are on the right

$$\begin{aligned} z &\in A_1^{-1}y_1, \\ \gamma^{-1}z &\in A_2y_2 + y_1 + \gamma^{-1}y_2, \\ T_A z &= y_2. \end{aligned}$$

If we define $y = (y_1, y_2) \in \mathcal{H}^2$ the first two lines can be written as

$$\begin{bmatrix} 1 \\ \gamma^{-1} \end{bmatrix} z \in \underbrace{\begin{bmatrix} A_1^{-1} & 0 \\ 0 & A_2 \end{bmatrix}}_{\Delta_{A,2}} y + \begin{bmatrix} 0 & 0 \\ 1 & \gamma^{-1} \end{bmatrix} y = \underbrace{\begin{bmatrix} A_1^{-1} & -1 \\ 1 & A_2 \end{bmatrix}}_{\Phi_{A,2}} y + \begin{bmatrix} 0 & 1 \\ 0 & \gamma^{-1} \end{bmatrix} y$$

which yields

$$\begin{aligned} y &= \left(\begin{bmatrix} 0 & 1 \\ 0 & \gamma^{-1} \end{bmatrix} + \Phi_{A,2} \right)^{-1} \begin{bmatrix} 1 \\ \gamma^{-1} \end{bmatrix} z, \\ T_A z &= z - [1] z + [0 \ 1] y, \end{aligned}$$

and the matrices M, N, U, V in the representation $(2, M, N, U, V)$ are then easily identified by comparing to Definition 4.1 which yields

$$M = \begin{bmatrix} 0 & 1 \\ 0 & \gamma^{-1} \end{bmatrix}, \quad N = \begin{bmatrix} 1 \\ \gamma^{-1} \end{bmatrix}, \quad V = [0 \ 1] \quad \text{and} \quad U = [1].$$

It is seen directly that

$$S = \begin{bmatrix} 1 \\ \gamma^{-1} \end{bmatrix} \quad \text{and} \quad P = [0 \ 1]$$

provides a factorization of this representation as $(2, SUP, SU, U, UP)$. For the convergence conditions, choosing $Q = [\gamma^{-1}]$ where I is the 2×2 identity matrix it is obvious that $Q \succ 0$ and we further have

$$\begin{aligned} (I - I_F)(P^T Q - S)U &= (I - I_F) \left(\gamma^{-1} \begin{bmatrix} 0 \\ 1 \end{bmatrix} - \begin{bmatrix} 1 \\ \gamma^{-1} \end{bmatrix} \right) \\ &= (I - I_F) \begin{bmatrix} -1 \\ 0 \end{bmatrix} \\ &= 0 \end{aligned}$$

and

$$\begin{aligned} W &= QU + (QU)^T - U^T QU - \frac{1}{2} U^T (P^T Q - S)^T B^\dagger (P^T Q - S)U \\ &= [\gamma^{-1}] + [\gamma^{-1}] - [\gamma^{-1}] - \frac{1}{2} [-1 \ 0] \begin{bmatrix} \beta_1^{-1} & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} -1 \\ 0 \end{bmatrix} \\ &= [\gamma^{-1} - \frac{1}{2\beta_1}]. \end{aligned}$$

The final condition for convergence is then $W \succ 0$ which yield the well known result $\gamma < 2\beta_1$.

A.2 Douglas–Rachford

The Douglas–Rachford splitting operator [2] is

$$T_{(A_1, A_2)} = \frac{1}{2} \text{Id} + \frac{1}{2} (2 J_{\gamma A_2} - \text{Id}) \circ (2 J_{\gamma A_1} - \text{Id})$$

where $(A_1, A_2) \in \mathcal{A}_2$ and $\gamma > 0$. Consider the evaluation $\hat{z} = T_{(A_1, A_2)} z$ for some $z \in \mathcal{H}$. This can be written as

$$\begin{aligned} x_1 &= J_{\gamma A_1} z, \\ y_2 &= J_{\gamma A_2} (2x_1 - z), \\ \hat{z} &= \frac{1}{2}z + \frac{1}{2}(2y_2 - (2x_1 - z)). \end{aligned}$$

We choose the primal index to $p = 2$ and apply the Moreau identity to the first resolvent,

$$\begin{aligned} y_1 &= J_{\gamma^{-1}A_1^{-1}}(\gamma^{-1}z), \\ x_1 &= z - \gamma y_1, \\ y_2 &= J_{\gamma A_2}(2x_1 - z), \\ \hat{z} &= \frac{1}{2}z + \frac{1}{2}(2y_2 - (2x_1 - z)). \end{aligned}$$

Eliminating x_1 gives

$$\begin{aligned} y_1 &= J_{\gamma^{-1}A_1^{-1}}(\gamma^{-1}z), \\ y_2 &= J_{\gamma A_2}(z - 2\gamma y_1), \\ \hat{z} &= \gamma y_1 + y_2. \end{aligned}$$

Using the definition of the resolvent yields

$$\begin{aligned} z &\in \gamma y_1 + A_1^{-1}y_1, \\ \gamma^{-1}z &\in 2y_1 + \gamma^{-1}y_2 + A_2y_2, \\ \hat{z} &= z - z + \gamma y_1 + y_2. \end{aligned}$$

Regrouping such that $\Phi_{(A_1, A_2), 2}$ can be identified finally gives

$$\begin{aligned} z &\in [\gamma y_1 + y_2] + [A_1^{-1}y_1 - y_2], \\ \gamma^{-1}z &\in [y_1 + \gamma^{-1}y_2] + [A_2y_2 + y_1], \\ \hat{z} &= z - [z] + [\gamma y_1 + y_2] \end{aligned}$$

and we can identify

$$M = \begin{bmatrix} \gamma & 1 \\ 1 & \gamma^{-1} \end{bmatrix}, \quad N = \begin{bmatrix} 1 \\ \gamma^{-1} \end{bmatrix}, \quad V = [\gamma \quad 1] \quad \text{and} \quad U = [1].$$

The representation can be factored as $(2, SUP, SU, U, UP)$ where

$$S = \begin{bmatrix} 1 \\ \gamma^{-1} \end{bmatrix} \quad \text{and} \quad P = [\gamma \quad 1].$$

For the convergence conditions, choosing $Q = [\gamma^{-1}]$ yield $Q \succ 0$ and

$$(I - I_F)(P^T Q - S) = (P^T Q - S) = \gamma^{-1} \begin{bmatrix} \gamma \\ 1 \end{bmatrix} - \begin{bmatrix} 1 \\ \gamma^{-1} \end{bmatrix} = 0$$

and

$$\begin{aligned} W &= QU + (QU)^T - U^T QU - \frac{1}{2}U^T (P^T Q - S)^T B^\dagger (P^T Q - S)U \\ &= Q + Q - Q = Q \end{aligned}$$

and hence is $W \succ 0$, i.e., fixed point iterations of the Douglas–Rachford splitting operator always converge.

A.3 Davis–Yin Three Operator Splitting

To find a representation of three operator splitting of Davis–Yin [8],

$$T_{(A_1, A_2, A_3)} = J_{\gamma A_3} \circ (2J_{\gamma A_1} - \text{Id} - \gamma A_2 \circ J_{\gamma A_1}) + \text{Id} - J_{\gamma A_1}.$$

where $\gamma > 0$ and $(A_1, A_2, A_3) \in \mathcal{A}_3^{\{2\}}$ we choose $p = 3$. Applying Moreau's identity to the resolvents of A_i for all $i \neq p$ (in this case only $J_{\gamma A_1}$) and defining the result of each forward and backward-step yields

$$\begin{aligned} y_1 &= (\text{Id} + \gamma^{-1} A_1^{-1})^{-1}(\gamma^{-1} z), \\ y_2 &= A_2(z - \gamma y_1), \\ y_3 &= (\text{Id} + \gamma A_3)^{-1}(2(z - \gamma y_1) - z - \gamma y_2), \\ T_A z &= y_3 + z - (z - \gamma y_1). \end{aligned}$$

Rearranging the first three lines such that we only have z on the left and y_1, y_2 and y_3 and unscaled operators on the right yields

$$\begin{aligned} z &\in A_1^{-1} y_1 + \gamma y_1, \\ z &\in A_2^{-1} y_2 + \gamma y_1, \\ \gamma^{-1} z &\in A_3 y_3 + 2y_1 + y_2 + \gamma^{-1} y_3, \\ T_A z &= \gamma y_1 + y_3. \end{aligned}$$

If we define $y = (y_1, y_2, y_3)$ we see that the first three lines can be written as

$$\begin{bmatrix} 1 \\ 1 \\ \gamma^{-1} \end{bmatrix} z \in \Delta_{A,3} y + \begin{bmatrix} \gamma & 0 & 0 \\ \gamma & 0 & 0 \\ 2 & 1 & \gamma^{-1} \end{bmatrix} y = \Phi_{A,3} y + \begin{bmatrix} \gamma & 0 & 1 \\ \gamma & 0 & 1 \\ 1 & 0 & \gamma^{-1} \end{bmatrix} y$$

and T_A can then be written as

$$\begin{aligned} y &= \left(\begin{bmatrix} \gamma & 0 & 1 \\ \gamma & 0 & 1 \\ 1 & 0 & \gamma^{-1} \end{bmatrix} + \Phi_{A,3} \right)^{-1} \begin{bmatrix} 1 \\ 1 \\ \gamma^{-1} \end{bmatrix} z, \\ T_A z &= z - [1] z + [\gamma \quad 0 \quad 1] y. \end{aligned}$$

From this we can easily identify the matrices M, N, U and V in the representation $(3, M, N, U, V)$ by comparing to Definition 4.1.

$$M = \begin{bmatrix} \gamma & 0 & 1 \\ \gamma & 0 & 1 \\ 1 & 0 & \gamma^{-1} \end{bmatrix}, \quad N = \begin{bmatrix} 1 \\ 1 \\ \gamma^{-1} \end{bmatrix}, \quad V = [\gamma \quad 0 \quad 1]. \quad \text{and} \quad U = [1]$$

This can be factored as $(3, SUP, SU, U, UP)$ where

$$S = \begin{bmatrix} 1 \\ 1 \\ \gamma^{-1} \end{bmatrix} \quad \text{and} \quad P = [\gamma \quad 0 \quad 1].$$

The convergence conditions are satisfied by $Q = [\gamma^{-1}]$. We see that $Q \succ 0$ and

$$\begin{aligned} (I - I_F)(P^T Q - S)U &= (I - I_F)(P^T Q - S) \\ &= (I - I_F) \left(\gamma^{-1} \begin{bmatrix} \gamma \\ 0 \\ 1 \end{bmatrix} - \begin{bmatrix} 1 \\ 1 \\ \gamma^{-1} \end{bmatrix} \right) \\ &= 0 \end{aligned}$$

and

$$\begin{aligned}
W &= QU + (QU)^T - U^T QU - \frac{1}{2} U^T (P^T Q - S)^T B^\dagger (P^T Q - S) U \\
&= Q - \frac{1}{2} (P^T Q - S)^T B^\dagger (P^T Q - S) \\
&= [\gamma^{-1}] - \frac{1}{2} \begin{bmatrix} 0 & -1 & 0 \end{bmatrix} B^\dagger \begin{bmatrix} 0 \\ -1 \\ 0 \end{bmatrix} \\
&= [\gamma^{-1} - \frac{1}{2\beta_2}]
\end{aligned}$$

and $W \succ 0$ if $\gamma < 2\beta_2$.

A.4 Forward-Backward with Polyak Momentum

Forward-Backward with Polyak momentum [30] can be written as a fixed point iteration of the frugal splitting operator

$$T_{(A_1, A_2)}(z_1, z_2) = \begin{pmatrix} \hat{z}_1 \\ \hat{z}_2 \end{pmatrix} = \begin{pmatrix} J_{\gamma A_2}(z_1 - \gamma A_1 z_1 + \theta z_2) \\ \hat{z}_1 - z_1 \end{pmatrix}$$

where $\gamma > 0$, $\theta \in \mathbb{R}$ and $(A_1, A_2) \in \mathcal{A}_2^{\{1\}}$. Note, there are other frugal splitting operators whose fixed point iteration are equivalent to forward-backward with Polyak momentum. We claim that $(2, SUP, SU, U, UP)$ where

$$U = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}, \quad S = \begin{bmatrix} 1 & 0 \\ \gamma^{-1}(1 - \theta) & \gamma^{-1}\theta \end{bmatrix} \quad \text{and} \quad P = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

is a representation of this frugal splitting operator. To show this we first note that

$$\begin{aligned}
V = UP &= \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}, \quad M = SUP = \begin{bmatrix} 0 & 1 \\ 0 & \gamma^{-1} \end{bmatrix} \quad \text{and} \\
N = SU &= \begin{bmatrix} 1 & 0 \\ \gamma^{-1}(1 - \theta) & \gamma^{-1}\theta \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ \gamma^{-1} & \gamma^{-1}\theta \end{bmatrix}
\end{aligned}$$

and

$$\begin{aligned}
Nz - My &\in \Phi_{A, 2}y, \\
\hat{z} &= z - Uz + Vy
\end{aligned}$$

can then be written as

$$\begin{aligned}
z_1 - y_2 &\in A_1^{-1}y_1 - y_2, \\
\gamma^{-1}(z_1 + \theta z_2) - \gamma^{-1}y_2 &\in A_2 y_2 + y_1, \\
\hat{z}_1 &= z_1 - z_1 + y_2, \\
\hat{z}_2 &= z_2 - z_1 - z_2 + y_2
\end{aligned}$$

which after some rearranging gives

$$\begin{aligned}
y_1 &\in A_1 z_1, \\
z_1 - \gamma y_1 + \theta z_2 &\in (\text{Id} + \gamma A_2) y_2, \\
\hat{z}_1 &= y_2, \\
\hat{z}_2 &= y_2 - z_1.
\end{aligned}$$

Rewriting the second line as a resolvent and combining the first and second gives

$$\begin{aligned} y_2 &= J_{\gamma A_2}(z_1 - \gamma A_1 z_1 + \theta z_2), \\ \hat{z}_1 &= y_2, \\ \hat{z}_2 &= y_2 - z_1 \end{aligned}$$

which is exactly the frugal splitting operator above. For the convergence, if we choose

$$Q = \gamma^{-1} \begin{bmatrix} 1 - \theta & \theta \\ \theta & a \end{bmatrix}$$

where $a > 0$ then

$$\begin{aligned} (I - I_F)(P^T Q - S)U &= (I - I_F)\gamma^{-1} \left(\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1 - \theta & \theta \\ \theta & a \end{bmatrix} - \begin{bmatrix} \gamma & 0 \\ 1 - \theta & \theta \end{bmatrix} \right) U \\ &= (I - I_F)\gamma^{-1} \left(\begin{bmatrix} 0 & 0 \\ 1 - \theta & \theta \end{bmatrix} - \begin{bmatrix} \gamma & 0 \\ (1 - \theta) & \theta \end{bmatrix} \right) U \\ &= (I - I_F) \begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix} \\ &= 0. \end{aligned}$$

Furthermore, we have

$$\begin{aligned} QU + (QU)^T - U^T QU &= \gamma^{-1} \begin{bmatrix} 1 - \theta & \theta \\ \theta & a \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} + (QU)^T - U^T QU \\ &= \gamma^{-1} \begin{bmatrix} 1 & \theta \\ \theta + a & a \end{bmatrix} + (QU)^T - U^T QU \\ &= \gamma^{-1} \begin{bmatrix} 2 & 2\theta + a \\ 2\theta + a & 2a \end{bmatrix} - \gamma^{-1} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & \theta \\ \theta + a & a \end{bmatrix} \\ &= \gamma^{-1} \begin{bmatrix} 2 & 2\theta + a \\ 2\theta + a & 2a \end{bmatrix} - \gamma^{-1} \begin{bmatrix} 1 + \theta + a & \theta + a \\ \theta + a & a \end{bmatrix} \\ &= \gamma^{-1} \begin{bmatrix} 1 - \theta - a & \theta \\ \theta & a \end{bmatrix} \end{aligned}$$

and

$$\begin{aligned} \frac{1}{2}U^T(P^T Q - S)^T B^\dagger(P^T Q - S)U &= \frac{1}{2} \begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \beta_1^{-1} & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix} \\ &= \gamma^{-1} \begin{bmatrix} \frac{\gamma}{2\beta_1} & \theta \\ \theta & a \end{bmatrix} \end{aligned}$$

which gives

$$\begin{aligned} W &= QU + (QU)^T - U^T QU - \frac{1}{2}U^T(P^T Q - S)^T B^\dagger(P^T Q - S)U \\ &= \gamma^{-1} \begin{bmatrix} 1 - \theta - a & \theta \\ \theta & a \end{bmatrix} - \gamma^{-1} \begin{bmatrix} \frac{\gamma}{2\beta_1} & \theta \\ \theta & a \end{bmatrix} \\ &= \gamma^{-1} \begin{bmatrix} 1 - \theta - a - \frac{\gamma}{2\beta_1} & \theta \\ \theta & a \end{bmatrix}. \end{aligned}$$

The final condition for convergence is then $Q \succ 0$ and $Q \succ 0$ which hold if

$$0 < 1 - \theta - \frac{\theta^2}{a} \quad \text{and} \quad 0 < 1 - \theta - \frac{\theta^2}{a} - a - \frac{\gamma}{2\beta_1}.$$

A.5 Forward-Backward with Nesterov Momentum

The update of forward-backward with Nesterov momentum [28, 29] can be seen as the application of the following frugal splitting operator

$$T_{(A_1, A_2)}(z_1, z_2) = \begin{pmatrix} \hat{z}_1 \\ \hat{z}_2 \end{pmatrix} = \begin{pmatrix} J_{\gamma A_2}(z_1 + \theta z_2 - \gamma A_1(z_1 + \theta z_2)) \\ \hat{z}_1 - z_1 \end{pmatrix}$$

where $\gamma > 0$, $\theta > 0$ and $(A_1, A_2) \in \mathcal{A}_2^{\{1\}}$. As with forward-backward with Polyak momentum, depending on how the Nesterov momentum is formulated there are other frugal splitting operator that also would yield a Nesterov momentum update. To derive a representation we choose the primal index $p = 2$ and note that the frugal splitting operator can be written as

$$\begin{aligned} y_1 &= A_1(z_1 + \theta z_2), \\ y_2 &= (\text{Id} + \gamma A_2)^{-1}(z_1 + \theta z_2 - \gamma y_1), \\ \hat{z}_1 &= y_2, \\ \hat{z}_2 &= y_2 - z_1. \end{aligned}$$

Inverting A_1 and $\text{Id} + \gamma A_2$ yield

$$\begin{aligned} z_1 + \theta z_2 &\in A_1^{-1}y_1, \\ \gamma^{-1}z_1 + \gamma^{-1}\theta z_2 - y_1 &\in A_2 y_2 + \gamma^{-1}y_2, \\ \hat{z}_1 &= z_1 - z_1 + y_2, \\ \hat{z}_2 &= z_2 - z_1 - z_2 + y_2. \end{aligned}$$

Rearranging so that $\Phi_{(A_1, A_2), 2}$ can be identified gives

$$\begin{aligned} z_1 + \theta z_2 &\in [y_2] + [A_1^{-1}y_1 - y_2], \\ \gamma^{-1}z_1 + \gamma^{-1}\theta z_2 &\in [\gamma^{-1}y_2] + [A_2 y_2 + y_1], \\ \hat{z}_1 &= z_1 - [z_1] + [y_2], \\ \hat{z}_2 &= z_2 - [z_1 + z_2] + [y_2] \end{aligned}$$

and we can identify

$$M = \begin{bmatrix} 0 & 1 \\ 0 & \gamma^{-1} \end{bmatrix}, \quad N = \begin{bmatrix} 1 & \theta \\ \gamma^{-1} & \gamma^{-1}\theta \end{bmatrix}, \quad V = \begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad U = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$$

This representation can be factored as $(2, SUP, SU, U, UP)$ where

$$S = \begin{bmatrix} 1 - \theta & \theta \\ \gamma^{-1}(1 - \theta) & \gamma^{-1}\theta \end{bmatrix} \quad \text{and} \quad P = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}.$$

For the convergence conditions we choose

$$Q = \gamma^{-1} \begin{bmatrix} 1 - \theta & \theta \\ \theta & a \end{bmatrix}$$

where $a > 0$. This yields

$$\begin{aligned}
0 &= (I - I_F)(P^T Q - S)U \\
&= (I - I_F) \left(\gamma^{-1} \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1-\theta & \theta \\ \theta & a \end{bmatrix} - \begin{bmatrix} 1-\theta & \theta \\ \gamma^{-1}(1-\theta) & \gamma^{-1}\theta \end{bmatrix} \right) U \\
&= (I - I_F) \left(\gamma^{-1} \begin{bmatrix} 0 & 0 \\ 1-\theta & \theta \end{bmatrix} - \begin{bmatrix} 1-\theta & \theta \\ \gamma^{-1}(1-\theta) & \gamma^{-1}\theta \end{bmatrix} \right) U \\
&= (I - I_F) \begin{bmatrix} -1+\theta & -\theta \\ 0 & 0 \end{bmatrix} U \\
&= (I - I_F) \begin{bmatrix} -1+\theta & -\theta \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \\
&= (I - I_F) \begin{bmatrix} -1 & -\theta \\ 0 & 0 \end{bmatrix} \\
&= 0
\end{aligned}$$

and

$$\begin{aligned}
QU + (QU)^T - U^T QU &= \gamma^{-1} \begin{bmatrix} 1-\theta & \theta \\ \theta & a \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} + (QU)^T - U^T QU \\
&= \gamma^{-1} \begin{bmatrix} 1 & \theta \\ \theta+a & a \end{bmatrix} + (QU)^T - U^T QU \\
&= \gamma^{-1} \begin{bmatrix} 2 & 2\theta+a \\ 2\theta+a & 2a \end{bmatrix} - \gamma^{-1} \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & \theta \\ \theta+a & a \end{bmatrix} \\
&= \gamma^{-1} \begin{bmatrix} 2 & 2\theta+a \\ 2\theta+a & 2a \end{bmatrix} - \gamma^{-1} \begin{bmatrix} 1+\theta+a & \theta+a \\ \theta+a & a \end{bmatrix} \\
&= \gamma^{-1} \begin{bmatrix} 1-\theta-a & \theta \\ \theta & a \end{bmatrix}
\end{aligned}$$

and

$$\begin{aligned}
\frac{1}{2}U^T(P^T Q - S)^T B^\dagger(P^T Q - S)U &= \frac{1}{2} \begin{bmatrix} -1 & 0 \\ -\theta & 0 \end{bmatrix} \begin{bmatrix} \beta_1^{-1} & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} -1 & -\theta \\ 0 & 0 \end{bmatrix} \\
&= \frac{1}{2\beta_1} \begin{bmatrix} -1 & 0 \\ -\theta & 0 \end{bmatrix} \begin{bmatrix} -1 & -\theta \\ 0 & 0 \end{bmatrix} \\
&= \frac{1}{2\beta_1} \begin{bmatrix} 1 & \theta \\ \theta & \theta^2 \end{bmatrix}.
\end{aligned}$$

Hence we have

$$\begin{aligned}
W &= QU + (QU)^T - U^T QU - \frac{1}{2}U^T(P^T Q - S)^T B^\dagger(P^T Q - S)U \\
&= \gamma^{-1} \begin{bmatrix} 1-\theta-a & \theta \\ \theta & a \end{bmatrix} - \frac{1}{2\beta_1} \begin{bmatrix} 1 & \theta \\ \theta & \theta^2 \end{bmatrix} \\
&= \gamma^{-1} \begin{bmatrix} 1-\theta-\hat{\gamma}-a & \theta(1-\hat{\gamma}) \\ \theta(1-\hat{\gamma}) & a-\theta^2\hat{\gamma} \end{bmatrix}
\end{aligned}$$

where $\hat{\gamma} = \frac{\gamma}{2\beta_1}$. The final condition for convergence is then $Q \succ 0$ and $W \succ 0$ which hold if

$$0 < 1 - \theta - \frac{\theta^2}{a}, \quad 0 < a - \theta^2\hat{\gamma} \quad \text{and} \quad 0 < 1 - \theta - \hat{\gamma} - a - \frac{\theta^2(1-\hat{\gamma})^2}{a-\theta^2\hat{\gamma}}.$$

A.6 A New Frugal Splitting Operator with Minimal Lifting

We will here derive a frugal splitting operator with minimal lifting that serves as the base of the algorithm in Theorem 8.1. This was also done in the proof of Theorem 8.1 but we will here be more detailed and verify the matrix calculations more carefully.

The frugal splitting operator will be over \mathcal{A}_n^F where $F = \{n - f, \dots, n - 1\}$ and $f = |F|$ and for notational convenience we let $R = \{2, \dots, n - f - 1\}$ with $r = |R|$. The representation of the new splitting is then (n, M, MK, HMK, HM) where

$$M = \left[\begin{array}{c|c|c|c} 1 & & & 1 \\ \hline 1 & \frac{1}{\theta}I & & 1 \\ \hline 1 & & & 1 \\ \hline 1 & & & 1 \end{array} \right] \in \mathbb{R}^{n \times n}, \quad K = \left[\begin{array}{c|c} \frac{1}{2} & \\ \hline & I \\ \hline & \\ \hline \frac{1}{2} & \end{array} \right] \in \mathbb{R}^{n \times (1+r)} \quad \text{and}$$

$$H = \theta \left[\begin{array}{c|c|c|c} \frac{1}{2+f} & & \frac{1}{2+f} \mathbf{1}^T & \frac{1}{2+f} \\ \hline & I & & \end{array} \right] \in \mathbb{R}^{(1+r) \times n}$$

where $\theta > 0$, I is the identity matrix in $\mathbb{R}^{r \times r}$, $\mathbf{1}$ are column vectors of ones with appropriate sizes and empty block denotes zero matrices. Since Proposition 6.1 is satisfied by these matrices, the operator corresponding to this representation is a frugal splitting operator. Let $T_{(\cdot)}: \mathcal{H}^{(1+r)} \rightarrow \mathcal{H}^{(1+r)}$ be this frugal splitting operator and consider the evaluation of $\hat{z} = T_A z$ where $A = (A_1, \dots, A_n) \in \mathcal{A}_n^F$. This can be written as

$$y = (M + \Phi_{A,n})^{-1} MKz$$

$$\hat{z} = z - HMKz + HMy.$$

Using the definition of inverse of $M + \Phi_{A,n}$ and writing out the matrices explicitly gives

$$\left[\begin{array}{c|c|c|c} 1 & & & \\ \hline 1 & \frac{1}{\theta}I & & \\ \hline 1 & & & \\ \hline 1 & & & \end{array} \right] z \in \left[\begin{array}{c|c|c|c} 1 & & & \\ \hline 1 & \frac{1}{\theta}I & & \\ \hline 1 & & & \\ \hline 2 & 1 & 1 & 1 \end{array} \right] y + \left[\begin{array}{c} A_1^{-1} \\ \vdots \\ A_{n-1}^{-1} \\ A_n \end{array} \right] y$$

$$\hat{z} = z - \left[\begin{array}{c|c} \theta & \\ \hline \theta \mathbf{1} & I \end{array} \right] z + \theta \left[\begin{array}{c|c|c|c} 1 & & & 1 \\ \hline 1 & \frac{1}{\theta}I & & 1 \end{array} \right] y$$

Setting $z = (z_1, \dots, z_{1+r})$, $\hat{z} = (\hat{z}_1, \dots, \hat{z}_{1+r})$, $y = (y_1, \dots, y_n)$ and writing out the corresponding equations for these matrix/operator expressions gives

$$z_1 \in (\text{Id} + A_1^{-1})y_1,$$

$$z_1 + \frac{1}{\theta}z_i \in y_1 + (\frac{1}{\theta} \text{Id} + A_i^{-1})y_i \quad \text{for all } i \in R,$$

$$z_1 \in y_1 + A_i^{-1}y_i \quad \text{for all } i \in F,$$

$$z_1 \in y_1 + \sum_{j=1}^{n-1} y_j + (\text{Id} + A_n)y_n,$$

$$\hat{z}_1 = (1 - \theta)z_1 + \theta(y_1 + y_n),$$

$$\hat{z}_i = -\theta z_1 + \theta(y_1 + y_n + \frac{1}{\theta}y_i) \quad \text{for all } i \in R.$$

Rearranging each line yields

$$y_1 = (\text{Id} + A_1^{-1})^{-1}z_1,$$

$$y_i = (\text{Id} + \theta A_i^{-1})^{-1}(\theta[z_1 - y_1] + z_i) \quad \text{for all } i \in R,$$

$$y_i = A_i(z_1 - y_1) \quad \text{for all } i \in F,$$

$$y_n = (\text{Id} + A_n)^{-1}([z_1 - y_1] - \sum_{j=1}^{n-1} y_j),$$

$$\hat{z}_1 = z_1 - \theta[z_1 - y_1] + \theta y_n,$$

$$\hat{z}_i = y_i - \theta[z_1 - y_1] + \theta y_n \quad \text{for all } i \in R.$$

Applying the Moreau identity to the first two lines gives and introducing variables x_i for all $x \in \{1, \dots, n\}$ that contains the results of all primal evaluations yields

$$\begin{aligned}
x_1 &= (\text{Id} + A_1)^{-1} z_1, \\
x_i &= (\text{Id} + \theta^{-1} A_i)^{-1} ([z_1 - y_1] + \theta^{-1} z_i) && \text{for all } i \in R, \\
x_i &= A_i(z_1 - y_1) && \text{for all } i \in F, \\
x_n &= (\text{Id} + A_n)^{-1} ([z_1 - y_1] - \sum_{j=1}^{n-1} y_j), \\
y_1 &= z_1 - x_1, \\
y_i &= \theta[z_1 - y_1] + z_i - \theta x_i = z_i + \theta(x_1 - x_i) && \text{for all } i \in R, \\
y_i &= x_i && \text{for all } i \in F, \\
y_n &= x_n, \\
\hat{z}_1 &= z_1 - \theta[z_1 - y_1] + \theta y_n, \\
\hat{z}_i &= y_i - \theta[z_1 - y_1] + \theta y_n && \text{for all } i \in R.
\end{aligned}$$

Eliminate the y_i variables

$$\begin{aligned}
x_1 &= (\text{Id} + A_1)^{-1} z_1, \\
x_i &= (\text{Id} + \theta^{-1} A_i)^{-1} (x_1 + \theta^{-1} z_i) && \text{for all } i \in R, \\
x_i &= A_i x_1 && \text{for all } i \in F, \\
\bar{x} &= \sum_{j \in F} x_j + \sum_{j \in R} z_j + \theta(x_1 - x_j), \\
x_n &= (\text{Id} + A_n)^{-1} (2x_1 - z_1 - \bar{x}), \\
\hat{z}_1 &= z_1 - \theta(x_1 - x_n), \\
\hat{z}_i &= z_i - \theta(x_i - x_n) && \text{for all } i \in R.
\end{aligned}$$

This concludes the derivations of the frugal splitting operator. To derive conditions for convergence we need a representation of the form (n, SUP, SU, U, UP) and since $U = HMK$ is invertible this is easily calculated as

$$\begin{aligned}
U = HMK &= \left[\begin{array}{c|c} \theta & I \\ \hline \theta \mathbf{1} & I \end{array} \right] \in \mathbb{R}^{(1+r) \times (1+r)}, & U^{-1} &= \left[\begin{array}{c|c} \theta^{-1} & I \\ \hline -\mathbf{1} & I \end{array} \right] \in \mathbb{R}^{(1+r) \times (1+r)}, \\
P = U^{-1}HM &= \left[\begin{array}{c|c|c|c} 1 & & & 1 \\ \hline & I & & \\ \hline & & & \\ \hline & & & 1 \end{array} \right] \in \mathbb{R}^{(1+r) \times n}, & S = MKU^{-1} &= \theta^{-1} \left[\begin{array}{c|c} 1 & I \\ \hline \mathbf{1} & \\ \hline 1 & \end{array} \right] \in \mathbb{R}^{n \times (1+r)}
\end{aligned}$$

Choosing

$$Q = \theta^{-1} \left[\begin{array}{c|c} 1 & I \\ \hline \mathbf{1} & \end{array} \right] \in \mathbb{R}^{(1+r) \times (1+r)}$$

then results in

$$\begin{aligned}
(I - I_F)(P^T Q - S)U &= (I - I_F) \left(\theta^{-1} \left[\begin{array}{c|c} 1 & \\ \hline & I \\ \hline & \\ \hline 1 & \end{array} \right] - \theta^{-1} \left[\begin{array}{c|c} 1 & \\ \hline & I \\ \hline 1 & \\ \hline 1 & \end{array} \right] \right) U \\
&= (I - I_F) \theta^{-1} \left[\begin{array}{c|c} & \\ \hline & \\ \hline -1 & \\ \hline & \end{array} \right] \left[\begin{array}{c|c} \theta & \\ \hline \theta \mathbf{1} & I \\ \hline \end{array} \right] \\
&= (I - I_F) \left[\begin{array}{c|c} & \\ \hline & \\ \hline -1 & \\ \hline & \end{array} \right] \\
&= 0
\end{aligned}$$

We further have

$$\begin{aligned}
QU + (QU)^T - U^T QU &= \theta^{-1}(U + U^T - U^T U) \\
&= \theta^{-1} \left(\left[\begin{array}{c|c} \theta & \\ \hline \theta \mathbf{1} & I \\ \hline \end{array} \right] + \left[\begin{array}{c|c} \theta & \theta \mathbf{1}^T \\ \hline & I \\ \hline \end{array} \right] - \left[\begin{array}{c|c} \theta & \theta \mathbf{1}^T \\ \hline & I \\ \hline \end{array} \right] \left[\begin{array}{c|c} \theta & \\ \hline \theta \mathbf{1} & I \\ \hline \end{array} \right] \right) \\
&= \theta^{-1} \left(\left[\begin{array}{c|c} 2\theta & \theta \mathbf{1}^T \\ \hline \theta \mathbf{1} & 2I \\ \hline \end{array} \right] - \left[\begin{array}{c|c} \theta^2(n-1-f) & \theta \mathbf{1}^T \\ \hline \theta \mathbf{1} & I \\ \hline \end{array} \right] \right) \\
&= \left[\begin{array}{c|c} 2 & \mathbf{1}^T \\ \hline \mathbf{1} & 2\theta^{-1}I \\ \hline \end{array} \right] - \left[\begin{array}{c|c} \theta(1+r) & \mathbf{1}^T \\ \hline \mathbf{1} & \theta^{-1}I \\ \hline \end{array} \right] \\
&= \left[\begin{array}{c|c} 2 - \theta(1+r) & \\ \hline & \theta^{-1}I \\ \hline \end{array} \right]
\end{aligned}$$

and

$$\begin{aligned}
\frac{1}{2}U^T(P^T Q - S)B^\dagger(P^T Q - S)U &= \frac{1}{2} \left[\begin{array}{c|c|c|c} & & -\mathbf{1}^T & \\ \hline & & & \\ \hline & & & \\ \hline & & & \end{array} \right] B^\dagger \left[\begin{array}{c|c} & \\ \hline & \\ \hline -1 & \\ \hline & \end{array} \right] \\
&= \frac{1}{2} \left[\begin{array}{c|c} \sum_{i \in F} \beta_i^{-1} & \\ \hline & \end{array} \right]
\end{aligned}$$

which, since $1 + r = n - 1 - f$, results in

$$\begin{aligned}
W &= QU + (QU)^T - U^T QU - \frac{1}{2}U^T(P^T Q - S)B^\dagger(P^T Q - S)U \\
&= \left[\begin{array}{c|c} 2 - \theta(1+r) & \\ \hline & \frac{1}{\theta}I \\ \hline \end{array} \right] - \frac{1}{2} \left[\begin{array}{c|c} \sum_{i \in F} \beta_i^{-1} & \\ \hline & \end{array} \right] \\
&= \left[\begin{array}{c|c} 2 - \theta(n-1-f) - \frac{1}{2} \sum_{i \in F} \beta_i^{-1} & \\ \hline & \frac{1}{\theta}I \\ \hline \end{array} \right].
\end{aligned}$$

We have $Q \succ 0$ and $W \succ 0$ if $\theta > 0$ and

$$0 < 2 - \theta(n-1-f) - \frac{1}{2} \sum_{i \in F} \beta_i^{-1}.$$

A.7 Minimal Lifting Method of Malitsky–Tam

Malitsky and Tam [6] presented a frugal splitting operator over \mathcal{A}_n with minimal lifting, $(\hat{z}_1, \dots, \hat{z}_{n-1}) = T_A(z_1, \dots, z_{n-1})$ where

$$\begin{aligned} x_1 &= J_{\gamma A_1}(z_1), \\ x_i &= J_{\gamma A_i}(z_i - z_{i-1} + x_{i-1}) && \text{for all } i \in \{2, \dots, n-1\}, \\ x_n &= J_{\gamma A_n}(-z_{n-1} + x_1 + x_{n-1}), \\ \hat{z}_i &= z_i + \theta(x_{i+1} - x_i) && \text{for all } i \in \{1, \dots, n-1\}. \end{aligned}$$

Note, this method this method was expanded by Briceño-Arias [31] to include inclusion problems that include compositions with linear operators.

Similarly to the derivation of our new method with minimal lifting, we derive a representation of this frugal splitting operator without any step-sizes, i.e., we set $\gamma = 1$ in the expressions above. As was done in Theorem 8.1, it is straightforward to modify the resulting convergence conditions to include step-sizes. In fact, since this frugal splitting operator have no forward evaluations, the convergence conditions for fixed point iteration with and without step-size will be the same. To start we select primal index $p = n$ and apply the Moreau identity to the $n - 1$ first resolvents which gives

$$\begin{aligned} y_1 &= J_{A_1^{-1}} z_1, \\ y_i &= J_{A_i^{-1}}(z_i - z_{i-1} + x_{i-1}) && \text{for all } i \in \{2, \dots, n-1\}, \\ y_n &= J_{A_n}(-z_{n-1} + x_1 + x_{n-1}), \\ x_1 &= z_1 - y_1, \\ x_i &= z_i - z_{i-1} + x_{i-1} - y_i && \text{for all } i \in \{2, \dots, n-1\}, \\ x_n &= y_n, \\ \hat{z}_i &= z_i + \theta(x_{i+1} - x_i) && \text{for all } i \in \{1, \dots, n-1\}. \end{aligned}$$

Looking at the expression of x_i for $i \in \{2, \dots, n-1\}$ we see

$$\begin{aligned} x_2 &= z_2 - z_1 + x_1 - y_2 = z_2 - y_1 - y_2 \\ x_3 &= z_3 - z_2 + x_2 - y_3 = z_3 - y_1 - y_2 - y_3 \\ &\vdots \\ x_i &= z_i - z_{i-1} + x_{i-1} - y_i = z_i - \sum_{j=1}^i y_j \end{aligned}$$

which gives

$$\hat{z}_i = z_i + \theta((z_{i+1} - \sum_{j=1}^{i+1} y_j) - (z_i - \sum_{j=1}^i y_j)) = z_i + \theta(z_{i+1} - z_i - y_{i+1})$$

for all $i \in \{1, \dots, n-2\}$ and

$$\hat{z}_{n-1} = z_{n-1} + \theta(y_n - (z_{n-1} - \sum_{j=1}^{n-1} y_j)) = z_{n-1} + \theta(-z_{n-1} + \sum_{j=1}^n y_j).$$

Inserting these expressions back in gives

$$\begin{aligned} y_1 &= J_{A_1^{-1}}(z_1), \\ y_i &= J_{A_i^{-1}}(z_i - \sum_{j=1}^{i-1} y_j) && \text{for all } i \in \{2, \dots, n-1\}, \\ y_n &= J_{A_n}(z_1 - y_1 - \sum_{j=1}^{n-1} y_j), \\ \hat{z}_i &= z_i - \theta(z_i - z_{i+1}) + \theta(-y_{i+1}) && \text{for all } i \in \{1, \dots, n-2\}, \\ \hat{z}_{n-1} &= z_{n-1} - \theta z_{n-1} + \theta \sum_{j=1}^n y_j. \end{aligned}$$

From this we can identify the representation (n, M, N, U, V) as

$$M = \begin{bmatrix} 1 & & & 1 \\ 1 & 1 & & 1 \\ \vdots & \ddots & \ddots & \vdots \\ 1 & \cdots & 1 & 1 \\ 1 & 0 & \cdots & 0 & 1 \end{bmatrix}, \quad N = \begin{bmatrix} 1 & & & \\ 0 & 1 & & \\ \vdots & \ddots & \ddots & \\ 0 & \cdots & 0 & 1 \\ 1 & 0 & \cdots & 0 \end{bmatrix}$$

and

$$U = \theta \begin{bmatrix} 1 & -1 & & & \\ & 1 & -1 & & \\ & & \ddots & \ddots & \\ & & & 1 & -1 \\ & & & & 1 \end{bmatrix}, \quad V = \theta \begin{bmatrix} 0 & -1 & & & \\ & 0 & -1 & & \\ & & \ddots & \ddots & \\ & & & 0 & -1 \\ 1 & 1 & \cdots & 1 & 1 & 1 \end{bmatrix}$$

where $M \in \mathbb{R}^{n \times n}$, $N \in \mathbb{R}^{n \times (n-1)}$, $U \in \mathbb{R}^{(n-1) \times (n-1)}$ and $V \in \mathbb{R}^{(n-1) \times n}$.

Since U is invertible a factorization of the form (n, SUP, SU, U, UP) must satisfy

$$S = NU^{-1} \quad \text{and} \quad P = U^{-1}V$$

which gives

$$S = \theta^{-1} \begin{bmatrix} 1 & 1 & \cdots & 1 \\ & 1 & \ddots & 1 \\ & & \ddots & \vdots \\ & & & 1 \\ 1 & 1 & \cdots & 1 \end{bmatrix} \in \mathbb{R}^{n \times (n-1)} \quad \text{and} \quad P = \begin{bmatrix} 1 & & & 1 \\ 1 & 1 & & 1 \\ \vdots & \ddots & \ddots & \vdots \\ 1 & \cdots & 1 & 1 & 1 \end{bmatrix} \in \mathbb{R}^{(n-1) \times n}.$$

The convergence conditions are then satisfied by $Q = \theta^{-1}I$ since

$$(I - I_F)(P^T Q - S)U = (S - S)U = 0$$

and

$$\begin{aligned} W &= QU + (QU)^T - U^T QU - \frac{1}{2}U^T (P^T Q - S)B^\dagger (P^T Q - S)U \\ &= QU + (QU)^T - U^T QU \\ &= \theta^{-1}(U + U^T - U^T U) \\ &= \begin{bmatrix} 2 & -1 & & & \\ -1 & 2 & -1 & & \\ & \ddots & \ddots & \ddots & \\ & & -1 & 2 & -1 \\ & & & -1 & 2 \end{bmatrix} - \theta \begin{bmatrix} 1 & & & & \\ -1 & 1 & & & \\ & \ddots & \ddots & & \\ & & -1 & 1 & \\ & & & -1 & 1 \end{bmatrix} \begin{bmatrix} 1 & -1 & & & \\ & 1 & -1 & & \\ & & \ddots & \ddots & \\ & & & 1 & -1 \\ & & & & 1 \end{bmatrix} \\ &= \begin{bmatrix} 2 & -1 & & & \\ -1 & 2 & -1 & & \\ & \ddots & \ddots & \ddots & \\ & & -1 & 2 & -1 \\ & & & -1 & 2 \end{bmatrix} - \theta \begin{bmatrix} 1 & -1 & & & \\ -1 & 2 & -1 & & \\ & \ddots & \ddots & \ddots & \\ & & -1 & 2 & -1 \\ & & & -1 & 2 \end{bmatrix} \\ &= (1 - \theta) \begin{bmatrix} 2 & -1 & & & \\ -1 & 2 & -1 & & \\ & \ddots & \ddots & \ddots & \\ & & -1 & 2 & -1 \\ & & & -1 & 2 \end{bmatrix} + \theta \begin{bmatrix} 1 & & & & \\ & 0 & & & \\ & & \ddots & & \\ & & & 0 & \end{bmatrix} \end{aligned}$$

The tridiagonal matrix is Toeplitz and have eigenvalues $2 + 2 \cos(\frac{k\pi}{n})$ for $k \in \{1, \dots, n-1\}$ and hence are $Q \succ 0$ and $W \succ 0$ for all $0 < \theta < 1$.

A.8 Minimal Lifting Method of Ryu

Ryu [5] presented a frugal splitting operator over \mathcal{A}_3 with minimal lifting, i.e., the lifting number is two. Let $T_{(\cdot)}: \mathcal{H}^2 \rightarrow \mathcal{H}^2$ be this frugal splitting operator and let $A \in \mathcal{A}_3$. The evaluation $(\hat{z}_1, \hat{z}_2) = T_A(z_1, z_2)$ is defined as

$$\begin{aligned} x_1 &= J_{A_1}(z_1), \\ x_2 &= J_{A_2}(z_2 + x_1), \\ x_3 &= J_{A_n}(-z_1 - z_2 + x_1 + x_2), \\ \hat{z}_1 &= z_1 + \theta(x_3 - x_1), \\ \hat{z}_2 &= z_2 + \theta(x_3 - x_2). \end{aligned}$$

To derive a representation of this we select the primal index $p = 3$ and apply the Moreau identity to the first two resolvents,

$$\begin{aligned} y_1 &= J_{A_1^{-1}}(z_1), \\ y_2 &= J_{A_2^{-1}}(z_1 + z_2 - y_1), \\ y_3 &= J_{A_n}(z_1 - 2y_1 - y_2), \\ \hat{z}_1 &= z_1 - \theta z_1 + \theta(y_1 + y_3), \\ \hat{z}_2 &= z_2 - \theta(z_1 + z_2) + \theta(y_1 + y_2 + y_3). \end{aligned}$$

From this we can identify

$$M = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix}, \quad N = \begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 0 \end{bmatrix}, \quad U = \theta \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \quad \text{and} \quad V = \theta \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix}$$

and $(3, M, N, U, V)$ is then a representation of $T_{(\cdot)}$. A factorization of the form $(3, SUP, SU, U, UP)$ needed for the convergence analysis is easily gotten since U is invertible and is then given by

$$S = \theta^{-1} \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 0 \end{bmatrix} \quad \text{and} \quad P = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}.$$

Choosing $Q = \theta^{-1}I$ yields

$$(I - I_F)(P^T Q - S)U = (P^T Q - S)U = (S - S)U = 0$$

and

$$\begin{aligned} W &= QU + (QU)^T - U^T QU - \frac{1}{2}U^T(P^T Q - S)^T B^\dagger(P^T Q - S)U \\ &= \theta^{-1}(U + U^T - U^T U) \\ &= \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} - \theta \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \\ &= (1 - \theta) \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} \end{aligned}$$

and it is clear that the final condition for convergence, $Q \succ 0$ and $W \succ 0$, hold if $0 < \theta < 1$.

A.9 Primal-Dual method of Chambolle–Pock

The primal-dual method made popular by Chambolle–Pock [7] can be seen as a fixed point iteration of the following frugal splitting operator over \mathcal{A}_2 ,

$$T_{(A_1, A_2)}(z_1, z_2) = \begin{pmatrix} \hat{z}_1 \\ \hat{z}_2 \end{pmatrix} = \begin{pmatrix} J_{\tau A_1}(z_1 - \tau z_2) \\ J_{\sigma A_2^{-1}}(z_2 + \sigma(2\hat{z}_1 - x_1)) \end{pmatrix}.$$

Since our generalized primal-dual resolvent also make use of a primal-dual formulation, deriving a representation of this frugal splitting operator is particularly easy. First we choose the primal index to $p = 1$ since we already have the first resolvent in primal form and the second in dual form. Introduce some intermediate variables

$$\begin{aligned} y_1 &= J_{\tau A_1}(z_1 - \tau z_2), \\ y_2 &= J_{\sigma A_2^{-1}}(z_2 - \sigma z_1 + 2\sigma y_1), \\ \hat{z}_1 &= y_1, \\ \hat{z}_2 &= y_2. \end{aligned}$$

Use the definition of a resolvent,

$$\begin{aligned} z_1 - \tau z_2 &\in y_1 + \tau A_1 y_1, \\ z_2 - \sigma z_1 + 2\sigma y_1 &\in y_2 + \sigma A_2^{-1} y_2, \\ \hat{z}_1 &= y_1, \\ \hat{z}_2 &= y_2, \end{aligned}$$

and rearrange to identify $\Phi_{A,1}$

$$\begin{aligned} \tau^{-1} z_1 - z_2 &\in [\tau^{-1} y_1 - y_2] + [A_1 y_1 + y_2], \\ \sigma^{-1} z_2 - z_1 &\in [-y_1 + \sigma^{-1} y_2] + [A_2^{-1} y_2 - y_1], \\ \hat{z}_1 &= z_1 - [z_1] + [y_1], \\ \hat{z}_2 &= z_2 - [z_2] + [y_2]. \end{aligned}$$

From this we can identify

$$M = \begin{bmatrix} \tau^{-1} & -1 \\ -1 & \sigma^{-1} \end{bmatrix}, \quad N = \begin{bmatrix} \tau^{-1} & -1 \\ -1 & \sigma^{-1} \end{bmatrix}, \quad V = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad U = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

This representation can be factorized as $(1, SUP, SU, U, UP)$ where

$$S = \begin{bmatrix} \tau^{-1} & -1 \\ -1 & \sigma^{-1} \end{bmatrix} \quad \text{and} \quad P = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$$

If we choose $Q = S$ we see that $Q \succ 0$ if $\sigma\tau < 1$ and

$$(I - I_F)(P^T Q - S)U = P^T Q - S = Q - S = 0$$

and

$$\begin{aligned} W &= QU + (QU)^T - U^T QU - \frac{1}{2}U^T (P^T Q - S)^T B^\dagger (P^T Q - S)U \\ &= Q + Q - Q \\ &= S. \end{aligned}$$

The convergence conditions with this choice of Q is then $Q \succ 0$ and $W \succ 0$ which hold as long as $S \succ 0$, i.e., as long as $\sigma\tau < 1$.

A.10 Projective Splitting

In [18] it was noted that the update of a synchronous version of projective splitting [32] can be seen as a generalized primal-dual resolvent over \mathcal{A}_n with representation $(n, M, M, \theta M, \theta M)$ where $\theta > 0$ and

$$M = \left[\begin{array}{c|c} \mathcal{T} & \mathbf{1} \\ \hline -\mathbf{1}^T & \tau_n^{-1} \end{array} \right] \in \mathbb{R}^{n \times n},$$

$\mathbf{1} \in \mathbb{R}^{n-1}$ is the vector of all ones, $\mathcal{T} \in \mathbb{R}^{(n-1) \times (n-1)}$ is a diagonal matrix with diagonal elements $\mathcal{T}_{i,i} = \tau_i$ for all $i \in \{1, \dots, n-1\}$ and $\tau_i > 0$ for all $i \in \{1, \dots, n\}$. In the projective splitting method the actual value of θ in each iteration is calculated based on the results of all resolvent evaluations but it is quite clear that the operator given by $(n, M, M, \theta M, \theta M)$ is a frugal splitting operator and we will show that fixed point iterations with a fixed θ converge for sufficiently small θ . With $T_{(\cdot)}: \mathcal{H}^n \rightarrow \mathcal{H}^n$ being this frugal splitting operator, the evaluation of $\hat{z} = T_A z$ for some $z \in \mathcal{H}^n$ and $A \in \mathcal{A}_n$ can be written as

$$\begin{aligned} Mz &\in My + \Phi_{A,n}y = (M + \Gamma_n)y + \Delta_{A,n}y \\ \hat{z} &= z - \theta M(z - y) = (I - \theta M)z + \theta My \end{aligned}$$

which explicitly becomes

$$\begin{aligned} \left[\begin{array}{c|c} \mathcal{T} & \mathbf{1} \\ \hline -\mathbf{1}^T & \tau_n^{-1} \end{array} \right] z &\in \left(\left[\begin{array}{c|c} \mathcal{T} & \mathbf{1} \\ \hline -\mathbf{1}^T & \tau_n^{-1} \end{array} \right] + \left[\begin{array}{c|c} A_1^{-1} & \\ \hline & \ddots \\ \hline & & A_{n-1}^{-1} \\ \hline & & & A_n \end{array} \right] \right) y \\ \hat{z} &= z - \theta \left[\begin{array}{c|c} \mathcal{T} & \mathbf{1} \\ \hline -\mathbf{1}^T & \tau_n^{-1} \end{array} \right] z + \theta \left[\begin{array}{c|c} \mathcal{T} & \mathbf{1} \\ \hline -\mathbf{1}^T & \tau_n^{-1} \end{array} \right] y. \end{aligned}$$

Setting $z = (z_1, \dots, z_n)$, $\hat{z} = (\hat{z}_1, \dots, \hat{z}_n)$, $y = (y_1, \dots, y_n)$ and writing out each line gives

$$\begin{aligned} \tau_i z_i + z_n &\in (\tau_i \text{Id} + A_i^{-1})y_i \quad \text{for all } i \in \{1, \dots, n-1\}, \\ \tau_n^{-1} z_n - \sum_{j=1}^{n-1} z_j &\in (\tau_n^{-1} \text{Id} + A_n)y_n, \\ \hat{z}_i &= z_i - \theta(\tau_i z_i + z_n) + \theta(\tau_i y_i + y_n) \quad \text{for all } i \in \{1, \dots, n-1\}, \\ \hat{z}_n &= z_n - \theta(\tau_n^{-1} z_n - \sum_{j=1}^{n-1} z_j) + \theta(\tau_n^{-1} y_n - \sum_{j=1}^{n-1} y_j). \end{aligned}$$

Rewriting these equations using resolvents gives

$$\begin{aligned} y_i &= \mathbf{J}_{\tau_i^{-1} A_i^{-1}}(z_i + \tau_i^{-1} z_n) && \text{for all } i \in \{1, \dots, n-1\}, \\ y_n &= \mathbf{J}_{\tau_n A_n}(z_n - \tau_n \sum_{j=1}^{n-1} z_j), \\ \hat{z}_i &= z_i - \theta(\tau_i z_i + z_n) + \theta(\tau_i y_i + y_n) && \text{for all } i \in \{1, \dots, n-1\}, \\ \hat{z}_n &= z_n - \theta(\tau_n^{-1} z_n - \sum_{j=1}^{n-1} z_j) + \theta(\tau_n^{-1} y_n - \sum_{j=1}^{n-1} y_j). \end{aligned}$$

Applying the Moreau identity and introducing variables for the primal evaluation yields

$$\begin{aligned}
x_i &= \mathbf{J}_{\tau_i A_i}(\tau_i z_i + z_n) && \text{for all } i \in \{1, \dots, n-1\}, \\
x_n &= \mathbf{J}_{\tau_n A_n}(z_n - \tau_n \sum_{j=1}^{n-1} z_j), \\
y_i &= z_i + \tau_i^{-1} z_n - \tau_i^{-1} x_i && \text{for all } i \in \{1, \dots, n-1\}, \\
y_n &= x_n, \\
\hat{z}_i &= z_i - \theta(\tau_i z_i + z_n) + \theta(\tau_i y_i + y_n) && \text{for all } i \in \{1, \dots, n-1\}, \\
\hat{z}_n &= z_n - \theta(\tau_n^{-1} z_n - \sum_{j=1}^{n-1} z_j) + \theta(\tau_n^{-1} y_n - \sum_{j=1}^{n-1} y_j).
\end{aligned}$$

Eliminating y_i for all $i \in \{1, \dots, n\}$ yields

$$\begin{aligned}
x_i &= \mathbf{J}_{\tau_i A_i}(\tau_i z_i + z_n) && \text{for all } i \in \{1, \dots, n-1\}, \\
x_n &= \mathbf{J}_{\tau_n A_n}(z_n - \tau_n \sum_{j=1}^{n-1} z_j), \\
\hat{z}_i &= z_i - \theta(x_i - x_n) && \text{for all } i \in \{1, \dots, n-1\}, \\
\hat{z}_n &= z_n - \theta(\tau_n^{-1} - \sum_{j=1}^{n-1} \tau_i^{-1}) z_n + \theta(\tau_n^{-1} x_n + \sum_{j=1}^{n-1} \tau_i^{-1} x_j).
\end{aligned}$$

To get convergence conditions we select $Q = \theta^{-1}I$ and factor the representations as (n, SUP, SU, U, UP) where

$$U = \theta M, \quad S = \theta^{-1}I \quad \text{and} \quad P = I$$

where $I \in \mathbb{R}^{n \times n}$ is the identity matrix. This results in

$$(I - I_F)(P^T Q - S)U = (P^T Q - S)U = (\theta^{-1}I - \theta^{-1}I)U = 0$$

and hence

$$\begin{aligned}
W &= QU + (QU)^T - U^T QU - \frac{1}{2}U^T(P^T Q - S)B^\dagger(P^T Q - S)U \\
&= \theta^{-1}(U + U^T - U^T U) \\
&= M + M^T - \theta M^T M \\
&= 2 \left[\begin{array}{c|c} \mathcal{T} & \\ \hline \mathbf{1}^T & \tau_n^{-1} \end{array} \right] - \theta \left[\begin{array}{c|c} \mathcal{T} & -\mathbf{1} \\ \hline \mathbf{1}^T & \tau_n^{-1} \end{array} \right] \left[\begin{array}{c|c} \mathcal{T} & \mathbf{1} \\ \hline -\mathbf{1}^T & \tau_n^{-1} \end{array} \right] \\
&= 2 \left[\begin{array}{c|c} \mathcal{T} & \\ \hline \mathbf{1}^T & \tau_n^{-1} \end{array} \right] - \theta \left[\begin{array}{c|c} \mathcal{T}^2 + \mathbf{1}\mathbf{1}^T & \mathcal{T}\mathbf{1} - \tau_n^{-1}\mathbf{1} \\ \hline \mathbf{1}^T \mathcal{T} - \tau_n^{-1}\mathbf{1}^T & \tau_n^{-2} + n - 1 \end{array} \right] \\
&= \theta \left[\begin{array}{c|c} 2\theta^{-1}\mathcal{T} & \\ \hline 2\theta^{-1}\tau_n^{-1} & \end{array} \right] + \theta \left[\begin{array}{c|c} -\mathcal{T}^2 - \mathbf{1}\mathbf{1}^T & -\mathcal{T}\mathbf{1} + \tau_n^{-1}\mathbf{1} \\ \hline -\mathbf{1}^T \mathcal{T} + \tau_n^{-1}\mathbf{1}^T & -\tau_n^{-2} - n + 1 \end{array} \right] \\
&= \theta \left[\begin{array}{c|c} 2\theta^{-1}\mathcal{T} - \mathcal{T}^2 - \mathbf{1}\mathbf{1}^T & -\mathcal{T}\mathbf{1} + \tau_n^{-1}\mathbf{1} \\ \hline -\mathbf{1}^T \mathcal{T} + \tau_n^{-1}\mathbf{1}^T & 2\theta^{-1}\tau_n^{-1} - \tau_n^{-2} - n + 1 \end{array} \right].
\end{aligned}$$

For convergence it is required that $Q \succ 0$ and $W \succ 0$. As long as $\theta > 0$ then $Q \succ 0$ and it is clear that $W \succ 0$ for sufficiently small θ . If $\tau_i = \tau_n^{-1}$ for all $i \in \{1, \dots, n-1\}$, then W simplifies to

$$W = \theta \tau_n^{-1} \left[\begin{array}{c|c} (2\theta^{-1} - \tau_n^{-1})I - \mathbf{1}\mathbf{1}^T & \\ \hline & 2\theta^{-1} - \tau_n^{-1} - n + 1 \end{array} \right].$$

and $W \succ 0$ if

$$\theta < \frac{2}{n-1 + \tau_n^{-1}}.$$

Note, although τ_i appear as step-sizes in the resolvents it could be argued that they are in fact not proper step-sizes since they appear outside the resolvents as well. If we introduce $\gamma > 0$ and apply this fixed point iteration to $\gamma A = (\gamma A_1, \dots, \gamma A_n)$ instead we get the following fixed point iteration

$$\begin{aligned}
 x_i &= J_{\tau_i \gamma A_i}(\tau_i z_i + z_n) && \text{for all } i \in \{1, \dots, n-1\}, \\
 x_n &= J_{\tau_n \gamma A_n}(z_n - \tau_n \sum_{j=1}^{n-1} z_j), \\
 \hat{z}_i &= z_i - \theta(x_i - x_n) && \text{for all } i \in \{1, \dots, n-1\}, \\
 \hat{z}_n &= z_n - \theta(\tau_n^{-1} - \sum_{j=1}^{n-1} \tau_i^{-1})z_n + \theta(\tau_n^{-1}x_n + \sum_{j=1}^{n-1} \tau_i^{-1}x_j)
 \end{aligned}$$

which converge for all $\gamma > 0$, as long as the choice of τ_i and θ are such that the Q and W above are positive definite. Hence, even if we might have to choose τ_i and θ small, the step-sizes in the resolvents can always be made arbitrarily large.