RECOLLEMENTS OF PARTIALLY WRAPPED FUKAYA CATEGORIES AND SURFACE CUTS

WEN CHANG, HAIBO JIN, AND SIBYLLE SCHROLL

ABSTRACT. In this paper we use recollements to investigate partially wrapped Fukaya categories of surfaces with marked points. In particular, we show that cutting surfaces gives rise to recollements of the corresponding partially wrapped Fukaya categories. Our approach is based on the fact that the partially wrapped Fukaya category of a surface with marked points is triangle equivalent to the perfect derived category of a homologically smooth and proper graded gentle algebra with zero differential as shown by Haiden, Katzarkov and Kontsevich. Using this, we study particular generators of partially wrapped Fukaya categories, namely full exceptional sequences, silting objects and simple-minded collections. In particular, we fully characterise the existence of full exceptional sequences and we give an example of a partially wrapped Fukaya category which does not admit a silting object, that is a generator with no positive self-extensions.

INTRODUCTION

Gentle algebras, first arising in the representation theory of associative algebras [AH, AS], connect to many other areas of mathematics such as Lie algebras, dimer models, cluster theory, N=2 Gauge theories and homological mirror symmetry. In the context of homological mirror symmetry, the partially wrapped Fukaya category of a graded marked surface is triangle equivalent to the bounded derived category of a homologically smooth graded gentle algebra with zero differential [HKK, LP].

On the other hand, the notion of recollements of triangulated categories was first introduced by A. A. Beilinson, J. Bernstein and P. Deligne in [BBD] (see Definition 1.5 for more details). Recollements play an important role in algebraic geometry and representation theory, in particular when comparing derived categories and their invariants (see for example [AKLY, Hap, Han, Ke2, Ko, KoN]).

In this paper, we use recollements to investigate partially wrapped Fukaya categories of marked surfaces. In particular, we show that cutting the surface gives rise to a recollement of the associated partially wrapped Fukaya categories. More precisely, we show

Theorem A (Theorem 4.1). Let $W(S, M, \eta(\Delta))$ be the partially wrapped Fukaya category associated to a graded surface dissection (S, M, Δ) with induced line field $\eta(\Delta)$. Let $L \subset \Delta$ be a subset of Δ such that the graded marked surface $W(S(L), M(L), \eta(L))$ generated by L (as graded ribbon subgraph of Δ) has no marked points in the interior. Then there is a recollement of partially wrapped Fukaya categories

$$\mathcal{W}(S_L, M_L, \eta(\Delta_L)) \longrightarrow \mathcal{W}(S, M, \eta(\Delta)) \longrightarrow \mathcal{W}(S(L), M(L), \eta(L))$$
(0.1)

where the left side corresponds to the graded cut marked surface along L.

Our proof is based on the fact that the partially wrapped Fukaya category $(S, M, \eta(\Delta))$ is triangle equivalent to the bounded derived category of a proper and homologically smooth differential

Date: April 4, 2023.

²⁰¹⁰ Mathematics Subject Classification. 16E35, 57M50.

graded gentle algebra with zero differential [HKK, LP]. In fact, we start with a more general setting. Namely, we first consider recollements arising from graded quadratic monomial algebras. For this, given a graded quadratic monomial algebra A = kQ/I, we introduce the notion of *partial cofibrant dg algebra resolution* A_J with respect to a subset J of I by 'resolving' only the relations in J (see Definition 2.1 for the details). Then we show the following

Theorem B (Theorem 2.5). Let A = kQ/I be a graded quadratic monomial algebra and let $J \subset I$. Let A_J be the partial cofibrant dg algebra resolution of A. Then there is a natural quasi-isomorphism $A_J \simeq A$ of dg k-algebras.

Moreover, if J is the set of relations at some vertices corresponding to an idempotent $e \in A$, the above quasi-isomorphism induces the following recollement (see Theorem 2.10)

$$\mathcal{D}(A_e) \xrightarrow{i^*} \mathcal{D}(A) \xrightarrow{?\otimes_{k}^{\mathbf{L}}eA} \mathcal{D}(eAe) , \qquad (0.2)$$

where A_e is again a quadratic monomial algebra explicitly described in terms of quiver and relations defined in Definition 2.8. We remark that there are serval different ways to construct dg algebras Bsuch that $\mathcal{D}(B)$ fits the left side of the recollement in (0.2), see for example [BCL, D, KaY2, Ke3, NS]. Here we give a new and concrete construction for any graded quadratic monomial algebra, which plays an important role in our study of partially wrapped Fukaya categories.

If in the recollement in (0.2) two out of the three algebras A, eAe, and A_e are homologically smooth and proper, so is the third one. Moreover, in this case (0.2) restricts to a recollement of perfect derived categories (see Proposition 4.2) and gives rise to the recollement (0.1). More generally we show that it can be extended to an unbounded ladder of recollements (see Proposition 4.11) as introduced in [AKLY].

The second aim of this paper is to study particular generators of partially wrapped Fukaya categories, namely, full exceptional sequences, silting objects and simple minded collections. Exceptional sequences originate in algebraic geometry [GR, Ru] and are instrumental in the construction of the Fukaya-Seidel categories [S]. Silting objects and simple-minded collections are some of the standard tools for studying triangulated categories and are important in Koszul duality [BGS, Ke1, KN, KoY, Ri].

We first note that in terms of the associated homologically smooth, proper graded gentle algebras, the recollement in (0.2) gives rise to reduction with respect to silting objects of the associated perfect derived categories. This induces a bijection (see Theorem 5.1)

{silting objects in $per(A_e)$ } \longleftrightarrow {silting objects in per(A) containing P},

which gives us a way to study silting theory of A in terms of the 'smaller' algebra A_e . We show that a similar result holds for reduction of simple-minded collections.

We then study which partially wrapped Fukaya categories admit full exceptional sequences, silting objects and simple-minded collections.

In general, it is difficult to determine whether a triangulated category has such generators or not. In the case of a partial wrapped Fukaya category, we first give a full characterization of the existence of full exceptional sequences.

Theorem C (Theorem 5.6). Let $\mathcal{W}(S, M, \eta(\Delta))$ be a partially wrapped Fukaya category. Then $\mathcal{W}(S, M, \eta(\Delta))$ has a full exceptional sequence if and only if (S, M) is not homeomorphic to a marked surface of genus ≥ 1 with exactly one boundary and one marked point on that boundary.

Next we study the existence of silting objects and simple-minded collections. Denote by $A^{(n)}$ the family of graded gentle algebras with parameters a_i, b_i for $1 \le i \le n$ associated to a surface of

genus n with exactly one boundary component and one marked point on that boundary component (see Section 5.2 for the precise definitions). We then show the following.

Theorem D (Theorem 5.10). Let $\mathcal{W}(S, M, \eta(\Delta))$ be a partially wrapped Fukaya category. Then $\mathcal{W}(S, M, \eta(\Delta))$ has a silting object if and only if it has a simple-minded collection and this is the case if

(1) $\mathcal{W}(S, M, \eta(\Delta))$ is not triangle equivalent to $\mathcal{D}^{\mathrm{b}}(A^{(n)})$, or

(2) $\mathcal{W}(S, M, \eta(\Delta))$ is triangle equivalent to $\mathcal{D}^{\mathrm{b}}(A^{(n)})$ with $a_i \neq 1$ or $b_i \neq 1$ for any $1 \leq i \leq n$.

We note that in case that n = 1, the converse of Theorem D holds. Namely, we show by hand that $A^{(1)}$ with the grading $a_1 = b_1 = 1$ does not admit a silting object by checking in a case by case analysis, that every indecomposable object in $\mathcal{D}^{\mathbf{b}}(A^{(1)})$ has a positive self-extension.

Acknowledgements. The authors would like to thank Bernhard Keller and Dong Yang for helpful comments regarding recollements and cofibrant dg algebra resolutions.

CONVENTIONS

In this paper, all algebras will be assumed to be over a base field k. Arrows in a quiver are composed from left to right as follows: for arrows a and b we write ab for the path from the source of a to the target of b. We adopt the convention that maps are composed from right to left, that is if $f: X \to Y$ and $g: Y \to Z$ then $gf: X \to Z$. In general, we consider right modules. We denote by \mathbb{Z} the set of integer numbers, and by \mathbb{Z}^* the set of non-zero integer numbers. We use notation |M| to represent the number of elements in a finite set M.

1. Preliminaries

1.1. **DG algebras.** Let A be a differential graded (dg) k-algebra, that is, a graded algebra endowed with a compatible structure of a k-complex. A (right) dg A-module is a graded A-module endowed with a compatible structure of a k-complex. Let $\mathcal{D}(A)$ be the derived category of right dg A-modules (see [Ke1, Ke4]). It is a triangulated category obtained from the category of dg A-modules by formally inverting all quasi-isomorphisms. The shift functor is given by the shift of complexes.

We denote by $\operatorname{per}(A)$ the *perfect category* of A, which is the smallest triangulated subcategory of $\mathcal{D}(A)$ generated by A and which is closed under direct summands. We denote by $\mathcal{D}^{\mathrm{b}}(A)$ the full subcategory of $\mathcal{D}(A)$ consisting of the objects M whose total cohomology is finite-dimensional (that is, $\dim_k \bigoplus_n \operatorname{H}^n(M) < \infty$). If $\operatorname{per}(A) \subset \mathcal{D}^{\mathrm{b}}(A)$, that is, the cohomology algebra $\mathcal{H}^*(A)$ is finite dimensional, we say A is *proper*. If $A \in \operatorname{per}(A \otimes_k A^{\mathrm{op}})$, we say A is *homologically smooth*. The following observations are well-known.

Lemma 1.1. Let A be a homologically smooth dg k-algebra. Then $\mathcal{D}^{\mathbf{b}}(A) \subset \mathsf{per}(A)$. If A is also proper, then $\mathcal{D}^{\mathbf{b}}(A) = \mathsf{per}(A)$.

Lemma 1.2. Let A be a dg k-algebra and let X be a dg A-module. Then the following are equivalent.

(1) $X \in \mathcal{D}^{\mathrm{b}}(A)$.

(2) For any $P \in per(A)$, $\mathbb{R}\mathscr{H}om_A(P, X)$ has finite dimensional total cohomology.

Lemma 1.3. [Ke1] Let A, B be two dg k-algebras. Let $F : \mathcal{D}(A) \to \mathcal{D}(B)$ be a triangle functor, which commutes with infinite direct sums and restricts to a functor $F|_{per} : per(A) \to per(B)$. Then F is a triangle equivalence if and only if $F|_{per}$ is a triangle equivalence. Moreover, in this case, Falso restricts to an equivalence $F : \mathcal{D}^{\mathbf{b}}(A) \xrightarrow{\simeq} \mathcal{D}^{\mathbf{b}}(B)$.

Lemma 1.4. [AKLY, Lemma 2.7] Let A, B be two dg k-algebras. Let $F : \mathcal{D}(A) \to \mathcal{D}(B)$ be a triangle functor with a right adjoint $G : \mathcal{D}(B) \to \mathcal{D}(A)$. If $F(\mathsf{per}(A)) \subset \mathsf{per}(B)$, then $G(\mathcal{D}^{\mathrm{b}}(B)) \subset \mathcal{D}^{\mathrm{b}}(A)$.

Proof. Let $Y \in \mathcal{D}^{b}(B)$. By Lemma 1.2, it suffices to show that, for any $P \in \text{per}(A)$, $\mathbb{R} \mathscr{H}om_{A}(P, G(Y))$ has finite dimensional total cohomology. Since (F, G) is an adjoint pair, we have

$$\mathrm{H}^{n}\mathbf{R}\mathscr{H}\mathrm{om}_{A}(P,G(Y)) = \mathrm{Hom}_{\mathcal{D}(A)}(P,G(Y)[n]) = \mathrm{Hom}_{\mathcal{D}(B)}(F(P),Y[n]) = \mathrm{H}^{n}\mathbf{R}\mathscr{H}\mathrm{om}_{B}(F(P),Y).$$

Since $F(P) \in \mathsf{per}(B)$ and $Y \in \mathcal{D}^{\mathsf{b}}(B)$, we have $\mathbf{R}\mathscr{H}\mathsf{om}_B(F(P), Y)$ has finite dimensional total cohomology by Lemma 1.2, and so does $\mathbf{R}\mathscr{H}\mathsf{om}_P(P, G(Y))$.

1.2. **Recollements.** In this section we recall some results on recollement. Recollements, introduced by Beilinson, Bernstein and Deligne in the 1980s, provide a powerful tool for studying problems in triangulated categories and algebraic geometry. Roughly speaking, a recollement consists of three triangulated categories linked by two triangulated functors both of which have left and right adjoint functors. The notion of recollement is an analogue of an exact sequence for triangulated categories, which generalizes derived equivalences and is closely related to homological ring epimorphisms.

Definition 1.5 ([BBD]). Let \mathcal{T}' , \mathcal{T} and \mathcal{T}'' be triangulated categories. A *recollement* of \mathcal{T} in terms of \mathcal{T}' and \mathcal{T}'' is the following diagram:

$$\mathcal{T}' \xrightarrow{i^*}_{i_*} \mathcal{T} \xrightarrow{j^*}_{j_*} \mathcal{T}''$$

$$(1.1)$$

satisfying

- (1) $(i^*, i_*, i^!)$ and $(j_!, j^*, j_*)$ are adjoint triples;
- (2) $i_*, j_*, j_!$ are fully faithful;

(3) $j^* \circ i_* = 0;$

(4) for any object X in \mathcal{T} , there are two triangles:

$$i_*i^!X \longrightarrow X \longrightarrow j_*j^*X \longrightarrow i_*i^!X[1]$$

 $j_!j^*X \longrightarrow X \longrightarrow i_*i^*X \longrightarrow j_!j^*X[1]$

where the morphisms are induced by the units and conuits of the adjunctions.

We give a well-known example of recollement, which follows from [CPS].

Example 1.6. Let A be a ring and let $e \in A$ be an idempotent. Assume the ideal AeA of A is stratifying (*i.e.* $Ae \otimes_{eAe}^{\mathbf{L}} eA \to AeA$ is an isomorphism). Then the associated ring epimorphism $A \to AeA$ induces a recollement of derived module categories of the form

$$\mathcal{D}(A/AeA) \xrightarrow{i_* = :: \otimes_A^{\mathbf{L}} A/AeA} \mathcal{D}(A) \xrightarrow{j_! = : \otimes_{eAe}^{\mathbf{L}} eA} \mathcal{D}(eAe)$$
$$i_! = \mathbf{R} \mathscr{H}_{\mathrm{Om}_A(A/AeA,?)} \qquad j_* = \mathbf{R} \mathscr{H}_{\mathrm{Om}_{eAe}(Ae,?)}$$

Most examples of recollements in the literature are known to be of this form, up to derived equivalence. We give a general version of Example 1.6 in the general dg setting in the next section.

Let \mathcal{X}, \mathcal{Y} and \mathcal{Z} be three full subcategories of a triangulated category \mathcal{T} . We call the triple $(\mathcal{X}, \mathcal{Y}, \mathcal{Z})$ a *TTF triple* of \mathcal{T} , if we have two *t*-structures $\mathcal{T} = \mathcal{X} \perp \mathcal{Y}$ and $\mathcal{T} = \mathcal{Y} \perp \mathcal{Z}$. It immediately follows that if $(\mathcal{X}, \mathcal{Y}, \mathcal{Z})$ is a TTF triple, then \mathcal{X}, \mathcal{Y} and \mathcal{Z} are all triangulated subcategories of \mathcal{T} . Note that given a recollement (1.1), we have an induced TTF triple $(j_!(\mathcal{T}''), i_*(\mathcal{T}')), j_*(\mathcal{T}''))$. We say two recollements are equivalent if the TTF triples associated to them coincide. The following observation is useful (see for example [BBD, Section 1.4.4] or [N3, Section 9.2]).

Proposition 1.7. Let \mathcal{T} be a k-linear triangulated category. There is a one-to-one correspondence between equivalence classes of recollements of \mathcal{T} and TTF triples of \mathcal{T} .

We need the following lemma for later use.

Lemma 1.8. Let (1.1) be a recollement. Then

- (1) i^* admits a left adjoint F if and only if $j_!$ admits a left adjoint. Moreover, in this case, F is fully faithful, and $(F(\mathcal{T}'), j_!(\mathcal{T}''), i_*(\mathcal{T}'))$ is a TTF triple.
- (2) i' admits a right adjoint G if and only if j* admits a right adjoint. Moreover, in this case, G is fully faithful and (i*(T'), j*(T''), G(T')) is a TTF triple.

Proof. See for example [N3, Proposition 9.1.18] and [M, Theorem 2.5], and their duals.

1.3. Homological epimorphisms of dg algebras. Homological epimorphisms were introduced by Geigle and Lenzing in [GL] to characterize when a homomorphism of rings $\phi : R \to S$ induces a full embedding of bounded derived categories $\mathcal{D}^{b}(\text{mod } S) \to \mathcal{D}^{b}(\text{mod } R)$. The dg generalization of this notion was studied by [P] and [NS]. In this section, we recall some basic facts on homological epimorphisms of dg algebras.

Let A and B be two dg k-algebras and let $f : A \to B$ be a homomorphism of dg algebras. We may regard B as a dg A-bimodule and any dg B-module M has a natural dg A-module structure induced by f. We call f a homological epimorphism if the induced functor $f_* : \mathcal{D}(B) \to \mathcal{D}(A)$ is fully faithful. Let $X \to A \xrightarrow{f} B \to X[1]$ be the triangle in $\mathcal{D}(A \otimes_k A^{\text{op}})$ induced by f. The following result characterizes when a dg homomorphism is a homological epimorphism.

Lemma 1.9. [NS, Lemma 4] Let $f : A \to B$ be a homomorphism of dg algebra. The following are equivalent

- (1) f is a homological epimorphism;
- (2) $B \otimes^{\mathbf{L}}_{A} B \cong B$ in $\mathcal{D}(B)$;
- (3) There is a recollement



where Loc(X) is the smallest triangulated subcategory of $\mathcal{D}(A)$ containing X and closed under infinite direct sums.

The following example generalizes Example 1.6.

Example 1.10. [KaY1, Proposition 2.10] Let A be a k-algebra and $e \in A$ be an idempotent. Then there is a dg k-algebra B and a morphism of dg algebras $f : A \to B$, such that there is a recollement



Moreover, B is also non-positive with $H^0(B) = A/AeA$.

1.4. Compactly generated triangulated categories. Let \mathcal{T} be a triangulated category with infinite direct sums. An object $X \in \mathcal{T}$ is called *compact*, if the functor $\operatorname{Hom}_{\mathcal{T}}(X, ?)$ commutes with infinite direct sums. We denote by \mathcal{T}^{c} the subcategory of \mathcal{T} consisting of compact objects. It is easy to see \mathcal{T}^{c} is a thick subcategory, that is, it is a triangulated subcategory of \mathcal{T} closed under direct summands.

We say that \mathcal{T} is *compactly generated* if \mathcal{T} admits a set of small generators \mathcal{C} , that is, if there is a subset \mathcal{C} of compact objects, such that \mathcal{T} coincides with the localising subcategory $\mathsf{Loc}(\mathcal{C})$ (which is the smallest sub triangulated category of \mathcal{T} containing \mathcal{C} and closed under infinite sums).

In a compactly generated triangulated category, all compact objects can be obtained from a set of compact generators, that is, we have the following.

Theorem 1.11. [N1, Section 2.2][Ke1, Theorem 5.3] Let \mathcal{T} be a compactly generated triangulated category and let \mathcal{C} be a set of compact generators. Then $\mathcal{T}^{c} = \text{thick}(\mathcal{C})$.

Example 1.12. Let A be a finite-dimensional algebra over a field k. Then the unbounded derived category $\mathcal{D}(A)$ is compactly generated by the free module A. So $\mathcal{D}(A)^{c} = \operatorname{per}(A)$ coincides with $\mathsf{K}^{\mathsf{b}}(\mathsf{proj} A)$.

We need the following lemma later.

Lemma 1.13. [N2, Theorems 4.1 and 5.1] Let $F : \mathcal{C} \to \mathcal{D}$ be a triangle functor between two compactly generated triangulated categories. Assume F admits a right adjoint $G : \mathcal{D} \to \mathcal{C}$. Then the following conditions are equivalent.

- (1) G admits a right adjoint.
- (2) F sends compact objects to compact objects.
- (3) G preserves coproducts.

The following result is important for our purpose.

Theorem 1.14. [N1, Theorem 2.1] Let \mathcal{T} be a compactly generated triangulated category and let \mathcal{R} be a set of compact objects. Let $\mathcal{T}'' := \mathsf{Loc}(\mathcal{R})$. Then there is a recollement

$$\mathcal{T}' \underbrace{\overset{i^*}{\underbrace{i_*}}}_{i^!} \mathcal{T} \underbrace{\overset{j_!}{\underbrace{j^*}}}_{j_*} \mathcal{T}'$$

where \mathcal{T}' is also compactly generated. Moreover, i^* and j_1 induce a short exact sequence

$$0 \leftarrow \mathcal{T}^{\prime c} \xleftarrow{i^*} \mathcal{T}^c \xleftarrow{j_!} \mathcal{T}^{\prime \prime c} \leftarrow 0,$$

that is, $\overline{j_!}$ is fully faithful, $\overline{i_*} \circ \overline{j_!} = 0$ and $\overline{i_*}$ induces a triangle equivalence $\mathcal{T}^c / \mathcal{T}''^c \to \mathcal{T}'^c$ up to direct summands.

We give the following useful observation.

Proposition 1.15. Let A, B, C be three homologically smooth and proper dg k-algebras. Assume there is a recollement

$$\mathcal{D}(B) \xrightarrow[i^*]{i^*} \mathcal{D}(A) \xrightarrow[j^*]{j^*} \mathcal{D}(C).$$

Then there is an induced recollement

$$\mathcal{D}^{\mathbf{b}}(B) \xrightarrow[i^{!}]{i_{*}} \mathcal{D}^{\mathbf{b}}(A) \xrightarrow[j_{*}]{j_{*}} \mathcal{D}^{\mathbf{b}}(C).$$

Proof. Note that by Theorem 1.14, the top functors i^* and $j_!$ restrict to the perfect categories, hence to the bounded derived categories (since they coincide by our assumption and Lemma 1.1). Also note that by Lemma 1.4, the middle functors i_* and j^* restrict to \mathcal{D}^{b} , hence to per. Then again by Lemma 1.4, we know that the bottom functors $i^!$ and j_* also restrict to \mathcal{D}^{b} . So the assertion is true.

1.5. Graded marked surfaces and graded gentle algebras. In this subsection we briefly recall some results on marked surfaces. We refer to [HKK, OPS, APS, Op, LP] for more details and examples.

Definition 1.16. A pair (S, M) is called a *marked surface* if

- (1) S is an oriented surface with boundary with connected components $\partial S = \bigcup_{i=1}^{b} \partial_i S$;
- (2) $M = M^{\circ} \cup M^{\bullet} \cup P^{\circ} \cup P^{\bullet}$ is a finite set of *marked points* on *S*. The elements of M° and M^{\bullet} are marked points on ∂S , which will be respectively represented by symbols \circ and \bullet . Each connected component $\partial_i S$ is required to contain at least one marked point of each colour, where in general the points \circ and \bullet are alternating on $\partial_i S$. The elements in $P = P^{\circ} \cup P^{\bullet}$ are marked points in the interior of *S*. We refer to these points as *punctures*, and we will also represent them by the symbols \circ and \bullet respectively.

Note that a marked surface is allowed to have empty boundary. If this happens, then we need both P° and P^{\bullet} to be non-empty. We adopt the convention that the following two special marked surfaces, the one corresponding to a disk without punctures which has only one \circ -point and one \bullet -point on the boundary, and the second corresponding to a sphere with only one \circ -puncture and one \bullet -puncture are 'zero' and if we cut the surface and obtain such a component, then we can disregard it.

Definition 1.17. Let (S, M) be a marked surface. An \circ -arc is a non-contractible curve, with endpoints in $M^{\circ} \cup P^{\circ}$, while an \bullet -arc is a non-contractible curve, with endpoints in $M^{\bullet} \cup P^{\bullet}$.

On the surface, all curves are considered up to homotopy, and all intersections of curves are required to be transversal.

Definition 1.18. Let (S, M) be a marked surface.

- (1) An admissible \circ -dissection Δ of (S, M) is a collection of pairwise non-intersecting (in the interior of S) and pairwise distinct \circ -arcs on the surface such that the arcs in Δ cut the surface into polygons each of which contains exactly one \bullet -point.
- (2) Let q be a common endpoint of two arcs ℓ_i and ℓ_j in an admissible \circ -dissection Δ , an oriented intersection from ℓ_i to ℓ_j at q is an anticlockwise angle locally from ℓ_i to ℓ_j based at q such that the angle is in the interior of the surface. We call an oriented intersection a minimal oriented intersection if it is not a composition of two oriented intersections of arcs from Δ .
- (3) A graded admissible \circ -dissection is a pair (Δ, G) , where Δ is an admissible \circ -dissection and G is a set of gradings, that is, for any minimal oriented intersection α , a grading is the data of an integer $g(\alpha) \in \mathbb{Z}$.
- (4) We call (S, M, Δ, G) a graded marked surface.

We similarly define admissible \bullet -dissections and graded admissible \bullet -dissections.

Remark 1.19. If there is no \circ -puncture in (S, M), then for any graded admissible dissection (Δ, G) , there exists a unique foliation (up to homotopy) on S (with a smooth structure) which induces the grading G, see [LP].

We introduce the following proposition-definition, whose proof is straightforward.

Proposition-Definition 1.20. Let (S, M, Δ, G) be a graded marked surface.

(1) There is a unique (up to homotopy) admissible \bullet -dissection Δ^* on (S, M) such that each arc ℓ^* in Δ^* intersects exactly one arc ℓ of Δ , which we will call the dual \bullet -dissection of Δ .



FIGURE 1. An example of an admissible \circ -dissection Δ of a marked surface and its dual Δ^* , where the arcs in Δ are colored by green and the arcs in Δ^* are colored by red.



FIGURE 2. Dual of minimal oriented intersections α and α^{op} . We set $g(\alpha^{op}) = 1 - g(\alpha)$ for the gradings.

- (2) For any minimal oriented intersection α of Δ from l_i to l_j, there is a unique minimal oriented intersection α^{op} of Δ^{*} from l^{*}_j to l^{*}_i, see Figure 2. Conversely, any minimal oriented intersection of Δ^{*} arises from this way. We call α^{op} the dual minimal oriented intersection of α.
- (3) The dual graded •-dissection (Δ^*, G^*) of (Δ, G) is the dual •-dissection Δ^* of Δ with dual of grading G^* defined by setting $g(\alpha^{op}) = 1 g(\alpha)$, for any minimal oriented intersection α and its dual α^{op} .
- We similarly define the dual graded \circ -dissection of an admissible graded \bullet -dissection.

Note that the dual graded dissection is unique (up to homotopy) and it is an involution, that is $(\Delta^{**}, G^{**}) = (\Delta, G)$. Moreover, the definition of the dual grading is motivated by the algebraic dual of a graded gentle algebra considered in subsection 2.2.

Example 1.21. See Figure 1 for an example of an admissible o-dissection of a marked surface and its dual dissection and Figure 2 for the dual of minimal oriented intersections.

For ease of notation, we will often omit the grading G of (Δ, G) and simply write Δ for a graded surface dissection.

We now recall how to construct a graded gentle algebra from an admissible graded dissection of a marked surface. We begin by recalling the definition of a graded gentle algebra.

Definition 1.22. A graded gentle algebra A is an algebra of the form A = KQ/I where $Q = (Q_0, Q_1)$ is a finite quiver and I an ideal of KQ satisfying the following conditions:

(1) Each vertex in Q_0 is the source of at most two arrows and the target of at most two arrows.



FIGURE 3. The gentle algebra associated to the admissible dissection in Figure 1, where the relations are depicted by dotted lines.

- (2) For each arrow α in Q_1 , there is at most one arrow β such that $0 \neq \alpha\beta \in I$; at most one arrow γ such that $0 \neq \gamma\alpha \in I$; at most one arrow β' such that $\alpha\beta' \notin I$; at most one arrow γ' such that $\gamma'\alpha \notin I$.
- (3) I is generated by paths of length two.
- (4) The grading of A is induced by associating to each arrow α an integer $|\alpha|$.

Note that A might be infinite dimensional. Sometimes such an algebra is also referred to as a graded locally gentle algebra.

We now recall how to associate a graded gentle algebra to a graded admissible dissection. For that, let (Δ, G) be a graded admissible \circ -dissection on a marked surface (S, M). Define the graded algebra $A(\Delta) = kQ(\Delta)/I(\Delta)$ by setting:

- (1) The vertices of $Q(\Delta)$ are given by the arcs in Δ .
- (2) Each minimal oriented intersection α from ℓ_i to ℓ_j gives rise to an arrow from ℓ_i to ℓ_j , which is still denoted by α . Furthermore, $|\alpha| = g(\alpha)$.
- (3) The ideal $I(\Delta)$ is generated by the following relations: whenever ℓ_i and ℓ_j intersect at a marked point which gives rise to an arrow $\alpha : \ell_i \to \ell_j$, and the other end of ℓ_j intersects ℓ_k at a marked point which gives rise to an arrow $\beta : \ell_j \to \ell_k$, then the composition $\alpha\beta$ is a relation.

Then $A(\Delta)$ is a graded gentle algebra. In particular this establishes a bijection between the set of homeomorphism classes of marked surfaces with graded admissible \circ -dissections and the set of isomorphism classes of graded gentle algebras [OPS, PPP]. Similarly, we construct a graded gentle algebra $A(\Delta^*) = kQ(\Delta^*)/I(\Delta^*)$ from a graded admissible \bullet -dissection Δ^* .

Definition 1.23. An \circ -arc is called an *exceptional* \circ -*arc* if γ has no self-intersections, that is, γ does not have any self-intersections neither in the interior of S nor at its endpoints. A \circ -dissection Δ is called *an* \circ -*exceptional dissection* if there is no oriented cycles in the quiver $Q(\Delta)$ associated to Δ . In particular, there is no loop in $Q(\Delta)$ and each arc in Δ is an exceptional \circ -arc.

Example 1.24. See Figure 3 for the gentle algebra $A(\Delta)$ associated to the admissible \circ -dissection Δ depicted in Figure 1. Note that here Δ is not an exceptional dissection since $Q(\Delta)$ has an oriented cycle.

2. Recollements arising from graded quadratic monomial algebras

2.1. Graded quadratic monomial algebras. Let Q be a graded quiver, that is, to every arrow in Q there is an associated integer $|\alpha|$. Furthermore we assume that Q_0 a finite set. Let I be a set of quadratic monomial relations and let $A = kQ/\langle I \rangle$. Then A is a graded k-algebra and we may regard A as a dg k-algebra with zero differential. Let J be a subset of I. In this subsection, we construct a new graded quadratic monomial algebra $A_J = KQ'/\langle I' \rangle$ such that A_J is quasiisomorphic to A by 'killing' the relations in J. Note that if J = I, then I' = 0 and A_J is the dg quiver algebra associated to A sometimes referred to as the cofibrant dg algebra resolution of A. Define the following sets of paths in Q.

$$J_1 := Q_1, \ J_2 := J, \ J_3 := \{\alpha_1 \alpha_2 \alpha_3 \mid \alpha_1 \alpha_2 \in J, \alpha_2 \alpha_3 \in J\},\$$

and, in general, we define

$$J_n := \{ \alpha_1 \cdots \alpha_n \mid \alpha_i \alpha_{i+1} \in J, \forall i = 1, 2, \dots, n-1 \}.$$

Note that J_n can be an empty set.

Definition 2.1. We define the partial cofibrant dg algebra resolution $A_J = kQ'/\langle I' \rangle$ as follows.

- (1) $Q'_0 = Q_0.$
- (2) The arrows of Q' are given by

$$Q'_1 = \{ [\alpha_1 \dots \alpha_n] : s(\alpha_1) \to t(\alpha_n) \mid \alpha_1 \dots \alpha_n \in J_n, n \ge 1 \},\$$

where $|[\alpha_1 ... \alpha_n]| := \sum_{i=1}^n |\alpha_i| - n + 1.$

(3) The quadratic relations are given by

$$I' := \{ [\alpha_1 \dots \alpha_n] [\beta_1 \dots \beta_m] \mid \alpha_n \beta_1 \in I \setminus J, n, m \ge 1 \}.$$

(4) The differential d' is defined as $d'([\alpha]) = 0$ for any $\alpha \in Q_1$ and

$$d'([\alpha_1 \dots \alpha_n]) := \sum_{i=1}^{n-1} (-1)^{|[\alpha_1 \dots \alpha_i]|} [\alpha_1 \dots \alpha_i] [\alpha_{i+1} \dots \alpha_n],$$

for any $\alpha_1 \ldots \alpha_n \in J_n$ and $n \ge 2$.

One can check that $d'^2 = 0$, so A_J is a well-defined dg k-algebra.

- **Remark 2.2.** (1) If J = I, then $I' = \emptyset$ and in this case, $A_J = (kQ', d')$ is a dg quiver algebra, which can be regarded as a cofibrant dg algebra resolution of A. Moreover, any algebra admits a dg algebra resolution (see for example [Opm]) and for a monomial algebra, there is an explicit minimal choice given in [T], which coincides with our construction above for the case J = I.
- (2) If $J = \emptyset$, then $J_n = \emptyset$ for any $n \ge 2$ and moreover, $A_J = A$.

We illustrate the construction above in an example.

Example 2.3. Let $Q: 1 \xrightarrow{\alpha} 2 \xrightarrow{\beta} 3 \xrightarrow{\delta} 4 \xrightarrow{\epsilon} 5$ be the graded quiver with zero grading and where $I = \{\alpha\beta, \beta\delta, \delta\epsilon\}$. Let $J = \{\alpha\beta, \beta\delta\}$. Then Q' is



where the grading is given by $|[\alpha\beta]| = -1 = |[\beta\delta]|$, $|[\alpha\beta\delta]| = -2$ and the differential is given by $d'([\alpha\beta]) = [\alpha][\beta], d'([\beta\delta]) = [\beta][\delta]$ and $d'([\alpha\beta\delta]) = [\alpha][\beta\delta] - [\alpha\beta][\delta]$ and where $I' = \{[\delta][\epsilon], [\beta\delta][\epsilon], [\alpha\beta\delta][\epsilon]\}$.

Note that in some cases A_J might be a quiver with infinitely many arrows. We give an example of such a case.

Example 2.4. Let Q be the graded quiver with zero grading $1 \bigcap \alpha$ and let $I = \{\alpha^2\}$. Let J = I, then Q' is

10



where $|[\alpha^n]| = 1 - n$ and $d'(|\alpha^n|) = \sum_{i=1}^{n-1} (-1)^{1-i} [\alpha^i] [\alpha^{n-i}].$

Our main result in this subsection is the following.

Theorem 2.5. There is a natural quasi-isomorphism $\phi : A_J \to A$ of dg k-algebras.

Proof. We first define a map $\phi : Q'_0 \cup Q'_1 \to kQ$ by $\phi(e_j) := e_j$, $\phi([\alpha]) := \alpha$ for any $\alpha \in Q_1$, $\phi([\alpha_1 \cdots \alpha_n]) := 0$ for $n \ge 2$. Then ϕ can be extended to an epimorphism $\phi : kQ' \to kQ$ of graded k-algebras. Note that we have $\phi(I') \subset I$ and $\phi(d'([\alpha_1 \cdots \alpha_n])) \in I$, for any $n \ge 1$. Thus ϕ also induces an epimorphism $\phi : A_J \to A$ of dg algebras.

Next we show that ϕ is a quasi-isomorphism of k-complexes. Let $P := \ker \phi$. It is enough to show P is acyclic. Define a map $N : Q_1 \times Q_1 \to \{0, 1\}$ by

$$N(\alpha, \beta) := \begin{cases} 1 & \text{if } \alpha\beta \in J; \\ 0 & \text{if } \alpha\beta \notin J. \end{cases}$$

We construct a morphism $\psi: P \to P[-1]$ of graded k-modules by

$$\psi([\alpha_{1}\cdots\alpha_{t_{1}}][\alpha_{t_{1}+1}\cdots\alpha_{t_{2}}]\cdots[\alpha_{t_{s}+1}\cdots\alpha_{n}])$$

:= $\frac{1}{m}\sum_{i=1}^{s}(-1)^{\sum_{j=1}^{i}|[\alpha_{t_{j-1}+1}\cdots\alpha_{t_{j}}]|}[\alpha_{1}\cdots\alpha_{t_{1}}][\alpha_{t_{1}+1}\cdots\alpha_{t_{2}}]\cdots[\alpha_{t_{i-1}+1}\cdots\alpha_{t_{i}}\alpha_{t_{i}+1}\cdots\alpha_{t_{i+1}}]\cdots[\alpha_{t_{s}+1}\cdots\alpha_{n}],$

where $m = n - s - 1 + \sum_{i=1}^{s} N(\alpha_{t_i}, \alpha_{t_i+1})$ and where we set $t_0 = 0$. If $\alpha_{t_i} \alpha_{t_i+1} \notin J$, we set above term to be zero.

We claim that $d'\psi + \psi d' = \text{Id}$. Then P is contractible, and hence acyclic. First note that any element $l_1 + \cdots + l_t$ of kQ' is in P if and only each l_i is in P. So we only need to check $d\psi + \psi d' = \text{Id}$ for paths in P.

Given $l = [\alpha_1 \cdots \alpha_{t_1}][\alpha_{t_1+1} \cdots \alpha_{t_2}] \cdots [\alpha_{t_s+1} \cdots \alpha_n] \in P$, l appears (n-s-1)/m times in $\psi d'(l)$ and appears $(\sum_{i=1}^s N(\alpha_{t_i}, \alpha_{t_i+1}))/m$ times in $d'\psi(l)$. For any other term l', if it appears in $\psi d'(l)$, one can check directly that the term -l' appears in $d'\psi(l)$. So $d'\psi(l) + \psi d'(l) = l$. \Box

Remark 2.6. Note that in the proof of Theorem 2.5 we need the assumption that the characteristic of k is zero.

Next we study the graded quadratic dual of A. Let $I^{\perp} := \{\beta^{\text{op}} \alpha^{\text{op}} \mid \alpha\beta \notin I\}$. Let $A^! = kQ^{\text{op}}/I^{\perp}$ such that $|\alpha^{\text{op}}| := 1 - |\alpha|$ for any arrow $\alpha^{\text{op}} \in Q_1^{\text{op}}$. It is obvious that $A^!$ is also a graded quadratic monomial algebra and $(A^!)^! = A$. Let S_i be the simple dg A-module corresponding to the vertex $i \in Q_0$ and let $S := \bigoplus_{i \in Q_0} S_i$. The following observation is known and we add a proof for the convenience of the reader.

Proposition 2.7. $A^!$ is isomorphic to the graded algebra $\bigoplus_{t \in \mathbb{Z}} \operatorname{Hom}_{\mathcal{D}^{\mathrm{b}}(A)}(S, S[t])$.

Proof. To calculate $\operatorname{Hom}_{\mathcal{D}^{\mathsf{b}}(A)}(S, S[t])$, we first give a projective resolution of S. Let $M_0 := \bigoplus_{i \in Q_0} Ae_i \otimes e_i A$ and

$$M_m := \bigoplus_{\alpha_1 \cdots \alpha_m \in \mathcal{F}_m} Ae_{s(\alpha_1)} \otimes e_{t(\alpha_m)} A(|\alpha_1| + \cdots + |\alpha_m|),$$

where $\mathcal{F}_m := \{\alpha_1 \cdots \alpha_m \text{ paths of } kQ \text{ such that } \alpha_i \alpha_{i+1} \in I\}$ and we denote by $(|\alpha_1| + \cdots + |\alpha_m|)$ the degree shift of $Ae_{s(\alpha_1)} \otimes e_{t(\alpha_m)}A$ as graded A-A-bimodule. There is an exact sequence of graded A-A-bimodules

$$\dots \to M_2 \xrightarrow{d_2} M_1 \xrightarrow{d_1} M_0 \xrightarrow{d_0} A \to 0$$
(2.1)

which gives a minimal A-A-bimodule resolution of A (see for example [B, Theorem 4.1]), where $d_m(a \otimes b) := a\alpha_1 \otimes b + (-1)^m a \otimes \alpha_n b$ and $d_0(a \otimes b) := ab$. Since (2.1) is contractible, applying $S_i \otimes_A$? to (2.1), we have a minimal resolution of S_i as graded A-module

$$P: \dots \to \bigoplus_{\alpha_1 \alpha_2 \in \mathcal{F}_2 \atop s(\alpha_1)=i} (e_{t(\alpha_2)}A(|\alpha_1|+|\alpha_2|)) \xrightarrow{\partial_2} \bigoplus_{\alpha \in Q_1 \atop s(\alpha)=i} (e_{t(\alpha)}A(|\alpha|)) \xrightarrow{\partial_1} e_i A \to 0,$$
(2.2)

whose total complex Tot(P) gives a projective resolution of S_i as dg A-module, where $\partial_m(x) =$ $(-1)^m x \alpha_m$. Then any non-zero morphism from S_i to $S_j[l]$ in $\mathcal{D}^{\mathbf{b}}(A)$ is induced by some non-zero map from top(P^m) to $S_i[l]$. Moreover, we see that $\operatorname{Hom}_{\mathcal{D}^b(A)}(S_i, S_i[l])$ has a k-basis

$$\{\alpha_1 \cdots \alpha_p \in \mathcal{F}_p \text{ such that } s(\alpha_1) = i, t(\alpha_p) = j \text{ and } p = l + |\alpha_1| + \cdots + |\alpha_p|\}.$$

In particular, any arrow $\alpha \in Q_1$ gives a non-zero morphism $f_\alpha : S_{s(\alpha)} \to S_{t(\alpha)}[1-|\alpha|]$. By the construction of P, we have $(f_{\beta}[1 - |\alpha|]) \circ f_{\alpha} \neq 0$ if and only if $0 \neq \alpha\beta \in I$. Thus the map $\phi: A^! \to \bigoplus_{t \in \mathbb{Z}} \operatorname{Hom}_{\mathcal{D}^{\mathrm{b}}(A)}(S, S[t])$ sending e_i to the identity morphism in $\operatorname{End}_{\mathcal{D}^{\mathrm{b}}(A)}(S_i)$ and α^{op} to f_{α} gives an isomorphism of dg algebras.

2.2. Recollements induced by idempotents of graded quadratic monomial algebras. Let A = kQ/I be a graded quadratic monomial algebra. Let e be the sum of m vertex idempotents. Without loss of generality, we may write $e = e_1 + \cdots + e_m$. Let J_e be the subset of I consisting of all the relations go through vertices $1, \ldots, m$.

Definition 2.8. We define a new algebra $A_e := kQ_e/\langle I_e \rangle$ as follows

- (1) The set of vertices of Q_e is $Q_0 \setminus \{1, 2, \ldots, m\}$.
- (2) The arrows of Q_e are of the form $[\alpha_1 \cdots \alpha_s] : s(\alpha_1) \to t(\alpha_s)$ with $\alpha_1 \cdots \alpha_s \in J_s$ and $s(\alpha_1), t(\alpha_s) \notin J_s$ $\{1,2,\cdots,m\}.$
- (3) $|[\alpha_1 \cdots \alpha_s]| := \sum_i^s |\alpha_i| s + 1.$ (4) $I_e := \{ [\alpha_1 \cdots \alpha_s] [\beta_1 \cdots \beta_t] \mid \alpha_s \beta_1 \in I \}.$

(5) The differential is 0.

Example 2.9. In case A is a graded gentle algebra, the algebra $A_e = kQ_e/I_e$ in Definition 2.8 is again a graded gentle algebra. In case $e = e_{\ell}$ is an idempotent corresponding to a single vertex ℓ , we obtain the following explicit description.

- (1) The vertex set of Q_e is given by $Q_0 \setminus \{\ell\}$;
- (2) The arrow set of Q_e is the union

$$\{ [\alpha] = \ell_i \xrightarrow{\alpha} \ell_j \mid \ell_i \neq \ell \text{ and } \ell_j \neq \ell \} \ \sqcup \ \{ [\alpha_1 \alpha_2] = \ell_i \xrightarrow{\alpha_1} \ell \xrightarrow{\alpha_2} \ell_j \mid \alpha_1 \alpha_2 \in I \text{ and } \ell_i \neq \ell \text{ and } \ell_j \neq \ell \}$$

$$\sqcup \{ [\alpha_1 \alpha_2 \alpha_3] = \ell_i \xrightarrow{\alpha_1} \ell \xrightarrow{\alpha_2} \ell \xrightarrow{\alpha_3} \ell_j \mid \alpha_1 \alpha_2, \alpha_2 \alpha_3 \in I \};$$

(3) $|[\alpha]| = |\alpha|, |[\alpha_1\alpha_2]| = |\alpha_1| + |\alpha_2| - 1 \text{ and } |[\alpha_1\alpha_2\alpha_3]| = |\alpha_1| + |\alpha_2| + |\alpha_3| - 2;$

(4) The ideal I_{ℓ} is generated by

$$\{ [\alpha_1] [\alpha_2] \mid \alpha_1 \alpha_2 \in I \} \ \sqcup \ \{ [\alpha_0] [\alpha_1 \alpha_2] \mid \alpha_0 \alpha_1 \in I \} \ \sqcup \ \{ [\alpha_1 \alpha_2] [\alpha_3] \mid \alpha_2 \alpha_3 \in I \}$$

 $\sqcup \{ [\alpha_0] [\alpha_1 \alpha_2 \alpha_3] \mid \alpha_0 \alpha_1 \in I \} \sqcup \{ [\alpha_1 \alpha_2 \alpha_3] [\alpha_4] \mid \alpha_3 \alpha_4 \in I \} \sqcup \{ [\alpha_0 \alpha_1] [\alpha_2 \alpha_3] \mid \alpha_1 \alpha_2 \in I \}.$

We show

Theorem 2.10. There is a recollement

$$\mathcal{D}(A_e) \xrightarrow{i^*} \mathcal{D}(A) \xrightarrow{j^* = ? \otimes_{eAe}^{\mathbf{L}} A e} \mathcal{D}(eAe) , \qquad (2.3)$$

such that i^* sends e_jA to e_jA_e and i_* sends $top(e_jA_e)$ to $top(e_jA)$ for any $j \neq 1, 2, \ldots, m$.

Remark 2.11. Note that if $J_e = I$, then (2.3) is given by [KaY2, Theorem 7.1]. One may also use Drinfeld's construction of dg quotient [D] to give a dg algebra C such that $\mathcal{D}(C)$ fits the left side of (2.3). We point out that C is quasi-isomorphic to A_e , but in the case that A is gentle, C is not gentle anymore and the structure of C may be complicated.

Recall that we constructed a dg algebra A_{J_e} from J_e (see Definition 2.1). To simplify the notation, we write $B = A_{J_e}$. The following observation is important for us.

Lemma 2.12. We have $A_e = B/BeB$ and the natural morphism $B \to A_e$ is a homological epimorphism of dg k-algebras. Moreover, there is a recollement

$$\mathcal{D}(A_e) \xrightarrow{\widetilde{i_*} = ? \otimes_B^{\mathbf{L}} A_e} \mathcal{D}(B) \xrightarrow{\widetilde{j_*} = ? \otimes_{B_e}^{\mathbf{L}} B_e} \mathcal{D}(eBe)$$

Proof. It suffices to show $A_e \otimes_B^{\mathbf{L}} A_e \cong A_e$ in $\mathcal{D}(A_e)$ by Lemma 1.9 (2). Notice that we have the following canonical triangle in $\mathcal{D}(B \otimes_k B^{\mathrm{op}})$

$$BeB \to B \xrightarrow{p} A_e \to BeB[1].$$

Applying $? \otimes_B^{\mathbf{L}} A_e$ to this triangle, we get a triangle in $\mathcal{D}(A_e)$

$$BeB \otimes_B^{\mathbf{L}} A_e \to A_e \to A_e \otimes_B^{\mathbf{L}} A_e \to BeB \otimes_B^{\mathbf{L}} A_e[1].$$

We claim that $BeB \otimes_{\mathbf{B}}^{\mathbf{L}} A_e = 0$, then $A_e \cong A_e \otimes_{\mathbf{B}}^{\mathbf{L}} A_e$ in $\mathcal{D}(A_e)$ holds. Note that $BeB = \bigoplus_{s \in Q_0} e_s BeB$ as right dg *B*-module. Let L_s be the set of paths from *s* to *i*, which are not in *I*. By our construction of Q', there is no relation at *i*, so we have $le_iB \cong e_iB$ as right dg *B*-module for any $l \in L_s$. So $BeB = \bigoplus_{s \in Q_0} \bigoplus_{l \in L_s} le_iB$ is a projective module. Then $BeB \otimes_{\mathbf{B}}^{\mathbf{L}} A_e = BeB \otimes_{\mathbf{B}} B/BeB = 0$. Then the claims follows. So the assertion is true.

Now we show Theorem 2.10.

Proof of Theorem 2.10. By Theorem 2.5, there is a quasi-isomorphism $\phi : B \to A$, it induces a quasi-isomorphism $eBe \to eAe$. Then we have triangle equivalences $\mathcal{D}(A) \xrightarrow{\phi^*}{\simeq} \mathcal{D}(B)$ and $\mathcal{D}(e_iAe_i) \xrightarrow{\phi^*}{\simeq} \mathcal{D}(e_iBe_i)$. Consider the recollement constructed in Lemma 2.12. Let $i^* := \tilde{i^*}\phi^*$, $i_* := (\phi^*)^{-1}\tilde{i_*}$ and $i^! := \tilde{i^!}\phi^*$. Then we get the recollement (2.3). Since ϕ induces a quasiisomorphism $e_jB \to e_jA$ and $\tilde{i_*}(\operatorname{top}(e_jA_e)) = \operatorname{top}(e_jB)$, we have

$$i^*(e_jA) = \widetilde{i^*}\phi^*(e_jA) \cong \widetilde{i^*}(e_jB) = e_jB \otimes_B^{\mathbf{L}} A_e = e_jA_e$$

and

$$i_*(\operatorname{top}(e_j A_e)) = (\phi^*)^{-1} i_*(\operatorname{top}(e_j A_e)) = (\phi^*)^{-1}(\operatorname{top}(e_j B)) = \operatorname{top}(e_j A)$$

for any $j \neq 1, 2, \ldots, m$. So the assertion is true.

Example 2.13. Let Q be the graded quiver $1 \xrightarrow{\alpha} 2 \xrightarrow{\gamma} 3$. Let $I := \{\alpha\beta, \beta^2, \beta\gamma\}$ and $A = kQ/\langle I \rangle$. Let $e = e_2$. Then $J_e = I$ and A_e is given by the following 'infinite Kronecker quiver'

$$1 \xrightarrow[[\alpha\beta^{3}\gamma]]{[\alpha\beta^{2}\gamma]} 3$$

where $|[\alpha\beta^n\gamma]| := |\alpha| + n|\beta| + |\gamma| - n - 1.$

We give a useful observation.

Proposition 2.14. Let e = e' + e''. Then we have isomorphisms $(A_{e'})_{e''} \cong A_e \cong (A_{e''})_{e'}$ of dg algebras.

Proof. We only show that there is an isomorphism $(A_{e'})_{e''} \cong A_e$ of dg k-algebras. We may assume $e' = e_1 + \cdots + e_l$ and $e'' = e_{l+1} + \cdots + e_m$ for $1 \leq l < m$. First note that the induced quivers Q_e and $(Q_{e'})_{e''}$ are isomorphic. In fact, they have same sets of vertices given by $Q_0 \setminus \{1, 2, \ldots, m\}$. Moreover, any arrow in A_e has the form $[\alpha_1 \cdots \alpha_m]$ such that $\alpha_i \alpha_{i+1} \in J_e$ for any $1 \leq i \leq m-1$. Set $p_0 = 0$, since $J_e = J_{e'} \sqcup J_{e''}$, we may write

$$[\alpha_1 \cdots \alpha_m] = [(\alpha_{p_0+1} \cdots \alpha_{p_1})(\alpha_{p_1+1} \cdots \alpha_{p_2}) \cdots (\alpha_{p_l+1} \cdots \alpha_m)], \qquad (2.4)$$

where $\alpha_{p_j+q(j)}\alpha_{p_j+q(j)+1} \in J_{e'}$ and $\alpha_{p_{j+1}}\alpha_{p_{j+1}+1} \in J_{e''}$ for any $0 \leq j \leq l-1$ and $1 \leq q(j) \leq p_{j+1}-1$. It is clear that (2.4) gives rise to a bijection between the set of arrows in Q_e and that in $(Q_{e'})_{e''}$. So Q_e and $(Q_{e'})_{e''}$ are isomorphic. Also it is easy to check that I_e and $(I_{e'})_{e''}$ are also isomorphic by Definition 2.8(4). Thus A_e is isomorphic to $(A_{e'})_{e''}$.

We can describe A_e by using $A^!$.

We have the following useful observations.

Proposition 2.15. There are isomorphisms of dg k-algebras.

(1) $A_e \cong ((1-e)A^!(1-e))^!.$ (2) $eAe \cong ((A^!)_{1-e})^!.$

Proof. We only show (1), since (2) follows directly from (1) and $(A^{!})^{!} = A$.

First note that both algebras have the same vertices. Assume $Q_0 = 1, 2, \ldots, n$. Then $1 - e = e_{m+1} + \cdots + e_n$. Any arrow l^{op} in $(1 - e)A^!(1 - e)$ is given by a path $l^{\text{op}} = \alpha_t^{\text{op}} \cdots \alpha_1^{\text{op}}$ with degree $(t - \sum_{i=1}^t |\alpha_i|, 1)$ such that $s(\alpha_t^{\text{op}}), t(\alpha_1^{\text{op}}) \in \{m + 1, \ldots, n\}$ and $\alpha_j^{\text{op}} \alpha_{j-1}^{\text{op}} \notin J_e^{\perp}$ for any $2 \leq j \leq t$. Then $l = \alpha_1 \cdots \alpha_t$ gives rise to an arrow of $((1 - e)A^!(1 - e))^!$ with deg $l = (\sum_{i=1}^t |\alpha_i| - t + 1, 1)$. Moreover, $s(\alpha_1), t(\alpha_t) \notin \{1, \ldots, m\}$ and $\alpha_j \alpha_{j+1} \in J_e$ for any $1 \leq j \leq t - 1$. Then l also gives rise to an arrow of A_e .

Let $l' = \beta_1 \cdots \beta_s$ be another arrow. Then ll' = 0 in $((1-e)A^!(1-e))^!$ if and only if $l'^{\text{op}}l^{\text{op}} \neq 0$ in $(1-e)A^!(1-e)$, if and only if $\alpha_t\beta_1 \in I \setminus J_e$, if and only if ll' = 0 in A_e .

So the two algebras have same arrows (with same grading) and same relations. Thus they are isomorphic. $\hfill \Box$

3. Recollements arising from cutting surfaces

In this section, we apply Theorem 2.10 to the class of graded gentle algebras. In particular, we show that both the left and right side of the recollement in Theorem 2.10 correspond to cutting the surface associated to the graded gentle algebra along arcs of the corresponding graded surface dissection and its dual, respectively.

More precisely, let $A(\Delta)$ be a graded gentle algebra associated to the graded marked surface (S, M, Δ) and let L be a subset of the arcs in the surface dissection Δ . In the following we define the graded cut marked surface (S_L, M_L, Δ_L) obtained from (S, M, Δ) by cutting along the arcs in L. We can also consider the graded marked surface obtained by cutting along arcs in Δ^* , for the dual graded surface dissection Δ^* of (S, M, Δ) .

We begin by stating the main result of this section as well as some consequences before introducing the necessary definitions and proving the stated results.

Theorem 3.1. Let (S, M, Δ) be a graded marked surface with associated graded gentle algebra $A(\Delta)$. Let L be a subset of Δ . Then we have the following recollement

$$\mathcal{D}(A(\Delta_L)) \longrightarrow \mathcal{D}(A(\Delta)) \longrightarrow \mathcal{D}(A(L)), \tag{3.1}$$

where Δ_L is the induced graded surface dissection obtained by cutting (S, M, Δ) along L and where L is realized as the graded surface dissection of the surface $(S_{\Delta^* \setminus L^*}, M_{\Delta^* \setminus L^*})$ which is obtained by cutting (S, M, Δ^*) along $\Delta^* \setminus L^*$ and dualising the induced surface dissection L^* .

We note that in [A, Section 1.3] a similar recollement to the above has been suggested by considering a cofibrant dg algebra resolution of $A(\Delta)$ which is also related to surface cuts.

We now give an example of Theorem 3.1.

Example 3.2. Consider the algebra $1 \xrightarrow{\alpha} 2 \xrightarrow{\beta} 3 \xrightarrow{\gamma} 4 \xrightarrow{\delta} 5$ with $\alpha\beta, \gamma\delta \in I$ corresponding to the surface dissection $\Delta = \{\ell_1, \ell_2, \ell_3, \ell_4, \ell_5\}$ in Figure 4 and with $L = \{\ell_2, \ell_4\}$. Suppose further that $|\alpha| = a, |\beta| = b, |\gamma| = c$ and $|\delta| = d$. Then the surfaces corresponding the left and the right side of the recollement in Theorem 3.1 are given in Figure 4. We note that the arrows $[\alpha\beta] : \overline{\ell_1} \to \overline{\ell_3}$ and $[\gamma\delta] : \overline{\ell_3} \to \overline{\ell_5}$ in $A(\Delta_L)$ have degrees $|[\alpha\beta]| = a + b - 1$ and $|[\gamma\delta]| = c + d - 1$, respectively. The induced arrow $\beta\gamma : \ell_2 \to \ell_4$ in A(L) has degree b + c.

- **Remark 3.3.** (1) Given a subset L of arcs of Δ , we can consider L as what is called a ribbon graph which is graded with a grading induced by the grading of Δ . From L we can construct a graded marked surface dissection (S(L), M(L), L), see for example [HKK, LP].
- (2) It is straightforward to see that the graded algebra A(L) associated to the graded sub-ribbon graph L of Δ is isomorphic to eAe, where $e \in A$ is the idempotent corresponding to L. Hence by Proposition 2.15 and Theorem 3.1, A(L) is isomorphic to $(A(\Delta^*_{\Delta^* \setminus L^*}))!$.
- (3) It is easy to see that L gives rise to an admissible dissection of $(S_{\Delta^* \setminus L^*}, M_{\Delta^* \setminus L^*})$ and the algebra associated to the surface dissection L is exactly the algebra A(L) associated to the graded ribbon graph L in (2).

Remark 3.3 implies the following.

Corollary 3.4. The marked surface (S(L), M(L)) obtained from the subribbon graph L of Δ is homeomorphic to the marked surface $(S_{\Delta^* \setminus L^*}, M_{\Delta^* \setminus L^*})$.

We now give the precise definitions of the terms above and then devote the rest of the Section to proving the stated results. We begin by defining the surface obtained from cutting arcs on a marked surface with a graded admissible dissection.

Definition 3.5. [APS, CS1] Let ℓ be an \circ -arc without self-intersection in the interior of a marked surface (S, M). Let p and q be the endpoints of ℓ which may coincide. We define the *cut surface* (S_{ℓ}, M_{ℓ}) to be the marked surface obtained from (S, M) by cutting and contracting along ℓ . In particular, we set $M_{\ell}^{\bullet} = M^{\bullet}, P_{\ell}^{\bullet} = P^{\bullet}$, and $M_{\ell}^{\circ} \cup P_{\ell}^{\circ} = ((M^{\circ} \cup P^{\circ}) \setminus \{p, q\}) \cup \{\overline{p}, \overline{q}\}$, where \overline{p} and \overline{q} are new \circ -marked points (on the boundary or in the interior) of S_{ℓ} obtained from cutting and contracting ℓ .

We similarly define the cut surface (S_{ℓ^*}, M_{ℓ^*}) of (S, M) by cutting along an \bullet -arc ℓ^* which has no self-intersection in the interior of the surface.



FIGURE 4. Cut surfaces induced by $L = \{\ell_2, \ell_4\}$ in Example 3.2. The surface in (B) is obtained by cutting the surface in (A) along L and the surface in (C) is obtained by cutting the surface in (A) along $\{\ell_1^*, \ell_3^*, \ell_5^*\}$.

Let ℓ' be an \circ -arc on (S, M), denote by $\overline{\ell'}$ the *induced arc* of ℓ' on (S_{ℓ}, M_{ℓ}) , see Figure 5 for an example. Note that if ℓ' intersects with ℓ , then $\overline{\ell'}$ is empty.

Example 3.6. Figure 5 gives an example of a cut surface and induced arcs. Namely, we cut an annulus along an arc ℓ connecting the two boundary components and then contract along the cut. From this we see that the cut surface is a disk.



FIGURE 5. An example of a cut surface, where the labels \overline{p} and \overline{q} are allowed to be exchanged. The arcs $\overline{\ell'}$ and $\overline{\ell''}$ are respectively the induced arcs of ℓ' and ℓ'' .

Note that the cut surface may not be connected, even if the original surface is. Recall that by our convention if the cut surface has a component corresponding to a sphere with a unique o-puncture and a unique o-puncture or an unpunctured disk with only one o-marked point and one o-marked point on the boundary, then this corresponds to the zero gentle algebra and we can ignore this component.

Proposition 3.7. Let (S, M, Δ, G) be a graded marked surface with associated graded gentle algebra $A(\Delta)$. Let ℓ be an arc in Δ and denote the associated idempotent in A also by ℓ . Denote by $\Delta_{\ell} = \{\overline{\ell_{\ell}}, \ell_{\ell} \in \Delta\}$ the set of arcs on the cut surface (S_{ℓ}, M_{ℓ}) induced from the arcs in Δ .

- (1) Then Δ_{ℓ} is an admissible dissection on (S_{ℓ}, M_{ℓ}) .
- (2) The algebra $A(\Delta_{\ell})$ associated to Δ_{ℓ} is isomorphic to $A(\Delta)_{\ell}$ as ungraded algebras, where $A(\Delta)_{\ell}$ is the (graded) algebra introduced in Definition 2.8.

With the notation of the above proposition we make the following definitions.

Definition 3.8. We call Δ_{ℓ} the *induced admissible* \circ -dissection of Δ on (S_{ℓ}, M_{ℓ}) and we call $(S_{\ell}, M_{\ell}, \Delta_{\ell}, G_{\ell})$ the graded cut surface of (S, M, Δ, G) at ℓ , where G_{ℓ} is the grading on Δ_{ℓ} such that $A(\Delta_{\ell}) \cong A(\Delta)_{\ell}$ as graded algebras.

Proof of Proposition 3.7. (1) Follows from [APS].

(2) Let (Q, I) be such that $A(\Delta) = kQ/I$ and let $A(\Delta_{\ell}) = kQ'/I'$. Since $A(\Delta)$ is a gentle algebra, $A(\Delta)_{\ell} = kQ_{\ell}/I_{\ell}$ is as defined in Example 2.9.

We now show that $kQ_{\ell}/I_{\ell} \cong kQ'/I'$. It follows from (1) that there is a canonical bijection from the vertex set of Q_{ℓ} to the vertex set of Q', which sends an arc $\ell_i \in \Delta \setminus \{\ell\}$ to an arc $\overline{\ell_i} \in \Delta_{\ell}$.

Under this bijection, we will further show that the set of arrows of Q_{ℓ} coincides with the set of arrows of Q'. For this, let $\overline{\alpha} : \overline{\ell_i} \to \overline{\ell_j}$ be an arrow of Q' corresponding to an oriented intersection at a common endpoint $\overline{\varrho}$ of $\overline{\ell_i}$ and $\overline{\ell_j}$. We claim that $\overline{\alpha}$ must correspond to an arrow of the form as described in Example 2.9 (2).

Suppose $\overline{\varrho} \notin \{\overline{p}, \overline{q}\}\$ where if ℓ has endpoints p and q, \overline{p} and \overline{q} are the induced marked points in (S_{ℓ}, M_{ℓ}) . Then $\overline{\varrho}$ can be lifted to a marked point on S, which we denoted by ϱ . Note that since the local configuration of ϱ does not change when cutting the surface, $\overline{\alpha}$ can be lifted to a minimal oriented intersection in (S, M, Δ) from ℓ_i to ℓ_j at ϱ , which gives rise to an arrow $\alpha : \ell_i \to \ell_j$ in Q. Thus in this case, $\overline{\alpha}$ corresponds to an arrow $[\alpha]$ in Q_{ℓ} , which belongs to the first subset as stated in Example 2.9 (2).

If $\overline{\varrho} \in \{\overline{p}, \overline{q}\}$, then both ℓ_i and ℓ_j share an endpoint with ℓ , and up to permutating the endpoints p and q of ℓ , we have the following three possibilities of lifting $\overline{\ell_i}$ and $\overline{\ell_j}$ to arcs ℓ_i and ℓ_j on (S, M).

(a) Both ℓ_i and ℓ_j share the same endpoint (for example p) with ℓ such that ℓ is not between ℓ_i and ℓ_j , see the following leftmost and middle pictures. In this case, note that the minimal oriented intersection $\overline{\alpha}$ is induced from a minimal oriented intersection α on (S, M, Δ) , which gives rise to an arrow of Q_{ℓ} . Thus in this case, $\overline{\alpha}$ also corresponds to an arrow $[\alpha]$ of Q_{ℓ} in the first subset as stated in Example 2.9 (2).



(b) Both ℓ_i and ℓ_j share the same endpoint p with ℓ such that ℓ is between ℓ_i and ℓ_j . Note that in this case, since $\overline{\alpha}$ arises from the intersection $\overline{\varrho}$, the other endpoint q of ℓ must be a \circ -puncture with ℓ the unique arc in Δ adjacent to it. That is, the local configuration must be as in the left picture of the following figure, with arrows $\alpha_1 : \ell_i \to \ell$, $\alpha_3 : \ell \to \ell_j$ and $\alpha_2 : \ell \to \ell$ in Q arising from minimal oriented intersections at p and q respectively. Then $\overline{\alpha}$ corresponds to an arrow $[\alpha_1 \alpha_2 \alpha_3]$ of Q_ℓ as in the third subset of Example 2.9 (2).



(c) The arcs ℓ_i and ℓ_j share different endpoints with ℓ , see the left picture below, where $\alpha_1 : \ell_i \to \ell$ and $\alpha_2 : \ell \to \ell_j$ are arrows in Q arising from minimal oriented intersections of q and p respectively. In this case we have $[\alpha_1 \alpha_2]$ as in the right picture, which belongs to the second subset in Example 2.9 (2).



We have thus established a bijection between the arrow set of Q_{ℓ} and the arrow set of Q', which induces an isomorphism of the quivers Q_{ℓ} and Q'. Furthermore one can check using the definition of a gentle algebra that the ideal I' of the algebra arising from the admissible dissection Δ_{ℓ} is generated exactly by the relations in I_{ℓ} described in Example 2.9 (4).

Lemma 3.9. Let (S, M, Δ, G) be a graded marked surface. Let ℓ and ℓ' be two arcs in Δ . Then there is a homeomorphism between $((S_{\ell})_{\ell'}, (M_{\ell})_{\ell'})$ and $((S_{\ell'})_{\ell}, (M_{\ell'})_{\ell})$, such that $((\Delta_{\ell})_{\ell'}, (G_{\ell})_{\ell'})$ and $((\Delta_{\ell'})_{\ell}, (G_{\ell'})_{\ell})$ coincide under this homeomorphism.

Proof. By Proposition 2.14 and Proposition 3.10, we have an isomorphism of graded gentle algebras associated to graded admissible \circ -dissections $((\Delta_{\ell})_{\ell'}, (G_{\ell})_{\ell'})$ and $((\Delta_{\ell'})_{\ell}, (G_{\ell'})_{\ell})$. Then the statement follows from the bijection between marked surfaces with graded admissible dissections up to homeomorphism and isomorphism classes of graded gentle algebras.

We now define the graded cut surface $(S_L, M_L, \Delta_L, G_L)$ of a graded marked surface (S, M, Δ, G) at a subset $L \subset \Delta$ to be the graded marked surface obtained by successively cutting (S, M, Δ, G) along the arcs in L. Note that by Lemma 3.9, the graded cut surface is independent up to homeomorphism of the order in which we cut the arcs in L. Thus it is well-defined. Similarly, one can cut along arcs $D^* \subset \Delta^*$ to obtain a graded \bullet -surface dissection $\Delta_{D^*}^*$ of the graded cut surface (S_{D^*}, M_{D^*}) .

The following observation naturally generalizes Proposition 3.7.

Proposition 3.10. Let (Δ, G) be a graded \circ -admissible dissection on (S, M) with associated graded gentle algebra A. Let L be a subset of Δ corresponding to an idempotent $L \in A$. Then we have isomorphism of graded gentle algebras $A(\Delta_L) \cong A(\Delta)_L$.

We now prove Theorem 3.1

Proof of Theorem 3.1. The left hand side of the theorem follows from Theorem 2.10 and Proposition 3.10. For the right hand side, let (S, M) be a marked surface with a graded admissible \circ -dissection Δ . Denote by $A = A(\Delta)$ the graded gentle algebra associated to Δ . Then it follows from Proposition 2.15 that $eAe \simeq ((A^{!})_{1-e})^{!}$ where e is the idempotent in A corresponding to the arcs in L. By our construction $(A^{!})_{1-e} \simeq A(L^{*}) \simeq (A(L))^{!}$ and thus $A(L) \simeq ((A^{!})_{1-e})^{!}$. Hence $A(L) \simeq eAe$. Thus by Theorem 2.10 the right hand side of the recollement is A(L).

4. Application to partially wrapped Fukaya categories

In this section we apply our results to the partially wrapped Fukaya category of a graded surface with stops as defined in [HKK]. More precisely, if a graded gentle algebra A with associated graded marked surface (S, M, Δ) is homologically smooth and proper then $\mathcal{D}^{b}(A)$ is triangle equivalent to the partially wrapped Fukaya category $\mathcal{W}(S, M, \eta(\Delta))$, see [HKK, Proposition 3.4] and [LP, Lemma 3.3], where $\eta(\Delta)$ is the line field corresponding to the grading of Δ . In this case we have the following recollement of partially wrapped Fukaya categories as a consequence of Theorem 3.1. For the statement of the next result we use Remark 3.3.

Theorem 4.1. Let $\mathcal{W}(S, M, \eta(\Delta))$ be the partially wrapped Fukaya category associated to a graded surface dissection (S, M, Δ) with induced line field $\eta(\Delta)$. Let $L \subset \Delta$ be a subset of Δ such that the graded marked surface $\mathcal{W}(S(L), M(L), \eta(L))$ generated by L (as graded ribbon subgraph of Δ) has no marked points in the interior. Then there is a recollement of partially wrapped Fukaya categories

$$\mathcal{W}(S_L, M_L, \eta(\Delta_L)) \longrightarrow \mathcal{W}(S, M, \eta(\Delta)) \longrightarrow \mathcal{W}(S(L), M(L), \eta(L))$$
(4.1)

where the left side corresponds to the graded cut marked surface along L.

Before proving Theorem 4.1, we show the following which is a consequence of Theorem 2.10.

Proposition 4.2. Let A = kQ/I be a graded gentle algebra and let e be an idempotent in A. If two of the three algebras A, eAe and A_e are homologically smooth and proper then so is the third one. Moreover, we have an induced recollement

$$\mathcal{D}^{\mathbf{b}}(A_{e}) \xrightarrow{i^{*}} \mathcal{D}^{\mathbf{b}}(A) \xrightarrow{? \otimes_{eAe}^{\mathbf{L}} eA} \mathcal{D}^{\mathbf{b}}(eAe) , \qquad (4.2)$$

$$\stackrel{i^{*}}{\underset{i^{!}}{\overset{i^{*}}}{\overset{i^{*}}{\overset{i^{*}}}{\overset{i^{*}}{\overset{i^{*}}{\overset{i^{*}}}{\overset{i^{*}}{\overset{i^{*}}{\overset{i^{*}}{\overset{i^{*}}}{\overset{i^{*}}{\overset{i^{*}}{\overset{i^{*}}}{\overset{i^{*}}{\overset{i^{*}}}{\overset{i^{*}}{\overset{i^{*}}}}}}}}}}}}}}}}}}}}$$

such that i^* sends e_jA to e_jA_e and i_* sends $\operatorname{top}(e_jA_e)$ to $\operatorname{top}(e_jA)$ for any $j \neq 1, 2, \ldots, m$. In this case, $\mathcal{D}^{\mathrm{b}}(eAe)$ is triangle equivalent to $\mathcal{D}^{\mathrm{b}}((A^!)_{(1-e)})$.

We point out that if we only assume one of A, A_e and eAe is homologically smooth and proper, Proposition 4.2 is not true in general. We give an example.

Example 4.3. Let $Q = 1 \xrightarrow{\alpha}_{\beta} 2$ and $I = \langle \alpha \beta \rangle$. Let A = KQ/I be the zero graded gentle

algebra and let $e = e_2$. Then $eAe = K[X]/(X^2)$ with degree |X| = 0 and $A_e = K[Y]$ with |Y| = -1. The functor $? \otimes_{eAe}^{\mathbf{L}} eA$ sends the simple eAe-module k to the complex

$$\cdots \xrightarrow{\beta\alpha} e_2 A \xrightarrow{\beta\alpha} e_2 A \xrightarrow{\beta\alpha} e_2 A \to 0 \to \cdots$$

which has infinite dimensional total cohomology. So ? $\otimes_{eAe}^{\mathbf{L}} eA$ does not restricts to $\mathcal{D}^{\mathbf{b}}$.

For a homologically smooth and proper graded gentle algebra A, we show that the bounded derived categories of A and its quadratic dual are equivalent.

Proposition 4.4. Let A = kQ/I be a graded gentle algebra which is homologically smooth and proper. Then there is a triangle equivalence $\mathcal{D}^{\mathbf{b}}(A) \simeq \mathcal{D}^{\mathbf{b}}(A^!)$.

Proof. Let (S, M, Δ, G) be the graded surface associated to A. Let $T := \bigoplus_{i \in Q_0} S_i$ where S_i is the simple dg A-modules corresponding to $i \in Q_0$. By [HKK], S_i corresponds to some curve γ_i

on S. It is well-known that in the ungraded case, the set $\Delta' := \{\gamma_i \mid i \in Q_0\}$ is exactly the dual dissection Δ^* of Δ up to rotation, where S_i can be realized as a complex of projective modules given by the curve γ_i , which is exactly (2.2) for the zero graded case. In general, when we take the grading into consideration, the arc S_i does not change, and the total complex Tot(P) of (2.2) is exactly the twisted object corresponding to γ_i constructed in [HKK]. So Δ' , which is Δ^* up to rotation, is also an admissible dissection of S.

Now we consider the graded surface dissection Δ' of (S, M) whose grading is induced by $\eta(\Delta)$. Let *B* be the graded gentle algebra corresponding to (S, M, Δ') . Since by definition $\eta(\Delta)$ is equal to $\eta(\Delta')$, the partially wrapped Fukaya categories $\mathcal{W}(S, M, \eta(\Delta))$ and $\mathcal{W}(S, M, \eta(\Delta'))$ are equal. Then by [HKK] the following are equivalent

$$\mathcal{D}^{\mathbf{b}}(A) \simeq \mathcal{W}(S, M, \eta(\Delta)) = \mathcal{W}(S, M, \eta(\Delta')) \simeq \mathcal{D}^{\mathbf{b}}(B),$$

where $S_i \in \mathcal{D}^{\mathbf{b}}(A)$ is sent to $\gamma_i \in \mathcal{W}(S, M, \eta(\Delta))$, and in turn it is send to $e_i B \in \mathcal{D}^{\mathbf{b}}(B)$. Now let $\gamma = \bigoplus_{i \in Q_0} \gamma_i$ in $\mathcal{W}(S, M, \eta(\Delta))$. Then we obtain isomorphisms of graded algebras:

$$B = \bigoplus_{j \in \mathbb{Z}} \operatorname{Hom}_{\mathcal{D}^{\mathsf{b}}(B)}(B, B[j]) \cong \bigoplus_{j \in \mathbb{Z}} \operatorname{Hom}_{\mathcal{W}(S, M, \eta(\Delta))}(\gamma, \gamma[j]) \cong \bigoplus_{j \in \mathbb{Z}} \operatorname{Hom}_{\mathcal{D}^{\mathsf{b}}(A)}(T, T[j]) \cong A^{!},$$

where the last isomorphism is due to Proposition 2.7. Thus we have triangle equivalences $\mathcal{D}^{\mathbf{b}}(A) \simeq \mathcal{D}^{\mathbf{b}}(B) \simeq \mathcal{D}^{\mathbf{b}}(A^{!})$.

It follows from the proof of Proposition 4.4 that the dual dissection gives rise to the geometric model for the quadratic dual. More precisely, we show the following.

Corollary 4.5. The surface (S, M) together with the graded dual surface dissection Δ^* is the geometric model for $\mathcal{D}^{\mathbf{b}}(A^!)$.

We now characterize when the algebras A, A_e and eAe are homologically smooth or proper, leading to the proof of Proposition 4.2.

Recall, for example from [HKK, Proposition 3.5], that A is homologically smooth if and only if there are no cyclic paths $\alpha_1 \cdots \alpha_n$ with $\alpha_i \alpha_{i+1} \in I$ for any $i \in \mathbb{Z}/n\mathbb{Z}$ and A is proper if and only if there are no cyclic paths $\alpha_1 \cdots \alpha_n$ with $\alpha_i \alpha_{i+1} \notin I$ for any $i \in \mathbb{Z}/n\mathbb{Z}$. We have the following immediate consequence.

Lemma 4.6. A is homologically smooth (resp. proper) if and only if $A^!$ is proper (resp. homologically smooth).

And we have the following observation.

Lemma 4.7. (1) If A is proper, then eAe is also proper. (2) If A is homologically smooth, then A_e is also homologically smooth.

Proof. Since eAe is a subalgebra of A, so it is clear that eAe is proper showing (1) and (2) directly follows using Lemma 4.6.

Notice that A is homologically smooth does not necessarily imply eAe homologically smooth, see for example Example 4.3. The same example shows that A proper does not imply A_e proper.

Lemma 4.8. Let A be homologically smooth and proper. Then the following are equivalent.

(1) eAe is homologically smooth.

(2) A_e is proper.

Proof. (1) \Rightarrow (2) If A_e is not proper, then there is a cyclic path $\alpha_1 \cdots \alpha_p$ with $\alpha_i \alpha_{i+1} \notin I$ for all $i \in \mathbb{Z}/p\mathbb{Z}$. We may assume without loss of generality that $\alpha_l = [\beta_{l_1} \cdots \beta_{l_{r(l)}}]$ for any $1 \leq l \leq q$ and $\alpha_l = [\alpha_l]$ for any $q < l \leq p$. Then $\beta_{1_1} \cdots \beta_{1_{r(1)}} \cdots \beta_{q_{r(q)}}$ is a cyclic path in *eAe* with any composition of successive paths is in I and *eAe* is not homologically smooth.

 $(2) \Rightarrow (1)$ Note that $A_e \cong ((1-e)A^!(1-e))^!$ and $eAe \cong ((A^!)_{1-e})^!$ by Proposition 2.15, and the result follows from the first part of the proof.

Note that for $(1) \Rightarrow (2)$, it is enough to assume that A is proper and dually, for $(2) \Rightarrow (1)$ it is enough to assume A is homologically smooth.

Proposition 4.9. (1) Assume A_e and eAe both are proper, then A is also proper. (2) Assume A_e and eAe both are homologically smooth, then A is also homologically smooth.

Proof. (1) If A is not proper, then there is a cyclic path $\alpha_1 \cdots \alpha_p$ with $\alpha_i \alpha_{i+1} \notin I$ for any $i \in \mathbb{Z}/p\mathbb{Z}$. Recall that $e = e_1 + \cdots e_m$. If $\{s(\alpha_1), \ldots, s(\alpha_p)\} \cap \{1, \ldots, m\} \neq \emptyset$, then *eAe* contains a cycle without relations, which contradicts the assumption that *eAe* is proper. So $\{s(\alpha_1), \ldots, s(\alpha_p)\} \cap \{1, \ldots, m\} = \emptyset$, and in this case, the cyclic path $\alpha_1 \cdots \alpha_p$ is contained in A_e , which implies A_e is non-proper, a contradiction. Thus A is proper.

(2) Since $A_e \cong ((1-e)A^!(1-e))^!$ and $eAe \cong ((A^!)_{1-e})^!$ by Proposition 2.15, we have that (2) directly follows from (1) and Lemma 4.6.

Now we prove Proposition 4.2.

Proof of Proposition 4.2. By Lemma 4.8 and Proposition 4.9, we know that if any two of A, A_e and eAe are homologically smooth and proper then the third one is also homologically smooth and proper. In this case, the recollement (2.3) restricts to bounded derived categories by Proposition 1.15. In this case, since $eAe \cong ((A^!)_{1-e})^!$ by Proposition 2.15, and since eAe is homologically smooth and proper, by Proposition 4.4, we have a triangle equivalence $\mathcal{D}^{\mathrm{b}}(eAe) \simeq \mathcal{D}^{\mathrm{b}}((A^!)_{(1-e)})$.

We have the following geometric version of Proposition 4.2, recall that if A is a graded gentle algebra associated to a graded marked surface (S, M, Δ, G) then A is homologically smooth if and only if there are no \bullet -punctures in M and A is proper if and only if there are no \circ -punctures in M.

Corollary 4.10. Let (S, M) be the marked surface with a graded admissible dissection (Δ, G) . Let L be a subset of Δ . If two out of the three surfaces (S, M), (S_L, M_L) and $(S_{\Delta^* \setminus L^*}, M_{\Delta^* \setminus L^*})$ have no punctures, then neither has the third.

Proof of Theorem 4.1. Recall that if a graded gentle algebra A with associated graded marked surface (S, M, Δ) is homologically smooth and proper then $\mathcal{D}^{\mathrm{b}}(A)$ is triangle equivalent to the partially wrapped Fukaya category $\mathcal{W}(S, M, \eta_{\Delta})$, see [HKK, Proposition 3.4] and [LP, Lemma 3.3]. Then the result directly follows from Proposition 4.2 and the above.

We end this section by showing the following result.

Proposition 4.11. The recollement (4.2) can be extended to an unbounded ladder.

Recall that an *unbounded ladder* introduced in [AKLY, Section 3] is an infinite diagram of triangulated categories and triangle functors

$$\mathcal{C}' \xrightarrow[i_2]{j_1} \mathcal{C} \xrightarrow[j_2]{j_1} \mathcal{C}'$$

such that any three consecutive rows form a recollement of C relative to C' and C''. To show Proposition 4.11, we need the following two observations.

Lemma 4.12. Assume A is a dg k-algebra which is homologically smooth and proper, then $\mathcal{D}^{\mathbf{b}}(A)$ admits a Serre functor which is given by the Nakayama functor $\nu_A := ? \otimes_A^{\mathbf{L}} DA$.

Proof. Since A is homologically smooth and proper, we have $\operatorname{per}(A) = \mathcal{D}^{\mathrm{b}}(A)$. In particular, $\operatorname{per}(A)$ is Hom-finite and ν_A induces a triangle equivalence from $\operatorname{per}(A)$ to $\operatorname{thick}(DA)$ (see [Ke1, Section 10]). Since we also have $\operatorname{per}(A^{\mathrm{op}}) = \mathcal{D}^{\mathrm{b}}(A^{\mathrm{op}})$, by using the k-duality D, we obtain that $\operatorname{thick}(DA) = \mathcal{D}^{\mathrm{b}}(A)$. So ν_A gives an auto-equivalence of $\mathcal{D}^{\mathrm{b}}(A)$. Moreover, for any $X \in \mathcal{D}(A)$ and $Y \in \operatorname{per}(A)$, we have

$$\operatorname{Hom}_{\mathcal{D}(A)}(X,\nu_A(Y)) \cong D\operatorname{Hom}_{\mathcal{D}(A)}(Y,X).$$

Thus ν_A is a Serre functor of $\mathcal{D}^{\mathrm{b}}(A)$.

Lemma 4.13. [ZZZZ, Lemma 2.8] Let C and D be two Hom-finite triangulated categories with Serre functors ν_C and ν_D respectively. Assume $F: C \to D$ is a triangle functor with a right adjoint $G: D \to C$. Then F admits a left adjoint $\nu_C^{-1}G\nu_D$ and G admits a right adjoint $\nu_D F\nu_C^{-1}$.

Now Proposition 4.11 is immediately from above lemmas.

Proof of Proposition 4.11. Since A, eAe and A_e are all homologically smooth and proper, so their bounded derived categories admit Serre functors ν_A , ν_{eAe} and ν_{A_e} respectively by Lemma 4.12. Since we already have a recollement (4.2), by Lemma 4.13, we can inductively extend it to obtain infinitely many adjoints both upwards and downwards, which gives rise to a ladder by Proposition 1.7 and Lemma 1.8.

5. Exceptional sequences, silting and simple-minded collections for graded Gentle Algebras

5.1. Silting reductions and simple-minded collection reductions. In this section we use Theorems 3.1 and 4.1 to study silting and simple-minded collection (SMC) reductions. Recall that an object P of a k-linear triangulated category \mathcal{T} is called *pre-silting* if $\operatorname{Hom}_{\mathcal{T}}(P, P[>0]) = 0$. We call P silting if it is pre-silting and $\mathcal{T} = \operatorname{thick}_{\mathcal{T}}(P)$. Let $T = \bigoplus_{i=1}^{m} T_i$ be a direct sum of indecomposable objects in \mathcal{T} . We say T is a *pre-simple-minded-collection* (or *pre-SMC* for short), if $\operatorname{Hom}_{\mathcal{T}}(T_i, T_j) = \delta_{i,j}k$ and $\operatorname{Hom}_{\mathcal{T}}(T_i, T_j[<0]) = 0$ for all i, j. We call T a simple-minded collection (or *SMC*) if it is a pre-SMC and $\mathcal{T} = \operatorname{thick}_{\mathcal{T}}(T)$. Let $X \in \mathcal{T}$. We denote by $\operatorname{Filt}(X)$ the smallest extension-closed subcategory of \mathcal{T} containing X. Note that if X is pre-silting, then $\operatorname{Filt}(X)$ coincides with add X, the smallest subcategory of \mathcal{T} containing X and closed under direct summands. We write ${}^{\perp_{\mathcal{T}}} X = \{Y \in \mathcal{T} \mid \operatorname{Hom}_{\mathcal{T}}(Y, X) = 0\}$ and $X^{\perp_{\mathcal{T}}} = \{Y \in \mathcal{T} \mid \operatorname{Hom}_{\mathcal{T}}(X, Y) = 0\}$ (or just write ${}^{\perp_{X}}$ and X^{\perp} for simplicity).

Let A = kQ/I be a graded gentle algebra. Let P be a pre-silting object of per(A). Silting reduction, introduced by [IY], is defined as the Verdier quotient $\mathcal{U} = per(A)/\text{thick}(P)$. Under the conditions,

- (P1) add P is covariantly finite in $^{\perp_{\mathsf{per}(A)}}P[>0]$ and contravariantly finite in $P[<0]^{\perp_{\mathsf{per}(A)}}$,
- (P2) For any $X \in per(A)$, we have $\operatorname{Hom}_{per(A)}(X, P[l]) = 0 = \operatorname{Hom}_{per(A)}(P, X[l])$ for $l \gg 0$,

we can realize \mathcal{U} as a certain subfactor category $\frac{\mathcal{Z}}{[\operatorname{add} P]}$ of $\operatorname{per}(A)$, where $\mathcal{Z} := ({}^{\perp}P[>0]) \cap (P[<0]^{\perp})$ (see [IY, Theorem 3.1]), and moreover, there is a bijection between silting objects in \mathcal{U} and those in $\operatorname{per}(A)$ containing $P([IY, \operatorname{Theorem 3.7}])$.

Let T be a pre-SMC of $\mathcal{D}^{\mathbf{b}}(A)$, the SMC reduction of $\mathcal{D}^{\mathbf{b}}(A)$ with respect to T is defined as $\mathcal{V} = \mathcal{D}^{\mathbf{b}}(A)/\operatorname{thick}(T)$ (see [J]). Similarly to silting reduction, under the conditions,

- (R1) Filt(T) is covariantly finite in $T[>0]^{\perp_{\mathcal{D}^{b}(A)}}$ and contravariantly finite in $^{\perp_{\mathcal{D}^{b}(A)}}T[<0]$,
- (R2) For any $X \in \mathcal{D}^{\mathbf{b}}(A)$, we have $\operatorname{Hom}_{\mathcal{D}^{\mathbf{b}}(A)}(X, T[l]) = 0 = \operatorname{Hom}_{\mathcal{D}^{\mathbf{b}}(A)}(T, X[l])$ for $l \ll 0$,

we can realize \mathcal{V} as a certain subcategory $\mathcal{S} := ({}^{\perp}T[<0]) \cap (T[>0]^{\perp})$ of $\mathcal{D}^{\mathbf{b}}(A)$ and there is a bijection between SMCs in \mathcal{V} and SMCs in $\mathcal{D}^{\mathbf{b}}(A)$ containing T ([J, Theorem 3.1]).

Theorem 5.1. Let A be a graded gentle algebra corresponding to a graded surface dissection (S, M, Δ) and let $e := \sum_{i=1}^{m} e_i$. Let P = eA be the corresponding projective dg A-module and let T = top(eA) be the corresponding semi-simple dg A-module.

(1) Assume A is proper and P is pre-silting, then the silting reduction per(A)/thick(P) is equivalent to $per(A_e)$. Moreover, it is obtained by cutting (S, M) along the arcs corresponding to e, and there is a bijection

 $\{silting \ objects \ in \ \mathsf{per}(A_e)\} \xleftarrow{1:1}{\longleftrightarrow} \{silting \ objects \ in \ \mathsf{per}(A) \ containing \ P\}.$

(2) Assume A and (1-e)A(1-e) are homologically smooth and proper, and T is a pre-simple minded collection of $\mathcal{D}^{\mathrm{b}}(A)$, then the simple minded collection reduction $\mathcal{D}^{\mathrm{b}}(A)/\operatorname{thick}(T)$ is equivalent to $\mathcal{D}^{\mathrm{b}}((1-e)A(1-e)) \simeq \mathcal{D}^{\mathrm{b}}((A^{!})_{e})$. Moreover, it is obtained by cutting (S, M)along the arcs corresponding to e in the dual dissection and there is a bijection

$$\{SMCs \text{ in } \mathcal{D}^{\mathrm{b}}((A^{!})_{e})\} \stackrel{1:1}{\longleftrightarrow} \{SMCs \text{ in } \mathcal{D}^{\mathrm{b}}(A) \text{ containing } T\}.$$

Remark 5.2. Let A be a homologically smooth and proper graded gentle algebra corresponding to a graded surface dissection (S, M, Δ) . Let Γ be a collection of pairwise non-intersecting \circ arcs in (S, M) corresponding to an object $X \in \mathcal{D}^{\mathbf{b}}(A) = \operatorname{per} A$. If X is pre-silting, then the silting reduction is also given by cutting the surface along Γ . In this case, X does not necessarily corresponds to a projective dg A-module.

The following two lemmas guarantee conditions (P1), (P2) and (R1), (R2) mentioned above hold, which are necessary for us.

Lemma 5.3. Let A be a proper dg k-algebra. For any $X, Y \in per(A)$, we have $Hom_{per(A)}(X, Y[l]) = 0 = Hom_{per(A)}(Y, X[l])$ for $l \gg 0$ and for $l \ll 0$.

Proof. Note that we have $\operatorname{Hom}_{\mathsf{per}(A)}(A, A[i]) = \operatorname{H}^{i}(A)$. Since A is proper, the assertion holds for X = Y = A. So it also holds for any $X, Y \in \operatorname{thick}(A) = \operatorname{per}(A)$.

Lemma 5.4. Let A be a graded gentle algebra which is homologically smooth and proper and let $e := \sum_{i=1}^{m} e_i$. Let $T = \operatorname{top}(eA)$ be the corresponding semi-simple dg A-module. Assume T is a pre-SMC in $\mathcal{D}^{\mathrm{b}}(A)$ and (1-e)A(1-e) is homologically smooth. Then $\operatorname{Filt}(T)$ is functorially finite in $\mathcal{D}^{\mathrm{b}}(A)$.

Proof. Let $e = e_1 + \cdots + e_m$ and let e' = 1 - e. Since e'Ae' is proper and by our assumption, it is homologically smooth, we have that $A_{e'}$ is also homologically smooth and proper by Proposition 4.2. Then we have a recollement by Proposition 4.2

 $\mathcal{D}^{\mathbf{b}}(A_{e'}) \xrightarrow{i^* \atop i_* = i_!} \mathcal{D}^{\mathbf{b}}(A) \xrightarrow{j^* = j^!} \mathcal{D}^{\mathbf{b}}(e'Ae') .$ (5.1)

Moreover, i_* sends $S'_i := \operatorname{top}(e_i A_{e'})$ to S_i for $1 \le i \le m$. So we have an equivalence $i_* : \mathcal{D}^{\mathrm{b}}(A_{e'}) \to \operatorname{thick}(T)$. By (5.1), there are two *t*-structures

$$\mathcal{D}^{\mathbf{b}}(A) = j_{!}(\mathcal{D}^{\mathbf{b}}(e'Ae')) \perp i_{*}(\mathcal{D}^{\mathbf{b}}(A_{e'})) = i_{*}(\mathcal{D}^{\mathbf{b}}(A_{e'})) \perp j_{*}(\mathcal{D}^{\mathbf{b}}(e'Ae')),$$

which implies that $\operatorname{thick}(T) = i_*(\mathcal{D}^{\mathrm{b}}(A_{e'}))$ is functorially finite in $\mathcal{D}^{\mathrm{b}}(A)$.

Next we show that $\operatorname{Filt}(T)$ is functorially finite in $\operatorname{thick}(T)$, then it is functorially finite in $\mathcal{D}^{\mathrm{b}}(A)$ and we are done. Since T is a pre-SMC in $\mathcal{D}^{\mathrm{b}}(A)$, it is a SMC in $\operatorname{thick}(T)$ and moreover, T' := $\bigoplus_{i=1}^{m} S'_{i} \text{ is a SMC in } \mathcal{D}^{\mathrm{b}}(A_{e'}). \text{ Then by Proposition 2.7, we have } (A_{e'})^{!} \cong \bigoplus_{t \in \mathbb{Z}} \operatorname{Hom}_{\mathcal{D}^{\mathrm{b}}(A_{e'})}(T', T'[t])$ is a non-negative dg algebra with $\operatorname{H}^{0}((A_{e'})^{!})$ semisimple. So $A_{e'}$ is a non-positive graded gentle algebra. Thus $\operatorname{Filt}(T')$ is functorially finite in $\mathcal{D}^{\mathrm{b}}(A_{e'})$ by [J, Proposition 3.11] and moreover, $\operatorname{Filt}(T) = i_{*}(\operatorname{Filt}(T'))$ is functorially finite in thick(T). So the assertion holds. \Box

Now we show Theorem 5.1.

Proof of Theorem 5.1. (1) Let $e = e_1 + \cdots + e_m$. Assume P = eA is a pre-silting object in $\operatorname{per}(A)$. Then by Theorems 1.14 and 2.10, the functor i^* in (2.3) induces a triangle equivalence $i^* : \operatorname{per}(A)/j_!(\operatorname{per}(eAe)) = \operatorname{per}(A)/\operatorname{thick}(P) \simeq \operatorname{per}(A_e)$ up to direct summands. Since A is proper, $\operatorname{per}(A)$ is Hom-finite and moreover, add P is functorially finite in $\operatorname{per}(A)$. So (P1) holds. Note that (P2) follows from Lemma 5.3. So we only need to show $\operatorname{per}(A)/\operatorname{thick}(P)$ is idempotent complete, then i^* is an equivalence and the bijection is given by [IY, Theorem 3.7].

We know that $\operatorname{per}(A)/\operatorname{thick}(P)$ is equivalent to the subfactor category $\frac{\mathcal{Z}}{[\operatorname{add} P]}$ of $\operatorname{per}(A)$, where $\mathcal{Z} := ({}^{\perp}P[>0]) \cap (P[<0]^{\perp})$ ([IY, Theorem 3.1]). Since $\operatorname{per}(A)$ is idempotent complete and Homfinite, so is \mathcal{Z} . Since it is well-known that any idempotent morphism in $\frac{\mathcal{Z}}{[\operatorname{add} P]}$ can be lifted to an idempotent morphism in \mathcal{Z} , which means $\frac{\mathcal{Z}}{[\operatorname{add} P]}$ is idempotent complete, we have $\operatorname{per}(A)/\operatorname{thick}(P)$ is idempotent complete. So the assertion is true.

(2) Let $e' = 1 - e_1 - \cdots - e_m$. Assume $T = \bigoplus_{i=1}^m S_i$ is a pre-SMC in $\mathcal{D}^{\mathbf{b}}(A)$. By Propositions 4.4 and 2.15, and the recollement (5.1), we have the following equivalences

 $\mathcal{D}^{\mathrm{b}}(A)/\operatorname{thick}(T) = \mathcal{D}^{\mathrm{b}}(A)/i_{*}(\mathcal{D}^{\mathrm{b}}(A_{e'})) \xrightarrow{\simeq} \mathcal{D}^{\mathrm{b}}(e'Ae') \xrightarrow{\simeq} \mathcal{D}^{\mathrm{b}}((A^{!})_{e}).$

Since by Lemmas 5.3 and 5.4, (R1) and (R2) hold, the assertion follows from [J, Theorem 3.1]. \Box

5.2. Existence of full exceptional sequences. In this section, we study the existence of full exceptional sequences in $\mathcal{D}^{\mathrm{b}}(A)$ for a graded gentle algebra A, which is homologically smooth and proper. Exceptional sequences originate in algebraic geometry [GR, Ru] and are instrumental in the construction of the Fukaya-Seidel categories [S]. Furthermore, exceptional sequences give rise to semi-orthogonal decompositions. Semi-orthogonal decompositions of gentle algebras are studied in [KS] in terms of surface cuts. We begin by recalling the definitions of exceptional sequences. An object X in $\mathcal{D}^{\mathrm{b}}(A)$ is exceptional if $\operatorname{Hom}_{\mathcal{D}^{\mathrm{b}}(A)}(X, X[\neq 0]) = 0$ and $\operatorname{Hom}_{\mathcal{D}^{\mathrm{b}}(A)}(X, X) = k$. We call a sequence (X_1, \ldots, X_n) exceptional if each X_i is an exceptional object in $\mathcal{D}^{\mathrm{b}}(A)$ and if $\operatorname{Hom}_{\mathcal{D}^{\mathrm{b}}(A)}(X_j, X_i[\mathbb{Z}]) = 0$ for all i < j. An exceptional sequence is full if $\operatorname{thick}_{\mathcal{D}^{\mathrm{b}}(A)}(\bigoplus X_i) = \mathcal{D}^{\mathrm{b}}(A)$.

We now define a special family of graded gentle algebras. Namely, let $Q^{(n)}$ be the following quiver

$$1 \xrightarrow{\alpha_{1}}{ \xrightarrow{\gamma_{1}}} 2 \xrightarrow{\delta_{1}}{ 3} \xrightarrow{\gamma_{2}}{ \xrightarrow{\gamma_{2}}} 4 \xrightarrow{\delta_{2}}{ \cdots} \xrightarrow{\delta_{n-1}}{ 2n-1} \xrightarrow{\alpha_{n}}{ \xrightarrow{\beta_{n}}{ \gamma_{n}}} 2n$$
(5.2)

and let $I^{(n)}$ be the ideal generated by the set $\{\alpha_i\beta_i, \beta_i\gamma_i, \gamma_j\delta_j, \delta_j\alpha_{j+1} \mid 1 \leq i \leq n, 1 \leq j \leq n-1\}$. Let $A^{(n)} := kQ^{(n)}/I^{(n)}$ be the graded gentle algebra, whose grading is induced by a map $|\cdot|_{A^{(n)}}$: $Q_1^{(n)} \to \mathbb{Z}$ (we will use $|\cdot|$ for simplicity). Write $a_i := |\alpha_i| + |\beta_i|$ and $b_i := |\beta_i| + |\gamma_i|$ for any $1 \leq i \leq n$. The marked surface associate to $A^{(n)}$ is a surface of genus n with exactly one boundary component with one marked point. So all curves in this surface correspond to a loop or a closed curve.

Note that in this section, whenever we say that two graded algebras A and B are derived equivalent, we mean that $\mathcal{D}^{\mathbf{b}}(A)$ and $\mathcal{D}^{\mathbf{b}}(B)$ are equivalent. We need the following observation.

Lemma 5.5. Let A = kQ/I be a graded gentle algebra which is homologically smooth and proper. Let (S, M, Δ) be the graded marked surface associated with A. Let $\ell_i \in \Delta$ be the arc corresponding to vertex $i \in Q_0$. Then the following are equivalent.

- (1) $e_i A e_i \neq k$ for any $i \in Q_0$,
- (2) ℓ_i is a loop for any $i \in Q_0$,
- (3) (S, M) has only one boundary and one marked \circ -point.

Moreover, if one of the three holds then A is derived equivalent to a graded gentle algebra of the form $A^{(n)}$, for some $n \ge 1$.

Proof. The equivalence of (1)-(3) directly follows from the definitions. Now we assume that (1)-(3) hold. Note (S, M) has a grading as a surface where the grading is induced by the graded gentle algebra A, see [LP]. Since A is homologically smooth and proper, $\mathcal{D}^{\mathrm{b}}(A)$ is derived equivalent to the partially wrapped Fukaya category associated to the graded marked surface (S, M), [HKK, LP]. Then since the surface associated to an algebra of the form $A^{(n)}$ is as in (3), by [HKK] there exists a grading on $A^{(n)}$ such that $\mathcal{D}^{\mathrm{b}}(A^{(n)})$ and $\mathcal{D}^{\mathrm{b}}(A)$ are equivalent since both are equivalent to the same partially wrapped Fukaya category associated to the graded marked surface (S, M).

The following result is a graded version of [CS2, Theorem A], which follows from the description of exceptional dissections in a marked surface in [CS2, Proposition 3.5] and Proposition 4.2.

Theorem 5.6. Let $\mathcal{W}(S, M, \eta(\Delta))$ be a partially wrapped Fukaya category associated to a graded surface dissection $(S, M, \eta(\Delta))$. Then $\mathcal{W}(S, M, \eta(\Delta))$ has a full exceptional sequence if and only if (S, M) is not homeomorphic to a marked surface of genus ≥ 1 with exactly one boundary and one marked point on that boundary.

Theorem 5.6 can be re-stated as follows.

Theorem 5.7. Let A be a graded gentle algebra which is proper and homologically smooth, then $\mathcal{D}^{\mathbf{b}}(A)$ has a full exceptional sequence if and only if it is not derived equivalent to $A^{(n)}$ for some $n \geq 1$.

Proof. We first show the 'if' part. Let (S, M, Δ) be the graded marked surface associated to A = kQ/I. Since A is not derived equivalent to $A^{(n)}$ for any grading of the arrows, then by Lemma 5.5 there are at least two marked points on S and thus by [CS2, Proposition 3.5], there exists an exceptional dissection Δ' of S. We may assume $\Delta = \Delta'$ for simplicity.

Let $Q_0 = \{1, 2, \dots, n\}$. Since Q is a quiver without oriented cycles (see Definition 1.23), we have $e_1Ae_1 = k$ and by Proposition 4.2, there is a recollement

$$\mathcal{D}^{\mathrm{b}}(A_{e_1}) \xrightarrow{F}{F_{\sigma}} \mathcal{D}^{\mathrm{b}}(A) \xrightarrow{? \otimes_{e_1 A e_1}^{L} e_1 A}{\mathcal{D}^{\mathrm{b}}(A)} \mathcal{D}^{\mathrm{b}}(k),$$

where $A_{e_1} = kQ_{e_1}/\langle I_e \rangle$ with $(Q_{e_1})_0 = \{2, 3, \ldots, n\}$ and kQ_{e_1} has no oriented cycles (see Definition 2.8). Let $A_0 := A$ and $A_1 := A_{e_1}$. We inductively construct a series of recollements

$$\mathcal{D}^{\mathbf{b}}(A_i) \xrightarrow{\longrightarrow} \mathcal{D}^{\mathbf{b}}(A_{i-1}) \xrightarrow{\longrightarrow} \mathcal{D}^{\mathbf{b}}(k).$$

where $A_i := (A_{i-1})_{e_i}$ for $1 \le i \le n-1$. Note that $A_{n-1} = k$ by our construction. So we get a full exceptional sequence by [Ka, Lemma 2.15].

For the 'only if' part, assume A is derived equivalent to $A^{(n)}$ and $\mathcal{D}^{\mathbf{b}}(A)$ has a full exceptional sequence. In particular, there exists an indecomposable object X such that $\operatorname{Hom}_{\mathcal{D}^{\mathbf{b}}(A^{(n)})}(X, X[i]) = 0$ for $i \neq 0$ and $\operatorname{Hom}_{\mathcal{D}^{\mathbf{b}}(A^{(n)})}(X, X) = k$. Since the surface of $A^{(n)}$ only has one boundary and one marked point, by [HKK] the admissible curve corresponding to X is a loop or a closed curve. It follows from [HKK] that $\dim \bigoplus_{i \in \mathbb{Z}} \operatorname{Hom}_{\mathcal{D}^{\mathbf{b}}(A)}(X, X[i]) \geq 2$. It is a contradiction. So A is not derived equivalent to $A^{(n)}$. \Box

A full exceptional sequence decomposes the derived category into derived categories of vector spaces. This decomposition in terms of the associated surface can be seen as repeatedly cutting off discs where the surface dissection is given by a single arc without self-intersection.

Example 5.8. An example of a decomposition of $\mathcal{D}^{\mathbf{b}}(A)$ as in the proof of Theorem 5.6. Let $Q: 1 \xrightarrow{\alpha} 2 \xrightarrow{\beta} 3 \xrightarrow{\gamma} 4$ and let $I = \langle \alpha\beta, \beta\gamma \rangle$. Let A = kQ/I be a graded gentle algebra. Then (P_4, P_3, P_2, P_1) is a full exceptional sequence of $\mathcal{D}^{\mathbf{b}}(A)$. And we have the following decomposition of $\mathcal{D}^{\mathbf{b}}(A)$:



where the relations are given by $\alpha\beta$ and $\beta\gamma$ and the grading of one arrow is the same as its grading in Q.

We note that any permutation of the projective indecomposables will give a decomposition of $\mathcal{D}^{\mathrm{b}}(A)$ into $\mathcal{D}^{\mathrm{b}}(k)$ induced by a full exceptional sequences in $\mathcal{D}^{\mathrm{b}}(A)$. Note that the full exceptional sequences in this case, do not have only projective indecomposables as their terms.

- **Remark 5.9.** (1) Let A be a homologically smooth and proper dg algebra (not necessarily gentle). Let X be an exceptional object in $\mathcal{D}^{\mathbf{b}}(A)$. We may regard the Verdier quotient $\mathcal{V} = \mathcal{D}^{\mathbf{b}}(A)/\operatorname{thick}(X)$ as the *exceptional reduction* of $\mathcal{D}^{\mathbf{b}}(A)$ at X. Then there are bijections between the full exceptional sequences in \mathcal{V} and full exceptional sequences in $\mathcal{D}^{\mathbf{b}}(A)$ ending at X, and those in $\mathcal{D}^{\mathbf{b}}(A)$ starting at X.
- (2) If A is a graded gentle algebra which is homologically smooth and proper, then the exceptional reduction of $\mathcal{D}^{b}(A)$ at X is obtained by cutting the surface associated to A along the arc corresponding to X.

5.3. Existence of silting objects and simple-minded collections. Recall the graded gentle algebra $A^{(n)}$ constructed from the quiver in (5.2) in the previous section. Our main result in this section is as follows.

Theorem 5.10. Let $\mathcal{W}(S, M, \eta(\Delta))$ be a partially wrapped Fukaya category. Then $\mathcal{W}(S, M, \eta(\Delta))$ has a silting object if and only if it has a simple-minded collection and this is the case if

(1) $\mathcal{W}(S, M, \eta(\Delta))$ is not triangle equivalent to $\mathcal{D}^{\mathrm{b}}(A^{(n)})$, or

(2) $\mathcal{W}(S, M, \eta(\Delta))$ is triangle equivalent to $\mathcal{D}^{\mathrm{b}}(A^{(n)})$ with $a_i \neq 1$ or $b_i \neq 1$ for any $1 \leq i \leq n$.

We re-state Theorem 5.10 as follows.

Theorem 5.11. Let A be a graded gentle algebra which is homologically smooth and proper. Then $\mathcal{D}^{b}(A)$ has a silting object if and only if it has a SMC and this is the case if

- (1) A is not derived equivalent to $A^{(n)}$, or
- (2) A is derived equivalent to $A^{(n)}$ and $a_i \neq 1$ or $b_i \neq 1$ holds for any $1 \leq i \leq n$.

The following shows the first part of Theorem 5.11.

Proposition 5.12. Let A be a homologically smooth and proper dg k-algebra. Then $\mathcal{D}^{b}(A)$ has silting objects if and only if it has SMCs.

Proof. If $\mathcal{D}^{\mathrm{b}}(A)$ has a silting object P, then we have a triangle equivalence $\mathbf{R}\mathscr{H}\mathrm{om}_A(P,?)$: $\mathcal{D}^{\mathrm{b}}(A) \xrightarrow{\simeq} \mathsf{per}(\mathscr{E}\mathrm{nd}(P))$ (see for example, [Ke4, Theorem 3.8]), where $\mathscr{E}\mathrm{nd}(P)$ is the endomorphism dg k-algebra which is non-positive because P is silting. Since A is homologically smooth and proper, then $\mathsf{per}(A) = \mathcal{D}^{\mathrm{b}}(A)$ by Lemma 1.1, so P is perfect and $\mathbf{R}\mathscr{H}\mathrm{om}_A(P,?)$ commutes with infinite direct sum. Moreover, $\mathbf{R}\mathscr{H}\mathrm{om}_A(P,?)$ gives an equivalence $\mathcal{D}(A) \simeq \mathcal{D}(\mathscr{E}\mathrm{nd}(P))$ which restricts to an equivalence $\mathcal{D}^{\mathrm{b}}(A) \simeq \mathcal{D}^{\mathrm{b}}(\mathscr{E}\mathrm{nd}(P))$ by Lemma 1.3. Thus we have $\mathsf{per}(\mathscr{E}\mathrm{nd}(P)) = \mathcal{D}^{\mathrm{b}}(\mathscr{E}\mathrm{nd}(P))$. Since $\mathcal{D}^{\mathrm{b}}(\mathscr{E}\mathrm{nd}(P))$ has a SMC given by simple dg modules, $\mathcal{D}^{\mathrm{b}}(A)$ also has a SMC.

On the other hand, assume $\mathcal{D}^{\mathrm{b}}(A)$ has a SMC X. Let $P_X \to X$ be a \mathcal{H} -projective resolution of X and let $B := \mathscr{E}\mathrm{nd}(P_X)$ be the Koszul dual dg algebra ([Ke1, Section 10]). Then we have a triangle equivalence $\mathbf{R}\mathscr{H}\mathrm{om}_A(P_X,?) : \mathcal{D}^{\mathrm{b}}(A) \simeq \mathrm{per}(B)$, which sends X to the free dg B-module B. Since $\mathcal{D}^{\mathrm{b}}(A) = \mathrm{per}(A)$, we have that $P_X \in \mathrm{per}(A)$ and $\mathbf{R}\mathscr{H}\mathrm{om}_A(R_X,?)$ can be extended to an equivalence $\mathcal{D}(A) \simeq \mathcal{D}(B)$, which restricts to $\mathcal{D}^{\mathrm{b}}(A) \simeq \mathcal{D}^{\mathrm{b}}(B)$ by Lemma 1.3. So we have $\mathrm{per}(B) = \mathcal{D}^{\mathrm{b}}(B)$. Note that $\mathrm{H}^{<0}(B) = 0$ and $\mathrm{H}^0(B)$ is semi-simple, so $\mathrm{H}^0(B)$ is a silting object in $\mathcal{D}^{\mathrm{b}}(B)$ by [KN, Corollary 4.1] and it gives a silting object in $\mathcal{D}^{\mathrm{b}}(A)$. Thus the assertion holds. \Box

Before proving Theorem 5.11, we need the following results.

Lemma 5.13. Let A and A' be two algebras of the form $A^{(1)}$ with different gradings. If $|\alpha|_A + |\beta|_A = |\alpha|_{A'} + |\beta|_{A'}$ and $|\beta|_A + |\gamma|_A| = |\beta|_{A'} + |\gamma|_{A'}$, then per(A) and per(A') are triangle equivalent.

Proof. Let P_1, P_2 be the indecomposable projective dg A-modules corresponding to the vertices 1 and 2, respectively. For any $i \in \mathbb{Z}$, the dg algebra $\operatorname{cnd}_A(P_1 \oplus P_2[i])$ is quasi-isomorphic to \widetilde{A} which as ungraded algebra is of the form $A^{(1)}$. Moreover, the grading of \widetilde{A} is given by $|\alpha|_{\widetilde{A}} = |\alpha| - i$, $|\beta|_{\widetilde{A}} = |\beta| + i$ and $|\gamma|_{\widetilde{A}} = |\gamma| - i$. Now let $i = |\alpha|_A - |\alpha|_{A'}$, then we have $\widetilde{A} = A'$. So we have triangle equivalences $\operatorname{per}(A) \simeq \operatorname{per}(\operatorname{cnd}_A(P_1 \oplus P_2[i])) \simeq \operatorname{per}(A')$.

Lemma 5.14. Let $A = A^{(1)}$. If $a_1 \neq 1$ or $b_1 \neq 1$, then $\mathcal{D}^{\mathbf{b}}(A)$ has a silting object.

Proof. If $a_1 \leq 0$ or $b_1 \leq 0$ holds, we may assume $a_1 \leq 0$. Then by Theorem 2.10, we have a recollement

$$\mathcal{D}(A_{e_2}) \xrightarrow{i^*} \mathcal{D}(A) \xrightarrow{?\otimes_{k}^{\mathbf{L}}e_2A} \mathcal{D}(e_2Ae_2) ,$$

$$i^! \qquad \mathbf{R}\mathscr{H}\mathrm{om}_k(Ae_2,?)$$

where $e_2Ae_2 = k[X]/(X^2)$ with deg $X = a_1 \leq 0$ and $A_{e_2} = k[Y]$ with deg $Y = [\alpha_1\beta_1] = a_1 - 1 < 0$. Since e_2Ae_2 is a silting object in $\operatorname{per}(e_2Ae_2)$, then $e_2A = (e_2Ae_2) \otimes_k^{\mathbf{L}} e_2A$ is pre-silting in $\operatorname{per}(A) = \mathcal{D}^{\mathrm{b}}(A)$. Since A_{e_2} is silting in $\operatorname{per}(A_{e_2})$, then by Theorem 5.1, we know $\mathcal{D}^{\mathrm{b}}(A)$ has a silting object. If $a_1 > 0$ and $b_1 > 0$, we consider $A^!$. Then at least one of $a'_1 := |\alpha_1^{\mathrm{op}}| + |\beta_1^{\mathrm{op}}| = 2 - a_1$ and $b'_1 = |\beta_1^{\mathrm{op}}| + |\gamma_1^{\mathrm{op}}| = 2 - b_1$ is non-positive since $a_1 \geq 2$ or $b_1 \geq 2$ holds. So $\mathcal{D}^{\mathrm{b}}(A^!)$ admits a silting object by above argument. Since $\mathcal{D}^{\mathrm{b}}(A) \simeq \mathcal{D}^{\mathrm{b}}(A^!)$ by Proposition 4.4, we have that $\mathcal{D}^{\mathrm{b}}(A)$ has a silting object.

Proof of Theorem 5.11. Thanks to Proposition 5.12, we only need to show the existence of silting objects. Assume A is derived equivalent to $A^{(n)}$ and $a_i \neq 1$ or $b_i \neq 1$ holds for any $1 \leq i \leq n$. We apply induction on n. The assertion is true for n = 1 by Lemma 5.14. Let $e = e_1 + e_2$. Assume n > 1 and the assertion is true for $1, 2, \ldots, n-1$. Note that we have a recollement by Proposition 4.2



where $A^{(1)} = eA^{(n)}e$ and $A^{(n-1)} = (A^{(n)})_e$ are given by the sub graded quivers of $Q^{(n)}$. Thus $\mathcal{D}^{\mathrm{b}}(A^{(n)})$ has silting objects by our assumption and Theorem 5.1.

If A is not derived equivalent to $A^{(n)}$, then $\mathcal{D}^{\mathbf{b}}(A)$ has a full exceptional sequence by Proposition 5.6. This exceptional sequence give rises to a silting object in $\mathcal{D}^{\mathbf{b}}(A)$ by [AI, Proposition 3.5]. So the assertion is true.

For $A = A^{(1)}$ with $a_1 = 1 = b_1$, we have checked, in a case by case analysis, that for every indecomposable object X in $\mathcal{D}^{\mathbf{b}}(A)$, we have that $\operatorname{Hom}(X, X[n])$ is non-zero, for some positive n. This shows that the converse of Theorem 5.11 is also true in this case and that $\operatorname{per}(A^{(1)})$ does not have a silting object.

Funding. The first author acknowledges support by Shaanxi Normal University and the NSF of China (Grant No. 12271321). The third author acknowledges support by the EPSRC Early Career Fellowship EP/P016294/1. The second and third author acknowledges partial support by the DFG through the project SFB/TRR 191 Symplectic Structures in Geometry, Algebra and Dynamics (Projektnummer 281071066–TRR 191).

References

- [A] C. Amiot, *Catégories triangulées, équivalences et modéles topologiques*, Mémoire d'habilitation à diriger des recherches, https://www-fourier.ujf-grenoble.fr/ amiot/HDR.pdf 15
- [AH] I. Assem and D. Happel, Generalized tilted algebras of type A_n , Comm. Algebra 9 (1981), no. 20, 2101–2125. 1
- [AS] I. Assem. and A. Skowroński, Iterated tilted algebras of type affine A_n , Math. Z. 195 (1987), no. 2, 269-290. 1 [AKLY] L. A. Hügel, S. Koenig, Q. H. Liu, D. Yang, Ladders and simplicity of derived module categories, J. Algebra
- 472 (2017), 15–66. 1, 2, 3, 21
- [AI] T. Aihara, O. Iyama, Silting mutation in triangulated categories, J. Lond. Math. Soc. (2) 85 (2012), no. 3, 633–668. 28
- [APS] C. Amiot, P-G. Plamondon and S. Schroll, A complete derived invariant for gentle algebras via winding numbers and Arf invariants, Selecta Math. (N.S.) 29 (2023), no. 2, 30. 7, 15, 17
- [B] Michael J. Bardzell, The alternating syzygy behavior of monomial algebras, J. Algebra 188 (1997), no. 1, 69–89.
 12
- [BBD] A. A. Beilinson, J. Bernstein and P. Deligne. Faisceaux pervers. Analysis and topology on singular spaces, I (Luminy, 1981), 5–171, Astérisque, 100, Soc. Math. France, Paris, 1982. 1, 4
- [BGS] A. A. Beilinson, V. Ginzburg, W. Soergel, Koszul duality patterns in representation theory, J. Amer. Math. Soc. 9 (1996), no. 2, 473–527. 2
- [BCL] C. Braun, J. Chuang, A. Lazarev, Derived localisation of algebras and modules, Adv. Math. 328 (2018), 555–622. 2
- [CPS] E. Cline, B. Parshall and L. L. Scott, Stratifying endomorphism algebras. Mem. Amer. Math. Soc. 124 (1996), no. 591, viii+119 pp. 4
- [CS1] W. Chang and S. Schroll, A geometric realization of silting theory for gentle algebras, Math. Z. 303 (2023), no. 3, 67. 15
- [CS2] W. Chang and S. Schroll, Exceptional sequences in the derived category of a gentle algebra, to appear in Selecta Math. arXiv:2205.15830, 2022. 25
- [D] V. Drinfeld, DG quotients of DG categories, J. Algebra 272 (2004), no. 2, 643–691. 2, 13
- [GL] W. Geigle, H. Lenzing, Perpendicular categories with applications to representations and sheaves, J. Algebra 144 (1991), no. 2, 273–343. 5
- [GR] A. L. Gorodentsev, A.N. Rudakov, Exceptional vector bundles on projective spaces, Duke Math. J. 54 (1987), no. 1, 115–130. 2, 24
- [Hap] D. Happel, Reduction techniques for homological conjectures, Tsukuba J. Math. 17 (1993), no. 1, 115–130. 1
 [Han] Y. Han, Yang, Recollements and Hochschild theory, J. Algebra 397 (2014), 535–547. 1
- [HKK] F. Haiden, L. Katzarkov, and M. Kontsevich, Flat surfaces and stability structures, Publ. Math. Inst. Hautes Études Sci. 126 (2017), 247-318. 1, 2, 7, 15, 19, 20, 21, 25
- [IY] O. Iyama, D. Yang, Silting reduction and Calabi-Yau reduction of triangulated categories, Trans. Amer. Math. Soc. 370 (2018), no. 11, 7861–7898. 22, 24
- [J] H. B. Jin, Reductions of triangulated categories and simple-minded collections, to appear in JLMS, arXiv:1907.05114, 2019. 22, 24

- [KN] B. Keller, P. Nicolás, Weight structures and simple dg modules for positive dg algebras, Int. Math. Res. Not. IMRN 2013, no. 5, 1028–1078. 2, 27
- [KaY1] M. Kalck, D. Yang, Relative singularity categories I: Auslander resolutions, Adv. Math. 301 (2016), 973– 1021. 5

[KaY2] M. Kalck, D. Yang, Relative singularity categories II: DG models, Preprint arXiv:1803.08192, 2018. 2, 13

- [KoY] S. Koenig, D. Yang, Silting objects, simple-minded collections, t-structures and co-t-structures for finitedimensional algebras, Doc. Math. 19 (2014), 403–438. 2
- [Ke1] B. Keller, Deriving DG categories, Ann. Sci. École Norm. Sup. (4) 27 (1994), no. 1, 63–102. 2, 3, 6, 22, 27
- [Ke2] B. Keller, Invariance and localization for cyclic homology of DG algebras, J. Pure Appl. Algebra 123 (1998), no. 1-3, 223–273. 1
- [Ke3] Keller, Bernhard On the cyclic homology of exact categories, J. Pure Appl. Algebra 136 (1999), no.1, 1–56. 2
- [Ke4] B. Keller, On differential graded categories, International Congress of Mathematicians. Vol. II, 151–190, Eur. Math. Soc., Zürich, 2006. 3, 23, 27
- [Ko] S, König, Tilting complexes, perpendicular categories and recollements of derived module categories of rings, J. Pure Appl. Algebra 73 (1991), no. 3, 211–232. 1
- [KoN] S. König, H. Nagase, Hochschild cohomology and stratifying ideals, J. Pure Appl. Algebra 213 (2009), no. 5, 886–891. 1
- [Ka] M. Kalck, Derived categories of quasi-hereditary algebras and their derived composition series, Representation theory – current trends and perspectives, 269–308, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2017. 25
- [KS] J. Kopřiva and J. Šťovíček, Semiorthogonal decompositions for bounded derived categories of gentle algebras, Preprint arXiv: 2209.14496, 2022. 24
- [LP] Y. Lekili and A. Polishchuk, Derived equivalences of gentle algebras via Fukaya categories. Math. Ann. 376 (2020), no. 1-2, 187-225. 1, 2, 7, 15, 19, 21, 25
- $[{\rm M}]\,$ J. Miyachi, Localization of triangulated categories and derived categories. J. Algebra 141 (1991), no. 2, 463-483. $_5$
- [N1] A. Neeman, The connection between the K-theory localization theorem of Thomason, Trobaugh and Yao and the smashing subcategories of Bousfield and Ravenel. Ann. Sci. École Norm. Sup. (4) 25 (1992), no. 5, 547–566.
- [N2] A. Neeman, The Grothendieck duality theorem via Bousfield's techniques and Brown representability, J. Amer. Math. Soc. 9(1)(1996), 205–236. 6
- [N3] A. Neeman, Triangulated Categories, Ann. of Math. Stud., vol. 148, Princeton University Press, Princeton, NJ, 2001. 4, 5
- [NS] P. Nicolás and M. Saorín, Parametrizing recollement data for triangulated categories. J. Algebra, 322(2009) 1220–1250. 2, 5
- [Opm] Steffen, Oppermann, Quivers for silting mutation, Adv. Math. 307 (2017), 684–714. 10
- [Op] S. Opper, On auto-equivalences and complete derived invariants of gentle algebras, Preprint, arXiv:1904.04859, 2019. 7
- [OPS] S. Opper G-P Plamondon and S. Schroll, A geometric model for the derived category of gentle algebras, Preprint, arXiv:1801.09659, 2018. 7, 9
- [P] D. Pauksztello, Homological epimorphisms of differential graded algebras, Comm. Algebra 37 (2009), no. 7, 2337–2350. 5
- [PPP] Y. Palu, V. Pilaud and P-G Plamondon, Non-kissing and non-crossing complexes for locally gentle algebras, J. Comb. Algebra 3 (2019), no. 4, 401–438. 9
- [Ri] J. Rickard, Equivalences of derived categories for symmetric algebras, J. Algebra 257 (2002), no. 2, 460–481. 2
- [Ru] A. N. Rudakov, Exceptional collections, mutations and helices, Helices and vector bundles, 1–6, London Math. Soc. Lecture Note Ser., 148, Cambridge Univ. Press, Cambridge, 1990. 2, 24
- [S] Paul Seidel. Vanishing cycles and mutation. In 3rd European congress of mathematics (ECM), Barcelona, Spain, July 10–14, 2000. Volume II, pages 65–85. Basel: Birkhäuser, 2001. 2, 24
- [T] P. Tamaroff, Minimal models for monomial algebras, Homology Homotopy Appl. 23 (2021), no. 1, 341–366. 10
 [ZZZZ] P. Zhang, Y. H. Zhang, G. D. Zhou, L. Zhu, Unbounded ladders induced by Gorenstein algebras, Colloq. Math. 151 (2018), no. 1, 37–56.

22

School of Mathematics and Statistics, Shaanxi Normal University, Xi'an 710062, China *Email address:* changwen1610163.com

Department of Mathematics, Universität zu Köln, Weyertal 86-90, 50931 Köln, Germany *Email address:* hjin@math.uni-koeln.de

Department of Mathematics, Universität zu Köln, Weyertal 86-90, 50931 Köln, Germany and Institutt for matematiske fag, NTNU, N-7491 Trondheim, Norway

Email address: schroll@math.uni-koeln.de

30