

A Possible Solution to the Helium Anomaly of EMPRESS VIII by Cuscuton Gravity Theory

Kazunori Kohri^{1,2,3,*} and Kei-ichi MAEDA^{4,5,†}

¹Theory Center, IPNS, KEK, 1-1 Oho, Tsukuba, Ibaraki 305-0801, Japan

²The Graduate University for Advanced Studies (SOKENDAI), 1-1 Oho, Tsukuba, Ibaraki 305-0801, Japan

³Kavli IPMU (WPI), UTIAS, The University of Tokyo, Kashiwa, Chiba 277-8583, Japan

⁴Department of Pure and Applied Physics, Graduate School of Advanced Science and Engineering, Waseda University, Okubo 3-4-1, Shinjuku, Tokyo 169-8555, Japan

⁵Center for Gravitational Physics and Quantum Information, Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8555, Japan

(Dated: June 24, 2022)

We discuss cosmology based on the cuscuton gravity theory to resolve the anomaly of the observational ${}^4\text{He}$ abundance reported by the EMPRESS collaboration. We find that the gravitational constant G_{cos} in Friedmann equation should be smaller than the Newton's constant G_{N} such that $\Delta G_{\text{N}}/G_{\text{N}} \equiv (G_{\text{cos}} - G_{\text{N}})/G_{\text{N}} = -0.086_{-0.028}^{+0.026}$ (68% C.L.) in terms of big-bang nucleosynthesis, which excludes $\Delta G_{\text{N}} = 0$ at more than 95 % C.L. To fit the data, we obtain a negative mass squared of a non-dynamical scalar field with the Planck-mass scale as $\sim -\mathcal{O}(1)M_{\text{PL}}^2(\mu/0.5M_{\text{PL}})^4$ with the cuscuton mass parameter μ . This fact could suggest the need for modified gravity theories such as the cuscuton gravity theory with a quadratic potential, which can be regarded as the low-energy Hořava-Lifshitz gravity, and might give a hint of quantum gravity.

I. INTRODUCTION

Quite recently, the EMPRESS collaboration (EMPRESS VIII) newly observed 10 Extremely Metal-Poor Galaxies (EMPGs) with metallicity ($< 0.1Z_{\odot}$) by using Subaru telescope, and obtaining data of the ${}^4\text{He}$ to hydrogen ratio (${}^4\text{He}/\text{H}$) by measuring the $\text{HeI}\lambda 10830$ near-infrared emission [1]. By analyzing the data of 64 galaxies in total with 13 EMPGs (including the new 10 EMPGs), they estimated the primordial mass fraction of ${}^4\text{He}$ to be $Y_p = 0.2379_{-0.0030}^{+0.0031}$, by extrapolating the data into the value at zero metallicity (the oxygen to hydrogen ratio $\text{O}/\text{H} \rightarrow 0$).

Comparing the data with theoretical predictions in the standard big-bang nucleosynthesis (BBN), they obtained the effective number of the neutrino species to be $N_{\nu,\text{eff}} = 2.41_{-0.21}^{+0.19}$ at 68 % C.L. It is remarkable that this means that the standard value of $N_{\nu,\text{eff}}$ predicted in the big-bang cosmology ($= N_{\nu,\text{eff, std}} \simeq 3.045 - 3.046$ [2, 3]), is observationally excluded at more than 2σ .

Seriously-taking this discrepancy of $N_{\nu,\text{eff}}$ in the standard big-bang cosmology, we must consider modified theories beyond the standard model. The EMPRESS collaboration extended their framework to a new theory beyond the standard model by adding one more free parameter, so-called “degeneracy parameter”, ξ_{ν_e} which means a non-zero lepton number in the electron neutrino sector. It has been known for a long time that a positive ξ_{ν_e} can reduce Y_p without changing the number of neutrino species much [4] by which the Helium anomaly can be solved this time. The best-fit value is $\xi_{\nu_e} \sim 0.05$ with

excluding $\xi_{\nu_e} = 0$ at more than 1σ [1]. Theoretically, such a large lepton number can be produced even after the cosmic temperature is smaller than the weak scale $\mathcal{O}(10^2)$ GeV in models with Q -balls (L -balls) [5, 6], oscillating sterile neutrinos [7], etc. In future, we can measure ξ_{ν_e} more precisely by planned observations of 21cm + CMB down to errors of $\Delta\xi_{\nu_e} \sim 5 \times 10^{-3}$ [8].

There is another way to solve this anomaly, which is a modification of the Einstein gravity. In terms of a modified gravity theory, recently an interesting model to realize dark energy has been proposed, called a cuscuton gravity theory [9, 10] or its extended version [11–13]. In the context of the beyond Horndeski theories, the original cuscuton gravity theory was extended to be a generalized one [14], in which the second-order time derivatives of a scalar field in the equation of motion disappears. Thus the scalar field appearing in the theories is just a non-dynamical shadowy mode. There is a new type of minimally modified gravity theory, which also has only two gravitational degrees of freedom [15]. It is called VCDM, which includes a cuscuton gravity theory and gives the equivalence in cosmological models [16]. As shown in Ref. [16, 17], both theories are related to each other.

As an attractive feature in the models of the cuscuton gravity theory, it is notable that the gravitational constant G_{cos} which appears in the Friedmann equations can be different from Newton's constant G_{N} .

In this *Letter*, we discuss how we can resolve the ${}^4\text{He}$ anomaly in the models of the cuscuton gravity theory. A modification on $N_{\nu,\text{eff}}$ from its standard value $N_{\nu,\text{eff, std}} = 3.046$ effectively has an identical effect on a modification on the gravitational constant without changing $N_{\nu,\text{eff}}$ in the Friedmann equations. Thus, we can look for a solution to the ${}^4\text{He}$ anomaly by modifying the gravitational constant in the cuscuton gravity theory. In particular,

* kohri-at-post.kek.jp

† maeda-at-waseda.jp

we show that we can concretely constrain the parameters in the models of the cuscuton gravity theory from the observations.

II. BOUNDS FROM BIG-BANG NUCLEOSYNTHESIS

In Ref. [1], the EMPRESS collaboration reported the primordial mass fraction of ${}^4\text{He}$,

$$Y_p = 0.2379_{-0.0030}^{+0.0031}, \quad (2.1)$$

at 68%C.L. According to the theoretical predictions in the BBN computation, it gives the effective number of neutrino species,

$$N_{\nu,\text{eff}} = 2.41_{-0.21}^{+0.19}, \quad (2.2)$$

which excludes the standard value, $N_{\nu,\text{eff,std}} = 3.046$ predicted in the big-bang cosmology more than 2σ .

The Friedmann equation in the ΛCDM model with vacuum energy V_0 is given by

$$H^2 = \frac{8\pi G_{\text{cos}}}{3}(\rho + V_0). \quad (2.3)$$

Since we can ignore the vacuum energy at the BBN stage, the Hubble parameter H is represented by a product of the energy density of the Universe ρ and G_{cos} which is the ‘‘effective’’ gravitational constant appeared in the Friedmann equations. It is notable that G_{cos} can be potentially different from the Newton’s constant, G_{N} , and we may write the difference to be $\Delta G_{\text{N}} \equiv G_{\text{cos}} - G_{\text{N}}$.

Then, we obtain a relation

$$\frac{\Delta G_{\text{N}}}{G_{\text{N}}} = \frac{7}{21 + \sqrt[3]{2 \cdot 11^4}} \Delta N_{\nu,\text{eff}}, \quad (2.4)$$

with $\Delta N_{\nu,\text{eff}} \equiv N_{\nu,\text{eff}} - N_{\nu,\text{eff,std}}$, and the prefactor of it approximately gives $7/(21 + \sqrt[3]{2 \cdot 11^4}) \simeq 0.135$ for $T \lesssim m_e$ with m_e being electron mass. On the other hand, we may have another value of the prefactor (~ 0.163) in case of $T \gtrsim m_e$. The difference between the former and the latter values comes from that neutrinos decoupled from the thermal bath just before $T \sim m_e$, and only photon was heated by the e^+e^- annihilation at around $T \sim m_e$. In this study, the decoupling temperature of the weak interaction between neutron and proton, T_{dec} , which mainly determines Y_p , tends to get delayed compared to the one in the standard big-bang cosmology ($T_{\text{dec}} \sim 0.8$ MeV) due to $N_{\nu,\text{eff}} < N_{\nu,\text{eff,std}}$. Thus, we adopt the former value ($=0.135$) in this study, which also gives a more conservative absolute value of the magnitude of $|\Delta G_{\text{N}}/G_{\text{N}}|$.

From the observational data (2.2), we obtain the bound on G_{cos} to be

$$\frac{G_{\text{cos}}}{G_{\text{N}}}\Big|_{\text{BBN}} = 0.914_{-0.028}^{+0.026} \quad (68\% \text{C.L.}), \quad (2.5)$$

or equivalently, $\Delta G_{\text{N}}/G_{\text{N}} = -0.086_{-0.028}^{+0.026}$ (68% C.L.).

III. BOUNDS FROM COSMIC MICROWAVE BACKGROUND

The modification of the gravitational constant can be constrained by observations of fluctuation and polarizations of the cosmic microwave background (CMB). Here is a remark about the invalidity of a translation of a bound on $N_{\nu,\text{eff}}$ from the CMB observation to the one on $G_{\text{cos}}/G_{\text{N}}$. Because background neutrinos are thermally produced and have adiabatic fluctuations, a bound on $N_{\nu,\text{eff}}$ from the CMB is obtained by both the total energy density and the evolution of adiabatic curvature perturbation. Therefore, there is no simple one-to-one mapping among the bounds on $N_{\nu,\text{eff}}$ and G_{cos} although an order-of-magnitude discussion would be still possible.

By using the data released by the Planck collaboration in 2018, the authors of Ref. [18] reported that the observational bound on $\Delta \equiv \sqrt{G_{\text{cos}}/G_{\text{N}}} - 1$ obtained by the CMB and the baryon acoustic oscillation (BAO) to be $\Delta = -0.022 \pm 0.023$ at 68% C.L. This gives the bounds on the gravitational constant,

$$\frac{G_{\text{cos}}}{G_{\text{N}}}\Big|_{\text{CMB+BAO}} = 0.956 \pm 0.045 \quad (68\% \text{C.L.}). \quad (3.1)$$

IV. MODELS OF THE CUSCUTON GRAVITY THEORY

The action of the cuscuton gravity theory is represented by [9, 10],

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2} M_{\text{PL}}^2 R + \mu^2 \sqrt{-X} - V(\phi) \right] + S_M(g_{\mu\nu}, \psi_M), \quad (4.1)$$

with M_{PL} being the reduced Planck mass ($\simeq 2.436 \times 10^{18}$ GeV) where R , $V = V(\phi)$, and $S_M(g_{\mu\nu}, \psi_M)$ mean the Ricci scalar, the potential energy of a scalar field ϕ , and the action of the matter field(s) ψ_M , respectively. Here the kinetic term X is defined by

$$X \equiv g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi. \quad (4.2)$$

and μ is the mass parameter of the cuscuton gravity theory.

In the cuscuton gravity theory, the potential $V(\phi)$ can be arbitrary, but if we assume the quadratic form of the potential such that

$$V = V_0 + \frac{1}{2} \alpha \phi^2, \quad (4.3)$$

which can be regarded as the low-energy Hořava-Lifshitz gravity [19], we find the ΛCDM cosmology with modification of the gravitational constant [10]. Here α means the mass squared parameter which has either a positive or negative signature. Then, the flat Friedmann equation is written by Eq. (2.3).

Note that cosmology in the VCDM theory is equivalent to that in the cuscuton gravity [16]. We can also show the

cuscuton gravity with a quadratic potential is equivalent to the Einstein-aether theory[20] with only one coupling constant c_2 [21]¹. In both theories (the Λ CDM and the Einstein-aether theory), we can take the Newtonian limit, which shows the Newtonian gravitational constant G_N is given by the reduced Planck mass as M_{PL} as $G_N = (8\pi M_{\text{PL}}^2)^{-1}$.

In those theories, the effective gravitational constant in the Friedmann equation is given by

$$\frac{G_F}{G_N} = \left(1 - \frac{3}{2} \frac{\mu^4}{\alpha M_{\text{PL}}^2}\right)^{-1} = \left(1 + \frac{3}{2} c_2\right)^{-1} = 1 - \frac{3}{2} \beta_2, \quad (4.4)$$

where the ‘‘potential’’ of the Λ CDM scalar field φ , which has the mass-dimension two, is chosen as

$$\mathcal{V}_{\Lambda\text{CDM}} = \frac{V_0}{M_{\text{PL}}^2} + \frac{1}{2} \beta_2 \varphi^2, \quad (4.5)$$

with the dimensionless mass-parameter β_2 .

In order to find $G_{\text{cos}} < G_N$, each parameter should satisfy $\alpha < 0$, $c_2 > 0$ and $\beta_2 > 0$. Note that we have the relation between these parameters as

$$\frac{\mu^4}{\alpha M_{\text{PL}}^2} = -c_2 \quad \text{and} \quad \beta_2 = \frac{c_2}{1 + \frac{3}{2} c_2}.$$

In what follows, we shall first discuss the constraints on c_2 just for simplicity, but we can translate them into the constraints on the other parameters.

From the observational bound from the EMPRESS VIII on G_{cos}/G_N shown in (2.5), we obtain the bound on c_2 ,

$$c_2 = 0.0626_{-0.0199}^{+0.0233} \quad (68\% \text{C.L.}), \quad (4.6)$$

with the BBN, which gives $0.0239 \leq c_2 \leq 0.1108$ at 95% C.L. It is remarkable that $c_2 = 0$ (general relativity) is excluded at more than 95% C.L. by BBN.

On the other hand, the CMB+BAO data gives

$$c_2 = 0.0307_{-0.0313}^{+0.0344} \quad (68\% \text{C.L.}). \quad (4.7)$$

which covers most of the parameter range in (4.6) allowed by EMPRESS VIII with BBN due to its larger error bars than those of (4.6). Then, we have the range $-0.0293 \leq c_2 \leq 0.1032$ at 95% C.L. Thus, contrary to the case of the BBN, general relativity ($c_2 = 0$) is allowed by the data of CMB+BAO.

We summarize the constraints on the parameters as follows: From the CMB+BAO data, we obtain

$$-0.0293 \leq c_2 \leq 0.1032,$$

$$\frac{\alpha M_{\text{PL}}^2}{\mu^4} \leq -9.690 \quad \text{or} \quad \frac{\alpha M_{\text{PL}}^2}{\mu^4} \geq 34.13, \\ -0.03065 \leq \beta_2 \leq 0.08937$$

at 95% C.L., while from the EMPRESS VIII data, we find

$$0.0239 \leq c_2 \leq 0.1108, \\ -41.84 \leq \frac{\alpha M_{\text{PL}}^2}{\mu^4} \leq -9.025, \\ 0.02307 \leq \beta_2 \leq 0.0950$$

at 95% C.L.

For the cuscuton gravity model, in Fig. 1, we show the (white) region allowed by EMPRESS VIII as well as (CMB + BAO) at 95% C.L. in the α - μ^2 plane. The red shaded regions are excluded by EMPRESS VIII with the BBN, while CMB + BAO data excludes the blue shaded region at 95% C.L.




FIG. 1. Regions excluded by EMPRESS VIII with the BBN (red) and CMB + BAO (blue) at 95% C.L. in the α - μ^2 plane. Here μ^2 and α are plotted in unit of the reduced Planck mass M_{PL} . $\mu = 0$, which corresponds to general relativity with a cosmological constant, is excluded by EMPRESS VIII with the BBN.

One may wonder about the negative value of α , because the potential is unbounded from below. However, it does not give any instability because the scalar field ϕ is non-dynamical.

V. CONCLUSION

We have studied a cosmological model in the cuscuton gravity theory with a quadratic potential $V = V_0 + \frac{1}{2} \alpha \phi^2$ to resolve the anomaly of the observational ⁴He abundance reported by the EMPRESS collaboration. This model is equivalent not only to the Λ CDM theory with a quadratic potential with the dimensionless coefficient β_2 but also to the Einstein-aether theory with only one coupling constant c_2 .

About the mass squared parameter α in the cuscuton gravity theory, we have obtained the allowed region as

¹ This model is a very special case of the Einstein-aether theory because aether field modes do not appear in the perturbation equations[22], which is consistent with the fact that the system has only two degrees of freedom.

$-41.84 \leq (\alpha/M_{\text{PL}}^2)(\mu/M_{\text{PL}})^{-4} \leq -9.690$, or equivalently $0.02307 \leq \beta_2 \leq 0.08937$ in the Λ CDM theory and $0.0239 \leq c_2 \leq 0.1032$ in the Einstein-aether theory. General relativity is excluded in the present approach.

Thus, this could suggest the need for modified gravity theories such as the cuscuton gravity theory with a quadratic potential, which can be regarded as the low-energy Hořava-Lifshitz gravity. In future, $N_{\nu, \text{eff}}$ can be measured more precisely by planned observations of 21cm + CMB down to errors of $\Delta N_{\nu, \text{eff}} \sim \mathcal{O}(10^{-2})$ [8]. Then, we will test the gravitational constant to be a precision within the order of $\Delta G_{\text{N}}/G_{\text{N}} \sim \mathcal{O}(10^{-3})$, which might give a hint of quantum gravity.

ACKNOWLEDGMENTS

We thank Niyaesh Afshordi and Shinji Mukohyama for useful discussions. K.M. also acknowledges the Yukawa Institute for Theoretical Physics at Kyoto University, where the present work was completed during the Visitors Program of FY2022. This work was supported in part by JSPS KAKENHI Grant Numbers JP17H01131 (K.K.), JP17H06359, JP19K03857 (K.M.), and by MEXT KAKENHI Grant Numbers JP20H04750, JP22H05270 (K.K.).

-
- [1] A. Matsumoto, M. Ouchi, K. Nakajima, M. Kawasaki, K. Murai, K. Motohara, Y. Harikane, Y. Ono, K. Kushibiki and S. Koyama, *et al.* [arXiv:2203.09617 [astro-ph.CO]].
- [2] G. Mangano, G. Miele, S. Pastor, T. Pinto, O. Pisanti and P. D. Serpico, Nucl. Phys. B **729**, 221-234 (2005) [arXiv:hep-ph/0506164 [hep-ph]].
- [3] P. F. de Salas and S. Pastor, JCAP **07**, 051 (2016) [arXiv:1606.06986 [hep-ph]].
- [4] K. Kohri, M. Kawasaki and K. Sato, Astrophys. J. **490**, 72-75 (1997) [arXiv:astro-ph/9612237 [astro-ph]].
- [5] M. Kawasaki, F. Takahashi and M. Yamaguchi, Phys. Rev. D **66**, 043516 (2002) [arXiv:hep-ph/0205101 [hep-ph]].
- [6] M. Kawasaki and K. Murai, [arXiv:2203.09713 [hep-ph]].
- [7] X. D. Shi, G. M. Fuller and K. Abazajian, Phys. Rev. D **60**, 063002 (1999) [arXiv:astro-ph/9905259 [astro-ph]].
- [8] K. Kohri, Y. Oyama, T. Sekiguchi and T. Takahashi, JCAP **09**, 014 (2014) [arXiv:1404.4847 [astro-ph.CO]].
- [9] N. Afshordi, D. J. H. Chung and G. Geshnizjani, Phys. Rev. D **75** (2007), 083513 [arXiv:hep-th/0609150 [hep-th]].
- [10] N. Afshordi, D. J. H. Chung, M. Doran and G. Geshnizjani, Phys. Rev. D **75** (2007), 123509 [arXiv:astro-ph/0702002 [astro-ph]].
- [11] A. Iyonaga, K. Takahashi and T. Kobayashi, JCAP **12** (2018), 002 [arXiv:1809.10935 [gr-qc]].
- [12] A. Iyonaga, K. Takahashi and T. Kobayashi, JCAP **07** (2020), 004 [arXiv:2003.01934 [gr-qc]].
- [13] K. Maeda and S. Panpanich, Phys. Rev. D **105**, 104022 (2022) [arXiv:2202.04908 [gr-qc]].
- [14] J. Gleyzes, D. Langlois, F. Piazza and F. Vernizzi, Phys. Rev. Lett. **114** (2015) no.21, 211101 [arXiv:1404.6495 [hep-th]].
- [15] A. De Felice, A. Doll and S. Mukohyama, JCAP **09** (2020), 034 doi:10.1088/1475-7516/2020/09/034 [arXiv:2004.12549 [gr-qc]].
- [16] A. De Felice, K. Maeda, S. Mukohyama, M. Pookkillath, Phys. Rev. D (to appear, 2022) [arXiv:22204.08294].
- [17] S. Mukohyama and K. Noui, JCAP **07**, 049 (2019) [arXiv:1905.02000 [gr-qc]].
- [18] M. Ballardini, F. Finelli and D. Sapone, [arXiv:2111.09168 [astro-ph.CO]].
- [19] N. Afshordi, Niayesh, Phys. Rev. D **80** (2009), 081502
- [20] T. Jacobson and D. Mattingly, Phys. Rev. D **64**, 024028.
- [21] J. Bhattacharyya, A. Coates, M. Colombo, A.E. Gümrükçüoğlu and T.P. Sotiriou Phys. Rev. D **97**, 064020 (2018)
- [22] S. Mukohyama, private communication.