

# A stable, efficient scheme for $\mathcal{C}^n$ function extensions on smooth domains in $\mathbb{R}^d$

Charles Epstein<sup>a</sup>, Shidong Jiang<sup>a</sup>

<sup>a</sup>Center for Computational Mathematics, Flatiron Institute, Simons Foundation, New York, New York 10010

---

## Abstract

A new scheme is proposed to construct a  $\mathcal{C}^n$  function extension for smooth functions defined on a smooth domain  $D \in \mathbb{R}^d$ . Unlike the PUX scheme, which requires the extrapolation of the volume grid via an expensive ill-conditioned least squares fitting, the scheme relies on an explicit formula consisting of a linear combination of function values in  $D$ , which only extends the function along the normal direction. To be more precise, the  $\mathcal{C}^n$  extension requires only  $n+1$  function values along the normal directions in the original domain and ensures  $\mathcal{C}^n$  smoothness by construction. When combined with a shrinking function and a smooth window function, the scheme can be made stable and robust for a broad class of domains with complex smooth boundary.

*Keywords:* function extension, the Vandermonde matrix, Chebyshev nodes, complex geometry, smooth domain

---

## 1. Introduction

In a classic paper by Robert Seeley [6], a simple formula is given for the  $\mathcal{C}^\infty$  extension of a  $\mathcal{C}^\infty$  function defined in a half space. To be more precise, let  $y \in \mathbb{R}^{d-1}$ ,  $x \in \mathbb{R}^1$ ,  $\mathbb{R}_+^d = \mathbb{R}^{d-1} \times \{x \geq 0\}$ , then  $\mathcal{C}^\infty(\mathbb{R}_+^d)$  consists of infinitely differentiable functions defined in  $\mathbb{R}^{d-1} \times (0, \infty)$  whose derivatives have continuous limits as  $x \rightarrow 0^+$ . For  $x < 0$ , define the extended function

$$E_\infty[f](y, x) = \sum_{j=0}^{\infty} w_j f(y, -t_j x) \phi(-t_j x), \quad (1)$$

where  $\{t_j\} \subset (0, \infty)$  is an unbounded, strictly increasing sequence, and  $\phi$  is a  $\mathcal{C}^\infty$  window function on  $\mathbb{R}^1$  with  $\phi(x) = 1$  for  $0 \leq x \leq 1$ , and  $\phi(x) = 0$  for  $x \geq 2$ . For any  $x < 0$ , the sum in (1) is finite and therefore  $E_\infty[f]$  has the same smoothness as the original function  $f$ . In order to ensure the smoothness of the extension across the boundary, one needs to require that the values of  $E_\infty[f]$  and  $f$  and all of their derivatives match  $x = 0$ . This leads to the following infinite system of linear equations for  $\{w_j\}$

$$\sum_{j=0}^{\infty} w_j t_j^i = (-1)^i, \quad i = 0, 1, \dots, \quad (2)$$

i.e., the weight vector  $w$  is the solution to an infinite Vandermonde system when the node vector  $t = (t_0, t_1, \dots)$  is given. Using  $t_j = 2^j$ , Seeley showed that  $\sum_{j=0}^{\infty} |w_j| |t_j|^i < \infty$  for all  $i \geq 0$ .

---

*Email addresses:* cepstein@flatironinstitute.org (Charles Epstein), sjiang@flatironinstitute.org (Shidong Jiang)

Thus, the extension operator  $E_\infty$  is actually a continuous linear operator from  $C^\infty(\mathbb{R}_+^d)$  to  $C^\infty(\mathbb{R}^d)$ .

Here we examine the extension formula (1) from the perspective of numerical computation. Function extensions have broad applications in numerical analysis and scientific computing, especially to boundary value problems for partial differential equations where the boundary does not have simple geometry. Most existing numerical schemes rely on a data fitting procedure using a choice of basis functions such as Fourier series, polynomials, radial basis functions, etc. The data fitting procedure is usually carried out on a volume grid to ensure the smoothness of the extended function in all variables, which is expensive for problems in two and higher dimensions. Another observation is that these schemes depend on the original data linearly, regardless of the basis functions chosen to fit the data. In other words, the extension operator is a linear operator. Obviously, any  $C^n$  extension must satisfy the condition that the extended function should match the original function in its value and derivatives up to order  $n$  on the boundary.

For applications to numerical analysis it is more appropriate to consider the finite differentiability analogue of  $E_\infty$ , given by

$$E_n[f](y, x) = \sum_{j=0}^n w_j f(y, -t_j x) \phi(-t_j x), \quad (3)$$

where now we have the finite Vandermonde system:

$$\sum_{j=0}^n w_j t_j^i = (-1)^i, \quad i = 0, 1, \dots, n. \quad (4)$$

One nice feature of the extension formulæ (1) and (3) is that the extension acts along one direction only, while the smoothness in the other directions is automatically guaranteed by construction. This feature can be extended to domains  $D \subset \mathbb{R}^d$ , with  $\partial D$  a smooth embedded hypersurface. For  $y \in \partial D$ , let  $\mathbf{n}_y$  denote the outer unit normal vector to  $\partial D$  at  $y$ . The tubular neighborhood theorem, see [1], ensures that for some  $\epsilon > 0$  the map from  $\partial D \times (-\epsilon, \epsilon)$  given by

$$(y, x) \mapsto y + x \mathbf{n}_y \quad (5)$$

is a diffeomorphism onto its image. For such domain  $D \subset \mathbb{R}^d$ , we can extend the definition of  $E_n$  by setting

$$E_n[f](y + x \mathbf{n}_y) = \sum_{j=0}^n w_j f(y - t_j x \mathbf{n}_y), \quad (6)$$

where  $y$  is a point on  $\partial D$ . The window function is dropped for now, but will usually be needed in real applications. Once again,  $E_n[f]$  has the same order of smoothness as  $f$  outside  $D$  and along the tangential directions on  $\partial D$  by construction. Thus, one only needs to ensure the continuity of the function value and its normal derivatives up to order  $n$  on  $\partial D$  in order for the extension to be in  $C^n$  in a small neighborhood of  $D$ . This leads to the following Vandermonde linear system for the nodes and weights

$$\begin{bmatrix} 1 & 1 & \cdots & 1 \\ t_0 & t_1 & \cdots & t_n \\ \vdots & \vdots & \cdots & \vdots \\ t_0^n & t_1^n & \cdots & t_n^n \end{bmatrix} \begin{bmatrix} w_0 \\ w_1 \\ \vdots \\ w_n \end{bmatrix} = \begin{bmatrix} 1 \\ -1 \\ \vdots \\ (-1)^n \end{bmatrix}, \quad (7)$$

which will be written as  $Aw = c$  for short. A very nice feature of this approach is that the nodes and weights are independent of the domain,  $D$ . By the nature of the extension operator, we need to have  $t_j \geq 0$  for  $j = 0, \dots, n$  and without loss of generality, we assume that they are arranged in increasing order, i.e.,  $0 \leq t_0 < t_1 < \dots < t_n$ .

## 2. Solution to the Vandermonde system (7)

It is well known that Vandermonde matrices are generally ill-conditioned (as is the problem of function extrapolation). In [5], it is shown that the condition number (in the maximum norm) of the Vandermonde matrix is bounded from below by  $(n+1)2^n$  when all nodes are nonnegative. The “quality” of the extension operator given by (6) is determined by the magnitude of the weights  $\{w_j\}$ . A good measure of this quality is the  $\ell^1$ -norm of  $w$  :

$$\|w\|_1 = \sum_{i=0}^n |w_i| \quad (8)$$

If the weights are calculated by solving (7) numerically, then the condition number of the Vandermonde matrix will affect the accuracy of the weights. However, (7) can be solved analytically by an elementary method, which we now present. Denote by  $B$  the inverse of the matrix  $A$ . We then have

$$\sum_{j=0}^n b_{ij} t_k^j = \delta_{ik}, \quad (9)$$

Consider the polynomial  $p_i(x) = \sum_{j=0}^n b_{ij} x^j$ . Then (9) is equivalent to

$$p_i(t_k) = \delta_{ik}. \quad (10)$$

Recall that the Lagrange basis functions  $\{l_i(x)\}$  for interpolation through the nodes  $\{t_k, k = 0, \dots, n\}$  are given by

$$l_i(x) = \prod_{m=0, m \neq i}^n \frac{x - t_m}{t_i - t_m}, \quad (11)$$

and the interpolant is then

$$Q(x) = \sum_{i=0}^n f_i l_i(x). \quad (12)$$

It is straightforward to check that  $l_i$  satisfies the condition (10). Thus,

$$p_i(x) = l_i(x), \quad (13)$$

and

$$\begin{aligned} w_i &= \sum_{j=0}^n b_{ij} (-1)^j = p_i(-1) = l_i(-1) \\ &= (-1)^i \prod_{m=0}^{i-1} \frac{1+t_m}{t_i-t_m} \cdot \prod_{m=i+1}^n \frac{1+t_m}{t_m-t_i}. \end{aligned} \quad (14)$$

That is,  $w_i$  is the value of the  $i$ th Lagrange basis function evaluated at  $-1$ .

### 3. Choice of interpolation nodes

As noted above, the quality of the function extension (6) depends critically on the magnitude of the weights  $\{w_i\}$ , which in turn is determined by the interpolation nodes  $t_i$  explicitly via the formula (14). To fix our discussion, let us assume that the nodes lie on the interval  $[0, a]$  and use the 1-norm to measure the quality of function extensions. In other words, we would like to choose the interpolation nodes  $t_i \in [0, a]$  such that  $\|w\|_1$  is minimized.

Combining (12) and (14), we obtain

$$\begin{aligned} \|w\|_1 &= \sum_{i=0}^n |w_i| \\ &= \sum_{i=0}^n (-1)^i l_i(-1) \\ &= P(-1), \end{aligned} \tag{15}$$

where  $P(x) = \sum_{i=0}^n (-1)^i l_i(x)$ .

**Definition 1.** A polynomial  $P$  is said to satisfy the *equioscillation property of modulus 1* on  $[0, a]$  if there exist  $t_i$  with  $0 \leq t_0 < t_1 < \dots < t_n \leq a$  such that  $P(t_i) = (-1)^i$  for  $i = 0, \dots, n$ .

**Lemma 1.** If  $P(x)$  is a polynomial of degree  $n$ , satisfying the equioscillation property of modulus 1, then its  $n$  roots  $x_i$  ( $i = 1, \dots, n$ ) satisfy the property  $t_0 < x_1 < t_1 < x_2 < t_2 < \dots < x_n < t_n$ .

*Proof.* This follows from the intermediate value theorem and the fact that  $P(x)$  changes sign alternatingly at  $t_i$  ( $i = 0, \dots, n$ ).  $\square$

**Lemma 2.** If  $P(x)$  is a polynomial of degree  $n$ , satisfying the equioscillation property of modulus 1, then  $P(x) > 0$  decreases monotonically for  $x < t_0$ .

*Proof.* This follows from Lemma 1 and the fact that  $P(t_0) = 1$ . That is,  $P(x) = c \prod_{i=1}^n (x_i - x)$  for some  $c > 0$ .  $\square$

It is easy to see that the choice of optimal interpolation nodes for the purpose of function extension is equivalent to the following problem:

*Find the minimal value of  $P(-1)$  among all polynomials of degree at most  $n$  that satisfy the equioscillation property of modulus 1 on  $[0, a]$ .*

Denote the minimal value by  $m_a$ , the corresponding polynomial by  $p^*(x)$ , and the associated nodes by  $\{t_j^*\}$ , respectively. In the following lemmas we give the properties of the optimal nodes and  $p^*$  itself.

**Lemma 3.**  $t_0^* = 0$ .

*Proof.* Suppose  $t_0^* > 0$ . Then  $q(x) = p^*(x + t_0^*)$  satisfies the alternating property  $q(t_i^* - t_0^*) = p^*(t_i^*) = (-1)^i$ . Furthermore,  $q(-1) = p^*(-1 + t_0^*) < p^*(-1)$  by Lemma 2, which leads to a contradiction.  $\square$

**Lemma 4.** If  $n > 0$ , then  $t_n^* = a$ .

*Proof.* Suppose  $t_n^* < a$ . Consider the polynomial  $q(x)$ , which is obtained by replacing  $t_n^*$  by  $a$  and keep all other nodes unchanged. We claim that  $q(-1) < p^*(-1)$ . This can be seen from (14), (15), and the facts that both  $\frac{1}{t_n - t_m}$  and  $\frac{1+t_n}{t_n - t_i} = 1 + \frac{1+t_i}{t_n - t_i}$  decrease as  $t_n$  increases. Hence the contradiction.  $\square$

**Lemma 5.** *If  $a_1 > a_2$ , then  $m_{a_1} < m_{a_2}$ .*

*Proof.* By the definition of the problem, we have  $m_{a_1} \leq m_{a_2}$  since the space is larger as  $a$  increases. The strict inequality is achieved by Lemma 4.  $\square$

**Lemma 6.**  $|p^*(x)| \leq 1$  for  $x \in (0, a)$ .

*Proof.* Suppose first that  $p^*(t^\circ) > 1$  at some point  $t^\circ \in (0, a)$ . Let  $t_j^*$  be the closest node to  $t^\circ$  such that  $p^*(t_j^*) = 1$ . It is clear that  $j$  has to be even and  $t^\circ < t_{j+1}^*$  by the alternating property. Consider the polynomial  $q(x)$  which is obtained by replacing  $t_j^*$  with  $t^\circ$ . Then  $p^*(t_i^*) - q(t_i^*) = 0$  for  $i \neq j$ . That is,

$$p^*(x) - q(x) = c \prod_{i=0}^{j-1} (x - t_i^*) \cdot \prod_{i=j+1}^n (t_i^* - x), \quad (16)$$

Furthermore, since  $p^*(t^\circ) - q(t^\circ) = p^*(t^\circ) - 1 > 0$ , we must have  $c > 0$ . It then follows that

$$\begin{aligned} p^*(-1) - q(-1) &= c \prod_{i=0}^{j-1} (-1 - t_i^*) \cdot \prod_{i=j+1}^n (t_i^* + 1) \\ &= c \prod_{i=0}^{j-1} (1 + t_i^*) \cdot \prod_{i=j+1}^n (t_i^* + 1) > 0 \end{aligned} \quad (17)$$

due to the facts that  $c > 0$  and  $j$  is even. Hence the contradiction.

Suppose that  $p^*(t^\circ) < -1$ . Then a similar argument leads to (17) again by the facts that  $c < 0$  and  $j$  is odd.  $\square$

**Lemma 7.**  $|p^*(x)| < 1$  for  $x \in (0, a)$  and  $x \neq t_i$ ,  $i = 0, 1, \dots, n$ .

*Proof.* This follows from Lemma 6, since otherwise there must be a point at which  $|p^*|$  is greater than 1.  $\square$

**Lemma 8.**  $p^{*'}(t_i^*) = 0$  for  $i = 1, \dots, n - 1$ .

*Proof.* By Lemmas 6 and 7,  $p^*(x)$  achieves its interior extreme values at  $t_i^*$ . Hence, its first derivative vanishes at those interior nodes.  $\square$

The following lemma is key to the uniqueness.

**Lemma 9.** *If  $p_1(x)$  and  $p_2(x)$  satisfy the equioscillation property of modulus 1 on  $[0, a]$  and  $|p_1(x)| \leq 1$ ,  $|p_2(x)| \leq 1$ , then  $p_1(x) \equiv p_2(x)$ .*

*Proof.* Suppose that  $p_1$  is not identically equal to  $p_2$ . Then there exists a point  $\xi < 0$  such that  $p_1(\xi) \neq p_2(\xi) > 0$ . Without loss of generality, assume that  $0 < p_1(\xi) < p_2(\xi)$ . Consider the polynomial

$$q(x) = p_1(x) - \frac{p_1(\xi)}{p_2(\xi)} p_2(x). \quad (18)$$

By construction,  $q(\xi) = 0$ . Furthermore,

$$\left| p_2(x) \frac{p_1(\xi)}{p_2(\xi)} \right| < 1, \quad x \in [0, a], \quad (19)$$

since  $p_1(\xi) < p_2(\xi)$  and  $|p_2(x)| \leq 1$  for  $x \in [0, a]$ . Thus,  $q$  has the same signs as  $p_1$  at its equioscillation nodes. By the intermediate value theorem,  $q$  has  $n$  zeros on  $(0, a)$ . Together with the zero at  $\xi$ ,  $q$  has  $n + 1$  zeros. Since  $q$  is a polynomial of degree at most  $n$ ,  $q$  has to be identically equal to zero everywhere, which leads to a contradiction.  $\square$

Using these results we can now give an explicit formula for  $p^*$  and  $\{t_j^*\}$ :

**Theorem 1.**  $p^*$  is unique and  $p^*(x) = T_n(1 - 2x/a)$ , where  $T_n$  is the Chebyshev polynomial of degree  $n$ .

*Proof.* The uniqueness follows from Lemma 9 and the existence follows from explicit construction. Namely, it is straightforward to check that  $T_n(1 - 2x/a)$  satisfies the equioscillation property of modulus 1 and  $|T_n(1 - 2x/a)| \leq 1$  on  $[0, a]$ .  $\square$

Since we only used the fact that  $-1$  is outside the interval  $[0, a]$  in our proofs and  $[0, a]$  can be translated to any interval  $[a, b]$ , we actually showed the following statement is true.

**Corollary 1.** Suppose that  $p(x)$  is a polynomial of degree at most  $n$  satisfying the property  $p(t_i) = (-1)^i$  or  $p(t_i) = (-1)^{i+1}$  for  $a \leq t_0 < t_1 < \dots < t_n \leq b$ . Then

$$|p(x)| \geq \left| T_n \left( \frac{2}{b-a} \left( x - \frac{a+b}{2} \right) \right) \right|, \quad x \notin [a, b]. \quad (20)$$

In other words, the translated and rescaled Chebyshev polynomials have the *least growth* outside  $[a, b]$  among all polynomials with the equioscillation property of modulus 1 on  $[a, b]$ . Using the explicit expressions of  $T_n$

$$T_n(x) = \begin{cases} \cos(n \arccos(x)), & |x| \leq 1 \\ \frac{1}{2} \left( (x - \sqrt{x^2 - 1})^n + (x + \sqrt{x^2 - 1})^n \right), & |x| \geq 1, \end{cases} \quad (21)$$

We may calculate the optimal weights explicitly. Recall that  $T_n(\cos \frac{i\pi}{n}) = (-1)^i$ ; consider the function

$$\phi(x) = \prod_{i=0}^n (x - t_i). \quad (22)$$

Then

$$l_i(x) = \frac{\phi(x)}{(x - t_i)\phi'(t_i)}, \quad (23)$$

and

$$w_i^* = l_i^*(-1) = -\frac{\phi^*(-1)}{(1 + t_i)\phi^{*'}(t_i)}. \quad (24)$$

We first work on the standard interval  $[-1, 1]$ . By Lemma 8, the  $n - 1$  interior nodes are the zeros of  $T_n'(x)$ . Thus,  $\phi(x) = c(x^2 - 1)T_n'(x)$  for the interval  $[-1, 1]$ . Back to  $[0, a]$ , we have

$$\phi^*(x) = x(x - a)T_n'(2x/a - 1), \quad (25)$$

and

$$\phi^{*'}(x) = \frac{2}{a}x(x - a)T_n''(2x/a - 1) + (2x - a)T_n'(2x/a - 1), \quad (26)$$

where the irrelevant constant  $c$  is dropped.

Table 1: Condition number of function extension formula (6) in  $l_1$  norm. The first row lists the order of the function extension. The first column lists the size of the interval.

$a \backslash n$	2	3	4	5	6	7	8	9
2	7.0	26.0	97	362	1351	5042	18 817	70 226
4	3.5	9.0	24	62	161	422	1104	2889
6	2.6	5.5	12	27	59	131	290	642
8	2.1	4.1	8	16	32	64	128	256
10	1.9	3.3	6	11	21	39	73	135
12	1.7	2.9	5	9	15	27	48	84
14	1.6	2.5	4	7	12	20	34	58
16	1.5	2.3	4	6	10	16	26	43

**Corollary 2.** *The optimal nodes on  $[0, a]$  for function extension formula (6) are Chebyshev nodes of the second kind shifted and scaled to the interval  $[0, a]$ :*

$$t_i^* = \frac{a}{2} \left( 1 - \cos \left( \frac{i\pi}{n} \right) \right), \quad i = 0, \dots, n. \quad (27)$$

And the associated optimal weights are

$$w_i^* = (-1)^i \frac{C_n(a)}{(1 + \delta_{i0} + \delta_{in})na(1 + t_i^*)}, \quad i = 0, \dots, n, \quad (28)$$

where

$$C_n(a) = \frac{1+a}{\sqrt{x_0^2-1}} \left( \left( x_0 + \sqrt{x_0^2-1} \right)^n - \left( x_0 - \sqrt{x_0^2-1} \right)^n \right), \quad x_0 = 1 + \frac{2}{a}. \quad (29)$$

Finally, the  $l_1$  norm of the associate optimal weights is

$$\begin{aligned} \|w^*\|_1 &= \sum_{i=0}^n |w_i^*| = T_n(1 + 2/a) \\ &= \frac{1}{2} \left( \left( 1 + \frac{2}{a} - 2\sqrt{\frac{1}{a} + \frac{1}{a^2}} \right)^n + \left( 1 + \frac{2}{a} + 2\sqrt{\frac{1}{a} + \frac{1}{a^2}} \right)^n \right). \end{aligned} \quad (30)$$

The function extension is, as expected, exponentially ill-conditioned. As  $a$  increases, the condition number decreases, due to the fact that  $-1$  is relatively closer to the origin with respect to the interval  $[0, a]$ . Since the extension formula (6) only imposes the minimal conditions to ensure  $C^n$  continuity across the boundary, (30) can be viewed as the *intrinsic condition number* of any linear function extension scheme.

#### 4. Stabilizing techniques

Table 1 lists condition numbers of the function extension formula (6), where the first row lists the extension order  $n$ , and the first column lists the size of the interval,  $D$ , where  $f$  is defined. While the condition numbers do increase exponentially fast as  $n$  increases, we observe that their numerical values are not very large, especially when  $a$  is sufficiently large. For finite difference/finite element methods,  $n$  is usually less than 5 and the condition number for  $a = 2$  seems rather benign.

#### 4.1. Shrinking function

For many practical applications, function extensions are carried out in a small neighborhood of  $D$  and the extended function is rolled off to zero via a window function. We first point out that we could modify the extension formula so that it uses the function values on the interval, say,  $[0, 1]$  instead of  $[0, a]$  with the weights and the extension order unchanged. To be more precise, we modify the extension formula as follows:

$$E_n[f](y + x\mathbf{n}) = \sum_{j=0}^n w_j f(y - t_j \psi(x)\mathbf{n}), \quad (31)$$

where  $\psi$  maps  $[0, 1]$  to  $[0, \delta]$ . Furthermore, we require

$$\begin{aligned} \psi(0) &= 0, \\ \psi'(0) &= 1, \\ \psi^{(j)}(0) &= 0, \quad j = 2, \dots, n. \end{aligned} \quad (32)$$

With the property (32) and the chain rule, it is easy to see that the derivatives of  $E_n[f]$  at  $x = 0$  up to order  $n$  are unchanged, thus ensuring  $\mathcal{C}^n$  smoothness of the extension. It is not hard to construct such a function. For example, one may choose

$$\psi(x) = s^{-1}(x), \quad s(x) = x + (x/\delta)^{n+1}(1 - \delta). \quad (33)$$

It is easy to see that (31) requires function values on the interval  $[0, a\delta]$ . Thus, if we set  $\delta = \frac{1}{a}$ , then it requires function values only on the interval  $[0, 1]$ .

##### 4.1.1. Evaluation of the shrinking function

The shrinking function  $\psi$  can be calculated by solving  $s(y) = x$  numerically via a root finding scheme. In practice, we find the secant method with initial guesses  $y_0 = 0$ ,  $y_1 = \delta$  always converges very rapidly. An alternative way is to build a piecewise polynomial approximation for  $\psi$  to allow even faster evaluation. The second method requires a precomputation step. But subsequently  $\psi(x)$  can be evaluated rapidly by evaluating a low-degree interpolating polynomial on each subinterval. In MATLAB, such piecewise polynomial approximation can be built using CHEBFUN [2].

##### 4.1.2. Effects of the shrinking function

When combined with large  $a$ , the shrinking function can lower the condition number of the function extension and the maximum norm of the extended function very effectively, without increasing the interpolation interval. On the other hand, the introduction of the shrinking function introduces high-frequency components into the extended function. One therefore needs to carefully balance these two effects in order to achieve better overall performance for the whole numerical scheme. When the underlying scheme is FFT based, it seems that the introduction of the shrinking function increases the number of equispaced grid points by a large factor. Hence its usage is not recommended in this case. When the underlying scheme is based on adaptive refinement, the introduction of the shrinking function increases the total number of intervals/boxes only mildly. In this case, it might be advantageous to use the shrinking function to lower the condition number and the magnitude of the extended part.



#### 4.2. Window function

We use a window function to roll the extension smoothly to zero. The choice of the window function is critical to the quality of the function extension. In practice, we find it is better to have two window functions in the extension formula. That is,

$$E_n[f](x) = \left( \sum_{i=0}^n w_i f(-t_i x) \phi_l(-t_i x) \right) \phi_g(-x), \quad (34)$$

where  $\phi_l$  is a *local* window function acting along  $f$ , and  $\phi_g$  is a global window function acting on the whole extended function. In order to keep the magnitude of the extended part under control,  $\phi$  should decay rapidly while avoiding introducing high frequency content. It is difficult to determine the optimal window function. In practice, we have observed that the following window function performs rather well.

$$\phi(x, r_0, r_1) = \begin{cases} 1, & x \leq r_0 \\ \frac{1}{2} \operatorname{erfc} \left( \frac{12}{\pi} \arcsin \left( \frac{x - (r_0 + r_1)/2}{r_1 - r_0} \right) \right), & x \in (r_0, r_1) \\ 0, & x \geq r_1, \end{cases} \quad (35)$$

where  $\operatorname{erfc}(x) = \frac{1}{\sqrt{\pi}} \int_x^\infty e^{-t^2} dt$  is the *complementary error function*, and  $0 < r_0 < r_1$  are two parameters to be specified.

### 5. Numerical experiments

We now demonstrate the quality of the function extension formula (34). We use  $D = [0, 0.5]$  as the interval where the original function is defined, and  $[-0.25, 0]$  as the interval to which the function is extended. We set  $\phi_g(x) = \phi(x, 10^{-6}, 0.25)$  and  $\phi_l(x) = \phi(x, 0.2, 1)$ . We measure the quality of the function extension via the following quantities.

- (a)  $\kappa = \|E_n[f]\|_\infty / \|f\|_\infty$ , i.e.,  $\kappa$  is the ratio of the maximum norm of  $E_n[f]$  to that of the original function  $f$ . It is clear that the lower the value of  $\kappa$ , the better quality of  $E_n[f]$ .
- (b) the power spectrum of the function

$$F(x) = \begin{cases} E_n[f](x), & x \in [-0.25, 0] \\ f(x)\phi_g(x), & x \in [0, 0.25], \end{cases} \quad (36)$$

We put a global window function on the original function as well so that the FFT can be used to calculate the power spectrum directly and the irrelevant behavior of the original function at  $x = 0.25$  does not pollute the power spectrum. It is clear that the narrower the power spectrum, the better quality of  $E_n[f]$ .

- (c) the number of chunks of the function via adaptive refinement

$$G(x) = \begin{cases} E_n[f](x), & x \in [-0.25, 0] \\ f(x), & x \in [0, 0.5]. \end{cases} \quad (37)$$

It is clear that the fewer number of chunks, the better. Here we choose the asymmetrical interval  $[-0.25, 0.5]$  so that the extension point, i.e., the origin, is not the middle point of the interval and thus the smoothness across the origin can be captured by the adaptive refinement.




Figure 1: Numerical results for extending the function  $f_1(x) = \frac{0.04}{0.04+x^2}$  from  $[0, 0.5]$  to  $[-0.25, 0]$ . The extension order is  $n = 9$  and the extension parameter  $a$  is set to 2. Left: the original function (in blue) and the extended function (in red), xticks show the endpoints of the chunks via adaptive refinement, i.e., three chunks are needed to resolve the function on the whole interval.  $\|E_n[f]\|_\infty/\|f\|_\infty = 1$ . Middle: the power spectrum of  $F(x)$  using 4000 points. Right: closeup of the power spectrum of  $F(x)$  using 200 points.

Similar to the widely used RBF-QR [3] and the PUX [4] schemes for function extensions, we use the following functions of increasing complexity in our numerical experiments.

$$\begin{aligned}
 f_1(x) &= \frac{0.04}{0.04+x^2}, \\
 f_2(x) &= \sin(2\pi(x+1)^2), \\
 f_3(x) &= (x^2-1)e^{-20x^2}, \\
 f_4(x) &= J_0(25(x+0.4)), \\
 f_5(x) &= \cos(7\arccos(4x-1)),
 \end{aligned} \tag{38}$$

where  $J_0$  is the first kind Bessel function of order zero.




Figure 2: Same as Figure 1, but for  $f_2(x) = \sin(2\pi(x+1)^2)$ .  $\|E_n[f]\|_\infty/\|f\|_\infty \approx 0.83$ .




Figure 3: Same as Figure 1, but for  $f_3(x) = (x^2-1)e^{-20x^2}$ .  $\|E_n[f]\|_\infty/\|f\|_\infty = 1$ .

Figures 1–4 present numerical results for the first four functions in (38). It turns out that the last function, i.e., the Chebyshev polynomial of degree 7 is the most difficult one for function extension. Figure 5 present numerical results for two experiments. The top row shows the




Figure 4: Same as Figure 1, but for  $f_4(x) = J_0(25(x+0.4))$ . Here  $n = 11$  and  $a = 2$ .  $\|E_n[f]\|_\infty/\|f\|_\infty \approx 1.78$ .

function extension with  $n = 9$  and  $a = 2$ . Though the power spectrum remains as good as the other four functions, the extended function has much larger magnitude. The bottom row shows the function extension when a shrinking function is used to control the magnitude of the extended function. Here  $n = 9$ ,  $a = 20$ , and the parameter for the shrinking function is  $\delta = 1/(2a)$ . As compared with the top row, the magnitude of the extended part is reduced by a large amount. However, the power spectrum deteriorates significantly, rendering the scheme much less efficient for the FFT-based schemes. However, the number of chunks is increased only mildly from 3 to 6 when adaptive refinement is used, which indicates that the second strategy might be useful for adaptive algorithms such as the volume fast multipole methods.




Figure 5: Same as Figure 1, but for  $f_5(x) = \cos(7 \arccos(4x-1))$ . Top:  $n = 9$ ,  $a = 2$ , and  $\|E_n[f]\|_\infty/\|f\|_\infty \approx 218$ . Bottom:  $n = 9$ ,  $a = 20$ , and  $\|E_n[f]\|_\infty/\|f\|_\infty \approx 7.365$ . Here a shrinking function  $\psi(x)$  with the parameter  $\delta = 1/(2a)$  is applied so that only function values on  $[0, 1/8]$  are used for extension.

## Acknowledgments

The authors would like to thank Travis Askham at New Jersey Institute of Technology for helpful discussions.

## References

- [1] R. Bott and L. Tu. *Differential forms in algebraic topology*, volume 82 of *GTM*. Springer-Verlag, 1982.

- [2] T. A. Driscoll, N. Hale, and L. N. Trefethen. *Chebfun guide*. Pafnuty Publications, Oxford, 2014.
- [3] B. Fornberg, E. Larsson, and N. Flyer. Stable computations with gaussian radial basis functions. *SIAM Journal on Scientific Computing*, 33(2):869–892, 2011.
- [4] F. Fryklund, E. Lehto, and A.-K. Tornberg. Partition of unity extension of functions on complex domains. *Journal of Computational Physics*, 375:57–79, 2018.
- [5] W. Gautschi and G. Inglese. Lower bounds for the condition number of vandermonde matrices. *Numerische Mathematik*, 52(3):241–250, 1987.
- [6] R. T. Seeley. Extension of  $C^\infty$  functions defined in a half space. *Proceedings of the American Mathematical Society*, 15(4):625–626, 1964.