

Sequential deconfinement and self-dualities in $4d \mathcal{N}=1$ gauge theories

Stephane Bajeot¹ Sergio Benvenuti²

¹*SISSA, Via Bonomea 265, 34136 Trieste, Italy*

²*INFN, Sezione di Trieste, Via Valerio 2, 34127 Trieste, Italy*

E-mail: sbajeot@sissa.it, benve79@gmail.com

ABSTRACT: We apply the technique of sequential deconfinement to the four dimensional $\mathcal{N}=1$ $Usp(2N)$ gauge theory with an antisymmetric field and $2F$ fundamentals. The fully deconfined frame is a length- N quiver. We use this deconfined frame to prove the known self-duality of $Usp(2N)$ with an antisymmetric field and 8 fundamentals. Along the way we encounter a subtlety: in certain quivers with degenerate holomorphic operators, a naive application of Seiberg duality rules leads to an incorrect superpotential or chiral ring.

We also consider the reduction to $3d \mathcal{N}=2$ theories, recovering known fully deconfined duals of $Usp(2N)$ and $U(N)$ gauge theories, and obtaining new ones.

Contents

1	Introduction and summary	1
2	Basic S-confining and dualities moves	5
3	Case study: $Usp(6)$ with $\begin{smallmatrix} \square \\ \square \end{smallmatrix} + 8\square$	6
3.1	Sequential deconfinement	6
3.2	Degenerate holomorphic operator ambiguity	9
3.3	Fully deconfined frame	11
3.4	Self-duality	12
4	Sequential deconfinement of $Usp(2N)$ with $\begin{smallmatrix} \square \\ \square \end{smallmatrix} + 2F\square$	14
4.1	Deconfine and dualize: first step	15
4.2	Second step	16
4.3	After k steps	17
4.4	After N-1 steps	18
4.5	Final step and fully deconfined frame	19
5	Reconfinement and self-duality for $Usp(2N)$ with $\begin{smallmatrix} \square \\ \square \end{smallmatrix} + 8\square$	20
6	Reduction to $3d \mathcal{N} = 2$ sequential deconfinement	25
A	R-charges and degenerate holomorphic operator ambiguity in generic frame \mathcal{T}_k	32

1 Introduction and summary

Dualities are among the most powerful tool to analyze quantum field theories at strong coupling. In this paper we are interested in four dimensional $\mathcal{N} = 1$ gauge theories with rank-2 matter. Various recent works derived dualities involving gauge theories with rank-2 matter using only basic dualities involving gauge theories with fundamental matter, like Seiberg [1], Intriligator-Pouliot [2] and Aharony dualities [3].

In three dimensions, [4, 5] proved a $3d \mathcal{N} = 2$ S-confining duality for $U(N)$ with adjoint and $(1, 1)$ fundamentals [6–8] iterating Aharony duality. [9] proved $3d \mathcal{N} = 2$ self-dualities of $U(N)$ with adjoint and $(2, 2)$ fundamentals and of $Usp(2N)$ with antisymmetric and 6 fundamentals [7, 8], iterating Aharony dualities. Iterative application of confining monopole dualities [10] have been proven very useful in $3d$ also in [11–14].

More recently, in four dimensions, [15, 16] proved the S-confinement of $4d \mathcal{N} = 1$ $Usp(2N)$ with antisymmetric and 6 fundamentals, and [15] proved that all S-confining quivers with one node (classified long time ago in [17, 18]) can be obtained from Seiberg and Intriligator-Pouliot S-confinements.

By *proving the duality* between theory \mathcal{T}_1 and theory \mathcal{T}_k , we mean constructing a sequence of quiver gauge theories \mathcal{T}_i , $i = 1, \dots, k$, such that \mathcal{T}_i is related to \mathcal{T}_{i+1} by the application of an elementary duality on a single node. So all the theories $\mathcal{T}_1, \dots, \mathcal{T}_k$ are infrared dual. Assuming, as is standard, that the renormalization group flows commute with dualizing a single node, this amounts to a proof of the non-elementary duality $\mathcal{T}_1 \leftrightarrow \mathcal{T}_k$. The strategy to construct the sequence of dual quivers requires the *deconfinement* of some rank-2 matter. The deconfinement we need in this paper was introduced in [15, 16], and uplifts the $3d$ deconfinements used in [4, 5, 9].¹

In a related context, [22, 23] proved the $4d$ mirrors dualities [24] (uplift of $3d \mathcal{N} = 4$ mirror symmetries [25, 26] for linear quivers) using only Intriligator-Pouliot dualities. One difference is that, in the mirror symmetry case, the intermediate steps are quasi-Lagrangian theories, that is they involve gauging symmetries which only emerge in the infrared, as in [27].

In this paper we work out the *sequentially deconfined* dual of $4d \mathcal{N} = 1$ $Usp(2N)$ with antisymmetric and $2F$ fundamentals, uplifting the $3d \mathcal{N} = 2$ results of [9]. This means that we step by step prove a duality with a linear quiver gauge theories with N nodes and a certain *saw* structure, full details of the theory are in (4.5). This fully deconfined dual frame enjoys the nice property that all the chiral ring operators are gauge singlet fields, similarly to what happens for Intriligator-Pouliot and Aharony dualities.²

As an application of this construction, we provide a simple proof of the *self-duality*

¹Similar deconfinements appear in [19, 20] in the study of orientifolded dimer models, and were used in [21] to construct $\mathcal{N} = 1$ Lagrangians for $4d \mathcal{N} = 2$ SCFTs.

²This property cannot be enjoyed by theories with general baryonic operators in the chiral ring, such as in $SU(N)$ SQCD or in the sequentially deconfined $3d \mathcal{N} = 2$ SO gauge theories [28].

*modulo flips*³ of $Usp(2N)$ with antisymmetric and 8 fundamentals, proposed long time ago in [29]. By *self-dual modulo flips* we mean that the electric and magnetic theory share the same gauge structure, but differ by gauge singlets fields of flip type. Self-dualities modulo flips have been discussed in [29–32], the simplest case is $SU(2)$ with 8 doublets. Interestingly, given a self-duality modulo flips, it is possible to move the singlets across the duality and construct *exactly self-dual theories* with enhanced infra-red global symmetry, see for instance [32–36]. In our case of $Usp(2N)$ with antisymmetric and 8 fundamentals, [35] discussed various exactly self-dual theories and discussed the associated symmetry enhancements. Sometimes these symmetry enhancements can be understood compactifying a $6d$ $(1, 0)$ SCFT on a Riemann surface. For $Usp(2N)$ with antisymmetric and 8 fundamentals, [36] related the self-duality and specific symmetry enhancements to a compactification of the rank- N E-string $6d$ SCFT on a 2-sphere.

We encounter one subtlety (which is also present in the $3d$ cases but was overlooked in [9]) during the process that we call *degenerate holomorphic operator ambiguity*. As the name suggests, this phenomenon appears when we reach a frame that contains more than one gauge invariant holomorphic operator with the same global symmetry quantum numbers (including $U(1)_R$), but only one combination is a chiral protected operator. If such an operator is flipped by a gauge singlet, only one specific combination appears in the superpotential. In the examples we encounter in this paper, it happens that if we follow the rules of Seiberg duality (as is usually done) we end up with the incorrect result. In some cases we can determine which is the precise combination of operators appearing in the chiral ring (equivalently, the combination that can be flipped) by going in a dual frame and using classical F-terms relations there. Hence, in the original theory with *degenerate holomorphic operator ambiguity*, the ambiguity is resolved by quantum relations. In the case of $F = 4$, the precise superpotential is crucial in the proof of the self-duality modulo flips, so this case provides a good consistency check of our procedure.

Future directions

One natural question is whether there is a relation between our sequentially deconfined dual of $Usp(2N)$ with an antisymmetric and $2F$ fundamentals and Kutasov-Schwimmer type dualities [38–40], which for Usp were proposed in [41]. Namely one can turn on a superpotential term $tr(A^j)$ on the electric side, such term maps to a singlet on the

³Flipping a gauge invariant operator \mathcal{O} means adding a gauge singlet field σ to the theory, and an interaction term of the form $\sigma\mathcal{O}$ to the superpotential. The gauge singlet σ is called a *flipper*.

magnetic, so a Higgsing process is induced. The study of this Higgsing might shed light on the dualities of [38–41]. We expect the *degenerate holomorphic operator ambiguity* encountered in this paper to play an important role.

There are quite a few *self-dualities modulo flips* proposed in the literature [29–32], such as $SU(2N)$ with antisymmetric, conjugate antisymmetric and $(4, 4)$ fundamentals, or $SU(6)$ with 2 antisymmetrics and $(2, 6)$ fundamentals, or $SU(8)$ with 2 antisymmetrics and $(0, 8)$ fundamentals. A natural question is if such self-dualities can be proven using only the basic Seiberg and Intriligator-Pouliot dualities, as done in this work for $Usp(2N)$ with antisymmetric and 8 fundamentals. Notice that many self-dual gauge theories have been constructed simply ‘adding one flavor’ to an S-confining gauge theory [29, 31], so the fact that the S-confining dualities can be proven [15] is encouraging.

Related to the above point, many S-confinements and many self-dualities have been proposed for $4d \mathcal{N} = 1$ $Spin(N)$ theories with spinors and vectors [29, 31, 32, 34]. It would be very interesting to find a way to deconfine spinorial matter and try to prove such proposals.

Limits to gauge theories with orthogonal and/or symplectic gauge groups. S-confinements for $3d \mathcal{N} = 2$ theories with SO/Usp gauge groups and adjoint matter were recently proposed in [42], and [43] pointed out a relation to $4d \mathcal{N} = 1$ S-confinements for Usp gauge group and antisymmetric matter. It might be interesting to deform the $4d \mathcal{N} = 1$ sequential deconfinements as in [43]. See [28] for the sequential deconfinement of $3d \mathcal{N} = 2$ rank-2 matter with SO/Usp gauge groups.

It would also be interesting to study *degenerate holomorphic operator ambiguity* in other examples, possibly involving different kind of gauge groups.

Structure of the paper

This paper is organized as follows.

In section 2 we recall our notation and the three basic dualities which will be iteratively used in the rest of the paper.

In section 3 we discuss in detail a simple but non-trivial example, namely we sequentially deconfine $Usp(6)$ with an antisymmetric and 8 fundamentals. This example is the simplest one where the issue of the *degenerate holomorphic operator ambiguity* appears. We construct the fully deconfined dual, then we reconfine the quiver tail in order to prove the self-duality of the theory.

In section 4 we present the general sequential deconfinement of $Usp(2N)$ with an antisymmetric and $2F$ fundamentals.

In section 5 we set $2F = 8$, which allows to sequentially reconfine the quiver tail, and prove the self-duality modulo flips of the theory.


In section 6 we show how to reduce our $4d \mathcal{N} = 1$ $Usp(2N)$ story to $3d \mathcal{N} = 2$, re-obtaining the results of for $U(N)$ and $Usp(2N)$ found in [9]. Along the way we also derive new sequentially deconfined duals, namely for $U(N)$ with adjoint and (F, F) fundamentals with monopole superpotentials.

2 Basic S-confining and dualities moves

In this section we present the basic ingredients that we use in this paper to obtain the more complicated dualities involving rank-2 matter field. The first basic move is the deconfinement of an antisymmetric field with a Usp gauge group as in [15, 16], which is a modification of the original Berkooz deconfinement [44]. This form was used in [15, 16] to tackle $Usp(2N)$ with 6 fundamentals, which is S-confining.

Throughout the paper, blue $2N$ circles denote $Usp(2N)$ gauge nodes.

Usp deconfinement:⁴



$$\mathcal{W} = v_1 b_1 d_1 + h_1 d_1 v_2 + \beta_1 \text{tr}(b_1 b_1) \quad (2.1)$$

⁴On the r.h.s quiver in (2.1), we didn't include the flipper β_1 on the drawing. Sometimes, as in [15] this flipper is represented on the quiver by a cross, \times , on the bifundamental field b_1 . Moreover, the trace "tr" is taken using the Usp invariant antisymmetric matrix.

Throughout the paper, a trace of an operator in the antisymmetric of a Usp group is taken using the Usp invariant antisymmetric matrix.

In order to get this deconfinement we have to use another basic move. It is the S-confining result of Intriligator-Pouliot (IP) [2] that involves $Usp(2N)$ gauge group and $2N + 4$ fields in the fundamental representation.

IP S-confining duality:

$$\begin{array}{ccc}
 \begin{array}{c} \text{Diagram 1: A blue circle with } 2N \text{ inside, connected by an arrow labeled } Q \text{ to a white box with } 2N + 4 \text{ inside.} \\ \mathcal{W} = 0 \end{array} & \longleftrightarrow & \begin{array}{c} \text{Diagram 2: A white box with } 2N + 4 \text{ inside, with a curved arrow labeled } \mu \text{ looping back to itself.} \\ \mathcal{W} = \text{Pfaff}(\mu) \end{array} \\
 \text{Mapping:} & & tr(QQ) \longleftrightarrow \mu
 \end{array} \tag{2.2}$$

The last move concerns the same Usp theory but with $2F \geq 2N + 4$ fundamentals now. It is the IP duality [2]. In quiver notation it reads

IP duality:


$$\begin{array}{ccc}
 \begin{array}{c} \text{Diagram 1: A blue circle with } 2N \text{ inside, connected by an arrow labeled } Q \text{ to a white box with } 2F \text{ inside.} \\ \mathcal{W} = 0 \end{array} & \longleftrightarrow & \begin{array}{c} \text{Diagram 2: A blue circle with } 2F - 2N - 4 \text{ inside, connected by an arrow labeled } q \text{ to a white box with } 2F \text{ inside. The white box has a curved arrow labeled } \Phi \text{ looping back to itself.} \\ \mathcal{W} = tr(qq) \Phi \end{array} \\
 \text{Mapping:} & & tr(QQ) \longleftrightarrow \Phi
 \end{array} \tag{2.3}$$

3 Case study: $Usp(6)$ with $\begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array} + 8\square$

In this section we study the $Usp(6)$ gauge theory with matter in the antisymmetric representation and 8 fundamentals, that is 4 flavors. It is the simplest example that contains all the ingredients that we want to exhibit. In the next sections, we will show the general case. It is known that this theory is self-dual *modulo flips* [45]. We will prove the self-duality using the basic moves of the previous section (2.1), (2.2) and (2.3).


3.1 Sequential deconfinement

The first step is the deconfinement of the antisymmetric with (2.1). We get the following two frames \mathcal{T}_0 and $\mathcal{T}_{0'}$




The global symmetry of the original theory is $SU(8) \times U(1)_A \times U(1)_R$. When we split the 8 chirals into $7+1$, we split the $SU(8)$ into $SU(7) \times U(1)_P$. So in the splitted form, the R-charges of the fields should be a function of two variables (for the two global $U(1)$'s that can mix with the $U(1)_R$). We choose to write the R-charges in terms of R_Q and R_A . Then the R-charges of the other fields are determined by the $U(1)_R$ ABJ anomaly⁵ and by the requirement that any superpotential term should have R-charge equal to 2. We have written the R-charges of the fields next to them.

Then we dualize the $Usp(6)$ node with (2.3). The fields d_1 and β_1 get a mass. After integrating them out and rescaling the fields to put a +1 in front of each term in the superpotential, we get



⁵This is the same as requiring the vanishing of the NSVZ β function.

$$\mathcal{T}_2:$$


3.2 Degenerate holomorphic operator ambiguity

We claim that the correct answer is that the operator flipped by H_2 is $\mathcal{O}_2 = V_1 R_1$, instead of the naive \mathcal{O}_1 . Our argument in favor of this statement comes from dualizing some nodes in the quiver, as we will explain soon. So in a sense the fact the the correct operator is not the naive one is due to quantum relations, which become classical relation after Seiberg duality.


– 9 –

The symbol \simeq in the last of equality in (3.6) means an equivalence in the chiral ring.

Therefore we conclude that the non-zero operator in the chiral ring is $\mathcal{O}_2 = V_1 R_1$ and so it should be this one that enters in the superpotential with H_2 .

3.3 Fully deconfined frame

We now go back to our deconfining procedure, and dualize the right $Usp(2)$ node in \mathcal{T}_2 using (2.3). We reach a frame that we call “fully deconfined” as in [9].



$$\begin{aligned}
\mathcal{W} = & m_1 \text{tr} (b_1 b_2 q_3 q_3 b_2 b_1) + m_2 \text{tr} (b_2 q_3 q_3 b_2) + m_3 q_3 q_3 + \alpha_1 \text{tr} (b_1 b_1) + \alpha_2 \text{tr} (b_2 b_2) \\
& + \text{tr} (b_1 a_2 b_1) + \text{tr} (b_2 a_2 b_2) + r_1 b_1 v_2 + r_2 b_2 v_3 + l_1 v_1 b_1 b_2 q_3 + l_2 v_2 b_2 q_3 + l_3 v_3 q_3 + h_1 v_1 b_1 r_2 + h_2 v_1 r_1
\end{aligned} \tag{3.8}$$

The antisymmetric field a_2 is traceless, as all the antisymmetric field of Usp that will appear in this paper. Once again there is the question of the operator flipped by h_2 because $v_2 r_2$ has the same quantum numbers as $v_1 r_1$. Using the same procedure of confining from the left, we would obtain that the operator $v_2 r_2$ is 0 on the chiral ring. Therefore we claim that the correct final superpotential is the one with this switching procedure and not the one that we would have got using naive iteration of IP dualities.

The final mapping of the chiral ring generators is

\mathcal{T}_1		$\mathcal{T}_{1'}$		\mathcal{T}_2		\mathcal{T}_{Dec}
M_1		m_1		M_1		m_1
$Q_2 Q_2$		$q_2 q_2$		M_2		m_2
$\text{tr}(Q_2 \Phi Q_2)$		$\text{tr}(q_2 (c_2 c_2) q_2)$		$Q_3 Q_3$		m_3
$Q_2 P_2$	\iff	$q_2 c_2 p_3$	\iff	$Q_3 P_3$	\iff	l_3
$\text{tr}(Q_2 \Phi P_2)$		$q_2 p_2$		L_2		l_2
L_1		l_1		L_1		l_1
$\text{tr}(\Phi^2)$		h_2		H_2		h_2
H_1		h_1		H_1		h_1

(3.9)


Combining the two mappings (3.2) and (3.9), we get the mapping between \mathcal{T}_0 and \mathcal{T}_{Dec}

\mathcal{T}_1		\mathcal{T}_{DEC}	
$\text{tr}(Q A^i Q)$	\iff	m_{i+1}	$i = 0, 1, 2$
$\text{tr}(Q A^j P)$		l_{3-j}	$j = 0, 1, 2$
$\text{tr}(A^k)$		h_{N+1-k}	$k = 2, 3$

(3.10)

3.4 Self-duality


We already said that this theory is self-dual [45]. Let us see now how we can use our \mathcal{T}_{Dec} frame to prove the self-duality. The strategy is to *reconfine* the quiver tail. We notice that the left $Usp(2)$ has 6 fundamentals, so we start confining from the left. The effect of this confinement is to kill the antisymmetric field a_2 . In addition, the fields α_1 , v_2 and h_2 get a mass and we produce a Pfaffian superpotential as in (3.5). We get



$$\begin{aligned}
\mathcal{W} = & m_1 \text{tr}(b_2 b_2 b_2 q_3 q_3 b_2) + m_2 \text{tr}(b_2 q_3 q_3 b_2) + m_3 q_3 q_3 + \alpha_2 \text{tr}(b_2 b_2) + r_2 b_2 v_3 \\
& + l_1 p_1 b_2 q_3 + l_3 v_3 q_3 + h_1 p_1 r_2 + l_2 p_1 b_2 b_2 q_3
\end{aligned}
\tag{3.11}$$

Then we can confine the $Usp(4)$ node. We will reach the self-dual frame of the original theory. Indeed, we produce a traceless antisymmetric field, B , for the $Usp(6)$ and the fields h_1 , α_2 and v_3 get a mass. The final quiver reads

$$\mathcal{R}_2 \equiv \mathcal{R}_{final} :$$




$$\mathcal{W} = \sum_{j=1}^3 m_j \text{tr} (q_3 B^{3-j} q_3) + \sum_{j=1}^3 l_j \text{tr} (p_2 B^{j-1} q_3)$$

(3.12)

We can repackage the final result into a manifestly $SU(8)$ invariant way

$$\mathcal{R}_{final} :$$



$$\mathcal{W} = \sum_{j=1}^3 \mu_j \text{tr} (\tilde{Q} B^{3-j} \tilde{Q})$$

(3.13)

Where we define

$$\mu_j = \left(\begin{array}{c} \overbrace{m_j \dots m_j}^7 \quad \overbrace{l_{4-j} \dots l_{4-j}}^1 \\ \vdots \\ \dots \quad \vdots \quad \dots \\ \vdots \quad 0 \end{array} \right) \left. \begin{array}{l} \vphantom{\left(} \right. 7 \\ \vphantom{\left(} \right. 1 \end{array} \right\} , \quad \tilde{Q} = \left(\begin{array}{c} \overbrace{q_N \dots q_N}^7 \quad \overbrace{p_{3-1} \dots p_{3-1}}^1 \end{array} \right)$$

(3.14)

During this sequential reconfinement, from \mathcal{T}_{Dec} and \mathcal{R}_{final} , the mapping is

\mathcal{T}_{Dec}	\mathcal{R}_1	\mathcal{R}_{final}
m_i	m_i	m_i
l_i	l_i	l_i
h_2	$\text{tr} (b_2 b_2)^2$	$\text{tr} (B^2)$
h_1	h_1	$\text{tr} (B^3)$

(3.15)

Comparing with the mapping (3.10), we read the mapping for the self-duality

$$\begin{array}{ccc}
\mathcal{T}_1 & & \mathcal{R}_{final} \\
\text{tr}(Q A^i Q) & \Longleftrightarrow & m_{i+1} \quad i = 0, \dots, 2 \\
\text{tr}(Q A^j P) & & l_{3-j} \quad j = 0, \dots, 2 \\
\text{tr}(A^k) & & \text{tr}(B^k) \quad k = 2, \dots, 3
\end{array} \tag{3.16}$$

This is precisely the mapping given in [45].

Notice that in the reconfinement the precise operator flipped by the singlet H_2 was crucial to obtain the duality with the correct amount of gauge singlets.

In the next section we generalize our discussion to arbitrary N and F .

4 Sequential deconfinement of $Usp(2N)$ with $\square + 2F\square$

In this section we study the general case. The $Usp(2N)$ gauge theory with a traceless antisymmetric field A and $2F$ complex chiral fields (the number of fundamentals should be even to avoid the global anomaly).

We proceed as in [9] to derive a chain of $2N$ dual frames consisting of quiver theories, with number of gauge nodes ranging from 1 to N . Let us start with the first \mathcal{T}_0 quiver

$$\begin{array}{ccc}
\mathcal{T}_0 : & & \\
\begin{array}{c} \text{A loop labeled } A \text{ on a blue circle with } 2N \\ \text{connected to a box labeled } 2F \\ \mathcal{W} = 0 \end{array} & \equiv & \begin{array}{c} \text{A loop labeled } A \text{ on a blue circle with } 2N \\ \text{connected to a box labeled } 2F - 1 \text{ via an arrow labeled } Q \\ \text{A vertical line labeled } P \text{ connects the blue circle to a red box labeled } 1 \\ \mathcal{W} = 0 \end{array}
\end{array} \tag{4.1}$$

4.1 Deconfine and dualize: first step

We start by using the deconfinement (2.1), we obtain

$\mathcal{T}_{0'} :$

$$\mathcal{W} = p_1 c_1 d_1 + h_1 d_1 p_2 + \beta_1 \text{tr}(c_1 c_1) \quad (4.2)$$

Then we dualize the $Usp(2N)$ node with (2.3). This step is the same as in (3.1).

$\mathcal{T}_1 :$

$$\mathcal{W} = M_1 Q_1 Q_1 + Q_1 C_1 Q_2 + L_1 Q_1 V_1 + \text{tr}(C_1 \Phi C_1) + H_1 V_1 C_1 P_2 \quad (4.3)$$

The mapping of the chiral ring generators after this first step is the following

\mathcal{T}_0	$\mathcal{T}_{0'}$	\mathcal{T}_1	
QQ	$q_1 q_1$	M_1	
$\text{tr}(QA^i Q)$	$\text{tr}(q_1 (c_1 c_1)^i q_1)$	$\text{tr}(Q_2 \Phi^{i-1} Q_2)$	$i = 1, \dots, N-1$
$\text{tr}(QA^j P)$	$\text{tr}(q_1 c_1 (c_1 c_1)^j p_2)$	$\text{tr}(Q_2 \Phi^j P_2)$	$j = 0, \dots, N-2$
$\text{tr}(QA^{N-1} P)$	$q_1 p_1$	L_1	
$\text{tr}(A^m)$	$\text{tr}((c_1 c_1)^m)$	$\text{tr}(\Phi^m)$	$m = 2, \dots, N-1$
$\text{tr}(A^N)$	h_1	H_1	

$$(4.4)$$

Now we iterate the procedure. We deconfine the traceless antisymmetric field, Φ and then we dualize. Let us write explicitly another step and then it will be enough to obtain the general story.

4.2 Second step

After the deconfinement we get

$\mathcal{T}_{1'} :$

$$\begin{aligned} \mathcal{W} = & m_1 q_1 q_1 + q_1 c_1 q_2 + l_1 v_1 q_1 \\ & + \text{tr}(c_1 c_2 c_2 c_1) + h_1 v_1 c_1 c_2 p_3 \\ & + h_2 d_2 p_3 + p_2 c_2 d_2 + \beta_2 \text{tr}(c_2 c_2) \end{aligned} \quad (4.5)$$

Now we dualize the $Usp(2N-2)$ node. The fields q_1 , d_2 and β_2 get a mass. In addition $\text{tr}(c_1 c_2 c_2 c_1)$ becomes a mass term therefore there will be no link between the $Usp(2F-6)$ and $Usp(2N-4)$ gauge group. After the integration of these massive fields and a rescaling we get

$\mathcal{T}_2 :$

$$\begin{aligned} \mathcal{W} = & M_1 \text{tr}(B_1 Q_2 Q_2 B_1) + M_2 Q_2 Q_2 + \text{tr}(C_2 \Phi C_2) + \text{tr}(B_1 A_1 B_1) + \alpha_1 \text{tr}(B_1 B_1) \\ & + \text{tr}(B_1 C_2 C_2 B_1) + R_1 B_1 V_2 + L_1 V_1 B_1 Q_2 + L_2 V_2 Q_2 + Q_2 C_2 Q_3 + H_1 V_1 B_1 C_2 P_3 + H_2 V_2 C_2 P_3 \end{aligned} \quad (4.6)$$


We recall that the antisymmetric field A_1 is traceless, as well as Φ .

The mapping after this second step is given by⁸

$$\begin{array}{ccccc}
\mathcal{T}_1 & & \mathcal{T}_{1'} & & \mathcal{T}_2 \\
M_1 & & m_1 & & M_1 \\
Q_2 Q_2 & & q_2 q_2 & & M_2 \\
\text{tr}(Q_2 \Phi^{i-1} Q_2) & & \text{tr}(q_2 (c_2 c_2)^{i-1} q_2) & & \text{tr}(Q_3 \Phi^{i-2} Q_3) & i = 2, \dots, N-1 \\
\text{tr}(Q_2 \Phi^j P_2) & \iff & \text{tr}(q_2 c_2 (c_2 c_2)^j p_3) & \iff & \text{tr}(Q_3 \Phi^j P_3) & j = 0, \dots, N-3 \\
\text{tr}(Q_2 \Phi^{N-2} P_2) & & q_2 p_2 & & L_2 \\
L_1 & & l_1 & & L_1 \\
\text{tr}(\Phi^m) & & \text{tr}((b_2 b_2)^m) & & \text{tr}(\Phi^m) & m = 2, \dots, N-2 \\
\text{tr}(\Phi^{N-1}) & & h_2 & & H_2 \\
H_1 & & h_1 & & H_1
\end{array} \quad (4.7)$$

4.3 After k steps

After the iteration of k steps, we get the following quiver



$$\begin{aligned}
\mathcal{W} = & \sum_{i=1}^k M_i \text{tr}(B_i B_{i+1} \dots B_{k-1} Q_k Q_k B_{k-1} \dots B_{i+1} B_i) + \text{tr}(C_k \Phi C_k) + \sum_{i=1}^{k-1} \alpha_i \text{tr}(B_i B_i) \\
& + \sum_{i=1}^{k-1} \text{tr}(B_i A_i B_i) + \sum_{i=1}^{k-2} \text{tr}(B_i A_{i+1} B_i) + \text{tr}(B_{k-1} C_k C_k B_{k-1}) + \sum_{i=1}^{k-1} R_i B_i V_{i+1} \\
& + \sum_{i=1}^k L_i V_i B_i B_{i+1} \dots B_{k-1} Q_k + Q_k C_k Q_{k+1} + \sum_{i=1}^k H_i V_i B_i B_{i+1} \dots B_{k-1} C_k P_{k+1}
\end{aligned} \quad (4.8)$$

The last term in the superpotential should be taken with a grain of salt. Indeed as explained in the appendix A, when k is great enough some operators become degenerate


⁸Using the table of R-charges of the fields given in appendix A in Table 1, it is easy to check that the mapping is consistent.

with $V_i B_i B_{i+1} \dots B_{k-1} C_k P_{k+1}$ and then the superpotential should be modified. This is the *degenerate holomorphic operator ambiguity* that we described in (3.4). Since it is a k -dependent statement, we decided to be cavalier when writing this term in (4.8) and write the modified version in \mathcal{T}_{N-1} .

The mapping of the chiral ring generators is the following

$$\begin{array}{ccc}
\mathcal{T}_{(k-1)'} & & \mathcal{T}_k \\
m_{1,\dots,k-1} & & M_{1,\dots,k-1} \\
l_{1,\dots,k-1} & & L_{1,\dots,k-1} \\
h_{1,\dots,k} & & H_{1,\dots,k} \\
q_k q_k & \iff & M_k \\
\text{tr}(q_k (c_k c_k)^{i-k+1} q_k) & & \text{tr}(Q_{k+1} \Phi^{i-k} Q_{k+1}) \quad i = k, \dots, N-1 \\
\text{tr}(q_k c_k (c_k c_k)^j p_{k+1}) & & \text{tr}(Q_{k+1} \Phi^j P_{k+1}) \quad j = 0, \dots, N-k-1 \\
q_k p_k & & L_k \\
\text{tr}((c_k c_k)^m) & & \text{tr}(\Phi^m) \quad m = 2, \dots, N-k
\end{array} \tag{4.9}$$

4.4 After N-1 steps




$$\begin{aligned}
\mathcal{W} = & \sum_{i=1}^{N-1} M_i \text{tr}(B_i B_{i+1} \dots B_{N-2} Q_{N-1} Q_{N-1} B_{N-2} \dots B_{i+1} B_i) + \sum_{i=1}^{N-2} \alpha_i \text{tr}(B_i B_i) \\
& + \sum_{i=1}^{N-2} \text{tr}(B_i A_i B_i) + \sum_{i=1}^{N-3} \text{tr}(B_i A_{i+1} B_i) + \text{tr}(B_{N-2} C_{N-1} C_{N-1} B_{N-2}) \\
& + \sum_{i=1}^{N-2} R_i B_i V_{i+1} + \sum_{i=1}^{N-1} L_i V_i B_i B_{i+1} \dots B_{N-2} Q_{N-1} + Q_{N-1} C_{N-1} Q_N \\
& + H_1 V_1 B_1 \dots B_{N-2} C_{N-1} P_N + \sum_{i=2}^{N-1} H_i V_1 B_1 \dots B_{N-1-i} R_{N-i}
\end{aligned} \tag{4.10}$$

The last term in the superpotential is the one after switching, see Appendix A. Since the last node is $Usp(2)$, there is no antisymmetric and we can directly use (2.3).

4.5 Final step and fully deconfined frame

\mathcal{T}_{Dec} :



$$\begin{aligned}
\mathcal{W} = & \sum_{i=1}^N m_i \text{tr} (b_i b_{i+1} \dots b_{N-1} q_N q_N b_{N-1} \dots b_i b_{i+1}) + \sum_{i=1}^{N-1} \alpha_i \text{tr} (b_i b_i) \\
& + \sum_{i=1}^{N-1} \text{tr} (b_i a_i b_i) + \sum_{i=1}^{N-2} \text{tr} (b_i a_{i+1} b_i) + \sum_{i=1}^{N-1} r_i b_i v_{i+1} \\
& + \sum_{i=1}^N l_i v_i b_i b_{i+1} \dots b_{N-1} q_N + \sum_{i=1}^{N-1} h_i v_1 b_1 \dots b_{N-1-i} r_{N-i}
\end{aligned} \tag{4.11}$$

We have obtained the fully “deconfined” frame. The mapping is

$$\begin{array}{ccc}
\mathcal{T}_{N-1} & & \mathcal{T}_{DEC} \\
M_{1,\dots,N-1} & & m_{1,\dots,N-1} \\
L_{1,\dots,N-1} & \Longleftrightarrow & l_{1,\dots,N-1} \\
H_{1,\dots,N-1} & & h_{1,\dots,N-1} \\
Q_N Q_N & & m_N \\
Q_N P_N & & l_N
\end{array} \tag{4.12}$$

Collecting all the mappings, we write the mapping from the \mathcal{T}_0 frame to \mathcal{T}_{Dec} ⁹


$$\begin{array}{ccc}
\mathcal{T}_0 & & \mathcal{T}_{DEC} \\
\text{tr} (Q A^i Q) & \Longleftrightarrow & m_{i+1} \quad i = 0, \dots, N-1 \\
\text{tr} (Q A^j P) & & l_{N-j} \quad j = 0, \dots, N-1 \\
\text{tr} (A^k) & & h_{N+1-k} \quad k = 2, \dots, N
\end{array} \tag{4.13}$$

One interesting property of the “deconfined” frame is that all chiral ring generators are elementary gauge singlets.

⁹The mapping is consistent with the R-charges given in Table 2.

5 Reconfinement and self-duality for $Usp(2N)$ with $\begin{smallmatrix} \square \\ \square \end{smallmatrix} + 8\square$

It was proposed in [45] that in the special case of $F = 4$ the theory studied in the previous section, $Usp(2N)$ with antisymmetric and $2F$ fundamentals, is self-dual modulo flips. In this section we use our \mathcal{T}_{Dec} frame (4.11) to prove this result. Let us rewrite it, specifying $F = 4$:




$$\begin{aligned}
\mathcal{W} = & \sum_{i=1}^N m_i \text{tr} (b_i b_{i+1} \dots b_{N-1} q_N q_N b_{N-1} \dots b_{i+1} b_i) + \sum_{i=1}^{N-1} \alpha_i \text{tr} (b_i b_i) \\
& + \sum_{i=2}^{N-1} \text{tr} (b_i a_i b_i) + \sum_{i=1}^{N-2} \text{tr} (b_i a_{i+1} b_i) + \sum_{i=1}^{N-1} r_i b_i v_{i+1} \\
& + \sum_{i=1}^N l_i v_i b_i b_{i+1} \dots b_{N-1} q_N + \sum_{i=1}^{N-1} h_i v_i b_1 \dots b_{N-1-i} r_{N-i}
\end{aligned} \tag{5.1}$$

$$\begin{array}{ccc}
\mathcal{T}_0 & & R_0 \\
\begin{array}{l} \text{tr} (Q A^j Q) \\ \text{tr} (Q A^j P) \\ \text{tr} (A^i) \end{array} & \Longleftrightarrow & \begin{array}{ll} m_{j+1} & j = 0, \dots, N-1 \\ l_{N-j} & j = 0, \dots, N-1 \\ h_{N+1-i} & i = 2, \dots, N \end{array}
\end{array} \tag{5.2}$$

We see that the $Usp(2)$ gauge group is coupled to $4+1+1$ chiral fields in the fundamental representation and so it confines (2.2). This step is similar as in (3.11). The confinement will give a mass to the traceless antisymmetric field a_2 as well as v_2 , α_1 and h_{N-1} . After integrating them out and computing the Pfaffian superpotential (See discussion below

(3.5)) we get

\mathcal{R}_1 :




$$\begin{aligned}
\mathcal{W} = & m_1 \text{tr} (b_2 b_2 b_2 \dots b_{N-1} q_N q_N b_{N-1} \dots b_2) + \sum_{i=2}^N m_i \text{tr} (b_i b_{i+1} \dots b_{N-1} q_N q_N b_{N-1} \dots b_{i+1} b_i) \\
& + \sum_{i=2}^{N-1} \alpha_i \text{tr} (b_i b_i) + \sum_{i=3}^{N-1} \text{tr} (b_i a_i b_i) + \sum_{i=2}^{N-2} \text{tr} (b_i a_{i+1} b_i) + \sum_{i=2}^{N-1} r_i b_i v_{i+1} \\
& + l_1 p_1 b_2 \dots b_{N-1} q_N + l_2 p_1 b_2 b_2 b_2 \dots b_{N-1} q_N + \sum_{i=3}^N l_i v_i b_i b_{i+1} \dots b_{N-1} q_N + \sum_{i=1}^{N-2} h_i p_1 b_2 \dots b_{N-1-i} r_{N-i}
\end{aligned} \tag{5.3}$$

The mapping of the chiral ring generators is

$$\begin{array}{ccc}
R_0 & & R_1 \\
m_j, l_j & \Longleftrightarrow & m_j, l_j \quad \text{it will stay like that} \\
h_i & & h_i \quad i = 1, \dots, N-2 \\
h_{N-1} & & \text{tr} (b_2 b_2)^2
\end{array} \tag{5.4}$$

We can now iterate. Indeed the $Usp(4)$ group is coupled to $6 + 1 + 1$ fundamentals and so it also confines


$\mathcal{R}_2 :$



$$\begin{aligned}
\mathcal{W} = & m_1 \text{tr}([b_3 b_3]^3 b_4 \dots b_{N-1} q_N q_N b_{N-1} \dots b_4) + m_2 \text{tr}([b_3 b_3]^2 b_4 \dots b_{N-1} q_N q_N b_{N-1} \dots b_4) \\
& + \sum_{i=3}^N m_i \text{tr}(b_i b_{i+1} \dots b_{N-1} q_N q_N b_{N-1} \dots b_{i+1} b_i) + \sum_{i=3}^{N-1} \alpha_i \text{tr}(b_i b_i) + \sum_{i=4}^{N-1} \text{tr}(b_i a_i b_i) \\
& + \sum_{i=3}^{N-2} \text{tr}(b_i a_{i+1} b_i) + \sum_{i=3}^{N-1} r_i b_i v_{i+1} + l_1 p_2 b_3 \dots b_{N-1} q_N + l_2 p_2 [b_3 b_3] b_3 \dots b_{N-1} q_N \\
& + l_3 p_2 [b_3 b_3]^2 b_3 \dots b_{N-1} q_N + \sum_{i=4}^N l_i v_i b_i b_{i+1} \dots b_{N-1} q_N + \sum_{i=1}^{N-3} h_i p_2 b_3 \dots b_{N-1-i} r_{N-i}
\end{aligned} \tag{5.5}$$

We can iterate. After k confinement, we get

$\mathcal{R}_k :$




$$\begin{aligned}
\mathcal{W} = & \sum_{j=1}^k m_j \text{tr}([b_{k+1} b_{k+1}]^{k+2-j} b_{k+2} \dots b_{N-1} q_N q_N b_{N-1} \dots b_{k+2}) \\
& + \sum_{i=k+1}^N m_i \text{tr}(b_i b_{i+1} \dots b_{N-1} q_N q_N b_{N-1} \dots b_{i+1} b_i) + \sum_{i=k+1}^{N-1} \alpha_i \text{tr}(b_i b_i) \\
& + \sum_{i=k+2}^{N-1} \text{tr}(b_i a_i b_i) + \sum_{i=k+1}^{N-2} \text{tr}(b_i a_{i+1} b_i) + \sum_{i=k+1}^{N-1} r_i b_i v_{i+1} \\
& + \sum_{j=1}^{k+1} l_j p_k [b_{k+1} b_{k+1}]^{j-1} b_{k+1} \dots b_{N-1} q_N + \sum_{i=k+2}^N l_i v_i b_i b_{i+1} \dots b_{N-1} q_N \\
& + \sum_{i=1}^{N-k-1} h_i p_k b_{k+1} \dots b_{N-1-i} r_{N-i}
\end{aligned} \tag{5.6}$$

The mapping between two successive reconfinement is

$$\begin{array}{ccc}
R_{k-1} & & R_k \\
h_i & & h_i \\
h_{N-k} & & \text{tr}(b_{k+1}b_{k+1})^{k+1} \\
\text{tr}(b_k b_k)^k & \Longleftrightarrow & \text{tr}(b_{k+1}b_{k+1})^k \\
\vdots & & \vdots \\
\text{tr}(b_k b_k)^2 & & \text{tr}(b_{k+1}b_{k+1})^2
\end{array} \quad i = 1, \dots, N - k - 1 \quad (5.7)$$

For $k = N - 2$, there won't be any antisymmetric field left. We get



$$\begin{aligned}
\mathcal{W} = & \sum_{j=1}^{N-2} m_j \text{tr}([b_{N-1}b_{N-1}]^{N-j} q_N q_N) + m_{N-1} \text{tr}(b_{N-1} q_N q_N b_{N-1}) + m_N q_N q_N \\
& + \alpha_{N-1} \text{tr}(b_{N-1}b_{N-1}) + r_{N-1} b_{N-1} v_N + \sum_{j=1}^{N-1} l_j p_{N-2} [b_{N-1}b_{N-1}]^{j-1} b_{N-1} q_N + l_N v_N q_N + h_1 p_{N-2} r_{N-1}
\end{aligned} \quad (5.8)$$

Now we can do the last confinement with the $Usp(2N - 2)$ group. It will produce the traceless antisymmetric field, B for $Usp(2N)$ (the trace part is killed by the flipper α_{N-1}). We get

\mathcal{R}_{N-1} :

$$\begin{aligned}
\mathcal{W} = & \sum_{j=1}^N m_j \text{tr}(q_N B^{N-j} q_N) \\
& + \sum_{j=1}^N l_j \text{tr}(p_{N-1} B^{j-1} q_N)
\end{aligned} \quad (5.9)$$

The mapping for this last reconfinement is given by

$$\begin{array}{ccc}
R_{N-2} & & R_{N-1} \\
h_1 & & \text{tr}(B^N) \\
\text{tr}(b_{N-1}b_{N-1})^{N-1} & \Longleftrightarrow & \text{tr}(B^{N-1}) \\
\text{tr}(b_{N-1}b_{N-1})^{N-2} & & \text{tr}(B^{N-2}) \\
\vdots & & \vdots \\
\text{tr}(b_{N-1}b_{N-1})^2 & & \text{tr}(B^2)
\end{array} \quad (5.10)$$

We can repackage the last frame into a manifestly $SU(8)$ invariant way to obtain the final frame

$$\mathcal{R}_{final} : \quad \begin{array}{c} \mu_1, \dots, \mu_N \\ \begin{array}{c} \text{---} B \text{---} \\ \text{---} \tilde{Q} \text{---} \end{array} \end{array} \quad \mathcal{W} = \sum_{j=1}^N \mu_j \text{tr}(\tilde{Q} B^{N-j} \tilde{Q}) \quad (5.11)$$

Where we define

$$\mu_j = \left(\begin{array}{c} \overbrace{m_j \dots m_j}^7 \quad \overbrace{l_{N+1-j} \dots l_{N+1-j}}^1 \\ \vdots \\ \dots \quad \vdots \quad \dots \\ \vdots \quad 0 \end{array} \right) \left. \begin{array}{l} \left. \vphantom{\begin{array}{c} \overbrace{m_j \dots m_j}^7 \quad \overbrace{l_{N+1-j} \dots l_{N+1-j}}^1} \right\} 7 \\ \left. \vphantom{\begin{array}{c} \vdots \\ \dots \quad \vdots \quad \dots \\ \vdots \quad 0} \right\} 1 \end{array} \right\} \end{array} \right. , \quad \tilde{Q} = \left(\begin{array}{c} \overbrace{q_N \dots q_N}^7 \quad \overbrace{p_{N-1} \dots p_{N-1}}^1 \end{array} \right) \quad (5.12)$$

Now combining the mappings, we see that the reconfinement procedure gives

$$\begin{array}{ccc}
R_0 & \Longleftrightarrow & R_{final} \\
h_i & & \text{tr}(B^{N+1-i}) \quad i = 1, \dots, N-1
\end{array} \quad (5.13)$$

Now if we compare the original frame \mathcal{T}_0 and the last frame after reconfinement \mathcal{R}_{final} we see the self-duality and we obtain the following mapping

$$\begin{array}{ccc}
T_0 & & R_{final} \\
\text{tr}(Q A^j Q) & \Longleftrightarrow & m_{j+1} \quad j = 0, \dots, N-1 \\
\text{tr}(Q A^j P) & & l_{N-j} \quad j = 0, \dots, N-1 \\
\text{tr}(A^i) & & \text{tr}(B^i) \quad i = 2, \dots, N
\end{array} \quad (5.14)$$

Which is precisely the mapping proposed in [45].

6 Reduction to $3d \mathcal{N} = 2$ sequential deconfinement

It is possible to reduce $4d \mathcal{N} = 1$ theories on a circle, obtaining $3d \mathcal{N} = 2$ theories. Generically, this step introduces a superpotential term linear in the basic monopole operator (exceptions are, for instance, theories with 8 supercharges). Once in $3d$, it is possible to turn on real mass deformations, that do not exist in $4d$. Starting from a $4d \, Usp(2N)$ gauge theory, $3d$ real masses allow to flow to $Usp(2N)$ or $U(N)$ gauge groups with or without various types of monopole superpotentials. This process has been discussed in detail in the case without rank-2 matter [10], and for the case of $2F = 8$ [7, 8]. A brane interpretation has been found in [46]. Examples of $4d \mathcal{N} = 1$ symplectic quivers reduced and deformed to $3d \mathcal{N} = 2$ or $3d \mathcal{N} = 4$ unitary quivers have been discussed in [5, 37], mostly from the superconformal index perspective.

On the electric side, the story is as follows:

(6.1)


Where the rank-2 field is a traceless antisymmetric for $Usp(2N)$ and a traceless adjoint for $U(N)$. The monopoles $\mathfrak{M}, \mathfrak{M}^\pm$ are the monopoles with minimal GNO charges. See [10] for more details.

We could also turn on different real masses, possibly leading to non-zero Chern-Simons terms as in [7–9], but we refrain to do this in the present paper.

In the remaining of this section we perform the reduction and deformation of the fully deconfined dual, recovering the results found in [9] working in $3d$.

Reduction to the deconfined dual of $Usp(2N)$ with antisymmetric and $2F+2$ fundamentals, $\mathcal{W} = \mathfrak{M}$

We put the $4d$ duality on a circle. On the electric side we get $3d \mathcal{N} = 2 Usp(2N)$ with antisymmetric and $(2F+1)_Q + 1_P$ fundamentals, $\mathcal{W} = \mathfrak{M}$, with global symmetry $SU(2F+2) \times U(1)$. On the magnetic side we obtain the same quiver as in $4d$, (4.11), with $F \rightarrow F+1$. The difference is that the superpotential now includes N additional terms, linear in the monopole operators with GNO charges for a single gauge group, $\sum_{i=1}^N \mathfrak{M}^{0^{i-1}, \bullet, 0^{N-i}}$:



$$\begin{aligned}
 \mathcal{W} = & \sum_{i=1}^N m_i \text{tr} (b_i b_{i+1} \dots b_{N-1} q_N q_N b_{N-1} \dots b_i b_{i+1}) + \sum_{i=1}^{N-1} \alpha_i \text{tr} (b_i b_i) + \sum_{i=1}^{N-1} \text{tr} (b_i a_i b_i) \\
 & + \sum_{i=1}^{N-2} \text{tr} (b_i a_{i+1} b_i) + \sum_{i=1}^{N-1} r_i b_i v_{i+1} + \sum_{i=1}^N l_i v_i b_i b_{i+1} \dots b_{N-1} q_N + \sum_{i=1}^{N-1} h_i v_1 b_1 \dots b_{N-1-i} r_{N-i} \\
 & + \sum_{i=1}^N \mathfrak{M}^{0^{i-1}, \bullet, 0^{N-i}}
 \end{aligned} \tag{6.2}$$

For convenience we reproduce also the mapping of the chiral ring generators:

$$\begin{aligned}
 \text{tr} (Q A^i Q) & \longleftrightarrow m_{i+1} & i = 0, \dots, N-1 \\
 \text{tr} (Q A^j P) & \longleftrightarrow l_{N-j} & j = 0, \dots, N-1 \\
 \text{tr} (A^k) & \longleftrightarrow h_{N+1-k} & k = 2, \dots, N
 \end{aligned} \tag{6.3}$$

This is the same mapping as in $4d$, at this level there are no monopoles in the chiral ring, due to the presence of linear monopole terms in the superpotential.

Flow to the deconfined dual of $Usp(2N)$ with antisymmetric and $2F$ fundamentals, $\mathcal{W} = 0$

We now discuss what happens on the fully deconfined quiver 6.2 upon turning on real masses. We first turn on a real mass of the form $(0^{2F}, +, -)$ (that is we are moving to

the left in the diagram 6.1), on the electric side Q_{2F+1} and P become massive. Notice that the rank of the global symmetry decrease by one unit. Accordingly the mesons $\text{tr}(Q_I A^i Q_{2F+1})$ and $\text{tr}(Q_I A^i P)$ have non-zero real mass, for $I = 1, \dots, 2F$. Notice that $\text{tr}(Q_{2F+1} A^i P)$ has zero total real mass. The electric theory becomes $3d \mathcal{N} = 2$ $Usp(2N)$ with antisymmetric and $2F$ fundamentals, $\mathcal{W} = 0$, with global symmetry $SU(2F) \times U(1) \times U(1)$.

It follows from the mapping 6.3 that the singlets $(l_i)_I$ and $(m_i)_{I,2F+1}$ become massive for $I = 1, \dots, 2F$, while $(l_i)_{2F+1}$ remain massless. The l_j 's and $(m_i)_{I,2F+1}$'s become massive imply that also the elementary gauge variant fields v_i, r_i and $(q_N)_{2F+1}$ become massive. In other words the saw structure in the fully deconfined quiver disappears, and we are left with

$$\begin{aligned}
\mathcal{W} = & \sum_{i=1}^N m_i \text{tr}(b_i b_{i+1} \dots b_{N-1} q q b_{N-1} \dots b_i b_{i+1}) + \sum_{i=1}^{N-1} \alpha_i \text{tr}(b_i b_i) + \sum_{i=1}^{N-1} \text{tr}(b_i a_i b_i) \\
& + \sum_{i=1}^{N-2} \text{tr}(b_i a_{i+1} b_i) + \sum_{i=1}^{N-1} h_i \mathfrak{M}^{\bullet^{N-i}, 0^i} + \sum_{i=1}^N (l_i)_{2F+1} \mathfrak{M}^{0^{i-1}, \bullet^{N-i+1}}
\end{aligned} \tag{6.4}$$

Notice that the linear monopole superpotential $\sum_{i=1}^N \mathfrak{M}^{0^{i-1}, \bullet^{N-i}}$ is lifted, while the massless gauge singlets h_i and $(l_i)_{2F+1}$ now flip monopole operators. (this is similar to the dimensional reduction of Seiberg and Intriligator-Pouliot dualities [47]). These interactions are generated dynamically, one way to understand them is that such interactions are allowed by all global symmetries, and if they are not generated the gauge singlets would be free fields, which cannot be correct.

Equation (6.4) agrees with the results of section 2.4 of [9] (modulo renaming $h_i \rightarrow \gamma_i$ and $(l_i)_{2F+1} \rightarrow \sigma_i$), obtained by sequentially deconfining in $3d$, using the deconfining duality $antisymm_{2N \times 2N} \leftrightarrow Usp(2N-2) - [2N], \mathcal{W} = \gamma \mathfrak{M}$. The only difference is the precise extended monopole flipped by h_i , the subtlety related to the degenerate holomorphic operators which can in principle be flipped by h_i was not appreciated in [9]. One can check that with the superpotential above, setting $F = 3$, the tail reconfines appropriately and it is possible to derive the self-duality modulo flips of $3d \mathcal{N} = 2$ $Usp(2N)$ with antisymmetric and 6 fundamentals, $\mathcal{W} = 0$, at each step one h_i singlet is eaten, while the extended monopoles flipped by $(l_i)_{2F+1}$ 'shorten' according

to the rules of [48].

Flow to the deconfined dual of $U(N)$ with adjoint and $F + 1$ flavors, $\mathcal{W} = \mathfrak{M}^+ + \mathfrak{M}^-$

We now start from the $3d$ duality $Usp(2N)$ with $2F + 2$, $\mathcal{W} = \mathfrak{M} \leftrightarrow 6.2$, and turn on a real mass of the form $(+^{F+1}, -^{F+1})$ (that is we are moving to the right in the diagram 6.1). This type of real mass induces a Higgsing of the form $Usp(2N_i) \rightarrow U(N_i)$ on both sides of the duality, the antisymmetric fields are replaced by adjoints and a pair of fundamentals is replaced by a fundamental plus an antifundamental. The Higgsing is induced by a vev of 'Coulomb branch type', that is, on both sides of the duality, we are going to a specific sublocus of the moduli space of vacua (inside the so called $\mathcal{N} = 2$ Coulomb branch) where there is the maximum amount of massless fields. Moreover, the monopole superpotentials \mathfrak{M} 's are replaced by $(\mathfrak{M}^+ + \mathfrak{M}^-)$'s. See [5, 10, 37] for more details.

On the electric side we flow to $U(N)$ with adjoint and $(F_Q + 1_P, (F + 1)_{\tilde{Q}})$ flavors and $\mathcal{W} = \mathfrak{M}^+ + \mathfrak{M}^-$, with global symmetry $SU(F + 1) \times SU(F + 1) \times U(1)^{10}$. The rank of the global symmetry decreases by one unit, as it should.

On the magnetic side, we end up with a fully deconfined quiver

$$\begin{aligned}
\mathcal{W} = & \sum_{i=1}^N M_i \text{tr} (b_i b_{i+1} \dots b_{N-1} q \tilde{q} \tilde{b}_{N-1} \dots \tilde{b}_i \tilde{b}_{i+1}) + \sum_{i=1}^{N-1} \alpha_i \text{tr} (b_i \tilde{b}_i) + \sum_{i=1}^{N-1} \text{tr} (b_i a_i \tilde{b}_i) \\
& + \sum_{i=1}^{N-2} \text{tr} (b_i a_{i+1} \tilde{b}_i) + \sum_{i=1}^{N-1} \text{tr} (r_i b_i v_{i+1}) + \sum_{i=1}^N l_i \text{tr} (\tilde{q} \tilde{b}_{N-1} \dots \tilde{b}_{i+1} \tilde{b}_i v_i) \\
& + \sum_{i=1}^{N-1} h_i \text{tr} (r_{N-i} \tilde{b}_{N-1-i} \dots \tilde{b}_1 v_1) + \sum_{i=1}^N \left(\mathfrak{M}^{0^{i-1}, +, 0^{N-i}} + \mathfrak{M}^{0^{i-1}, -, 0^{N-i}} \right)
\end{aligned} \tag{6.5}$$

Where we did not draw the h_i singlets¹¹. The global symmetry of (6.5) is $SU(F + 1) \times SU(F) \times U(1) \times U(1)$.

¹⁰We split the global symmetry artificially into $SU(F + 1)_{\tilde{Q}} \times SU(F)_Q \times U(1)_P \times U(1)_A$ to match the visible symmetries in the magnetic side.

¹¹The double arrows $U(N_1) \leftrightarrow U(N_2)$ stand for a pair of bifundamentals with opposite orientation.

The mapping is¹²

$$\begin{aligned} \text{tr}(\tilde{Q}_I A^i Q^J) & \quad (M_{i+1})_I^J & i = 0, \dots, N-1 \\ \text{tr}(\tilde{Q}_I A^i P) & \iff (l_{N-i})_I & i = 0, \dots, N-1 \\ \text{tr}(A^k) & h_{N+1-k} & k = 2, \dots, N \end{aligned} \quad (6.6)$$

In the special case $F = 3$, we can use the confining duality for $U(N)$ with $(N+2, N+2)$ flavors and $\mathcal{W} = \mathfrak{M}^- + \mathfrak{M}^+$ [10] and reconfine the tail in 6.5, deriving the self-duality modulo flips of $3d \mathcal{N} = 2$ $U(N)$ with adjoint and $(4, 4)$ fundamentals, $\mathcal{W} = \mathfrak{M}^+ + \mathfrak{M}^-$, discussed in [7].

Flow to the deconfined dual of $U(N)$ with adjoint and F flavors, $\mathcal{W} = \mathfrak{M}^+$

We now turn on real masses $(0^F, +; 0^F, -)$ in the previous duality.

On the electric side one monopole superpotential is lifted and we flow to $U(N)$ with adjoint and $(F_Q, F_{\tilde{Q}})$ flavors and $\mathcal{W} = \mathfrak{M}^+$, with global symmetry $SU(F) \times SU(F) \times U(1) \times U(1)$. The rank of the global symmetry decreases by one unit, as expected.

On the magnetic side, using 6.6, the singlets $(l_j)_I$ and $(M_i)_{F+1}^J$ become massive. This in turn implies that \tilde{q}_{F+1} gets a mass, together with the v_i 's and the r_i 's. Hence the saw disappears and we end up with

$$\begin{aligned} \mathcal{W} = & \sum_{i=1}^N M_i \text{tr}(b_i b_{i+1} \dots b_{N-1} q \tilde{q} \tilde{b}_{N-1} \dots \tilde{b}_i \tilde{b}_{i+1}) + \sum_{i=1}^{N-1} \alpha_i \text{tr}(b_i \tilde{b}_i) + \sum_{i=1}^{N-1} \text{tr}(b_i a_i \tilde{b}_i) \\ & + \sum_{i=1}^{N-2} \text{tr}(b_i a_{i+1} \tilde{b}_i) + \sum_{i=1}^N (l_i)_{2F+1} \mathfrak{M}^{0^{i-1}, +^{N-i+1}} + \sum_{i=1}^{N-1} h_i \mathfrak{M}^{+, N-i, 0^i} + \sum_{i=1}^N \mathfrak{M}^{0^{i-1}, -, 0^{N-i}} \end{aligned} \quad (6.7)$$

Notice that half of the linear monopole superpotential disappeared and that the massless gauge singlets $(l_i)_F$ and h_i now flip monopole operators instead of mesons constructed with the saw.

In the special case $F = 3$, we can use the confining duality for $U(N)$ with $(N+1, N+1)$ flavors and $\mathcal{W} = \mathfrak{M}^- + h\mathfrak{M}^+$ [10] to reconfine the tail in 6.7, deriving the self-

¹²We have checked that the mapping is consistent by computing the R-charges of the operators (as a function of two variables corresponding to the two $U(1)$ symmetries) on both sides.


duality modulo flips of $3d \mathcal{N} = 2 U(N)$ with adjoint and $(3, 3)$ fundamentals, $\mathcal{W} = \mathfrak{M}^+$ [7].

Flow to the deconfined dual of $U(N)$ with adjoint and $F - 1$ flavors, $\mathcal{W} = 0$

We now turn on real masses $(0^{F-1}, +; 0^{F-1}, -)$ in the previous duality.

On the electric side one monopole superpotential is lifted and we flow to $U(N)$ with adjoint and $F - 1, F - 1$ flavors and $\mathcal{W} = 0$, with global symmetry $SU(F - 1) \times SU(F - 1) \times U(1)^3$. Again, the rank of the global symmetry decreases by one unit.

On the magnetic side,



$$\begin{aligned}
\mathcal{W} = & \sum_{i=1}^N M_i \text{tr}(b_i b_{i+1} \dots b_{N-1} q \tilde{q} \tilde{b}_{N-1} \dots \tilde{b}_i \tilde{b}_{i+1}) + \sum_{i=1}^{N-1} \alpha_i \text{tr}(b_i \tilde{b}_i) + \sum_{i=1}^{N-1} \text{tr}(b_i a_i \tilde{b}_i) \\
& + \sum_{i=1}^{N-2} \text{tr}(b_i a_{i+1} \tilde{b}_i) + \sum_{i=1}^N \left((l_i)_{2F+1} \mathfrak{M}^{0^{i-1}, +^{N-i+1}} + (M_i)_{F,F} \mathfrak{M}^{0^{i-1}, -^{N-i+1}} \right) \\
& + \sum_{i=1}^{N-1} h_i \mathfrak{M}^{+, 0^{N-i}} + \sum_{i=1}^{N-1} \mathfrak{M}^{0^{i-1}, -, 0^{N-i}}
\end{aligned} \tag{6.8}$$

The result 6.8 agrees with section 3.2 of [9] (modulo renaming $h_i \rightarrow \gamma_i$, $(l_i)_{2F+1} \rightarrow \sigma_i^+$, $(M_i)_{F,F} \rightarrow \sigma_i^-$ and $F \rightarrow F + 1$). Also here, the difference is the precise extended monopole flipped by h_i , the subtlety related to the degenerate holomorphic operators which can in principle be flipped by h_i was not appreciated in [9]. In the special case $F = 3$, we can use the confining duality for $U(N)$ with $(N + 1, N + 1)$ flavors and $\mathcal{W} = \mathfrak{M}^- + h\mathfrak{M}^+$ to reconfine the tail in 6.8 and derive the self-duality modulo flips of $3d \mathcal{N} = 2 U(N)$ with adjoint and $(2, 2)$ fundamentals, $\mathcal{W} = 0$, discussed in [7].¹³

¹³At each reconfining step, one linear monopole term disappears, one h_i is eaten, while the $(M_i)_I^J$'s ($I, J = 1, 2$) and the two towers $(l_i)_{2F+1}$ and $(M_i)_{F,F}$ survive. The self-duality reads

$$U(N) + \Phi + (2_Q, 2_{\bar{Q}}), \mathcal{W} = 0. \Leftrightarrow U(N) + \phi + (2_q, 2_{\bar{q}}), \mathcal{W} = \sum_i \left(M_i \text{tr}(\tilde{q} \phi^i q) + (l_i)_{2F+1} \mathfrak{M}_{\phi^i}^+ + (M_i)_{F,F} \mathfrak{M}_{\phi^i}^- \right). \tag{6.9}$$

Acknowledgments

We are grateful to Simone Giacomelli, Sara Pasquetti and Shlomo Razamat for useful conversations.

Stephane Bajeot is partially supported by the INFN Research Project STEFI. Sergio Benvenuti is partially supported by the INFN Research Project GAST.

A R-charges and degenerate holomorphic operator ambiguity in generic frame \mathcal{T}_k

We start by writing the R-charges of the fields in the generic frame \mathcal{T}_k (4.8).

Fields	R-charges in \mathcal{T}_k
V_i	$R_{V_i} = 5 - 2F + (N - \frac{5}{2} + \frac{3i}{2})R_A + (2F - 1)R_Q$
L_i	$R_{L_i} = 2F - 4 - (N - 2 + i)R_A - (2F - 2)R_Q$
M_i	$R_{M_i} = 2R_Q + (i - 1)R_A$
H_i	$(N - i + 1)R_A$
R_j	$R_{R_j} = 2F - 3 - (N + \frac{3j}{2} - \frac{1}{2})R_A - (2F - 1)R_Q$
B_j	$R_{B_j} = \frac{1}{2}R_A$
A_j	$R_{A_j} = 2 - R_A$
C_k	$R_{C_k} = 1 - \frac{1}{2}R_A$
Q_k	$R_{Q_k} = 1 + \frac{1}{2}(1 - k)R_A$
Q_{k+1}	$R_{Q_{k+1}} = R_Q + \frac{k}{2}R_A$
P_{k+1}	$R_{P_{k+1}} = 2F - 4 - (2N - 2 + \frac{k}{2})R_A - (2F - 1)R_Q$
Φ	$R_\Phi = R_A$

Table 1. R-charges in the frame \mathcal{T}_k with $k = 1, \dots, N - 1$, $i = 1, \dots, k$ and $j = 1, \dots, k - 1$.

Now we want to find the degenerate operators that can couple to H_i , as we did in (3.4) in the case of $Usp(6)$. In order to so, we look at the R-charges of the operator $V_i B_i \dots B_{k-1} C_k P_{k+1}$ which is the natural candidate to be coupled to H_i . We find

$$R(V_i B_i \dots B_{k-1} C_k P_{k+1}) = 2 - (N - i + 1)R_A \quad (\text{A.1})$$

The potential degenerate operators should be a singlet under the non-abelian global symmetry and should have the same R-charges (A.1). We can build the degenerate

operators from the fields V_a , B_b and R_c . Indeed, we start from V_m , then we put some B_b and end with R_n ($n \geq m$)¹⁴. The form of these operators is then: $V_m B_m \dots B_{n-1} R_n$. The number of fields B is $n - m$. The R-charge is

$$R_{V_m} + (n - m)R_{B_b} + R_{R_n} = 2 - (2 + (n - m))R_A \quad (\text{A.2})$$

If we compare with (A.1), we find the following condition

$$n - m = N - i - 1 \quad (\text{A.3})$$

with $1 \leq m \leq n \leq k - 1$, $i = 1, \dots, k$ and $k = 1, \dots, N - 1$.

We can already make two remarks:

- For $i = 1$, the constraint becomes $n - m = N - 2$ but the maximal value of $n - m$ is $k - 2$ and k satisfies $k \leq N - 1$. We conclude that there is never a solution for $i = 1$. Therefore there is never a degenerate operator associated to H_1 .
- If $k = 1$ then n and m don't exist. Conclusion, in order to get degenerate operators we should have $N \geq 3$ which means that degenerate operators will pop up in frames with at least 3 gauge groups (which correspond to $k = 2$).

We can ask the more precise question: What is the first frame, $\mathcal{T}_{k_{min}}$, when some operators degenerate? In order to answer that we have to try to maximize the l.h.s of (A.3) and minimize the r.h.s. Therefore it is enough to look at $n = k_{min} - 1$, $m = 1$ and $i = k_{min}$ to determine k_{min} (it could also have degenerate operators in $\mathcal{T}_{k_{min}}$ not associated to $n = k_{min} - 1$, $m = 1$ and $i = k_{min}$). We obtain

$$k_{min} - 2 = N - k_{min} - 1 \quad \Rightarrow \quad k_{min} = \left\lceil \frac{N + 1}{2} \right\rceil \quad (\text{A.4})$$

So when we reach $\mathcal{T}_{k_{min}}$ we start having this issue of degenerate operators.

Now let us solve (A.3) in the case $k = N - 1$. In this case, $1 \leq m \leq n \leq N - 2$

$$\begin{aligned} i = 2 : \quad & n = N + m - 3 \\ & \exists \text{ solution for } m = 1 \\ i = 3 : \quad & n = N + m - 4 \\ & \exists \text{ solution for } m = 1, 2 \\ & \vdots \\ i = N - 1 : \quad & n = m \\ & \exists \text{ solution for } m = 1, \dots, N - 2 \end{aligned}$$

¹⁴We cannot end with V_{j+1} because the F-term equation of R_j sets the combination $B_j V_{j+1}$ to 0.

Conclusion, the degenerate operators (with respect to $V_i B_i \dots B_{N-2} C_{N-1} F_N$) that potentially coupled to H_i in \mathcal{T}_{N-1} are:

$$\begin{aligned}
H_2 : & \quad V_1 B_1 \dots B_{N-3} R_{N-2} & 1 \text{ operator} \\
H_3 : & \quad V_1 B_1 \dots B_{N-4} R_{N-3}, V_2 B_2 \dots B_{N-3} R_{N-2} & 2 \text{ operators} \\
& \quad \vdots \\
H_{N-1} : & \quad V_1 R_1, V_2 R_2, \dots, V_{N-2} R_{N-2} & N-2 \text{ operators}
\end{aligned}$$

Finally, we can study the final frame \mathcal{T}_{Dec} . The R-charges are the following

Fields	R-charges in \mathcal{T}_{Dec}
v_i	$5 - 2F + (N - \frac{5}{2} + \frac{3i}{2})R_A + (2F - 1)R_Q$
l_i	$2F - 4 - (N - 2 + i)R_A - (2F - 2)R_Q$
m_i	$2R_Q + (i - 1)R_A$
h_j	$(N - i + 1)R_A$
r_j	$2F - 3 - (N + \frac{3j}{2} - \frac{1}{2})R_A - (2F - 1)R_Q$
b_j	$\frac{1}{2}R_A$
a_j	$2 - R_A$
q_N	$1 - R_Q - \frac{1}{2}(N - 1)R_A$

Table 2. R-charges in \mathcal{T}_{Dec} with $i = 1, \dots, N$ and $j = 1, \dots, N - 1$.

Now using the previous result for \mathcal{T}_{N-1} , we can summarize all the candidates to be coupled to h_i in the final frame \mathcal{T}_{Dec}

$$\begin{aligned}
h_1 : & \quad v_1 b_1 \dots b_{N-2} r_{N-1} & \text{No degenerate operator} \\
h_2 : & \quad v_1 b_1 \dots b_{N-3} r_{N-2}, v_2 b_2 \dots b_{N-2} r_{N-1} & 2 \text{ operators} \\
h_3 : & \quad v_1 b_1 \dots b_{N-4} r_{N-3}, v_2 b_2 \dots b_{N-3} r_{N-2}, v_3 b_3 \dots b_{N-2} r_{N-1} & 3 \text{ operators} \\
& \quad \vdots \\
h_{N-1} : & \quad v_1 r_1, v_2 r_2, \dots, v_{N-2} r_{N-2}, v_{N-1} r_{N-1} & N-1 \text{ operators}
\end{aligned}$$

Now the question is obvious, which operator is the correct one?

In the special case of $F = 4$, we could use the same argument that we used in Section 3. It goes as follows. When we reach the frame $\mathcal{T}_{k_{min}}$ some operators become degenerate. In order to decide the correct operator, we start confining from the left (it is possible because in the case of $F = 4$ the gauge group becomes $Usp(2)$). Then at some point we will discover that using the F-term equation for H_1 (which is never associated to a

degenerate operator as we saw) we can select the correct operator associated to some H_a (as in (3.6)). Then, the procedure is iterative meaning that we should re-use our previous results for H_a and do more and more reconfinement to select all the correct operators associated to the other H_b . All in all, we end up in the frame \mathcal{T}_{N-1} with the following superpotential term

$$\delta\mathcal{W}_{N-1} = H_1 V_1 B_1 \dots B_{N-2} C_{N-1} P_N + \sum_{i=2}^{N-1} H_i V_1 B_1 \dots B_{N-1-i} R_{N-i} \quad (\text{A.5})$$

which becomes in the final frame \mathcal{T}_{Dec}

$$\delta\mathcal{W}_{Dec} = \sum_{i=1}^{N-1} h_i v_1 b_1 \dots b_{N-1-i} r_{N-i} \quad (\text{A.6})$$

Unfortunately, in the case of $F > 4$ the previous argument fails because we cannot reconfine from the left. In this case we can deconfine the antisymmetric field A_1 but we didn't manage to find constraints and remove the degeneracy. Therefore, in this case the superpotential that we wrote in (4.10) and (4.11) are ambiguous. We wrote them with the results (A.5), (A.6) obtain in the case $F = 4$ but it is logically possible that they are wrong for $F > 4$.

References

- [1] N. Seiberg, “Electric - magnetic duality in supersymmetric nonAbelian gauge theories,” *Nucl. Phys. B* **435** (1995) 129–146, [arXiv:hep-th/9411149](#).
- [2] K. A. Intriligator and P. Pouliot, “Exact superpotentials, quantum vacua and duality in supersymmetric $\text{SP}(N(c))$ gauge theories,” *Phys. Lett. B* **353** (1995) 471–476, [arXiv:hep-th/9505006](#).
- [3] O. Aharony, “IR duality in $d = 3$ $N=2$ supersymmetric $\text{USp}(2N(c))$ and $\text{U}(N(c))$ gauge theories,” *Phys. Lett. B* **404** (1997) 71–76, [arXiv:hep-th/9703215](#).
- [4] S. Pasquetti and M. Sacchi, “From $3d$ dualities to $2d$ free field correlators and back,” *JHEP* **11** (2019) 081, [arXiv:1903.10817 \[hep-th\]](#).
- [5] S. Pasquetti and M. Sacchi, “ $3d$ dualities from $2d$ free field correlators: recombination and rank stabilization,” *JHEP* **01** (2020) 061, [arXiv:1905.05807 \[hep-th\]](#).
- [6] N. Aghaei, A. Amariti, and Y. Sekiguchi, “Notes on Integral Identities for $3d$ Supersymmetric Dualities,” *JHEP* **04** (2018) 022, [arXiv:1709.08653 \[hep-th\]](#).

- [7] S. Benvenuti, “A tale of exceptional 3d dualities,” *JHEP* **03** (2019) 125, [arXiv:1809.03925 \[hep-th\]](#).
- [8] A. Amariti and L. Cassia, “USp(2N_c) SQCD₃ with antisymmetric: dualities and symmetry enhancements,” *JHEP* **02** (2019) 013, [arXiv:1809.03796 \[hep-th\]](#).
- [9] S. Benvenuti, I. Garozzo, and G. Lo Monaco, “Sequential deconfinement in 3d $\mathcal{N} = 2$ gauge theories,” *JHEP* **07** (2021) 191, [arXiv:2012.09773 \[hep-th\]](#).
- [10] F. Benini, S. Benvenuti, and S. Pasquetti, “SUSY monopole potentials in 2+1 dimensions,” *JHEP* **08** (2017) 086, [arXiv:1703.08460 \[hep-th\]](#).
- [11] S. Benvenuti and S. Giacomelli, “Supersymmetric gauge theories with decoupled operators and chiral ring stability,” *Phys. Rev. Lett.* **119** no. 25, (2017) 251601, [arXiv:1706.02225 \[hep-th\]](#).
- [12] S. Benvenuti and S. Giacomelli, “Abelianization and sequential confinement in 2 + 1 dimensions,” *JHEP* **10** (2017) 173, [arXiv:1706.04949 \[hep-th\]](#).
- [13] S. Benvenuti and S. Giacomelli, “Lagrangians for generalized Argyres-Douglas theories,” *JHEP* **10** (2017) 106, [arXiv:1707.05113 \[hep-th\]](#).
- [14] S. Giacomelli and N. Mekareeya, “Mirror theories of 3d $\mathcal{N} = 2$ SQCD,” *JHEP* **03** (2018) 126, [arXiv:1711.11525 \[hep-th\]](#).
- [15] S. Bajeot and S. Benvenuti, “S-confinements from deconfinements,” [arXiv:2201.11049 \[hep-th\]](#).
- [16] L. E. Bottini, C. Hwang, S. Pasquetti, and M. Sacchi, “Dualities from dualities: the sequential deconfinement technique,” [arXiv:2201.11090 \[hep-th\]](#).
- [17] C. Csaki, M. Schmaltz, and W. Skiba, “A Systematic approach to confinement in N=1 supersymmetric gauge theories,” *Phys. Rev. Lett.* **78** (1997) 799–802, [arXiv:hep-th/9610139](#).
- [18] C. Csaki, M. Schmaltz, and W. Skiba, “Confinement in N=1 SUSY gauge theories and model building tools,” *Phys. Rev. D* **55** (1997) 7840–7858, [arXiv:hep-th/9612207](#).
- [19] I. Garcia-Etxebarria, B. Heidenreich, and T. Wrase, “New N=1 dualities from orientifold transitions. Part I. Field Theory,” *JHEP* **10** (2013) 007, [arXiv:1210.7799 \[hep-th\]](#).
- [20] I. n. García-Etxebarria, B. Heidenreich, and T. Wrase, “New N=1 dualities from orientifold transitions - Part II: String Theory,” *JHEP* **10** (2013) 006, [arXiv:1307.1701 \[hep-th\]](#).

- [21] I. n. G. Etxebarria, B. Heidenreich, M. Lotito, and A. K. Sorout, “Deconfining $\mathcal{N} = 2$ SCFTs, or the Art of Brane Bending,” [arXiv:2111.08022 \[hep-th\]](#).
- [22] L. E. Bottini, C. Hwang, S. Pasquetti, and M. Sacchi, “4d S -duality wall and $SL(2, \mathbb{Z})$ relations,” [arXiv:2110.08001 \[hep-th\]](#).
- [23] C. Hwang, S. Pasquetti, and M. Sacchi, “Rethinking mirror symmetry as a local duality on fields,” [arXiv:2110.11362 \[hep-th\]](#).
- [24] C. Hwang, S. Pasquetti, and M. Sacchi, “4d mirror-like dualities,” *JHEP* **09** (2020) 047, [arXiv:2002.12897 \[hep-th\]](#).
- [25] K. A. Intriligator and N. Seiberg, “Mirror symmetry in three-dimensional gauge theories,” *Phys. Lett. B* **387** (1996) 513–519, [arXiv:hep-th/9607207](#).
- [26] A. Hanany and E. Witten, “Type IIB superstrings, BPS monopoles, and three-dimensional gauge dynamics,” *Nucl. Phys. B* **492** (1997) 152–190, [arXiv:hep-th/9611230](#).
- [27] A. Gadde, S. S. Razamat, and B. Willett, ““Lagrangian” for a Non-Lagrangian Field Theory with $\mathcal{N} = 2$ Supersymmetry,” *Phys. Rev. Lett.* **115** no. 17, (2015) 171604, [arXiv:1505.05834 \[hep-th\]](#).
- [28] S. Benvenuti and G. Lo Monaco, “Sawing an adjoint: sequential deconfinement in ortho-symplectic gauge theories,” [to appear](#).
- [29] C. Csaki, M. Schmaltz, W. Skiba, and J. Terning, “Selfdual $N=1$ SUSY gauge theories,” *Phys. Rev. D* **56** (1997) 1228–1238, [arXiv:hep-th/9701191](#).
- [30] J. Distler and A. Karch, “ $N=1$ dualities for exceptional gauge groups and quantum global symmetries,” *Fortsch. Phys.* **45** (1997) 517–533, [arXiv:hep-th/9611088](#).
- [31] A. Karch, “More on $N=1$ selfdualities and exceptional gauge groups,” *Phys. Lett. B* **405** (1997) 280–286, [arXiv:hep-th/9702179](#).
- [32] S. S. Razamat, O. Sela, and G. Zafrir, “Curious patterns of IR symmetry enhancement,” *JHEP* **10** (2018) 163, [arXiv:1809.00541 \[hep-th\]](#).
- [33] S. S. Razamat, O. Sela, and G. Zafrir, “Between Symmetry and Duality in Supersymmetric Quantum Field Theories,” *Phys. Rev. Lett.* **120** no. 7, (2018) 071604, [arXiv:1711.02789 \[hep-th\]](#).
- [34] O. Sela and G. Zafrir, “Symmetry enhancement in 4d Spin(n) gauge theories and compactification from 6d,” *JHEP* **12** (2019) 052, [arXiv:1910.03629 \[hep-th\]](#).
- [35] C. Hwang, S. Pasquetti, and M. Sacchi, “Flips, dualities and symmetry enhancements,” *JHEP* **05** (2021) 094, [arXiv:2010.10446 \[hep-th\]](#).

- [36] C. Hwang, S. S. Razamat, E. Sabag, and M. Sacchi, “Rank Q E-string on spheres with flux,” *SciPost Phys.* **11** no. 2, (2021) 044, [arXiv:2103.09149 \[hep-th\]](#).
- [37] S. Pasquetti, S. S. Razamat, M. Sacchi, and G. Zafrir, “Rank Q E-string on a torus with flux,” *SciPost Phys.* **8** no. 1, (2020) 014, [arXiv:1908.03278 \[hep-th\]](#).
- [38] D. Kutasov, “A Comment on duality in $N=1$ supersymmetric nonAbelian gauge theories,” *Phys. Lett. B* **351** (1995) 230–234, [arXiv:hep-th/9503086](#).
- [39] D. Kutasov and A. Schwimmer, “On duality in supersymmetric Yang-Mills theory,” *Phys. Lett. B* **354** (1995) 315–321, [arXiv:hep-th/9505004](#).
- [40] D. Kutasov, A. Schwimmer, and N. Seiberg, “Chiral rings, singularity theory and electric - magnetic duality,” *Nucl. Phys. B* **459** (1996) 455–496, [arXiv:hep-th/9510222](#).
- [41] J. H. Brodie and M. J. Strassler, “Patterns of duality in $N=1$ SUSY gauge theories, or: Seating preferences of theater going nonAbelian dualities,” *Nucl. Phys. B* **524** (1998) 224–250, [arXiv:hep-th/9611197](#).
- [42] S. Benvenuti and G. Lo Monaco, “A toolkit for ortho-symplectic dualities,” [arXiv:2112.12154 \[hep-th\]](#).
- [43] A. Amariti and S. Rota, “3d $\mathcal{N} = 2$ SO/USp adjoint SQCD: s-confinement and exact identities,” [arXiv:2202.06885 \[hep-th\]](#).
- [44] M. Berkooz, “The Dual of supersymmetric $SU(2k)$ with an antisymmetric tensor and composite dualities,” *Nucl. Phys. B* **452** (1995) 513–525, [arXiv:hep-th/9505067](#).
- [45] C. Csaki, W. Skiba, and M. Schmaltz, “Exact results and duality for $SP(2N)$ SUSY gauge theories with an antisymmetric tensor,” *Nucl. Phys. B* **487** (1997) 128–140, [arXiv:hep-th/9607210](#).
- [46] A. Amariti, D. Orlando, and S. Reffert, “Monopole Quivers and new 3D $N=2$ dualities,” *Nucl. Phys. B* **924** (2017) 153–177, [arXiv:1705.09297 \[hep-th\]](#).
- [47] O. Aharony, S. S. Razamat, N. Seiberg, and B. Willett, “3d dualities from 4d dualities,” *JHEP* **07** (2013) 149, [arXiv:1305.3924 \[hep-th\]](#).
- [48] S. Benvenuti, I. Garozzo, and G. Lo Monaco, “Monopoles and dualities in 3d $\mathcal{N} = 2$ quivers,” *JHEP* **10** (2021) 191, [arXiv:2012.08556 \[hep-th\]](#).