

# Homotopy cartesian squares in extriangulated categories<sup>★</sup>

Jing He, Chenbei Xie and Panyue Zhou

## Abstract

Let  $(\mathcal{C}, \mathbb{E}, \mathfrak{s})$  be an extriangulated category. Given a composition of two commutative squares in  $\mathcal{C}$ , if two commutative squares are homotopy cartesian, then their composition is also a homotopy cartesian. This covers the result by Mac Lane (1998) for abelian categories and the result by Christensen and Frankland (2022) for triangulated categories.

**Keywords:** extriangulated categories; homotopy cartesian squares; triangulated categories; abelian categories; push-out and pull-back

**2020 Mathematics Subject Classification:** 18G80; 18E10

## 1 Introduction

Most of homological algebra can be carried out in the setting of abelian categories, a class of categories which includes all categories of modules. We recall the properties of pull-back and push-out squares in an abelian category. Given two morphisms  $x': A' \rightarrow B'$  and  $b: B \rightarrow B'$ . Then a commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{x} & B \\ a \downarrow & & \downarrow b \\ A' & \xrightarrow{x'} & B' \end{array}$$

is *pull-back* if and only if  $0 \rightarrow A \xrightarrow{\begin{bmatrix} a \\ x \end{bmatrix}} A' \oplus B \xrightarrow{[x' \ -b]} B' \rightarrow 0$  is left exact. Given two morphisms  $x: A \rightarrow B$  and  $a: A \rightarrow A'$ . Then a commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{x} & B \\ a \downarrow & & \downarrow b \\ A' & \xrightarrow{x'} & B' \end{array}$$

is *push-out* if and only if  $A \xrightarrow{\begin{bmatrix} a \\ x \end{bmatrix}} A' \oplus B \xrightarrow{[x' \ -b]} B' \rightarrow 0$  is right exact. By the above properties, we can obtain that the commutative diagram

$$\begin{array}{ccc} A & \xrightarrow{x} & B \\ a \downarrow & & \downarrow b \\ A' & \xrightarrow{x'} & B' \end{array}$$

---

<sup>★</sup>Panyue Zhou was supported by the National Natural Science Foundation of China (Grant No. 11901190).

is *pull-back and push-out* if and only if

$$0 \rightarrow A \xrightarrow{\begin{bmatrix} a \\ x \end{bmatrix}} A' \oplus B \xrightarrow{[x' \ -b]} B' \rightarrow 0$$

is exact.

Triangulated categories were introduced in the mid 1960's by Verdier in his thesis. Having their origins in algebraic geometry and algebraic topology, triangulated categories have by now become indispensable in many different areas of mathematics. Neeman [N, Definition 1.4.1] introduced the notion of homotopy cartesian squares in triangulated categories. More precisely, let  $\mathcal{C}$  be a triangulated category with shift functor  $\Sigma$ . A commutative square

$$\begin{array}{ccc} A & \xrightarrow{x} & B \\ a \downarrow & & \downarrow b \\ A' & \xrightarrow{x'} & B' \end{array}$$

is called *homotopy cartesian* if there exists a triangle

$$A \xrightarrow{\begin{bmatrix} a \\ x \end{bmatrix}} A' \oplus B \xrightarrow{[x' \ -b]} B' \xrightarrow{\partial} \Sigma A$$

The morphism  $\partial$  is called a *differential* of the homotopy cartesian square. We can think of a homotopy cartesian square as the triangulated analogue of a pull-back and push-out square in an abelian category.

Recently, extriangulated categories were recently introduced by Nakaoka and Palu [NP] by extracting those properties of  $\text{Ext}^1(-, -)$  on exact categories and on triangulated categories that seem relevant from the point of view of cotorsion pairs. In particular, triangulated categories and exact categories are extriangulated categories. There are a lot of examples of extriangulated categories which are neither triangulated categories nor exact categories, see [NP, ZZ, HZZ, ZhZ, NP1]. Hence, many results on exact categories (abelian categories are exact categories) and triangulated categories can be unified in the same framework. Motivated by this idea, we study homotopy cartesian squares in an extriangulated category. Let us recall the definition of homotopy cartesian squares in an extriangulated category.

**Definition 1.1.** [H, Definition 3.1] Let  $(\mathcal{C}, \mathbb{E}, \mathfrak{s})$  be an extriangulated category. Then a commutative square

$$\begin{array}{ccc} A & \xrightarrow{x} & B \\ a \downarrow & & \downarrow b \\ A' & \xrightarrow{x'} & B' \end{array}$$

is called *homotopy cartesian* if there exists an  $\mathbb{E}$ -triangle

$$A \xrightarrow{\begin{bmatrix} a \\ x \end{bmatrix}} A' \oplus B \xrightarrow{[x' \ -b]} B' \xrightarrow{\theta} \mathbb{E}(B', A)$$

The  $\mathbb{E}$ -extension  $\theta \in \mathbb{E}(B', A)$  is called a *differential* of the homotopy cartesian square.

This gives a simultaneous generalization of push-out and pullback squares in an abelian

category and homotopy cartesian squares in triangulated categories.

The main result of this article is the following, which is a simultaneous generalization of a result of Mac Lane [M] for abelian categories and of a result of Christensen-Frankland [CF] for triangulated categories.

**Theorem 1.2.** (see Theorem 3.2 for more details) *Let  $(\mathcal{C}, \mathbb{E}, \mathfrak{s})$  be an extriangulated category. Consider the following commutative diagram*

$$\begin{array}{ccccc} A & \xrightarrow{x} & B & \xrightarrow{y} & C \\ a \downarrow & & \downarrow b & & \downarrow c \\ & \text{(I)} & & \text{(II)} & \\ A' & \xrightarrow{x'} & B' & \xrightarrow{y'} & C' \end{array}$$

*If the two squares (I) and (II) are homotopy cartesian in  $\mathcal{C}$ , then the outside rectangle is a homotopy cartesian in  $\mathcal{C}$ .*

*Moreover, given differentials  $\theta_L \in \mathbb{E}(B', A)$  and  $\theta_R \in \mathbb{E}(C', B)$  of the left square and right square respectively, there exists a differential  $\theta_P$  for the outside rectangle satisfying*

$$\theta_R = x_* \theta_P \quad \text{and} \quad \theta_L = (y')^* \theta_P.$$

This article is organized as follows. In Section 2, we recall the definition of an extriangulated category. In Section 3, we give a composition of two commutative squares, if both squares are homotopy cartesian, then their composition is also a homotopy cartesian.

## 2 Preliminaries

Let us briefly recall some definitions and basic properties of extriangulated categories from [NP]. We omit some details here, but the reader can find them in [NP].

Let  $\mathcal{C}$  be an additive category equipped with an additive bifunctor

$$\mathbb{E} : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \text{Ab},$$

where Ab is the category of abelian groups. For any objects  $A, C \in \mathcal{C}$ , an element  $\delta \in \mathbb{E}(C, A)$  is called an  $\mathbb{E}$ -extension. Let  $\mathfrak{s}$  be a correspondence which associates an equivalence class

$$\mathfrak{s}(\delta) = [A \xrightarrow{x} B \xrightarrow{y} C]$$

to any  $\mathbb{E}$ -extension  $\delta \in \mathbb{E}(C, A)$ . This  $\mathfrak{s}$  is called a *realization* of  $\mathbb{E}$ , if it makes the diagrams in [NP, Definition 2.9] commutative. A triplet  $(\mathcal{C}, \mathbb{E}, \mathfrak{s})$  is called an *extriangulated category* if it satisfies the following conditions.

- (1)  $\mathbb{E} : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \text{Ab}$  is an additive bifunctor.
- (2)  $\mathfrak{s}$  is an additive realization of  $\mathbb{E}$ .
- (3)  $\mathbb{E}$  and  $\mathfrak{s}$  satisfy the compatibility conditions in [NP, Definition 2.12].

**Remark 2.1.** Note that both exact categories and triangulated categories are extriangulated categories, see [NP, Example 2.13] and extension closed subcategories of extriangulated categories are again extriangulated, see [NP, Remark 2.18]. Moreover, there are extriangulated categories which are neither exact categories nor triangulated categories, see [NP, Proposition 3.30], [ZZ, Example 4.14], [HZZ, Remark 3.3], [ZhZ, Lemma 20] and [NP1, Theorem 4.22].

We collect the following terminology from [NP].

**Definition 2.2.** [NP, Definition 2.19] Let  $(\mathcal{C}, \mathbb{E}, \mathfrak{s})$  be an extriangulated category.

- (1) A sequence  $A \xrightarrow{x} B \xrightarrow{y} C$  is called a *conflation* if it realizes some  $\mathbb{E}$ -extension  $\delta \in \mathbb{E}(C, A)$ . In this case,  $x$  is called an *inflation* and  $y$  is called a *deflation*.
- (2) If a conflation  $A \xrightarrow{x} B \xrightarrow{y} C$  realizes  $\delta \in \mathbb{E}(C, A)$ , we call the pair  $(A \xrightarrow{x} B \xrightarrow{y} C, \delta)$  an  $\mathbb{E}$ -triangle, and write it in the following way.

$$A \xrightarrow{x} B \xrightarrow{y} C \overset{\delta}{\dashrightarrow}$$

We usually do not write this “ $\delta$ ” if it is not used in the argument.

- (3) Let  $A \xrightarrow{x} B \xrightarrow{y} C \overset{\delta}{\dashrightarrow}$  and  $A' \xrightarrow{x'} B' \xrightarrow{y'} C' \overset{\delta'}{\dashrightarrow}$  be any pair of  $\mathbb{E}$ -triangles. If a triplet  $(a, b, c)$  realizes  $(a, c): \delta \rightarrow \delta'$ , then we write it as

$$\begin{array}{ccccc} A & \xrightarrow{x} & B & \xrightarrow{y} & C \overset{\delta}{\dashrightarrow} \\ \downarrow a & & \downarrow b & & \downarrow c \\ A' & \xrightarrow{x'} & B' & \xrightarrow{y'} & C' \overset{\delta'}{\dashrightarrow} \end{array}$$

and call  $(a, b, c)$  a *morphism of  $\mathbb{E}$ -triangles*.

### 3 Proof of the Main Result

First we prove the following crucial lemma.

**Lemma 3.1.** *Let  $(\mathcal{C}, \mathbb{E}, \mathfrak{s})$  be an extriangulated category. Consider a homotopy cartesian square*

$$\begin{array}{ccc} A & \xrightarrow{x} & B \\ \downarrow a & & \downarrow b \\ A' & \xrightarrow{x'} & B' \xrightarrow{y'} C' \overset{\delta'}{\dashrightarrow} \end{array}$$

*with given differential  $\theta \in \mathbb{E}(B', A)$  and the bottom row is an  $\mathbb{E}$ -triangle in  $\mathcal{C}$ . Then the square extends to a morphism of  $\mathbb{E}$ -triangles of the form  $(a, b, 1)$ , as illustrated in the diagram*

$$\begin{array}{ccc} A & \xrightarrow{x} & B \xrightarrow{y} C' \overset{\delta}{\dashrightarrow} \\ \downarrow a & & \downarrow b \quad \parallel \\ A' & \xrightarrow{x'} & B' \xrightarrow{y'} C' \overset{\delta'}{\dashrightarrow} \end{array},$$

*satisfying  $\theta = (y')^* \delta$ .*

*Proof.* Since the square

$$\begin{array}{ccc} A & \xrightarrow{x} & B \\ a \downarrow & & \downarrow b \\ A' & \xrightarrow{x'} & B' \end{array}$$

is homotopy cartesian, then there exists an  $\mathbb{E}$ -triangle

$$A \xrightarrow{\begin{bmatrix} a \\ x \end{bmatrix}} A' \oplus B \xrightarrow{[x' \ -b]} B' \xrightarrow{-\theta} \gg$$

By the dual of Proposition 3.17 in [NP], we obtain the following commutative diagram of  $\mathbb{E}$ -triangles

$$\begin{array}{ccccccc} & & A & \xlongequal{\quad} & A & & \\ & & \downarrow \begin{bmatrix} a \\ x \end{bmatrix} & & \downarrow x & & \\ A' & \xrightarrow{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} & A' \oplus B & \xrightarrow{[0 \ 1]} & B & \xrightarrow{0} & \gg \\ \parallel & & \downarrow [x' \ -b] & & \downarrow -y & & \\ A' & \xrightarrow{x'} & B' & \xrightarrow{y'} & C' & \xrightarrow{-\delta'} & \gg \\ & & \downarrow \theta & & \downarrow -\delta & & \\ & & \Downarrow & & \Downarrow & & \end{array}$$

which satisfies  $\theta = (y')^*\delta$ . From the diagram, we have  $y = y'b$ . The commutative diagram

$$\begin{array}{ccccc} A & \xrightarrow{x} & B & \xrightarrow{-y} & C' & \xrightarrow{-\delta} \gg \\ \parallel & & \parallel & & \downarrow -1 & \\ A & \xrightarrow{x} & B & \xrightarrow{y} & C & \xrightarrow{-\delta} \gg \end{array}$$

shows that  $A \xrightarrow{x} B \xrightarrow{y} C \xrightarrow{-\delta} \gg$  is an  $\mathbb{E}$ -triangle. Thus we obtain the following morphism of  $\mathbb{E}$ -triangles

$$\begin{array}{ccccc} A & \xrightarrow{x} & B & \xrightarrow{y} & C' & \xrightarrow{-\delta} \gg \\ \downarrow a & & \downarrow b & & \parallel & \\ A' & \xrightarrow{x'} & B' & \xrightarrow{y'} & C' & \xrightarrow{-\delta'} \gg \end{array}$$

This completes the proof.  $\square$

We now prove our main result.

**Theorem 3.2.** *Let  $(\mathcal{C}, \mathbb{E}, \mathfrak{s})$  be an extriangulated category. Consider the following commutative diagram*

$$\begin{array}{ccccc} A & \xrightarrow{x} & B & \xrightarrow{y} & C \\ a \downarrow & \text{(I)} & \downarrow b & \text{(II)} & \downarrow c \\ A' & \xrightarrow{x'} & B' & \xrightarrow{y'} & C' \end{array}$$

*If the two squares (I) and (II) are homotopy cartesian in  $\mathcal{C}$ , then the outside rectangle is a homotopy cartesian in  $\mathcal{C}$ .*

*Moreover, given differentials  $\theta_L \in \mathbb{E}(B', A)$  and  $\theta_R \in \mathbb{E}(C', B)$  of the left square and right*

square respectively, there exists a differential  $\theta_P$  for the outside rectangle satisfying

$$\theta_R = x_*\theta_P \quad \text{and} \quad \theta_L = (y')^*\theta_P.$$

*Proof.* Since the square (I) is homotopy cartesian, then there exists an  $\mathbb{E}$ -triangle

$$A \xrightarrow{\begin{bmatrix} a \\ x \end{bmatrix}} A' \oplus B \xrightarrow{[x' \ -b]} B' \xrightarrow{\theta_L} \gg$$

with differential  $\theta_L \in \mathbb{E}(B', A)$ . Note that the following diagram

$$\begin{array}{ccccc} A & \xrightarrow{\begin{bmatrix} a \\ x \end{bmatrix}} & A' \oplus B & \xrightarrow{[x' \ -b]} & B' \xrightarrow{\theta_L} \gg \\ \parallel & & \downarrow \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} & & \downarrow -1 \\ A & \xrightarrow{\begin{bmatrix} x \\ a \end{bmatrix}} & B \oplus A' & \xrightarrow{[b \ -x']} & B' \xrightarrow{\theta_L} \gg \end{array}$$

is commutative and all vertical morphisms are isomorphisms, thus we have that

$$A \xrightarrow{\begin{bmatrix} x \\ a \end{bmatrix}} B \oplus A' \xrightarrow{[b \ -x']} B' \xrightarrow{\theta_L} \gg \quad (3.1)$$

is also an  $\mathbb{E}$ -triangle. By the direct sum of the  $\mathbb{E}$ -triangle (3.1) and the trivial  $\mathbb{E}$ -triangle  $0 \rightarrow C \xrightarrow{-1} C \xrightarrow{0} \gg$ , we obtain that

$$A \xrightarrow{\begin{bmatrix} x \\ a \\ 0 \end{bmatrix}} B \oplus A' \oplus C \xrightarrow{\begin{bmatrix} b \ -x' \ 0 \\ 0 \ 0 \ -1 \end{bmatrix}} B' \oplus C \xrightarrow{[\theta_L \ 0]} \gg$$

is an  $\mathbb{E}$ -triangle. Note that the following diagram

$$\begin{array}{ccccc} A & \xrightarrow{\begin{bmatrix} x \\ a \\ 0 \end{bmatrix}} & B \oplus A' \oplus C & \xrightarrow{\begin{bmatrix} b \ -x' \ 0 \\ 0 \ 0 \ -1 \end{bmatrix}} & B' \oplus C \xrightarrow{[\theta_L \ 0]} \gg \\ \parallel & & \downarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ y & 0 & 1 \end{bmatrix} & & \parallel \\ A & \xrightarrow{\begin{bmatrix} x \\ a \\ yx \end{bmatrix}} & B \oplus A' \oplus C & \xrightarrow{\begin{bmatrix} b \ -x' \ 0 \\ y \ 0 \ -1 \end{bmatrix}} & B' \oplus C \xrightarrow{[\theta_L \ 0]} \gg \end{array}$$

is commutative and all vertical morphisms are isomorphisms, thus we have that

$$A \xrightarrow{\begin{bmatrix} x \\ a \\ yx \end{bmatrix}} B \oplus A' \oplus C \xrightarrow{\begin{bmatrix} b \ -x' \ 0 \\ y \ 0 \ -1 \end{bmatrix}} B' \oplus C \xrightarrow{[\theta_L \ 0]} \gg$$

is also an  $\mathbb{E}$ -triangle. This shows that the square

$$\begin{array}{ccc} A & \xrightarrow{\begin{bmatrix} a \\ yx \end{bmatrix}} & A' \oplus C \\ x \downarrow & & \downarrow \begin{bmatrix} x' \ 0 \\ 0 \ 1 \end{bmatrix} \\ B & \xrightarrow{\begin{bmatrix} b \\ y \end{bmatrix}} & B' \oplus C \end{array}$$

is homotopy cartesian.

Since the square (II) is homotopy cartesian, then there exists an  $\mathbb{E}$ -triangle

$$B \xrightarrow{\begin{bmatrix} b \\ y \end{bmatrix}} B' \oplus C \xrightarrow{[y' \ -c]} C' \xrightarrow{\theta_R} \gg$$

with differential  $\theta_R \in \mathbb{E}(C', B)$ . Consider the following commutative diagram

$$\begin{array}{ccccc} A & \xrightarrow{\begin{bmatrix} a \\ yx \end{bmatrix}} & A' \oplus C & & \\ \downarrow x & & \downarrow \begin{bmatrix} x' & 0 \\ 0 & 1 \end{bmatrix} & & \\ B & \xrightarrow{\begin{bmatrix} b \\ y \end{bmatrix}} & B' \oplus C & \xrightarrow{[y' \ -c]} & C' \xrightarrow{\theta_R} \gg, \end{array}$$

by Lemma 3.1, we have the following morphism of  $\mathbb{E}$ -triangles

$$\begin{array}{ccccc} A & \xrightarrow{\begin{bmatrix} a \\ yx \end{bmatrix}} & A' \oplus C & \xrightarrow{[y'x' \ -c]} & C' \xrightarrow{\theta_P} \gg \\ \downarrow x & & \downarrow \begin{bmatrix} x' & 0 \\ 0 & 1 \end{bmatrix} & & \parallel \\ B & \xrightarrow{\begin{bmatrix} b \\ y \end{bmatrix}} & B' \oplus C & \xrightarrow{[y' \ -c]} & C' \xrightarrow{\theta_R} \gg. \end{array} \quad (3.2)$$

It follows that the  $\mathbb{E}$ -triangle

$$A \xrightarrow{\begin{bmatrix} a \\ yx \end{bmatrix}} A' \oplus C \xrightarrow{[y'x' \ -c]} C' \xrightarrow{\theta_P} \gg$$

shows that the outside rectangle is a homotopy cartesian. Moreover, the diagram (3.2) also shows that  $\theta_R = x_*\theta_P$  holds.

Note that the following diagram

$$\begin{array}{ccccc} A & \xrightarrow{\begin{bmatrix} a \\ x \end{bmatrix}} & A' \oplus B & \xrightarrow{[x' \ -b]} & B' \xrightarrow{\theta_L} \gg \\ \parallel & & \downarrow \begin{bmatrix} 1 & 0 \\ 0 & y \end{bmatrix} & & \downarrow y' \\ A & \xrightarrow{\begin{bmatrix} a \\ yx \end{bmatrix}} & A' \oplus C & \xrightarrow{[y'x' \ -c]} & C' \xrightarrow{\theta_P} \gg \end{array}$$

is a morphism of  $\mathbb{E}$ -triangles, this shows that  $\theta_L = (y')^*\theta_P$  holds.

This completes the proof.  $\square$

**Remark 3.3.** Our main result was proved in [M], when  $\mathcal{C}$  is an abelian category. It also was proved in [CF], when  $\mathcal{C}$  is an triangulated category.

## References

- [CF] J. D. Christensen, M. Frankland. On good morphisms of exact triangles. J. Pure Appl. Algebra 226 (2022), no. 3, 106846, 33 pp.
- [H] J. He. Extensions of covariantly finite subcategories revisited. Czechoslovak Math. J. 69(144) (2019), no. 2, 403–415.

- [HZZ] J. Hu, D. Zhang, P. Zhou. Proper classes and Gorensteinness in extriangulated categories. *J. Algebra* 551 (2020), 23–60.
- [M] S. Mac Lane. *Categories for the working mathematician*. Second edition. Graduate Texts in Mathematics, 5. Springer-Verlag, New York, 1998.
- [N] A. Neeman. *Triangulated categories*. Annals of Mathematics Studies, 148. Princeton University Press, Princeton, NJ, 2001.
- [NP] H. Nakaoka, Y. Palu. Extriangulated categories, Hovey twin cotorsion pairs and model structures. *Cah. Topol. Géom. Différ. Catég.* 60 (2019), no. 2, 117–193.
- [NP1] H. Nakaoka, Y. Palu. External triangulation of the homotopy category of exact quasi-category. arXiv: 2004.02479, 2020.
- [V] J. Verdier. *Des catégories dérivées des catégories abéliennes*. (French. French summary) [On derived categories of abelian categories] With a preface by Luc Illusie. Edited and with a note by Georges Maltsiniotis. Astérisque No. 239 (1996).
- [ZZ] P. Zhou, B. Zhu. Triangulated quotient categories revisited. *J. Algebra* 502 (2018), 196–232.
- [ZhZ] B. Zhu, X. Zhuang. Tilting subcategories in extriangulated categories. *Front. Math. China* 15 (2020), no. 1, 225–253.

**Jing He**

College of Science, Hunan University of Technology and Business, 410205 Changsha, P. R. China

E-mail: jinghe1003@163.com

**Chenbei Xie**

College of Mathematics, Hunan Institute of Science and Technology, 414006 Yueyang, Hunan, P. R. China.

E-mail: xcb19980313@163.com

**Panyue Zhou**

College of Mathematics, Hunan Institute of Science and Technology, 414006 Yueyang, Hunan, P. R. China.

E-mail: panyuezhou@163.com