

Kramers-Moyal analysis of interplanetary magnetic field fluctuations at sub-ion scales

Simone Benella^{1*}, Mirko Stumpo^{2,1}, Giuseppe Consolini¹, Tommaso Alberti¹, Monica Laurenza¹ and Emiliya Yordanova³

¹*INAF - Istituto di Astrofisica e Planetologia Spaziali, Roma, Italy.

²Dipartimento di Fisica, Università degli Studi di Roma Tor Vergata, Roma, Italy.

³Swedish Institute for Space Physics, Uppsala, Sweden.

*Corresponding author(s). E-mail(s): simone.benella@inaf.it;
Contributing authors: mirko.stumpo@inaf.it;
giuseppe.consolini@inaf.it; tommaso.alberti@inaf.it;
monica.laurenza@inaf.it; eya@irfu.se;

Abstract

In the framework of statistical time series analysis of complex dynamics we present a multiscale characterization of solar wind turbulence in the near-Earth environment. The data analysis, based on the Markov-process theory, is meant to estimate the Kramers-Moyal coefficients associated with the measured magnetic field fluctuations. In fact, when the scale-to-scale dynamics can be successfully described as a Markov process, first- and second-order Kramers-Moyal coefficients provide a complete description of the dynamics in terms of Langevin stochastic process. The analysis is carried out by using high-resolution magnetic field measurements gathered by Cluster during a fast solar wind period on January 20, 2007. This analysis extends recent findings in the near-Sun environment with the aim of testing the universality of the Markovian nature of the magnetic field fluctuations in the sub-ion/kinetic domain.

Keywords: solar wind turbulence – Cluster – interplanetary magnetic field – Markov processes

1 Introduction

During last decades, high-resolution magnetic field measurements gathered in the inner Heliosphere have increased the interest in investigating the nature of the magnetic field fluctuations observed at scales below the ion inertial length [1]. Observations gathered in the near-Earth environment were recently accompanied by high-resolution observations in the vicinity of the Sun, thus providing a comprehensive picture of the principal solar wind properties as well as their radial evolution [2, 3]. For interplanetary magnetic field (IMF) fluctuations observed at scales below 0.1-1 Hz, the physical phenomena show the universal features of fully developed turbulence with an important role played by the coupling between magnetic and velocity fields near the Sun and a fluid-like turbulence above ~ 0.6 astronomical units. All these features are consistent with results obtained in the framework of fluid-like approximation of the solar wind turbulence from the integral scale of the energy injection to the ion inertial length d_i [1]. Conversely, there is no general consensus on what are the mechanisms responsible for the IMF fluctuations observed in the range below d_i , namely sub-ion/kinetic range, and thus alternative approaches based on basics of non-equilibrium statistical mechanics might be successful in unveiling important characteristics of such processes [4]. An interesting method for investigating complex time series within the framework of stochastic processes has been developed and widely employed to characterize the energy transfer across the turbulent cascade [5–9]. These authors suggest that the turbulent cascade can be described as a Markov process whose two-points transition probabilities enable us to reproduce the evolution of the Probability Distribution Functions (PDFs) of the longitudinal velocity increments u_ℓ associated with the redistribution of energy across scales. More specifically, the amplitude of u_ℓ at subsequent scales within the inertial range can be represented as a Langevin process with a drift force and a diffusion strength that are scale-dependent. In the field of space plasma turbulence, Strumik and Macek [10, 11] shown that radial IMF fluctuations at typical time scales within the inertial range appear to satisfy the Markov property and the dynamics is accurately reproduced by means of the Fokker-Planck (FP) equation.

An important result by Kiyani *et al.* [12] using ESA-Cluster mission data, pointed out the existence of a global scale-invariance at scales below the ion-inertial length, thus suggesting an intriguing difference with respect to the anomalous scaling, which is ubiquitous in fully developed turbulence, being the fingerprint of intermittency. Analogous results were reported also in the case of the near-Sun environment through the high-resolution observations gathered by the Parker Solar Probe mission during the perihelia [e.g., see 13, and references therein]. Regarding the Markov property in the sub-ion/kinetic range, Benella *et al.* [14] recently showed that IMF fluctuations at these scales keep satisfy such condition in the near-Sun environment, thus sharing with the inertial range the local structure of the energy transfer across scales. However, the two regimes appear to be independent in a statistical sense, i.e. there is no correspondence between the IMF fluctuations measured in the inertial range and

the corresponding fluctuations observed in the sub-ion/kinetic domain, thus supporting the idea that they are originated by different physical mechanisms. It has been shown as the evolution of the statistics within the sub-ion/kinetic range can be modeled through the FP equation. In this case, the observed global-scale invariance can be viewed as the stationary solution of the equation that governs the evolution in scale of the probability distribution functions (PDFs) of the rescaled IMF fluctuations (i.e., IMF fluctuations divided by their standard deviations).

The aims of this work are mainly two:

- The first aim is to extend the analysis of Markov properties proposed in [14], here referred as Kramers-Moyal (KM) analysis, to high-frequency near-Earth observations of solar wind gathered by the ESA-Cluster mission. In fact, the comparison between results obtained in different solar wind conditions at different heliocentric distances is crucial in order to look for an universality of these statistical properties at the sub-ion/kinetic scales;
- The second aim is to provide a general method for estimating the master curve of the rescaled IMF PDFs in the sub-ion/kinetic range by only using the KM coefficients evaluated on the IMF timeseries.

2 Data and Methods

2.1 Data

The data we used in this study were gathered by the ESA-Cluster B spacecraft on 20 January 2007 in a 75 minutes time interval from 12:00 UT to 13:15 UT with a sampling rate of ~ 450 samples/s. This period was characterized by a fast solar wind stream ($v \sim 600$ km⁻¹) with a mean interplanetary magnetic field (IMF) of $B_0 \sim 4$ nT and a plasma density of $n \sim 2$ cm⁻³. Since we are interested in IMF fluctuations parallel and perpendicular with respect to the mean field, we rotated the IMF components in the minimum variance reference frame. Therefore, we focus on the minimum (B_1) and maximum (B_3) variance components, which are mainly aligned along directions parallel and perpendicular to the mean field, respectively. The IMF components are reported in Figure 1 (left panel) along with their power spectral densities (PSDs; right panel). These PSDs have an $f^{-\beta}$ behavior with two distinct regimes separated by a spectral break located around the ion-frequency $f_i \simeq 0.6$ Hz. For frequencies below f_i we observe that the PSD exhibit a scaling *à la* Kolmogorov (i.e., $\beta \sim -5/3$) that is in agreement with the fluid-like description of the solar wind turbulence, whereas for frequencies larger than f_i the spectral trend present the usual sub-ion/kinetic exponent $\beta \in [-7/3, -8/3]$. In the following we perform the analysis in the time domain by assuming that we are exploring spatial scales r via Taylor's hypothesis, i.e., $r = \tau U_0$, being U_0 the average plasma bulk speed.

2.2 Methods

The method we use in this work is based on the Markov process theory. By introducing the increments of the IMF $b_i(\tau) \doteq B(t + \tau) - B(t)$ (where $i = \{1, 2, 3\}$), we can interpret $b_i(\tau)$ as a stochastic processes that evolve across the time scales τ . If the probability of observing the increment $b_{1,i}$ at the scale τ_1 given the increments $b_{2,i}$ at the scale τ_2 until $b_{\tau_n,i}$ at the scale τ_n , with $\tau_1 < \tau_2 < \dots < \tau_n$ does not depend on increments at scales larger than τ_3 the process satisfy the Markov condition, i.e.,

$$p(b_{1,i}, \tau_1 | b_{2,i}, \tau_2; \dots; b_{\tau_n,i}, \tau_n) = p(b_{1,i}, \tau_1 | b_{2,i}, \tau_2). \quad (1)$$

For a Markov process, the transition probability between $b_{1,i}$ and $b_{3,i}$, i.e., the probability of observing the fluctuation $b_{1,i}$ at scale τ_1 given $b_{3,i}$ at scale τ_3 , can be written in terms of the Chapman-Kolmogorov (CK) equation by integrating over fluctuations $b_{2,i}$ at the intermediate scales τ_2 such that $\tau_1 < \tau_2 < \tau_3$ [15],

$$p(b_{1,i}, \tau_1 | b_{3,i}, \tau_3) = \int_{-\infty}^{+\infty} p(b_{1,i}, \tau_1 | b_{2,i}, \tau_2) p(b_{2,i}, \tau_2 | b_{3,i}, \tau_3) db_{2,i}. \quad (2)$$

The time evolution of the transition probability is governed by the *master equation*

$$-\frac{\partial}{\partial \tau} p(b_i, \tau | b'_i, \tau') = \mathcal{L}_{KM}(b_i, \tau) p(b_i, \tau | b'_i, \tau'), \quad (3)$$

where the operator $\mathcal{L}_{KM}(b_i)$ is the Kramers-Moyal (KM) operator and it is defined as

$$\mathcal{L}_{KM}(b_i, \tau) = \sum_{k=1}^{\infty} \left(-\frac{\partial}{\partial b_i} \right)^k D^{(k)}(b_i, \tau). \quad (4)$$

The functions $D^{(k)}(b_i, \tau)$ are called KM coefficients and are defined as

$$D^{(k)}(b_i, \tau) = \frac{1}{k!} \lim_{\delta \rightarrow 0} \frac{1}{\delta} \mathbb{E}[(b_i(\tau - \delta) - b_i(\tau))^k | b_i(\tau)]. \quad (5)$$

The minus sign on the l.h.s. of Equation (3) is due to the direction of the time evolution from larger towards smaller scales [8]. In principle, the KM expansion encloses an infinite number of terms that contribute to the time evolution of the statistics of the process. An important class of stochastic processes is represented by those for which the Pawula's theorem is valid. In fact, the Pawula's theorem states that if $D^{(k)}(b_i, \tau) = 0$, then all the coefficients of order $k \geq 3$ vanish and the KM expansion reduces to the Fokker-Planck (FP) equation [15]

$$-\frac{\partial}{\partial \tau} p(b_i, \tau | b'_i, \tau') = \left[-\frac{\partial}{\partial b_i} D^{(1)}(b_i, \tau) + \frac{\partial^2}{\partial b_i^2} D^{(2)}(b_i, \tau) \right] p(b_i, \tau | b'_i, \tau'). \quad (6)$$




Fig. 1 Left: IMF minimum (blue) and maximum (red) variance components on 20 January 2007 between 12:00 and 13:15. Right: PSD of B_1 (blue) and B_3 (red). Black dotted lines are $-5/3$ and $-8/3$ slopes, whereas the red dotted line indicates the ion-frequency f_i .

The first-order KM coefficient $D^{(1)}(b_i, \tau)$ is representative of the *drift* term, whereas the second-order KM coefficient $D^{(2)}(b_i, \tau)$ is related to the *diffusion* strength, which modulates the amplitude of the delta-correlated Gaussian noise, $\Gamma(\tau)$, in the corresponding Langevin equation

$$-\frac{\partial b_i}{\partial \tau} = D^{(1)}(b_i, \tau) + \sqrt{2D^{(2)}(b_i, \tau)} \Gamma(\tau). \quad (7)$$

Since the true values of KM coefficients are not accessible from data, we will approximate them as their finite-time version for a sufficient small value of δ

$$D_{\delta}^{(k)}(b_i, \tau) = \frac{1}{k! \delta} \mathbb{E}[(b_i(\tau - \delta) - b_i(\tau))^k | b_i(\tau)]. \quad (8)$$

Here, we applied the above analysis to the small-scale increments of the magnetic field at the sub-ion/kinetic scales, starting from the verification of the CK equation (2), and successively evaluating the KM coefficients up to the fourth order.

3 Results

The first step of the analysis is to check the Markov property of B_1 and B_3 fluctuations at sub-ion/kinetic scales. By inspecting the PSD we observe that the instrumental noise affects magnetic field measurements around 100 Hz, thus we set the minimum time scale separation value to $\delta = 0.02$ s. In order to test the validity of the Markov property we evaluate both the l.h.s. and the r.h.s. of the CK equation (Eq. 2) and we compare the resulting transition probabilities. The l.h.s. of Equation (2) represents the empirical transition probability p_E , whereas the r.h.s. is the CK transition probability p_{CK} . The results of the CK test are reported in Figure 2 for three different values of the scale separation: $\tau_3 - \tau_1 = 0.02$ s, $\tau_3 - \tau_1 = 0.2$ s and $\tau_3 - \tau_1 = 2$ s, being $\tau_1 = 0.02$ s. For sake of simplicity, the intermediate scale τ_2 of the CK integral is chosen as the half of the scale separation, although the results are not affected by any particular choice of $\tau_2 \in (\tau_1, \tau_3)$. Figures 2a and 2d show


6 *Kramers-Moyal analysis of IMF fluctuations at sub-ion scales*

Fig. 2 Comparison between p_E (red curves) and p_{CK} (blue curves) at different timescales for minimum (upper panels) and maximum (lower panels) variance directions. The timescale differences for the CK test are 0.02 s (panels a and d), 0.2 s (panels b and e), and 1.0 s (panels c and f).

an excellent agreement between p_E and p_{CK} at 0.02 s scale separation for b_1 and b_3 , respectively. A similar result can be observed at 0.2 s scale separation as reported in Figures 2b and 2e, thus pointing out that the observed process in scale $b_i(\tau)$ can be successfully approximated as a Markov process until the smaller scales that are not affected by instrumental noise. If the scale separation is pushed to 2 s, that corresponds to approach the low end of the inertial range, the CK equation is still valid, but values of the fluctuations b_{i,τ_1} observed at the smaller scale no longer depend on the correspondent values b_{i,τ_3} observed at larger scales. This result support the idea that IMF fluctuations observed in the sub-ion/kinetic and inertial ranges are statistically independent, being perfectly in agreement with previous findings in the near-Sun environment [14].

The next step in the analysis is the evaluation of the first-, second- and fourth-order KM coefficients in order to check whether or not the Pawula's theorem is fulfilled. In Figure 3 are reported the first-, second- and fourth-order KM coefficients for both minimum and maximum variance components with $\tau = 0.06$ s and $\delta = 0.02$. The fourth-order coefficient vanishes for B_1 and B_3 in the case of Figure 3 (i.e., $\tau = 0.06$ s) and for any τ in the sub-ion/kinetic range (not shown), thus confirming that the evolution is scale of statistics of the process $b_i(\tau)$ is governed by the FP Equation (6) with scale-dependent drift and diffusion terms.

One of the striking features of the sub-ion/kinetic range is the global-scale invariance which is generally shown by rescaling the PDFs by introducing the




Fig. 3 First-, second- and fourth-order finite-scale KM coefficients for minimum (left) and maximum (right) variance components of the IMF.

following transformation [12, 13, 16]

$$b_{i,\tau} \rightarrow x_{i,\tau} = b_{i,\tau}/\sigma_{i,\tau} \quad (9)$$

$$p(b_{i,\tau}) \rightarrow p(x_{i,\tau}) = \sigma_{i,\tau} p(b_{i,\tau}). \quad (10)$$

The existence of a master curve, i.e., a PDF invariant in shape, can be interpreted in terms of stationary solution of the FP Equation (6) across sub-ion/kinetic scales [14]. We aim to derive such master curve starting from the KM coefficients evaluated for $b_{i,\tau}$. Since we are dealing with finite-time KM coefficients instead of their “true” value (i.e. taking the limit $\delta \rightarrow 0$), the relation between the KM coefficients $D^{(k)}(b_i, \tau)$ and $D^{(k)}(x_i, \tau)$ is not straightforward. By substituting the transformation (10) in Equation (8), we obtain the following relations between the KM coefficients

$$D_\delta^{(1)}(x_i, \tau) = \frac{D_\delta^{(1)}(b_i, \tau)}{\sigma} + \frac{1}{\delta} \left(\frac{1}{\sigma'} - \frac{1}{\sigma} \right) \mathbb{E}[b_i(\tau - \delta)|b_i(\tau)], \quad (11)$$

$$D_\delta^{(2)}(x_i, \tau) = \frac{D_\delta^{(2)}(b_i, \tau)}{\sigma^2} + \frac{1}{2\delta} \left(\frac{1}{\sigma'^2} - \frac{1}{\sigma^2} \right) \mathbb{E}[b_i^2(\tau - \delta)|b_i(\tau)] + \frac{1}{\delta\sigma} \left(\frac{1}{\sigma'} - \frac{1}{\sigma} \right) \mathbb{E}[b_i(\tau - \delta)b(\tau)|b_i(\tau)], \quad (12)$$

where σ and σ' are the standard deviations of the IMF fluctuations at the scales τ and $\tau - \delta$, respectively. Starting from the calculation of all the finite-scale KM coefficients at different time scales within the sub-ion/kinetic range,


8 *Kramers-Moyal analysis of IMF fluctuations at sub-ion scales*

Fig. 4 Comparison between the empirical PDF of the rescaled variable x_i (circles) and the stationary solution of the FP equation (black lines) for the minimum (a) and maximum (b) variance components. The time scales used in the sub-ion/kinetic range range from 0.02 s to 0.2 s.

Equations (11) and (12) enable us to evaluate all the corresponding finite-scale KM coefficients of the rescaled variables. By solving the FP equation under for the stationary PDF, the general solution reads

$$p_{ST}(x_i) = \mathcal{N}_i \exp \left\{ -\ln D^{(2)}(x_i) + \int_{-\infty}^{x_i} \frac{D^{(1)}(\xi)}{D^{(2)}(\xi)} d\xi \right\} \quad (13)$$

where \mathcal{N}_i is the normalization factor. A fundamental remark of our reasoning is that by evaluating the functions $D^{(1)}(x_i, \tau)$ and $D^{(2)}(x_i, \tau)$ through Equations (11) and (12) at any τ and by substituting them in Equation (13), the obtained stationary solution $p_{ST}(x_i)$ approximates the empirical master curve and does not depend on τ . Figure 4ab shows the average of the functions $p_{ST}(x_i)$ obtained through Equation (13) within the sub-ion/kinetic range (black lines) along with the empirical rescaled PDFs (circles). The stationary PDFs exhibit an excellent agreement with observations within $x_i \in [-5\sigma_{x_i}, 5\sigma_{x_i}]$ for both minimum and maximum variance components. We want to emphasize that the steady state solution has to be interpreted as the analogous of the *PDF-invariant shape* observed in the sub-ion/kinetic range, which stems from the fact that the FP Equation (6) describe the evolution of PDFs as a function of the time scale τ . It has been shown that by parametrizing the KM coefficients as a linear ($D^{(1)}(x_i)$) and quadratic ($D^{(2)}(x_i)$) functions of x_i , the master curve of Figure 4 can be written as a Kappa distribution [14]. In this work we present a derivation of the steady state solution in the rescaled variable x_i that is based on the evaluation of KM coefficients on the variable b_i at any τ the sub-ion/kinetic regime.

4 Discussion and conclusions

An universal framework of the solar wind processes originating magnetic field fluctuations at sub-ion/kinetic scales is still lacking. The analysis of Markov properties and the KM coefficients associated with such fluctuations can therefore provide an important method enabling us to get new insights on the characteristics of the physical processes acting at these scales. In this work we consider a fast solar wind stream in the near-Earth environment and we have given evidence that the description of the PDF evolution as a function of the time scale τ by means of Markov processes is successful at sub-ion/kinetic scales. Such property has been also verified for IMF fluctuations within the inertial range of magneto-fluid turbulence [10, 11], then supporting the idea that the energy flow from one scale to another has a local structure in both inertial and sub-ion/kinetic regimes. Although there is a net energy transfer between them, we provided evidence that a correspondence in terms of IMF fluctuations between inertial and sub-ion/kinetic ranges does not seem to emerge, confirming recent findings in the near-Sun environment [14]. This statistical independence between the observed fluctuations, together with the transition between different dynamical regimes observed in the PSD, supports the idea that the physical processes originating IMF fluctuations have a different nature in the two regimes.

The main result reported in this work is represented by the clear Markovian nature of the IMF fluctuations in the sub-ion/kinetic range in the near-Earth solar wind. As pointed out in several recent works, in fact, many solar wind characteristics, i.e. IMF fluctuations, exhibit a radial dependence in terms of spectral features [2], intermittency [3], entropic character [17] and so forth. By performing the KM analysis in the near-Earth IMF fluctuations we obtained similar results in comparison with the near-Sun solar wind conditions. This allows us to assert that such Markovian property of IMF fluctuations does not depend on the heliocentric distance and thus might represent a universal property in the sub-ion/kinetic domain.

The solar wind inertial range shows the typical anomalous scaling of fully developed turbulence, with an intermittent character of the energy transfer, whose strength depends on the heliocentric distance from the Sun [3]. In terms of scale dynamics of the IMF fluctuation statistics, the intermittency is associated with a modification of the PDF across the inertial range. This process affects both IMF-fluctuation amplitude as well as the shape of their PDFs. Indeed, the higher probability of observing enhanced IMF fluctuations towards smaller scales is reflected in the modification of the Gaussian PDF at the integral scale into a leptokurtic function through the energy cascade. In the framework of Markov processes, the PDF dynamics across the inertial scales can be modeled through the general FP equation, accounting for the variation of the shape of the PDF as a function of τ . As a consequence, the absence of modification of the initial PDF across different scales can be due to the lack of intermittency in the physical processes accounting for the energy transfer. In this work we confirmed that the global scaling observed

at sub-ion/kinetic scales can be interpreted by means of the stationary FP equation (i.e., $\partial p/\partial \tau \rightarrow 0$) since the PDFs of IMF fluctuations exhibit a well-defined shape-invariant form in this domain. From the physical point of view our results can be resembled in the following scenario: the energy transfer mechanisms in the inertial range and in the sub-ion/kinetic domain have a different nature, therefore they originate IMF fluctuations exhibiting a statistical independence between the two regimes and the intermittent nature of IMF fluctuations in the inertial range is lost at sub-ion/kinetic scales. A joint analysis involving both high-resolution magnetic and velocity field is desirable and will provide a more comprehensive picture of the physical processes acting at sub-ion/kinetic scales.

Acknowledgments. This work is funded by the Italian MIUR-PRIN grant 2017APKP7T on “Circumterrestrial Environment: Impact of Sun-Earth Interaction”. The authors acknowledge the Cluster FGM and STAFF P.I.s and teams and the ESA-Cluster Archive for making available the data used in this work. The FGM and STAFF data were combined using the IRFU-Matlab analysis package available at <https://github.com/irfu/irfu-matlab>. M.S. acknowledges the PhD course in Astronomy, Astrophysics and Space Science of the University of Rome “Sapienza”, University of Rome “Tor Vergata” and Italian National Institute for Astrophysics (INAF), Italy. E.Y. is supported by SNSA grants 86/20 and 145/18.

References

- [1] Bruno, R., Carbone, V.: Turbulence in the Solar Wind vol. 928, (2016). <https://doi.org/10.1007/978-3-319-43440-7>
- [2] Chen, C., Bale, S., Bonnell, J., Borovikov, D., Bowen, T., Burgess, D., Case, A., Chandran, B., De Wit, T.D., Goetz, K., *et al.*: The evolution and role of solar wind turbulence in the inner heliosphere. The Astrophysical Journal Supplement Series **246**(2), 53 (2020)
- [3] Alberti, T., Laurenza, M., Consolini, G., Milillo, A., Marcucci, M.F., Carbone, V., Bale, S.D.: On the scaling properties of magnetic-field fluctuations through the inner heliosphere. The Astrophysical Journal **902**(1), 84 (2020). <https://doi.org/10.3847/1538-4357/abb3d2>
- [4] Carbone, V., Telloni, D., Lepreti, F., Vecchio, A.: High-frequency magnetic fluctuations in space plasmas and the role of electron Landau damping. The Astrophysical Journal Letters **924**(2), 26 (2022). <https://doi.org/10.3847/2041-8213/ac4740>
- [5] Pedrizzetti, G., Novikov, E.A.: On Markov modelling of turbulence. Journal of Fluid Mechanics **280**, 69–93 (1994)

- [6] Friedrich, R., Peinke, J.: Description of a turbulent cascade by a fokker-planck equation. *Physical Review Letters* **78**(5), 863 (1997)
- [7] Friedrich, R., Peinke, J.: Statistical properties of a turbulent cascade. *Physica D: Nonlinear Phenomena* **102**(1-2), 147–155 (1997)
- [8] Renner, C., Peinke, J., Friedrich, R.: Experimental indications for markov properties of small-scale turbulence. *Journal of Fluid Mechanics* **433**, 383 (2001)
- [9] Peinke, J., Tabar, M.R., Wächter, M.: The fokker–planck approach to complex spatiotemporal disordered systems. *Annual Review of Condensed Matter Physics* **10**, 107–132 (2019)
- [10] Strumik, M., Macek, W.: Statistical analysis of transfer of fluctuations in solar wind turbulence. *Nonlinear Processes in Geophysics* **15**(4), 607–613 (2008)
- [11] Strumik, M., Macek, W.M.: Testing for markovian character and modeling of intermittency in solar wind turbulence. *Physical Review E* **78**(2), 026414 (2008)
- [12] Kiyani, K., Chapman, S., Khotyaintsev, Y.V., Dunlop, M., Sahraoui, F.: Global scale-invariant dissipation in collisionless plasma turbulence. *Physical Review Letters* **103**(7), 075006 (2009)
- [13] Chhiber, R., Matthaeus, W.H., Bowen, T.A., Bale, S.D.: Subproton-scale intermittency in near-sun solar wind turbulence observed by the parker solar probe. *The Astrophysical Journal Letters* **911**(1), 7 (2021)
- [14] Benella, S., Stumpo, M., Consolini, G., Alberti, T., Carbone, V., Laurenza, M.: Markovian features of the solar wind at subproton scales. *The Astrophysical Journal Letters* **928**(2), 21 (2022). <https://doi.org/10.3847/2041-8213/ac6107>
- [15] Risken, H., Caughey, T.K.: The Fokker-Planck Equation: Methods of Solution and Application, 2nd ed. *Journal of Applied Mechanics* **58**(3), 860 (1991). <https://doi.org/10.1115/1.2897281>
- [16] Alberti, T., Consolini, G., Carbone, V., Yordanova, E., Marcucci, M.F., De Michelis, P.: Multifractal and chaotic properties of solar wind at mhd and kinetic domains: An empirical mode decomposition approach. *Entropy* **21**(3), 320 (2019)
- [17] Stumpo, M., Quattrociochi, V., Benella, S., Alberti, T., Consolini, G.: Self-organization through the inner heliosphere: Insights from parker solar probe. *Atmosphere* **12**(3), 321 (2021)