ON THE NON-EXISTENCE OF SPECIAL GENERIC MAPS ON COMPLEX PROJECTIVE SPACES

NAOKI KITAZAWA

ABSTRACT. We prove the non-existence of *special generic* maps on complex projective space as our extended new result. Simplest special generic maps are Morse functions with exactly two singular points on spheres, or Morse functions in Reeb's theorem, and canonical projections of unit spheres.

Manifolds represented as connected sums of products of manifolds diffeomorphic to unit spheres admit such maps in considerable cases.

Real and complex projective spaces have been shown to admit no such maps in most cases by the author. This gives a complete answer for complex projective spaces as a corollary to a more general result, which is also our main result.

1. INTRODUCTION.

Morse theory, or theory of Morse functions, gives fundamental and strong tools in singularity theory and its applications to geometry of manifolds.

A *singular* point of a smooth map means a point in the manifold of the domain where the rank of the differential is smaller than both the manifolds of the domain and the target.

Theorem 1. A closed manifold admits a Morse function with exactly two singular points if and only if it is homeomorphic to a sphere in the case where the manifold is not 4-dimensional. In the 4-dimensional case, this holds if it is diffeomorphic to the unit sphere.

These functions are generalized as follows.

Definition 1. A smooth map from an *m*-dimensional manifold with no boundary into an *n*-dimensional manifold with no boundary is called a *special generic* map if $m \ge n \ge 1$ and for suitable coordinates, at each singular point it is represented by the form $(x_1, \dots, x_m) \to (x_1, \dots, x_{n-1}, \sum_{j=1}^{m-n+1} x_{n-1+j}^2)$

The definition seems to restrict the topology and the differntiable structure of the manifold strongly. This is true.

Before explaining about this, we introduce fundamental terminologies and notation on smooth manifolds. \mathbb{R}^k denotes the k-dimensional Euclidean space with the canonical differentiable structure. This is a simplest smooth manifold. This is also regarded as the Riemannian manifold endowed with the standard Euclidean metric. For $x \in \mathbb{R}^k$, ||x|| denotes the distance between x and the origin $0 \in \mathbb{R}^k$ under the standard metric or the canonical Euclidean metric. For a positive integer

Key words and phrases. Special generic maps. (Co)homology. (Complex) projective spaces. Closed and simply-connected manifolds.

²⁰²⁰ Mathematics Subject Classification: Primary 57R45. Secondary 57R19.

 $k, D^k := \{x \mid ||x|| \leq 1\} \subset \mathbb{R}^k$ denotes the k-dimensional unit disk, which is a k-dimensional compact, connected and smooth closed submanifold. $S^{k-1} := \{x \mid ||x|| = 1\} \subset \mathbb{R}^k$ denotes the (k-1)-dimensional unit sphere, which is a (k-1)-dimensional closed smooth submanifold with no boundary. The 0-dimensional unit sphere is a two-point set endowed with the discrete topology for k = 1 (k-1=0). The (k-1)-dimensional unit sphere is connected for $k \geq 2$ $(k-1 \geq 1)$.

Canonical projections of unit spheres are special generic. *Exotic* spheres, which are defined as manifolds which are homeomorphic to (unit) spheres and which are not diffeomorphic to unit spheres, do not admit such maps in considerable cases. [2, 20, 21, 22, 23, 27] give related answers for example.

On the other hands, for example, manifolds represented as connected sums of manifolds diffeomorphic to unit spheres admit these maps in considerable cases.

It is important and difficult problems to know the (non-)existence of special generic maps.

For example, it seems to be natural to consider such problems for projective spaces, 2, 3 or 4-dimensional closed manifolds and higher dimensional closed (and simply-connected) manifolds for example. In fact related results such as [20, 21, 22, 23, 25] have been obtained by Saeki and Sakuma since 1990. Recently Nishioka ([19]), Wrazidlo ([27, 28, 29]) and the author ([13, 14, 15, 16]) have obtained related results. Our paper concerns such problems for complex projective spaces. The (non-)existence for real-projective spaces has been announced to be completely solved by the author. In short, existence is rare.

Main Theorem 1. Let $k \ge 2$ be an arbitrary integer. The k-dimensional complex projective space, which is a 2k-dimensional closed and simply-connected smooth manifold, admits no special generic maps into \mathbb{R}^{2k-1} . Furthermore, by Theorem 4, it admits no special generic maps into any Euclidean space.

This was an open problem for an arbitrary odd integer $k \ge 3$ in [15]. This follows from the following more general theorem. Terminologies and notation on homology groups and cohomology groups and rings are in section 3. However, we suppose elementary knowledge on these notions.

Here we note that \mathbb{R}^1 is generally denoted by \mathbb{R} , that the ring of all integers (rational numbers) is denoted by \mathbb{Z} (resp. \mathbb{Q}), that the relation $\mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R}$ holds and that \mathbb{Q} is the field of quotients (fractions) of the free commutative group \mathbb{Z} , whose rank is 1.

Main Theorem 2. Let k_1 and k_2 be arbitrary integers greater than 1. Let M be a closed and simply-connected smooth manifold of dimension $m = k_1k_2$ satisfying the following conditions.

- (1) The rational cohomology group $H^j(M; \mathbb{Q})$ is isomorphic to \mathbb{Q} if $0 \leq j \leq k_1k_2$ and j is divisible by k_1 and it is the trivial group if j is not as before.
- (2) There exists a generator u_0 of $H^{k_1}(M;\mathbb{Q})$ and the cup product $\bigcup_{j_1=1}^{j_2} u_0$, or the j_2 -th power of u_0 in the rational cohomology ring $H^*(M;\mathbb{Q})$ is a generator of $H^{j_2k_1}(M;\mathbb{Q})$ for every integer $1 \le j_2 \le k_2 - 1$.

Then M admits no special generic maps into \mathbb{R}^{m-1} .

The second section is for fundamental properties of special generic maps. We also review results related to our present study more precisely.

The third section is devoted to expositions on Main Theorems 1 and 2. In fact, a key ingredient is merely a slight generalization of a family of arguments used by the author in [14, 15, 16]. We close our paper by related examples and remarks.

2. Preliminaries.

Throughout the present paper, a smooth manifold homeomorphic to a unit sphere is called a *homotopy sphere*. A homotopy sphere which is (not) diffeomorphic to a unit sphere is a *standard* sphere (resp. an *exotic* sphere).

Every smooth manifold is well-known to admit the structure of a PL manifold canonically. In several scenes, we regard smooth manifolds as the canonical PL manifolds. More generally, topological manifolds are known to be regarded as CW complexes.

For a manifold or a polyhedron X, let dim X denote the dimension of X.

A diffeomorphism is a smooth map having no singular points being also a homeomorphism. A diffeomorphism from a given manifold X onto the same manifold is said to be a diffeomorphism on X. The diffeomorphism group of the smooth manifold X is the group of all diffeomorphisms on X. We also regard this as the topological space endowed with the so-called Whitney C^{∞} topology. Whitney C^{∞} topologies are natural topologies of sets of smooth maps between smooth manifolds. See [7] for systematic expositions for example.

We also need terminologies and notions on bundles. For systematic expositions, see [18, 26].

A bundle is said to be *smooth* if the fiber is a smooth manifold and the structure group is a subgroup of the diffeomorphism group. A *linear* bundle is a smooth bundle whose fiber is an Euclidean space, a unit sphere, or a unit disk and whose structure group consists of diffeomorphisms regarded as linear transformations naturally.

Proposition 1 (E. g. [20]). Let $m \ge n \ge 1$ be integers. A special generic map $f : M \to N$ on an m-dimensional closed manifold M into an n-dimensional manifold N with no boundary always enjoys the following properties.

- (1) There exist an n-dimensional compact manifold W_f , a smooth immersion $\overline{f}: W_f \to N$ and a smooth surjection $q_f: M \to W_f$ satisfying the relation $f = \overline{f} \circ q_f$.
- (2) q_f maps the singular set S(f) of f onto the boundary $\partial W_f \subset W_f$ as a diffeomorphism. Moreover, S(f) is a smooth closed submanifold in M with no boundary.
- (3) We have the following two bundles.
 - (a) For a suitable small collar neighborhood N(∂W_f) ⊂ W_f, we have a linear bundle whose fiber is the (m − n + 1)-dimensional unit disk and whose projection is given by the composition of the map q_f|_{q_f-1(N(∂W_f))} onto N(∂W_f) with the canonical projection to ∂W_f.
 - (b) We have a smooth bundle over W_f Int $N(\partial W_f)$ whose fiber is an (m-n)-dimensional standard sphere and whose projection is given by the restriction of q_f to the preimage of W_f Int $N(\partial W_f)$.

Hereafter, |X| denotes the size of a finite set X.

Example 1. Let $m \ge n \ge 2$ be integers. Let $\{S^{k_j} \times S^{m-k_j}\}_{j \in J}$ be a family of finitely many products of two unit spheres where k_j is an integer satisfying

 $1 \leq k_j \leq n-1$. We consider a connected sum of these |J| manifolds in the smooth category. Let M_0 be an *m*-dimensional closed and connected manifold diffeomorphic to the obtained manifold. We have a special generic map f_0 so that W_{f_0} is diffeomorphic to a manifold represented as a boundary connected sum of |J| manifolds in $\{S^{k_j} \times D^{n-k_j}\}_{j \in J}$ just before. Here the boundary connected sum is considered in the smooth category. Furthermore, here we can do in such a way that $N = \mathbb{R}^n$, that $f_0|_{S(f_0)}$ is an embedding and that the two bundles of Proposition 1 (3) are trivial bundles.

Proposition 2 (E. g. [20]). Let $m > n \ge 1$ be integers. A special generic map $f: M \to N$ on an m-dimensional closed and connected manifold M into an n-dimensional manifold N with no boundary always enjoys the following properties. We abuse the notation of Proposition 1 such as $q_f: M \to W_f$.

- (1) We have an (m + 1)-dimensional compact and connected topological manifold (PL) manifold W bounded by M, (resp. where we discuss in the PL category), and which collapses to an n-dimensional compact and smooth manifold W_f in the situation of Proposition 1 and identified with a suitable CW subcomplex (resp. subpolyhedron) of W. If m - n = 1, 2, 3 in addition, then W can be chosen as a smooth manifold and r as a smooth map.
- (2) The canonical inclusion $i_M : M \to W$ and a continuous (resp. PL) map $r_f : W \to W_f$ giving a collapsing to W_f satisfy the relation $q_f = r \circ i_M$.
- (3) If M is oriented, then W can be chosen as an oriented manifold. If m-n = 1, 2, 3 in addition, then W can be chosen as a smooth manifold and r as a smooth map as just before.

We give some remarks on Proposition 2 as Remark 1 in the following.

Remark 1. Let $m > n \ge 1$ be integers again. Suppose that we have a smooth immersion $\overline{f}_0 : W_{f_0} \to N$ of an *n*-dimensional compact and connected smooth manifold W_{f_0} into an *n*-dimensional manifold N with no boundary. We have a special generic map $f_0 : M_0 \to N$ on a suitable *m*-dimensional closed and connected manifold M_0 into N enjoying the three properties in Proposition 1 and we can have an (m+1)-dimensional compact and connected smooth manifold W_0 and a smooth map $r_0 : W_0 \to W_{f_0}$ as in Proposition 2 where $f : M \to N$, W and $r : W \to W_f$ are replaced by $f_0 : M_0 \to N$, W_0 and $r_0 : W_0 \to W_{f_0}$ for example. Furthermore, we can do in such a way that the bundles in Proposition 1 (3) are trivial bundles here.

For more generalized versions of Proposition 2, see [24] and see papers [9, 10, 11] by the author.

Note also that we concentrate on the case m - n = 1 in main ingredients and we discuss problems in the smooth category in situations of Proposition 2 in main ingredients of our paper.

Theorem 2. Let m > 0 be an integer. We have the following conditions for closed manifolds to admit special generic maps into fixed Euclidean spaces where connected sums and boundary connected sums are taken in the smooth category. We also abuse the notation in Propositions 1 and 2 such as the manifold W_f .

(1) ([20]) Let $m \ge 2$. An m-dimensional closed and connected manifold admits a special generic map into \mathbb{R}^2 if and only if either of the following cases holds.

- (a) $m \neq 4$ and M is homeomorphic to the m-dimensional unit sphere.
- (b) m = 4 and M is an m-dimensional standard sphere.
- (c) M is diffeomorphic to a manifold represented as a connected sum of the total spaces of smooth bundles over a circle or the 1-dimensional unit sphere whose fibers are diffeomorphic to homotopy spheres which are not 4-dimensional exotic spheres.

Furthermore, for a special generic map f here, W_f is diffeomorphic to a boundary connected sum of finitely many copies of $S^1 \times D^1$.

- (2) ([20]) Let m = 4, 5. An *m*-dimensional closed and simply-connected manifold admits a special generic map into \mathbb{R}^3 if and only if either of the following two holds.
 - (a) M is an m-dimensional standard sphere.
 - (b) M is diffeomorphic to a manifold represented as a connected sum of the total spaces of smooth (linear) bundles over the 2-dimensional unit sphere whose fibers are 3-dimensional standard spheres.

Furthermore, for a special generic map f here, W_f is diffeomorphic to a boundary connected sum of finitely many copies of $S^2 \times D^1$.

- (3) ([19]) Let m = 5. An m-dimensional closed and simply-connected manifold admits a special generic map into ℝ⁴ if and only if either of the following two holds.
 - (a) M is an m-dimensional standard sphere.
 - (b) M is diffeomorphic to a manifold represented as a connected sum of the total spaces of smooth (linear) bundles over the 2-dimensional unit sphere whose fibers are 3-dimensional standard spheres.

For studies on special generic maps, see also [1, 6] as pioneering ones, for example.

3. The proof of Main Theorems.

3.1. Complex projective spaces whose dimensions are greater than 1 do not admit special generic maps into the Euclidean spaces whose codimensions are -1. $\mathbb{C}P^k$ denotes the k-dimensional complex projective space, which is a k-dimensional complex space and 2k-dimensional smooth manifold.

Theorem 3. Let m = 2k > 2 be an even integer greater than 2. $\mathbb{C}P^k$ does not admit special generic maps into \mathbb{R}^{m-1} .

3.2. Fundamental algeberaic topology. We introduce notation and terminologies on *homology groups*, *cohomology groups* and *rings* and *homotopy groups*. For elementary or advanced theory, consult [8] for example.

Let A be a commutative ring. Let (X, X') be a pair of topological spaces satisfying $X' \subset X$ where X' can be the empty set.

The k-th homology group (cohomology group) whose coefficient ring is A is denoted by $H_k(X, X'; A)$ (resp. $H^k(X, X'; A)$). If A is isomorphic to \mathbb{Z} (resp. \mathbb{Q}), then we add "integral" (resp. "rational") after "k-th". If X' is empty, then we may omit ",X'" in the notation and the k-th homology group (cohomology group) of the pair (X, X') is also called the k-th homology group (resp. cohomology group) of X. We can define integral and rational ones similarly. (j-th) (co)homology classes of (X, X') (or X) whose coefficient ring is A are elements of the (j-th) (resp. co)homology groups whose coefficient rings are A. We can define integral and rational (co)homology classes similarly.

 $\pi_k(X)$ denotes the k-th homotopy group of a topological space X.

Let (X_1, X_1') and (X_2, X_2') be pairs of topological spaces satisfying $X_1' \subset X_1$ and $X_2' \subset X_2$ where the second topological space of each pair can be empty as before. For a continuous map $c : X_1 \to X_2$ satisfying $c(X_1') \subset X_2'$, $c_* :$ $H_*(X_1, X_1'; A) \to H_*(X_2, X_2'; A)$ and $c^* : H^*(X_2, X_2'; A) \to H^*(X_1, X_1'; A)$ denote the homomorphisms defined in a canonical way. For a continuous map c : $X_1 \to X_2, c_* : \pi_k(X_1) \to \pi_k(X_2)$ also denotes the homomorphism between the homotopy groups of degree k, which is also defined in a canonical way.

Let $H^*(X; A)$ denote the direct sum $\bigoplus_j H^j(X; A)$ for every integer $j \ge 0$. The cup product for a sequence $\{c_j\}_{j=1}^l \subset H^*(X; A)$ of l > 0 cohomology classes is important. $\bigcup_{j=1}^l c_j$ denotes this. This gives $H^*(X; A)$ the structure of a graded commutative algebra over A and this is the cohomology ring of X whose coefficient ring is A. For a pair $c_1, c_2 \in H^*(X; A)$, we also use the notation such as $c_1 \sqcup c_2$ for example. We define integral and rational cohomology rings similarly.

The fundamental class of a compact, connected and oriented manifold Y is the $(\dim Y)$ -th homology class, defined as the generator of the group $H_{\dim Y}(Y, \partial Y; A)$, isomorphic to A, being compatible with the orientation.

Let $i_{Y,X} : Y \to X$ be a smooth, PL or topologically flat embedding mapping the boundary ∂Y into the boundary ∂X and the interior Int Y into the interior Int X. In short, Y is embedded *properly* in X. Let $h \in H_{\dim Y}(X, \partial X; A)$. If the value of the homomorphism i_{Y,X_*} induced by the embedding $i_{Y,X} : Y \to X$ at the fundamental class of Y is equal to h, then h is said to be *represented* by the (oriented) submanifold Y.

A may be $\mathbb{Z}/2\mathbb{Z}$, the field of order 2, or more generally, the commutative ring consisting of elements whose orders are at most 2. In this case, we may ignore orientations.

We also need the notions of *Poincaré duals* in the following four.

- The Poincaré dual $PD(c_h) \in H^{\dim Y-j}(Y, \partial Y; A)$ to a homology class $c_h \in H_j(Y; A)$.
- The *Poincaré* dual $PD(c_c) \in H_{\dim Y-j}(Y, \partial Y; A)$ to a cohomology class $c_c \in H^j(Y; A)$.
- The *Poincaré* dual $PD(c_h) \in H^{\dim Y-j}(Y; A)$ to a homology class $c_h \in H_j(Y, \partial Y; A)$.
- The Poincaré dual $PD(c_c) \in H^{\dim Y-j}(Y; A)$ to a cohomology class $c_c \in H^j(Y, \partial Y; A)$.

Related to this, Poincaré duality theorem for the manifold Y is also important.

We can define the *dual* to a homology class of a basis of a free submodule of a homology group of $(X, X' \subset X)$ where we consider modules over A. Note that X' may be empty. It is a uniquely defined cohomology class in the cohomology group of the pair (X, X') in such a way that the degrees of the homology class and the cohomology class are same.

3.3. **Proving Theorem 3 and Main Theorems.** Theorem 3 with the following theorem yields Main Theorem 1.

Theorem 4 ([13]). Let $m > n \ge 1$ and l > 0 be integers. Let A be commutative ring. Let M be an m-dimensional closed and connected manifold. We suppose the existence of a sequence $\{a_j\}_{j=1}^l \subset H^*(M; A)$ of cohomology classes satisfying the following two conditions.

 $\mathbf{6}$

- The cup product $\cup_{j=1}^{l} a_j$ is not the zero element.
- The degree of each cohomology class in the sequence $\{a_j\}_{j=1}^l$ is smaller than or equal to m n and the sum of the degrees of these l cohomology classes is greater than or equal to n.

Then M does not admit special generic maps into any n-dimensional connected manifold which is not compact and which has no boundary.

We review the original proof in [13]. An essential ingredient of the proof has been shown in [12] first. This is also reviewed in [14, 15, 16].

We omit precise expositions on *handles* and their *indice* for polyhedra including PL manifolds.

Reviewing the proof of Theorem 4. We abuse the notation in Propositions 1 and 2 for example.

Suppose that M admits a special generic map into an n-dimensional connected manifold N which is non-closed and which has no boundary. We can take an (m+1)-dimensional compact and connected PL manifold W in Proposition 2: Remark 1 says that we consider the case m - n = 1, 2, 3 only and discuss problems in the smooth category and and we may restrict the case as this.

 W_f is an n-dimensional compact and connected manifold smoothly immersed into the connected and non-closed manifold N with no boundary via $\overline{f}: W_f \to N$. As a result it is (simple) homotopy equivalent to an (n-1)-dimensional compact and connected polyhedron. W is (simple) homotopy equivalent to W_f . We can see that W is a PL manifold obtained by attaching handle to $M \times \{0\} \subset M \times [-1, 0]$ whose indices are greater than $(m+1) - \dim W_f = m - n + 1$ where the boundary $\partial W = M$ can be identified with $M \times \{-1\} \subset M \times [-1, 0]$ naturally. We can take a unique cohomology class $b_j \in H^*(W; A)$ satisfying the relation $a_j = i_M^*(b_j)$. W is (simple) homotopy equivalent to an (n-1)-dimensional polyhedron. This means that the cup product $\cup_{j=1}^l a_j$ is the zero element. This contradicts the assumption. This completes the proof.

We show Theorem 3 (Main Theorem 1) and Main Theorem 2

A proof of Theorem 3 (Main Theorem 1) and Main Theorem 2. We prove results through several STEPs.

STEP 1 Preparations.

We prove theorems by adopting methods in [14, 15, 16, 17] together with fundamental theory in [20] for example. However, we adopt shorter expositions or different ones. We abuse the notation in Propositions 1 and 2 such as a special generic map $f: M \to N := \mathbb{R}^n = \mathbb{R}^{m-1}$, the surjective smooth map $q_f: M \to W_f$ onto an (m-1)-dimensional compact and connected manifold W_f smoothly immersed into \mathbb{R}^{m-1} via a smooth immersion $\overline{f}: W_f \to \mathbb{R}^n$, an (m+1)-dimensional compact and connected smooth manifold W with a smooth map $r: W \to W_f$. Assume the existence of such a map $f: M \to \mathbb{R}^n$.

The argument on handles to obtain W from M or $M \times [-1,0]$ implies that M, W and W_f are simply-connected. The canonical inclusion $i_M : M \to W$ and $r : W \to W_f$ induce isomorphisms $i_{M*} : \pi_1(M) \to \pi_1(W)$ and $q_{f*} : \pi_1(M) \to \pi_1(W_f)$.

For $r: W \to W_f$, we have the following objects and properties.

- There exists an (m+1)-dimensional compact and connected smooth manifold $\tilde{W} \supset W$ which is cornered.
- $\tilde{r}: \tilde{W} \to W_f$ is regarded as the projection of a bundle over W_f whose fiber is (PL) homeomorphic to the unit disk of dimension m n + 1 = m (m 1) + 1 = 2 and whose structure group consists of (resp. PL) homeomorphisms of the disk.
- W collapses to W, which is regarded as a CW subcomplex (resp. subpolyhedron) of \tilde{W} , and these two manifolds are (resp. PL) homeomorphic.
- $\tilde{r}^{-1}(W_f \text{Int } N(\partial W_f)) = r^{-1}(W_f \text{Int } N(\partial W_f))$ in Proposition 2.
- The preimage of W_f Int $N(\partial W_f)$ by r in Proposition 2 is unchanged in the collapsing of $\tilde{W} \supset \tilde{r}^{-1}(W_f - \operatorname{Int} N(\partial W_f)) = r^{-1}(W_f - \operatorname{Int} N(\partial W_f))$ to $W \supset r^{-1}(W_f - \operatorname{Int} N(\partial W_f)) = \tilde{r}^{-1}(W_f - \operatorname{Int} N(\partial W_f))$. The restrictions of r and \tilde{r} to these subsets agree.

STEP 2 Supposing that the singular set S(f) is not connected and a proof of the fact that this is connected in the case $k_1 > 2$.

Suppose that the singular set S(f) is not connected. Then there exist a 1-dimensional compact and connected manifold L smoothly and properly embedded into W_f and two distinct connected components C_1 and C_2 of the boundary ∂W_f and $L \cap C_j$ is not empty for j = 1, 2. In other words, the boundary is embedded in the subset $C_1 \bigcup C_2$ of the boundary ∂W_f and the interior is embedded in the interior Int W_f . The preimage $q_f^{-1}(L)$ can be regarded as a 2-dimensional standard sphere. It is regarded as the 2-dimensional manifold of the domain of a Morse function for the Reeb's theorem. The (m-2)-th integral homology class represented by S(f) is regarded as the Poincaré dual to the integral cohomology class regarded as the dual to a homology class represented by the 2-dimensional sphere if the rank of $H^2(M;\mathbb{Q})$ is not 0. Note that this is the case of $k_1 = 2$. More precisely, in this case the rank is 1. We can define the dual to the 2nd integral homology class in a suitable way since the ranks of $H_2(M;\mathbb{Z})$ and $H_2(M;\mathbb{Q})$ are 1 here.

STEP 3 A proof of the fact that in the case $k_1 > 2$ the rank of the integral homology group $H_2(W_f; \mathbb{Z})$ is 0.

Suppose that the rank of $H_2(W_f; \mathbb{Z})$ is not 0. By the structure of the map, there exists a smoothly embedded copy S_f^2 of the 2-dimensional unit sphere in M enjoying the following properties.

- $q_f(S_f^2) \cap \partial W_f$ is a finite set.
- The value of the homomorphism q_{f_*} at the 2nd integral homology class represented by the sphere is some element of $H_2(W_f; \mathbb{Z})$ whose order is infinite. This is seen as a homology class represented by a 2-dimensional sphere smoothly embedded in Int W_f . S_f^2 can be seen as a kind of variants of the images of sections of the bundle over the sphere in Int W_f whose projection is given by the restriction of q_f .

If $k_1 > 2$, then we can see that the rank of $H_2(M; \mathbb{Z})$ must be not 0. This is a contradiction.

STEP 4 A proof of the fact that in the case $k_1 = 2 S(f)$ is connected.

We use contradiction as in STEP 3. Suppose that S(f) is not connected in the case $k_1 = 2$. The rank of $H_2(W_f; \mathbb{Z})$ must be 0 due to a related argument in STEP 2 and by the assumption that the ranks of $H_2(M; \mathbb{Q})$ and $H_2(M; \mathbb{Z})$ must be at

8

most 1. This means that the two bundles in Proposition 1 (3) must be trivial. This is due to well-known classifications of bundles whose fibers are circles. A smooth bundle whose fiber is a circle is well-known to be regarded as a linear bundle and such bundles over a fixed space which is a CW complex (or a space of some suitable wider class of topological spaces) are classified by the 2nd integral cohomology classes of the base space. See also [18, 26] for example.

We can see that the so-called *self-intersection* of S(f) can be taken as the empty set. In other words, we can consider a smooth isotopy moving the original S(f)to a place apart from the original place by the structures of the bundles in Proposition 1 (3). STEP 2 implies that the square $u_0 \cup u_0$ of u_0 is the zero element of $H^4(M;\mathbb{Q})$ due to Poincaré duality for M. We can also see that even if we discuss the case where the coefficient ring is \mathbb{Z} , then the cup product is the zero element of $H^4(M;\mathbb{Z})$. The assumption that S(f) is not connected is shown to be false.

STEP 5 A proof of the fact that in the case $k_1 > 2$ the ranks of the integral homology groups $H_i(W_f; \mathbb{Z})$ and $H_{j-1}(W_f, \partial W_f; \mathbb{Z})$ are 0 for $2 \le j \le k_1 - 1$.

Suppose that the rank of the integral homology group $H_j(W_f; \mathbb{Z})$ is not 0 here. STEP 3 and the classification of smooth bundles whose fibers are circles, reviewed in STEP 4, imply that the bundles of Proposition 1 (3) are trivial. This implies that the rank of the integral homology group $H_j(M;\mathbb{Z})$ is not 0. This is a contradiction. Suppose that the rank of the integral homology group $H_{j-1}(W_f, \partial W_f;\mathbb{Z})$ is not 0 here. We consider the bundle whose projection is $\tilde{r}: \tilde{W} \to W_f$, discussed in STEP 1. For each element of $H_{j-1}(W_f, \partial W_f;\mathbb{Z})$, we can consider a kind of *prism operators* or *Thom classes* to obtain a unique element of $H_{(j-1)+(m-n+1)}(W;\mathbb{Z}) =$ $H_{m-n+j}(W;\mathbb{Z}) = H_{j+1}(W;\mathbb{Z})$ and consider the boundary to obtain a unique element of $H_j(M;\mathbb{Z})$ in the context of arguments on the homology groups. Note that n = m - 1 and that \tilde{W} is (PL) homeomorphic to and collapses to W here for example. This gives a monomorphism and implies that the rank of the integral homology group $H_j(M;\mathbb{Z})$ is not 0. This is also a contradiction.

STEP 6 A proof of the fact that for any integer $k_1 \ge 2$ the ranks of the integral homology groups $H_{k_1}(W_f; \mathbb{Z})$ and $H_{k_1-1}(W_f, \partial W_f; \mathbb{Z})$ are 1 and 0 respectively.

S(f) has been shown to be connected in STEPs 2 and 4. For example Proposition 1 (2) implies that ∂W_f is connected. We have a homology exact sequence for W_f as

$$H_1(W_f;\mathbb{Z}) \to H_1(W_f,\partial W_f;\mathbb{Z}) \to H_0(\partial W_f;\mathbb{Z}) \to H_0(W_f;\mathbb{Z})$$

and $H_1(W_f, \partial W_f; \mathbb{Z})$ is the trivial group by the facts that W_f is simply-connected and that ∂W_f is connected. This allows us to consider the case $k_1 > 2$ only in proving that the rank of $H_{k_1-1}(W_f, \partial W_f; \mathbb{Z})$ is 0.

Note that in STEP 6 (and throughout the present proof), the fact that M, W and W_f are simply-connected is important to apply Poincaré duality for any coefficient ring of the (co)homology groups.

Suppose that the rank of $H_{k_1-1}(W_f, \partial W_f; \mathbb{Z})$ is greater than 1. By the canonical way presented in STEP 5, we can obtain elements of $H_{k_1-1+m-n}(M; \mathbb{Z}) = H_{k_1}(M; \mathbb{Z})$. We can also see that the ranks of $H_{k_1}(M; \mathbb{Z})$ and $H_{k_1}(M; \mathbb{Q})$ are greater than 1. This is a contradiction. Suppose that the rank of $H_{k_1-1}(W_f, \partial W_f; \mathbb{Z})$ is 1. Similarly, we can see that the ranks of $H_{k_1}(M; \mathbb{Z})$ and $H_{k_1}(M; \mathbb{Q})$ are at least 1 and that the ranks are 1 by the assumption. Consider the Poincaré dual to the dual

to a generator of $H_{k_1-1}(W_f, \partial W_f; \mathbb{Q})$, whose rank is 1 of course and the resulting element is an element of $H_{m-k_1}(W_f; \mathbb{Q})$ and a generator of this vector space over \mathbb{Q} . We may regard that this is an element of a generator of $H_{m-k_1}(W_f; \mathbb{Z})$, whose rank is 1 of course. The bundles in Proposition 1 (3) are trivial by the condition $k_1 > 2$ and STEP 3. By considering a section of the trivial bundle, we have the Poincaré dual to the dual to a generator of $H_{k_1}(M; \mathbb{Q})$. This means that the square of a generator of $H^{k_1}(M; \mathbb{Q})$ or the cup product of two same element here is the zero element. This is a contradiction. The ranks of $H_{k_1-1}(W_f, \partial W_f; \mathbb{Z})$ and $H_{k_1-1}(W_f, \partial W_f; \mathbb{Q})$ are shown to be 0.

Suppose that the rank of $H_{k_1}(W_f; \mathbb{Z})$ is greater than 1. Arguments on sections or variants of bundles over spheres or more general subspaces of W_f in STEP 3 and STEP 5 imply that the rank of $H_{k_1}(M; \mathbb{Z})$ is greater than 1. This is a contradiction. Suppose that the rank of $H_{k_1}(W_f; \mathbb{Z})$ is 0. The rank of $H_{k_1}(W_f; \mathbb{Q})$ is 0 of course then. We have a homology exact sequence for W as

$$H_{k_1+1}(W;\mathbb{Q}) \to H_{k_1+1}(W,\partial W;\mathbb{Q}) \to H_{k_1}(M;\mathbb{Q}) \to H_{k_1}(W;\mathbb{Q})$$

and by Poincaré duality theorem for W, $H_{k_1+1}(W, \partial W; \mathbb{Q})$ and $H_{m-k_1}(W; \mathbb{Q})$ are isomorphic as vector spaces over \mathbb{Q} . Since W collapses to W_f , they are also isomorphic to $H_{m-k_1}(W_f; \mathbb{Q})$ and Poincaré duality theorem for W_f implies that they are also isomorphic to $H_{(m-1)-(m-k_1)}(W_f, \partial W_f; \mathbb{Q}) = H_{k_1-1}(W, \partial W_f; \mathbb{Q})$. The rank of $H_{k_1-1}(W_f, \partial W_f; \mathbb{Z})$ is shown to be 0 for $k_1 \geq 2$ and this implies that the rank of $H_{k_1}(M; \mathbb{Q})$ is 0 from the related argument in STEP 5. This is a contradiction. We have that the ranks of $H_{k_1}(W_f; \mathbb{Q})$ and $H_{k_1}(W_f; \mathbb{Z})$ are 1.

STEP 7 Investigating the (k_2-1) -th power $\bigcup_{j=1}^{k_2-1} u_0$ of a generator $u_0 \in H^{k_1}(M; \mathbb{Q})$ and showing that this is the zero element and that this is a contradiction.

The ranks of $H_{k_1}(W_f; \mathbb{Z})$ and $H_{k_1}(W_f; \mathbb{Q})$ have been shown to be 1. We consider a generator of a free subgroup of $H_{k_1}(W_f; \mathbb{Z})$ whose rank is 1, the dual to this, which we can define in a suitable way, and the Poincaré dual to the integral cohomology class, which is an element of the $(m - k_1 - 1)$ -th integral homology group $H_{m-k_1-1}(W_f; \mathbb{Z})$. We canonically have a unique element of $H_{m-k_1}(M; \mathbb{Z})$ by the method in STEP 5. We may regard that this is also the Poincaré dual PD $(u_0) \in H_{m-k_1}(M; \mathbb{Q})$ to u_0 .

We consider intersection theory and Poincaré duality for W_f and M. We can consider (variants of) generic intersections for copies of the homology class $PD(u_0)$. Here we may discuss where the coefficient ring is \mathbb{Z} . Remember that $m = k_1k_2$ and that the dimension of W_f is m - 1. We have the relation

$$(m-k_1-1) + (k_2-2)\{(m-k_1-1) - (m-1)\} = (m-k_1-1) - k_1(k_2-2) = k_1 - 1$$

and this relation enables us to regard that the Poincaré dual to the (k_2-1) -th power $\bigcup_{j=1}^{k_2-1} u_0$ of u_0 is represented by the element obtained by the method in STEP 5 from an element of $H_{k_1-1}(W_f, \partial W_f; \mathbb{Q})$, which is the trivial vector space. The (k_2-1) -th power is the zero element of $H^{k_1(k_2-1)}(M; \mathbb{Q})$. This is a contradiction.

This completes our proof.

A proof of Main Theorem 1. Theorem 4 and Main Theorem 2 completes the proof for the case $n \neq m$, For the case m = n, we can know from the theory of Eliashberg [3] and well-known facts on *Pontrjagin classes* of complex projective spaces, explained about in [18] for example.

Example 2. The k_2 -dimensional quaternionic projective space is for $k_1 = 4$ in Main Theorem 2. Its (*j*-th) integral homology group is always free for any integer *j*. [5] presents some meaningful examples for Main Theorem 2 more. See Theorem C there for example.

We close our paper by a remark related to a fact discussed by the author first in [17] and another several remarks.

Remark 2. We consider a situation in Remark 1.

Assume that a smooth immersion $f_0: W_{f_0} \to N$ is given similarly. According to [17], we have a special generic map $f_0: M_0 \to N$ on a suitable *m*-dimensional closed and connected manifold M_0 enjoying the following two properties.

- (1) The two bundles in Proposition 1 (3) are trivial bundles where the notation is as in Remark 1.
- (2) There exists a special generic map $f_k : M_0 \to N \times \mathbb{R}^k$ such that the composition of f_k with the canonical projection to N is the original map f_0 for any integer $1 \le k \le m n$.

More precisely, in Main Theorem 1 of [17], a special generic map $f: M \to N$ on an *m*-dimensional closed and connected manifold M enjoying some good topological properties is given and f_0 is given as a map on the manifold M_0 , obtained as a manifold whose cohomology ring is isomorphic to that of M.

It is an open problem whether M in Main Theorem 2 admits special generic maps into \mathbb{R}^n for $m - k_1 < n \leq m$ or n = m where we may regard the case n = m is another problem. Note also that it does not admit ones into \mathbb{R}^n for $1 \leq n \leq m - k_1$ by Theorem 4. We can also say that the special generic maps are not ones presented here even if partial answers to the open problem are given affirmatively.

We also note that real projective spaces admit no special generic maps in most cases ([13]). These cases are also suitable for exercises on Theorem 4.

Remark **3.** In the present paper, we may drop conditions on homology and homotopy groups for example. For example, we concentrate on studies on closed and simply-connected manifolds whose dimensions are at least 5, which are central objects in classical algebraic topology and differential topology. As pioneers, we and the author are trying to understand these manifolds geometric and constructive ways via special generic maps and more general good smooth maps such as so-called *fold* maps, generalizing the class of Morse functions and special generic maps. Related papers are [13, 14, 15, 16] for example. For our remark here, see also Remark 4 of [14] for example. For systematic singularity theory of so-called *generic* smooth maps including fold maps, we introduce [7] as a related book.

Remark 4. In Main Theorems, it is not meaningful to discuss the case where the Euler number of the manifold is odd. For example, the case where k is even in Main Theorem 1 and the case where k_2 is even in Main Theorem 2 are such cases. This is due to the existence theory of fold maps in [3, 4]

As another remark, in Main Theorem 2, k_1 must be even to make the problem meaningful.

4. Acknowledgments and declarations.

Naoki Kitazawa is a member of the project supported by JSPS KAKENHI Grant Number JP17H06128 "Innovative research of geometric topology and singularities of differentiable mappings" (Principal investigator: Osamu Saeki). Our present study is supported by the project.

We declare that data essentially supporting the study are all in the present paper.

References

- O. Burlet and G. de Rham, Sur certaines applications génériques d'une variété close a 3 dimensions dans le plan, Enseign. Math. 20 (1974). 275–292.
- [2] E. Calabi, Quasi-surjective mappings and a generation of Morse theory, Proc. U.S.-Japan Seminar in Differential Geometry, Kyoto, 1965, pp. 13–16.
- [3] Y. Eliashberg, On singularities of folding type, Math. USSR Izv. 4 (1970). 1119–1134.
- Y. Eliashberg, Surgery of singularities of smooth mappings, Math. USSR Izv. 6 (1972). 1302– 1326.
- J. Fowler and Z. Su, Smooth manifolds with prescribed rational cohomology ring, Geometriae Dedicata 182 (2016), 215–232, arXiv:1403.1801.
- [6] Y. K. S. Furuya and P. Porto, On special generic maps from a closed manifold into the plane, Topology Appl. 35 (1990), 41–52.
- [7] M. Golubitsky and V. Guillemin, Stable mappings and their singularities, Graduate Texts in Mathematics (14), Springer-Verlag (1974).
- [8] A. Hatcher, Algebraic Topology, A modern, geometrically flavored introduction to algebraic topology, Cambridge: Cambridge University Press (2002).
- [9] N. Kitazawa, On round fold maps (in Japanese), RIMS Kokyuroku Bessatsu B38 (2013), 45–59.
- [10] N. Kitazawa, On manifolds admitting fold maps with singular value sets of concentric spheres, Doctoral Dissertation, Tokyo Institute of Technology (2014).
- [11] N. Kitazawa, Fold maps with singular value sets of concentric spheres, Hokkaido Mathematical Journal Vol.43, No.3 (2014), 327–359.
- [12] N. Kitazawa Notes on explicit smooth maps on 7-dimensional manifolds into the 4dimensional Euclidean space, submitted to a refereed journal, arxiv:1911.11274.
- [13] N. Kitazawa, Closed manifolds admitting no special generic maps whose codimensions are negative and their cohomology rings, submitted to a refereed journal, arxiv:2008.04226v5.
- [14] N. Kitazawa, Restrictions on special generic maps on 6-dimensional or higher dimensional closed and simply-connected manifolds, submitted to a refereed journal, arxiv:2201.09437v4.
- [15] N. Kitazawa, Proofs of the non-existence of special generic maps on the 3-dimensional complex projective space, submitted to a refereed journal, arxiv:2202.00883.
- [16] N. Kitazawa, Characterizing certain classes of 6-dimensional closed and simply-connected manifolds via special generic maps, submitted to a refereed journal, arxiv:2205.04048.
- [17] N. Kitazawa, A note on cohomological structures of special generic maps, a revised version is submitted based on positive comments by referees and editors after the second submission to a refereed journal.
- [18] J. Milnor and J. Stasheff, *Characteristic classes*, Annals of Mathematics Studies, No. 76. Princeton, N. J; Princeton University Press (1974).
- [19] M. Nishioka, Special generic maps of 5-dimensional manifolds, Revue Roumaine de Math'ematiques Pures et Appliquèes, Volume LX No.4 (2015), 507–517.
- [20] O. Saeki, Topology of special generic maps of manifolds into Euclidean spaces, Topology Appl. 49 (1993), 265–293.
- [21] O. Saeki, Topology of special generic maps into ℝ³, Workshop on Real and Complex Singularities (Sao Carlos, 1992), Mat. Contemp. 5 (1993), 161–186.
- [22] O. Saeki and K. Sakuma, On special generic maps into ℝ³, Pacific J. Math. 184 (1998), 175–193.
- [23] O. Saeki and K. Sakuma, Special generic maps of 4-manifolds and compact complex analytic surfaces, Math. Ann. 313, 617–633, 1999.
- [24] O. Saeki and K. Suzuoka, Generic smooth maps with sphere fibers J. Math. Soc. Japan Volume 57, Number 3 (2005), 881–902.

- [25] K. Sakuma, On special generic maps of simply connected 2n-manifolds into R³, Topology Appl. 50 (1993), 249–261.
- [26] N. Steenrod, The topology of fibre bundles, Princeton University Press (1951).
- [27] D. J. Wrazidlo, Standard special generic maps of homotopy spheres into Eucidean spaces, Topology Appl. 234 (2018), 348–358, arxiv:1707.08646.
- [28] D. J. Wrazidlo (including a joint work with O. Saeki and K. Sakuma), The Milnor 7-sphere does not admit a special generic map into \mathbb{R}^3 , http://math.ac.vn/conference/FJV2018/images/slides/Wardzilo_FJV2018.pdf, 2018.
- [29] D. J. Wrazidlo, On special generic maps of rational homology spheres into Euclidean spaces, arxiv:2009.05928.

INSTITUTE OF MATHEMATICS FOR INDUSTRY, KYUSHU UNIVERSITY, 744 MOTOOKA, NISHI-KU FUKUOKA 819-0395, JAPAN, TEL (OFFICE): +81-92-802-4402, FAX (OFFICE): +81-92-802-4405,

Email address: n-kitazawa@imi.kyushu-u.ac.jp

 $Webpage:\ https://naokikitazawa.github.io/NaokiKitazawa.html$