

Stochastic Langevin Differential Inclusions with Applications to Machine Learning

Fabio V. Difonzo^{1*}, Vyacheslav Kungurtsev² and Jakub Mareček²

¹Department of Mathematics, University of Bari Aldo Moro, Via E. Orabona 4, Bari, 70125, Italy.

²Department of Computer Science, Faculty of Electrical Engineering, Czech Technical University in Prague, Karlovo namesti 13, Prague, 11000, hlavní mesto Praha, Czech Republic.

*Corresponding author(s). E-mail(s): fabio.difonzo@uniba.it;
Contributing authors: vyacheslav.kungurtsev@fel.cvut.cz;
jakub.marecek@fel.cvut.cz;

Abstract

Stochastic differential equations of Langevin-diffusion form have received significant recent, thanks to their foundational role in both Bayesian sampling algorithms and optimization in machine learning. In the latter, they serve as a conceptual model of the stochastic gradient flow in training over-parametrized models. However, the literature typically assumes smoothness of the potential, whose gradient is the drift term. Nevertheless, there are many problems, for which the potential function is not continuously differentiable, and hence the drift is not Lipschitz-continuous everywhere. This is exemplified by robust losses and Rectified Linear Units in regression problems. In this paper, we show some foundational results regarding the flow and asymptotic properties of Langevin-type Stochastic Differential Inclusions under assumptions appropriate to the machine-learning settings. In particular, we show strong existence of the solution, as well as asymptotic minimization of the canonical Free Energy Functional.

Keywords: Sampling, Langevin diffusion, Differential Inclusions, Robust Regression, Fokker-Planck Equation

1 Introduction

In this paper, we study the following stochastic differential inclusion,

$$dX_t \in -F(X_t)dt + \sqrt{2\sigma} dB_t \quad (1)$$

wherein $F(x) : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is a set-valued map. We are particularly interested in the case where $F(x)$ is the Clarke subdifferential of some continuous tame function $f(x)$. This is motivated by the recent interest in studying Langevin-type diffusions in the context of machine learning applications, both as a scheme for sampling in a Bayesian framework (as spurred by the seminal work [48]) and as a model of the trajectory of stochastic gradient descent, with a view towards understanding the asymptotic properties of training deep neural networks [20]. It is typically assumed that $F(x)$ above is Lipschitz, and as such its potential $f(x)$ is continuously differentiable. In many problems of relevance, including empirical risk minimization with a robust (e.g., l_1 or Huber) loss and neural networks with ReLU activations, this is not the case, and yet there is at least partial empirical evidence suggesting the long-run behavior of a numerically similar operation is similar in its capacity to generate a stochastic process which minimizes a Free Energy associated with the learning problem.

This paper is organized as follows. In Section 2 we study the functional analytical properties of $F(x)$ as it appears in (1) when it represents a noisy estimate of a subgradient element of an empirical loss function that itself satisfies the conditions of a *definable* potential, especially as it appears in the context of deep learning applications. Note that the research program undertaken relates to a recent conjecture of Bolte and Pauwels [9, Remark 12] which suggests the strong convergence of iterates in a stochastic subgradient type sequence to stationary points for this class of potentials. Subsequently, in Section 3 we prove that there exists a strong solution to (1), confirming the existence of a trajectory in the general case. In this sense, we extend the work of [30, 29] studying diffusions with discontinuous drift to set-valued drift. Next in Section 4 we prove the correspondence of a Fokker-Planck type equation to modeling the probability law associated with this stochastic process, and show that it asymptotically minimizes a Free Energy functional corresponding to the loss function of interest, extending the seminal work of [22] which had proven the same result in the case of continuous $F(x)$. We present some numerical results confirming the expected asymptotic behavior of (1) in Section 5 and summarize our findings and their implications in Section 6.

1.1 Related Work

Stochastic differential inclusions are the topic of the monograph [25], which presents the associated background of stochastic differential equations (SDEs) as well as set-valued analysis and differential inclusions, providing a notion of a weak and strong solution to equations of the form (1), and ones with more general expressions, especially in regards to the noise term. In this work,

and others studying stochastic differential inclusions in the literature, such as, e.g., [24, 26], it is assumed that the set-valued drift term $F(x)$ is Lipschitz continuous. In this paper we are interested in the more general case, where $F(x)$ may not be Lipschitz.

Langevin diffusions have had three distinct significant periods of development. To begin with, the original elegant correspondence of SDEs with semi-groups associated with parabolic differential operators was explored in depth by [42] (first edition published in the 1970's), following the seminal paper [21]. See also [23].

Later, they served as a canonical continuous Markovian process with the height of activity on ergodicity theory, the long run behavior of stochastic processes, associated with the famous monograph [35], first appearing in the 1990's. See, e.g. [34].

Finally, most recently, the Langevin diffusion has been a model to study the distributional dynamics of noisy training of contemporary machine learning models [20]. At this point, we have the closest works to this one. In Deep Neural Networks (DNNs), Rectified Linear Units (ReLUs), which involve a component-wise maximum of zero and a linear expression, are standard functional components in predictors appearing in both regression and classification models. Mean-field analyses seek to explain the uncanny generalization abilities of DNNs by considering the distributional dynamics of idealized networks with infinitely many hidden layer neurons by considering both the stochastic equations corresponding to said dynamics, as well as Fokker-Planck-type PDEs modeling the flow of the distribution of the weights in the network. Analyses along these lines in the contemporary literature include [31, 39]. Although this line of works does use limiting arguments involving stochastic and distributional dynamics, they do not directly consider the Langevin differential inclusion and the potential PDE solution for the distribution, as we do here.

2 Tame functions, o-minimal structures and the exceptional set

Deep learning raises a number of important questions on the interface of optimization theory and stochastic analysis [10]. In particular, there are still major gaps in our understanding of the applications of plain stochastic gradient descent (SGD) in deep learning, leaving aside the numerous related recent optimization algorithms for deep learning [38]. A particularly challenging aspect of deep learning is the composition of functions that define the objective landscape. These functions are typically recursively evaluated piecewise non-linear maps. The nonlinearities are due to the sigmoidal function, exponentiation and related operations, which are neither semialgebraic nor semianalytic, and the piece-wise nature of the landscape is due to the common appearance of component-wise max.

We consider Euclidean space \mathbb{R}^n with the canonical Euclidean scalar product $\langle \cdot, \cdot \rangle$ and a locally Lipschitz continuous function $f: \mathbb{R}^n \rightarrow \mathbb{R}$. For any

$x \in \mathbb{R}^n$, the Clarke subgradient of f [12]:

$$\partial^c f(x) = \text{conv} \left\{ v \in \mathbb{R}^n : \exists y_k \xrightarrow[k \rightarrow \infty]{} x \text{ with } y_k \in \mathbb{R}, v_k = \nabla f(y_k) \xrightarrow[k \rightarrow \infty]{} v \right\}.$$

Following a long history of work [33, e.g.], we utilize a lesser known function class of definable functions, which are known [44] to include restricted analytic fields with exponentiation. We refer to Macintyre, McKenna, and van den Dries [32] and Knight, Pillay, and Steinhorn [27] for the original definitions, and to van den Dries-Miller [45, 43] and Coste [13] for excellent book-length surveys. In particular, we use the following:

Definition 2.1 (Structure, cf. Coste [13], Knight et al. [27]) A *structure* on $(\mathbb{R}, +, \cdot)$ is a collection of sets $\mathcal{O} = (\mathcal{O}_p)_{p \in \mathbb{N}}$, where each \mathcal{O}_p is a family of subsets of \mathbb{R}^n such that for each $p \in \mathbb{N}$:

1. \mathcal{O}_p contains the set $\{x \in \mathbb{R}^n : g(x) = 0\}$, where g is a polynomial on \mathbb{R}^n , i.e., $g \in \mathbb{R}[X_1, X_2, \dots, X_p]$, and the set is often known as the family of real-algebraic subsets of \mathbb{R}^n ;
2. if any A belongs to \mathcal{O}_p , then $A \times \mathbb{R}$ and $\mathbb{R} \times A$ belong to \mathcal{O}_{p+1} ;
3. if any A belongs to \mathcal{O}_{p+1} , then $\pi(A) \in \mathcal{O}_p$, where $\pi: \mathbb{R}^{n+1} \times \mathbb{R} \rightarrow \mathbb{R}^n$ is the coordinate or “canonical” projection onto \mathbb{R}^n , i.e., the projection to the first p coordinates;
4. \mathcal{O}_p is stable by complementation, finite union, finite intersection and contains \mathbb{R}^n , which defines a Boolean algebra of subsets of \mathbb{R}^n .

Definition 2.2 (Definable functions, cf. Coste [13] and van den Dries-Miller[45]) A *structure* on $(\mathbb{R}, +, \cdot)$ is called *o-minimal* when the elements of \mathcal{O}_1 are exactly the finite unions of (possibly infinite) intervals and points. The sets $A \subseteq \mathbb{R}^n$ belonging to an o-minimal structure \mathcal{O}_p , for some $p \in \mathbb{N}$, are called *definable in the o-minimal structure* \mathcal{O} . One often shortens this to *definable* if the structure \mathcal{O} is clear from the context. A set-valued mapping is said to be definable in \mathcal{O} , whenever its graph is definable in \mathcal{O} .

A subset of \mathbb{R}^n is called *tame* if there exists an o-minimal structure such that the subset is definable in the o-minimal structure. Notice that this notion of tame geometry goes back to Grothendieck’s *topologie modérée* [18].

Next, we consider two more regularity properties. First, we consider functions that have conservative set-valued fields of [9], or equivalently, are path differentiable [9]. This includes convex, concave, Clarke regular, and Whitney stratifiable functions.

Definition 2.3 (Conservative set valued fields of [9]) Let $D: \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ be a set-valued map. D is a *conservative (set-valued) field* whenever it has closed graph,

nonempty compact values and for any absolutely continuous loop $\gamma: [0, 1] \rightarrow \mathbb{R}^n$, that is $\gamma(0) = \gamma(1)$, we have

$$\int_0^1 \max_{v \in D(\gamma(t))} \langle \dot{\gamma}(t), v \rangle dt = 0$$

in the Lebesgue sense.

A function f defined through any of the equivalent forms

$$f(x) = f(0) + \int_0^1 \max_{v \in D(\gamma(t))} \langle \dot{\gamma}(t), v \rangle dt \quad (2)$$

$$= f(0) + \int_0^1 \min_{v \in D(\gamma(t))} \langle \dot{\gamma}(t), v \rangle dt \quad (3)$$

$$= f(0) + \int_0^1 \langle \dot{\gamma}(t), D(\gamma(t)) \rangle dt \quad (4)$$

for any γ absolutely continuous with $\gamma(0) = 0$ and $\gamma(1) = x$, is called a *potential function for D* . Let us note that f is well and uniquely defined up to a constant.

We shall also say that D admits f as a potential or that D is a conservative field for f .

The second notion of regularity, which we consider, is the notion of piecewise Lipschitzianity on \mathbb{R}^n from [30]. The associated exception(al) set [29] is the subset of that function's domain where the function is not Lipschitz. We recall the definition here for convenience, and we state it for set-valued maps.

Definition 2.4 A function $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ is *piecewise Lipschitz continuous* if there exists a hypersurface Θ , which we call the *exceptional set* for f , such that Θ has finitely many connected components θ_i , $i = 1, 2, \dots$, such that $f|_{\mathbb{R}^n \setminus \Theta}$ is intrinsic Lipschitz (cf. [30, Definition 3.2]).

A set-valued function $F: \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is *piecewise Lipschitz continuous* if there exists an exceptional set Θ , defined analogously as for single-valued functions, such that F is Lipschitz on $\mathbb{R}^n \setminus \Theta$ with respect to Hausdorff H -metric (cf. [4]).

Let us recall that, given a set-valued map with compact convex values $F: \mathbb{R}^n \rightrightarrows \mathbb{R}^n$, we say that F is H -Lipschitz, or just Lipschitz, if and only if there exists a constant $l > 0$ such that for any $x, y \in \mathbb{R}^n$ we have

$$\rho_H(F(x), F(y)) \leq l \|x - y\|,$$

where, for arbitrary compact sets $A, B \subseteq \mathbb{R}^n$,

$$\rho_H(A, B) := \max\left\{\max_{a \in A} \text{dist}(a, B), \max_{b \in B} \text{dist}(b, A)\right\},$$

with $\text{dist}(a, B) := \min_{b \in B} \|a - b\|_2$.

Definition 2.5 ([45, 8]) A C^r stratification of a closed (sub)manifold M of \mathbb{R}^n is a locally finite partition $(M_i)_{i \in I}$ of M into C^r submanifolds (called *strata*) having the property that for $i \neq j$, $\text{cl}(M_i) \cap M_j \neq \emptyset$ implies that M_j is entirely contained in $\text{cl}(M_i) \setminus M_i$ (called the *frontier* of M_i) and $\dim(M_j) < \dim(M_i)$.

Moreover, a C^r stratification $M = (M_i)_{i \in I}$ of M has the Whitney-(a) property if, for each $x \in M_i \cap M_j$ (with $i \neq j$) and for each sequence $x_k \subseteq M_i$ we have

$$\begin{cases} \lim_{k \rightarrow \infty} x_k = x, \\ \lim_{k \rightarrow \infty} T_{x_k} M_i = \mathcal{T}, \end{cases} \implies T_x M_j \subseteq \mathcal{T},$$

where $T_x M_j$ (respectively, $T_{x_k} M_i$) denotes the tangent space of the manifold M_j at x (respectively, of M_i at x_k).

If, moreover, a Whitney stratification $(M_i)_{i \in I}$ satisfies for all $i \in I$ and $x \in M_i$ the transversality condition

$$e_{n+1} \notin T_x M_i, \quad (5)$$

where $e_{n+1} = [0 \cdots 0 \ 1] \in \mathbb{R}^{n+1}$, then it is called a *nonvertical* Whitney stratification.

A function $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is said to be *stratifiable*, in any of its connotations, if the graph of f , denoted by $\text{gr}f$, admits a corresponding C^r stratification.

Remark 2.1 Let us note that, from Definition 2.5, if $(M_i)_{i \in I}$ is a stratification of a stratifiable submanifold M , then it follows that $M = \bigcup_{i \in I} M_i$. Therefore, by the Hausdorff maximality principle, there must exist $\bar{i} \in I$ such that $\dim M_{\bar{i}} = \max_{i \in I} \dim M_i$ and, thus, $M = M_{\bar{i}}$ since all the other subspaces M_i , $i \neq \bar{i}$ have zero Lebesgue measure. The same argument holds, a fortiori, for every finite subset $J \subseteq I$.

In this paper we establish the existence guarantees of strong solutions to the equation (1) and study the PDE describing the flow of the probability mass of $X(t)$. To set up the subsequent exposition, we must first establish the groundwork of linking Whitney stratifiable potential functions, which describe the loss landscape of neural networks and other statistical models on the one hand, to set valued piecewise Lipschitz continuous maps, which describe the distributional flow of stochastic subgradient descent training, modeled by (1), on the other.

Theorem 2.1 *Let $F : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ be a definable conservative field admitting a tame potential $f : \mathbb{R}^n \rightarrow \mathbb{R}$. Let F be Lipschitz on $\mathbb{R}^n \setminus B_\delta(0)$ with constant L , for some $L, \delta > 0$. Then F is piecewise Lipschitz continuous.*

Proof As f is tame, by definition, it is also definable. Since F is conservative, from [9] f is locally Lipschitz and, since it is tame, [7, Theorem 1] implies that f is semismooth. Therefore, by [8, Corollary 9], letting $r \geq 2$ be an arbitrarily fixed integer, f admits a nonvertical C^r Whitney stratification $(M_i)_{i \in I}$. With abuse of notation, let $(M_i)_{i \in I}$ denote the stratification relative to the domain of f . So, on the account of local finiteness, there must exist a maximal finite subset of indices $J \subseteq I$ such that

$$B_\delta(0) \cap M_j \neq \emptyset, \quad j \in J.$$

Since f is semismooth we deduce that its directional derivative $f'(\cdot; v)$ is C^{r-1} for all $v \in \mathbb{N}$ and, thus, is Lipschitz on each $B_\delta(0) \cap M_j$, $j \in J$; hence, the Riemannian gradient $\nabla_R f(x)$ is Lipschitz on $B_\delta(0) \cap M_j$, $x \in M_j$, $j \in J$, since $\nabla_R f(x)$ is the restriction of the directional derivative to tangent directions onto M_j [8]. Let us note that the Clarke subgradient of f at x , denoted by $\partial^\circ f(x)$, is such that (see [8, Proposition 4])

$$\text{Proj}_{T_x M_j} \partial^\circ f(x) \subseteq \{\nabla_R f(x)\},$$

where M_j is the stratum such that $x \in M_j$, $j \in J$.

Now, on the account of Remark 2.1, for some $\bar{j}_x \in J$, we have $\dim M_{\bar{j}_x} = \dim B_\delta(0) = n$. By compactness, there exists a finite covering of $B_\delta(0)$ of maximal dimension, denoted by $(M_{\bar{j}})_{\bar{j} \in \bar{J}}$, such that $B_\delta(0) \subseteq \bar{M}$, where $\bar{M} := \bigcup_{\bar{j} \in \bar{J}} M_{\bar{j}}$. Therefore $\nabla_R f(x) = \nabla f(x)$ on $B_\delta(0)$.

Then, as a consequence of [9, Theorem 1], there exists a zero-measure set $S \subseteq \mathbb{R}^n$ such that $F(x) = \{\nabla f(x)\}$ for all $x \in B_\delta(0) \setminus S$. Now, for $x \in B_\delta(0) \setminus S$, from [9, Corollary 1], we have that

$$\partial^\circ f(x) \subseteq \text{conv} F(x) = \text{conv}\{\nabla f(x)\} = \{\nabla f(x)\},$$

so that

$$\text{Proj}_{T_x M_j} \partial^\circ f(x) \subseteq \text{Proj}_{T_x M_j} \{\nabla f(x)\} = \{\nabla f(x)\}.$$

It then follows that $\nabla f(x)$, and so F , is Lipschitz on $B_\delta(0) \setminus S$. Since $B_\delta(0)$ is compact, there exists a finite family $\{\theta_k\}_{k \in K}$ such that $B_\delta(0) \cap S \subseteq \bigcup_{k \in K} \theta_k$. Thus F is Lipschitz on $B_\delta(0) \setminus \bigcup_{k \in K} \theta_k$, and the claim is proved. \square

3 Existence and Uniqueness of Solution to the SDI

In this section we will prove that (1) admits a strong solution. More precisely, it will be proven that there exists a suitable selection of $F(X_t)$ such that the corresponding SDE has a strong solution: this will in turn imply that the original stochastic differential inclusion has a solution as well. Our result will rely on an existence and uniqueness pertaining to SDEs with discontinuous drift in [30].

3.1 Piecewise Lipschitz selections of upper semicontinuous set-valued maps

In our setting, assuming that F is the Clarke subdifferential of some continuous tame function guarantees that F is an upper semi-continuous set-valued map. We further assume that F is bounded with compact convex values. Our aim is to prove that, under these assumptions, F has a piecewise Lipschitz selection.

We are going to need the following result from [2, Theorem 9.4.3].

Theorem 3.1 *Consider a Lipschitz set-valued map F from a metric space to nonempty closed convex subsets of \mathbb{R}^n . Then F has a Lipschitz selection, called Steiner selection.*

Moreover, we recall the classical Kirszbraun's Theorem (see [16]).

Theorem 3.2 *If $S \subseteq \mathbb{R}^n$ and $f : S \rightarrow \mathbb{R}^n$ is Lipschitz, then f has a Lipschitz extension $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$ with the same Lipschitz constant.*

In the next Theorem, which is the main result of this section, we prove that, under some mild assumptions, the set-valued map F in (1) has a piecewise Lipschitz selection for any suitable compact covering of \mathbb{R}^n .

Theorem 3.3 *Let $F : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ be an upper semi-continuous set-valued map with closed convex values, and piecewise Lipschitz continuous, with exceptional set Θ . Let us assume that there exists $b > 0$ such that $F(x) \subseteq bB_n$ for all $x \in \mathbb{R}^n$. Then F has a piecewise Lipschitz selection with exceptional set Θ , arbitrarily smooth on the interior of each connected component.*

Proof Let $\{R_i\}_{i=1}^r$ be the finite family of closed subsets of \mathbb{R}^n such that

$$\bigcup_{i=1}^r R_i = \mathbb{R}^n \setminus \Theta.$$

For each $i = 1, \dots, r$, Theorem 3.1 on $F|_{R_i}$ implies that there exists a finite sequence of equi-Lipschitz, and thus continuous, selection functions $\{f_i\}$ where each f_i is defined on R_i , that is $f_i : R_i \rightarrow \mathbb{R}^n$.

Let us note that, since $F(x) \subseteq bB_n$ for all $x \in \mathbb{R}^n$ then $\{f_i\}_{i=1}^r$ is uniformly bounded. We can now extend each f_i on the whole \mathbb{R}^n , to some function, still denoted by f_i , which is Lipschitz with the same constant as the original function's on the account of Kirszbraun's Theorem 3.2; moreover, the family $\{f_i\}$ can be assumed to retain uniform boundedness.

Let now $\varepsilon > 0$ be given and let $i = 1, \dots, r$ be fixed. Let us consider a partition of unity $\{\varphi^\varepsilon\}$ as in [40, Lemma 1.2]. Now, following a classical construction (e.g., see [3]), we let

$$g_i^\varepsilon(x) := \int_{\mathbb{R}^n} f_i(x) \varphi^\varepsilon(y - x) dy.$$

It then follows that $g_i^\varepsilon \in \mathcal{C}^\infty(\mathbb{R}^n)$ is a Lipschitz function, with the same Lipschitz constant as f_i 's, and is such that $g_i^\varepsilon \in F(x)$ by straightforward computations. Let us stress that the Lipschitz constant of each g_i^ε is independent of ε , so that $\{g_i^\varepsilon\}_{\varepsilon > 0}$ is equi-Lipschitz on R_i .

Let now $x \in \mathbb{R}^n$ and $i_x \in \mathbb{N}$ be unique index such that $x \in R_{i_x}$. We then define

$$f(x) := g_{i_x}^\varepsilon(x), \quad x \in \mathbb{R}^n.$$

It then follows that $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is countably piecewise Lipschitz on \mathbb{R}^n according to Definition 2.4 and its exceptional set is Θ . Obviously $f(x) \in F(x)$, and this proves the claim. \square

Remark 3.1 Theorem 3.3 still holds if we replace \mathbb{R}^n with any $A \subseteq \mathbb{R}^n$: in fact, we can always extend F to an upper semi-continuous set-valued map defined on the whole \mathbb{R}^n by virtue of [41, Theorem 2.6].

Finally, we have the following.

Corollary 3.1 *Let $F : \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ be an upper semi-continuous set-valued map with closed convex values, and piecewise Lipschitz continuous with a \mathcal{C}^3 exceptional set Θ . Assume that there exists $b > 0$ such that $F(x) \subseteq bB_n$ for all $x \in \mathbb{R}^n$. Then there exists a unique strong solution for the SDI (1).*

Proof From Theorem 3.3 there exists a drift $\mu : \mathbb{R}^n \rightarrow \mathbb{R}^n$, piecewise Lipschitz selection of the set-valued map F , which satisfies Assumptions 3.4 – 3.6 from [30]. Therefore, on the account of [30, Theorem 3.21], we obtain that the SDE

$$dX_t = -\mu(X_t)dt + \sqrt{2\sigma} dB_t \quad (6)$$

has a unique global strong solution, which in turn represents a strong solution to the stochastic differential inclusion (1), and this proves the claim. \square

Remark 3.2 Let us stress that the Lipschitz selection of the set-valued map F may not be Lipschitz on $\Theta_s \setminus \Theta$. This does not affect the result of Corollary 3.1 since, in order for it to hold, it suffices to provide a strong solution to (6).

4 Fokker-Planck Equation and Free Energy Minimization

4.1 Fokker-Planck Equation

The Fokker-Planck equation describes the evolution of the probability density associated with the random process modeled by the diffusion flow. Classical results deriving the Fokker-Planck equation from SDEs can be reviewed in [15, 21, 23, 42]. In particular, for a drift diffusion of the form $dX_t = -\nabla f(X_t) dt + \sqrt{2\sigma} dB_t$ it holds that, from any initial distribution $\rho(0) = \rho_0$, the density $\rho(x, t)$ of X_t is given by

$$\frac{\partial \rho}{\partial t} = -\operatorname{div}(\nabla f(x)\rho) + \frac{1}{2}\Delta(\sigma^2\rho) \quad (7)$$

and has a limiting stationary distribution governed by the Gibbs form $\exp\{-f(x)\}/Z$. In [22] it was shown that this corresponds to the gradient flow of a variational problem of minimizing the free energy.

These classical results, as already can be gleaned from the statements of the form of the PDE, require smoothness of F . The equation can be derived from applying integration by parts after Itô's Lemma on the diffusion process. Following [17, 4.3.5], for arbitrary g , we have,

$$\begin{aligned} \frac{\langle g(x(t)) \rangle}{dt} &= \frac{d}{dt} \langle g(X(t)) \rangle \\ &= \langle \nabla f(x) \partial_x f + \frac{1}{2} \sigma \partial_{xx}^2 g \rangle \end{aligned}$$

$$\begin{aligned}
&= \int dx \left[\nabla f(x) \partial_x g + \frac{1}{2} \sigma \partial_{xx}^2 g \right] \rho(x, t | x_0, t_0) \\
&= \int dx g(x) \partial \rho(x, t | x_0, t_0)
\end{aligned}$$

Replacing the expression $\nabla f(x)$ with an arbitrary coefficient function of form $a(x)$, we can see that in the case that $a(x)$ is a selection of the Clarke subdifferential, and thus not a continuous function of x , and as such even the weak distributional sense derivative of $a(x)$ does not exist and integration by parts cannot be applied.

4.2 Generic Solution Existence Results

Nevertheless, a solution $\rho(x, t)$ can be shown to exist for the system as stated in weak form. To this end, we follow [6]. Let $\Omega_T \subseteq \mathbb{R}^n \times [0, T]$ open and

$$L_{A,b}\varphi = \sum_{i=1}^n a_i \frac{\partial \varphi}{\partial x_i} + \sum_{i,j=1}^n \sigma_{ij} \frac{\partial \varphi}{\partial x_i \partial x_j}, \quad \varphi \in \mathcal{C}_0^\infty(\Omega).$$

Moreover, let n' be such that $\frac{1}{n} + \frac{1}{n'} = 1$. It holds the following.

Theorem 4.1 (Corollary 3.2,[6]) *Let μ be a locally finite Borel measure on Ω such that $a_i, \sigma_{ij} \in L_{loc}^1(\Omega_T, \mu)$ and,*

$$\int_{\Omega_T} \left[\frac{\partial \phi}{\partial t} + \sum_{i=1}^n a_i \frac{\partial \phi}{\partial x_i} + \sum_{i,j=1}^n \sigma_{ij} \frac{\partial \phi}{\partial x_i \partial x_j} \right] d\mu = 0$$

for all nonnegative $\phi \in \mathcal{C}_0^\infty(\Omega_T)$. Furthermore let σ_{ij} be uniformly bounded, nondegenerate and Hölder continuous, then $\mu = \rho(x, t) dx dt$ with $\rho \in L_{loc}^p(\Omega_T)$ for every $p \in [1, (n+2)')$.

Remark 4.1 As observed in [6], one cannot expect that the density of μ is continuous even for infinitely differentiable σ_{ij} under these conditions. However, we note that in [37] a continuous solution is shown under the assumption that $a_i \in L^p(\Omega_T, \mu)$, i.e. it is globally integrable. Again, however, this is very restrictive in the case of studying the evolution of diffusion operators on tame nonsmooth potentials of interest.

Existence of an invariant measure μ for the probability flow, i.e., a solution for the purely elliptic part, is guaranteed by [5, Theorem 1.6] and [1, Theorem 1.2].

4.3 Fokker-Planck With Boundary Conditions

Note, however, that we have an expectation as to what the stationary distribution for (1), namely, of Gibbs form proportional to $\exp\{-f(x)\}$. This measure

is even absolutely continuous, and thus has higher regularity than Theorem 4.1 suggests.

To this end, consider [17, Chapter 5.1.1] which considers boundary conditions at a discontinuity for the FP equation. We can consider that the space is partitioned into connected components R_i , and in each region, the continuous SDE and associated FP (7) holds. However, for boundaries between regions $S_{ij} \subseteq \Theta$ we have,

$$n \cdot J_i(x, t)|_{S_{ij}^i} = n \cdot J_j(x, t)|_{S_{ij}^j}, \quad \rho(x, t)|_{S_{ij}^i} = \rho(x, t)|_{S_{ij}^j} \quad (8)$$

where the probability current J is defined as,

$$J_i(x, t) = F_i(x)\rho(x, t) + \frac{1}{2}\sigma \frac{\partial \rho(x, t)}{\partial x_i}$$

as defined on region R_i

Note that in one dimension this is simple: we have, at a point of non-differentiability x_c ,

$$\begin{aligned} \rho(x, t)|_{x_c^+} &= \rho(x, t)|_{x_c^-}, \\ \left(F(x)\rho(x, t) + \frac{1}{2}\sigma \frac{\partial \rho(x, t)}{\partial x_i} \right) \Big|_{x_c^+} &= - \left(F(x)\rho(x, t) + \frac{1}{2}\sigma \frac{\partial \rho(x, t)}{\partial x_i} \right) \Big|_{x_c^-} \end{aligned}$$

where we abuse notation to indicate the partial directional derivative of ρ from either side of x_c .

We can consider writing a stationary solution $\pi(x)$ to this system as we have a suspected ansatz, similarly as done in [17, Section 5.3]. Generically stationary implies $\nabla \cdot J(x) = 0$, or that J is divergence-free with respect to x . We have,

$$\sum_{i=1}^n \frac{\partial}{\partial x_i} \left[F_i(x)\rho(x, t) + \frac{1}{2}\sigma \frac{\partial \rho(x, t)}{\partial x_i} \right] = 0 \quad (9)$$

Indeed, let

$$\pi(x) = \exp\{-f(x)/\sigma\} / Z.$$

On the domain $\mathbb{R}^n \setminus \Theta$ where $\nabla F(x)$ is well-defined, this can be seen immediately to solve (9). Since Θ is of measure zero with respect to the ambient space, we can define,

$$Z = \int_{\mathbb{R}^n} \pi(x) dx = \int_{\cup_i R_i} \pi(x) dx$$

Of course, a constructive ansatz for the stationary solution, the elliptic form, neither shows its uniqueness, nor the existence, uniqueness and regularity of the probability mass flow $\rho(x, t)$. To the best of our knowledge, no such result exists for the network of parabolic first-order systems under consideration, and

at the same time, we shall see that the continuity of $\rho(x, t)$ is important in the next section for the variational conception of the FP equation as a minimizing flow for the free energy.

To this end, we can consider two lines of work as a foundation to formulate the requisite results. In particular [36] shows regularity conditions of solutions to second-order parabolic equations defined on a Lipschitz domain. As seen in Section 2, for the problems of interest, the exceptional set can be parameterized in a smooth way, which implies that for compact sets, the boundary is Lipschitz. We must, however, take care to translate the results appropriately to the potentially unbounded domains a potential R_i could correspond to.

The closest to our line of work is considering graph-structured networks of PDEs, such as modeling Kirchhoff's laws. A prominent and representative work along these lines is [47]. In this setting, there is a network of one-dimensional paths embedded in some ambient space in \mathbb{R}^n , with the paths connected at a series of vertices, with boundary conditions connecting a collection of linear parabolic PDEs governing the flow of a quantity across the network. Thus, the spirit of a network structure of PDEs with connecting boundary connections is analogous to our problem, however, with the caveat that they are one-dimensional embedded domains, even if embedded in a larger space. Existence and smoothness regularity conditions are shown.

We consider an approach using domain decomposition methods for the solutions of PDEs based on optimal control theory [19] (see also, e.g., [14]) and consider reformulating the boundary conditions as a control to establish a variational formulation.

First let $\delta > 0$ be a regularization parameter and consider the optimization problem

$$\begin{aligned} \min_{\{\rho_i \in H^1(R_i)\}\{g_{ij} \in L^2(S_{ij})\}} \mathcal{J}_\delta(\{\rho_i\}, \{g_{ij}\}) \\ \text{subject to } \frac{\partial \rho_i}{\partial t} + F(x) \cdot \nabla \rho_i + \sigma \nabla \cdot \nabla \rho_i = 0, \text{ on } R_i, \forall i, \quad (10) \\ n \cdot J_i = g_{ij}, \text{ on } S_{ij}, \\ n \cdot J_j = -g_{ij}, \text{ on } S_{ij}, \end{aligned}$$

where

$$\mathcal{J}_\delta(\{\rho_i\}, \{g_{ij}\}) := \sum_{ij} \left[\int_{S_{ij}} (\rho_i - \rho_j)^2 dS_{ij} + \frac{\delta}{2} \int_{S_{ij}} g_{ij}^2 dS_{ij} \right].$$

Let us also note the weak form of the PDE constraints,

$$\begin{aligned} \int_{R_i} [\partial_t \rho_i v + \sigma \nabla \rho_i \cdot \nabla v + F \cdot \nabla \rho v] dx = \\ \sum_j \int_{S_{ij}} g_{ij} dS_{ij} - \sum_j \int_{S_{ji}} g_{ji} dS_{ji}, \forall v \in H^1(R_i). \quad (11) \end{aligned}$$

We have the following, akin to [19, Theorem 2.1]. Let

$$\mathcal{U} = \{ \{ \rho_i \in H^1(R_i) \} \{ g_{ij} \in L^2(S_{ij}) \} \text{ satisfying (11), } \mathcal{J}_\delta(\{ \rho_i \}, \{ g_{ij} \}) < \infty \}$$

Theorem 4.2 *There exists a unique solution $(\{ \rho_i \in H^1(R_i) \} \{ g_{ij} \in L^2(S_{ij}) \})$ to (10) in \mathcal{U} .*

Proof Let $\{ \rho_i^{(n)}, g_{ij}^{(n)} \}$ be a minimizing sequence in \mathcal{U} , i.e.,

$$\lim_{n \rightarrow \infty} \mathcal{J}_\delta(\{ \rho_i^{(n)}, g_{ij}^{(n)} \}) = \inf_{\{ \rho_i, g_{ij} \} \in \mathcal{U}} \mathcal{J}_\delta(\{ \rho_i, g_{ij} \})$$

By the definition of \mathcal{U} we have that $g_{ij}^{(n)}$ are uniformly bounded in $L^2(S_{ij})$.

Now, we argue that by [28, Theorem IV.5.3] the PDEs given by (11) have unique solutions ρ_i continuous with respect to the inputs, i.e.,

$$\| \rho_i \|_{H^{l+2, l/2+1}} \leq C \sum_j \left(\| g_{ij} \|_{H^{l+1, (l+1)/2}(S_{ij})} + \| g_{ji} \|_{H^{l+1, (l+1)/2}(S_{ji})} \right) \quad (12)$$

for some non-integral l , when the norms on the right-hand side are well defined.

When we consider the conditions for the application of this Theorem, we see that the only unsatisfied assumption is the integrability of the coefficients with respect to an appropriate dual Sobolev space. However, we can see from the proof of the result, in particular [28, Equation (IV.7.1)], that this is used only to show that the operators,

$$\left[\partial_t + \sigma \nabla^2 + F \cdot \nabla \right] \text{ and } [\sigma \nabla + F \cdot]$$

are bounded. However, with σ constant and $F \in L^\infty(R_i)$, this also clearly holds and these are operators from $H^{l+2, l/2+1}$ to $H^{l, l/2+1}$ on R_i and $H^{l+1, l/2+1}$ on S_{ij} respectively.

For any l , we can apply the Poincaré inequality on the left and Sobolev embedding on the right of (12) to obtain,

$$\| \rho_i \|_{H^1(R_i)} \leq C \sum_j \left(\| g_{ij} \|_{L^2(S_{ij})} + \| g_{ji} \|_{L^2(S_{ji})} \right) \quad (13)$$

Thus, by the uniform boundedness of $g_{ij}^{(n)}$ from the definition of \mathcal{U} , we get the boundedness of $\rho_i^{(n)}$ and the existence of a convergent subsequence $\{ \{ \rho_i^{(n_k)} \}, \{ g_{ij}^{(n_k)} \} \}$ convergent to $(\{ \hat{\rho}_i \}, \hat{g}_{ij})$ with every $\hat{\rho}_i$ in $H^1(R_i)$ and \hat{g}_{ij} in $L^2(S_{ij})$. Passing to the limit we see that they also satisfy (11) and by the lower semicontinuity of \mathcal{J}_δ we have that

$$\inf_{\{ \rho_i \}, \{ g_{ij} \}} \mathcal{J}_\delta(\rho_i, g_{ij}) = \liminf_k \mathcal{J}_\delta(\rho_i^{(n_k)}, g_{ij}^{(n_k)}) \geq \mathcal{J}_\delta(\{ \hat{\rho}_i, g_{ij} \})$$

and $(\{ \hat{\rho}_i \}, \hat{g}_{ij})$ is optimal. Since \mathcal{J}_δ is convex and \mathcal{U} is linear, it is uniquely optimal. \square

Now consider the following weak system of PDEs, now for a unique ρ ,

$$\int_{R_i} [\partial_t \rho v + \sigma \nabla \rho \cdot \nabla v + F \cdot \nabla \rho v] dx = \sum_j \int_{S_{ij}} (n \cdot (F \rho + \frac{1}{2} \sigma \nabla \rho)) v dS_{ij} - \sum_j \int_{S_{ji}} (n \cdot (F \rho + \frac{1}{2} \sigma \nabla \rho)) v dS_{ji}, \forall v \in H^1(R_i) \quad (14)$$

We then prove the following.

Theorem 4.3 *For each $\delta > 0$ denote $(\{\rho_i^\delta\}\{g_{ij}^\delta\})$ to be the solutions to (10) as given by Theorem 4.2. We have that for any convergent subsequence as $\delta \rightarrow 0$, it holds that there exists ρ such that with $\rho_i = \rho|_{R_i}$ and $g_{ij} = n \cdot (F \rho + \frac{1}{2} \sigma \nabla \rho)|_{S_{ij}}$,*

$$\sum_i \|\rho_i^\delta - \rho_i\|_{H^1(R_i)} + \sum_{ij} \|g_{ij}^\delta - g_{ij}\|_{L^2(S_{ij})} \rightarrow 0$$

and ρ solves (14).

Proof From the definition of $(\{\rho_i^\delta\}\{g_{ij}^\delta\})$ we have that

$$\mathcal{J}_\delta(\{\rho_i^\delta\}\{g_{ij}^\delta\}) \leq \mathcal{J}_\delta(\{\rho_i\}\{g_{ij}\}),$$

which is,

$$\sum_{ij} \left[\int_{S_{ij}} (\rho_i^\delta - \rho_j^\delta)^2 dS_{ij} + \frac{\delta}{2} \int_{S_{ij}} (g_{ij}^\delta)^2 dS_{ij} \right] \leq \frac{\delta}{2} \int_{S_{ij}} g_{ij}^2 dS_{ij},$$

implying, by the uniform boundedness of g_{ij} in $L^2(S_{ij})$ that $\|\rho_i^\delta - \rho_j^\delta\|_{L^2(S_{ij})} \rightarrow 0$ as $\delta \rightarrow 0$. Furthermore, from (13) we get that $\|\rho_i^\delta\|_{H^1(R_i)}$ are uniformly bounded. Thus, with $\delta \rightarrow 0$ there is a subsequence converging to $\{\rho_i^*, g_{ij}^*\}$ over $\{H^1(R_i), L^2(S_{ij})\}$ and passing to the limit implies they satisfy (11). Furthermore $\|\rho_i^\delta - \rho_j^\delta\|_{H^1(S_{ij})} \rightarrow 0$ implies that $\rho_i^*|_{S_{ij}} = \rho_j^*|_{S_{ij}}$. Defining $\rho \in H^1(\mathbb{R}^n)$ by $\rho|_{R_i \cup (\cup_j \{S_{ij}\})} = \rho_i$ we obtain the unique solution to (14). \square

We have now proven the existence of $\rho \in H^1(\mathbb{R}^n)$ satisfying the weak form of the PDE (7) over $\cup_i R_i$, corresponding to where the coefficients are smooth almost everywhere, and with the boundary conditions (8).

4.4 Variational Flow for the Free Energy

In [22] the FP equation (7) is shown to be the gradient flow of the functional,

$$\mathcal{F}(\rho) = E(\rho) + S(\rho; \sigma) = \int f(x) \rho dx + \int \rho \log \rho dx \quad (15)$$

when $F(x) = \nabla f(x)$ and the stationary distribution is given by its minimizer. To this effect, one can consider a scheme,

$$\rho^{(k)} := \arg \min_{\rho} \frac{1}{2} W(\rho^{(k-1)}, \rho)^2 + h\mathcal{F}(\rho) \quad (16)$$

for some small h , where W refers to the Wasserstein distance. We have the following Proposition, whose proof is unchanged in our setting,

Proposition 4.1 [22, Proposition 4.1] *Given $\rho \in M$, there exists a unique solution to (16).*

Above, M is defined as

$$M := \left\{ \rho : \mathbb{R}^n \rightarrow [0, \infty) \text{ measurable, and } \int \rho(x) dx = 1, \int |x|^2 \rho(x) dx < \infty \right\}$$

Now we extend the classical main result, Theorem 5.1 in [22], to our setting, which requires a few modifications to account for the nonsmooth potential in the free energy.

Theorem 4.4 *Let $\rho^0 \in M$ with $F(\rho^0) < \infty$ and $\rho_h^{(k)}$ the solution for (16) and the interpolation $\rho_h : (0, \infty) \times \mathbb{R}^n \rightarrow [0, \infty)$ by,*

$$\rho_h(t) = \rho_h^{(k)} \text{ for } t \in [kh, (k+1)h) \text{ and } k \in \mathbb{N} \cup \{0\}$$

Then as $h \rightarrow 0$, $\rho_h(t) \rightarrow \rho(t)$ weakly in $L^1(\mathbb{R}^n)$ for all $t \in (0, \infty)$ where $\rho \in H^1((0, \infty) \times \mathbb{R}^n)$ solves (9) with initial condition $\rho(t) \rightarrow \rho^0$ strongly in $L^1(\mathbb{R}^n)$ for $t \rightarrow 0$.

Proof Let $\xi \in C_0^\infty(\mathbb{R}^n, \mathbb{R}^n)$ be a smooth vector field with bounded support, and its flux Φ_τ as $\partial_\tau \Phi_\tau = \xi \circ \Phi_\tau$ for all $\tau \in \mathbb{R}$ and $\Phi_0 = id$. The measure $\rho_\tau(y) dy$ is the push forward of $\rho^{(k)}(y) dy$ under Φ_τ . This means that

$$\int_{\mathbb{R}^n} \rho_\tau(y) \zeta(y) dy = \int_{\mathbb{R}^n} \rho^{(k)}(y) \zeta(\Phi_\tau(y)) dy$$

for all $\zeta \in C_0^0(\mathbb{R}^n)$, which implies that $\det \nabla \Phi_\tau \rho_t \circ \Phi_\tau = \rho^{(k)}$. By the properties of (16) we have that

$$\frac{1}{\tau} \left[\left(\frac{1}{2} W(\rho^{(k-1)}, \rho_\tau)^2 + h\mathcal{F}(\rho_\tau) \right) - \left(\frac{1}{2} W(\rho^{(k-1)}, \rho^{(k)})^2 + h\mathcal{F}(\rho^{(k)}) \right) \right] \geq 0. \quad (17)$$

Now, consider $\zeta = f$ and so

$$\int_{\mathbb{R}^n} \rho_\tau(y) f(y) dy = \int_{\mathbb{R}^n} \rho^{(k)}(y) f(\Phi_\tau(y)) dy$$

and also

$$\frac{1}{\tau} \left(E(\rho_\tau) - E(\rho^{(k)}) \right) = \int_{\mathbb{R}^n} \frac{1}{\tau} (f(\Phi_\tau(y)) - f(y)) \rho^{(k)}(y) dy$$

Recalling the consideration of a conservative vector field, cf. Definition 2.3 above. In the original, the term $\nabla f(y) \cdot \xi(y)$ appears, instead, we have a vector $\chi(y)$ that represents directional change in f . Specifically, we can write that

$$\frac{d}{d\tau}[E(\rho_\tau)]_{\tau=0} = \int_{\mathbb{R}^n} \chi(y) \cdot \xi(y) \rho^{(k)}(y) dy.$$

We can continue similarly as in the proof of [22, Theorem 5.1] to obtain,

$$\frac{d}{d\tau}[S(\rho_\tau)]_{\tau=0} = - \int_{\mathbb{R}^n} \rho^{(k)} \operatorname{div} \xi dy$$

and subsequently the following a priori estimates, wherein for any $T < \infty$ there exists C such that for all $N \in \mathbb{N}$ and $h \in [0, T]$ with $Nh \leq T$ there holds,

$$\begin{aligned} M(\rho_h^{(N)}) &\leq C, \\ \int_{\mathbb{R}^n} \max\{\rho_h^{(N)} \log \rho_h^{(N)}, 0\} dx &\leq C, \\ E(\rho_h^{(N)}) &\leq C, \\ \sum_{k=1}^N W(\rho_h^{(k-1)}, \rho_h^{(k)})^2 &\leq Ch \end{aligned} \tag{18}$$

which implies the existence of a convergent subsequence,

$$\rho_h \rightarrow \rho \text{ weakly in } L^1((0, T) \times \mathbb{R}^n) \text{ for all } T < \infty$$

with $\rho(t) \in M$ and $E(\rho) \in L^\infty((0, T))$, and ρ satisfying

$$- \int_{(0, \infty) \times \mathbb{R}^n} \rho(\partial_t \zeta - \chi \cdot \nabla \zeta + \Delta \zeta) dx dt = \int_{\mathbb{R}^n} \rho^0 \zeta(0) dx \text{ for all } \zeta \in C_0^\infty(\mathbb{R} \times \mathbb{R}^n) \tag{19}$$

We can proceed similarly as in the proof of Theorem 5.1 in [22] using appropriate test functions to conclude that $\rho \in L^p_{loc}((0, \infty), \mathbb{R}^n)$. Now, clearly this ρ solves (14), which has a unique solution in $H^1(\mathbb{R}^n, (0, T))$ by Theorem 4.3.

The final statements of the Theorem follow as in Theorem 5.1 in [22]. \square

5 Numerical Illustration

For an illustration, we took, the one dimensional function

$$f(x) := \begin{cases} -x - 1 & x < -1 \\ x + 1 & -1 \leq x < 0 \\ 1 - x & 0 \leq x < 1 \\ x - 1 & x \geq 1 \end{cases}, \quad F(x) := \begin{cases} -1 & x < -1 \\ [-1, 1] & x = -1 \\ 1 & -1 < x < 0 \\ [-1, 1] & x = 0 \\ -1 & 0 < x < 1 \\ [-1, 1] & x = 1 \\ 1 & x > 1 \end{cases} \tag{20}$$

The probability density function associated with minimizing the free energy is shown in Figure 1 and the result of one hundred thousand samples generated by the Metropolis Algorithm in Figure 2.




Fig. 1 Stationary Distribution $e^{-f(x)}$ Associated with (20).




Fig. 2 Histogram of 100k Metropolis Samples of (20)

As an illustration, we ran unadjusted Langevin dynamics with the Euler-Maruyama discretization, i.e., generated samples with the following iteration,

$$x_{k+1} = x_k - \epsilon g_k + \sqrt{2\epsilon} B_k, \quad g \in F(x_k), \quad B_k \sim N(0, 1). \quad (21)$$

We generated ten million samples with $\epsilon \in \{0.01, 0.001, 0.0001\}$. We plot the histograms of the final sample count in Figure 3 and plot the Wasserstein distance (computed with `scipy`) in Figure 4.




Fig. 3 Histogram for (21) at $\epsilon \in \{0.01, 0.001, 0.0001\}$ in this order.




Fig. 4 Wasserstein Distance of the Langevin-type iteration (21) and Metropolis-generated samples for $\epsilon \in \{0.01, 0.001, 0.0001\}$, in this order.

We can observe that indeed the iteration does recreate the posterior, i.e., it appears to be ergodic, although the quantity of samples required is fairly large, indeed it seems to converge in probability distance after about 3 million samples, suggesting geometric ergodicity is unlikely.

6 Discussion and Implications

The standard potential gradient diffusion process has featured prominently in both the theoretical analysis meant to give insight as to the approximation and generalization performance associated with the long-term behavior of SGD as applied to neural networks, for example, [20]. It has also featured prominently in algorithms for sampling high-dimensional data sets, e.g. [11]. It is standard for these studies to require that the drift term is Lipschitz, and as such, that the potential is continuously differentiable. In many contemporary applications, such as in (Bayesian, in the case of sampling) Neural Networks with ReLU or convolutional layers, this is not the case, and the presence of points wherein the loss function is not continuously differentiable is endemic. As such, the primary results of this paper, the existence of a solution of the stochastic differential inclusion drift as well as the existence of a Fokker-Planck equation characterizing the evolution of the probability distribution which asymptotically converges to a Gibbs distribution of the free energy, provide some basic insights into these processes without unrealistic assumptions. Specifically, even with these nonsmooth elements as typically arising in Whitney-stratifiable compositions of model and loss criteria, the overall understanding of the asymptotic macro behavior of the algorithmic processes remains as expected.

Additional insights gathered from studies of learning from studying an approximating SDE are not as straightforward to extend to the differential-inclusion setting. Namely, geometric ergodicity which requires convexity and hypocoercivity properties [46] does not clearly extend to notions of fast convergence in the non-smooth setting. The wide and shallow basins around minima become nuances, when considering that a Hessian of a loss function may not exist at certain points. Thus, considerations of mixing rate for sampling and qualitative properties of limiting distributions are interesting topics for further study in specific cases of specific stochastic differential inclusions.

Acknowledgments

We'd like to thank Thomas Surowiec for helpful discussions regarding PDE theory as related to the arguments in this paper. FVD has been supported by *REFIN* Project, grant number 812E4967 funded by Regione Puglia. VK and JM were supported by the OP VVV project CZ.02.1.01/0.0/0.0/16_019/0000765 "Research Center for Informatics" and the Czech Science Foundation (22-15524S).

References

- [1] S. Albeverio, V. Bogachev, and M. Röckner. On uniqueness of invariant measures for finite- and infinite-dimensional diffusions. *Communications on Pure and Applied Mathematics*, 52(3):325–362, 1999.
- [2] J.P. Aubin and H. Frankowska. *Set-Valued Analysis*. Birkhauser, Boston, 1990.
- [3] D. Azagra, Juan Ferrera, Fernando López-Mesas, and Yenny Rangel Oliveros. Smooth Approximation of Lipschitz functions on Riemannian manifolds. *Journal of Mathematical Analysis and Applications*, 326:1370–1378, 02 2007.
- [4] Robert Baier and Elza Farkhi. Regularity of set-valued maps and their selections through set differences. Part 1: Lipschitz continuity. *Serdica Mathematical Journal*, 39:365–390, 01 2013.
- [5] V. Bogachev and M. Röckner. A Generalization of Khasminskii's Theorem on the Existence of Invariant Measures for Locally Integrable Drifts. *Theory of Probability and Its Applications*, 45:363–378, 2001.
- [6] V.I. Bogachev, N.V. Krylov, and M. Röckner. On regularity of transition probabilities and invariant measures of singular diffusions under minimal conditions. *Communications in Partial Differential Equations*, 26(11-12):2037–2080, 2001.
- [7] Jérôme Bolte, Aris Daniilidis, and Adrian Lewis. Tame functions are semismooth. *Mathematical Programming*, 117(1):5–19, Mar 2009.
- [8] Jérôme Bolte, Aris Daniilidis, Adrian Lewis, and Masahiro Shiota. Clarke subgradients of stratifiable functions. *SIAM Journal on Optimization*, 18(2):556–572, 2007.

- [9] Jérôme Bolte and Edouard Pauwels. Conservative set valued fields, automatic differentiation, stochastic gradient methods and deep learning. *Mathematical Programming*, 188(1):19–51, 2021.
- [10] Léon Bottou, Frank E Curtis, and Jorge Nocedal. Optimization methods for large-scale machine learning. *Siam Review*, 60(2):223–311, 2018.
- [11] Giovanni Bussi and Michele Parrinello. Accurate sampling using Langevin dynamics. *Physical Review E*, 75(5):056707, 2007.
- [12] Frank H Clarke. *Optimization and nonsmooth analysis*. SIAM, 1990.
- [13] M. Coste. An introduction to o-minimal geometry. *Univ. de Rennes*, 1999.
- [14] V. Dolean, P. Jolivet, and F. Nataf. *An Introduction to Domain Decomposition Methods: Algorithms, Theory, and Parallel Implementation*. Other Titles in Applied Mathematics. Society for Industrial and Applied Mathematics, 2015.
- [15] Neil A Eklund. Boundary behavior of solutions of parabolic equations with discontinuous coefficients. *Bulletin of the American Mathematical Society*, 77(5):788–792, 1971.
- [16] Herbert Federer. *Geometric measure theory*. Classics in mathematics. Springer, 1996.
- [17] Crispin Gardiner. *Stochastic methods*, volume 4. Springer Berlin, 2009.
- [18] Alexandre Grothendieck. *Around Grothendieck's Esquisse D'un Programme*, volume 1. Cambridge University Press, 1997.
- [19] M.D. Gunzburger, J.S. Peterson, and H. Kwon. An optimization based domain decomposition method for partial differential equations. *Computers & Mathematics with Applications*, 37(10):77–93, 1999.
- [20] Wenqing Hu, Chris Junchi Li, Lei Li, and Jian-Guo Liu. On the diffusion approximation of nonconvex stochastic gradient descent. *arXiv preprint arXiv:1705.07562*, 2017.
- [21] Seizô Itô. The fundamental solution of the parabolic equation in a differentiable manifold. *Osaka Mathematical Journal*, 5(1):75–92, 1953.
- [22] Richard Jordan, David Kinderlehrer, and Felix Otto. The Variational Formulation of the Fokker-Planck Equation. *SIAM J. Math. Anal.*, 29(1):1–17, January 1998.
- [23] John Kent. Time-reversible diffusions. *Advances in Applied Probability*, 10(4):819–835, 1978.
- [24] Michał Kisielewicz. Stochastic representation of partial differential inclusions. *Journal of mathematical analysis and applications*, 353(2):592–606, 2009.
- [25] Michał Kisielewicz. *Stochastic differential inclusions and applications*. Springer, 2013.
- [26] Michał Kisielewicz. *Set-Valued Stochastic Integrals and Applications*. Springer, 2020.
- [27] Julia F Knight, Anand Pillay, and Charles Steinhorn. Definable sets in ordered structures. ii. *Transactions of the American Mathematical Society*, 295(2):593–605, 1986.
- [28] Olga Alexandrovna Ladyzhenskaya, Vsevolod Alekseevič Solonnikov,

- and Nina Nikolaevna Ural'ceva. *Linejnye i kvazilinejnye uravneniâ parabolicheskogo tipa*. Izdatel'stvo" Nauka", Glavnaâ redakciâ fiziko-matematičeskoj literatury, 1967.
- [29] Gunther Leobacher and Alexander Steinicke. Exception sets of intrinsic and piecewise Lipschitz functions. *arXiv preprint arXiv:2105.12004*, 2021.
- [30] Gunther Leobacher and Michaela Szölgényi. A strong order 1/2 method for multidimensional SDEs with discontinuous drift. *The Annals of Applied Probability*, 27(4):2383–2418, 2017.
- [31] Tao Luo, Zhi-Qin John Xu, Zheng Ma, and Yaoyu Zhang. Phase diagram for two-layer relu neural networks at infinite-width limit. *Journal of Machine Learning Research*, 22(71):1–47, 2021.
- [32] Angus Macintyre, Kenneth McKenna, and Lou van den Dries. Elimination of quantifiers in algebraic structures. *Advances in Mathematics*, 47(1):74–87, 1983.
- [33] Angus Macintyre and Eduardo D Sontag. Finiteness results for sigmoidal “neural” networks. In *Proceedings of the twenty-fifth annual ACM symposium on Theory of computing*, pages 325–334, 1993.
- [34] Sean P Meyn and Richard L Tweedie. Stability of Markovian processes III: Foster-Lyapunov criteria for continuous-time processes. *Advances in Applied Probability*, pages 518–548, 1993.
- [35] Sean P Meyn and Richard L Tweedie. *Markov chains and stochastic stability*. Springer Science & Business Media, 2012.
- [36] Robin Nittka. Regularity of solutions of linear second order elliptic and parabolic boundary value problems on Lipschitz domains. *Journal of Differential Equations*, 251(4-5):860–880, 2011.
- [37] Nikolai Ivanovich Portenko. *Generalized diffusion processes*, volume 83. American Mathematical Soc., 1990.
- [38] Robin M Schmidt, Frank Schneider, and Philipp Hennig. Descending through a crowded valley-benchmarking deep learning optimizers. In *International Conference on Machine Learning*, pages 9367–9376. PMLR, 2021.
- [39] Alexander Shevchenko, Vyacheslav Kungurtsev, and Marco Mondelli. Mean-field analysis of piecewise linear solutions for wide relu networks. *arXiv preprint arXiv:2111.02278*, 2021. Accepted to Journal of Machine Learning Research.
- [40] M. A. Shubin. Weak Bloch property and weight estimates for elliptic operators. *Séminaire Équations aux dérivées partielles (Polytechnique) dit aussi "Séminaire Goulaouic-Schwartz"*, 1989-1990.
- [41] G.V. Smirnov. *Introduction to the Theory of Differential Inclusions*. American Mathematical Society, Rhode Island, 2002.
- [42] Daniel W Stroock and SR Srinivasa Varadhan. *Multidimensional diffusion processes*. Springer, Berlin Heidelberg, 2007.
- [43] Lou van den Dries. *Tame topology and o-minimal structures*, volume 248 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, United Kingdom, 1998.

22 REFERENCES

- [44] Lou van den Dries, Angus Macintyre, and David Marker. The elementary theory of restricted analytic fields with exponentiation. *Annals of Mathematics*, 140(1):183–205, 1994.
- [45] Lou Van den Dries and Chris Miller. Geometric categories and o-minimal structures. *Duke Mathematical Journal*, 84(2):497–540, 1996.
- [46] Cédric Villani. Hypocoercivity. *arXiv preprint math/0609050*, 2006.
- [47] Joachim von Below. Classical solvability of linear parabolic equations on networks. *Journal of Differential Equations*, 72(2):316–337, 1988.
- [48] Max Welling and Yee W Teh. Bayesian learning via stochastic gradient Langevin dynamics. In *Proceedings of the 28th international conference on machine learning (ICML-11)*, pages 681–688. Citeseer, 2011.