

# Soft theorems in curved spacetime

Peng Cheng<sup>a,b</sup> and Pujian Mao<sup>a</sup>

<sup>a</sup> *Center for Joint Quantum Studies and Department of Physics,  
School of Science, Tianjin University, 135 Yaguan Road, Tianjin 300350, China*

<sup>b</sup> *Lanzhou Center for Theoretical Physics,  
Key Laboratory of Theoretical Physics of Gansu Province,  
Lanzhou University, 222 South Tianshui Road, Lanzhou 730000, Gansu, China*

ABSTRACT. In this paper, we derive a soft photon theorem in the near horizon region of the Schwarzschild black hole from the Ward identity of the near horizon large gauge transformation. The flat spacetime soft photon theorem can be recovered as a limiting case of the curved spacetime. The soft photons on the horizon are indeed soft electric hairs. This accomplishes the triangle equivalence on the black hole horizon.

## 1 Introduction

It has recently been shown that black holes can carry soft hairs [1] which provides a possible resolution of the black hole information paradox, see, e.g., [2] for a review and [3,4] for more recent developments. The discovery of the black hole soft hair is originally inspired by a fascinating equivalence relation in the infrared sector of gravity where the BMS supertranslation Ward identity is equivalent to Weinberg's soft graviton theorem [5, 6]. In the language of S-matrix, an alternative interpretation of such equivalence relation is that the soft graviton is generated by the action of a supertranslation on the quantum state. If one implements the supertranslation action on an asymptotically flat black hole, soft gravitons can be created on the horizon. This is the original proposal for soft hairs on black holes [1].

Though supertranslations are well defined in any asymptotically flat spacetime, the soft theorem is highly relying on a flat spacetime background where the scattering amplitudes are defined. Hence it is questionable if a supertranslation action can really create soft gravitons on black hole horizons. Even though one can consider that supertranslation actions should always be conceptually related to soft gravitons, the property of such soft gravitons and their relations to soft graviton theorems in curved spacetime are not clear at all and it is the aim of the present work to explore soft theorems in curved spacetime and to address their connections to soft hairs.

In this paper, we derive a soft photon theorem from the Ward identity of the near horizon large gauge transformation. In [7, 8], it was argued that the asymptotic symmetries of gauge theories are symmetries of the quantum S-matrix. We assume that this argument should be valid for any boundary of the spacetime. Considering the horizon as the inner boundary of the Schwarzschild black hole, the near horizon symmetries should be symmetries of S-matrix in the curved spacetime, at least in the near horizon region. We solve the Maxwell theory which is coupled with a general source current in the near horizon region. The near horizon charge associated to the near horizon large gauge transformation splits into a linear (hard) piece and a non-linear (soft) piece. Then the connection of soft photon theorems and asymptotic symmetries in [8] can be extended to the near horizon region of the Schwarzschild black hole with two crucial adaptations in the mode expansion of the non-linear piece of the charge. The first one is that we need to deal with the dispersion relation of the wave vector in curved spacetime. The second one is that the stationary phase approximation at null infinity (see, e.g., [2] for details) can not be applied in the near horizon region. We first write the mode expansion in the isotropic coordinates. Then the integration of the momentum in the mode expansion can be performed by the usual angular variables. The mode expansion is divergent on the horizon. We use the Feynman prescription to regularize the radial coordinate on the horizon. Thus the action of the non-linear on the state is obtained without using the stationary phase approximation. The action of the linear piece of the charge on the state can be obtained simply by introducing the commutation relation in the near horizon region. In the end, we derive the soft photon theorem in position space from the Ward identity of the near horizon large gauge transformation. Then we transform the soft factor into momentum space and discuss its relation to the one in flat spacetime. We further show that the near horizon soft photons have precisely the forms of soft electric hairs. This is a direct proof of the argument in [1] that soft particles implant soft hairs on black hole horizons. Moreover, the near horizon soft photon theorem, if it is verified from amplitudes side in curved spacetime, would accomplish the triangle relation at the horizon. Hopefully, the soft theorem derived in the present work could be a stepping stone for a fuller understanding of physics in the near horizon region.

## 2 Near horizon Maxwell theory

We consider Maxwell theory coupled to a conserved matter current in the near horizon region of Schwarzschild black hole. We adopt the retarded coordinates  $(u, r, z, \bar{z})$  where  $(z, \bar{z})$  are the complex stereographic coordinates and they related to the usual angular variables  $(\theta, \phi)$  by  $z = \cot \frac{\theta}{2} e^{i\phi}$ . The line element of Schwarzschild black hole is

$$ds^2 = -f(r)du^2 - 2dudr + 2\Omega^2\gamma_{z\bar{z}}dzd\bar{z}, \quad \gamma_{z\bar{z}} = \frac{2}{(1+z\bar{z})^2}, \quad (1)$$

where  $f(r) = \frac{r}{r+2M}$  and  $\Omega = (r + 2M)$ . The horizon  $\mathcal{H}$  is precisely the submanifold  $r = 0$  in the retarded spherical coordinates, with topology  $S^2 \times \mathbb{R}$ .




Figure 1: The Penrose diagram of the outside region of the Schwarzschild black hole.

The retarded coordinates can only cover half of the horizon, i.e.,  $\mathcal{H}^-$  as demonstrated in Figure 1. The other half  $\mathcal{H}^+$  can be described by advanced coordinates  $(v, r, z, \bar{z})$ . The crossing point of  $\mathcal{H}^-$  and  $\mathcal{H}^+$  is the bifurcation point  $B$ . In this work, we will only paid attention to the  $\mathcal{H}^-$  part while everything can be similarly repeated on  $\mathcal{H}^+$ . The near horizon symmetries on  $\mathcal{H}^-$  and  $\mathcal{H}^+$  can be connected by the near bifurcation symmetries [9].

We consider a gauge field  $A_\mu$  and a conserved matter current  $J_\mu$  with the radial gauge and the near horizon fall-off conditions as follows

$$A_r = 0, \quad A_u = \mathcal{O}(r), \quad A_z = \mathcal{O}(1), \quad J_r = 0, \quad J_u = \mathcal{O}(1), \quad J_z = \mathcal{O}(1). \quad (2)$$

The radial component of the current is set to zero by the ambiguities of a conserved current.<sup>1</sup> This is a more consistent choice of working in the radial gauge. We assume the following ansatz as initial data for the near horizon expansion of the gauge fields

$$A_{z(\bar{z})} = A_{z(\bar{z})}^0(u, z, \bar{z}) + \sum_{m=1}^{\infty} A_{z(\bar{z})}^m(u, z, \bar{z})r^m, \quad (3)$$

and the components of the current

$$J_{z(\bar{z})} = J_{z(\bar{z})}^0(u, z, \bar{z}) + \sum_{m=1}^{\infty} J_{z(\bar{z})}^m(u, z, \bar{z})r^m. \quad (4)$$

<sup>1</sup>When a conserved current is derived from a global symmetry, it is naturally defined up to the equivalence  $J^\mu \sim J^\mu + \nabla_\nu k^{[\mu\nu]}$ , so it makes more sense to consider equivalence classes of currents  $[J^\mu]$  (see [10, 11] for details).

From the Maxwell's equations,  $\nabla_\mu F^{\mu\nu} = J^\nu$ , and the current conservation condition,  $\nabla_\mu J^\mu = 0$ , in a  $r$ -expansion, we obtain

$$J_u = \frac{J_u^0(u, z, \bar{z})}{\Omega^2} + \frac{1}{\Omega^2} \int_0^r dr' [\gamma_{z\bar{z}}^{-1}(\partial_z J_{\bar{z}} + \partial_{\bar{z}} J_z)] , \quad (5)$$

$$A_u = \frac{A_u^0(u, z, \bar{z})}{2M} - \frac{A_u^0(u, z, \bar{z})}{\Omega} + \int_0^r dr' \frac{1}{\Omega^2} \int_0^{r'} dr'' [\gamma_{z\bar{z}}^{-1}(\partial_z \partial_{r''} A_{\bar{z}} + \partial_{\bar{z}} \partial_{r''} A_z)] , \quad (6)$$

$$\partial_u A_u^0 = -\gamma_{z\bar{z}}^{-1} \partial_u (\partial_z A_{\bar{z}}^0 + \partial_{\bar{z}} A_z^0) - J_u^0 , \quad (7)$$

$$2\partial_u \partial_r A_z = \partial_z \partial_r A_u + \partial_r (f \partial_r A_z) + \partial_z \left[ \frac{1}{\Omega \gamma_{z\bar{z}}} (\partial_{\bar{z}} A_z - \partial_z A_{\bar{z}}) \right] - J_z , \quad (8)$$

where  $J_u^0(u, z, \bar{z})$  and  $A_u^0(u, z, \bar{z})$  are integration constants. We have omitted one equation that can be obtained from the last one above by simply swapping  $z \rightleftharpoons \bar{z}$ . The time evolution of the coefficients in the expansion of  $A_{z(\bar{z})}$  are uniquely determined except the leading  $A_{z(\bar{z})}^0$ . We refer to  $\partial_u A_z^0$  as the electromagnetic “news” since it reflects the propagation of electromagnetic waves.

To summarize, the general solution to the Maxwell system in spacetime (1) with the prescribed conditions (2) is completely determined in terms of the initial data  $A_u^0(u_0, z, \bar{z})$ ,  $A_z^m(u_0, z, \bar{z})$ ,  $A_{\bar{z}}^m(u_0, z, \bar{z})$  ( $m \geq 1$ ), the functions  $A_z^0(u, z, \bar{z})$ ,  $A_{\bar{z}}^0(u, z, \bar{z})$  and the matter current. The latter is characterized by the source functions  $J_u^0(u, z, \bar{z})$ ,  $J_z^m(u, z, \bar{z})$ ,  $J_{\bar{z}}^m(u, z, \bar{z})$  ( $m \geq 0$ ).

The radial gauge condition in (2) leaves residual gauge transformations of the form  $\delta A_\mu = \partial_\mu \varepsilon(z, \bar{z})$ , where  $\varepsilon(z, \bar{z})$  is an arbitrary function on the horizon. These transformations can be interpreted as symmetries on the horizon

$$\delta A_z^0 = \partial_z \varepsilon(z, \bar{z}) , \quad \delta A_{\bar{z}}^0 = \partial_{\bar{z}} \varepsilon(z, \bar{z}) , \quad (9)$$

We define the associated charge at the bifurcation point on the horizon

$$Q_\varepsilon = \int_B dz d\bar{z} \gamma_{z\bar{z}} \varepsilon(z, \bar{z}) \Omega^2 F_{ru} = \int_{\mathcal{H}^-} dz d\bar{z} du \gamma_{z\bar{z}} \varepsilon(z, \bar{z}) \Omega^2 \partial_u F_{ru} , \quad (10)$$

where in the second equality we assumed  $A_u^0|_{i^-} = 0$ , meaning that in the far past the system contains no bulk electric charge. Inserting the solutions in the charge yields

$$Q_\varepsilon = - \int_{\mathcal{H}^-} dz d\bar{z} du \varepsilon(z, \bar{z}) (\gamma_{z\bar{z}} J_u^0 + \partial_u \partial_z A_{\bar{z}}^0 + \partial_u \partial_{\bar{z}} A_z^0) . \quad (11)$$

### 3 Near horizon soft photon theorem

If a symmetry is generated by a charge  $Q$ , the associated Ward identity in the S-matrix language reads as

$$\langle \text{out} | Q^{\text{out}} - Q^{\text{in}} | \text{in} \rangle = 0 . \quad (12)$$

For a spontaneously broken symmetry, the charge must act non-linearly on the states. Otherwise it would annihilate the vacuum. Hence one can decompose the charge into linear and non-linear pieces  $Q = Q_L + Q_{NL}$ . The Ward identity for a broken charge can be written as

$$\langle \text{out} | Q_{NL}^{\text{out}} - Q_{NL}^{\text{in}} | \text{in} \rangle = -\langle \text{out} | Q_L^{\text{out}} - Q_L^{\text{in}} | \text{in} \rangle. \quad (13)$$

The analysis in the previous section is carried out only near  $\mathcal{H}^-$ . We assume that all *in* states in the S-matrix are defined on  $\mathcal{H}^-$ . Particles are created on  $\mathcal{H}^-$  and then propagated to the bulk and ended on  $\mathcal{H}^+$ . In this sense,  $\mathcal{H}^-$  and  $\mathcal{H}^+$  are similar to the past and future null infinity  $I^-$  and  $I^+$ . We will only paid attention to the *in* part in the present work. The analysis of the *out* part can be carried out analogously up to a identification at the bifurcation point. For notational brevity, we suppress the *in* label in the following text.

Let us now decompose the charge (11) into a piece containing the “news” and the rest containing the sources<sup>2</sup>

$$Q_{NL} = \int_{\mathcal{H}^-} dz d\bar{z} du \varepsilon \partial_u \partial_{\bar{z}} A_z^0, \quad (14)$$

$$Q_L = \frac{1}{2} \int_{\mathcal{H}^-} dz d\bar{z} du \gamma_{z\bar{z}} \varepsilon J_u^0. \quad (15)$$

These are respectively the non-linear and linear pieces above. We keep only the anti-holomorphic terms for the news part. The sources are split into two parts. That is where the half factor comes in (15). Such split is consistent with the treatment in [8] where the authors consider only the sector of the phase space with no long-range magnetic fields. In our case, such conditions lead to  $F_{z\bar{z}} = 0$  at  $B$  and  $i^0$ . Correspondingly, the helicity of soft photon is decoupled. Hence one can only deal with either the holomorphic part or the anti-holomorphic part.

The soft photon theorem is obtained from the Ward identity by a concrete choice of  $\varepsilon(z, \bar{z})$  at null infinity [8]. We follow the same choice for the near horizon case

$$\varepsilon(z, \bar{z}) = \frac{1}{w - z}, \quad (16)$$

for which  $\partial_{\bar{z}} \varepsilon = -2\pi \delta^2(z - w)$ . The derivation of the soft theorem is parallel with the cases at null infinity. In order to obtain the proper action of the charges on the *in* states, one has to define canonical commutation relations on the horizon.

### 3.1 Mode expansion in Schwarzschild spacetime

The mode expansion of fields in curved spacetime needs to be adapted to the dispersion relation of the momentum of a moving particle in curved spacetime. It is more convenient to work in the isotropic coordinates system  $(t, \rho, z, \bar{z})$  where the line element of the

<sup>2</sup>We drop the total minus sign in the charge as it won't affect the Ward identity.

Schwarzschild black hole is

$$ds^2 = - \left( \frac{2\rho - M}{2\rho + M} \right)^2 dt^2 + \frac{(2\rho + M)^4}{16\rho^4} (d\rho^2 + \rho^2 d\mathcal{S}^2), \quad (17)$$

where  $d\mathcal{S}^2$  is the line element of a unit sphere. The isotropic coordinates are connected to the retarded coordinates in (1) by

$$t = u + r + 2M + 2M \ln \frac{r}{2M}, \quad \left(1 + \frac{M}{2\rho}\right)^2 \rho = r + 2M. \quad (18)$$

The constant time slices of the line element (17) are conformally flat. It would be convenient to rewrite the conformally flat part in the Cartesian coordinates  $\vec{x} = (x^1, x^2, x^3)$ , with coordinate transformation

$$x^1 = \rho \frac{(z + \bar{z})}{1 + z\bar{z}}, \quad x^2 = \rho \frac{-i(z - \bar{z})}{1 + z\bar{z}}, \quad x^3 = \rho \frac{1 - z\bar{z}}{1 + z\bar{z}}. \quad (19)$$

Then the line element (17) becomes

$$ds^2 = - \left( \frac{2\rho - M}{2\rho + M} \right)^2 dt^2 + \frac{(2\rho + M)^4}{16\rho^4} d\vec{x}^2. \quad (20)$$

For a massless particle moving along a trajectory parametrized by  $\lambda$ , the four momentum is defined as

$$p_\mu = g_{\mu\nu} \frac{dx^\nu}{d\lambda}. \quad (21)$$

The Schwarzschild solution (20) has one timelike Killing vector

$$K^\mu = (\partial_t)^\mu = (1, 0, 0, 0). \quad (22)$$

One can define the corresponding conserved energy as

$$\omega = -p_\mu K^\mu = -p_0. \quad (23)$$

Thus, the dispersion relation for the massless particle moving along a null trajectory is

$$- \left( \frac{2\rho + M}{2\rho - M} \right)^2 \omega^2 + \frac{16\rho^4}{(2\rho + M)^4} \vec{p}^2 = 0, \quad (24)$$

where  $\vec{p}$  is the three momentum of the conformally flat part.

For a free massless scalar field  $\Phi(x)$ , we can write the field operator in isotropic coordinates as [12]

$$\Phi(x^\mu) = -\frac{1}{(2\pi)^4} \int d\omega d^3\vec{p} \tilde{\phi}(p_\mu) e^{ip \cdot x}. \quad (25)$$

Inserting the dispersion relation as a delta function in the integral and working out the integration of  $\omega$

$$- \int d\omega 2\pi\delta \left[ - \left( \frac{2\rho + M}{2\rho - M} \right)^2 \omega^2 + \frac{16\rho^4}{(2\rho + M)^4} \vec{p}^2 \right] = \left( \frac{2\rho - M}{2\rho + M} \right)^2 \frac{1}{2\omega}, \quad (26)$$

the expansion (25) can be further written as

$$\Phi(x^\mu) = \frac{1}{(2\pi)^3} \left( \frac{2\rho - M}{2\rho + M} \right)^2 \int \frac{d^3\vec{p}}{2\omega} \tilde{\phi}(p_\mu) e^{ip \cdot x} \Big|_{-p_0=\omega} \quad (27)$$

$$= \frac{1}{(2\pi)^3} \left( \frac{2\rho - M}{2\rho + M} \right)^2 \int \frac{d^3\vec{p}}{2\omega} [\mathbf{a}(p_\mu) e^{ip \cdot x} + \mathbf{a}^\dagger(p_\mu) e^{-ip \cdot x}] \Big|_{\omega>0}. \quad (28)$$

In the flat spacetime case one only has  $\frac{1}{2\omega}$  in the mode expansion. Now we have an extra factor  $\frac{(2\rho-M)^2}{(2\rho+M)^2}$  because of the dispersion relation (24).

For a  $U(1)$  gauge field  $A_\mu$ , the mode expansion will involve two polarization vectors orthogonal to the propagating direction. The mode expansion of  $A_\mu$  respect to the dispersion relation (24) is

$$A_\mu(x) = \sum_{\alpha=\pm} \frac{1}{(2\pi)^3} \left( \frac{2\rho - M}{2\rho + M} \right)^2 \int \frac{d^3\vec{p}}{2\omega} [\epsilon_\mu^{*\alpha} \mathbf{a}_\alpha(p) e^{ip \cdot x} + \epsilon_\mu^\alpha \mathbf{a}_\alpha^\dagger(p) e^{-ip \cdot x}]. \quad (29)$$

The polarization vectors satisfy the normalization condition  $\epsilon_\alpha^\mu \epsilon_{\beta\mu}^* = \delta_{\alpha\beta}$ . It is of convenience to parametrize the photon four-momentum by

$$p^\mu = \frac{\omega}{1+z\bar{z}} \frac{4\rho^2}{4\rho^2 - M^2} \left( \frac{(2\rho + M)^3}{4\rho^2(2\rho - M)} (1+z\bar{z}), (z+\bar{z}), -i(z-\bar{z}), (1-z\bar{z}) \right), \quad (30)$$

and

$$p_\mu = \frac{\omega}{1+z\bar{z}} \frac{(2\rho + M)^3}{4\rho^2(2\rho - M)} \left( -\frac{4\rho^2(2\rho - M)}{(2\rho + M)^3} (1+z\bar{z}), (z+\bar{z}), -i(z-\bar{z}), (1-z\bar{z}) \right). \quad (31)$$

In contrast to the flat spacetime case where a null momentum can be fully characterized by the energy and a point on the celestial sphere, the null momentum in Schwarzschild spacetime always has the radial dependence. The polarization vector can be written as

$$\begin{aligned} \epsilon^{+\mu} &= \frac{1}{\sqrt{2}} \frac{4\rho^2}{(2\rho + M)^2} \left( \frac{(2\rho + M)^3}{4\rho^2(2\rho - M)} \bar{z}, 1, -i, -\bar{z} \right), \\ \epsilon^{-\mu} &= \frac{1}{\sqrt{2}} \frac{4\rho^2}{(2\rho + M)^2} \left( \frac{(2\rho + M)^3}{4\rho^2(2\rho - M)} z, 1, i, -z \right) \end{aligned} \quad (32)$$

It's easy to check that

$$p_\mu \epsilon^{\pm\mu} = 0, \quad \epsilon_\alpha^\mu \epsilon_{\beta\mu}^* = \delta_{\alpha\beta}. \quad (33)$$

In the  $(z, \bar{z})$  coordinates, we have

$$\epsilon_z^+ = \frac{(M + 2\rho)^2}{2\sqrt{2}\rho(1+z\bar{z})}, \quad \epsilon_{\bar{z}}^- = \frac{(M + 2\rho)^2}{2\sqrt{2}\rho(1+z\bar{z})}. \quad (34)$$

Inserting the expression

$$\exp[ip_\mu x^\mu] = \exp \left[ -i\omega t + i\omega \frac{(2\rho + M)^3}{4\rho(2\rho - M)} \hat{p} \cdot \hat{x} \right], \quad (35)$$

where  $\hat{p}$  and  $\hat{x}$  are unit vectors along  $\vec{p}$  and  $\vec{x}$ , the mode expansion of  $A_\mu$  can be written as

$$A_\mu = \sum_{\alpha=\pm} \frac{(2\rho - M)^2}{(2\rho + M)^2} \int \frac{d^3\vec{p}}{(2\pi)^3} \frac{1}{2\omega} \epsilon_\mu^\alpha \mathbf{a}_\alpha \exp \left[ -i\omega t + i\omega \frac{(2\rho + M)^3}{4\rho(2\rho - M)} \hat{p} \cdot \hat{x} \right] + c.c.. \quad (36)$$

The spatial part of the metric is conformally flat, so there is no extra Jacobian in the momentum integral. We can further write the integration measure as

$$\int \frac{d^3\vec{p}}{(2\pi)^3} \frac{1}{2\omega} = \frac{(2\rho + M)^9}{(2\rho)^6(2\rho - M)^3} \int \frac{\omega d\omega}{8\pi^2} \int_0^\pi d\Theta \sin \Theta, \quad (37)$$

where  $\Theta$  is the angle between  $\hat{p}$  and  $\hat{x}$ . Note that we have used the dispersion relation (24). With the integration measure (37), the mode expansion can be rewritten as

$$A_\mu = \sum_{\alpha=\pm} \frac{(2\rho + M)^7}{(2\rho)^6(2\rho - M)} \int \frac{\omega d\omega}{8\pi^2} \int_0^\pi d\Theta \sin \Theta \epsilon_\mu^\alpha \mathbf{a}_\alpha e^{-i\omega t + i\omega \frac{(2\rho + M)^3}{4\rho(2\rho - M)} \cos \Theta} + c.c.. \quad (38)$$

The mode expansion in the flat spacetime (see, e.g., in [2, 8, 13]) can be recovered by setting  $M = 0$ . Near the horizon, we have  $\rho \rightarrow \frac{M}{2}$ . Let  $R = \rho - \frac{M}{2}$ . Hence  $R = r + O(r^2)$  in the near horizon region. The  $\Theta$  integration in the mode expansion can be worked out directly with the help of the relation

$$\int_0^\pi d\Theta \sin \Theta e^{i\omega \frac{2(M+R)^3}{R(M+2R)} \cos \Theta} = \frac{R(M+2R)}{\omega(M+R)^3} \sin \left[ \frac{2\omega(M+R)^3}{R(M+2R)} \right]. \quad (39)$$

The exponential in the integrand blows up as  $R \rightarrow 0$ . At null infinity, a stationary-phase approximation has been applied in this step to control the divergence when  $r \rightarrow \infty$ , see, e.g., in [2, 8, 13]. Here we follow the Feynman prescription to introduce a small imaginary part to the radial coordinate  $R \rightarrow R + i\tau$ . This will keep the radial coordinate  $R$  continuous on the real axis. Then we will trace all the contributions without any approximation. For notational brevity, we will momentarily suppress the imaginary part  $i\tau$ . In terms of  $R$ , we obtain the mode expansion as

$$A_\mu = \sum_{\alpha=\pm} \frac{8(M+R)^4}{\pi^2(M+2R)^5} \left( \frac{R^2}{M^2 + 2MR} \right)^{-2iM\omega} \times \int d\omega \epsilon_\mu^\alpha \mathbf{a}_\alpha e^{-i\omega u - i\omega \frac{2(M+R)^2}{M+2R}} \sin \left[ \frac{2\omega(M+R)^3}{R(M+2R)} \right] + c.c.. \quad (40)$$

Note that we have adopted the retarded time  $u$  using the transformation in (18). The near horizon field  $A_z^0$  is related to the plane wave modes by

$$A_z^0 = \frac{16\sqrt{2}}{\pi^2(1+z\bar{z})} \frac{(M+R)^6}{(M+2R)^6} \left( \frac{R^2}{M^2 + 2MR} \right)^{-2iM\omega} \times \int d\omega \sin \left[ \frac{2\omega(M+R)^3}{R(M+2R)} \right] \left( \mathbf{a}_+ e^{-i\omega u - i\omega \frac{2(M+R)^2}{M+2R}} + \mathbf{a}_-^\dagger e^{i\omega u + i\omega \frac{2(M+R)^2}{M+2R}} \right). \quad (41)$$



### 3.2 Soft photon theorem in position space

The non-linear part of the conserved charge in (14) can be expressed as

$$Q_{\text{NL}} = \lim_{R \rightarrow 0} \int_{\mathcal{H}^-} dz d\bar{z} du \varepsilon(z, \bar{z}) \partial_u \partial_{\bar{z}} A_z. \quad (42)$$

Inserting  $\varepsilon(z, \bar{z}) = \frac{1}{w-z}$  which leads to  $\partial_{\bar{z}} \varepsilon = 2\pi \delta^2(z-w)$  and using the Fourier relation

$$\int_{-\infty}^{\infty} du \partial_u F(u) = 2\pi i \lim_{\omega \rightarrow 0} \left[ \omega \tilde{F}(\omega) \right], \quad (43)$$

where we define  $F(u) = \int_{-\infty}^{\infty} d\omega e^{i\omega u} \tilde{F}(\omega)$ , the non-linear part of charge with the mode expansion (41) is reduced to

$$Q_{\text{NL}} = \frac{-128\sqrt{2}i}{(1+w\bar{w})} \frac{M^2}{R} \lim_{\omega \rightarrow 0} \left[ \omega^2 \mathbf{a}_+ + \omega^2 \mathbf{a}_-^\dagger \right]. \quad (44)$$

Hence, the action of this piece of the charge on the *in* state is

$$\langle \text{out} | Q_{\text{NL}} | \text{in} \rangle = \frac{-128\sqrt{2}i}{(1+w\bar{w})} \frac{M^2}{R} \lim_{\omega \rightarrow 0} \langle \text{out} | \omega^2 \mathbf{a}_-^\dagger | \text{in} \rangle \quad (45)$$

Regarding the linear piece (15), for simplicity, we restrict ourselves to scalar charged (with charge  $Q_e$ ) matter with current  $J_\mu = iQ_e(\bar{\Phi} \nabla_\mu \Phi - \Phi \nabla_\mu \bar{\Phi})$ .<sup>3</sup> Since the derivatives act on complex scalar field  $\Phi$ , it makes no difference whether using covariant derivative or normal derivative. With respect to (5), the current at leading order is  $J_u^0 = i\Omega^2 Q_e(\bar{\Phi} \partial_u \Phi - \Phi \partial_u \bar{\Phi})$ . Then one can apply the canonical commutation relation as the flat spacetime case [13]

$$[\partial_u \bar{\Phi}^0(u, z, \bar{z}), \Phi^0(u', w, \bar{w})] = \frac{i}{4} \gamma_{w\bar{w}}^{-1} \Theta(u-u') \delta^2(z-w), \quad (46)$$

to get the action of the linear charge on the *in* state as

$$\langle \text{out} | Q_{\text{L}}^{(0)} | \text{in} \rangle = \sum_{k=1}^n \frac{M^2 Q_k}{(w-w_k)} \langle \text{out} | \text{in} \rangle, \quad (47)$$

where  $Q_k$  is the electric charge of the  $k$ -th particle. We have assumed there are  $n$  incoming particles.

Finally the Ward identity (13) can yield a soft photon theorem in Schwarzschild space-time as

$$\lim_{\omega \rightarrow 0} \langle \text{out} | \mathbf{a}_-^\dagger | \text{in} \rangle = \frac{1}{128} \frac{R}{i\omega} \left( \frac{1+w\bar{w}}{\sqrt{2}\omega} \sum_{k=1}^n \frac{Q_k}{w-w_k} \right) \langle \text{out} | \text{in} \rangle. \quad (48)$$

We have several remarks on this soft theorem as follows:

<sup>3</sup>Note that the determinant of the metric is omitted in the current with respect to the equation of motion of the Maxwell theory in previous section.

- The square of the soft energy shows up in the denominator of the soft factor. But this does not mean that the soft factor is of order  $\frac{1}{\omega^2}$  in the soft expansion. The soft theorem is derived from the Ward identity of the near horizon symmetry which is restricted in the near horizon region, i.e.,  $R \rightarrow 0$ .
- We use the Feynman prescription to regularize the radial coordinate  $R$  on the horizon. A similar treatment is not needed at null infinity. Because the null infinity is conformally introduced. The divergence from  $r \rightarrow \infty$  can be absorbed into the conformal factor. Hence the quantities defined at the null infinity should be finite. But the horizon is part of the spacetime and particles can freely cross it. So the divergence from the horizon should be regularized. On the horizon, we set  $R = i\tau$  and  $\tau$  being infinitesimal.
- Let  $k = \frac{\tau}{128\omega}$  be a finite constant. The soft factor in (48) recovers the flat spacetime soft photon factor when  $k = 1$ .
- A curious point is that the soft factor does not include the mass parameter  $M$  which indicates the curvature of the spacetime. Nevertheless, it may not be a surprising point since we derive the soft theorem in position space. The hard particles locally can not see the spacetime curvature due to the equivalent principle. The soft factor is built on the data of hard particles. So the soft factor in position space may not see the spacetime curvature. But the dispersion relation of the null momenta involve  $M$  and we will show in the next page that the soft factor in momentum space does include the mass parameter  $M$ .

### 3.3 Soft photon theorem in momentum space

We consider the *in* states (the hard particles) are created on the horizon. Then the hard momenta in the near horizon region can be parametrized as

$$q_k^\mu = \frac{E_k}{1 + z_k \bar{z}_k} \frac{M}{4R} \left( \frac{4M}{R} (1 + z_k \bar{z}_k), (z_k + \bar{z}_k), -i(z_k - \bar{z}_k), (1 - z_k \bar{z}_k) \right), \quad (49)$$

and

$$q_{k\mu} = \frac{E_k}{1 + z_k \bar{z}_k} \frac{4M}{R} \left( -\frac{R}{4M} (1 + z_k \bar{z}_k), (z_k + \bar{z}_k), -i(z_k - \bar{z}_k), (1 - z_k \bar{z}_k) \right). \quad (50)$$

Using the soft momentum in (31) and the polarization vectors in (32), one can show that

$$\sum_{k=1}^n \frac{Q_k q_k \cdot \epsilon}{q_k \cdot p} = \frac{R}{\omega M} \left( \frac{1 + z\bar{z}}{\sqrt{2}} \sum_{k=1}^n \frac{Q_k}{z - z_k} \right). \quad (51)$$

One can see that the expression  $\sum_{k=1}^n \frac{Q_k q_k \cdot \epsilon}{q_k \cdot p}$  which is singular in the soft limit in flat spacetime can be regular on the horizon as  $R \rightarrow 0$ . But it is indeed singular in the

flat limit  $M \rightarrow 0$ . Applying the relation in (51), we obtain the soft photon theorem in momentum space as

$$\lim_{\omega \rightarrow 0} \langle \text{out} | \mathbf{a}_-^\dagger | \text{in} \rangle = \frac{1}{128} \frac{M}{i\omega} \sum_{k=1}^n \frac{Q_k q_k \cdot \epsilon}{q_k \cdot p} \langle \text{out} | \text{in} \rangle. \quad (52)$$

Naively, one can read from (52) that  $M = 0$  will kill the soft factor and hence the soft theorem. Though setting  $M = 0$  will not bring any divergence in the computation, it will lead to the fact that  $R = 0$  is just a point rather than a three dimensional null hypersurface. There is no horizon at all at  $R = 0$  in such case. Somehow the regularization in  $R \rightarrow R + i\tau$  when  $R = 0$  prevents one taking the limit  $M \rightarrow 0$ . Consequently, we need to introduce a corresponding regularization  $M \rightarrow M + im$  where  $m$  is infinitesimal. Then the flat spacetime soft photon theorem yields that  $m = 128\omega$  in the soft limit.

## 4 Soft photons as soft electric hair on the horizon

### 4.1 Near horizon electromagnetic memories and soft electric hairs

We follow the prescription in [14] to formulate the near horizon electromagnetic memories. Consider that an electromagnetic shockwave passes through the horizon of the black hole. The information about the electromagnetic wave is encoded in the change of the surface charge. Suppose that there is initially no electromagnetic field and the system finally settles into a stationary solution with flat gauge connections on the horizon  $\partial_z A_{\bar{z}}^0 = \partial_{\bar{z}} A_z^0$  [8]. According to (11), the change of the surface charge is just the charge evaluated at a later time

$$\Delta Q_\epsilon = - \int_{u=u_f} dz d\bar{z} \epsilon(z, \bar{z}) (\partial_z A_{\bar{z}}^0 + \partial_{\bar{z}} A_z^0), \quad (53)$$

where we only consider the contribution from electromagnetic shockwave which will yield the near horizon electromagnetic memory. The permanent change of the surface charge (53) is called the near horizon electromagnetic memory and it is fully characterized by the change of  $A_z^0$  and  $A_{\bar{z}}^0$ . The zero mode of the charge (53) is vanishing. So the electromagnetic shockwave has implanted soft electric hairs on the black hole horizon [1].

Alternatively, the near horizon electromagnetic memory can be derived from the near horizon large transformation as a precise example of the equivalence between memories and asymptotic symmetries [1, 15]. The action of the infinitesimal near horizon large transformation, denoted by  $\chi(z, \bar{z})$ , on the electromagnetic fields is

$$\delta_\chi A_z = \partial_z \chi(z, \bar{z}), \quad \delta_\chi A_{\bar{z}} = \partial_{\bar{z}} \chi(z, \bar{z}). \quad (54)$$

For the case of initially no electromagnetic field, the final electromagnetic fields are just

$$A_z^0 = \partial_z \chi(z, \bar{z}), \quad A_{\bar{z}}^0 = \partial_{\bar{z}} \chi(z, \bar{z}). \quad (55)$$

The change of the surface charge (53) is

$$\Delta Q_\varepsilon = -2 \int_{u=u_f} dz d\bar{z} \varepsilon(z, \bar{z}) \partial_z \partial_{\bar{z}} \chi, \quad (56)$$

As we will show in the next subsection that the soft electric hairs can be implanted by soft photons on the horizon.

## 4.2 Near horizon soft photons as soft hairs

As a common understanding, the low-energy limit of a quantum result should be connected to a classical result. Soft photon theorems connect a  $n + 1$  particles state to a  $n$  particles state in a low-energy expansion of the extra soft photon in the form

$$M_{n+1}(p_1, \dots, p_n, [q; \varepsilon^\pm]) = \frac{1}{\omega} S^{(0)\pm} M_n(p_1, \dots, p_n) + \mathcal{O}(1). \quad (57)$$

In the low-energy limit, the soft theorem should have certain relevance to classical computation. Following the proposal in [16, 17] at null infinity, one can read from the soft theorem that the soft factors can be interpreted as the expectation value of near horizon fields fluctuation

$$S^{(0)\pm} = \lim_{\omega \rightarrow 0} \omega \frac{M_{n+1}^\pm}{M_n}. \quad (58)$$

Then the expectation value of near horizon fields fluctuation with soft photon emission is simply related to the change of classical fields as

$$A_\mu \varepsilon^{\mu\pm} = S^{(0)\pm}. \quad (59)$$

Hence

$$A_z = \varepsilon_z^{*+} S^{(0)+} + \varepsilon_z^{*-} S^{(0)-}, \quad (60)$$

where we assume the system initially has no electromagnetic field.

Inserting the soft photon factors (48) into (60) (recall the complex conjugate), we obtain

$$A_z = 2k \sum_{k=1}^n \frac{Q_k}{z - z_k}, \quad A_{\bar{z}} = 2k \sum_{k=1}^n \frac{Q_k}{\bar{z} - \bar{z}_k}. \quad (61)$$

Consider  $k = \frac{\tau}{128\omega}$  as a constant factor, the above electromagnetic fields are exactly the form of a memory discussed previously in this section. Hence the soft photons in the near horizon region naturally has the form of soft electric hairs. Acting with  $\partial_{\bar{z}(z)}$  on  $A_{z(\bar{z})}$  and considering the real part, we get

$$\partial_{\bar{z}} A_z + \partial_z A_{\bar{z}} = 8\pi k \sum_{k=1}^n Q_k \delta^2(z - z_k). \quad (62)$$

Inserting this into the charge (53), one obtains

$$\Delta Q_\varepsilon = -8\pi k \sum_{k=1}^n Q_k \varepsilon(z_k, \bar{z}_k), \quad (63)$$

where  $(z_k, \bar{z}_k)$  are the locations of the hard particles projected on the horizon which generate the electromagnetic shockwave.

## 5 Conclusion and discussion

We derive a soft photon theorem in the near horizon region of the Schwarzschild black hole. The soft factor can recover the flat spacetime soft photon factor in the zero mass limit  $M \rightarrow 0$ . The soft factor interpreted as the expectation value of near horizon fields fluctuation is in the form of soft electric hairs on the black hole horizon. Hence this verifies that the soft particles on black hole horizons can implant soft hairs.

In the derivation of the soft photon theorem, we restrict ourselves to the case that all particles are in the near horizon region. But we expect our analysis to be generic to the formalism and relevant issues can be addressed elsewhere. We point out some for future directions. The most direct one is to extend our derivation to other curved spacetime with horizon, such as Kerr black hole or cosmological solution. Another glaring one is to study the scattering with particles in both near horizon region and asymptotic region. In particular, one may need to enhance the near horizon or asymptotic symmetries to symplectic symmetries [18]. Inspired by the sub-leading soft photon theorem [19] (see also [13]) in the flat spacetime, one may wonder if there are more soft photon theorems in the near horizon region. It is more promising in the case of gravity where one has two types of near horizon symmetries, namely near horizon supertranslations and superrotations [20]. Naively, one should be able to derive two soft graviton theorems in the near horizon region. Another interesting point in the gravity case is to include higher derivative terms. They are order-suppressed in the asymptotic region, see also discussions in [21] from the soft theorem side. However in the near horizon region, the higher derivative terms have significant influence on the near horizon charges [22]. If they can affect the near horizon soft graviton theorem would be a very meaningful investigation.

## Acknowledgments

The authors thank Laura Donnay, Zhengwen Liu, Shahin Sheikh-Jabbari and Jun-Bao Wu for useful discussions. This work is supported in part by the National Natural Science Foundation of China (NSFC) under Grant No. 11905156 and No. 11935009. P.C. is also supported in part by NSFC Grant No. 12047501.

## References

- [1] S. W. Hawking, M. J. Perry, and A. Strominger, “Soft Hair on Black Holes,” *Phys. Rev. Lett.* **116** no. 23, (2016) 231301, arXiv:1601.00921 [hep-th].
- [2] A. Strominger, “Lectures on the infrared structure of gravity and gauge theory,” 1703.05448v2.
- [3] S. Pasterski and H. Verlinde, “Hps meets amps: How soft hair dissolves the firewall,” *JHEP* **09** (2021) 099, arXiv:2012.03850 [hep-th].
- [4] P. Cheng and Y. An, “Soft black hole information paradox: Page curve from maxwell soft hair of a black hole,” *Phys. Rev. D* **103** no. 12, (2020) 126020, arXiv:2012.14864 [hep-th].
- [5] A. Strominger, “On BMS Invariance of Gravitational Scattering,” *JHEP* **07** (2014) 152, arXiv:1312.2229 [hep-th].
- [6] T. He, V. Lysov, P. Mitra, and A. Strominger, “BMS supertranslations and Weinberg’s soft graviton theorem,” *JHEP* **05** (2015) 151, arXiv:1401.7026 [hep-th].
- [7] A. Strominger, “Asymptotic Symmetries of Yang-Mills Theory,” *JHEP* **07** (2014) 151, arXiv:1308.0589 [hep-th].
- [8] T. He, P. Mitra, A. P. Porfyriadis, and A. Strominger, “New Symmetries of Massless QED,” *JHEP* **10** (2014) 112, arXiv:1407.3789 [hep-th].
- [9] H. Adami, D. Grumiller, S. Sadeghian, M. M. Sheikh-Jabbari, and C. Zwikel, “T-Witts from the horizon,” *JHEP* **04** (2020) 128, arXiv:2002.08346 [hep-th].
- [10] G. Barnich and F. Brandt, “Covariant theory of asymptotic symmetries, conservation laws and central charges,” *Nucl. Phys.* **B633** (2002) 3–82, arXiv:hep-th/0111246 [hep-th].
- [11] E. Conde and P. Mao, “Remarks on asymptotic symmetries and the subleading soft photon theorem,” *Phys. Rev. D* **95** no. 2, (2017) 021701, arXiv:1605.09731 [hep-th].
- [12] V. Mukhanov and S. Winitzki, “Mode expansions cheat sheet,” in *Introduction to quantum effects in gravity*, pp. 216–217. Cambridge University Press, 2007.
- [13] V. Lysov, S. Pasterski, and A. Strominger, “Low’s Subleading Soft Theorem as a Symmetry of QED,” *Phys. Rev. Lett.* **113** no. 11, (2014) 111601, arXiv:1407.3814 [hep-th].

- [14] H. Adami, D. Grumiller, M. M. Sheikh-Jabbari, V. Taghiloo, H. Yavartanoo, and C. Zwickel, “Null boundary phase space: slicings, news & memory,” *JHEP* **11** (2021) 155, arXiv:2110.04218 [hep-th].
- [15] L. Donnay, G. Giribet, H. A. González, and A. Puhm, “Black hole memory effect,” *Phys. Rev. D* **98** no. 12, (2018) 124016, arXiv:1809.07266 [hep-th].
- [16] A. Strominger and A. Zhiboedov, “Gravitational Memory, BMS Supertranslations and Soft Theorems,” *JHEP* **01** (2016) 086, arXiv:1411.5745 [hep-th].
- [17] P. Mao and H. Ouyang, “Note on soft theorems and memories in even dimensions,” *Phys. Lett. B* **774** (2017) 715–722, arXiv:1707.07118 [hep-th].
- [18] G. Compère, P. Mao, A. Seraj, and M. M. Sheikh-Jabbari, “Symplectic and Killing symmetries of AdS<sub>3</sub> gravity: holographic vs boundary gravitons,” *JHEP* **01** (2016) 080, arXiv:1511.06079 [hep-th].
- [19] F. E. Low, “Bremsstrahlung of very low-energy quanta in elementary particle collisions,” *Phys. Rev.* **110** (1958) 974–977.
- [20] L. Donnay, G. Giribet, H. A. Gonzalez, and M. Pino, “Supertranslations and Superrotations at the Black Hole Horizon,” *Phys. Rev. Lett.* **116** no. 9, (2016) 091101, arXiv:1511.08687 [hep-th].
- [21] H. Elvang, C. R. T. Jones, and S. G. Naculich, “Soft Photon and Graviton Theorems in Effective Field Theory,” *Phys. Rev. Lett.* **118** no. 23, (2017) 231601, arXiv:1611.07534 [hep-th].
- [22] H.-S. Liu and P. Mao, “Near horizon gravitational charges,” *JHEP* **05** (2022) 123, arXiv:2201.10308 [hep-th].