

PROJECTIVE FRAÏSSÉ LIMITS OF GRAPHS WITH CONFLUENT EPIMORPHISMS

WŁODZIMIERZ J. CHARATONIK, ALEKSANDRA KWIATKOWSKA,
AND ROBERT P. ROE

ABSTRACT. We show that the class of finite connected graphs with confluent epimorphism is a projective Fraïssé class and we investigate the continuum obtained as the topological realization of the projective Fraïssé limit. In particular, we show that this continuum is indecomposable, but not hereditarily indecomposable, one-dimensional, Kelley, pointwise self-homeomorphic, but not homogeneous. Moreover, it is hereditarily unicoherent and each point is the top of the Cantor fan.

1. INTRODUCTION

In [9] Irwin and Solecki introduced the concept of a projective Fraïssé limit analogous to injective Fraïssé limit from model theory. They considered finite linear (combinatorial) graphs, where the only constraint on the mappings was that they preserve the edge relation, and showed that the topological realization of the Fraïssé limit is the pseudo-arc. Bartořova and Kwiatkowska continued this study in [1], where they considered finite (combinatorial) rooted trees and found that in that case the topological realization of the Fraïssé limit is the Lelek fan. In [12], Panagiotopoulos and Solecki introduced appropriate definitions for connectedness and monotone maps on combinatorial graphs using the edge relation. They considered finite connected graphs with monotone epimorphisms and showed that the topological realization of the Fraïssé limit is the Menger curve. In [7], Charatonik and Roe considered finite trees with epimorphisms, which are monotone, confluent, order-preserving, retractions, light, etc. As topological realizations of the various Fraïssé limits they obtained known continua, such as the Cantor fan and the generalized Ważewski dendrite D_3 , as well as previously unknown continua.

Date: June 27, 2022.

2020 Mathematics Subject Classification. 03C98, 54D80, 54E40, 54F15, 54F50.

Key words and phrases. projective Fraïssé limit, continuum theory, topological graphs, confluent maps, homogeneous.

In the present article, we study topological properties of the continuum obtained as the topological realization of the Fraïssé limit of the class of finite connected graphs with confluent epimorphisms. We summarize our results in the following theorem.

Theorem 1.1. *The class \mathcal{G} of finite connected graphs and confluent epimorphisms is a projective Fraïssé class. Its projective Fraïssé limit \mathbb{G} has transitive edge relation and its topological realization $|\mathbb{G}|$ is a continuum, which has the following properties:*

- (1) *it is a Kelley continuum;*
- (2) *it is one-dimensional;*
- (3) *it is an indecomposable continuum;*
- (4) *all arc components are dense;*
- (5) *each point is the top of the Cantor fan;*
- (6) *it is pointwise self-homeomorphic;*
- (7) *it is hereditarily unicoherent;*
- (8) *it is not homogeneous.*

A major and somewhat surprising result of the article is that the continuum $|\mathbb{G}|$ is not homogeneous, which we accomplish in Section 9. Rogers [13] proved that each homogeneous, hereditarily indecomposable curve (i.e. continuum of dimension 1) is tree-like. Note that we cannot apply his theorem since $|\mathbb{G}|$ is not hereditarily indecomposable. Interestingly, it turns out that $|\mathbb{G}|$ is indecomposable and hereditarily unicoherent.

2. DEFINITIONS AND BACKGROUND RESULTS

A *graph* is an ordered pair $A = (V(A), E(A))$, where $E(A) \subseteq V(A)^2$ is a reflexive and symmetric relation on $V(A)$. The elements of $V(A)$ are called *vertices* of graph A and elements of $E(A)$ are called *edges*. If E is a set of edges, the symmetrization of E is $E^S = \{\langle x, y \rangle \in E \text{ or } \langle y, x \rangle \in E\}$. If A and B are graphs then $A \setminus B$ denotes a graph with $V(A \setminus B) = V(A) \setminus V(B)$ and $\langle x, y \rangle \in E(A \setminus B)$ if and only if $x, y \in V(A \setminus B)$ and $\langle x, y \rangle \in E(A)$.


A *topological graph* K is a graph $(V(K), E(K))$, whose domain $V(K)$ is a 0-dimensional, compact, second-countable (thus has a metric d) space and $E(K)$ is a closed, reflexive and symmetric subset of $V(K)^2$. A topological graph is an example of a topological \mathcal{L} -structure, in the sense of Irwin-Solecki [9].

Given two graphs A and B , a function $f: A \rightarrow B$ is a *homomorphism* if it maps edges to edges, i.e. $\langle a, b \rangle \in E(A)$ implies $\langle f(a), f(b) \rangle \in E(B)$. A homomorphism f is an *epimorphism* if it is moreover surjective on both vertices and edges. An *isomorphism* is an injective epimorphism.

The theory of projective Fraïssé limits was developed in [9] and further refined in [12]. We literally recall their definition here.

Definition 2.1. Let \mathcal{F} be a class of finite graphs with a fixed class of epimorphisms between the structures in \mathcal{F} . We say that \mathcal{F} is a *projective Fraïssé class* if

- (1) \mathcal{F} is countable up to isomorphism, that is, any sub-collection of pairwise non-isomorphic structures of \mathcal{F} is countable;
- (2) epimorphisms are closed under composition and each identity map is an epimorphism;
- (3) for $B, C \in \mathcal{F}$ there exist $D \in \mathcal{F}$ and epimorphisms $f: D \rightarrow B$ and $g: D \rightarrow C$; and
- (4) for $A, B, C \in \mathcal{F}$ and for every two epimorphisms $f: B \rightarrow A$ and $g: C \rightarrow A$, there exist $D \in \mathcal{F}$ and epimorphisms $f_0: D \rightarrow B$ and $g_0: D \rightarrow C$ such that $f \circ f_0 = g \circ g_0$, i.e. the diagram (D1) commutes.

(D1) 

The last property is known as the *projective amalgamation property*.

Definition 2.2. Given $\varepsilon > 0$, a mapping $f: G \rightarrow H$, where G is a metric space with metric d , is said to be an ε -map if for each $x \in H$ the diameter of $f^{-1}(x)$ is less than ε .

The following, from [7], is a slight modification of the basic result that first appeared in [9].

Theorem 2.3. *Let \mathcal{F} be a projective Fraïssé class with a fixed class of epimorphisms between the structures of \mathcal{F} . There exists a unique topological graph \mathbb{F} such that*

- (1) *for each $A \in \mathcal{F}$, there exists an epimorphism from \mathbb{F} onto A ;*
- (2) *for $A, B \in \mathcal{F}$ and epimorphisms $f: \mathbb{F} \rightarrow A$ and $g: \mathbb{F} \rightarrow B$ there exists an epimorphism $h: \mathbb{F} \rightarrow B$ such that $f = g \circ h$.*
- (3) *For every $\varepsilon > 0$ there is a graph $G \in \mathcal{F}$ and an epimorphism $f: \mathbb{F} \rightarrow G$ such that f is an ε -map.*

The topological graph \mathbb{F} from Theorem 2.3 is called the *projective Fraïssé limit* of \mathcal{F} .

Notation 2.4. Given a sequences F_n of finite graphs and epimorphisms $\alpha_n: F_{n+1} \rightarrow F_n$ we denote the inverse sequence by $\{F_n, \alpha_n\}$. For $m > n$ we let $\alpha_n^m = \alpha_n \circ \dots \circ \alpha_{m-1}$ and note that $\alpha_n^{n+1} = \alpha_n$. Given an inverse sequence $\{F_n, \alpha_n\}$ the associated inverse limit space is the subspace of the product space $\prod F_i$ determined by $\{(x_1, x_2, \dots) : x_i \in F_i \text{ and } x_i = \alpha_i(x_{i+1})\}$ and is denoted as $\mathbf{F} = \varprojlim \{F_n, \alpha_n\}$. Further, we denote the canonical projection from the inverse limit space \mathbf{F} to the n th factor space F_n by α_n^∞ .

We define the edge relation on \mathbf{F} as follows. If $\mathbf{x} = (x_1, x_2, \dots)$ and $\mathbf{y} = (y_1, y_2, \dots)$ are points in \mathbf{F} then $\langle \mathbf{x}, \mathbf{y} \rangle$ is an edge in \mathbf{F} if and only if for all n , $\langle x_n, y_n \rangle$ is an edge in F_n .

We will frequently use the following observations:

Lemma 2.5. *Let $\{F_n, \alpha_n\}$ be an inverse sequence of finite graphs, where each α_n is an epimorphism. Let F denote its inverse limit.*

- (1) *If A is a finite graph and $f: F \rightarrow A$ is an epimorphism, then there is n and an epimorphism $\gamma: F_n \rightarrow A$ such that $f = \gamma \circ \alpha_n^\infty$.*
- (2) *Suppose that $\{G_m, \beta_m\}$ is another inverse sequence of finite graphs, where each β_m is an epimorphism, such that F is its inverse limit. Then there are subsequences (n_i) and (m_j) and there are epimorphisms $g_i: F_{n_i} \rightarrow G_{m_i}$ and $h_i: G_{m_{i+1}} \rightarrow F_{n_i}$ such that $\alpha_{n_i}^{n_{i+1}} = h_i g_{i+1}$ and $\beta_{m_i}^{m_{i+1}} = g_i h_i$.*

Proof. For (1), take any n such that the partition $\{(\alpha_n^\infty)^{-1}(x) : x \in F_n\}$ refines the partition $\{f^{-1}(x) : x \in A\}$. We let for $z \in F_n$, $\gamma(z)$ to be $f(t)$ for any $t \in (\alpha_n^\infty)^{-1}(z)$. It is straightforward to see that γ is an epimorphism. For (2), if we already constructed (g_i) and (h_i) for $i < i_0$, to get g_{i_0} apply part (1) to $f_1 = \beta_{m_{i_0}}^\infty$ and get n_{i_0} and g_{i_0} such that $f_1 = g_{i_0} \circ \alpha_{n_{i_0}}^\infty$. Next, we apply part (1) to $f_2 = \alpha_{n_{i_0}}^\infty$ and get m_{i_0+1} and h_{i_0} such that $f_2 = h_{i_0} \circ \beta_{m_{i_0+1}}^\infty$. Those g_i and h_i are as required. \square

Definition 2.6. Given a projective Fraïssé class \mathcal{F} an inverse sequence $\{F_n, \alpha_n\}$ where $F_n \in \mathcal{F}$ and $\alpha_n: F_{n+1} \rightarrow F_n$ are epimorphisms is said to be a *Fraïssé sequence* for \mathcal{F} if the following two conditions hold.

- (1) For any $G \in \mathcal{F}$ there is an n and an epimorphism from F_n onto G ;
- (2) For any n , any pair $G, H \in \mathcal{F}$, and any epimorphisms $g: H \rightarrow G$ and $f: F_n \rightarrow G$ there exists $m > n$ and an epimorphism $h: F_m \rightarrow H$ such that $g \circ h = f \circ \alpha_n^m$ i.e the diagram (D2) commutes.

$$(D2) \quad \begin{array}{ccc} F_n & \xleftarrow{\alpha_n^m} & F_m \\ f \downarrow & & \downarrow h \\ G & \xleftarrow{g} & H \end{array}$$

The following follows from [9, Theorem 2.4].

Theorem 2.7. *Given a projective Fraïssé class \mathcal{F} there is a Fraïssé sequence $\{F_i, g_i\}$ for \mathcal{F} such that the inverse limit $\mathbf{F} = \varprojlim \{F_i, g_i\}$ is isomorphic to the projective Fraïssé limit \mathbb{F} .*

The following result is Proposition 2.3 of [3].

Proposition 2.8. *Let \mathcal{F} be a projective Fraïssé class. Let $\{A_n, \alpha_n\}$ be an inverse sequence in \mathcal{F} . Assume that for each $A \in \mathcal{F}$, $n \in \mathbb{N}$, and epimorphism $f: A \rightarrow A_n$, there exists $m \geq n$ and an epimorphism $g: A_m \rightarrow A$ such that $fg = \alpha_n^m$. Then $\{A_n, \alpha_n\}$ is a Fraïssé sequence for \mathcal{F} . The converse holds as well.*

Definition 2.9. A finite graph T is a *tree* if for every two distinct vertices $a, b \in T$ there is a unique finite sequence $v_0 = a, v_1, \dots, v_n = b$ of vertices in T such that for every $i \in \{0, 1, \dots, n-1\}$ we have $\langle v_i, v_{i+1} \rangle \in E(T)$ and $v_i \neq v_{i+1}$. Let n be a natural number, a vertex $p \in T$ has *order* n ($\text{ord}(p) = n$) if there are n non-degenerate edges in T that contain p . If $\text{ord}(p) = 1$ then p is an *end-vertex*, if $\text{ord}(p) = 2$ then p is an *ordinary vertex*, and if $\text{ord}(p) \geq 3$ then p a *ramification vertex*.

Definition 2.10. Given a topological graph K , if the edge relation $E(K)$ is also a transitive relation, then it is an equivalence relation and K is known as a *prespace*. The quotient space $K/E(K)$ is called the *topological realization* of K and is denoted by $|K|$. Throughout we will use π to denote the quotient map $\pi: K \rightarrow |K|$.

The following is Theorem 2.18 of [7]. It allows us to show, in some cases, transitivity of the edge relation, and thus the existence of the topological realization.

Theorem 2.11. *Suppose that \mathcal{F} is a projective Fraïssé class of graphs and for every $G \in \mathcal{F}$, for all pairwise different $a, b, c \in V(G)$ such that $\langle a, b \rangle \in E(G)$ and $\langle b, c \rangle \in E(G)$ there is a graph H and an epimorphism $f_G^H: H \rightarrow G$ such that for all vertices $p, q, r \in V(H)$ such that $f_G^H(p) = a$, $f_G^H(q) = b$, and $f_G^H(r) = c$ we have $\langle p, q \rangle \notin E(H)$ or $\langle q, r \rangle \notin E(H)$. Then the edge relation of the projective Fraïssé limit \mathbb{F} of \mathcal{F} is transitive.*

Remark 2.12. Note that Theorem 2.11 in fact implies that for every $a \in \mathbb{F}$ there is at most one $b \in \mathbb{F}$, $b \neq a$, such that $\langle a, b \rangle \in E(\mathbb{F})$. Hence, $\text{card}(\pi^{-1}(x)) \leq 2$ for each $x \in |\mathbb{F}|$.

Below we give definitions for connectedness properties for topological graphs analogous to similarly named properties for continua. The first definition below originally appeared in [9] and the remainder were introduced in [7].

Definition 2.13. Given a topological graph G , a subset S of $V(G)$ is *disconnected* if there are two nonempty closed subsets P and Q of S such that $P \cup Q = S$ and if $a \in P$ and $b \in Q$, then $\langle a, b \rangle \notin E(G)$. A subset S of $V(G)$ is *connected* if it is not disconnected. A graph G is connected if $V(G)$ is connected.

Definition 2.14. Given a graph G , a subset S of $V(G)$, and a vertex $a \in S$, the *component of S containing a* is the largest connected subset C of S that contains a ; in other words $C = \bigcup\{P \subseteq S : a \in P \text{ and } P \text{ is connected}\}$.

Definition 2.15. Let F be a graph, A and B be subgraphs of F and $a, b \in F$. We say that A is *adjacent to B* if $A \cap B = \emptyset$ and there are $a \in V(A)$ and $b \in V(B)$ such that $\langle a, b \rangle \in E(F)$. We say that a is *adjacent to A* if $\{a\}$ is adjacent to A , and similarly, a is *adjacent to b* if $\{a\}$ is adjacent to $\{b\}$.

Definition 2.16. We say that a topological graph G is an *arc* if it is connected and there are at most two vertices $a, b \in V(G)$ such that for every $x \in V(G) \setminus \{a, b\}$ the graph $G \setminus \{x\}$ is not connected. The vertices a and b are called *end-vertices* of the arc and we say that G *joins a and b* . Note, we allow a single vertex to be an arc.

Definition 2.17. A topological graph G is called *arcwise connected* if for every two vertices $a, b \in V(G)$ there is a subgraph of G that is an arc and contains a and b . If $p \in V(G)$ then the *arc-component of G at p* is $\bigcup\{A : A \text{ is an arc in } G \text{ and } p \in A\}$.

Definition 2.18. A topological graph G is called *unicoherent* if for every two closed connected subsets P and Q of $V(G)$ with $P \cup Q = V(G)$ the intersection $P \cap Q$ is connected. The G is called *hereditarily unicoherent* if for every two closed connected subsets P and Q of $V(G)$ the intersection $P \cap Q$ is connected.

Definition 2.19. A hereditarily unicoherent and arcwise connected topological graph is called a *dendroid*.

Definition 2.20. By a *rooted graph* we mean a topological graph G with a distinguished vertex $r(G)$, called the *root*. On a rooted graph we define an order \leq by $x \leq y$ if every arc containing $r(G)$ and y contains x .

Definition 2.21. A rooted dendroid T is a *fan* if for every $s, t \in T$ which are incomparable in the order from Definition 2.20, if $p \notin \{s, t\}$ such that $p \leq s$ and $p \leq t$, then p is the root of T .

Definition 2.22. A rooted graph X with the root $r(X)$ is called a *smooth dendroid* if X is a dendroid and the order \leq , defined in 2.20, is closed. A smooth dendroid which is a fan is called a *smooth fan*.

3. CONFLUENT EPIMORPHISMS

Let us start with recalling the following definition introduced in [7].

Definition 3.1. Given two topological graphs G and H , an epimorphism $f: G \rightarrow H$ is called *confluent* if for every connected subset Q of $V(H)$ and every component C of $f^{-1}(Q)$ we have $f(C) = Q$. Equivalently, if for every connected subset Q of $V(H)$ and every vertex $a \in V(G)$ such that $f(a) \in Q$, there is a connected set C of $V(G)$ such that $a \in C$ and $f(C) = Q$. It is called *monotone* if preimages of connected sets are connected. Clearly, every monotone epimorphism is confluent.

Observation 3.2. *If $f: X \rightarrow Y$ and $g: Y \rightarrow Z$ are confluent epimorphisms between topological graphs, then $g \circ f: X \rightarrow Z$ is a confluent epimorphism.*

Let us recall some results shown in [7]

Theorem 3.3. *Given two finite graphs G and H , the following conditions are equivalent for an epimorphism $f: G \rightarrow H$:*

- (1) f is confluent;
- (2) for every edge $P \in E(H)$ and for every vertex $a \in V(G)$ such that $f(a) \in P$, there is an edge $E \in E(G)$ and a connected set $R \subseteq V(G)$ such that $E \cap R \neq \emptyset$, $f(E) = P$, $a \in R$, and $f(R) = \{f(a)\}$;
- (3) for every edge $P \in E(H)$ and every component C of $f^{-1}(P)$ there is an edge E in C such that $f(E) = P$.

The following theorem is an analogue to [4, I, p. 213]. Its proof is a direct consequence of definitions.

Theorem 3.4. *Let $f: F \rightarrow G$ be a confluent epimorphism between topological graphs, let Q be a closed connected subgraph of G , and let C be a component of $f^{-1}(Q)$. Then the restriction $f|_C$ is confluent.*

Lemma 3.5. *Let $\{F_n, \alpha_n\}$ be an inverse sequence of finite graphs, where each α_n is a confluent epimorphism. Let F denote its inverse limit. Then for each n , the map $\alpha_n^\infty: F \rightarrow F_n$ is a confluent epimorphism.*

Proof. Compare with Theorem 2.1.23 in [10]. See also the proof of Theorem 6.2. □

We also recall [7, Proposition 4.13]

Lemma 3.6. *Consider the epimorphisms $f: F \rightarrow G$ and $g: G \rightarrow H$ between topological graphs. If the composition $g \circ f: F \rightarrow H$ is confluent, then g is confluent.*

It is not the case, as the following example shows, that if $f: G \rightarrow H$ is a confluent epimorphism and K is a subgraph of G then $f|_K: K \rightarrow H$ must be confluent.

Example 3.7. Let K consist of the vertices $a_1, c_1, a_2, d_1, a_3, b_1$ and edges $\langle a_1, c_1 \rangle, \langle c_1, a_2 \rangle, \langle a_2, d_1 \rangle, \langle d_1, a_3 \rangle, \langle a_3, b_1 \rangle, \langle b_1, a_1 \rangle$. Let G be the graph K with the additional vertices b_2, c_2, d_2 and edges $\langle b_2, a_2 \rangle, \langle b_2, b_1 \rangle, \langle c_2, a_3 \rangle, \langle c_2, c_1 \rangle, \langle d_2, a_1 \rangle, \langle d_2, d_1 \rangle$. Finally, let H be the triod with vertices A, B, C, D and edges $\langle A, B \rangle, \langle A, C \rangle, \langle A, D \rangle$. Define $f: G \rightarrow H$ by $f(a_i) = A, f(b_i) = B, f(c_i) = C, f(d_i) = D$. One can use Theorem 3.3 to check that $f: G \rightarrow H$ is confluent but $(f|_K)^{-1}(\langle A, B \rangle)$ contains the component $\{a_2\}$ so condition (3) of Theorem 3.3 is not satisfied by $f|_K: K \rightarrow H$.

The construction below was introduced in [7].

Definition 3.8. For given graphs A, B, C and epimorphisms $f: B \rightarrow A, g: C \rightarrow A$ by *standard amalgamation procedure* we mean the graph D defined by $V(D) = \{(b, c) \in B \times C : f(b) = g(c)\}; E(D) = \{ \langle (b, c), (b', c') \rangle : \langle b, b' \rangle \in E(B) \text{ and } \langle c, c' \rangle \in E(C) \};$ and $f_0((b, c)) = b, g_0((b, c)) = c$. Note that f_0 and g_0 are epimorphisms and $f \circ f_0 = g \circ g_0$.

Theorem 3.9. *The class of finite connected graphs with confluent epimorphisms is a projective Fraïssé class.*

Proof. We need to see that Definition 2.1 holds for this class. Conditions (1) and (2) are easy to check. By [7, Corollary 4.11] and the

standard amalgamation procedure we see that condition (4) holds. Finally, condition (3) follows from condition (4) by noting that given finite connected graphs B and C if we let A the graph consisting of a single vertex and $f: B \rightarrow A$, $g: C \rightarrow A$ be constant maps, which are confluent, then condition (4) gives the existence of a connected graph D and the required confluent epimorphisms from D onto B and C . \square

Theorem 3.10. *If \mathcal{G} is the class of finite connected graphs with confluent epimorphisms and \mathbb{G} is the projective Fraïssé limit of \mathcal{G} then the edge relation $E(\mathbb{G})$ is transitive.*

Proof. To see that \mathbb{G} has a transitive edge relation note that given G , a finite connected graph containing edges $\langle a, b \rangle$ and $\langle b, c \rangle$, we may let H be the graph with $V(H) = V(G) \cup \{s\}$ and $E(H) = E(G) \cup \{\langle a, s \rangle, \langle s, b \rangle\} \setminus \{\langle a, b \rangle\}$ and $f_G^H: H \rightarrow G$ be the confluent epimorphism defined by $f_G^H(s) = a$ and $f_G^H = id$ otherwise. Then H satisfies conditions of Theorem 2.11, so \mathbb{G} has a transitive edge relation. \square

Notation 3.11. Throughout the remainder of the article we denote the class of finite connected graphs with confluent epimorphisms by \mathcal{G} and the projective Fraïssé limit of the class \mathcal{G} by \mathbb{G} . We denote the topological realization of \mathbb{G} by $|\mathbb{G}|$.

4. KELLEY CONTINUUM AND DIMENSION

Here we make a couple of observations about $|\mathbb{G}|$, which follow easily from known results.

The following definition adapts the idea of a Kelley continuum to topological graphs. It was first given in [7, Definition 6.3].

Definition 4.1. A topological graph X is called *Kelley* if X is connected and for every closed and connected set $K \subseteq V(X)$, every vertex $p \in K$, and every sequence $p_n \rightarrow p$ of vertices in X there are closed and connected sets K_n such that $p_n \in K_n$ and $\lim K_n = K$. By compactness, this definition is equivalent to saying that for every $\varepsilon > 0$ there is $\delta > 0$ such that for each two vertices $p, q \in X$ satisfying $d(p, q) < \delta$, and for each connected set $K \subseteq X$ such that $p \in K$ there is a connected set $L \subseteq X$ satisfying $q \in L$ and $H(K, L) < \varepsilon$. Note that every finite graph is Kelley.

Theorem 4.2. *The continuum $|\mathbb{G}|$ is a Kelley continuum.*

Proof. That $|\mathbb{G}|$ is a Kelley continuum follows from [7, Theorem 6.5] and [7, Observation 6.4]. \square

Theorem 4.3. *Suppose G is a topological graph with a transitive edge relation and such that $\text{card}(\pi^{-1}(x)) \leq 2$, where π is the quotient map. Then $\dim |G| \leq 1$.*

Proof. It is known, see e.g. [11, Theorem 22.2], that for a surjective map $f: X \rightarrow Y$ between compact sets, if $\text{card}(\text{bd } f^{-1}(y)) \leq m$ for all $y \in Y$, then $\dim(Y) \leq \dim(X) + m - 1$. In our case $\dim(V(G)) = 0$ and $\text{card}(\text{bd}(\pi^{-1}(x))) = \text{card}(\pi^{-1}(x)) \leq 2$, by Remark 2.12, so the conclusion follows. \square

Corollary 4.4. *The continuum $|G|$ is one-dimensional.*

5. INDECOMPOSABLE TOPOLOGICAL GRAPHS


Definition 5.1. A connected topological graph G is called *decomposable* if there are two closed connected subgraphs A and B such that $V(G) = V(A) \cup V(B)$, $V(A) \neq V(G)$, and $V(B) \neq V(G)$. A topological graph G is called *indecomposable* if it is not decomposable.

Theorem 5.2. *Suppose \mathcal{F} is a projective Fraïssé class of connected graphs and for every graph $F \in \mathcal{F}$ there is a graph $G \in \mathcal{F}$ and a confluent epimorphism $f_F^G: G \rightarrow F$ such that for every two connected graphs $A, B \subseteq G$ such that $V(G) = V(A) \cup V(B)$ we have $f_F^G(V(A)) = V(F)$ or $f_F^G(V(B)) = V(F)$. Then the projective Fraïssé limit \mathbb{F} of \mathcal{F} is an indecomposable graph.*

Proof. Suppose the projective Fraïssé limit \mathbb{F} of \mathcal{F} is a decomposable graph, i.e. there are two closed connected subgraphs A and B such that $V(\mathbb{F}) = V(A) \cup V(B)$, $V(A) \neq V(\mathbb{F})$, and $V(B) \neq V(\mathbb{F})$. Consider vertices $\mathbf{a} \in V(A) \setminus V(B)$ and $\mathbf{b} \in V(B) \setminus V(A)$ and let $\varepsilon > 0$ be such that $\varepsilon < d(\mathbf{x}, \mathbf{a})$ for every $\mathbf{x} \in V(B)$ and $\varepsilon < d(\mathbf{x}, \mathbf{b})$ for every $\mathbf{x} \in V(A)$. Let $F \in \mathcal{F}$ be a graph and let $f_F: \mathbb{F} \rightarrow F$ be a confluent epimorphism, which is an ε -map. Then $f_F(V(A)) \cup f_F(V(B)) = V(F)$, $f_F(V(A)) \neq V(F)$, and $f_F(V(B)) \neq V(F)$. Let the graph G and the epimorphism f_F^G be as in assumption of the Theorem, and let $f_G: \mathbb{F} \rightarrow G$ be a confluent epimorphism such that $f_F = f_F^G \circ f_G$. Then $f_F^G(f_G(V(A))) = V(F)$ or $f_F^G(f_G(V(B))) = V(F)$, contrary to the established properties of $f_F(V(A))$ and $f_F(V(B))$. \square

Theorem 5.3. *The class \mathcal{G} satisfies assumptions of Theorem 5.2, so its projective Fraïssé limit \mathbb{G} is indecomposable.*

Proof. Given graph F , let $\{E_1, E_2, \dots, E_n\}$ be the set of all nondegenerate edges and let $E_i = \langle a_i, b_i \rangle$ for each $i \in \{1, 2, \dots, n\}$.



Define the graph G by the following conditions (see the picture above, each ellipse represents a copy of the graph F):

- (1) $V(G) = V(F) \times \{1, 2, \dots, 2n + 1\}$;
- (2) $E(G) = E(F) \times \{1, 2, \dots, 2n + 1\} \cup \bigcup_{i=1}^n \{ \langle (a_i, i), (b_i, i + 1) \rangle, \langle (a_i, n + i), (b_i, n + i + 1) \rangle \}$.

Define f_F^G to be the projection onto the first coordinate and note that f_F^G is confluent. Now suppose that A and B are connected subgraphs of G such that $V(A) \cup V(B) = V(G)$. Note that if A intersects $F \times \{i\}$ and $F \times \{i + j\}$ for some $j \geq 1$ then, since A is connected, it must contain the edges between $F \times \{i + k\}$ and $F \times \{i + k + 1\}$, for each $k = 0, 1, \dots, j - 1$.

One of A or B , say A , must intersect at least $n + 1$ of the copies of F . It follows that $f_F^G(A) = F$. Hence, by Theorem 5.2, \mathbb{G} is indecomposable. \square

The following observations follow from the fact that a topological graph G is connected if and only if its topological realization $|G|$ is connected.

Observation 5.4. *Suppose G is an indecomposable topological graph with a transitive edge relation; then the topological realization $|G|$ is an indecomposable continuum.*

Corollary 5.5. *The continuum $|\mathbb{G}|$ is indecomposable.*

6. ARC COMPONENTS, PSEUDO-ARCS, AND CANTOR FANS

We show in this section that arc components are dense in both \mathbb{G} and $|\mathbb{G}|$, $|\mathbb{G}|$ contains a pseudo-arc, and each point of $|\mathbb{G}|$ is a vertex of the Cantor fan.

We start with the following lemma.

Lemma 6.1. *Given finite topological graphs G and H , a confluent epimorphism $f: G \rightarrow H$, an arc $A \subseteq H$ with an end-vertex a , and vertex $b \in V(G)$ such that $f(b) = a$ then there is an arc $B \subseteq G$ with one of the end-vertices b such that $f|_B: B \rightarrow A$ is a monotone epimorphism.*

Proof. Let $A = \{a_1 = a, a_2, \dots, a_n\}$ be an arc with $a = a_1$ and $\langle a_i, a_{i+1} \rangle \in E(H)$, $i = 1, \dots, n-1$. Let C_1 be the component of $f^{-1}(\{a_1 = a, a_2\})$ that contains the vertex b and note that C_1 contains an arc B_1 which has b as an end-vertex, whose image is the set $\{a_1, a_2\}$, such that $f|_{B_1}$ is monotone, and $|f^{-1}(a_2) \cap B_1| = 1$. In a similar way we may construct an arc B_2 in the component C_2 of $f^{-1}(\{a_2, a_3\})$ such that B_2 contains the vertex from $f^{-1}(a_2) \cap B_1$ as an end-vertex, $f|_{B_2}$ is monotone, $f(B_2) = \{a_2, a_3\}$, and $|f^{-1}(a_3) \cap B_2| = 1$. Continuing the same way, we construct an arc $B = B_1 \cup B_2 \cup \dots \cup B_{n-1}$ that has the required properties. \square

Theorem 6.2. *Each arc component of \mathbb{G} is dense in \mathbb{G} .*

Proof. Let $\mathbf{a}, \mathbf{b} \in \mathbb{G}$ and let $\varepsilon > 0$ be given. Write \mathbb{G} as an inverse limit of an inverse sequence $\{G_n, g_n\}$. Let N be such that for $n > N$, g_n^∞ is an ε -map. Write $\mathbf{a} = (a_0, a_1, a_2, \dots)$ with $a_i \in G_i$, and let $b_N = g_N^\infty(\mathbf{b})$. Let A_N be an arc in G_N with end-vertices a_N and b_N . Apply Lemma 6.1 to g_N , a_N , a_{N+1} and the arc A_N to obtain an arc A_{N+1} in G_{N+1} . Next, apply Lemma 6.1 to g_{N+1} , a_{N+1} , a_{N+2} and the arc A_{N+1} to obtain an arc A_{N+2} in G_{N+2} . Continue in the same way. Finally, for each $n \geq N$ we have an arc A_n with an end-vertex equal to a_n , such that $g_n|_{A_{n+1}}$ is monotone and onto A_n . Then the arc obtained as the inverse limit of $\{A_n, g_n|_{A_{n+1}}\}$ has \mathbf{a} as one of the end-vertices and the second end-vertex is within ε from \mathbf{b} .

Corollary 6.3. *The continuum $|\mathbb{G}|$ has all arc-components dense.*

\square

In [9] it was shown that the topological realization of the projective Fraïssé limit of the class of finite linear graphs, which are called arcs in this article, with epimorphisms which were not necessarily confluent, is the pseudo-arc. Our next result shows, somewhat surprisingly, that $|\mathbb{G}|$ contains a pseudo-arc.

Lemma 6.4. *Let $\{I_n, \beta_n\}$, where β_n 's are epimorphisms (not necessarily confluent) and I_n 's are arcs, be an inverse sequence with the following property. For every arc J , $k > 0$, and monotone epimorphism $g: J \rightarrow I_k$, there is $l > k$ and an epimorphism (not necessarily confluent) $f: I_l \rightarrow J$ with $g \circ f = \beta_k^l$. Then the inverse limit of $\{I_n, \beta_n\}$ can be embedded in \mathbb{G} .*

Proof. Let $\{F_n, \alpha_n\}$ be a Fraïssé sequence for \mathcal{G} and $\{I_n, \beta_n\}$ be the inverse sequence of finite arcs. We may assume that $F_1 = I_1$. Let $J_1 = J_2 = F_1$ and take $f_1: I_1 \rightarrow J_1$ and $g_1: J_2 \rightarrow I_1$ to be identity maps. Take $n_1 = m_1 = 1 = n_2$.

Suppose that we have $n_1 < \dots < n_{k+1}$, $m_1 < \dots < m_k$, and we have epimorphisms $f_i: I_{m_i} \rightarrow J_i$ (not necessarily confluent) and monotone epimorphisms $g_i: J_{i+1} \rightarrow I_{m_i}$, for $i = 1, 2, \dots, k$. We also have arcs $J_i \subseteq F_{n_i}$ such that $g_{i-1} \circ f_i = \beta_{m_{i-1}}^{m_i}$ and $f_i \circ g_i = \alpha_{n_i}^{n_{i+1}}|_{J_{i+1}}$, for $i = 1, \dots, k$.

We now obtain $n_{k+1} < n_{k+2}$, $m_k < m_{k+1}$, $f_{k+1}, g_{k+1}, J_{k+2}$. Using the property of $\{I_n, \beta_n\}$ we find m_{k+1} and f_{k+1} such that $f_{k+1}: I_{m_{k+1}} \rightarrow J_{k+1} \subseteq F_{n_{k+1}}$ and $g_k \circ f_{k+1} = \beta_{m_k}^{m_{k+1}}$.

We claim that we can extend the epimorphism $f_{k+1}: I_{m_{k+1}} \rightarrow J_{k+1}$, which may be not confluent, to a confluent epimorphism $F: A \rightarrow F_{n_{k+1}}$, for some connected graph A . We let $V(A)$ to be the disjoint union of $V(I_{m_{k+1}})$ and $V(F_{n_{k+1}})$. Let

$$E(A) = \{\langle x, y \rangle : \langle x, y \rangle \in E(F_{n_{k+1}}) \cup E(I_{m_{k+1}}) \text{ or} \\ (x \in V(I_{m_{k+1}}), y \in V(F_{n_{k+1}}), \langle f_{k+1}(x), y \rangle \in E(F_{n_{k+1}}))\}^S.$$

The map F defined by $F|_{I_{m_{k+1}}} = f_{k+1}$ and is the identity otherwise is a confluent epimorphism extending f_{k+1} .

Finally use that $\{F_n, \alpha_n\}$ is the Fraïssé sequence to find n_{k+2} and $G: F_{n_{k+2}} \rightarrow A$ such that $F \circ G = \alpha_{n_{k+1}}^{n_{k+2}}$. Let $J_{k+2} \subseteq F_{n_{k+2}}$ be an arc such that $G(J_{k+2}) = I_{m_{k+1}}$ and $G|_{J_{k+2}}$ is monotone. We let $g_{k+1} = G|_{J_{k+2}}$.

Note that the inverse limit of $\{J_n, \alpha_n|_{J_{n+1}}\}$ is isomorphic to the inverse limit of $\{I_n, \beta_n\}$. Since the inverse limit of $\{F_{n_i}, \alpha_{n_i}^{n_{i+1}}\}$ is isomorphic to \mathbb{G} , we get the required embedding of inverse limit of $\{I_n, \beta_n\}$ into \mathbb{G} . □

Corollary 6.5. *The pseudo-arc can be embedded in $|\mathbb{G}|$.*

Proof. Apply Irwin-Solecki [9], where it is shown that the pseudo-arc is the topological realization of the Fraïssé class of finite arcs. Then form any Fraïssé sequence $\{I_n, \beta_n\}$ for that Fraïssé class. □

We next want to show the following result.

Theorem 6.6. *Each point of the continuum $|\mathbb{G}|$ is the top of the Cantor fan in $|\mathbb{G}|$.*

First we recall from continuum theory.

Definition 6.7. A continuum homeomorphic to the cone over the Cantor set is called the *Cantor fan*, that is, it is homeomorphic to the quotient space $C \times [0, 1] / \sim$ where C is the Cantor set and the points $(x, 1)$ are identified. We will call the point $(x, 1)$ the *top* of the fan. Note that this top point is often called the vertex of the fan but we want to only use the term vertex for elements of graphs. The Cantor

fan X can be characterized as a smooth fan whose set of endpoints is homeomorphic to the Cantor set (see Theorem 1 in [6]).

Definition 6.8. Given $G \in \mathcal{G}$ and a vertex $p \in G$ let

$$\Delta(G, p) = G \times \{0, 1\} / (\{p\} \times \{0, 1\})$$

and $\delta: \Delta(G, p) \rightarrow G$ be just the projection $\delta(x, i) = x$. Observe that δ is a confluent epimorphism.

Proof. (Proof of Theorem 6.6).

Let $\{F_i, \alpha_i\}$ be a Fraïssé sequence for \mathcal{G} . Choose $\mathbf{a} \in \mathbb{G}$ and write $\mathbf{a} = (a_i)$. Let $\pi: \mathbb{G} \rightarrow |\mathbb{G}|$ be the quotient map. We are going to construct the Cantor fan in $|\mathbb{G}|$ with the top $\pi(\mathbf{a})$.

Let K_0 be a graph in \mathcal{G} containing at least two vertices. Using that $\{F_i, \alpha_i\}$ is a Fraïssé sequence, take n_0 and a confluent epimorphism $f_0: F_{n_0} \rightarrow K_0$. Suppose that we have already constructed $f_i: F_{n_i} \rightarrow K_i$ and we have $\delta_i: K_i \rightarrow F_{n_{i-1}}$, for $i < m$, such that $\delta_i \circ f_i = \alpha_{n_{i-1}}^{n_i}$. Let $K_m = \Delta(F_{n_{m-1}}, a_{n_{m-1}})$ and let $\delta_m: K_m \rightarrow F_{n_{m-1}}$ be the projection map of Definition 6.8, which we are subscripting to help keep track of which graph it is operating on. By Proposition 2.8, there is n_m and a confluent epimorphism $f_m: F_{n_m} \rightarrow K_m$ such that $\delta_m \circ f_m = \alpha_{n_{m-1}}^{n_m}$. Let $\beta_i: K_{i+1} \rightarrow K_i$ be $\beta_i = f_i \circ \delta_{i+1}$. See Diagram (D3).

$$(D3) \quad \begin{array}{ccc} F_{n_i} & \xleftarrow{\alpha_{n_i}^{n_{i+1}}} & F_{n_{i+1}} \\ f_i \downarrow & & \downarrow f_{i+1} \\ K_i & \xleftarrow{\beta_i} & K_{i+1} \end{array}$$

All maps in Diagram (D4) are confluent epimorphisms and $\varprojlim \{K_i, \beta_i\}$ is isomorphic to \mathbb{G} .

$$(D4) \quad \begin{array}{ccccccc} & & \beta_0 & & \beta_1 & & \\ & & \curvearrowleft & & \curvearrowleft & & \\ K_0 & \xleftarrow{f_0} & F_{n_0} & \xleftarrow{\delta_0} & K_1 = \Delta(F_{n_0}, a_{n_0}) & \xleftarrow{f_1} & F_{n_1} & \xleftarrow{\delta_1} & K_2 = \dots \\ & & \uparrow & & \uparrow & & \uparrow & & \\ & & & & \alpha_{n_0}^{n_1} & & & & \end{array}$$

For any 0,1-sequence s we use $\text{len}(s)$ to denote the length of the sequence s and, for $n < \text{len}(s)$, $s|_n$ to denote the sequence consisting of the first n entries of s .

Now we are going to construct arcs in the graphs K_n whose inverse limits are arcs in \mathbb{G} . Let A be an arc in K_0 such that $f_0(a_{n_0})$ is an end-vertex of A . Let $A_\emptyset = A$; by Lemma 6.1 there is an arc B_\emptyset in F_{n_0} such that $f_0(B_\emptyset) = A_\emptyset$ and a_{n_0} is an end-vertex of B_\emptyset . By the construction of K_1 there are two arcs A_0 and A_1 in K_1 with $A_0 \cap A_1 = f_1(a_{n_1})$ and $\delta_0(A_0) = \delta_0(A_1) = B_\emptyset$; consequently $\beta_0(A_0) = \beta_0(A_1) = A_\emptyset$. By Lemma 6.1, there are arcs B_k , $k = 0, 1$, in F_{n_1} such that $f_1(B_k) = A_k$ and a_{n_1} is an end-vertex of each B_k .

Using this procedure repeatedly, we may construct, for any finite 0,1-sequence s an arc A_s in $K_{\text{len}(s)}$ such that the following conditions are satisfied:

- (1) $A_\emptyset = A$;
- (2) if $\text{len}(s) = n + 1$, then $A_{s|_n} = \beta_n(A_s)$;
- (3) if $\text{len}(s) = \text{len}(s') = k$ and $s \neq s'$, then $A_s \cap A_{s'} = f_k(a_{n_k})$.

For an infinite 0,1-sequence s , we define $s_n = s|_n$ and $A_s = \varprojlim \{A_{s_n}, \beta_n|_{A_{s_{n+1}}}\}$. Then the union

$$L = \bigcup \{A_s : s \text{ is an infinite 0,1-sequence}\}$$

is, by [7, Corollary 5.16], a smooth dendroid in \mathbb{G} . The vertex \mathbf{a} is the only vertex such that $L \setminus \{\mathbf{a}\}$ has more than 2 components, hence L is a smooth fan. The set of end vertices of L is closed and has no isolated vertices.

It follows that the set $\pi(L)$ is a smooth fan in the topological sense, with the top $\pi(\mathbf{a})$. The set of endpoints is closed and has no isolated points. Thus $\pi(L)$ is homeomorphic to the Cantor fan. \square

7. POINTWISE SELF-HOMEOMORPHIC GRAPHS

Definition 7.1. A topological space X is called *pointwise self-homeomorphic at a point* $x \in X$ if for any neighborhood U of x there is a set V such that $x \in V \subseteq U$ and V is homeomorphic to X . A space is called *pointwise self-homeomorphic* if it is pointwise self-homeomorphic at each of its points. A topological graph is *pointwise self-isomorphic* if in the above we consider topological graphs and instead of requiring a homeomorphism we require a graph isomorphism.

In this section, show that \mathbb{G} is pointwise self-isomorphic and use this to show that $|\mathbb{G}|$ is pointwise self-homeomorphic.

Lemma 7.2. *Suppose U and W are finite connected graphs and $f: W \rightarrow U$ is a confluent epimorphism, and $U \subseteq G$, where G is again a finite connected graph. Then there is a finite connected graph H such that*

$W \subseteq H$ and a confluent epimorphism $f^*: H \rightarrow G$ satisfying $f^*|_W = f$ and $W = (f^*)^{-1}(U)$.

Proof. Take $V(H)$ to be the disjoint union of $V(W)$ and $V(G \setminus U)$ and let

$$E(H) = \{\langle x, y \rangle : \langle x, y \rangle \in E(W) \cup E(G \setminus U) \text{ or} \\ (x \in V(W), y \in V(G \setminus U), \langle f(x), y \rangle \in E(G))\}^S.$$

Let f^* be equal to f on W and to the identity on $G \setminus U$. □

Theorem 7.3. *Let $\mathbf{a} \in \mathbb{G}$ and let U be an open neighborhood of \mathbf{a} . Then there is a topological graph $\mathbb{G}_{\mathbf{a}}$ with $\mathbf{a} \in \mathbb{G}_{\mathbf{a}} \subseteq U$, which is isomorphic with \mathbb{G} .*

Proof. Let $\{F_n, \alpha_n\}$ be a Fraïssé sequence for \mathcal{G} . Write \mathbf{a} as (a_i) with $a_i \in F_i$. Take k such that $(\alpha_k^\infty)^{-1}(a_k) \subseteq U$. For $i \leq k$, take $C_i = \{a_i\}$, and for $i > k$, let C_i be the connected component of $(\alpha_k^i)^{-1}(a_k)$ that contains a_i . By $\bar{\alpha}_m^n$ we denote $\alpha_m^n|_{C_n}$. Using Proposition 2.8, we show that $\{C_n, \bar{\alpha}_n\}$ is a Fraïssé sequence for \mathcal{G} .

For that, pick l and a confluent epimorphism $f: A \rightarrow C_l$, where $A \in \mathcal{G}$. Apply Lemma 7.2 to $U = C_l$, $W = A$, and $G = F_l$. Obtain $H \in \mathcal{G}$ and a confluent epimorphism $f^*: H \rightarrow G$, which extends f and $A = (f^*)^{-1}(C_l)$. Since $\{F_n, \alpha_n\}$ is a Fraïssé sequence for \mathcal{G} , we get n and a confluent epimorphism $g: F_n \rightarrow H$ such that $\alpha_l^n = (f^*)g$. By confluence of g , since A is connected, and using $A = (f^*)^{-1}(C_l)$, we have that the restriction of g to C_n (which we denote by \hat{g}) is onto A and, by Theorem 3.4, it is confluent. Clearly $\bar{\alpha}_l^n = f\hat{g}$.

Let $\mathbb{G}_{\mathbf{a}}$ be the inverse limit of $\{C_n, \bar{\alpha}_n\}$. □

Theorem 7.4. *The Fraïssé limit \mathbb{G} is pointwise self-isomorphic.*

Proof. Let $\mathbf{a} \in \mathbb{G}$ and $\varepsilon > 0$. Then, by condition (3) of Theorem 2.3, there is a graph $G \in \mathcal{G}$ and a confluent epimorphism $f_G: \mathbb{G} \rightarrow G$ such that $\text{diam } f_G^{-1}(f_G(\mathbf{a})) < \varepsilon$. By Theorem 7.3 the component of $f_G^{-1}(f_G(\mathbf{a}))$ that contains \mathbf{a} is a subgraph of \mathbb{G} isomorphic to \mathbb{G} . □

Observation 7.5. *If G is a pointwise self-isomorphic graph with a transitive edge relation, then its topological realization $|G|$ is a pointwise self-homeomorphic compactum.*

Corollary 7.6. *The continuum $|\mathbb{G}|$ is a pointwise self-homeomorphic.*

8. HEREDITARILY UNICOHERENCE

The goal of this section is to show that the graph \mathbb{G} and, consequently, its topological realization are hereditarily unicoherent (see Definition 2.18).

We start with a definition and then we prove a general theorem on unicoherence of Fraïssé limits.

Definition 8.1. Given a topological graph G a quadruple $\langle H, K, C, D \rangle$ is called a *cycle division in G* if the following conditions are satisfied:

- (1) H and K are closed connected subsets of $V(G)$;
- (2) C and D are nonempty subsets of $V(G)$ which are closed in $H \cup K$;
- (3) $H \cap K = C \cup D$;
- (4) $C \cap D = \emptyset$;
- (5) if $c \in C$ and $d \in D$ then $\langle c, d \rangle \notin E(G)$,

Note that Condition (4) follows from Condition (5).

In what follows, if $\langle H, K, C, D \rangle$ is a cycle division in G , we will sometimes instead say that G contains a cycle division $\langle H, K, C, D \rangle$.

Observation 8.2. *If a topological graph G contains a cycle division then G is not hereditarily unicoherent. If there is a cycle division $\langle H, K, C, D \rangle$ such that $H \cup K = V(G)$ then G is not unicoherent.*

Definition 8.3. If G and F are topological graphs containing cycle divisions $C(G) = \langle H_G, K_G, C_G, D_G \rangle$ and $C(F) = \langle H_F, K_F, C_F, D_F \rangle$ respectively, and f is a mapping from G to F , then f is said to *map $C(G)$ onto $C(F)$* if $f(H_G) = H_F$, $f(K_G) = K_F$, $f(C_G) = C_F$, and $f(D_G) = D_F$.

Observation 8.4. *Let $f: \mathbb{G} \rightarrow F$ be a confluent epimorphism and let $\langle H, K, C, D \rangle$ be a cycle division in \mathbb{G} such that if $c \in f(C)$ and $d \in f(D)$ then $\langle c, d \rangle \notin E(F)$, and $f(H) \cap f(K) = f(H \cap K)$. Then $\langle f(H), f(K), f(C), f(D) \rangle$ is a cycle division in F .*

Indeed, this suffices because we always have $f(C \cup D) = f(C) \cup f(D)$, and therefore $f(C) \cup f(D) = f(C \cup D) = f(H \cap K) = f(H) \cap f(K)$.

This observation leads us to the following lemma.

Lemma 8.5. *Let $\langle H, K, C, D \rangle$ be a cycle division in \mathbb{G} and let*

$$\{U_x: x \in K \setminus (C \cup D)\} \cup \{U_y: y \in H \setminus (C \cup D)\} \cup \{V_1, V_2, \mathbb{G} \setminus (K \cup H)\}$$

by an open cover of \mathbb{G} where for $x \in K \setminus (C \cup D)$ take U_x to be a neighborhood of x disjoint from H , for $y \in H \setminus (C \cup D)$ take U_y to be a neighborhood of y disjoint from K and V_1 and V_2 are open sets

containing C and D such that $(V_1 \times V_2) \cap E(\mathbb{G}) = \emptyset$. If $f: \mathbb{G} \rightarrow F$ is a confluent epimorphism that refines this open cover (i.e. for every $a \in F$, $f^{-1}(a)$ is contained in some set from the cover), then $\langle f(H), f(K), f(C), f(D) \rangle$ is a cycle division in F .

Moreover, if $g: \mathbb{G} \rightarrow G$ and $h: G \rightarrow F$ are confluent epimorphisms such that $h \circ g = f$, then $\langle g(H), g(K), g(C), g(D) \rangle$ is a cycle division in G .

In the following lemma we do not require the epimorphisms to be confluent.

Lemma 8.6. *Suppose \mathcal{F} is a Fraïssé class of graphs such that for each graph $F \in \mathcal{F}$ and for each cycle division $\langle H, K, C, D \rangle$ in F (respectively, such that $H \cup K = V(F)$) there is a graph $G \in \mathcal{F}$ such that no cycle division in G is mapped onto $\langle H, K, C, D \rangle$. Then the projective Fraïssé limit \mathbb{F} of \mathcal{F} is hereditarily uncoherent (respectively, uncoherent).*

Proof. Suppose \mathbb{F} is not hereditarily uncoherent (uncoherent). Then there is a cycle division $\langle H, K, C, D \rangle$ in \mathbb{F} (such that $V(H) \cup V(K) = V(\mathbb{F})$). Let F be a graph in \mathcal{F} such that the epimorphism $f_F: \mathbb{F} \rightarrow F$ is such that $\langle f_F(H), f_F(K), f_F(C), f_F(D) \rangle$ is a cycle division in F . Take G as a graph satisfying the assumption and let $f_F^G: G \rightarrow F$ be an epimorphism. Let $f_G: \mathbb{F} \rightarrow G$ be an epimorphism such that $f_F^G \circ f_G = f_F$. Then $\langle f_G(H), f_G(K), f_G(C), f_G(D) \rangle$ is a cycle division in G that f_F^G maps onto $\langle f_F(H), f_F(K), f_F(C), f_F(D) \rangle$, contrary to the definition of G . \square

Theorem 8.7. *\mathbb{G} is hereditarily uncoherent.*

Proof. Suppose that \mathbb{G} is not hereditarily uncoherent. Then there is a cycle division $\langle H, K, C, D \rangle$ in \mathbb{G} . Let F be a graph in \mathcal{G} and $f_F: \mathbb{G} \rightarrow F$ be an epimorphism such that $\langle f_F(H), f_F(K), f_F(C), f_F(D) \rangle$ is a cycle division in F .

Let G be the graph where

$$V(G) = \{(v, i) : v \in V(F) \text{ and } i \in \{0, 1\}\}.$$

Let

$$E_{i,j} = \{\langle (x, i), (y, j) \rangle : x \in f_F(K) \setminus f_F(C), y \in f_F(C) \text{ and } \langle x, y \rangle \in E(F)\},$$

for $i, j \in \{0, 1\}$. Let

$$E(G) = (\{\langle (x, i), (y, i) \rangle : i \in \{0, 1\}, \langle x, y \rangle \in E(F)\} \cup E_{0,1} \cup E_{1,0}) \setminus (E_{0,0} \cup E_{1,1}).$$

For simplicity, let us denote $K_i = f_F(K) \times \{i\}$, $i = \{0, 1\}$, viewed as graphs induced from G , and we similarly define $H_i, C_i, D_i, K_i \setminus C_i$ and

let $F_i = (K_i \setminus C_i) \cup H_i$. Clearly K_0 is isomorphic to K_1 , and the same for the other pairs of graphs.

We claim that the graph G is connected. Since H_i are connected, $D_i \subseteq H_i, K_i \setminus C_i$, and there is at least one edge joining F_1 and F_2 , it suffices to show that for any $k, l \in K_0 \setminus C_0$ there is a path in G from k to l . (Note that K_0 and $K_0 \setminus C_0$ are graphs in G and may not be connected.) Since $K \subseteq F$ is connected take a path P from k to l in K . In the case that $P \cap C = \emptyset$, this is the required path. Otherwise take c, c' such that c is the first element in $P \cap C$ and c' is the last element in $P \cap C$. It may be the case that $c = c'$. If

$$P = \{p_1 = k, p_2, \dots, p_{n_1}, c, \dots, c', p_{n_2}, \dots, p_m = l\},$$

then the required path from k to l in G is

$$P' = \{k, p_2 \times \{0\}, \dots, p_{n_1} \times \{0\}, c \times \{1\}, h_1, h_2, \dots, h_n, c' \times \{1\}, p_{n_2} \times \{0\}, \dots, l\},$$

where $Q = \{c \times \{1\}, h_1, h_2, \dots, h_n, c' \times \{1\}\}$ is a path in H_1 joining $c \times \{1\}$ and $c' \times \{1\}$.

The projection $\alpha: G \rightarrow F$ given by $\alpha((x, i)) = x$ is an epimorphism. In particular, it is confluent.

There exists an epimorphism $f_G: \mathbb{G} \rightarrow G$ such that $\alpha \circ f_G = f_F$. Then $\langle f_G(H), f_G(K), f_G(C), f_G(D) \rangle$ is a cycle division in G (see Lemma 8.5). Clearly, $f_G(H) \subseteq f_F(H) \times \{0, 1\}$ and $\alpha(f_G(H)) = f_F(H)$; and similarly for K, C, D .

Since $f_G(H)$ is connected, and there is no edge in G between $f_F(H) \times \{0\}$ and $f_F(H) \times \{1\}$, as well as $\alpha(f_G(H)) = f_F(H)$, we have $f_G(H) = f_F(H) \times \{0\}$ or $f_G(H) = f_F(H) \times \{1\}$. Without loss of generality, suppose that $f_G(H) = f_F(H) \times \{0\}$ holds. This implies that $f_G(C) = f_F(C) \times \{0\}$ and $f_G(D) = f_F(D) \times \{0\}$. Therefore $(f_F(C) \cup f_F(D)) \times \{0\} \subseteq f_G(K) \subseteq f_F(K) \times \{0, 1\}$. However, there is no edge between the set $((f_F(K) \setminus f_F(C)) \times \{0\}) \cup (f_F(C) \times \{1\})$ (this set contains $f_F(D) \times \{0\}$) and the set $((f_F(K) \setminus f_F(C)) \times \{1\}) \cup (f_F(C) \times \{0\})$ (this set contains $f_F(C) \times \{0\}$). This gives a contradiction with connectedness of $f_G(K)$. □

Theorem 8.8. $|\mathbb{G}|$ is hereditarily unicoherent.

Proof. Suppose $|\mathbb{G}|$ is not hereditarily unicoherent. Then there exist subcontinua H and K in $|\mathbb{G}|$ such that $H \cap K$ is disconnected. So $H \cap K = C \cup D$ where $C \neq \emptyset \neq D$ and $\overline{C} \cap D = \emptyset = C \cap \overline{D}$.

We claim that $\langle \pi^{-1}(H), \pi^{-1}(K), \pi^{-1}(C), \pi^{-1}(D) \rangle$ is a cycle division in \mathbb{G} . To see this, note that the sets H and K are closed since the quotient map π is continuous. The set $\pi^{-1}(H)$ is connected otherwise,

the image of the sets, under π , that form the disconnection of $\pi^{-1}(H)$ would disconnect H . Likewise, $\pi^{-1}(K)$ is connected. Finally, $\pi^{-1}(H) \cap \pi^{-1}(K) = \pi^{-1}(C) \cup \pi^{-1}(D)$ and, since π maps the vertices of a non-degenerate edge to a single point, there is no edge between $\pi^{-1}(C)$ and $\pi^{-1}(D)$. \square

9. NON-HOMOGENEITY

A topological graph A is said to be *homogeneous* if for any $x, y \in V(A)$ there is an isomorphism $f: A \rightarrow A$, i.e. an automorphism, such that $f(x) = y$. Likewise, a continuum C is said to be *homogeneous* if for any $x, y \in C$ there is a homeomorphism $f: C \rightarrow C$ such that $f(x) = y$.

The goal of this section is to prove the following theorem.

Theorem 9.1. *The continuum $|\mathbb{G}|$ is not homogeneous.*

We will accomplish that by showing that there are points in $|\mathbb{G}|$, which belong to a solenoid, as well as there are points in $|\mathbb{G}|$, which do not belong to a solenoid.

9.1. Solenoids, graph-solenoids, and almost graph-solenoids.

A *solenoid* is the continuum $\varprojlim \{S^1, z^{n_i}\}$ where S^1 is viewed as the unit circle in the complex plane. Hagopian showed in [8] that a solenoid can be characterized as a homogeneous continuum where every proper non-degenerate subcontinuum is an arc.

For $A \in \mathcal{G}$ we will say that $C \subseteq A$ is a *cycle* in A if $|V(C)| > 2$ and we can enumerate the vertices of C as $(c_0, c_1, \dots, c_n = c_0)$ in a way that $c_i \neq c_j$ whenever $0 \leq i < j < n$ and $\langle c_i, c_j \rangle \in E(A)$ if and only if $|j - i| \leq 1$ or $\{i, j\} \subseteq \{0, n\}$. A *path* in A is a sequence $p_0, p_1, p_2, \dots, p_n$ of vertices in A such that $n > 0$ and for every $0 \leq i < n$ it holds $\langle p_i, p_{i+1} \rangle \in E(A)$ and $p_i \neq p_{i+1}$.

Note that if $f: D \rightarrow C$ is a confluent epimorphism between cycles then for every $a \in C$ and every component A of $f^{-1}(a)$, if $x, y \in D$, $x \neq y$, are adjacent to A , then $f(x) \neq f(y)$. Confluent epimorphisms between cycles will often be referred to as *wrapping maps*. The *winding number* of a wrapping map f is n if for every (equivalently: some) $c \in C$, $f^{-1}(c)$ has exactly n components.

We call the inverse limit of an inverse sequence of cycles $\{C_n, p_n\}$, where p_n are confluent epimorphisms, a *graph-solenoid* if for infinitely many n the winding number of p_n is greater than 1 and for every $x \in V(C_n)$ every component of $p_n^{-1}(x)$ contains at least 2 vertices.

Let \mathcal{C} be the class of all cycles with confluent epimorphisms.

Lemma 9.2. *The class \mathcal{C} is a projective Fraïssé class.*

Proof. Conditions (1) and (2) of Definition 2.1 are clear. For condition (3), given cycles B and C having m and n vertices, $m \geq n$, let D be a cycle with m vertices and f, g be the obvious epimorphisms.

For condition (4), given cycles A, B , and C with $A = (a_0, a_1, \dots, a_k = a_0)$ and confluent epimorphisms $f: B \rightarrow A$ and $g: C \rightarrow A$, let N be the maximum of the cardinalities of the components of $f^{-1}(a_i)$ and of $g^{-1}(a_i)$, $0 \leq i \leq k$, and let W_f and W_g be the winding numbers for f and g . Let $D = (d_0, d_1, \dots, d_M)$ be a cycle where $M = N \cdot W_f \cdot W_g \cdot k$. Define $f_0: D \rightarrow B$ and $g_0: D \rightarrow C$ as follows. For $0 \leq i \leq k$ and $1 \leq j \leq W_f$, let B_i^j be a component of $f^{-1}(a_i)$ such that B_i^j and B_{i+1}^j are adjacent, B_k^j and B_0^{j+1} are adjacent and $B_k^{W_f}$ and B_0^1 are adjacent. Similarly define C_i^j as components of $g^{-1}(a_i)$. Since $|B_0^1|$ and $|C_0^1|$ have cardinality less than or equal to N there are a monotone epimorphisms f_0 and g_0 from the arc $[d_0, d_{N-1}]$ onto B_0^1 and C_0^1 respectively such that $f_0(d_{N-1})$ and $g_0(d_{N-1})$ are adjacent to B_1^1 and C_1^1 respectively. Next define $f_0: [d_N, d_{2N-1}] \rightarrow B_1^1$ and $g_0: [d_N, d_{2N-1}] \rightarrow C_1^1$ to be monotone epimorphisms such that $f_0(d_N)$ and $g_0(d_N)$ are adjacent to B_0^1 and C_0^1 respectively. Then $f_0(d_{2N-1})$ and $g_0(d_{2N-1})$ will be adjacent to B_2^1 and C_2^1 respectively. Continue this process to obtain the required confluent epimorphisms f_0 and g_0 from D onto B and C . \square

A necessary and sufficient condition for \mathcal{D} to have a topological realization (transitive set of edges) is the following.

(\ddagger) for every $C \in \mathcal{D}$ there is $f: D \rightarrow C$ in \mathcal{D} such that for every $x \in C$ every component of $f^{-1}(x)$ has at least two vertices;

The sufficiency follows from Theorem 2.18 in [7] and one can easily check that this condition is also necessary.

Lemma 9.3. *Let $\{C_n, g_n\}$ be an inverse sequence of cycles with confluent epimorphisms which satisfies (\ddagger) and for every k, N there is $l > k$ such that the winding number of g_k^l is divisible by N . Then $\{C_n, g_n\}$ is a Fraïssé sequence for the class \mathcal{C} .*

Proof. We check the condition in the statement of Proposition 2.8. Let $f: T \rightarrow C_m$, where T is a cycle, be a confluent epimorphism. Let W_f be the winding number for f and let N be the maximum cardinality of the components of $f^{-1}(x)$, $x \in C_m$. By hypothesis, we can find an n such that W_f divides the winding number of g_m^n and for every $x \in C_m$ the minimum cardinality of the components of $(g_m^n)^{-1}(x)$ is at least N . Then similarly as in Lemma 9.3 we can find a confluent epimorphism $h: C_n \rightarrow T$ such that $fh = g_m^n$.

□

In the following we consider possibly other projective Fraïssé classes of cycles.

Example 9.4. Let p_1, p_2, \dots, p_k be prime numbers and let \mathcal{D} be the class of cycles having an even number of vertices and with confluent epimorphisms whose winding numbers are of the form $p_1^{n_1} p_2^{n_2} \dots p_k^{n_k}$, where $n_1, n_2, \dots, n_k \in \mathbb{N}$. Then \mathcal{D} is a projective Fraïssé class.

Let \mathcal{D} be a projective Fraïssé class of cycles with confluent epimorphisms. Note that \mathcal{D} need not contain all possible cycles or all possible confluent epimorphisms. Denote the projective Fraïssé limit of \mathcal{D} by \mathbb{D} . Let $W_{\mathcal{D}}$ be the set of winding numbers of epimorphisms in \mathcal{D} . Therefore, by the amalgamation property, if $m, n \in W_{\mathcal{D}}$, then there is $k \in W_{\mathcal{D}}$, which is divisible by both m and n such that also $\frac{k}{n}, \frac{k}{m} \in W_{\mathcal{D}}$. Moreover, since epimorphisms are closed under compositions, $W_{\mathcal{D}}$ is closed under taking products, i.e. it is a multiplicative semigroup, and since the identity maps are in \mathbb{D} , we have $1 \in W_{\mathcal{D}}$.

Now let W be a multiplicative semigroup of positive integers containing 1 such that if $m, n \in W$, then there is $k \in W$, which is divisible by both m and n , and $\frac{k}{n}, \frac{k}{m} \in W$. Let \mathcal{D}_W be the class that consists of all cycles and all confluent epimorphisms such that winding numbers are in W . Therefore \mathcal{D}_W is the maximal (with respect to containment) class of confluent epimorphisms such that $W_{\mathcal{D}_W} = W_{\mathcal{D}}$. Then \mathcal{D}_W is a projective Fraïssé class. Furthermore we have the following.

Lemma 9.5. *Let \mathcal{D} be a projective Fraïssé class of cycles with confluent epimorphisms which satisfies (\ddagger) . Then the projective Fraïssé limit of $\mathcal{D}_{W_{\mathcal{D}}}$ is isomorphic to the projective Fraïssé limit of \mathcal{D} .*

Proof. Let $\{D_n, \alpha_n\}$ be a Fraïssé sequence for \mathcal{D} and let $\{F_n, \beta_n\}$ be a Fraïssé sequence for $\mathcal{D}_{W_{\mathcal{D}}}$. We have to show that there are subsequences (n_k) and (m_k) and confluent epimorphisms $f_k: F_{n_k} \rightarrow D_{m_k}$ and $g_n: D_{m_{k+1}} \rightarrow F_{n_k}$ such that $\alpha_{m_k}^{m_{k+1}} = f_n \circ g_n$ and $\beta_{n_k}^{n_{k+1}} = g_n \circ f_{n+1}$.

Since $D_1 \in \mathcal{D}_{W_{\mathcal{D}}}$ there exist n and an epimorphism $f_1: F_n \rightarrow D_1$. Take $n_1 = n$ and $m_1 = 1$. Note that F_{n_1} and f_1 may not be in \mathcal{D} . From property (2) for Fraïssé sequences there exists $m > m_1$ such that the winding number of $\alpha_{m_1}^m$ is a multiple of the winding number of f_1 . By (\ddagger) there is $m_2 \geq m$ such that for any $x \in D_1$ the cardinality of any component of $(\alpha_{m_1}^{m_2})^{-1}(x)$ is greater than the cardinality of any component of $f_1^{-1}(x)$. In the same manner as in the proof of Lemma 9.3 we may construct a confluent epimorphism $g_1: D_{m_2} \rightarrow F_{n_1}$ such that $f_1 \circ g_1 = \alpha_{m_1}^{m_2}$. Continuing this process we find the required sequence of epimorphisms.

□

Proposition 9.6. *Let \mathcal{D} be a projective Fraïssé class of cycles with confluent epimorphisms, which satisfies (\ddagger) and $W_{\mathcal{D}}$ is infinite. Then the topological realization $|\mathbb{D}|$ exists and is a solenoid.*

We first need a lemma.

Lemma 9.7. *Let \mathcal{D} be a projective Fraïssé class of cycles with confluent epimorphisms, which satisfies (\ddagger) and let $a_0, a_1 \in \mathbb{D}$ be such that $\langle a_0, a_1 \rangle \in E(\mathbb{D})$. Then the topological graph \mathbb{D}_0 obtained from \mathbb{D} by identifying a_0 and a_1 is isomorphic to the projective Fraïssé limit \mathbb{D} .*

Proof. Without loss of generality, \mathcal{D} is of the form \mathcal{D}_W , as in Lemma 9.5. We check that conditions (1), (2), (3) from Theorem 2.3 hold for \mathbb{D}_0 using that they do hold for \mathbb{D} .

Given $X \in \mathcal{D}$, let $X_0 \in \mathcal{D}$ be such that $|V(X_0)| = |V(X)| + 1$. There exists a confluent epimorphism $f: \mathbb{D} \rightarrow X_0$. If $f(a_0) \neq f(a_1)$ let $p: X_0 \rightarrow X$ be a confluent epimorphism such that $p(f(a_0)) = p(f(a_1))$ and let $f_0: \mathbb{D}_0 \rightarrow X$ be defined by $f_0(x) = p(f(a_0))$ if $x \in f^{-1}(f(a_0)) \cup f^{-1}(f(a_1))$ and $f_0(x) = p(f(x))$ otherwise. If $f(a_0) = f(a_1)$ then p can be taken to be any confluent epimorphism from X_0 to X and f_0 is defined the same as previously.

Now let $f_0: \mathbb{D}_0 \rightarrow C$ and $g: D \rightarrow C$ be confluent epimorphisms, where C, D are cycles in \mathcal{D} . Let a be the vertex in \mathbb{D}_0 that is the identification of a_0 and a_1 . Define $\rho: \mathbb{D} \rightarrow \mathbb{D}_0$ by $\rho(a_0) = \rho(a_1) = a$ and $\rho = \text{id}$ otherwise. Clearly ρ is a confluent epimorphism and let $f = f_0 \circ \rho$. By property (2) for \mathbb{D} there is a confluent epimorphism $w: \mathbb{D} \rightarrow D$ such that $g \circ w = f$. If $w(a_0) = w(a_1)$ then $w': \mathbb{D}_0 \rightarrow D$ defined as $w'(a) = w(a_0) = w(a_1)$ and $w' = w$ otherwise satisfies $g \circ w' = f_0$.

In the case $w(a_0) \neq w(a_1)$ take cycle E such that $|V(E)| = 2|V(D)|$ and $h: E \rightarrow D$ let be a wrapping map that has the winding number equal to 1 and preimage of each vertex is of cardinality 2. By property (2) for \mathbb{D} we get $p: \mathbb{D} \rightarrow E$ such that $f = (gh) \circ p$. Let E_0 be equal to E with $p(a_0)$ and $p(a_1)$ identified and let $\rho_1: E \rightarrow E_0$ be the map that identifies $p(a_0)$ and $p(a_1)$. Let $p_0: \mathbb{D}_0 \rightarrow E_0$ be the quotient of p . Then we have $p_0 \circ \rho = \rho_1 \circ p$. Denote $t = \rho_1(p(a_0))$ and let $h_0: E_0 \rightarrow D$ be equal to h outside t , and let $h_0(t) = h(p(a_0))$. Then the map $p_1 = h_0 \circ p_0: \mathbb{D}_0 \rightarrow D$ is as required, that is, $g \circ p_1 = f_0$.

Given $\epsilon > 0$ there is $C \in \mathcal{D}$ and a confluent epimorphism $f: \mathbb{D} \rightarrow C$ such that for all $x \in C$ the $\text{diam}(f^{-1}(x)) < \epsilon/2$. If $f(a_0) \neq f(a_1)$ let C_0 be the cycle obtained from C by identifying $f(a_0)$ and $f(a_1)$; otherwise

let $C_0 = C$. Let $f_0: \mathbb{D}_0 \rightarrow C_0$ be the quotient of f . Then f_0 is an ϵ -map. □

Proof of Proposition 9.6. By the result of Hagopian we have to show that the topological realization is homogeneous and that every proper non-degenerate subcontinuum is an arc.

Let $\{C_n, g_n\}$ be a Fraïssé sequence for \mathcal{D} . Let $K \subseteq |\mathbb{D}|$ be a proper subcontinuum and consider $\pi_D^{-1}(K)$, where $\pi_D: \mathbb{D} \rightarrow |\mathbb{D}|$ is the quotient map. This is a proper topological subgraph of \mathbb{D} , which is connected. Since any proper connected subgraph of a cycle is an arc, in fact $\pi_D^{-1}(K)$ is the inverse limit of $\{I_n, g_n|I_{n+1}\}$, where I_n is an arc from some point on. Note also that if I_n is an arc, then $g_n|I_{n+1}$ is monotone, and hence $\pi_D^{-1}(K)$ is an arc. Moreover, π_D takes endpoints of $\pi_D^{-1}(K)$ to distinct points. Therefore $K = \pi_D(\pi_D^{-1}(K))$ is an arc.

Let $A, B \in |\mathbb{D}|$. Applying twice Lemma 9.7, we can assume that $|\pi_D^{-1}(A)| = 1$ and $|\pi_D^{-1}(B)| = 1$. Let $a \in \mathbb{D}$ be the only point such that $\pi_D(a) = A$ and similarly let $b \in \mathbb{D}$ be the only point such that $\pi_D(b) = B$. We show that there is an automorphism of \mathbb{D} , which takes a to b . This automorphism will induce a homeomorphism of $|\mathbb{D}|$ that takes A to B . It suffices to show the following.

Claim: Let $a = (a_n) \in \mathbb{D}$, i.e. $a_n \in C_n$, be such that $|\pi_D^{-1}(\pi_D(a))| = 1$. Let m , a cycle T , and a confluent epimorphism $f: T \rightarrow C_m$, be given. Take $t \in T$ with $f(t) = a_m$. Then there is $n > m$ and $h: C_n \rightarrow T$ such that $h(a_n) = t$ and $fh = g_m^n$.

Given the Claim, we can find sequences (n_i) and (m_j) and confluent epimorphisms $f_i: C_{n_i} \rightarrow C_{m_i}$ and $h_i: C_{m_{i+1}} \rightarrow C_{n_i}$ such that $g_{n_i}^{n_{i+1}} = h_i f_{i+1}$ and $g_{m_i}^{m_{i+1}} = f_i h_i$ and $f_i(a_{n_i}) = b_{m_i}$ and $h_i(b_{m_{i+1}}) = a_{n_i}$. These induces an automorphism of \mathbb{D} that takes a to b . □

Proof of Claim. Let I_n for $n \geq m$ be the component of $(g_m^n)^{-1}(a_m)$ containing a_n . Since $|\pi_D^{-1}(\pi_D(a))| = 1$ there is n such that a_n is not an end-vertex of I_n . Therefore for any k there is n_k such that the cardinality of each sub-arc of I_{n_k} between a_{n_k} and an end-vertex of I_{n_k} is at least k . Hence for large enough n there is a monotone epimorphism from I_n onto J which takes a_n to t , where J is the component of $f^{-1}(a_m)$ containing t . Using this last property we find the required h similarly as in Lemma 9.3. □

Solenoids may be topological realizations of topological graphs other than graph-solenoids. We will also need generalizations of graph-solenoids, see Proposition 9.16.

A surjective homomorphism $g: D \rightarrow C$, where C, D are cycles with $C = (c_0, c_1, c_2, \dots, c_k = c_0)$ is an *almost wrapping map* if there is a confluent epimorphism $f: D \rightarrow C$ such that for every $y \in D$ and $x = c_i \in C$, if $f(y) = c_i$ then $g(y) \in \{c_i, c_{i+1}\}$. We call f a *confluent witness* for g . Clearly a confluent witness is not unique. If for every $x \in V(C)$ every component of $f^{-1}(x)$ contains at least 2 vertices, then we call f a *proper confluent witness* for g . A *winding number* of g is the winding number of f . We call the inverse limit of an inverse sequence of cycles $\{C_n, p_n\}$, where p_n are almost wrapping maps, an *almost graph-solenoid* if for infinitely many n the winding number of p_n is greater than 1 and every p_n has a proper confluent witness.

Whenever we consider almost wrapping maps between cycles, we will assume that these cycles are oriented and that almost wrapping maps are orientation preserving. If $C = (c_0, c_1, c_2, \dots, c_k = c_0)$, then we let the *orientation* $<_C$ on C to be the binary relation $<_C = \{(c_i, c_{i+1}) : i = 0, \dots, k-1\}$, and we write $c_i <_C c_{i+1}$. An almost wrapping map $g: D \rightarrow C$, where C, D are cycles, is *orientation preserving* if for some (equivalently any) confluent witness $f: D \rightarrow C$ of g we have: for every $x, y \in D$, if $x <_D y$, then $f(x) <_C f(y)$ or $f(x) = f(y)$. For a cycle C and $a, b \in C$, an *oriented arc*, which we simply denote by $[a, b]$, is the proper subset of C , $\{a = x_0, x_1, \dots, x_m = b\}$ such that for each $j = 0, 1, \dots, m-1$, it holds $x_j <_C x_{j+1}$. We allow that $a = b$ and in that case $[a, b] = \{a\}$. Note that unless $a = b$, then $[a, b] \neq [b, a]$. We also let $(a, b) = [a, b] \setminus \{a, b\}$, and similarly define $[a, b)$ and $(a, b]$.

The following remark is an immediate consequence of the definition of an almost wrapping map.

Remark 9.8. Let $w: D \rightarrow C$ be an almost wrapping map. Then for any $x \in C$ we can partition D into a sequence of arcs: $A_i = [a_i, b_i]$, $B_i = [c_i, d_i]$ with b_i adjacent to c_i and d_i adjacent to a_{i+1} such that for every i , $w(A_i) \subseteq \{x^-, x, x^+\}$, where $x^- \neq x^+$ are adjacent to x , $w(a_i) = w(b_i) = x$, and $w(B_i) = C \setminus \{x\}$.

Remark 9.9. Let $w: D \rightarrow C$ be an almost wrapping map. It is possible that for $x, y \in D$ we have $x <_D y$ and $w(y) <_C w(x)$. However, there are no x, y, z with $x <_D y$, $y <_D z$, $x \neq z$ such that $w(y) <_C w(x)$ and $w(z) <_C w(y)$.

We also have the following description of almost wrapping maps. It characterizes almost wrapping maps as those surjective homomorphisms, which do not reverse the order of triples.

Lemma 9.10. *Let $w: D \rightarrow C$ be a surjective homomorphism, where C, D are cycles. Then w is an almost wrapping map if and only if there*

are no non-adjacent $x \neq z \in C$ and no $a, c \in D$ such that $w(c) = x$, $w(a) = z$, and $w([a, c]) = [x, z]$.

Proof. Suppose first that $w: D \rightarrow C$ is an almost wrapping map. Write $C = (c_0, c_1, \dots, c_n = c_0)$ and let $f: D \rightarrow C$ be a confluent epimorphism such that for every $y \in D$ and $c_l \in C$, if $f(y) = c_l$ then $w(y) \in \{c_l, c_{l+1}\}$, i.e. if $w(y) = c_l$, then $f(y) \in \{c_{l-1}, c_l\}$. Assume there are non-adjacent $x = c_i \neq z = c_j \in C$ and $a, c \in D$ such that $w(c) = x$, $w(a) = z$, and $w([a, c]) = [x, z]$. Then we have $f(c) \in \{c_{i-1}, c_i\}$, $f(a) \in \{c_{j-1}, c_j\}$, and $f([a, c])$ is one of $[c_{i-1}, c_{j-1}]$, $[c_{i-1}, c_j]$, $[c_i, c_{j-1}]$, $[c_i, c_j]$. As f is order preserving, $f(a) \in \{c_{i-1}, c_i\}$ and $f(c) \in \{c_{j-1}, c_j\}$. However, this is impossible since c_i and c_j are non-adjacent.

To prove the converse, suppose that there are no non-adjacent $x \neq z \in C$ and no $a, c \in D$ such that $w(c) = x$, $w(a) = z$, and $w([a, c]) = [x, z]$, and write $C = (c_1, \dots, c_n = c_1)$. Then there are no $x, y, z \in C$ with $x <_C y$ and $y <_C z$, and $a = b_0, b_1, \dots, c = b_m$ with $b_j <_C b_{j+1}$ such that $w(c) = x$, $w(b_j) = y$ for $j = 1, \dots, m-1$, and $w(a) = z$. This implies that we can partition D (in a possibly non-unique way) into arcs $A_i = [a_i, b_i]$, $1 \leq i \leq kn$, for some k , with b_i adjacent to a_{i+1} , and b_{kn} adjacent to a_1 , such that $w(A_i) \in \{c_{(i \bmod n)}, c_{(i+1 \bmod n)}\}$. Therefore f such that $f(A_i) = \{c_i\}$ is a confluent witness for w . \square

Almost wrapping maps are closed under taking compositions.

Lemma 9.11. *Let C, D, E be cycles and let $g_1: D \rightarrow C$ and $g_2: E \rightarrow D$ be almost wrapping maps, which have proper confluent witnesses. Then $g = g_1 \circ g_2$ is an almost wrapping map, which has a proper confluent witness.*

Proof. Write $D = (d_0, \dots, d_n = d_0)$ and $C = (c_0, \dots, c_m = c_0)$. Let $f_1: D \rightarrow C$ and $f_2: E \rightarrow D$ be proper confluent witnesses for g_1 and g_2 , and consider the confluent epimorphism $f_0 = f_1 \circ f_2$. Note first that for $e \in E$ if $f_2(e) = d_i$ and $f_1(d_i) = c_j$, then $g_2(e) \in \{d_i, d_{i+1}\}$, $g_1(d_i) \in \{c_j, c_{j+1}\}$ and $g_1(d_{i+1}) \in \{c_{j+1}, c_{j+2}\}$, hence $g(e) \in \{c_j, c_{j+1}, c_{j+2}\}$. Therefore we have to modify f_0 to get a proper confluent witness for g .

For each j and a component $Y = [y, z]$ of $f_0^{-1}(c_j)$, let first $X = [w, x]$ denote the component of $f_0^{-1}(c_{j-1})$ adjacent to Y . If there is $s \in X$ such that $g(s) = c_{j+1}$, let s_X be the first such vertex, that is, $g(s_X) = c_{j+1}$ and there is no $s' \in [w, s_X)$ such that $g(s') = c_{j+1}$. If there is no $s \in X$ such that $g(s) = c_{j+1}$, let $s_X = y$. Analogously, if there is $s \in Y$ such that $g(s) = c_{j+2}$ let s_Y be the first such vertex in Y , otherwise let s_Y be the vertex not in Y that is adjacent to z .

Define a map from E to C by declaring that f assumes value c_j on $[s_X, s_Y)$. To see that f is a confluent epimorphism, it suffices to see that $s_Y \neq y$, since then $[s_X, s_Y) \neq \emptyset$. We show that in fact $s_Y \neq y^+$, where y^+ is such that $y <_E y^+$, which will imply that for every $a \in V(C)$ every component of $f^{-1}(a)$ contains at least 2 vertices. We have that $Y = [y, z]$ is a component of $f_0^{-1}(c_j)$. Let $Y_1 = [y_1, z_1]$ be the component of $f_1^{-1}(c_j)$ such that $f_2(Y) = Y_1$ and let y_1^+ be such that $y_1 <_D y_1^+$. Then $g_2(y) \in \{y_1, y_1^+\}$ and since f_2 is a proper confluent witness for g_2 , we have $f_2(y^+) = f_2(y) = y_1$, and therefore $g_2(y^+) \in \{y_1, y_1^+\}$. Since f_1 is a proper confluent witness for g_1 , we have $f_1(y_1^+) = f_1(y_1) = c_j$, and therefore $g_1(y_1), g_1(y_1^+) \in \{c_j, c_{j+1}\}$. This implies that $g(y), g(y^+) \in \{c_j, c_{j+1}\}$, and hence $s_Y \neq y, y^+$.

Finally, we check that for any $t \in [s_X, s_Y)$, $g(t) \in \{c_j, c_{j+1}\}$. We have that f_0 assumes value c_j on Y and c_{j-1} on X . Therefore g can only assume values from $\{c_j, c_{j+1}, c_{j+2}\}$ on Y and values from $\{c_{j-1}, c_j, c_{j+1}\}$ on X . Since $[s_X, s_Y) \subseteq X \cup Y$, and by the choice of s_Y there is no $t \in [y, s_Y)$ such that $g(t) = c_{j+2}$, we only have to show that there is no $t \in [s_X, y)$ such that $g(t) = c_{j-1}$. If $s_X = y$ there is nothing to show. In the case $s_X \neq y$, suppose towards the contradiction that there is $t \in [s_X, x)$ such that $g(t) = c_{j-1}$. Apply Lemma 9.10 to the almost wrapping map g_1 , to c_{j-1}, c_{j+1} , and $g_2(s_X), g_2(t)$. This gives a contradiction. \square

Let \mathbb{F} be a topological graph such that for every $x \in \mathbb{F}$ there is at most one $y \neq x$, $y \in \mathbb{F}$ such that $\langle x, y \rangle \in E(\mathbb{F})$, and let $\delta: \mathbb{F} \rightarrow |\mathbb{F}|$ be the quotient map. Let

$$O_{\mathbb{F}} = \{x \in \mathbb{F}: \text{there is no } y \neq x \text{ with } \langle x, y \rangle \in E(\mathbb{F})\}$$

be the set of *singletons* of \mathbb{F} and let

$$T_{\mathbb{F}} = \{x \in \mathbb{F}: \text{there is exactly one } y \neq x \text{ with } \langle x, y \rangle \in E(\mathbb{F})\}$$

be the set of *doubletons* of \mathbb{F} . We have $\mathbb{F} = O_{\mathbb{F}} \cup T_{\mathbb{F}}$.

The set $O_{\mathbb{F}}$ is G_{δ} , i.e. it is the intersection of countably many open sets. Indeed, it is the intersection of

$$A_n = \{x \in \mathbb{F}: d(x, y) < \frac{1}{n} \text{ for every } y \text{ such that } \langle x, y \rangle \in E(\mathbb{F})\},$$

where d is some fixed metric on \mathbb{F} . Each $\mathbb{F} \setminus A_n$ is closed since the edge relation $E(\mathbb{F})$ is closed in \mathbb{F}^2 and \mathbb{F} is compact.

Note that if $\mathbb{F}_1 \subseteq \mathbb{F}_2$ are topological graphs then $\mathbb{F}_1 \cap O_{\mathbb{F}_2} \subseteq O_{\mathbb{F}_1}$, but the equality may not hold.

Remark 9.12. The set $O_{\mathbb{G}}$, where \mathbb{G} is the Fraïssé limit of graphs with confluent epimorphisms, is a dense G_{δ} in \mathbb{G} .

Lemma 9.13. *Let \mathbb{F}_0 be a topological graph such that $\mathbb{F}_0 = O_{\mathbb{F}_0} \cup T_{\mathbb{F}_0}$ and let $\delta: \mathbb{F}_0 \rightarrow |\mathbb{F}_0|$ be the quotient map. Then there is a non-empty open set $U \subseteq \mathbb{F}_0$ such that if $\mathbb{F} = \mathbb{F}_0 \setminus U$ then $O_{\mathbb{F}}$ is dense in \mathbb{F} and $\delta(\mathbb{F}) = |\mathbb{F}_0|$.*

Proof. Let U_1, U_2, U_3, \dots be a countable open base for \mathbb{F}_0 with $U_i \neq \emptyset$. We will proceed by the following recursive procedure. At step $n-1$ we have constructed a closed subset $\mathbb{F}_{n-1} \subseteq \mathbb{F}_0$. If for every $x \in \mathbb{F}_{n-1} \cap U_n$ there is $y \in \mathbb{F}_{n-1} \setminus U_n$ such that $\langle x, y \rangle \in E(\mathbb{F}_{n-1})$, then take $\mathbb{F}_n = \mathbb{F}_{n-1} \setminus U_n$. Otherwise, let $\mathbb{F}_n = \mathbb{F}_{n-1}$. Clearly $\delta(\mathbb{F}_n) = |\mathbb{F}_0|$.

We claim that the set $\mathbb{F} = \bigcap_n \mathbb{F}_n$ is as required. Clearly $\delta(\mathbb{F}) = |\mathbb{F}_0|$. Suppose towards a contradiction that $O_{\mathbb{F}}$ is not dense in \mathbb{F} . Let U be an open nonempty set in \mathbb{F} such that for every $x \in U$ there is $y \in \mathbb{F}$, $y \neq x$, so that $\langle x, y \rangle \in E(\mathbb{F})$. Fix a metric d on \mathbb{F} and let

$$D_n = \{x \in U : \text{there is } y \in \mathbb{F} \text{ with } \langle x, y \rangle \in E(\mathbb{F}) \text{ and } d(x, y) \geq \frac{1}{n}\}.$$

Each D_n is closed in U and we have $\bigcup_n D_n = U$. By the Baire category theorem for the open set U there is n_0 such that D_{n_0} has a nonempty interior. Take $V \subseteq \text{int}(D_{n_0})$ open nonempty of diameter $< \frac{1}{n}$. Therefore, for every $x \in V$, there is $y \in \mathbb{F} \setminus V$ with $\langle x, y \rangle \in E(\mathbb{F})$. Let n be such that $U_n \cap \mathbb{F} \subseteq V$, and without loss of generality by shrinking V , we have $U_n \cap \mathbb{F} = V$. This means that U_n was subtracted in the construction of \mathbb{F} , which gives a contradiction. Indeed, if U_n was not subtracted at stage n , then one of the following would have to happen. The first possibility is that there is $x \in U_n \cap O_{\mathbb{F}_{n-1}}$. However, then $x \in \mathbb{F} \setminus V$, which is impossible. The second possibility is that there are $x \neq y \in U_n$ with $\langle x, y \rangle \in E(\mathbb{F}_{n-1})$. However, then $x, y \notin V$, hence also $x, y \notin \mathbb{F}$, which cannot be the case as $\delta(\mathbb{F}) = |\mathbb{F}_0|$. \square

Example 9.14. Let \mathbb{F} be a topological graph such that $\mathbb{F} = O_{\mathbb{F}} \cup T_{\mathbb{F}}$ and let $|\mathbb{F}|$ be its topological realization. Suppose that $O_{\mathbb{F}}$ is uncountable and dense in \mathbb{F} . Then there is $\mathbb{F}_1 \supseteq \mathbb{F}$ with the same topological realization, $\mathbb{F}_1 = O_{\mathbb{F}_1} \cup T_{\mathbb{F}_1}$, but $O_{\mathbb{F}_1}$ is not dense in \mathbb{F}_1 .

Indeed, $O_{\mathbb{F}}$ is an uncountable dense G_{δ} in \mathbb{F} . Therefore it contains a copy C of the Cantor set. Let D be another copy of the Cantor set and let $h: C \rightarrow D$ be a homeomorphism. We take a topological graph \mathbb{F}_1 obtained as follows. Take $V(\mathbb{F}_1) = V(\mathbb{F}) \cup D$ and let $\langle x, y \rangle \in E(\mathbb{F}_1)$ iff $x = y$ or $(x, y \in \mathbb{F} \text{ and } \langle x, y \rangle \in E(\mathbb{F}))$ or $(x \in D, y \in C, \text{ and } x = h(y))$ or $(y \in D, x \in C, \text{ and } y = h(x))$. We declare D to be a clopen in \mathbb{F}_1 . Then \mathbb{F}_1 is as required as $D \subseteq T_{\mathbb{F}_1}$.

For the moment, see Section 9.2, assume that $|\mathbb{G}|$ contains a solenoid S . Write S as the inverse limit of $\{X_n, w_n\}$, where X_n is the unit circle S^1 and $w_n: X_{n+1} \rightarrow X_n$ is given by $w_n(e^{it}) = e^{il_n t}$, for some $l_n \in \mathbb{N} \setminus \{0, 1\}$. Further, suppose that S is a topological realization of a topological graph $\mathbb{S} \subseteq \mathbb{G}$ such that $O_{\mathbb{S}}$ is dense in \mathbb{S} . Recall that $\pi: \mathbb{G} \rightarrow |\mathbb{G}|$ is the quotient map and denote $\pi_S = \pi|_{\mathbb{S}}$.

For $A \subseteq \mathbb{S}$ we let $k(A)$ to be the closure in \mathbb{S} of the set of the singletons of A , i.e. $k(A) = \overline{O_A}$. We will apply this operation to sets of the form $(w_n^\infty \pi_S)^{-1}(P)$, where P is an arc $[a = e^{it_1}, b = e^{it_2}]$ (respectively an open arc (a, b)) in S^1 . Denote $(w_n^\infty \pi_S)^{-1}(P)$ by $[a, b]_n$ (respectively $(a, b)_n$).

- Lemma 9.15.** (1) *If $A = [a, b]_n$, then $k(A) = \overline{(a, b)_n}$. In that case $k(A)$ is a regular closed set.*
(2) *If $[c, d] \subseteq [a, b]$ and $n \in \mathbb{N}$, then $k([c, d]_n) \subseteq k([a, b]_n)$.*
(3) *Suppose that $[a_1, b_1], [a_2, b_2], \dots, [a_n, b_n]$ is a cover of S^1 by arcs such that $b_i = a_{i+1}$, $i = 1, \dots, n-1$ and $b_n = a_1$ and the interiors of the arcs are disjoint. Then for every $n \in \mathbb{N}$, $k([a_1, b_1]_n), k([a_2, b_2]_n), \dots, k([a_n, b_n]_n)$ is a cover of \mathbb{S} .*

Proof. (1) Since $O_{[a, b]_n} \cap (a, b)_n$ is dense in $(a, b)_n$, we get $\overline{(a, b)_n} \subseteq k(A)$. To obtain the reverse inclusion we additionally have to show that any vertex in $x \in O_{[a, b]_n} \setminus (a, b)_n$ is a limit of vertices in $(a, b)_n$. For that note that $(w_n^\infty)^{-1}((a, b))$ is dense in $(w_n^\infty)^{-1}([a, b])$. Therefore $\pi_S(x)$ is a limit of vertices (z_n) in $(w_n^\infty)^{-1}((a, b))$. Let $y_n \in (a, b)_n$ be such that $\pi_S(y_n) = z_n$. After passing to a subsequence we can assume that (y_n) is convergent to some $y \in [a, b]_n$. Since $\pi_S(x) = \pi_S(y)$ and $x \in O_{[a, b]_n}$, we obtain $x = y$.

(2) It follows immediately from (1) since $(c, d)_n \subseteq (a, b)_n$.

(3) Let $x \in \mathbb{S}$ and let (x_n) be a sequence of vertices in $O_{\mathbb{S}}$ converging to x . Then by the continuity of π_S and w_n^∞ , we have that $(w_n^\infty \pi_S(x_n))$ converges to $w_n^\infty \pi_S(x)$. Passing to a subsequence, we have that (from some point on) all $(w_n^\infty \pi_S(x_n))$ lie in a single $[a_{i_0}, b_{i_0}]$. Then $x \in k([a_{i_0}, b_{i_0}]_n)$. \square

Proposition 9.16. *The topological graph \mathbb{S} is an almost graph-solenoid.*

Proof. Let S be the solenoid $\varprojlim \{X_n, w_n\}$, where $X_n = S^1$ and $w_n: X_{n+1} \rightarrow X_n$ is given by $w_n(e^{it}) = e^{il_n t}$, for some $l_n \in \mathbb{N} \setminus \{0, 1\}$. Let \mathcal{P}_1 be the cover of X_1 :

$$\{[e^0, e^{i2\frac{\pi}{5}}], [e^{i2\frac{\pi}{5}}, e^{i2\frac{2\pi}{5}}], \dots, [e^{i2\frac{4\pi}{5}}, e^{i2\frac{5\pi}{5}}]\},$$

and for every $n > 1$ let \mathcal{P}_n be the cover of X_n :

$$\{[e^{i\frac{m2\pi}{L_n}}, e^{i\frac{(m+1)2\pi}{L_n}}]: m = 0, 1, \dots, L_n - 1\},$$

where $L_n = 5 \times 2^{n-1} \times l_1 l_2 \dots l_{n-1}$. Let \mathcal{D}_n be the cover of S defined as

$$\mathcal{D}_n = \{(w_n^\infty)^{-1}(P) : P \in \mathcal{P}_n\}.$$

For every n consider the cover $\mathcal{E}_n = \{k(E) : E \in \pi_S^{-1}(\mathcal{D}_n)\}$ of \mathbb{S} by closed sets. The graph C_n such that $V(C_n) = \mathcal{E}_n$ and $\langle E, F \rangle \in E(C_n)$ iff there is an edge between a vertex in E and a vertex in F , is a cycle. Moreover, whenever $m < n$, the cover \mathcal{E}_n refines \mathcal{E}_m and this refinement induces a confluent epimorphism from C_n onto C_m .

Additionally, for every $\epsilon > 0$ there is N such that for every $n > N$ and every $E \in \mathcal{E}_n$, $\text{diam}(E) < \epsilon$. If $x \in S$ then x may be written as $x = (e^{it_1^x}, e^{it_2^x}, \dots)$. Let d_S denote the metric on S defined as $d_S(x, y) = \sum_{n=1}^{\infty} |t_n^x - t_n^y|/2^n$ and let d denote a metric on \mathbb{S} . Note that $\pi_S|_{O_S}$ is a homeomorphism onto its image. Therefore there is M such that for any $x, y \in O_S$ we have $d(x, y) \leq M d_S(\pi_S(x), \pi_S(y))$. By the definition of d_S there is N such that for every $n > N$ and every $D \in \mathcal{D}_n$, $\text{diam}(D) < \frac{\epsilon}{M}$. Then $\text{diam}(O_{\pi_S^{-1}(D)}) < \epsilon$. Since a set and its closure have the same diameter, we have $\text{diam}(k(\pi_S^{-1}(D))) < \epsilon$.

Now we enlarge the sets in \mathcal{E}_n to obtain a cover of clopen sets $\mathcal{U}_n = \{U_E : E \in \mathcal{E}_n\}$. We require that $E \subseteq U_E$ and for every n and $U_E, U_F \in \mathcal{U}_n$, there is an edge between a vertex in U_E and a vertex in U_F iff $\langle E, F \rangle \in E(C_n)$. Moreover, let $\text{diam}(U_E) < 2\text{diam}(E)$, and let whenever $m < n$, the cover \mathcal{U}_n refine \mathcal{U}_m .

Enumerate \mathcal{U}_n into U_1, U_2, \dots, U_{k_n} so that for every i , there is an edge between a vertex in U_i and a vertex in U_{i+1} (we let $U_{k_n+1} = U_1$). We now let $V_i = U_i \setminus U_{i+1}$ and $\mathcal{V}_n = \{V_i : i = 1, \dots, k_n\}$. Then \mathcal{V}_n is a clopen partition of \mathbb{S} .

There is an edge between a vertex in V_i and a vertex in V_{i+1} , for each i . Indeed, since \mathbb{S} is connected, the only other possibility would be that there is a single i_0 such that there is no edge between a vertex in V_{i_0} and a vertex in V_{i_0+1} . However, this is impossible, since the same would have to be true for the cover \mathcal{V}_{n+1} . However, the winding number of the confluent epimorphism obtained from the inclusion of \mathcal{E}_{n+1} into \mathcal{E}_n is at least 2.

Since for every $\epsilon > 0$ there is N such that for every $n > N$ and every $U \in \mathcal{U}_n$, $\text{diam}(U) < \epsilon$, using the Lebesgue covering lemma, we pass to a subsequence of covers (\mathcal{V}_n) so that for each n , \mathcal{V}_{n+1} refines \mathcal{V}_n . This gives a surjective homomorphism $g_n : C_{n+1} \rightarrow C_n$. If $f_n : C_{n+1} \rightarrow C_n$ is the confluent epimorphism obtained from the inclusion of \mathcal{E}_{n+1} into \mathcal{E}_n , then for every $y \in C_{n+1}$ and $x \in C_n$, if $f_n(y) = c_i$, then $g_n(y) \in \{c_i, c_{i+1}\}$, where $C_n = (c_0, c_1, \dots, c_{k_n} = c_0)$. \square

Remark 9.17. Proposition 1.4.3 in Basso's PhD thesis [2] implies that there is a graph-solenoid \mathbb{S}_1 , given by the inverse limit of cycles with confluent epimorphisms, whose topological realization is equal to S .

9.2. There are points that belong to a solenoid contained in $|\mathbb{G}|$.
In this section we show the following theorem.

Theorem 9.18. *There is a dense set of points in $|\mathbb{G}|$ that belong to a solenoid.*

We start with several lemmas.

Lemma 9.19. *Let $A, B \in \mathcal{G}$ and let $f: B \rightarrow A$ be a confluent epimorphism. Let $C = (c_0, c_1, \dots, c_n = c_0)$ be a cycle in A . Then there is a cycle $D \subseteq B$ such that $f(D) = C$.*

Proof. Note first that by the confluence of f , for every $x \in B$ and $0 \leq i < n$, if $f(x) = c_i$ then there is a path $x = x_0, x_1, \dots, x_k, z$ in B , for some k , such that for every j , $f(x_j) = c_i$ and $f(z) = c_{i+1}$. Using this simple observation, and that B is a finite set, we obtain that for any $y \in B$ with $f(y) = c_0$ there is a path $y = y_0, y_1, \dots, y_l$ such that: 1) for $i < l$ the y_i 's are pairwise different; 2) there is $i_0 < l$ such that $y_l = y_{i_0}$ and $f(\{y_{i_0}, \dots, y_l\}) = \{c_0, \dots, c_{n-1}\}$; 3) the sequence $f(y_0), f(y_1), \dots, f(y_l)$ is of the form $c_0, \dots, c_{n-1}, c_0, \dots, c_{n-1}, \dots, c_0, \dots, c_t$, for some t . Therefore $D = (y_{i_0}, \dots, y_l)$ is the required cycle. \square

Remark 9.20. Note that it may not be the case that for every $b \in B$ with $f(b) = c_0$ there is a cycle $D \subseteq B$ with $b \in D$ such that $f(D) = C$. Compare with Lemma 6.1.

Lemma 9.21. *For any $A \in \mathcal{G}$, $m > 0$ and cycle C in A there is $B \in \mathcal{G}$, a cycle D in B and a confluent epimorphism $f: B \rightarrow A$ such that $f(D) = C$ and the winding number of $f|_D$ equals m .*

Proof. Consider m disjoint copies of A : A_1, \dots, A_m . Pick any $x, y \in V(C)$ with $\langle x, y \rangle \in E(A)$. The copy of x in A_i we call x_i , the copy of y in A_i we call y_i , and the copy of C in A_i we call C_i . From the disjoint union $A_1 \cup A_2 \cup \dots \cup A_m$ remove edges: $\langle x_1, y_1 \rangle, \langle x_2, y_2 \rangle, \dots, \langle x_m, y_m \rangle$. Add edges: $\langle x_1, y_2 \rangle, \langle x_2, y_3 \rangle, \langle x_3, y_4 \rangle, \dots, \langle x_m, y_1 \rangle$. This is the required B . Note that B is connected and that the canonical projection $p: B \rightarrow A$ is a confluent epimorphism. Let D be the finite graph where $V(D) = V(C_1) \cup \dots \cup V(C_m)$ and edges of D being all of the edges in the union of the C_i 's except for the deletions and additions above. \square

Lemma 9.22. *For any $A \in \mathcal{G}$ and $a \in A$ there is $B \in \mathcal{G}$, a confluent epimorphism $f: B \rightarrow A$, and $b \in B$ with $f(b) = a$ such that b belongs to a cycle in B .*

Proof. Let $\langle a, c \rangle$ be a non-degenerate edge in A . To define B let $b \notin V(A)$, $V(B) = V(A) \cup \{b\}$ and $E(B) = E(A) \cup \{\langle b, a \rangle, \langle b, c \rangle\}$. The map that takes b to a , $\langle b, a \rangle$ to $\langle a, a \rangle$ and $\langle b, c \rangle$ to $\langle a, c \rangle$ and is the identity otherwise is confluent and (a, b, c, a) is a cycle in B . \square

Lemma 9.23. *For any $A \in \mathcal{G}$ and cycle C in A there is $B \in \mathcal{G}$, a cycle D in B and a confluent epimorphism $f: B \rightarrow A$ such that $f(D) = C$, the winding number is one, and for every $x \in C$, $|f^{-1}(x) \cap D| = 2$.*

Proof. Write $C = (c_0, c_1, \dots, c_n = c_0)$. Let D be the cycle $(c_0, c'_0, c_1, c'_1, \dots, c_n, c'_n = c_0)$. First take B_0 , the disjoint union of the graphs $A \setminus C$ and D and let $\alpha: D \rightarrow C$ be the map that takes c_i and c'_i to c_i , for every i . We then obtain B from B_0 by adding edges $\langle x, y \rangle$ and $\langle y, x \rangle$ iff $x \in V(A \setminus C)$, $y \in V(D)$, and $\langle x, \alpha(y) \rangle \in E(A)$ \square

We say that $\mathbf{x} \in \mathbb{G}$ belongs to a graph-solenoid (an almost graph-solenoid) if there is a graph-solenoid (an almost graph-solenoid) \mathbb{S} and an embedding $h: \mathbb{S} \rightarrow \mathbb{G}$ such that $\mathbf{x} \in h(\mathbb{S})$.

Fix $\{F_n, \alpha_n\}$ a Fraïssé sequence for the class \mathcal{G} .

Theorem 9.24. *The set of vertices in \mathbb{G} that belong to a graph-solenoid, which is a Fraïssé limit of the class \mathcal{C} of cycles with confluent epimorphisms, is dense.*

Proof. Given any m and $a \in F_m$ apply Lemma 9.22, with $A = F_m$ to obtain $B \in \mathcal{G}$ and a confluent epimorphism $f: B \rightarrow F_m$ where B contains a cycle C such that $a \in f(C)$. From the properties of the Fraïssé sequence, there is $n > m$ and a confluent epimorphism $g: F_n \rightarrow B$ such that $fg = \alpha_m^n$. By Lemma 9.19 there is a cycle $E \subseteq F_n$ which satisfies $g(E) = C$. Let $a_1 \in E$ be such that $fg(a_1) = a$. Write $m_1 = n$ and $C_1 = E$.

Applying Lemmas 9.21, 9.23 and 9.19, construct a sequence $m_1 < m_2 < \dots$ as well as $a_i \in F_{m_i}$, which belongs to a cycle C_i in F_{m_i} , $\alpha_{m_i}^{m_{i+1}}(C_{i+1}) = C_i$ and the winding number of $\alpha_{m_i}^{m_{i+1}}|_{C_{i+1}}$ is a multiple of i . Then the sequence (a_i) induces a vertex $\mathbf{x} \in (\alpha_m^\infty)^{-1}(a) \subseteq \mathbb{G}$ which belongs to a graph-solenoid.

By the construction and Lemma 9.3, the inverse sequence $\{C_i, \alpha_{m_i}^{m_{i+1}}|_{C_{i+1}}\}$ is a Fraïssé sequence for \mathcal{C} . \square

Proof of Theorem 9.18. Apply Theorem 9.24 and Proposition 9.6. \square

9.3. There are points that do not belong to a solenoid contained in $|\mathbb{G}|$.

Theorem 9.25. *There exists a point in $|\mathbb{G}|$ that does not belong to a solenoid.*

We first present a construction in which for an arbitrary finite graph B and an increasing sequence of subsets of B we obtain a confluent epimorphism $g: C \rightarrow B$. Then using this construction we obtain a Fraïssé sequence $\{K_n, \beta_n\}$ for \mathcal{G} together with a vertex $\bar{x} = (x_n)$ with $x_n \in K_n$. Finally, we show that $\pi(\bar{x})$, where $\pi: \mathbb{G} \rightarrow |\mathbb{G}|$ is the quotient map, does not belong to a solenoid.

Construction. Let B be a finite connected graph and let $\emptyset \neq X_0 \subsetneq X_1 \subsetneq \dots \subsetneq X_{n-1} \subsetneq X_n = B$ be an increasing sequence of connected subgraphs of B . Given the sequence (X_i) , we construct a confluent epimorphism g from a connected graph C to B as follows. First the construction of C .

Step 0: Take $Y_0 = X_0$.

Step 1: We construct Y_1 . For every $x \in V(Y_0)$ and $t \in B$ with $\langle x, t \rangle \in E(B)$ take $X_{i+1}^{x,t}$ to be an isomorphic copy of X_{i+1} , where i is such that $t \in V(X_{i+1}) \setminus V(X_i)$. Then Y_1 is the graph Y_0 together with all of the graphs $X_{i+1}^{x,t}$ for $x \in Y_0$ and corresponding vertices t and with edges joining the copy of t in $X_{i+1}^{x,t}$ with $x \in Y_0$.

Step $1 < k < n$: We have already constructed $Y_0 \subseteq Y_1 \subseteq \dots \subseteq Y_k$.

To construct Y_{k+1} note that $Y_k \setminus Y_{k-1}$ is the disjoint union of isomorphic copies of the graphs X_j for various $j \geq k-1$. For a vertex $x \in Y_k \setminus Y_{k-1}$, denote by ψ_x the isomorphism between X_j and the isomorphic copy of X_j that contains x . If $t \in V(B) \setminus V(X_j)$ is such that $\langle \psi_x^{-1}(x), t \rangle \in E(B)$ take $X_{i+1}^{x,t}$ to be an isomorphic copy of the graph X_{i+1} , where i is such that $t \in V(X_{i+1}) \setminus V(X_i)$, together with an edge joining the copy of t in $X_{i+1}^{x,t}$ with x . Then Y_{k+1} is the graph Y_k together with all of the graphs $X_{i+1}^{x,t}$ for $x \in V(Y_k) \setminus V(Y_{k-1})$ and corresponding vertices t . Precisely, let Y_{k+1} be the following graph. Take

$$V(Y_{k+1}) = V(Y_k) \cup \bigcup_{t,x} V(X_{i+1}^{x,t})$$

and

$$E(Y_{k+1}) = E(Y_k) \cup \bigcup_{t,x} E(X_{i+1}^{x,t}) \cup \{\langle x, t \rangle : x \in V(X_j), t \in V(X_{i+1}^{x,t})\}.$$

Finally let $C = Y_n$ and let $g: C \rightarrow B$ be the obvious projection. Then g is a confluent epimorphism.


Call the graph C the *unfolding* of B with respect to the sequence (X_i) . Also, the graph Y_1 will be called the *kernel* of C .


Let A be a connected finite graph and let T be a connected subgraph of A . We say that $T \subseteq A$ is *adjacently disconnecting* A if for every $x \neq y \in V(A \setminus T)$, adjacent to T , any path joining x and y intersects T . In other words, T is adjacently disconnecting A if any homomorphism from a finite arc into A , which maps one end-vertex of the arc to x and the other to y , is such that the intersection of the image of the arc with T is nonempty.

Lemma 9.26. *Let B be a finite connected graph and let $\emptyset \neq X_0 \subsetneq \dots \subsetneq X_{n-1} \subsetneq X_n = B$ be an increasing sequence of connected subgraphs of B , and let $g: C \rightarrow B$ be the epimorphism obtained in the construction above. For every $k = 0, 1, \dots, n$, let C_k denote the component of $g^{-1}(X_k)$, which contains the kernel of C . Then C_k is adjacently disconnecting C .*

Proof. The $V(C)$ is the disjoint union of multiple copies of $V(X_0), V(X_1), \dots, V(X_n)$. For each $k = 0, 1, \dots, n$ the C_k is the induced subgraph of C whose set of vertices is the union of all disjoint copies of $V(X_0), V(X_1), \dots, V(X_k)$, which were attached at some stage of the construction. Finally, note that if we were to identify the vertices in each of the disjoint copies of the $V(X_i)$'s in C to single vertices, then the resulting graph is a tree. \square

Example 9.27. Let B be the graph having vertices a, b, c, d, e and edges $\langle a, b \rangle, \langle a, c \rangle, \langle a, d \rangle, \langle b, c \rangle, \langle c, d \rangle, \langle c, e \rangle, \langle d, e \rangle$. Let X_0 be the subgraph having vertices a, b and the edge $\langle a, b \rangle$, X_1 be the subgraph having vertices a, b, c and edges $\langle a, b \rangle, \langle a, c \rangle, \langle b, c \rangle$, X_2 be the subgraph having vertices a, b, c, d and edges $\langle a, b \rangle, \langle a, c \rangle, \langle a, d \rangle, \langle b, c \rangle, \langle c, d \rangle$, and $X_3 = B$. The graphs Y_0, Y_1 and part of Y_2 are shown below.





Fraïssé sequence. We now construct a Fraïssé sequence $\{K_n, \beta_n\}$ for \mathcal{G} and we construct $\bar{x} = (x_n)$ in its inverse limit (so $x_n \in K_n$), which have the property stated in the proposition below.

Proposition 9.28. *There exists a Fraïssé sequence $\{K_n, \beta_n\}$ for \mathcal{G} and $\bar{x} = (x_n) \in O_{\mathbb{G}}$ with $x_n \in K_n$ such that for every n , $\{x_n\}$ is adjacently disconnecting K_n , and for every $m < n$ the component $C_{m,n}$ of $(\beta_m^n)^{-1}(st(x_m))$ that contains x_n , where*

$$st(x_n) = \{x_n\} \cup \{y \in K_n : y \text{ is adjacent to } x_n\}$$

is adjacently disconnecting K_n .

Proof. Let $\{F_n, \alpha_n\}$ be a Fraïssé sequence for \mathcal{G} , where F_1 is not a singleton. We will construct an inverse sequence $\{K_n, \beta_n\}$ which is also a Fraïssé sequence for \mathcal{G} , and sequences of confluent epimorphisms g_n and f_n . Each K_n is the unfolding of F_{k_n} with respect to a prescribed increasing sequence of connected graphs (X_i) , $g_n: K_n \rightarrow F_{k_n}$ is the mapping from the unfolding construction, $f_n: F_{k_{n+1}} \rightarrow K_n$ is obtained using that $\{F_n, \alpha_n\}$ is a Fraïssé sequence, and $\beta_n = f_n \circ g_{n+1}$. We also construct a vertex $\bar{x} \in \mathbb{G}$ where $\bar{x} = (x_n)$ with $x_n \in K_n$ and $\beta_n(x_{n+1}) = x_n$.

Let z be any vertex in F_1 , K_1 be the unfolding of F_1 with respect to the sequence $(\{z\}, F_1)$, and $g_1: K_1 \rightarrow F_1$ be the mapping from the unfolding construction. Let $x_1 \in K_1$ be the unique vertex in the kernel of K_1 . Create a graph G by replacing each edge $\langle x_1, y \rangle \in E(K_1)$ by the two edges $\langle x_1, y_1 \rangle$ and $\langle y_1, y \rangle$. Then $g: G \rightarrow K_1$ that maps each y_1 to x_1 and is the identity otherwise is a confluent epimorphism. Let $k_1 = 1$ and since $\{F_n, \alpha_n\}$ is a Fraïssé sequence there is a k_2 and a confluent epimorphism $h: F_{k_2} \rightarrow G$ such that $g_1 \circ g \circ h = \alpha_{k_1}^{k_2}$. Let $f_1 = g \circ h$.

Note that for some $v \in f_1^{-1}(x_1)$ any vertex in F_{k_2} that is adjacent to v is mapped by f_1 to x_1 .

Suppose that $\{K_n, \beta_n\}$ and $x_n \in K_n$ with $\beta_{n-1}(x_n) = x_{n-1}$, $n \leq N$, are constructed. We also have an increasing sequence (k_n) , $n \leq N+1$, and maps $f_n: F_{k_{n+1}} \rightarrow K_n$, $g_n: K_n \rightarrow F_{k_n}$, $n \leq N$, with $\beta_n = f_n \circ g_{n+1}$ for $n < N$, and $\alpha_{k_n}^{k_{n+1}} = g_n \circ f_n$ for $n \leq N$. We now construct K_{N+1} , $x_{N+1} \in K_{N+1}$, f_{N+1} , and g_{N+1} with the corresponding properties for $N+1$ instead of N . Then we let $\beta_N = f_N \circ g_{N+1}$.

Let $X_{N+1} = F_{k_{N+1}}$ and pick a vertex $v_{N+1} \in f_N^{-1}(x_N)$ such that all vertices adjacent to v_{N+1} are also mapped to x_N , and denote $X_0 = \{v_{N+1}\}$. Let X_1 be the component of $f_N^{-1}(st(x_N))$ that contains X_0 . Let for $1 \leq i < N$, X_{i+1} be the component of $(\beta_{N-i}^N \circ f_N)^{-1}(st(x_{N-i}))$ that contains X_0 . Let K_{N+1} be the unfolding of $F_{k_{N+1}}$ with respect to the sequence (X_i) , let $g_{N+1}: K_{N+1} \rightarrow F_{k_{N+1}}$ be the resulting epimorphism, and let x_{N+1} be the unique vertex in the kernel of K_{N+1} . Take $\beta_N = f_N \circ g_{N+1}$. Because $\{F_n, \alpha_n\}$ is a Fraïssé sequence there exists k_{N+2} and a confluent epimorphism $f_{N+1}: F_{k_{N+2}} \rightarrow K_{N+1}$ such that $\alpha_{k_{N+1}}^{k_{N+2}} = g_{N+1} \circ f_{N+1}$. Similarly as in the first step we can and we do assume that some vertex in $F_{k_{N+2}}$ mapped to x_{N+1} by f_{N+1} is such that all vertices adjacent to it are also mapped to x_{N+1} .

The $\{K_n, \beta_n\}$ is a Fraïssé sequence, which follows from the fact that $\{F_n, \alpha_n\}$ is a Fraïssé sequence and diagram chasing.

This finishes the construction. Now we check that $\{K_n, \beta_n\}$ and $\bar{x} = (x_n)$ have the required properties.

By the construction, $\beta_n^{n+1}(st(x_{n+1})) = x_n$ for all n , which implies $\bar{x} \in O_{\mathbb{G}}$. Furthermore, for every $m < n$, the $X_{n-m} \subseteq F_{k_n}$ is the component of $(\beta_m^{n-1} \circ f_{n-1})^{-1}(st(x_m))$ that contains $X_0 = \{v_n\} \subseteq F_{k_n}$, $\beta_m^n = \beta_m^{n-1} \circ f_{n-1} \circ g_n$, and $g_n(x_n) = v_n$. Therefore $C_{m,n}$ is the component of $g_n^{-1}(X_{n-m})$ that contains x_n , which by Lemma 9.26 is adjacently disconnecting K_n . Since $\{x_n\} = g_n^{-1}(X_0)$, Lemma 9.26 also implies that $\{x_n\}$ is adjacently disconnecting K_n . □

Proposition 9.29. *Let $\{K_n, \beta_n\}$ and $\bar{x} \in \mathbb{G}$ be as constructed in Proposition 9.28. Then $\bar{x} = (x_n)$ does not belong to an almost graph-solenoid.*

Proof. Let \mathbb{S} be an almost graph-solenoid and write $\mathbb{S} = \varprojlim \{C_i, w_i\}$, where C_i 's are cycles and $w_i: C_{i+1} \rightarrow C_i$ are almost wrapping maps. Towards the contradiction suppose that there is an embedding $\tau: \mathbb{S} \rightarrow \mathbb{G} (= \varprojlim \{K_n, \beta_n\})$ such that $\bar{x} \in \tau(\mathbb{S})$. Therefore $\tau(\mathbb{S}) = \varprojlim \{S_n, \beta_n | S_{n+1}\}$, where $V(S_n) = V(\beta_n^\infty(\tau(\mathbb{S})))$ and $\langle x, y \rangle \in E(S_n)$ iff $\langle x, y \rangle \in E(K_n)$.

As in the proof of Lemma 2.5, we obtain increasing sequences (n_i) and (m_i) and surjections $p_i: S_{n_i} \rightarrow C_{m_i}$, $q_i: C_{m_{i+1}} \rightarrow S_{n_i} \subseteq K_{n_i}$ with $p_i \circ q_i = w_{m_i}^{m_{i+1}}$ and $q_i \circ p_{i+1} = \beta_{n_i}^{m_{i+1}}|_{S_{n_{i+1}}}$.

Since almost wrapping maps are closed under compositions (see Lemma 9.11), $w_{m_i}^{m_{i+1}}$ is an almost wrapping map. To simplify the notation, we assume that $m_i = i$. Without loss of generality, we may assume that C_1 has at least 6 vertices.

Note that each q_i is a homomorphism. Indeed, let $\langle a, b \rangle \in E(C_{i+1})$. Then there is $\langle a', b' \rangle \in E(\mathbb{S})$ with $w_i^\infty(a') = a$ and $w_i^\infty(b') = b$. Then $q_i(a) = \beta_{n_i}^\infty \tau(a')$, $q_i(b) = \beta_{n_i}^\infty \tau(b')$, and $\langle \beta_{n_i}^\infty \tau(a'), \beta_{n_i}^\infty \tau(b') \rangle \in E(K_{n_i})$, as needed.

Let $T = (\{x_{n_1}\} \cup \{y \in K_{n_1} : y \text{ is adjacent to } x_{n_1}\})$, $T_0 = T \cap S_{n_1}$, and $A = (q_1)^{-1}(T) = (q_1)^{-1}(T_0)$.

Claim 1. There is an arc in $C_2 \setminus A$, which consists of at least two vertices.

Proof of Claim 1. We first show that $|w_1(A)| \leq 3$ and any vertex in $w_1(A)$ is joined by an edge with $p_1(x_{n_1})$. Indeed, if $s \in T_0 \setminus \{x_{n_1}\}$, then since $\{x_{n_1}\}$ is adjacently disconnecting K_{n_1} , there are adjacent $a, b \in A$ such that $q_1(a) = s$ and $q_1(b) = x_{n_1}$. Therefore, since w_1 is a homomorphism, $\langle w_1(a) = p_1(s), w_1(b) = p_1(x_{n_1}) \rangle \in E(C_1)$. Since there are exactly two vertices in C_1 adjacent to $p_1(x_{n_1})$, we get $|w_1(A)| \leq 3$.

Suppose towards the contradiction that any arc in $C_2 \setminus A$ is a singleton. Then since w_1 is a homomorphism, any vertex $w_1(y)$ for a $y \in C_2 \setminus A$, is adjacent to $w_1(A)$. But this is impossible since C_1 has at least 6 vertices. □

Denote $\beta = \beta_{n_1}^{n_2}$, let D be the component of $\beta^{-1}(T)$ containing x_{n_2} , and let $E = (q_2)^{-1}(D)$. We have that D is adjacently disconnecting K_{n_2} . Let $\{L_i = [k_i, l_i], J_i = [u_i, v_i], i \leq m\}$ be the partition of C_3 into arcs such that $E = \bigcup_i J_i$, l_i is adjacent to u_i , and v_i is adjacent to k_{i+1} . Write also A as the disjoint union of arcs $M_j = [a_j, b_j]$ in a way that $b_j \in [a_j, a_{j+1}]$.

Let us make several observations.

- (1) For every i , $w_2(J_i) \subseteq A$, in particular, $w_2(u_i), w_2(v_i) \in A$. Indeed, from $\beta \circ q_2 = q_1 \circ w_2$ it follows that $w_2(E) \subseteq A$.
- (2) For every i , $w_2(k_i), w_2(l_i) \notin A$. Indeed, suppose towards the contradiction that say $w_2(l_i) \in A$. That implies by $\beta \circ q_2 = q_1 \circ w_2$ that $\beta \circ q_2(l_i) \in T$, hence $q_2(l_i) \in \beta^{-1}(T)$. By the choice of the partition u_i is adjacent to l_i , therefore since q_2 is a homomorphism, $q_2(u_i)$ is adjacent to $q_2(l_i)$. This along with

$q_2(l_i) \in \beta^{-1}(T)$ and $q_2(u_i) \in D$ implies that $q_2(l_i) \in D$. A contradiction.

- (3) For every i , $w_2(l_i)$ is adjacent to $w_2(u_i)$, and $w_2(v_i)$ is adjacent to $w_2(k_{i+1})$. This simply follows from the fact that w_2 is a homomorphism.
- (4) For every i there is j such that $w_2(J_i) = \{a_j\}$ or $w_2(J_i) = \{b_j\}$ or $w_2(J_i) = M_j$. Moreover, in the case $w_2(J_i) = M_j$ and $|M_j| \geq 2$, we have $a_j = w_2(u_i) \neq w_2(v_i) = b_j$.

It follows from (1), (2) and (3) that there is j such that $w_2(\{u_i, v_i\}) = \{a_j\}$ or $w_2(\{u_i, v_i\}) = \{b_j\}$ or $w_2(\{u_i, v_i\}) = \{a_j, b_j\}$. In the scenario $w_2(\{u_i, v_i\}) = \{a_j, b_j\}$ we get in fact $w_2(J_i) = M_j$, as w_2 is a homomorphism. In the case $w_2(\{u_i, v_i\}) = \{a_j\}$ and $b_j \neq a_j$, then by (2) and (3), we obtain $w_2(l_i) = w_2(k_{i+1})$, and denote $x = w_2(k_{i+1})$. Let $y \in M_j$ be such that $w_2(J_i) = [a_j, y]$, and take $z \in J_i$ with $w_2(z) = y$. Since w_2 is an almost wrapping map, by Lemma 9.10 applied to x, y and z, k_{i+1} , we must have $y = a_j$, and hence $w_2(J_i) = \{a_j\}$. We similarly conclude that $w_2(J_i) = \{b_j\}$ in the case when $w_2(\{u_i, v_i\}) = \{b_j\}$. Moreover if $w_2(u_i) \neq w_2(v_i)$, again by Lemma 9.10, this time applied to a_j, b_j and v_i, u_i , we must have $a_j = w_2(u_i) \neq w_2(v_i) = b_j$ and not $b_j = w_2(u_i) \neq w_2(v_i) = a_j$.

The following claim will finish the proof of the proposition.

Claim 2. There is an arc $I_i = [k_i, l_i]$ in $C_3 \setminus E$ such that $w_2(k_i) \neq w_2(l_i)$.

Indeed, denoting $k_i = k$ and $l_i = l$ for simplicity, if $w_2(k) \neq w_2(l)$, then $q_2(k) \neq q_2(l)$ as $w_2 = p_2 \circ q_2$. As k and l are adjacent to E , $q_2(k)$ and $q_2(l)$ are adjacent to D . Then, since D is adjacently disconnecting K_{n_2} , the image by q_2 of the arc in $C_3 \setminus E$ with end-vertices k and l intersects D , which is a contradiction.

Proof of Claim 2. Suppose towards the contradiction that for every j , $w_2(k_j) = w_2(l_j)$.

Case 1. For every i , we have $w_2(u_i) = w_2(v_i)$.

Then by (4) any vertex in $w_2(E)$ is adjacent to some vertex in $C_2 \setminus A$. Since q_1 is a homomorphism this implies that any vertex in $q_1 \circ w_2(E)$ is adjacent to some vertex in $K_{n_1} \setminus T$. However, as $x_{n_1} \in q_1 \circ w_2(E)$, this is impossible.

Case 2. For some i_0 , we have $w_2(u_{i_0}) \neq w_2(v_{i_0})$.

To simplify the notation let $i_0 = m$, so $w_2(u_m) \neq w_2(v_m)$. Let j be such that $w_2(J_m) = [a_j, b_j]$ and let $m_0 \leq m$ be the least such that

$w_2(J_{m_0}) = w_2(J_m)$. By (4), $a_j = w_2(u_{m_0}) = w_2(u_m) \neq w_2(v_m) = w_2(v_{m_0}) = b_j$.

Consider the sequence:

$$w_2(v_m) = w_2(v_{m_0}), w_2(k_1) = w_2(l_1), w_2(J_1), \dots, w_2(k_i) = w_2(l_i), w_2(J_i), \dots, \\ w_2(k_{m_0-1}) = w_2(l_{m_0-1}), w_2(J_{m_0-1}), w_2(k_{m_0}) = w_2(l_{m_0}), w_2(u_{m_0}).$$

By (3), any two consecutive elements in this sequence are adjacent, therefore it is a path joining $w_2(v_{m_0})$ and $w_2(u_{m_0})$; we denote it by P . The $w_2(J_m)$ contains at least two vertices and P does not contain edges from $w_2(J_m)$. Therefore $C_2 = w_2(J_m) \cup P$, which implies that $w_2(E) = A$ and $C_2 \setminus A$ is the union of singletons $\{w_2(k_i) = w_2(l_i)\}$. But this is a contradiction since there are arcs in the complement of A , which are not singletons.

□

□

Proof of Theorem 9.25. Let $\bar{x} \in \mathbb{G}$ be the vertex constructed in Proposition 9.28, along with the Fraïssé sequence $\{K_n, \beta_n\}$. We claim that $z = \pi(\bar{x})$, where $\pi: \mathbb{G} \rightarrow |\mathbb{G}|$ is the projection map, does not belong to a solenoid. Suppose towards a contradiction that there is a solenoid S such that $z \in S \subseteq |\mathbb{G}|$. By Lemma 9.13 there is a topological graph $\mathbb{S} \subseteq \pi^{-1}(S)$ such that $O_{\mathbb{S}}$ is dense in \mathbb{S} and $\pi(\mathbb{S}) = S$. Since $\bar{x} \in O_{\pi^{-1}(S)}$, we have $\bar{x} \in \mathbb{S}$. By Proposition 9.16, \mathbb{S} is an almost graph-solenoid. On the other hand, by Proposition 9.29, \bar{x} does not belong to an almost graph-solenoid. We obtain a contradiction.

□

10. OPEN PROBLEMS

We have shown many properties of the continuum $|\mathbb{G}|$. We do not know if these properties characterize $|\mathbb{G}|$.

Problem 10.1. Find a topological characterization of $|\mathbb{G}|$.

The following problem is probably quite difficult given the amount of work involved in showing that $|\mathbb{G}|$ is not homogeneous.

Problem 10.2. Investigate the orbits of homogeneity for $|\mathbb{G}|$.

In investigating the homogeneity of $|\mathbb{G}|$ we spent time thinking about the following combinatorial question which we believe is interesting.

Question 10.3. Given a finite connected graph H does there exist a homogeneous finite connected graph G and a confluent epimorphism $f: G \rightarrow H$?

REFERENCES

- [1] D. Bartošová and A. Kwiatkowska, *Lelek fan from a projective Fraïssé limit*, Fund. Math. 231 (2015), no. 1, 57–79.
- [2] G. Basso, *Compact Metrizable Structures via Projective Fraïssé Theory with an Application to the Study of Fences*, PhD thesis, 2020, University of Lausanne and University of Torino.
- [3] G. Basso and R. Camerlo, *Fences, their endpoints, and projective Fraïssé limit*, Trans. Amer. Math. Soc. **374** (2021), 4501–4535.
- [4] J. J. Charatonik, *Confluent mappings and unicoherence of continua*, Fund. Math. 56 (1964), 213–220.
- [5] J. J. Charatonik, *Open Mappings of Universal Dendrites*, Bull. Acad. Polon. Sci., Ser. Sci. Math. 27(1980), 489–494.
- [6] J. J. Charatonik, W. J. Charatonik, *Fans with the property of Kelley*, Topology Appl. 29 (1988), no. 1, 73–78.
- [7] W. J. Charatonik and R. P. Roe, *Projective Fraïssé limits of trees*, preprint.
- [8] C. L. Hagopian, *A characterization of solenoids*, Pac. J. of Math. 68 (1977), no. 2, 425–435.
- [9] T. Irwin and S. Solecki, *Projective Fraïssé Limits and the Pseudo-arc*, Tran. Amer. Math. Soc. **358**, (2006), 3077–3096.
- [10] S. Macías, *Topics on continua*, Second edition, Springer, Cham, 2018, xvii+440 pp.
- [11] S. B. Nadler, Jr. *Dimension theory: an introduction with exercises*, Aportaciones Matemáticas: Textos [Mathematical Contributions: Texts], 18. Sociedad Matemática Mexicana, México, 2002. viii+180 pp. ISBN: 970-32-0026-5.
- [12] A. Panagiotopoulos and S. Solecki, *A combinatorial model for the Menger curve*, accepted to Journal of Topology and Analysis.
- [13] J. T. Rogers Jr. *Homogeneous hereditarily indecomposable continua are tree-like*, Houston J. Math. 8 (1982), no. 3, 421–428.

W. J. CHARATONIK, DEPARTMENT OF MATHEMATICS AND STATISTICS, MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY, 400 W 12TH ST, ROLLA MO 65409-0020

A. KWIATKOWSKA, INSTITUT FÜR MATHEMATISCHE LOGIK UND GRUNDLAGENFORSCHUNG, UNIVERSITÄT MÜNSTER, EINSTEINSTRASSE 62, 48149 MÜNSTER, GERMANY **and** INSTYTUT MATEMATYCZNY, UNIWERSYTET WROCŁAWSKI, PL. GRUNWALDZKI 2/4, 50-384 WROCŁAW, POLAND

Email address: `kwiatkoa@uni-muenster.de`

R. P. ROE, DEPARTMENT OF MATHEMATICS AND STATISTICS, MISSOURI UNIVERSITY OF SCIENCE AND TECHNOLOGY, 400 W 12TH ST, ROLLA MO 65409-0020

Current address: Robert P. Roe, 1407 Athene Dr, Lafayette CO 80026 USA

Email address: `rroe@mst.edu`