

UNCERTAINTY PRINCIPLE VIA VARIATIONAL CALCULUS ON MODULATION SPACES

NUNO C. DIAS, FRANZ LUEF AND JOÃO N. PRATA

ABSTRACT. We approach uncertainty principles of Cowling-Price-Heisenberg-type as a variational principle on modulation spaces. In our discussion we are naturally led to compact localization operators with symbols in modulation spaces. The optimal constant in these uncertainty principles is the smallest eigenvalue of the inverse of a compact localization operator. The Euler-Lagrange equations for the associated functional provide equations for the eigenfunctions of the smallest eigenvalue of these compact localization operators. As a by-product of our proofs we derive a generalization to mixed-norm spaces of an inequality for Wigner and Ambiguity functions due to Lieb.

1. INTRODUCTION

A cornerstone of Fourier analysis is the uncertainty principle about the time and frequency localization of a function. Loosely speaking it states that a function $f(x)$ and its Fourier transform $\widehat{f}(\omega)$ cannot both be sharply localized. One of the first quantitative versions of the uncertainty principle to be put forward was the Heisenberg-Pauli-Weyl inequality:

$$(1.1) \quad \left(\int_{\mathbb{R}} (x - x_0)^2 |f(x)|^2 dx \right)^{1/2} \left(\int_{\mathbb{R}} (\omega - \omega_0)^2 |\widehat{f}(\omega)|^2 d\omega \right)^{1/2} \geq \frac{\|f\|_{L^2(\mathbb{R})}^2}{4\pi},$$

for any $x_0, \omega_0 \in \mathbb{R}$. In what follows, we will set $x_0 = \omega_0 = 0$, as this can be easily achieved by a phase-space translation, which does not alter the results.

This inequality can be shown to be equivalent to [FoSi]:

$$(1.2) \quad \|xf\|_{L^2(\mathbb{R})}^2 + \|\omega\widehat{f}\|_{L^2(\mathbb{R})}^2 \geq \frac{\|f\|_{L^2(\mathbb{R})}^2}{2\pi}.$$

The constant is sharp and we obtain an equality if and only if f is a generalized Gaussian [Gr01].

Cowling and Price [CoPr] have generalized (1.2) by considering other norms and other weights. Here we wish to consider the following result [CoPr, FoSi].

Theorem 1 (Cowling-Price). *Suppose $p, q \in [1, \infty]$ and $a, b > 0$. There is a constant $K > 0$ such that:*

$$(1.3) \quad \| |x|^a f \|_{L^p(\mathbb{R})} + \| |\omega|^b \widehat{f} \|_{L^q(\mathbb{R})} \geq K \|f\|_{L^2(\mathbb{R})},$$

for all tempered distributions f and \widehat{f} , which are represented by locally integrable functions, if and only if:

$$(1.4) \quad a > \frac{1}{2} - \frac{1}{p} \quad \text{and} \quad b > \frac{1}{2} - \frac{1}{q} .$$

In the present work we intend to extend the Cowling-Price uncertainty principle in the following sense. We shall prove, in arbitrary dimension $d \geq 1$, that inequalities of the form

$$(1.5) \quad \| |x|^a f \|_{L^p(\mathbb{R}^d)} + \| |\omega|^b \widehat{f} \|_{L^q(\mathbb{R}^d)} \geq K \| f \|_{M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)} ,$$

hold for all $f \in M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)$ and where the left-hand side may be infinite. Here $M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)$ is a certain modulation space to be specified below. Moreover, we shall prove that there exist functions f_0 for which we obtain an equality and which are the solutions of certain nonlinear functional equations. This latter result is missing in Cowling and Price's original work.

In addition, we shall also prove that, for certain weight functions ψ, ϕ , we have an inequality of the form:

$$(1.6) \quad \| \psi f \|_{L^2(\mathbb{R}^d)}^2 + \| \phi \widehat{f} \|_{L^2(\mathbb{R}^d)}^2 \geq K \| f \|_{M_{m_0}^2(\mathbb{R}^d)} ,$$

where $M_{m_0}^2(\mathbb{R}^d)$ is another modulation space.

Inequalities (1.5,1.6) can be regarded as generalizations of the Heisenberg-Pauli-Weyl inequality (1.2). Moreover, we shall show that we obtain an equality in (1.6) for functions f_0 which are eigenfunctions of a certain positive compact localization operator and that the optimal constant is related with its eigenvalues.

One ingredient of our treatment is the work of Galperin and Gröchenig [Ga, GG, Gr96], where Heisenberg-type inequalities are interpreted as embeddings of weighted Lebesgue spaces $L_a^p(\mathbb{R}^d)$ and weighted Fourier images $\mathcal{FL}_b^q(\mathbb{R}^d)$ of weighted $L_b^q(\mathbb{R}^d)$ spaces into modulation spaces $M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)$.

We denote by $L_a^p(\mathbb{R}^d)$ the space of distributions f in the Lebesgue space $L^p(\mathbb{R}^d)$ with finite $\| \cdot \|_{L_a^p(\mathbb{R}^d)}$, where $\| f \|_{L_a^p(\mathbb{R}^d)} = \left(\int_{\mathbb{R}^d} |f(x)|^p (1+|x|)^{ap} dx \right)^{1/p}$.

The space $\mathcal{FL}_b^q(\mathbb{R}^d)$ consists of all tempered distributions $f \in \mathcal{S}'(\mathbb{R}^d)$ such that \widehat{f} is in $L_b^q(\mathbb{R}^d)$ with finite norm $\| f \|_{\mathcal{FL}_b^q(\mathbb{R}^d)} = \| \widehat{f} \|_{L_b^q(\mathbb{R}^d)}$.

Modulation spaces were introduced by Feichtinger in 1983 [Fe81, Fe06, Fe83B]. Here we restrict the discussion to the following classes of modulation spaces: $M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)$ and $M_m^2(\mathbb{R}^d)$, to be defined below. Loosely speaking, modulation spaces allow one to encode the decay and integrability properties of a function on the phase-space.

There are various ways to measure the phase-space content of a function, e.g. Wigner transform, Rihazcek transform [Co]. We use the matrix coefficients of the Schrödinger representation π of the Heisenberg group $\mathbb{H}(d)$, $\pi(x, \omega)g(t) = e^{2\pi i t \cdot \omega} g(t - x)$:

$$(1.7) \quad V_g f(x, \omega) = (f, \pi(x, \omega)g)_{L^2} = \int_{\mathbb{R}^d} f(t) \overline{g(t-x)} e^{-2\pi i t \cdot \omega} dt,$$

for f, g in $L^2(\mathbb{R}^d)$. In signal analysis $V_g f$ is known as the short-time-Fourier transform. If $g \in \mathcal{S}(\mathbb{R}^d)$, then the short-time Fourier transform extends to tempered distributions $f \in \mathcal{S}'(\mathbb{R}^d)$.

For $\alpha, \beta \in \mathbb{R}$ and $r, s \in [1, 2]$ we define the modulation space $M_{\alpha, \beta}^{r, s}(\mathbb{R}^d)$ as the space of all functions such that

$$(1.8) \quad \|f\|_{M_{\alpha, \beta}^{r, s}(\mathbb{R}^d)} = \left(\int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} |V_g f(x, \omega)|^r (1 + |x|)^{\alpha r} dx \right)^{s/r} (1 + |\omega|)^{\beta s} d\omega \right)^{1/s}$$

is finite with respect to a fixed non-zero g in $\mathcal{S}(\mathbb{R}^d)$. Note that different choices of the non-zero window $g \in \mathcal{S}(\mathbb{R}^d)$ lead to equivalent norms for $M_{\alpha, \beta}^{r, s}(\mathbb{R}^d)$. If we want to emphasize the choice of window g , we may sometimes write $\|\cdot\|_{g, M_{\alpha, \beta}^{r, s}(\mathbb{R}^d)}$.

We shall also consider the modulation spaces with norm

$$(1.9) \quad \|f\|_{M_m^2(\mathbb{R}^d)} = \left(\int_{\mathbb{R}^d} \int_{\mathbb{R}^d} |m(x, \omega) V_g f(x, \omega)|^2 dx d\omega \right)^{1/2},$$

where $m(x, \omega)$ is an appropriate time-frequency weight (see e.g. [AF]).

We need the following particular case of the results in [Ga, GG]:

Theorem 2 (Galperin-Gröchenig). *Let $\alpha, \beta \geq 0$, $0 < r, s \leq 2$ and $1 \leq p, q \leq \infty$. Suppose that $r \leq p$ and $s \leq q$. If*

$$(1.10) \quad \left(\frac{a-\alpha}{d} + \frac{1}{p} - \frac{1}{r} \right) \left(\frac{b-\beta}{d} + \frac{1}{q} - \frac{1}{s} \right) >$$

$$\max \left\{ \left(\frac{1}{r} - \frac{1}{q'} + \frac{\alpha}{d} \right), \left(\frac{1}{r} - \frac{1}{2} + \frac{\alpha}{d} \right) \right\} \times \max \left\{ \left(\frac{1}{s} - \frac{1}{p'} + \frac{\beta}{d} \right), \left(\frac{1}{s} - \frac{1}{2} + \frac{\beta}{d} \right) \right\}$$

with all factors positive, then $L_a^p(\mathbb{R}^d) \cap \mathcal{FL}_b^q(\mathbb{R}^d)$ is compactly embedded in $M_{\alpha, \beta}^{r, s}(\mathbb{R}^d)$. Thus, there exists $C > 0$ such that:

$$(1.11) \quad \|f\|_{L_a^p(\mathbb{R}^d)} + \|\widehat{f}\|_{L_b^q(\mathbb{R}^d)} \geq C \|f\|_{M_{\alpha, \beta}^{r, s}(\mathbb{R}^d)}.$$

As usual p', q', \dots denote the Hölder duals of $p, q, \dots \in [1, \infty]$: $\frac{1}{p} + \frac{1}{p'} = 1$, etc. We also remark that the condition $r' \leq p, p' \leq r$ is equivalent to $\left| \frac{1}{p} - \frac{1}{2} \right| \leq \left| \frac{1}{r} - \frac{1}{2} \right|$.

We will resort to this theorem to prove (Theorem 11) that inequality (1.5) holds if $1 < r, s \leq 2$; $\alpha, \beta \geq 0$; $0 < a, b \leq r'$, and $r \leq p \leq r'$; $s \leq q \leq r'$ and (1.10) is valid.

That this extends the Cowling-Price uncertainty principle is now obvious. If we set $r = s = 2$, $\alpha = \beta = 0$, $d = 1$ in (1.10), we obtain (1.4). Likewise (1.3) follows from (1.5) and the fact that $M_{0,0}^{2,2}(\mathbb{R}) = L^2(\mathbb{R})$ [Gr01].

We shall prove the second inequality (eq.(1.6)) in Theorem 12. For this, we shall require another compact embedding theorem due to Boggiatto and Toft [BoTo] (see also [PfTo]) for modulation spaces of the form $M_m^{p,q}(\mathbb{R}^d)$ (to be defined below in eq.(2.6)):

Theorem 3 (Boggiatto-Toft). *Assume that $m_1, m_2 \in \mathcal{P}(\mathbb{R}^{2d})$, and that $p, q \in [1, \infty]$. Then the embedding*

$$(1.12) \quad i : M_{m_1}^{p,q}(\mathbb{R}^d) \rightarrow M_{m_2}^{p,q}(\mathbb{R}^d)$$

is compact if and only if $m_2/m_1 \in L_0^\infty(\mathbb{R}^{2d})$.

Here $\mathcal{P}(\mathbb{R}^{2d})$ is a class of weights which are polynomially moderate (see Definition 1) and $L_0^\infty(\mathbb{R}^{2d})$ is the set of all $f \in L^\infty(\mathbb{R}^{2d})$ such that

$$(1.13) \quad \lim_{R \rightarrow \infty} \left(\text{ess sup}_{|z| \geq R} |f(z)| \right) = 0.$$

Let us remark that the compactness of such embeddings between modulation spaces also follows from Theorem 9.4 in [FeGr].

The strategy of the proofs consists of approaching this problem from the point of view of variational calculus. We will minimize the functional

$$(1.14) \quad f \mapsto \| |x|^a f \|_{L^p(\mathbb{R}^d)} + \| |\omega|^b \hat{f} \|_{L^q(\mathbb{R}^d)}$$

on the constraint set:

$$(1.15) \quad \Omega = \left\{ f \in L_a^p(\mathbb{R}^d) \cap \mathcal{F}L_b^q(\mathbb{R}^d) : \|f\|_{M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)} = 1 \right\}.$$

In a similar fashion, we will seek to minimize

$$(1.16) \quad f \mapsto \|\psi f\|_{L^2(\mathbb{R}^d)}^2 + \|\phi \hat{f}\|_{L^2(\mathbb{R}^d)}^2,$$

for f in an appropriate modulation space $M_m^2(\mathbb{R}^d)$ and such that $\|f\|_{M_{m_0}^2(\mathbb{R}^d)} = 1$, where $M_{m_0}^2(\mathbb{R}^d)$ is another modulation space, such that $M_m^2(\mathbb{R}^d) \subset M_{m_0}^2(\mathbb{R}^d)$ (see Corollary 3).

These variational problems will naturally lead to Euler-Lagrange equations (Theorem 13 and Theorem 15, Theorem 17).

To prove the existence of minimizers, we start by showing that the functionals (1.14, 1.16) are weakly lower semicontinuous (Lemma 2 and Lemma 4). Then we prove that certain subsets of the constraint sets¹ are weakly sequentially compact (Proposition 3 and Proposition 4). This means that the functionals attain a minimum on these subsets. Finally, we show that they are in fact minima on the entire constraint sets (Theorems 11 and 12).

An essential tool in this last step is a generalization of an inequality for Wigner and radar ambiguity functions due to E. Lieb. In [L1] Lieb proved that

$$(1.17) \quad \|A(f, g)\|_{L^r(\mathbb{R}^2)} \geq C \|f\|_{L^p(\mathbb{R})} \|g\|_{L^{p'}(\mathbb{R})},$$

where $1 \leq r < 2$; $r \leq p, p' \leq r'$, $C > 0$ is a constant and $A(f, g)(x, \omega)$ is the radar ambiguity function:

$$(1.18) \quad A(f, g)(x, \omega) = e^{-i\pi\omega \cdot x} V_g f(-x, \omega).$$

In Theorem 6 and Corollary 1 we shall prove an extension of this inequality to arbitrary dimension $d \geq 1$ and mixed-norm spaces:

$$(1.19) \quad \|A(f, g)\|_{L_{\omega,x}^{r,s}(\mathbb{R}^{2d})} \geq C \|f\|_{L^u(\mathbb{R}^d)} \|g\|_{L^v(\mathbb{R}^d)}$$

for appropriate values of u, v, r, s and where

$$(1.20) \quad \|F\|_{L_{\omega,x}^{r,s}(\mathbb{R}^{2d})} = \left(\int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} |F(x, \omega)|^r d\omega \right)^{\frac{s}{r}} dx \right)^{\frac{1}{s}}.$$

We also determine the sharp constants and the functions which saturate (1.19).

¹Each of these subsets is the intersection of a closed ball and the constraint set.

NOTATION

We denote by $\mathcal{S}(\mathbb{R}^d)$ the space of Schwartz test functions and by $\mathcal{S}'(\mathbb{R}^d)$ its dual - the space of tempered distributions. The duality bracket is $\langle f, g \rangle$ for g in some functional space \mathcal{B} and f in its dual \mathcal{B}' . The inner product in a Hilbert space \mathcal{H} is $(\cdot, \cdot)_{\mathcal{H}}$, which is linear in the first argument. The Fourier transform of $f \in L^2(\mathbb{R}^d)$ is given by

$$(1.21) \quad \widehat{f}(\omega) = \int_{\mathbb{R}^d} f(x) e^{-2\pi i x \cdot \omega} dx$$

where

$$(1.22) \quad x \cdot \omega = x_1 \omega_1 + \cdots + x_d \omega_d,$$

for all $x = (x_1, \dots, x_d)$, $\omega = (\omega_1, \dots, \omega_d) \in \mathbb{R}^d$. We shall also use the notation:

$$(1.23) \quad \langle x \rangle = 1 + |x|.$$

Notice that the notation $\langle x \rangle$ is more commonly used in the literature for the Japanese bracket $\langle x \rangle = (1 + |x|^2)^{1/2}$, which is equivalent to our weight (1.23). We shall nevertheless use the weight (1.23) as it will make our derivations simpler.

If there is a constant $C > 0$ such that $A(f) \leq CB(f)$ for all f in some set, then we write $A(f) \lesssim B(f)$. If $A(f) \lesssim B(f)$ and $B(f) \lesssim A(f)$, then we write $A(f) \asymp B(f)$. If a sequence $(u_n)_n$ in some normed space X converges strongly to some $u \in X$, then we write $u_n \rightarrow u$, and if it converges weakly, then we write $u_n \rightharpoonup u$.

We denote the compact embedding of a functional space \mathcal{A} into \mathcal{B} by $\mathcal{A} \subset\subset \mathcal{B}$.

2. MODULATION SPACES

Definition 1. A weight in \mathbb{R}^d is a positive and continuous function $m \in L_{loc}^\infty(\mathbb{R}^d)$. Given two weights m and v , m is said to be v -moderate, if

$$(2.1) \quad m(x+y) \leq Cm(x)v(y), \quad \forall x, y \in \mathbb{R}^d,$$

for some $C > 0$. We denote by $\mathcal{P}(\mathbb{R}^d)$ the set of all weights m which are v -moderate for some polynomial weight v .

Definition 2. Given a window $g \in \mathcal{S}(\mathbb{R}^d) \setminus \{0\}$, we define the short-time Fourier transform of $f \in \mathcal{S}(\mathbb{R}^d)$ by

$$(2.2) \quad V_g f(x, \omega) = (f, \pi(x, \omega)g)_{L^2(\mathbb{R}^d)} = \int_{\mathbb{R}^d} f(t) \overline{g(t-x)} e^{-2\pi i t \cdot \omega} dt.$$

This extends to $f \in \mathcal{S}'(\mathbb{R}^d)$, if we use the duality bracket:

$$(2.3) \quad V_g f(x, \omega) = \langle f, \pi(x, \omega)\overline{g} \rangle.$$

To study the time-frequency content of a function or tempered distribution, we shall consider the mixed norms:

$$(2.4) \quad \|F\|_{L_{x,\omega}^{r,s}(\mathbb{R}^{2d})} = \left(\int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} |F(x, \omega)|^r dx \right)^{\frac{s}{r}} d\omega \right)^{\frac{1}{s}},$$

and

$$(2.5) \quad \|F\|_{L_{\omega,x}^{r,s}(\mathbb{R}^{2d})} = \left(\int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} |F(x,\omega)|^r d\omega \right)^{\frac{s}{r}} dx \right)^{\frac{1}{s}},$$

for $F \in \mathcal{S}'(\mathbb{R}^{2d})$ and $1 \leq r, s < \infty$, with the obvious modification for r or $s = \infty$.

Given a weight $m \in \mathcal{P}(\mathbb{R}^{2d})$, the modulation space $M_m^{r,s}(\mathbb{R}^d)$ is defined as the set of all $f \in \mathcal{S}'(\mathbb{R}^d)$ such that

$$(2.6) \quad \|f\|_{M_m^{r,s}(\mathbb{R}^d)} = \|mV_g f\|_{L_{x,\omega}^{r,s}(\mathbb{R}^{2d})} = \left(\int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} |V_g f(x,\omega)m(x,\omega)|^r dx \right)^{\frac{s}{r}} d\omega \right)^{\frac{1}{s}} < \infty.$$

We shall write M_m^r , when $r = s$ and $M^{r,s}$, when $m \equiv 1$.

If instead of the order of integration (2.4), we use the alternative order (2.5), we obtain the so-called Wiener amalgam spaces $\mathbf{W}(\mathcal{FL}^p, \ell^q)(\mathbb{R}^d)$, which were introduced in [Fe83A]. The Fourier properties of these spaces for separable weights are explained in [Fe90].

Among the modulation spaces we find for instance $L^2(\mathbb{R}^d)$, or the Sobolev spaces $H^s(\mathbb{R}^d)$, the Feichtinger algebra $\mathbf{S}_0(\mathbb{R}^d) = M^{1,1}(\mathbb{R}^d)$ [Fe81], but also the spaces $M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)$ mentioned in the Introduction. Indeed, for $m(x,\omega) = \langle x \rangle^\alpha \langle \omega \rangle^\beta$, then $M_m^{r,s}(\mathbb{R}^d) = M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)$ as in (1.8). For the particular choice of weights $v_s(x,\omega) = (1 + |x|^2 + |\omega|^2)^{s/2}$, and $r = s = 2$ we obtain the so-called Shubin class spaces $(\mathbf{Q}_s(\mathbb{R}^d), \|\cdot\|_{\mathbf{Q}_s})$. Notice that these spaces are denoted by $M_{v_s}^2$ in [Gr01]. Special versions of the form M_m^2 are also given in [AF].

For future reference, we state the following Proposition (Proposition 11.3.1 in [Gr01]):

Proposition 1. *Let $g \in \mathcal{S}(\mathbb{R}^d) \setminus \{0\}$.*

- (1) *If $m(x,\omega) = m(x)$, then $M_m^2 = L_m^2$.*
- (2) *If $m(x,\omega) = m(\omega)$, then $M_m^2 = \mathcal{FL}_m^2$.*

3. LIEB'S UNCERTAINTY PRINCIPLE FOR MODULATION SPACES

The radar ambiguity function is given by:

$$(3.1) \quad A(f,g)(x,\omega) = \int_{\mathbb{R}^d} f\left(t - \frac{x}{2}\right) \overline{g\left(t + \frac{x}{2}\right)} e^{-2\pi i \omega \cdot t} dt.$$

Let us remark that the ambiguity function and its Fourier transform - the Wigner transform - can be extended to $(\mathbf{S}'_0(\mathbb{R}^d), \|\cdot\|_{\mathbf{S}'_0})$, since it is evaluated by taking a tensor product, then an automorphism of phase space \mathbb{R}^{2d} and then a partial Fourier transform. All these are well defined operators for the Feichtinger algebra $(\mathbf{S}_0(\mathbb{R}^d), \|\cdot\|_{\mathbf{S}_0})$, and extend, by duality, to $\mathbf{S}'_0(\mathbb{R}^d)$ in a unique way (see [FeHo90, FeHo14, Ho]).

It is straightforward to obtain the following properties.

$$(3.2) \quad A(f,g)(x,\omega) = A(\widehat{f}, \widehat{g})(-\omega, x).$$

From (3.1) it follows that

$$(3.3) \quad A(f, g)(x, \omega) = e^{-i\pi\omega \cdot x} V_g f(-x, \omega) .$$

Lieb proved (Theorem 1 in [L1]) the following theorem:

Theorem 4 (Lieb). *Let $r > 2$. If $f \in L^p(\mathbb{R})$ and $g \in L^{p'}(\mathbb{R})$ for $r' \leq p, p' \leq r$, then:*

$$(3.4) \quad \|A(f, g)\|_{L^r(\mathbb{R}^2)} \leq [H(r, p)]^{1/r} \|f\|_{L^p(\mathbb{R})} \|g\|_{L^{p'}(\mathbb{R})},$$

where $H(r, p) > 0$ is given by:

$$(3.5) \quad [H(r, p)]^2 = \frac{pp'}{r^2} |r - 2|^{2-r} |r - p|^{-1+r/p} |r - p'|^{-1+r/p'},$$

with the convention $0^0 \equiv 1$ when $r = p$ or $r = p'$.

This inequality is sometimes called *Lieb's uncertainty principle*, as it implies a lower bound on the measure of the support of $A(f, g)$ [Gr01].

Lieb also proved a reverse inequality (Theorem 2 in [L1]).

Theorem 5 (Lieb). *Assume that, for a.e. fixed $x \in \mathbb{R}$, the function $t \mapsto f(t - \frac{x}{2}) \overline{g(t + \frac{x}{2})}$ is in $L^1(\mathbb{R})$ for almost every $x \in \mathbb{R}$. Let $1 \leq r < 2$ and assume that $0 < \|A(f, g)\|_{L^r(\mathbb{R}^2)} < \infty$. Then $f, g \in L^u(\mathbb{R})$ for every $r \leq u \leq r'$. Moreover, for any p with $r \leq p, p' \leq r'$, we have that:*

$$(3.6) \quad \|A(f, g)\|_{L^r(\mathbb{R}^2)} \geq [H(r, p)]^{1/r} \|f\|_{L^p(\mathbb{R})} \|g\|_{L^{p'}(\mathbb{R})}.$$

Remark 1. The constant $H(r, p)$ can be expressed as:

$$(3.7) \quad H(r, p) = C_{r'}^r \left(\frac{C_{\frac{p}{r'}}^{\frac{p}{r'}} C_{\frac{p'}{r'}}^{\frac{p'}{r'}}}{C_{\frac{r}{r'}}^{\frac{r}{r'}}} \right)^{r/r'},$$

where, for $0 < p \leq \infty$, C_p is the Babenko-Beckner [Ba, Be] constant, which reads:

$$(3.8) \quad C_p = \sqrt{\frac{p^{1/p}}{|p'|^{1/p'}}},$$

for $p \neq 1$ and $p \neq \infty$, and

$$(3.9) \quad C_1 = C_\infty = 1.$$

Remark 2. The constants in (3.4, 3.6) are sharp. Lieb also proved that one can obtain an equality in (3.4) for $p, p' > r'$ and in (3.6) for $1 < r < 2$ if and only if f, g are certain matched Gaussians.

The purpose of this section is to generalize inequality (3.6) to arbitrary dimension $d > 1$ and for the mixed norms $\|\cdot\|_{L_{x,\omega}^{r,s}(\mathbb{R}^{2d})}$ and $\|\cdot\|_{L_{\omega,x}^{r,s}(\mathbb{R}^{2d})}$ (cf. (2.4, 2.5)).

Remark 3. A generalization of inequality (3.4) for modulation spaces was derived in [CoNi].

We shall prove the following generalization of the reverse Lieb inequality:

Theorem 6. Assume that $t \mapsto \widehat{f}\left(t + \frac{\omega}{2}\right) \overline{\widehat{g}\left(t - \frac{\omega}{2}\right)}$ is in $L^1(\mathbb{R}^d)$ for a.e. $\omega \in \mathbb{R}^d$. Notice that under these circumstances the definition of $A(f, g)$ makes sense, in view of (3.2). Let $1 \leq r, s \leq 2$ and assume that $0 < \|A(f, g)\|_{L_{x, \omega}^{r, s}(\mathbb{R}^{2d})} < \infty$. Then $\widehat{f}, \widehat{g} \in L^u(\mathbb{R}^d)$ for every $0 < u \leq r'$. Moreover, for every pair u, v such that $0 < u, v \leq r'$ and

$$(3.10) \quad \frac{1}{u} + \frac{1}{v} = \frac{1}{s} + \frac{1}{r'},$$

we have:

$$(3.11) \quad \|A(f, g)\|_{L_{x, \omega}^{r, s}(\mathbb{R}^{2d})} \geq B(r, s, u, v) \|\widehat{f}\|_{L^u(\mathbb{R}^d)} \|\widehat{g}\|_{L^v(\mathbb{R}^d)},$$

where

$$(3.12) \quad B(r, s, u, v) = C_{r'}^d \left(\frac{C_{u/r'} C_{v/r'}}{C_{s/r'}} \right)^{d/r'}.$$

The constant in (3.11) is sharp and for $1 < r < 2$, $0 < u, v < r'$, we have an equality if and only if \widehat{f}, \widehat{g} are Gaussians of the form:

$$(3.13) \quad \begin{aligned} \widehat{f}(\omega) &= \exp[-\omega \cdot (|m'|A + iB)\omega + c \cdot \omega + \gamma] \\ \widehat{g}(\omega) &= \exp[-\omega \cdot (|n'|A - iB)\omega + \tilde{c} \cdot \omega + \tilde{\gamma}] \end{aligned}$$

where A is a real, symmetric, positive-definite $d \times d$ matrix, B is a real, symmetric $d \times d$ matrix, $c, \tilde{c} \in \mathbb{C}^d$, $\gamma, \tilde{\gamma} \in \mathbb{C}$, and:

$$(3.14) \quad m' = \frac{u(r-1)}{ur - u - r}, \quad n' = \frac{v(r-1)}{vr - v - r}.$$

Here m' and n' are the Hölder duals of $m = \frac{u}{r}$ and $n = \frac{v}{r}$, respectively.

From this theorem and its proof we obtain the following two corollaries:

Corollary 1. Assume that $t \mapsto f\left(t - \frac{x}{2}\right) \overline{g\left(t + \frac{x}{2}\right)}$ is in $L^1(\mathbb{R}^d)$ for a.e. $x \in \mathbb{R}^d$. Let $1 \leq r, s \leq 2$ and assume that $0 < \|A(f, g)\|_{L_{\omega, x}^{r, s}(\mathbb{R}^{2d})} < \infty$. Then $f, g \in L^u(\mathbb{R}^d)$ for every $0 < u \leq r'$. Moreover, for every pair u, v such that $0 < u, v \leq r'$ and (3.10) holds, we have:

$$(3.15) \quad \|A(f, g)\|_{L_{\omega, x}^{r, s}(\mathbb{R}^{2d})} \geq B(r, s, u, v) \|f\|_{L^u(\mathbb{R}^d)} \|g\|_{L^v(\mathbb{R}^d)},$$

where

$$(3.16) \quad B(r, s, u, v) = C_{r'}^d \left(\frac{C_{u/r'} C_{v/r'}}{C_{s/r'}} \right)^{d/r'}.$$

The constant in (3.15) is sharp and for $1 < s < 2$ and $0 < u, v < r'$, we have an equality if and only if f, g are matched Gaussians of the form (3.13, 3.14) with \widehat{f}, \widehat{g} replaced by f, g , respectively and ω replaced by x .

The proof of this corollary follows exactly the same steps of the proof of Theorem 6 (see section 3.2).

Corollary 2. Let $1 \leq r, s \leq 2$; $1 < p, q < \infty$ and $g \in \mathcal{S}(\mathbb{R}^d) \setminus \{0\}$. Assume that f is such that $0 < \|f\|_{g, M^{r, s}(\mathbb{R}^d)} < \infty$ and $\|f\|_{L^p(\mathbb{R}^d)} < \infty$, $\|\widehat{f}\|_{L^q(\mathbb{R}^d)} < \infty$.

∞ . Then $f, \widehat{f} \in L^u(\mathbb{R}^d)$ for every $0 < u \leq r'$. Moreover, for every $0 < v \leq r'$, such that (3.10) holds, we have:

$$(3.17) \quad \|\widehat{f}\|_{L^u(\mathbb{R}^d)} \leq \frac{\|f\|_{g, M^{r,s}(\mathbb{R}^d)}}{B(r, s, u, v) \|\widehat{g}\|_{L^v(\mathbb{R}^d)}}$$

and

$$(3.18) \quad \|f\|_{L^u(\mathbb{R}^d)} \leq \frac{\|f\|_{g, M^{r,s}(\mathbb{R}^d)}}{B(r, s, u, v) \|g\|_{L^v(\mathbb{R}^d)}}$$

where $B(r, s, u, v)$ is given by (3.12).

Proof. By assumption, we have $\widehat{f} \in L^q(\mathbb{R}^d)$ and $\widehat{g} \in L^{q'}(\mathbb{R}^d)$. It follows by Hölder's inequality that the function $t \mapsto \widehat{f}(t + \frac{\omega}{2}) \overline{\widehat{g}(t - \frac{\omega}{2})}$ is in $L^1(\mathbb{R}^d)$ for a.e. $\omega \in \mathbb{R}^d$. Also, from (3.3) we have

$$(3.19) \quad \begin{aligned} \infty &> \|f\|_{g, M^{r,s}(\mathbb{R}^d)} = \\ &= \left(\int_{\mathbb{R}^d} \left(\int_{\mathbb{R}^d} |V_g f(x, \omega)|^r dx \right)^{s/r} d\omega \right)^{1/s} = \\ &= \|V_g f\|_{L_{x,\omega}^{r,s}(\mathbb{R}^{2d})} = \|A(f, g)\|_{L_{x,\omega}^{r,s}(\mathbb{R}^{2d})} \end{aligned}$$

By Theorem 6 we obtain (3.17). Eq. (3.18) is an immediate consequence of Corollary 1 and the fact that (cf.(3.2)):

$$(3.20) \quad \|A(f, g)\|_{L_{x,\omega}^{r,s}(\mathbb{R}^{2d})} = \|A(\widehat{f}, \widehat{g})\|_{L_{\omega,x}^{r,s}(\mathbb{R}^{2d})}.$$

□

To prove Theorem 6 we shall need to recapitulate some classical theorems in harmonic analysis.

3.1. Some results in harmonic analysis. The first result is the Hausdorff-Young inequality:

Theorem 7 (Hausdorff-Young). *Let $r \in [1, 2]$. If $f \in L^r(\mathbb{R}^d)$, then $\widehat{f} \in L^{r'}(\mathbb{R}^d)$, and:*

$$(3.21) \quad \|\widehat{f}\|_{L^{r'}(\mathbb{R}^d)} \leq C_r^d \|f\|_{L^r(\mathbb{R}^d)},$$

where C_r is the Babenko-Beckner constant (3.8, 3.9). There is obviously a converse result. Suppose that \widehat{f} exists and $\widehat{f} \in L^r(\mathbb{R}^d)$. Since $(\widehat{f})^\widehat{(x)} = f(-x)$ then $f \in L^{r'}(\mathbb{R}^d)$ and

$$(3.22) \quad \|f\|_{L^{r'}(\mathbb{R}^d)} \leq C_r^d \|\widehat{f}\|_{L^r(\mathbb{R}^d)}.$$

Equality is achieved in (3.21) and in (3.22) when $1 < r < 2$ if and only if f is a Gaussian of the form

$$(3.23) \quad f(x) = \gamma \exp(-x \cdot Ax + c \cdot x),$$

with $\gamma \in \mathbb{C}$, A is any real, symmetric, positive-definite $d \times d$ matrix and $c \in \mathbb{C}^d$. For $r = 1$ we can obtain an equality for many different functions. For $r = r' = 2$ an equality holds for all functions (Plancherel's Theorem).

The sharp constant in (3.21) is due to Beckner [Be] and under some more restrictive conditions to Babenko [Ba]. The criterion for equality was obtained by Lieb [L2].

The second inequality is due to Young. It applies to the convolution of two functions:

$$(3.24) \quad (f \star g)(x) = \int_{\mathbb{R}^d} f(x-y)g(y)dy.$$

Theorem 8 (Young). *Let $1 \leq m, n, r \leq \infty$ with $\frac{1}{m} + \frac{1}{n} = 1 + \frac{1}{r}$. If $f \in L^m(\mathbb{R}^d)$ and $g \in L^n(\mathbb{R}^d)$, then $f \star g \in L^r(\mathbb{R}^d)$ and*

$$(3.25) \quad \|f \star g\|_{L^r(\mathbb{R}^d)} \leq \left(\frac{C_m C_n}{C_r} \right)^d \|f\|_{L^m(\mathbb{R}^d)} \|g\|_{L^n(\mathbb{R}^d)}.$$

Here C_m, C_r , etc denote the Babenko-Beckner constant as defined in (3.8, 3.9). Equality in (3.25) holds for $m, n > 1$ if and only if

$$(3.26) \quad \begin{aligned} f(x) &= \gamma \exp[-m'(x - \bar{x}) \cdot A(x - \bar{x}) - ik \cdot x] \\ g(x) &= \tilde{\gamma} \exp[-n'(x - \tilde{x}) \cdot A(x - \tilde{x}) + ik \cdot x] \end{aligned}$$

where $\gamma, \tilde{\gamma} \in \mathbb{C}$, $\bar{x}, \tilde{x}, k \in \mathbb{R}^d$ and A is any real, symmetric, positive-definite $d \times d$ matrix.

The sharp constant in Young's inequality was obtained independently by Beckner [Be] and by Brascamp and Lieb [BrL].

The reverse inequality was first obtained by Leindler [Le]. Its sharp version was derived by Brascamp and Lieb [BrL]. A simple proof can be found in [Bar].

Theorem 9 (Leindler). *Let f and g be non-negative, real-valued functions on \mathbb{R}^d that are not identically zero and assume that $f \star g \in L^r(\mathbb{R}^d)$ for $0 < r \leq 1$. Let $0 < m, n \leq 1$ be such that $\frac{1}{m} + \frac{1}{n} = 1 + \frac{1}{r}$. Then $f \in L^m(\mathbb{R}^d)$ and $g \in L^n(\mathbb{R}^d)$, and we have:*

$$(3.27) \quad \|f \star g\|_{L^r(\mathbb{R}^d)} \geq \left(\frac{C_m C_n}{C_r} \right)^d \|f\|_{L^m(\mathbb{R}^d)} \|g\|_{L^n(\mathbb{R}^d)}.$$

Equality holds in (3.27) when $0 < m, n < 1$, if and only if

$$(3.28) \quad \begin{aligned} f(x) &= \gamma \exp[-|m'|(x - \bar{x}) \cdot A(x - \bar{x})] \\ g(x) &= \tilde{\gamma} \exp[-|n'|(x - \tilde{x}) \cdot A(x - \tilde{x})] \end{aligned}$$

where $\gamma, \tilde{\gamma} \in \mathbb{R}^+$, $\bar{x}, \tilde{x} \in \mathbb{R}^d$ and A is any real, symmetric, positive-definite $d \times d$ matrix.

Finally, we shall also use the following extension of Cauchy's functional equation to quadratics:

Theorem 10 (Lieb). *Let ξ, η be complex-valued, Lebesgue measurable functions on \mathbb{R}^d that satisfy $|\xi(t)| = |\eta(t)| = 1$, for a.e. $t \in \mathbb{R}^d$. Suppose there exist functions $\mu : \mathbb{R}^d \rightarrow \mathbb{R}^d$ and $\nu : \mathbb{R}^d \rightarrow \mathbb{R}$ (which are not a priori measurable) such that for a.e. $\omega \in \mathbb{R}^d$ the following holds for a.e. $t \in \mathbb{R}^d$:*

$$(3.29) \quad \xi\left(t + \frac{\omega}{2}\right) \eta\left(t - \frac{\omega}{2}\right) = \exp[i\mu(\omega) \cdot t + i\nu(\omega)].$$

Then there exist a real, symmetric $d \times d$ matrix A , $\bar{\xi}, \bar{\eta} \in \mathbb{R}^d$ and $\gamma, \delta \in \mathbb{R}$ such that:

$$(3.30) \quad \begin{aligned} \xi(t) &= \exp(it \cdot At + i\bar{\xi} \cdot t + i\gamma) \\ \eta(t) &= \exp(-it \cdot At + i\bar{\eta} \cdot t + i\delta) \end{aligned}$$

Conversely, if ξ, η are of the form (3.30), then (3.29) holds.

The proof of this theorem is basically the same as that of Lemma 4 in [L1] with the obvious adaptations to $d > 1$.

3.2. Proof of Theorem 6. We are now in a position to prove Theorem 6. The proof follows essentially the same steps as in the proof of Theorem 2 of [L1].

Since by assumption, for a.e. fixed $\omega \in \mathbb{R}^d$, the function $t \mapsto \widehat{f}\left(t + \frac{\omega}{2}\right) \overline{\widehat{g}\left(t - \frac{\omega}{2}\right)}$ is in $L^1(\mathbb{R}^d)$, then by (3.1,3.2) we conclude that $A(f, g)$ is the Fourier transform of this L^1 function. We can thus use (3.22) in Theorem 7 (with $p = 1$) to obtain:

$$(3.31) \quad \int_{\mathbb{R}^d} |A(f, g)(x, \omega)|^r dx \geq C_{r'}^{dr} \left(\int_{\mathbb{R}^d} |\widehat{f}\left(t + \frac{\omega}{2}\right) \widehat{g}\left(t - \frac{\omega}{2}\right)|^{r'} dt \right)^{\frac{r}{r'}}$$

for a.e. $x \in \mathbb{R}^d$. Notice that the left-hand side of (3.31) is finite for a.e. $x \in \mathbb{R}^d$, since by assumption $A(f, g) \in L_{x, \omega}^{r, s}(\mathbb{R}^{2d})$. The integral on the right-hand side can be written as:

$$(3.32) \quad J(\omega) = \int_{\mathbb{R}^d} |\widehat{f}\left(t + \frac{\omega}{2}\right)|^{r'} |\widehat{g}\left(t - \frac{\omega}{2}\right)|^{r'} dt = \left(|(\widehat{f})^\vee|^{r'} \star |\widehat{g}|^{r'} \right)(-\omega),$$

where $(\widehat{f})^\vee(\xi) = \widehat{f}(-\xi)$.

It follows from (3.31,3.32):

$$(3.33) \quad \|A(f, g)\|_{L_{x, \omega}^{r, s}(\mathbb{R}^{2d})} \geq C_{r'}^d \left(\int_{\mathbb{R}^d} |J(\omega)|^{\frac{s}{r'}} d\omega \right)^{\frac{1}{s}} = C_{r'}^d \| |(\widehat{f})^\vee|^{r'} \star |\widehat{g}|^{r'} \|_{L^{s/r'}(\mathbb{R}^d)}^{\frac{1}{r'}}$$

From the Leindler inequality, we thus get:

$$(3.34) \quad \begin{aligned} \|A(f, g)\|_{L_{x, \omega}^{r, s}(\mathbb{R}^{2d})} &\geq C_{r'}^d \left(\frac{C_m C_n}{C_{s/r'}} \right)^{\frac{d}{r'}} \| |\widehat{f}|^{r'} \|_{L^m(\mathbb{R}^d)}^{\frac{1}{r'}} \| |\widehat{g}|^{r'} \|_{L^n(\mathbb{R}^d)}^{\frac{1}{r'}} = \\ &= C_{r'}^d \left(\frac{C_m C_n}{C_{s/r'}} \right)^{\frac{d}{r'}} \| \widehat{f} \|_{L^{mr'}(\mathbb{R}^d)} \| \widehat{g} \|_{L^{nr'}(\mathbb{R}^d)} \end{aligned}$$

where $0 < m, n \leq 1$ with

$$(3.35) \quad \frac{1}{m} + \frac{1}{n} = 1 + \frac{r'}{s}.$$

We thus recover (3.11) with $0 < u = mr' \leq r'$, $0 < v = nr' \leq r'$ and $\frac{1}{u} + \frac{1}{v} = \frac{1}{r'} + \frac{1}{s}$. Notice that if $r = s$, then u, v are conjugate to each other, as in Lieb's inequality (3.6).

From Theorem 7, we have an equality in (3.31) for $1 < r < 2$, if and only if (cf.(3.23)):

$$(3.36) \quad \widehat{f}\left(t + \frac{\omega}{2}\right) \overline{\widehat{g}\left(t - \frac{\omega}{2}\right)} = \lambda(\omega) \exp[-t \cdot C(\omega)t + c(\omega) \cdot t]$$

for a.e. $\omega \in \mathbb{R}^d$, and where $\lambda(\omega) \in \mathbb{C}$, $C(\omega)$ is any real, symmetric, positive-definite $d \times d$ matrix and $c(\omega)$ is any vector in \mathbb{C}^d .

On the other hand, we have an equality in (3.34) for $0 < u, v < r'$ (or equivalently, for $0 < n, m < 1$) if and only if

$$(3.37) \quad \begin{aligned} |\widehat{f}(\omega)| &= \exp[-|m'|(\omega - \zeta) \cdot A(\omega - \zeta) + \delta] \\ |\widehat{g}(\omega)| &= \exp[-|n'|(\omega - \chi) \cdot A(\omega - \chi) + \widetilde{\delta}] \end{aligned}$$

where A is a real, symmetric positive-definite matrix, $\zeta, \chi \in \mathbb{R}^d$, $\delta, \widetilde{\delta} \in \mathbb{R}$, and:

$$(3.38) \quad \begin{aligned} m' &= \frac{m}{m-1} = \frac{u}{u-r'} = \frac{u(r-1)}{u(r-1)-r} \\ n' &= \frac{n}{n-1} = \frac{v}{v-r'} = \frac{v(r-1)}{v(r-1)-r} \end{aligned}$$

Since these functions never vanish, we may safely define

$$(3.39) \quad \xi(\omega) \equiv \frac{\widehat{f}(\omega)}{|\widehat{f}(\omega)|}, \quad \eta(\omega) \equiv \frac{\overline{\widehat{g}(\omega)}}{|\widehat{g}(\omega)|}.$$

By a simple inspection, we conclude that ξ and η satisfy all the conditions stated in Theorem 10 for $\mu(\omega) = \text{Im}(c(\omega))$ and $\nu(\omega) = -i \ln(\lambda(\omega)/|\lambda(\omega)|)$. We conclude that:

$$(3.40) \quad \begin{aligned} \xi(\omega) &= \exp(-i\omega \cdot B\omega + ig \cdot \omega + i\rho) \\ \eta(\omega) &= \exp(i\omega \cdot B\omega + ih \cdot \omega + i\widetilde{\rho}) \end{aligned}$$

where B is any real, symmetric $d \times d$ matrix, $g, h \in \mathbb{R}^d$ and $\rho, \widetilde{\rho} \in \mathbb{R}$. From $\widehat{f}(\omega) = \xi(\omega)|\widehat{f}(\omega)|$ and $\widehat{g}(\omega) = \overline{\eta(\omega)}|\widehat{g}(\omega)|$, and (3.37,3.40) we recover (3.13), which concludes the proof.

4. COWLING-PRICE TYPE UNCERTAINTY PRINCIPLE

In the sequel we shall consider functionals of the form:

$$(4.1) \quad \mathfrak{F}_{a,b}^{p,q}[f] = \| |x|^a f \|_{L^p(\mathbb{R}^d)} + \| |\omega|^b \widehat{f} \|_{L^q(\mathbb{R}^d)}$$

Proposition 2. *Let $1 \leq p, q \leq \infty$ and $a, b \geq 0$. The functional $\mathfrak{F}_{a,b}^{p,q}$ is continuous and convex in $L_a^p(\mathbb{R}^d) \cap \mathcal{FL}_b^q(\mathbb{R}^d)$.*

Proof. Convexity is obvious. The rest is an immediate consequence of the fact that the functional is just the natural norm (in the spirit of H. Triebel [Tr83]) of the function space $L_a^p(\mathbb{R}^d) \cap \mathcal{FL}_b^q(\mathbb{R}^d)$, and is therefore continuous. \square

Before we continue let us make the following observation:

Lemma 1. *Let $1 < p, q < \infty$ and $a, b \geq 0$. The space $L_a^p(\mathbb{R}^d) \cap \mathcal{FL}_b^q(\mathbb{R}^d)$ is reflexive.*

Proof. Since $L_a^p(\mathbb{R}^d)$ and $\mathcal{F}L_b^q(\mathbb{R}^d)$ are reflexive, so is $\mathcal{G} \equiv L_a^p(\mathbb{R}^d) \times \mathcal{F}L_b^q(\mathbb{R}^d)$, when we use the usual product norm $\|(x, y)\|_{X \times Y} = \|x\|_X + \|y\|_Y$ for the product of normed vector spaces $X \times Y$ and dual bracket $\langle (x^*, y^*), (x, y) \rangle_{X \times Y} = \langle x^*, x \rangle_X + \langle y^*, y \rangle_Y$, for $(x^*, y^*) \in X' \times Y'$, $(x, y) \in X \times Y$. Then the set $\mathcal{K} \equiv \{(f, g) \in \mathcal{G} : f = g\}$ is a closed subspace of \mathcal{G} with respect to the previous product topology. Therefore, \mathcal{K} is reflexive. Since $L_a^p(\mathbb{R}^d) \cap \mathcal{F}L_b^q(\mathbb{R}^d)$ is isomorphic to \mathcal{K} , it is also reflexive. \square

Notice that this also follows from an old result concerning dual spaces of sums and intersections of Banach spaces (when both are embedded continuously into $\mathcal{S}'(\mathbb{R}^d)$) [LiRo].

To avoid a proliferation of indices, we shall use in the sequel the notation

$$(4.2) \quad \mathcal{B}_1(\mathbb{R}^d) = L_a^p(\mathbb{R}^d) \cap \mathcal{F}L_b^q(\mathbb{R}^d), \quad \mathcal{B}_2(\mathbb{R}^d) = M_{\alpha, \beta}^{r, s}(\mathbb{R}^d)$$

with indices $a, b, p, q, r, s, \alpha, \beta$ to be specified. Moreover, we shall assume a fixed window $g \in \mathcal{S}(\mathbb{R}^d) \setminus \{0\}$ for the norm of the modulation space $M_{\alpha, \beta}^{r, s}(\mathbb{R}^d)$.

We denote by $\overline{\mathcal{B}_1}(R)$ the closed ball of radius $R > 0$ in $\mathcal{B}_1(\mathbb{R}^d)$:

$$(4.3) \quad \overline{\mathcal{B}_1}(R) = \left\{ f \in \mathcal{B}_1(\mathbb{R}^d) : \|f\|_{\mathcal{B}_1(\mathbb{R}^d)} \leq R \right\}.$$

Proposition 3. *Let $0 < r, s \leq 2$, $\alpha, \beta, a, b \geq 0$ and $1 < p, q < \infty$. Suppose that $r \leq p$ and $s \leq q$, and that inequality (1.10) holds with all factors non-negative. If, for $R > 0$, the set*

$$(4.4) \quad \mathcal{U}(R) = \left\{ f \in \overline{\mathcal{B}_1}(R) : \|f\|_{\mathcal{B}_2(\mathbb{R}^d)} = 1 \right\}$$

is nonempty, then it is a weakly sequentially compact subset of $\mathcal{B}_1(\mathbb{R}^d)$.

Proof. Suppose that $\mathcal{U}(R)$ is nonempty. Let $(f_n)_n$ be an arbitrary sequence in $\mathcal{U}(R)$. Since $\mathcal{B}_1(\mathbb{R}^d)$ is reflexive (Lemma 1), we conclude that $(f_n)_n$ has a weakly convergent subsequence, say $(g_k)_k$ with $g_k \rightharpoonup g$ for some $g \in \mathcal{B}_1(\mathbb{R}^d)$. Since $\overline{\mathcal{B}_1}(R)$ is convex and closed, we have by Mazur's Theorem that $g \in \overline{\mathcal{B}_1}(R)$. It remains to prove that $\|g\|_{\mathcal{B}_2(\mathbb{R}^d)} = 1$. As $\mathcal{B}_1(\mathbb{R}^d)$ is compactly embedded in $\mathcal{B}_2(\mathbb{R}^d)$ (see Theorem 2), the sequence $(g_k)_k$ has a subsequence $(h_l)_l$ converging strongly in $\mathcal{B}_2(\mathbb{R}^d)$ to some $h \in \mathcal{B}_2(\mathbb{R}^d)$. By the continuity of the norm, we have: $\|h\|_{\mathcal{B}_2(\mathbb{R}^d)} = \lim_{l \rightarrow \infty} \|h_l\|_{\mathcal{B}_2(\mathbb{R}^d)} = 1$. We conclude the proof by showing that $h = g$ a.e.. Since $(h_l)_l$ is a subsequence of $(g_k)_k$, we have for all $u \in \mathcal{B}_2'(\mathbb{R}^d)$:

$$(4.5) \quad \begin{aligned} \lim_{l \rightarrow \infty} \langle u, g - h_l \rangle_2 &= \langle u, g \rangle_2 - \lim_{l \rightarrow \infty} \langle u, h_l \rangle_2 = \\ &= \langle u, g \rangle_1 - \lim_{l \rightarrow \infty} \langle u, h_l \rangle_1 = \langle u, g \rangle_1 - \langle u, g \rangle_1 = 0, \end{aligned}$$

where $\langle \cdot, \cdot \rangle_1$, $\langle \cdot, \cdot \rangle_2$ denote the duality brackets in $\mathcal{B}_1(\mathbb{R}^d)$, $\mathcal{B}_2(\mathbb{R}^d)$, respectively, and where we used the fact that $u \in \mathcal{B}_2'(\mathbb{R}^d) \subset \mathcal{B}_1'(\mathbb{R}^d)$ and $g, h_l \in \mathcal{B}_1(\mathbb{R}^d) \subset \mathcal{B}_2(\mathbb{R}^d)$.

On the other hand, since $(h_l)_l$ converges to h strongly in $\mathcal{B}_2(\mathbb{R}^d)$, it also converges weakly, and thus:

$$(4.6) \quad \lim_{l \rightarrow \infty} \langle u, h_l \rangle_2 = \langle u, h \rangle_2.$$

From (4.5,4.6), we conclude that

$$(4.7) \quad \langle u, g - h \rangle_2 = 0,$$

for all $u \in \mathcal{B}'_2(\mathbb{R}^d)$, and hence $g = h$ a.e. \square

Lemma 2. *Assume the same conditions as in the previous Proposition. If $\mathcal{U}(R)$ is nonempty, then the functional $\mathfrak{F}_{a,b}^{p,q}$ is weakly lower semicontinuous in $\mathcal{U}(R)$.*

Proof. The closed ball $\overline{\mathcal{B}_1}(R)$ is a convex and closed subset of $\mathcal{B}_1(\mathbb{R}^d)$. Since $\mathfrak{F}_{a,b}^{p,q}$ is convex and continuous, it is weakly lower semicontinuous on $\overline{\mathcal{B}_1}(R)$. On the other hand, $\mathcal{U}(R)$ is a weakly sequentially compact subset of $\mathcal{B}_1(\mathbb{R}^d)$. Then the restriction of $\mathfrak{F}_{a,b}^{p,q}$ to $\mathcal{U}(R)$ is also weakly lower semicontinuous. \square

Theorem 11. *Let $1 < r, s \leq 2$, $\alpha, \beta \geq 0$, $0 < a, b \leq r'$ and $r \leq p \leq r'$, $s \leq q \leq r'$. If, in addition, inequality (1.10) holds, with all factors non-negative, then the functional*

$$(4.8) \quad \mathfrak{F}_{a,b}^{p,q}[f] = \| |x|^a f \|_{L^p(\mathbb{R}^d)} + \| |\omega|^b \widehat{f} \|_{L^q(\mathbb{R}^d)}$$

attains a minimum in

$$(4.9) \quad \Omega = \left\{ f \in L_a^p(\mathbb{R}^d) \cap \mathcal{F}L_b^q(\mathbb{R}^d) : \|f\|_{M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)} = 1 \right\}.$$

Proof. Let $C_G > 0$ denote the sharp constant in the Galperin-Gröchenig inequality:

$$(4.10) \quad C_G = \inf \frac{\|f\|_{\mathcal{B}_1(\mathbb{R}^d)}}{\|f\|_{\mathcal{B}_2(\mathbb{R}^d)}},$$

where the infimum is taken over all $f \in \mathcal{B}_1(\mathbb{R}^d) \setminus \{0\}$. Consequently, there exists $f_1 \in \Omega$ such that

$$(4.11) \quad C_G < \|f_1\|_{\mathcal{B}_1(\mathbb{R}^d)} < 2C_G.$$

Let

$$(4.12) \quad M = \max \left\{ \frac{ap-1}{p}, \frac{bq-1}{q}, 0 \right\},$$

and

$$(4.13) \quad R_G = 2^M \left(\frac{1}{B(r, s, q, u) \|\widehat{g}\|_{L^u(\mathbb{R}^d)}} + \frac{1}{B(r, s, p, v) \|g\|_{L^v(\mathbb{R}^d)}} + 2C_G \right),$$

where $0 < u, v \leq r'$ are such that:

$$(4.14) \quad \frac{1}{q} + \frac{1}{u} = \frac{1}{r'} + \frac{1}{s}, \quad \frac{1}{p} + \frac{1}{v} = \frac{1}{r'} + \frac{1}{s}.$$

From (4.11), we conclude that $f_1 \in \overline{\mathcal{B}_1}(R_G)$, and hence $\mathcal{U}(R_G)$, as defined in (4.4) is nonempty. By Proposition 3 it is weakly sequentially compact and $\mathfrak{F}_{a,b}^{p,q}$ is weakly lower semicontinuous therein. Consequently, there exists a minimizer f_0 of $\mathfrak{F}_{a,b}^{p,q}$ in $\mathcal{U}(R_G)$. It remains to prove that f_0 is a minimizer everywhere in Ω .

Suppose that $ap \geq 1$. We then have from the inequality $|x + y|^t \leq 2^{t-1}(|x|^t + |y|^t)$ for $t \geq 1$:

$$\begin{aligned} \|f\|_{L_a^p(\mathbb{R}^d)} &= \left(\int_{\mathbb{R}^d} (1 + |x|)^{ap} |f(x)|^p dx \right)^{1/p} \\ &\leq 2^{(ap-1)/p} \left(\|f\|_{L^p(\mathbb{R}^d)}^p + \| |x|^a f \|_{L^p(\mathbb{R}^d)}^p \right)^{1/p} \leq 2^M \left(\|f\|_{L^p(\mathbb{R}^d)} + \| |x|^a f \|_{L^p(\mathbb{R}^d)} \right) \end{aligned} \quad (4.15)$$

Alternatively, if $ap < 1$, then we obtain again:

$$\begin{aligned} \|f\|_{L_a^p(\mathbb{R}^d)} &= \left(\int_{\mathbb{R}^d} (1 + |x|)^{ap} |f(x)|^p dx \right)^{1/p} \\ &\leq \left(\|f\|_{L^p(\mathbb{R}^d)}^p + \| |x|^a f \|_{L^p(\mathbb{R}^d)}^p \right)^{1/p} \leq 2^M \left(\|f\|_{L^p(\mathbb{R}^d)} + \| |x|^a f \|_{L^p(\mathbb{R}^d)} \right) \end{aligned} \quad (4.16)$$

In the same fashion, we have:

$$\|\widehat{f}\|_{L_b^q(\mathbb{R}^d)} \leq 2^M \left(\|\widehat{f}\|_{L^q(\mathbb{R}^d)} + \| |\omega|^b \widehat{f} \|_{L^q(\mathbb{R}^d)} \right) \quad (4.17)$$

Altogether, we conclude that:

$$\begin{aligned} \|f\|_{\mathcal{B}_1(\mathbb{R}^d)} &= \|f\|_{L_a^p(\mathbb{R}^d)} + \|\widehat{f}\|_{L_b^q(\mathbb{R}^d)} \\ &\leq 2^M \left(\|f\|_{L^p(\mathbb{R}^d)} + \|\widehat{f}\|_{L^q(\mathbb{R}^d)} + \mathfrak{F}_{a,b}^{p,q}[f] \right) \end{aligned} \quad (4.18)$$

If f is such that $\|f\|_{\mathcal{B}_1(\mathbb{R}^d)} > R_G$, then:

$$\begin{aligned} \|f\|_{L^p(\mathbb{R}^d)} + \|\widehat{f}\|_{L^q(\mathbb{R}^d)} + \mathfrak{F}_{a,b}^{p,q}[f] &\geq 2^{-M} \|f\|_{\mathcal{B}_1(\mathbb{R}^d)} \\ &> \frac{1}{B(r,s,q,u)\|\widehat{g}\|_{L^u(\mathbb{R}^d)}} + \frac{1}{B(r,s,p,v)\|g\|_{L^v(\mathbb{R}^d)}} + 2C_G \end{aligned} \quad (4.19)$$

But, in view of the trivial inequality $\|f\|_{g,M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)} \geq \|f\|_{g,M^{r,s}(\mathbb{R}^d)}$ for all $\alpha, \beta \geq 0$ and of Corollary 2, this entails:

$$\mathfrak{F}_{a,b}^{p,q}[f] > 2C_G > \|f_1\|_{\mathcal{B}_1(\mathbb{R})} \geq \mathfrak{F}_{a,b}^{p,q}[f_1] \geq \mathfrak{F}_{a,b}^{p,q}[f_0] \quad (4.20)$$

which concludes the proof. \square

5. AN UNCERTAINTY PRINCIPLE IN THE HILBERT CASE

In this section, we deal with the more specific case where $r = s = p = q = 2$. All the functional spaces involved become Hilbert spaces. The advantage is that we can slightly generalize our construction and consider weights other than the powers $|x|^a$ and $|\omega|^b$. Moreover, as we shall see later, the minimizers are given by the eigenvalue equation of the inverse of a certain compact localization operator.

Let us start with the following definition.

Definition 3. *Given some weight $m_0 \in \mathcal{P}(\mathbb{R}^{2d})$, we call a pair of continuous functions $\psi, \phi \in L_{loc}^\infty(\mathbb{R}^d)$ m_0 -admissible weights if*

$$m(x, \omega) = \sqrt{|m_0(x, \omega)|^2 + |\psi(x)|^2 + |\phi(\omega)|^2}$$

satisfies $m \in \mathcal{P}(\mathbb{R}^{2d})$ and $\frac{m_0}{m} \in L^\infty(\mathbb{R}^{2d})$.

Let us remark that the conditions of the previous definition imply that $|\psi(x)| \rightarrow \infty$ and $|\phi(x)| \rightarrow \infty$ at a polynomial rate as $|x| \rightarrow \infty$.

A simple consequence of this definition and of the compactness theorem 3 is the following.

Corollary 3. *Let (ψ, ϕ) be m_0 -admissible weights. Then we have the compact embedding*

$$(5.1) \quad M_m^2(\mathbb{R}^d) \subset\subset M_{m_0}^2(\mathbb{R}^d).$$

For future reference, we consider the following alternative norm.

Lemma 3. *Let (ψ, ϕ) be m_0 -admissible weights. Then the norm $\|\cdot\|_{M_m^2(\mathbb{R}^d)}$ is equivalent to*

$$(5.2) \quad \|f\|_{\psi, \phi, m_0}^2 = \|f\|_{M_{m_0}^2(\mathbb{R}^d)}^2 + \|\psi f\|_{L^2(\mathbb{R}^d)}^2 + \|\phi \widehat{f}\|_{L^2(\mathbb{R}^d)}^2.$$

Proof. Following the same steps of the proof which leads to Proposition 1 (see the proof of Proposition 11.3.1 in [Gr01]), we obtain

$$(5.3) \quad \begin{aligned} \|f\|_{\psi, \phi, m_0}^2 &= \|f\|_{M_{m_0}^2(\mathbb{R}^d)}^2 + \|\psi f\|_{L^2(\mathbb{R}^d)}^2 + \|\phi \widehat{f}\|_{L^2(\mathbb{R}^d)}^2 \\ &\asymp \|f\|_{M_{m_0}^2(\mathbb{R}^d)}^2 + \|f\|_{M_{|\psi|}^2(\mathbb{R}^d)}^2 + \|f\|_{M_{|\phi|}^2(\mathbb{R}^d)}^2 = \\ &= \int_{\mathbb{R}^{2d}} m^2(z) |V_g f(z)|^2 dz = \|f\|_{M_m^2(\mathbb{R}^d)}^2. \end{aligned}$$

□

Notice that we can associate the norm $\|\cdot\|_{\psi, \phi, m_0}$ to the inner product:

$$(5.4) \quad \begin{aligned} (u, v)_{\psi, \phi, m_0} &= \int_{\mathbb{R}^{2d}} m_0(z)^2 V_g u(z) \overline{V_g v(z)} dz + \\ &+ \int_{\mathbb{R}^d} |\psi(x)|^2 u(x) \overline{v(x)} dx + \int_{\mathbb{R}^d} |\phi(\omega)|^2 \widehat{u}(\omega) \overline{\widehat{v}(\omega)} d\omega, \end{aligned}$$

such that $(u, u)_{\psi, \phi, m_0} = \|u\|_{\psi, \phi, m_0}^2$. Thus, the space $\mathcal{B}^{\psi, \phi, m_0}(\mathbb{R}^d)$ of measurable functions f with finite $\|f\|_{\psi, \phi, m_0} < \infty$, endowed with the inner product $(\cdot, \cdot)_{\psi, \phi, m_0}$, is a Hilbert space and:

$$\mathcal{B}^{\psi, \phi, m_0}(\mathbb{R}^d) = M_m^2(\mathbb{R}^d).$$

We shall now follow *mutatis mutandis* our proof of weak sequential compactness (Proposition 3) for this case. In the sequel we shall always assume that (ψ, ϕ) are a m_0 -admissible set of weights with $m_0 \in \mathcal{P}(\mathbb{R}^{2d})$. Let us denote by $\overline{B}_R^{\psi, \phi, m_0}$ the following closed ball of radius $R > 0$:

$$(5.5) \quad \overline{B}_R^{\psi, \phi, m_0} := \left\{ f \in M_m^2(\mathbb{R}^d) : \|f\|_{\psi, \phi, m_0} \leq R \right\}.$$

Proposition 4. *Let $R > 0$. If the set*

$$(5.6) \quad U_R^{\psi, \phi, m_0} = \left\{ f \in \overline{B}_R^{\psi, \phi, m_0} : \|f\|_{M_{m_0}^2(\mathbb{R}^d)} = 1 \right\}$$

is nonempty, then it is a weakly sequentially compact subset of $M_m^2(\mathbb{R}^d)$.

Proof. Suppose that U_R^{ψ, ϕ, m_0} is nonempty. Let $(f_n)_n$ be an arbitrary sequence in U_R^{ψ, ϕ, m_0} . Since $M_m^2(\mathbb{R}^d)$ is reflexive (it is a Hilbert space), and the norms $\|\cdot\|_{M_m^2(\mathbb{R}^d)}$ and $\|\cdot\|_{\psi, \phi, m_0}$ are equivalent, we conclude that $(f_n)_n$ has a weakly convergent subsequence $(g_k)_k$, say

$$(5.7) \quad g_k \rightharpoonup g,$$

for some $g \in M_m^2(\mathbb{R}^d)$, and by Mazur's Theorem $g \in \overline{B_R^{\psi, \phi, m_0}}$. It remains to prove that $\|g\|_{M_{m_0}^2(\mathbb{R}^d)} = 1$. From Corollary 3, we conclude that the sequence $(g_k)_k$ has a subsequence $(h_l)_l$ converging strongly in $M_{m_0}^2(\mathbb{R}^d)$, say $\|h_l - h\|_{M_{m_0}^2(\mathbb{R}^d)} \rightarrow 0$, for some $h \in M_{m_0}^2(\mathbb{R}^d)$. By the continuity of the norm, we also have $\|h\|_{M_{m_0}^2(\mathbb{R}^d)} = 1$. The proof is complete if we show that $g = h$.

Recall that $M_{m_0}^2(\mathbb{R}^d)$ is a Hilbert space with inner product

$$(5.8) \quad (u, v)_{M_{m_0}^2(\mathbb{R}^d)} = \int_{\mathbb{R}^{2d}} m_0^2(z) V_g u(z) \overline{V_g v(z)} dz.$$

The mapping $(u, v) \mapsto (u, v)_{M_{m_0}^2(\mathbb{R}^d)}$ is a sesquilinear form on $\mathcal{B}^{\psi, \phi, m_0}(\mathbb{R}^d) \times \mathcal{B}^{\psi, \phi, m_0}(\mathbb{R}^d) = M_m^2(\mathbb{R}^d) \times M_m^2(\mathbb{R}^d)$. Moreover, it is bounded as we now prove. Since $m_0(z) \leq m(z)$ for a.e. $z \in \mathbb{R}^{2d}$, we have from the Cauchy-Schwarz inequality and Lemma 3:

$$(5.9) \quad \begin{aligned} & \left| (u, v)_{M_{m_0}^2(\mathbb{R}^d)} \right| \leq \int_{\mathbb{R}^{2d}} m^2(z) \left| V_g u(z) \overline{V_g v(z)} \right| dz \\ & \leq \|V_g u\|_{L_m^2(\mathbb{R}^{2d})} \|V_g v\|_{L_m^2(\mathbb{R}^{2d})} = \|u\|_{M_m^2(\mathbb{R}^d)} \|v\|_{M_m^2(\mathbb{R}^d)} \asymp \|u\|_{\psi, \phi, m_0} \|v\|_{\psi, \phi, m_0}. \end{aligned}$$

By the Riesz representation theorem, there exists a bounded linear operator $A : M_m^2(\mathbb{R}^d) \rightarrow M_{m_0}^2(\mathbb{R}^d)$, such that

$$(5.10) \quad (u, v)_{M_{m_0}^2(\mathbb{R}^d)} = (Au, v)_{\psi, \phi, m_0},$$

for all $u, v \in M_m^2(\mathbb{R}^d)$.

Let $u \in \mathcal{S}(\mathbb{R}^d)$. Since $m_0 \in \mathcal{P}(\mathbb{R}^{2d})$, we have that $\mathcal{S}(\mathbb{R}^d)$ is dense in $M_{m_0}^2(\mathbb{R}^d)$. We then have:

$$(5.11) \quad (h_l - g, u)_{M_{m_0}^2(\mathbb{R}^d)} = (A(h_l - g), u)_{\psi, \phi, m_0}.$$

If we take the limit $l \rightarrow \infty$, the right-hand side of the previous equation vanishes, while the left-hand side becomes $(h - g, u)_{M_{m_0}^2(\mathbb{R}^d)}$. Since $\mathcal{S}(\mathbb{R}^d)$ is dense in $M_{m_0}^2(\mathbb{R}^d)$, we conclude that $h = g$ a.e. \square

Remark 4. Since $\mathcal{B}^{\psi, \phi, m_0}(\mathbb{R}^d)$ may be identified with the modulation space $M_m^2(\mathbb{R}^d)$, we have a concrete realization of the operator A in the preceding proof as a localization operator. Recall that a localization operator with symbol σ and window $g \in \mathcal{S}(\mathbb{R}^d)$ is of the form

$$A_\sigma^g f = \int_{\mathbb{R}^{2d}} \sigma(x, \omega) \langle f, \pi(x, \omega)g \rangle \pi(x, \omega)g \, dx d\omega.$$

If we take for σ the weight m_0/m , then $A_{m_0/m}^g$ is an isomorphism between $M_m^2(\mathbb{R}^d)$ and $M_{m_0}^2(\mathbb{R}^d)$ by the results in [GT1, GT2].

Next, we define the functional $\mathfrak{F}^{\psi,\phi,m_0} : M_m^2(\mathbb{R}^d) \rightarrow [0, \infty)$,

$$(5.12) \quad \mathfrak{F}^{\psi,\phi,m_0}[f] = \|\psi f\|_{L^2(\mathbb{R}^d)}^2 + \|\phi \widehat{f}\|_{L^2(\mathbb{R}^d)}^2 = \|f\|_{\psi,\phi,m_0}^2 - \|f\|_{M_{m_0}^2(\mathbb{R}^d)}^2.$$

We then have:

Lemma 4. *Let $R > 0$ be such that U_R^{ψ,ϕ,m_0} as defined in Proposition 4 is nonempty. Then, the functional $\mathfrak{F}^{\psi,\phi,m_0}$ is weakly lower semicontinuous in U_R^{ψ,ϕ,m_0} .*

Proof. Consider the maps

$$(5.13) \quad (\mathcal{M}_\psi f)(x) := \psi(x)f(x), \quad (\mathcal{U}_\phi f)(\omega) := (\mathcal{M}_\phi \mathcal{F}f)(\omega) = \phi(\omega)\widehat{f}(\omega),$$

for every $x, \omega \in \mathbb{R}^d$ and $f \in \mathcal{S}(\mathbb{R}^d)$. They extend to bounded linear operators $M_m^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R}^d)$: $\|\mathcal{M}_\psi f\|_{L^2(\mathbb{R}^d)} = \|\psi f\|_{L^2(\mathbb{R}^d)} \leq \|f\|_{\psi,\phi,m_0} \asymp \|f\|_{M_m^2(\mathbb{R}^d)}$. Thus, the map $f \mapsto \|\mathcal{M}_\psi f\|_{L^2(\mathbb{R}^d)}$ is a continuous and convex functional on the closed ball $\overline{B}_R^{\psi,\phi,m_0}$, which is a convex and closed subset of $M_m^2(\mathbb{R}^{2d})$. This means that this map is weakly lower semicontinuous in $\overline{B}_R^{\psi,\phi,m_0}$.

On the other hand, by Proposition 4, U_R^{ψ,ϕ,m_0} is a weakly sequentially compact subset of $\mathcal{B}^{\psi,\phi,m_0}(\mathbb{R}^d)$. Hence the restriction of $\|\mathcal{M}_\psi f\|_{L^2(\mathbb{R}^d)}$ to $U_R^{\psi,\phi,m_0} \subset \overline{B}_R^{\psi,\phi,m_0}$ is weakly lower semicontinuous. The product of two nonnegative weakly lower semicontinuous functionals is again weakly lower semicontinuous, which entails that $\|\mathcal{M}_\psi f\|_{L^2(\mathbb{R}^d)}^2$ is weakly lower semicontinuous. The same can be said about $\|\mathcal{U}_\phi f\|_{L^2(\mathbb{R}^d)}^2$. Consequently, $\mathfrak{F}^{\psi,\phi,m_0}$, being the sum of these two functionals, is weakly lower semicontinuous in U_R^{ψ,ϕ,m_0} . \square

We next prove the existence of minimizers.

Theorem 12. *Let $R > 1$ be such that U_R^{ψ,ϕ,m_0} as defined in Proposition 4 is nonempty. Then there exists $f_0 \in U_R^{\psi,\phi,m_0}$ such that*

$$(5.14) \quad \mathfrak{F}^{\psi,\phi,m_0}[f_0] \leq \mathfrak{F}^{\psi,\phi,m_0}[f],$$

for all

$$(5.15) \quad f \in \Omega := \left\{ f \in M_m^2(\mathbb{R}^d) : \|f\|_{M_{m_0}^2(\mathbb{R}^d)} = 1 \right\}.$$

Proof. The set U_R^{ψ,ϕ,m_0} is weakly sequentially compact (cf. Proposition 4). Moreover, the functional $\mathfrak{F}^{\psi,\phi,m_0}$ is weakly lower semicontinuous (Lemma 4). Consequently, there exists a minimizer f_0 of $\mathfrak{F}^{\psi,\phi,m_0}$ in U_R^{ψ,ϕ,m_0} . It remains to prove that f_0 is in fact a minimizer on the whole set Ω .

From (5.12) and (5.15), we have:

$$(5.16) \quad \mathfrak{F}^{\psi,\phi,m_0}[f] = \|f\|_{\psi,\phi,m_0}^2 - 1,$$

for all $f \in \Omega$.

Since $f_0 \in U_R^{\psi,\phi,m_0}$, it follows:

$$(5.17) \quad \mathfrak{F}^{\psi,\phi,m_0}[f_0] \leq R^2 - 1.$$

On the other hand, if $f \in \Omega \setminus U_R^{\psi,\phi,m_0}$:

$$(5.18) \quad \mathfrak{F}^{\psi,\phi,m_0}[f] > R^2 - 1,$$

and the result follows. \square

6. EULER-LAGRANGE EQUATIONS

6.1. The Banach case. In this section, we shall derive an equation satisfied by the minimizer f_0 of the functional (4.8) in the constraint set (4.9). We thus consider the functional

$$(6.1) \quad \begin{aligned} \mathfrak{L}_{a,b}^{p,q}[f, \lambda] &:= \mathfrak{F}_{a,b}^{p,q}[f] + \lambda \left(1 - \|f\|_{M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)} \right) = \\ &= \| |x|^a f \|_{L^p(\mathbb{R}^d)} + \| |\omega|^b \widehat{f} \|_{L^q(\mathbb{R}^d)} + \lambda \left(1 - \|f\|_{M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)} \right), \end{aligned}$$

where λ is a Lagrange multiplier.

Theorem 13. *The minimizers f_0 of (4.8) in the constraint set (4.9) are solutions of the equation:*

$$(6.2) \quad \begin{aligned} &\| |x|^a f \|_{L^p(\mathbb{R}^d)}^{1-p} \int_{\mathbb{R}^d} |x|^{ap} |f(x)|^{p-2} f(x) \overline{u(x)} dx + \\ &+ \| |\omega|^b \widehat{f} \|_{L^q(\mathbb{R}^d)}^{1-q} \int_{\mathbb{R}^d} |\omega|^{bq} |\widehat{f}(\omega)|^{q-2} \widehat{f}(\omega) \overline{\widehat{u}(\omega)} d\omega = \\ &= \lambda \int_{\mathbb{R}^d} \|V_g f(\cdot, \omega)\|_{L_\alpha^r(\mathbb{R}^d)}^{s-r} \left[\int_{\mathbb{R}^d} |V_g f(x, \omega)|^{r-2} V_g f(x, \omega) \overline{V_g u(x, \omega)} \langle x \rangle^{\alpha r} dx \right] \langle \omega \rangle^{\beta s} d\omega, \end{aligned}$$

for all $u \in L_a^p(\mathbb{R}^d) \cap \mathcal{FL}_b^q(\mathbb{R}^d)$, and where

$$(6.3) \quad \|V_g f(\cdot, \omega)\|_{L_\alpha^r(\mathbb{R}^d)} = \left(\int_{\mathbb{R}^d} |V_g f(x, \omega)|^r \langle x \rangle^{\alpha r} dx \right)^{1/r}.$$

The minimizers satisfy this equation with the smallest possible value of λ .

Proof. Let $u \in \mathcal{S}(\mathbb{R}^d)$. Recall that $\mathcal{S}(\mathbb{R}^d)$ is a dense subset of $L_a^p(\mathbb{R}^d) \cap \mathcal{FL}_b^q(\mathbb{R}^d)$. For $t \in \mathbb{R}$, we have:

$$(6.4) \quad \begin{aligned} &\| |x|^a (f + tu) \|_{L^p(\mathbb{R}^d)} = \| |x|^a f \|_{L^p(\mathbb{R}^d)} + \\ &+ t \| |x|^a f \|_{L^p(\mathbb{R}^d)}^{1-p} \int_{\mathbb{R}^d} |x|^{ap} |f(x)|^{p-2} \mathcal{R} \left(f(x) \overline{u(x)} \right) dx + \mathcal{O}(t^2), \end{aligned}$$

where \mathcal{R} denotes the real part of a complex number.

Similarly:

$$(6.5) \quad \begin{aligned} &\| |\omega|^b (\widehat{f} + t\widehat{u}) \|_{L^q(\mathbb{R}^d)} = \| |\omega|^b \widehat{f} \|_{L^q(\mathbb{R}^d)} + \\ &+ t \| |\omega|^b \widehat{f} \|_{L^q(\mathbb{R}^d)}^{1-q} \int_{\mathbb{R}^d} |\omega|^{bq} |\widehat{f}(\omega)|^{q-2} \mathcal{R} \left(\widehat{f}(\omega) \overline{\widehat{u}(\omega)} \right) d\omega + \mathcal{O}(t^2). \end{aligned}$$

Finally:

$$\begin{aligned}
(6.6) \quad & \|f + tu\|_{M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)} = \|V_g(f + tu)\|_{L_{\alpha,\beta}^{r,s}(\mathbb{R}^{2d})} = \\
& = \|f\|_{M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)} + t\|f\|_{M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)}^{1-s} \times \\
& \times \int_{\mathbb{R}^d} \|V_g f(\cdot, \omega)\|_{L_{\alpha}^{s-r}(\mathbb{R}^d)}^{s-r} \left[\int_{\mathbb{R}^d} |V_g f(x, \omega)|^{r-2} \mathcal{R} \left(V_g f(x, \omega) \overline{V_g u(x, \omega)} \right) \langle x \rangle^{\alpha r} dx \right] \langle \omega \rangle^{\beta s} d\omega + \\
& + \mathcal{O}(t^2)
\end{aligned}$$

where $\|V_g f(\cdot, \omega)\|_{L_{\alpha}^r(\mathbb{R}^d)}$ is as in eq.(6.3).

The stationarity condition for (6.1) is obtained from the Fréchet derivative:

$$\begin{aligned}
(6.7) \quad & 0 = \lim_{t \rightarrow 0} \frac{1}{t} \left(\mathfrak{L}_{a,b}^{p,q}[f_0 + tu, \lambda] - \mathfrak{L}_{a,b}^{p,q}[f_0, \lambda] \right) \\
& \Leftrightarrow \| |x|^a f_0 \|_{L^p(\mathbb{R}^d)}^{1-p} \int_{\mathbb{R}^d} |x|^{ap} |f_0(x)|^{p-2} \mathcal{R} \left(f_0(x) \overline{u(x)} \right) dx + \\
& + \| |\omega|^b \widehat{f_0} \|_{L^q(\mathbb{R}^d)}^{1-q} \int_{\mathbb{R}^d} |\omega|^{bq} |\widehat{f_0}(\omega)|^{q-2} \mathcal{R} \left(\widehat{f_0}(\omega) \overline{\widehat{u}(\omega)} \right) d\omega = \\
& = \lambda_0 \int_{\mathbb{R}^d} \|V_g f_0(\cdot, \omega)\|_{L_{\alpha}^{s-r}(\mathbb{R}^d)}^{s-r} \left[\int_{\mathbb{R}^d} |V_g f_0(x, \omega)|^{r-2} \mathcal{R} \left(V_g f_0(x, \omega) \overline{V_g u(x, \omega)} \right) \langle x \rangle^{\alpha r} dx \right] \langle \omega \rangle^{\beta s} d\omega,
\end{aligned}$$

where we used the constraint $\|f_0\|_{M_{\alpha,\beta}^{r,s}(\mathbb{R}^d)} = 1$. The result then extends to $L_a^p(\mathbb{R}^d) \cap \mathcal{F}L_b^q(\mathbb{R}^d)$ by density.

Since (6.7) holds for all $u \in L_a^p(\mathbb{R}^d) \cap \mathcal{F}L_b^q(\mathbb{R}^d)$, then we get in particular for $u = f_0$:

$$(6.8) \quad \lambda_0 = \| |x|^a f_0 \|_{L^p(\mathbb{R}^d)} + \| |\omega|^b \widehat{f_0} \|_{L^q(\mathbb{R}^d)} = \mathfrak{L}_{a,b}^{p,q}[f_0].$$

If f_1 is some other solution of (6.7) with λ_0 replaced by λ_1 and f_0 is the minimizer, we would get:

$$(6.9) \quad \lambda_0 = \mathfrak{L}_{a,b}^{p,q}[f_0] \leq \mathfrak{L}_{a,b}^{p,q}[f_1] = \lambda_1.$$

Finally, if we multiply (6.7) by i and add it to the same equation with u replaced by $-iu$, we obtain (6.2). \square

6.2. The Hilbert case. We want to minimize the functional

$$\mathfrak{F}^{\psi,\phi,m_0}[f] = \|\psi f\|_{L^2(\mathbb{R}^d)}^2 + \|\phi \widehat{f}\|_{L^2(\mathbb{R}^d)}^2 = \|f\|_{\psi,\phi,m_0}^2 - \|f\|_{M_{m_0}^2(\mathbb{R}^d)}^2$$

in $M_m^2(\mathbb{R}^d)$, subject to the constraint:

$$(6.10) \quad \|f\|_{M_{m_0}^2(\mathbb{R}^d)} = 1.$$

We thus optimize the functional

$$(6.11) \quad \mathfrak{L}^{\psi,\phi,m_0}[f, \lambda] = \mathfrak{F}^{\psi,\phi,m_0}[f] + \lambda \left(1 - \|f\|_{M_{m_0}^2(\mathbb{R}^d)}^2 \right),$$

where λ is a Lagrange multiplier. In this section we shall always think of $M_m^2(\mathbb{R}^d)$ as $\mathcal{B}^{\psi,\phi,m_0}(\mathbb{R}^d)$, with the inner product $(\cdot, \cdot)_{\psi,\phi,m_0}$ and the norm $\|\cdot\|_{\psi,\phi,m_0}$.

Before we proceed, let us recall that the operator A defined in eq. (5.10) is such that:

$$(6.12) \quad (u, v)_{M_{m_0}^2(\mathbb{R}^d)} = (Au, v)_{\psi, \phi, m_0},$$

for all $u, v \in M_m^2(\mathbb{R}^d) = \mathcal{B}^{\psi, \phi, m_0}(\mathbb{R}^d)$.

Theorem 14. *The operator A is positive-definite, compact and closed. It has empty residual spectrum, 0 belongs to the continuous spectrum and it is a point of accumulation. Moreover, all remaining spectral values are eigenvalues.*

Proof. From the definition of A , we have for all $u, v \in M_m^2(\mathbb{R}^d)$:

$$(6.13) \quad (Au, v)_{\psi, \phi, m_0} = (u, v)_{M_{m_0}^2(\mathbb{R}^d)} = \overline{(v, u)}_{M_{m_0}^2(\mathbb{R}^d)} = \overline{(Av, u)}_{\psi, \phi, m_0} = (u, Av)_{\psi, \phi, m_0}.$$

Hence $A = A^*$.

Similarly

$$(6.14) \quad (Au, u)_{\psi, \phi, m_0} = \|u\|_{M_{m_0}^2(\mathbb{R}^d)}^2 > 0,$$

for all $u \in M_m^2(\mathbb{R}^d) \setminus \{0\}$. Consequently, A is positive-definite.

That A is closed is a simple consequence of the fact that it is bounded and defined on the whole of $M_m^2(\mathbb{R}^d)$.

Next, we prove compactness. Let $(u_n)_{n \in \mathbb{N}}$ be a bounded sequence in $M_m^2(\mathbb{R}^d)$:

$$(6.15) \quad \|u_n\|_{\psi, \phi, m_0} \leq C$$

for some constant $C > 0$ and all $n \in \mathbb{N}$. Since $M_m^2(\mathbb{R}^d) \subset\subset M_{m_0}^2(\mathbb{R}^d)$, $(u_n)_{n \in \mathbb{N}}$ has a subsequence $(v_n)_{n \in \mathbb{N}}$ which converges in $M_{m_0}^2(\mathbb{R}^d)$. It follows that

$$\begin{aligned} \|Av_n - Av_m\|_{\psi, \phi, m_0}^2 &= (v_n - v_m, A(v_n - v_m))_{M_{m_0}^2(\mathbb{R}^d)} \leq \\ &\leq \|A(v_n - v_m)\|_{M_{m_0}^2(\mathbb{R}^d)} \|v_n - v_m\|_{M_{m_0}^2(\mathbb{R}^d)} \leq \\ (6.16) \quad &\leq \|A(v_n - v_m)\|_{M_m^2(\mathbb{R}^d)} \|v_n - v_m\|_{M_{m_0}^2(\mathbb{R}^d)} \asymp \\ &\asymp \|A(v_n - v_m)\|_{\psi, \phi, m_0} \|v_n - v_m\|_{M_{m_0}^2(\mathbb{R}^d)} \leq \\ &\leq 2C \|A\|_{Op} \|v_n - v_m\|_{M_{m_0}^2(\mathbb{R}^d)}, \end{aligned}$$

where we used (6.15) in the last inequality. This shows that $(Av_n)_{n \in \mathbb{N}}$ is a Cauchy sequence. Since $M_m^2(\mathbb{R}^d)$ is complete, we conclude that $(Av_n)_{n \in \mathbb{N}}$ converges. Since the bounded sequence $(u_n)_{n \in \mathbb{N}}$ was chosen arbitrarily, the operator A is compact.

That all non-zero elements of the spectrum are eigenvalues is an immediate consequence of the fact that A is compact. Eq. (6.14) shows that A is injective. Since A is compact and injective, we conclude that 0 is in the continuous spectrum and the residual spectrum is empty.

If the spectrum of A were finite, then A would have to be of finite rank. But since $A : M_m^2(\mathbb{R}^d) \rightarrow \text{Ran}(A)$ is bijective, this is impossible. Hence, the spectrum is infinite and 0 must be an accumulation point. \square

We are now in a position to obtain the Euler-Lagrange equation for the minimizer f_0 .

Theorem 15. *The minimizer f_0 of $\mathfrak{F}^{\psi,\phi,m_0}$ in $M_m^2(\mathbb{R}^d)$, subject to the constraint (6.10) is a solution of the eigenvalue equation:*

$$(6.17) \quad Af_0 = \frac{1}{\lambda+1}f_0.$$

Moreover, we have that

$$(6.18) \quad \lambda = \mathfrak{F}^{\psi,\phi,m_0}[f_0]$$

is such that $\frac{1}{\lambda+1}$ is the largest eigenvalue of the operator A .

Proof. Notice that we can reexpress the functional (6.11) as:

$$(6.19) \quad \mathfrak{L}^{\psi,\phi,m_0}[f, \lambda] = (f, f)_{\psi,\phi,m_0} - (Af, f)_{\psi,\phi,m_0} + \lambda \left(1 - (Af, f)_{\psi,\phi,m_0} \right),$$

The stationarity condition is:

$$(6.20) \quad f_0 - Af_0 - \lambda Af_0 = 0,$$

which yields (6.17).

If we compute the inner product $(\cdot, \cdot)_{\psi,\phi,m_0}$ of the previous equation with f_0 and use the constraint condition (6.10), we obtain:

$$(6.21) \quad \begin{aligned} & (f_0, f_0)_{\psi,\phi,m_0} - (Af_0, f_0)_{\psi,\phi,m_0} - \lambda (Af_0, f_0)_{\psi,\phi,m_0} = 0 \\ \Leftrightarrow & \|f_0\|_{\psi,\phi,m_0}^2 - \|f_0\|_{M_{m_0}^2(\mathbb{R}^d)}^2 - \lambda \|f_0\|_{M_{m_0}^2(\mathbb{R}^d)}^2 = 0 \\ \Leftrightarrow & \mathfrak{F}^{\psi,\phi,m_0}[f_0] - \lambda = 0, \end{aligned}$$

which is (6.18).

It remains to prove that $\frac{1}{\lambda+1}$ is the largest eigenvalue of A . Let $\mu \in \mathbb{R}$ be some other eigenvalue of A and $v_\mu \in M_m^2(\mathbb{R}^d)$ an eigenvector associated with it, which we assume to be in Ω :

$$(6.22) \quad Av_\mu = \mu v_\mu, \quad \|v_\mu\|_{M_{m_0}^2(\mathbb{R}^d)} = 1.$$

From the previous equation, we conclude that:

$$(6.23) \quad (1 - \mu)Av_\mu = \mu(v_\mu - Av_\mu),$$

and if we compute the inner product of this equation with v_μ , we obtain:

$$(6.24) \quad \begin{aligned} & (1 - \mu)(Av_\mu, v_\mu)_{\psi,\phi,m_0} = \mu \left((v_\mu, v_\mu)_{\psi,\phi,m_0} - (Av_\mu, v_\mu)_{\psi,\phi,m_0} \right) \\ \Leftrightarrow & (1 - \mu)\|v_\mu\|_{M_{m_0}^2(\mathbb{R}^d)}^2 = \mu \left(\|v_\mu\|_{\psi,\phi,m_0}^2 - \|v_\mu\|_{M_{m_0}^2(\mathbb{R}^d)}^2 \right) \\ \Leftrightarrow & 1 - \mu = \mu \mathfrak{F}^{\psi,\phi,m_0}[v_\mu]. \end{aligned}$$

It follows that:

$$(6.25) \quad \mathfrak{F}^{\psi, \phi, m_0} [v_\mu] \geq \mathfrak{F}^{\psi, \phi, m_0} [f_0] \Leftrightarrow \frac{1}{\mu} - 1 \geq \lambda \Leftrightarrow \mu \leq \frac{1}{\lambda + 1},$$

which concludes the proof. \square

It may be more convenient to express the Euler-Lagrange equations in terms of the inverse A^{-1} rather than A .

Theorem 16. A^{-1} is densely defined in $M_m^2(\mathbb{R}^d)$, closed and positive-definite. Its spectrum consists only of eigenvalues which can be written as a sequence $0 < \nu_1 \leq \nu_2 \leq \dots$, with $\nu_j \rightarrow +\infty$. Moreover, all the eigenspaces are finite dimensional.

Proof. We start by proving that $\text{Ran}(A)$ is dense in $M_m^2(\mathbb{R}^d)$. Since $m \in \mathcal{P}(\mathbb{R}^{2d})$, then $\mathcal{S}(\mathbb{R}^d) \subset M_m^2(\mathbb{R}^d)$. Let $u \in M_m^2(\mathbb{R}^d)$ be such that:

$$(6.26) \quad 0 = (Av, u)_{\psi, \phi, m_0} = (v, u)_{M_{m_0}^2(\mathbb{R}^d)},$$

for all $v \in \mathcal{S}(\mathbb{R}^d)$. Since $\mathcal{S}(\mathbb{R}^d)$ is dense in $M_{m_0}^2(\mathbb{R}^d)$, we conclude that $u = 0$, and thus $\{Av : v \in \mathcal{S}(\mathbb{R}^d)\}$ is dense in $\mathcal{B}^{\psi, \phi, m_0}(\mathbb{R}^d) = M_m^2(\mathbb{R}^d)$.

Under these circumstances and taking into account Theorem 14, we conclude that A^{-1} is densely defined, closed and that $(A^{-1})^* = (A^*)^{-1}$. Thus A^{-1} is self-adjoint and positivity follows immediately.

The statements regarding the spectrum are also an immediate consequence of Theorem 14. We just remark that 0 is a regular value of A^{-1} , since its inverse A exists, is bounded and defined on the whole $M_m^2(\mathbb{R}^d)$. \square

Theorem 17. The minimizer f_0 is an eigenvector of A^{-1} associated with the eigenvalue $\lambda + 1 = \mathfrak{F}^{\psi, \phi, m_0} [f_0] + 1$, which is the smallest eigenvalue of A^{-1} .

Proof. The proof follows *mutatis mutandis* that of Theorem 15. \square

Example 1. As a particular example, consider a $d = 1$ system with $M_{m_0}^2(\mathbb{R}) = L^2(\mathbb{R})$ ($m_0 \equiv 1$), $\psi(x)$ and $\phi(\omega) = \omega$ admissible weights.

We thus have, for all $u \in \text{Ran}(A)$, $v \in \mathcal{B}^{\psi, \omega, 1}(\mathbb{R})$:

$$(6.27) \quad (A^{-1}u, v)_{L^2(\mathbb{R})} = (u, v)_{\psi, \omega, 1}$$

$$\Leftrightarrow \int_{\mathbb{R}} (A^{-1}u)(x) \overline{v(x)} dx = \int_{\mathbb{R}} (1 + |\psi(x)|^2) u(x) \overline{v(x)} dx + \int_{\mathbb{R}} |\omega|^2 \widehat{u}(\omega) \overline{\widehat{v}(\omega)} d\omega$$

$$\Leftrightarrow \int_{\mathbb{R}} (A^{-1}u)(x) \overline{v(x)} dx = \int_{\mathbb{R}} (1 + |\psi(x)|^2) u(x) \overline{v(x)} dx + \int_{\mathbb{R}} \mathcal{D}u(x) \overline{\mathcal{D}v(x)} dx,$$

where \mathcal{D} denotes the distributional derivative.

Since $\mathcal{S}(\mathbb{R}) \subset \mathcal{B}^{\psi, \phi, 1}(\mathbb{R})$, we conclude from Theorem 17 and (6.27) that the minimizer f_0 satisfies:

$$(6.28) \quad -\frac{1}{4\pi^2} \mathcal{D}^2 f_0(x) + |\psi(x)|^2 f_0(x) = \lambda f_0(x),$$

with λ the smallest eigenvalue of $A^{-1} - I = -\frac{1}{4\pi^2} \mathcal{D}^2 + |\psi(x)|^2$.

If, in addition, $\psi(x)$ is such that $|\psi(x)|^2 \in C^2(\mathbb{R})$, then a well known theorem shows that the distributional and classical solutions coincide, so

that f_0 is a solution of the time-independent Schrödinger equation

$$(6.29) \quad -\frac{1}{4\pi^2}f_0''(x) + |\psi(x)|^2 f_0(x) = \lambda f_0(x).$$

If $\psi(x) = x$, then all the aforementioned conditions are satisfied, and we obtain the Schrödinger equation for the simple harmonic oscillator:

$$(6.30) \quad -\frac{1}{4\pi^2}f_0''(x) + x^2 f_0(x) = \lambda f_0(x).$$

It is well known that a solution associated with the lowest eigenvalue is:

$$(6.31) \quad f_0(x) = 2^{1/4} e^{-\pi x^2},$$

which corresponds to

$$(6.32) \quad \lambda = \frac{1}{2\pi}.$$

And we thus obtain the inequality

$$(6.33) \quad \|xu\|_{L^2(\mathbb{R})}^2 + \|\omega\widehat{u}\|_{L^2(\mathbb{R})}^2 \geq \frac{1}{2\pi}$$

for all u with $\|u\|_{L^2(\mathbb{R})} = 1$, for which the left-hand side makes sense. This is equivalent to the standard Heisenberg uncertainty principle.

Notice that we have chosen to obtain the minimizer with the inverse operator A^{-1} instead of the compact operator A . The latter alternative would have been much harder to solve [BonTorr, CaRoTo].

ACKNOWLEDGEMENTS

The work of N.C. Dias and J.N. Prata is supported by the Portuguese Science Foundation (FCT) grant PTDC/MAT-CAL/4334/2014.

REFERENCES

- [AF] R. Aceska and H. G. Feichtinger. Functions with variable bandwidth via time-frequency analysis tools. *J. Math. Anal. Appl.*, 382(1):275-289, 2011.
- [Ba] K.I. Babenko. An inequality in the theory of Fourier integrals. *Izv. Akad. Nauk SSSR Ser. Mat.* **25** (1961), 531-542. English trans. Amer. Math. Soc. Transl. (2) **44**, 115-128.
- [Bar] F. Barthe. Optimal Young's inequality and its converse: a simple proof. *Geom. Func. Anal.* **8** (1998), 234-242.
- [BoTo] P. Boggiatto, J. Toft. Embeddings and compactness for generalized Sobolev-Shubin spaces and modulation spaces. *Appl. Anal.* **84** (2005), 269-282.
- [BonTorr] B. Bongioanni and J.-L. Torrea. Sobolev spaces associated to the harmonic oscillator. *Proc. Indian Acad. Sci., Math. Sci.* **116** (2006), 337-360.
- [BrL] H.J. Brascamp, E.H. Lieb. Best constants in Young's inequality, its converse, and its generalization to more than three functions. *Adv. Math.* **20** (1976), 151-173.
- [Be] W. Beckner. Inequalities in Fourier analysis. *Ann. Math.* **102** (6) (1975), 159-182.
- [CaRoTo] M. Cappiello, L. Rodino, J. Toft. On the inverse to the harmonic oscillator. *Comm. Par. Diff. Eq.* **40** (2015) 1096-1118.
- [Co] L. Cohen. Time-frequency distributions—A review, *Proc. IEEE* **77** (1989) 941-981.
- [CoNi] E. Cordero, F. Nicola. Sharp integral bounds for Wigner distributions. *Int. Math. Res. Notices* **2018** (2018) 1779-1807.
- [CoPr] M.G. Cowling, J.F. Price. Bandwidth versus time concentration: the Heisenberg-Pauli-Weyl inequality. *SIAM J. Math. Anal.* **15** (1984) 151-165.
- [Fe81] H.G. Feichtinger. On a new Segal algebra. *Monatshefte Math.* **92** (1981), 269-289.

- [Fe83A] H. G. Feichtinger. A new family of functional spaces on the Euclidean n -space. In Proc.Conf. on Theory of Approximation of Functions, Teor. Priblizh., 1983.
- [Fe83B] H. G. Feichtinger. Modulation spaces on locally compact Abelian groups. Technical report, University of Vienna, January 1983.
- [Fe90] H. G. Feichtinger. Generalized amalgams, with applications to Fourier transform. *Canad. J. Math.* **42**(3) (1990), 395-409.
- [Fe06] H.G. Feichtinger. Modulation spaces: Looking back and ahead. *Sampling Theory in Signal and Image Processing* **5** (2006), 109-140.
- [FeGr] H. G. Feichtinger and K. Gröchenig. Banach spaces related to integrable group representations and their atomic decompositions, II. *Monatsh. Math.* **108** (1989).
- [FeHo90] H. G. Feichtinger and W. Hörmann. Harmonic analysis of generalized stochastic processes on locally compact abelian groups, 1990.
- [FeHo14] H. G. Feichtinger and W. Hörmann. A distributional approach to generalized stochastic processes on locally compact abelian groups. In G. Schmeisser and R. Stens, editors, New Perspectives on Approximation and Sampling Theory. Festschrift in honor of Paul Butzer's 85th birthday, pages 423-446. Cham: Birkhäuser/Springer, 2014.
- [FoSi] G.B. Folland, A. Sitaram, The uncertainty principle: A mathematical survey. *J. Fourier Anal. and Appl.* **3** (1997) 207-238.
- [Ga] Y.V. Galperin. On compactness of embeddings of Fourier-Lebesgue spaces into modulation spaces. *Int. J. Anal.* **2013** (2013), 681573.
- [GG] Y.V. Galperin, K. Gröchenig. Uncertainty principles as embeddings of modulation spaces. *J. Math. Anal. Appl.* **274** (2002), 181-202.
- [Gr01] K. Gröchenig. Foundations of Time-Frequency Analysis. Birkhäuser Boston (2001).
- [Gr96] K. Gröchenig, An uncertainty principle related to the Poisson summation formula, *Stu. Math.* **121** (1996), 87-104.
- [GT1] K. Gröchenig, J. Toft. Isomorphism properties of Toeplitz operators and pseudo-differential operators between modulation spaces. *J. Anal. Math.* **114** (2011), 255-283.
- [GT2] K. Gröchenig, J. Toft. The range of localization operators and lifting theorems for modulation and Bargmann-Fock spaces. *Trans. Amer. Math. Soc.* **365** (2013), 4475-4496.
- [Ho] W. Hörmann. Generalized Stochastic Processes and Wigner Distribution. PhD thesis, University of Vienna, (AUSTRIA), 1989.
- [Le] L. Leindler. On a certain converse of Hölder's inequality. II. *Acta Math. Sci. Szeged* **33** (1972), 217-223.
- [L1] E.H. Lieb. Integral bounds for radar ambiguity functions and Wigner distributions. *J. Math. Phys.* **31** (1990), 594-599.
- [L2] E.H. Lieb. Gaussian kernels have only Gaussian maximizers. *Invent. Math.* **102** (1990), 179-208.
- [LiRo] T. S. Liu and A. C. M. van Rooij. Sums and intersections of normed linear spaces. *Math. Nachr.*, 42(1-3):29-42, 1969.
- [PfTo] C. Pfeuffer, J. Toft, Compactness properties for modulation spaces. *Complex Analysis and Operator Theory* **13** (2019) 3521-3548.
- [Tr83] H. Triebel, Theory of Function Spaces. Basel, Birkhäuser (1983).

N.C. DIAS - GRUPO DE FÍSICA MATEMÁTICA, FACULDADE DE CIÊNCIAS DA UNIVERSIDADE DE LISBOA, CAMPO GRANDE, EDIFÍCIO C6 1749-016 LISBOA, PORTUGAL, AND ESCOLA SUPERIOR NÁUTICA INFANTE D. HENRIQUE, AV. ENGENHEIRO BONNEVILLE FRANCO, 2770-058 PAÇO DE ARCOS, PORTUGAL

Email address: `ncdias@meo.pt`

F. LUEF - DEPARTMENT OF MATHEMATICAL SCIENCES, NTNU TRONDHEIM, 7041 TRONDHEIM, NORWAY

Email address: `franz.luef@ntnu.no`

J.N. PRATA - GRUPO DE FÍSICA MATEMÁTICA, FACULDADE DE CIÊNCIAS DA UNIVERSIDADE DE LISBOA, CAMPO GRANDE, EDIFÍCIO C6 1749-016 LISBOA, PORTUGAL, AND ESCOLA SUPERIOR NÁUTICA INFANTE D. HENRIQUE, AV. ENGENHEIRO BONNEVILLE FRANCO, 2770-058 PAÇO DE ARCOS, PORTUGAL

Email address: joao.prata@mail.telepac.pt