

# SMOOTH FANS THAT ARE ENDPOINT RIGID

RODRIGO HERNÁNDEZ-GUTIÉRREZ AND LOGAN C. HOEHN

*Dedicated to Professor Włodzimierz Charatonik (1957–2021).*

ABSTRACT. Let  $X$  be a smooth fan and denote its set of endpoints by  $E(X)$ . Let  $E$  be one of the following spaces: the natural numbers, the irrational numbers, or the product of the Cantor set with the natural numbers. We prove that there is a smooth fan  $X$  such that  $E(X)$  is homeomorphic to  $E$  and for every homeomorphism  $h: X \rightarrow X$ , the restriction of  $h$  to  $E(X)$  is the identity. On the other hand, we also prove that if  $X$  is any smooth fan such that  $E(X)$  is homeomorphic to complete Erdős space, then  $X$  is necessarily homeomorphic to the Lelek fan; this adds to a 1989 result by Włodzimierz Charatonik.

## 1. INTRODUCTION

Recall that a space  $X$  is *homogeneous* if for every two points  $x, y \in X$  there exists a homeomorphism  $h: X \rightarrow X$  with  $h(x) = y$ . The study of homogeneous spaces in topology has a long history and it is still a relevant topic nowadays. For example, see the survey on homogeneity [3].

Many of the classical examples of spaces are homogeneous; for example: the real line, the circle, the Cantor set, the space of irrational numbers and the Hilbert cube. However, most of the continua studied by continuum theorists are not homogeneous. Indeed, even the arc, which is considered the simplest non-degenerate continuum, is not homogeneous. Nevertheless, even when they are not homogeneous, many of these continua share the common feature that, informally speaking, homeomorphisms can only distinguish a finite number of different points.

To formalize this notion, let  $\mathcal{H}(X)$  denote the group of autohomeomorphisms of the space  $X$ . Notice that  $\mathcal{H}(X)$  acts on  $X$  by  $\langle h, x \rangle \mapsto h(x)$ . The orbit of a point  $x \in X$  under this action is  $\{h(x) : h \in \mathcal{H}(X)\}$ . The *homogeneity degree* of  $X$  is defined to be the number of these orbits. It turns out that many relevant examples of spaces in continuum theory have a finite degree of homogeneity. See the discussion in the introduction of [1] for a comprehensive list of results about continua with low homogeneity degree.

An important class of continua is the class of dendroids. Two interesting subclasses of dendroids are dendrites and smooth fans. In [2] and [1] the authors gave complete lists of all dendrites with homogeneity degree 3 and all smooth fans with homogeneity degree 3 and 4, respectively.

---

*Date:* June 28, 2022.

*2020 Mathematics Subject Classification.* Primary: 54F50; Secondary: 54F15, 54G20, 54F65.

*Key words and phrases.* smooth fan, rigidity, Lelek fan, Erdős space.

In this paper we will go to the exact opposite of homogeneity. Recall that a topological space is called *rigid* if its only autohomeomorphism is the identity. Equivalently,  $X$  is rigid if every orbit is a point. When we say that a homeomorphism is trivial we mean that it is equal to the identity function.

For zero-dimensional separable metric spaces, it known from [13] that an infinite rigid space cannot be Borel (in any Polish space in which it is embedded) and in fact, as shown in [14], the statement “there are infinite zero-dimensional rigid separable metric spaces that are analytic” cannot be decided from the axioms of ZFC. However, for continua, rigidity is not uncommon. Well-known examples of rigid continua are Cook continua [6]. Of course Cook continua are complicated continua but we can make the following observations.

**1. Observation** There exists a rigid dendrite.

To prove Observation 1 one may follow the construction of Wazewski’s universal dendrite: it is possible to construct a dendrite with a dense set of ramification points such that any two ramification points have different orders. Then any autohomeomorphism is trivial when restricted to the dense set of ramification points, so it is trivial.

The next observation concerns smooth fans; see Section 2.1 below for the relevant definitions.

**2. Observation** There is no smooth fan that is rigid.

*Proof.* For a smooth fan  $X$ , consider the set  $E(X)$  of endpoints of  $X$ . If  $E(X)$  is dense in  $X$ , then  $X$  is homeomorphic to the Lelek fan according to [5]; as observed in [1] in this case  $E(X)$  is an uncountable orbit. Now, assume that  $E(X)$  is not dense in  $X$ . We may assume that  $X$  is a subset of the Cantor fan (see [4] and [9]) so it is easy to see that there is an open set  $U \subset X \setminus E(X)$  that is homeomorphic to  $C \times (0, 1)$  where  $C$  is some non-empty, zero-dimensional space. Then it is not difficult to construct a homeomorphism  $h: X \rightarrow X$  that is non-trivial inside  $U$ .  $\square$

So it would seem that there is nothing remarkable about smooth fans and rigidity. However, in this paper we would like to argue that this is not the case. We cannot reach rigidity for smooth fans but we can reach it in the endpoints in the sense that there exist smooth fans  $X$  such that every homeomorphism  $h: X \rightarrow X$  is such that  $h$  is trivial when restricted to the set of endpoints of  $X$ . A smooth fan with this property will be called *endpoint rigid*.

In Section 3, we will produce several examples of endpoint rigid smooth fans, beginning with Proposition 7, in which such fans are constructed whose set of endpoints is homeomorphic to (1) the natural numbers, (2) the irrationals, and (3) the product of the Cantor set with the natural numbers. Notice that each of these three endpoint sets is homogeneous. So even if the set of endpoints of a smooth fan is homogeneous, in some sense this homogeneity is completely lost by the way it is placed inside the smooth fan.

It remains an open problem (see Questions 16, 17 and 18) exactly which spaces may be obtained as the set of endpoints of an endpoint rigid smooth fan. Motivated by this question, we show in Section 5 that the set of endpoints of an endpoint rigid smooth fan cannot be homeomorphic to complete Erdős space (defined in Section

4 below). In fact, we prove in Theorem 19 below that the Lelek fan is the only smooth fan whose set of endpoints is homeomorphic to complete Erdős space, thus obtaining a new characterization of the Lelek fan (see Corollary 21).

## 2. PRELIMINARIES

All the spaces considered in this paper are separable and metric. A *continuum* is a compact, connected metric space. A continuum is *non-degenerate* if it contains more than one point. An *arc* is a continuum which is homeomorphic to the interval  $[0, 1]$ .

We denote the natural numbers by  $\omega = \{0, 1, 2, \dots\}$ . Given a point  $x$  in a continuum  $X$  and a natural number  $n \in \omega \setminus \{0\}$ , we say the *order* of  $x$  is at least  $n$  if there exist  $n$  arcs  $A_1, \dots, A_n \subset X$ , all of which have  $x$  as one endpoint, and which are otherwise pairwise disjoint. We say  $x$  has order (exactly)  $n$  if the order of  $x$  is at least  $n$  but not at least  $n + 1$ .

Suppose  $X$  is a continuum in which for any two distinct points  $x, y \in X$ , there is a unique arc in  $X$  whose endpoints are  $x$  and  $y$ . In such a case, we denote this arc by  $xy$ . In particular, we will use this notation with (smooth) fans throughout this paper.

**2.1. Smooth fans.** Let  $C$  denote the Cantor set. The *Cantor fan*  $F_C$  is defined to be the cone over the Cantor set. A *smooth fan* is any non-degenerate continuum which is homeomorphic to a subcontinuum of the Cantor fan (see [4] and [9]). We will always assume that an embedding of a smooth fan  $X$  in the Cantor fan  $F_C$  contains the cone point of  $F_C$ , and that this point has order at least 2 in  $X$ . The point of  $X$  corresponding to the cone point of  $F_C$  is called the *vertex* of  $X$ . It is uniquely defined if it has order at least 3. A point  $e \in X$  is an *endpoint* if it has order 1. We denote by  $E(X)$  the set of all endpoints of  $X$ .

The *Lelek fan*, constructed in [12], is the unique smooth fan  $X$  with the property that  $E(X)$  is dense in  $X$  [5] (see also Corollary 21 below).

**2.2. USC functions.** For the sake of explicit constructions of smooth fans, it will be convenient to make use of the language of upper semi-continuous functions defined on the Cantor set. Recall that a function  $\varphi: C \rightarrow [0, \infty)$  is *upper semi-continuous* (USC) if for every  $t \in (0, \infty)$  the set  $\varphi^{-1}((-\infty, t])$  is open. Following [8], we define

$$G_0^\varphi = \{\langle x, \varphi(x) \rangle : x \in C, \varphi(x) > 0\}, \text{ and}$$

$$L_0^\varphi = \{\langle x, y \rangle : x \in C, 0 \leq y \leq \varphi(x)\}.$$

Observe that  $\varphi$  is USC if and only if  $L_0^\varphi$  is closed in  $C \times \mathbb{R}$ .

A USC function  $\varphi: C \rightarrow [0, \infty)$  will be called *non-degenerate* if there are at least two distinct points  $x_1, x_2 \in C$  such that  $\varphi(x_1) > 0$  and  $\varphi(x_2) > 0$ .

**3. Lemma** Given any non-degenerate USC function  $\varphi: C \rightarrow [0, \infty)$  defined on the Cantor set  $C$ , the space  $L_0^\varphi / (C \times \{0\})$  is a smooth fan whose endpoint set is the image of  $G_0^\varphi$  under the quotient projection. Conversely, given any smooth fan  $X$ , there exists a non-degenerate USC function  $\varphi: C \rightarrow [0, 1]$  defined on the Cantor set  $C$  such that  $L_0^\varphi / (C \times \{0\}) \approx X$  and  $G_0^\varphi \approx E(X)$ .

*Proof.* The forward direction is clear, and is left to the reader.

For the converse, let us construct an explicit embedding of the Cantor fan  $F_C$  in  $\mathbb{R}^2$ : namely, fix a Cantor set  $C \subset \mathbb{R}$ , let  $p = \langle 0, 0 \rangle$ , and let  $F_C$  be the union of all straight-line segments from  $p$  to points of  $C \times \{1\}$ .

Let  $X$  be a smooth fan, and assume that  $X \subseteq F_C$ , and that  $p$  is the vertex of  $X$  (and the order of  $p$  in  $X$  is at least 2). Define  $\varphi: C \rightarrow [0, 1]$  as follows: given  $x \in C$ , let  $\varphi(x)$  be the maximum  $y$ -coordinate of a point of  $X$  on the straight-line segment from  $p$  to  $\langle x, 1 \rangle$ . We leave it to the reader to confirm that this function  $\varphi$  satisfies the desired properties.  $\square$

### 3. ENDPOINT RIGID FANS

**4. Definition** A smooth fan  $X$  will be called *endpoint rigid* if for each homeomorphism  $h: X \rightarrow X$ ,  $h(x) = x$  for all  $x \in E(X)$ .

In order to construct endpoint rigid smooth fans, we will make use of the notion of a  $\mathcal{K}$ -fan, defined next.

Two subsets  $A_1, A_2$  of  $(0, 1)$  will be called *order isomorphic* if there is an order preserving autohomeomorphism  $h: [0, 1] \rightarrow [0, 1]$  such that  $h(A_1) = A_2$ .

**5. Definition** Let  $\mathcal{K} = \{K_n : n \in \omega\}$  be a countably infinite set of compact subsets of  $(0, 1)$ , no two of which are order isomorphic. A smooth fan  $X$  with vertex  $p$  is a  $\mathcal{K}$ -fan if there exists a subset  $E_{\mathcal{K}} = \{e_n : n \in \omega\} \subseteq E(X)$ , such that:

- (i) For each  $n \in \omega$ ,  $\overline{E(X)} \cap pe_n \setminus \{p, e_n\}$  is order isomorphic to  $K_n$ ;
- (ii) For each  $e \in E(X) \setminus E_{\mathcal{K}}$ ,  $\overline{E(X)} \cap pe \setminus \{p, e\}$  is not order isomorphic to any element of  $\mathcal{K}$ ; and
- (iii) For each  $e \in E(X) \setminus E_{\mathcal{K}}$ ,  $\overline{E_{\mathcal{K}}} \cap pe \setminus \{p\} \neq \emptyset$ .

**6. Proposition** Let  $\mathcal{K}$  be a collection as in Definition 5. Any  $\mathcal{K}$ -fan is endpoint rigid.

*Proof.* Let  $X$  be a  $\mathcal{K}$ -fan with vertex  $p$ , and let  $E_{\mathcal{K}} = \{e_n : n \in \omega\}$  be as in Definition 5. Let  $h: X \rightarrow X$  be a homeomorphism. It follows from (i) and (ii) of Definition 5 that  $E_{\mathcal{K}}$  is invariant under  $h$ . Now, since no two of the spaces in  $\mathcal{K}$  are order isomorphic, by (i) of Definition 5 we have that  $h$  is the identity on  $E_{\mathcal{K}}$ . From (iii) of Definition 5, we conclude that  $h$  fixes at least one point other than  $p$  on each leg of  $X$ , hence it is the identity on  $E(X)$ . Thus  $X$  is endpoint rigid.  $\square$

**7. Proposition** Let  $\mathcal{K}$  be a collection as in Definition 5. There exists a  $\mathcal{K}$ -fan  $X$ . Moreover, if  $E_{\mathcal{K}} = \{e_n : n \in \omega\}$  is as in Definition 5 and  $p$  is the vertex of  $X$ , then  $X$  can be constructed so that either:

- (1) (Assuming  $\emptyset \notin \mathcal{K}$ )  $E(X)$  is homeomorphic to the natural numbers and  $E(X) = E_{\mathcal{K}}$ ; or
- (2)  $\overline{E(X)}$  is homeomorphic to the irrationals, and for each  $e \in E(X) \setminus E_{\mathcal{K}}$ ,  $\overline{E(X)} \cap pe = \{e\}$ ; or
- (3)  $E(X)$  is homeomorphic to the product of the Cantor set with the natural numbers, and for each  $e \in E(X) \setminus E_{\mathcal{K}}$ ,  $\overline{E(X)} \cap pe = \{p, e\}$ .




FIGURE 1. An illustration of a few of the points  $x_s$  and intervals  $I_s$  from the proof of Proposition 7.

*Proof.* To begin, we recursively construct a particular Cantor set  $C \subset [0, 1]$ . The set  $C$  will be indexed by the set  $\omega^{\leq \omega}$  of countable sequences of natural numbers (possibly finite or even empty).

Recall that  $\omega^{\leq \omega} = \bigcup \{\omega^\alpha : \alpha \leq \omega\}$ , that is, the set of functions with domain either a natural number or  $\omega$ , and image contained in  $\omega$ . If  $s \in \omega^\alpha$  then  $\alpha$  is the *length* of  $s$  and will be denoted by  $|s|$ . Given  $s \in \omega^{\leq \omega}$  and  $i < |s|$  then the  $i$ -th term of  $s$  is  $s(i)$ . If  $s \in \omega^{\leq \omega}$  and  $n \leq |s|$  then  $s \upharpoonright n$  denotes the restriction of  $s$  to  $n$ . We will also use the notation  $\omega^{< \omega} = \bigcup \{\omega^n : n < \omega\}$ . If  $n < \omega$ ,  $s \in \omega^n$  and  $i \in \omega$  then  $s \frown i$  denotes the sequence such that  $(s \frown i) \upharpoonright n = s$  and  $(s \frown i)(n) = i$ . Given  $s \in \omega^{< \omega}$ , its *immediate successors* are the sequences  $s \frown i$  for  $i \in \omega$ .

We start by recursively defining a countable sequence of points  $\{x_s : s \in \omega^{< \omega}\} \subset [0, 1]$  and a countable sequence of closed intervals  $\{I_s : s \in \omega^{< \omega}\}$  with the following properties:

- (i)  $x_\emptyset = 0$  and  $I_\emptyset = [0, 1]$ ;
- (ii) For every  $s \in \omega^{< \omega}$ ,  $\{x_{s \frown n} : n \in \omega\}$  is a decreasing sequence of real numbers contained in  $\text{int}_{\mathbb{R}}(I_s)$  converging to  $x_s$ ; and
- (iii) Given  $s \in \omega^{< \omega}$ , assume that  $I_s = [x_s, b]$ . Then  $I_{s \frown 0} = [x_{s \frown 0}, \frac{1}{2}(x_{s \frown 0} + b)]$  and  $I_{s \frown i} = [x_{s \frown (i+1)}, \frac{1}{2}(x_{s \frown (i+1)} + x_{s \frown i})]$  for every  $i \in \omega$ .

See Figure 1 for an illustration of an example of such points  $x_s$  and intervals  $I_s$  as described above.

Now, let  $f \in \omega^\omega$ . Notice that the sequence  $\{x_{f \upharpoonright n} : n \in \omega\}$  is an increasing sequence bounded by 1; let  $x_f$  be the limit of this sequence. So define  $C = \{x_s : s \in \omega^{\leq \omega}\}$ . It should be clear that  $C$  is a Cantor set.

We make two remarks regarding convergence in  $C$ : given a sequence  $s_1, s_2, \dots$  in  $\omega^{\leq \omega}$ ,

- (a)  $x_{s_k} \rightarrow x_f$  for some  $f \in \omega^\omega$  if and only if  $\max\{n : f \upharpoonright n = s_k \upharpoonright n\} \rightarrow \infty$  as  $k \rightarrow \infty$ ; and
- (b)  $x_{s_k} \rightarrow x_s$  for some  $s \in \omega^{< \omega}$  if and only if for all but finitely many  $k$ , either  $s_k = s$ , or  $s_k$  extends  $s$ , and for such  $k$  in the latter case,  $s_k(|s|) \rightarrow \infty$  as  $k \rightarrow \infty$ .

Fix a bijection  $\theta: \omega^{< \omega} \rightarrow \omega$  with the property that

- (c)  $\theta(s_1) < \theta(s_2)$  whenever  $s_2$  extends  $s_1$ .

It follows that:

- (d) If  $s_1, s_2, \dots \in \omega^{< \omega}$  are pairwise distinct, then  $\theta(s_k) \rightarrow \infty$  as  $k \rightarrow \infty$ .

We next define an upper semi-continuous function  $\varphi: C \rightarrow [0, 1]$ . We begin by recursively defining  $\varphi(x_s)$  for each  $s \in \omega^{< \omega}$ . Let  $\varphi(x_\emptyset) = 1$ .

Let  $s \in \omega^{< \omega}$ , and suppose  $\varphi(x_s) > 0$  has been defined. We choose a set  $Y_s \subset (0, \varphi(x_s))$  which is an order isomorphic copy of  $K_{\theta(s)}$ , and choose values  $\varphi(x_{s \frown i})$

for  $i < \omega$ , with details depending on which of the properties (1), (2), or (3) we wish to achieve:

- For (1), we choose  $Y_s \subset (0, \min\{\varphi(x_s), 2^{-\theta(s)}\})$ , and for  $i = 0, 1, \dots$  we choose values  $\varphi(x_{s \smallfrown i}) \in Y_s$  in such a way that  $\overline{\{\varphi(x_{s \smallfrown i}) : i \geq n\}} = Y_s$  for each  $n \in \omega$ .
- For (2), we choose  $Y_s \subset (\max\{0, \varphi(x_s) - 2^{-\theta(s)}\}, \varphi(x_s))$ , and for  $i = 0, 1, \dots$  we choose values  $\varphi(x_{s \smallfrown i}) \in Y_s \cup \{\varphi(x_s)\}$  in such a way that  $\overline{\{\varphi(x_{s \smallfrown i}) : i \geq n\}} = Y_s \cup \{\varphi(x_s)\}$  for each  $n \in \omega$ .
- For (3), we choose  $Y_s \subset (0, \min\{\varphi(x_s), 2^{-\theta(s)}\})$ , and for  $i = 0, 1, \dots$  we choose values  $\varphi(x_{s \smallfrown i}) \in Y_s \cup \{\varphi(x_s)\}$  in such a way that  $\overline{\{\varphi(x_{s \smallfrown i}) : i \geq n\}} = Y_s \cup \{\varphi(x_s)\}$  for each  $n \in \omega$ .

We remark that by construction,

- (e)  $\varphi(x_{s_1}) \geq \varphi(x_{s_2})$  whenever  $s_2$  extends  $s_1$ .

We make further observations about  $\varphi$  depending on the variant (1), (2), or (3) of the construction:

- (f1) For (1), for any  $\emptyset \neq s \in \omega^{<\omega}$ ,  $\varphi(x_s) < 2^{-\theta(s||s|-1)}$ .

This holds because for any  $s \in \omega^{<\omega}$  and any  $i \in \omega$ ,  $\varphi(x_{s \smallfrown i}) \in Y_s$ , hence by choice of  $Y_s$ ,  $\varphi(x_{s \smallfrown i}) < 2^{-\theta(s)}$ .

- (f2) For (2), for any  $s_1, s_2 \in \omega^{<\omega}$  such that  $s_2$  extends  $s_1$ ,  $\varphi(x_{s_1}) - \varphi(x_{s_2}) < 2^{-\theta(s_1)+1}$ .

To prove this, note that for any  $s \in \omega^{<\omega}$  and any  $i \in \omega$ ,  $\varphi(x_{s \smallfrown i}) \in Y_s \cup \{\varphi(x_s)\}$ , hence by choice of  $Y_s$ ,  $\varphi(x_{s \smallfrown i}) > \varphi(x_s) - 2^{-\theta(s)}$ . So

$$\begin{aligned} \varphi(x_{s_1}) - \varphi(x_{s_2}) &= \sum_{n=|s_1|}^{|s_2|-1} \varphi(x_{s_2 \upharpoonright n}) - \varphi(x_{s_2 \upharpoonright n+1}) \\ &< \sum_{n=|s_1|}^{|s_2|-1} 2^{-\theta(s_2 \upharpoonright n)} \\ &\leq \sum_{m=0}^{|s_2|-|s_1|-1} 2^{-\theta(s_1)-m} \quad \text{by (c)} \\ &< 2^{-\theta(s_1)+1}. \end{aligned}$$

- (f3) For (3), for any  $\emptyset \neq s \in \omega^{<\omega}$ , either  $\varphi(x_s) < 2^{-\theta(s||s|-1)}$  or  $\varphi(x_s) = \varphi(x_{s||s|-1})$ .

This holds because for any  $s \in \omega^{<\omega}$  and any  $i \in \omega$ , either  $\varphi(x_{s \smallfrown i}) \in Y_s$  or  $\varphi(x_{s \smallfrown i}) = \varphi(x_s)$ . In the former case, the estimate follows by the choice of  $Y_s$  as in (1).

Finally, for  $f \in \omega^\omega$ , define  $\varphi(x_f) = \lim_{n \rightarrow \infty} \varphi(x_{f \upharpoonright n})$ ; this limit exists by (e).

**7.1. Claim**  $\varphi$  is upper semi-continuous.

*Proof of Claim 7.1.* Suppose  $s_1, s_2, \dots \in \omega^{<\omega}$  are such that  $x_{s_k}$  converges in  $C$  and  $\varphi(x_{s_k})$  converges to  $y \in [0, 1]$ .

If  $x_{s_k} \rightarrow x_f$  for some  $f \in \omega^\omega$ , then for each  $k$  let  $n(k)$  be maximal such that  $s_k \upharpoonright n(k) = f \upharpoonright n(k)$ . Then  $n(k) \rightarrow \infty$  as  $k \rightarrow \infty$ . By (e) we have  $\varphi(x_{s_k}) \leq \varphi(x_{s_k \upharpoonright n(k)}) = \varphi(x_f \upharpoonright n(k))$ . The latter quantity converges to  $\varphi(x_f)$  by definition of  $\varphi$ . Thus  $y \leq \varphi(x_f)$  in this case.

If  $x_{s_k} \rightarrow x_s$  for some  $s \in \omega^{<\omega}$ , then for all but finitely many  $k$ , either  $s_k = s$  or  $s_k$  extends  $s$ , and for all such  $k$  we have  $\varphi(x_{s_k}) \leq \varphi(x_s)$  by (e). Thus  $y \leq \varphi(x_s)$  in this case as well.  $\square$ (Claim 7.1)

Let  $X = L_0^\varphi / (C \times \{0\})$ , and let  $p$  represent the point in the quotient corresponding to  $C \times \{0\}$ . Then  $X$  is a smooth fan with vertex  $p$ .

For points  $\langle x_s, y \rangle \in L_0^\varphi$  with  $y > 0$ , for convenience we identify  $\langle x_s, y \rangle$  with its image in  $X$  under the quotient projection. The set of endpoints  $E(X)$  is thus the set  $\{\langle x_s, \varphi(x_s) \rangle : s \in \omega^{\leq \omega} \text{ and } \varphi(s) > 0\}$ , and given any endpoint  $e = \langle x_s, \varphi(x_s) \rangle$  in  $X$ , the set  $pe \setminus \{p\}$  is identified with  $\{x_s\} \times (0, \varphi(x_s)]$ .

For each  $s \in \omega^{<\omega}$ , let  $e_{\theta(s)} \in X$  be the endpoint  $\langle x_s, \varphi(x_s) \rangle$ , and let  $E_{\mathcal{K}} = \{e_{\theta(s)} : s \in \omega^{<\omega}\}$ . For any  $f \in \omega^\omega$  such that  $\varphi(x_f) > 0$ , it is immediate from the definition of  $\varphi(x_f)$  that the endpoint  $\langle x_f, \varphi(x_f) \rangle$  is in  $\overline{E_{\mathcal{K}}}$ . This already confirms (iii) of Definition 5. Moreover, we have  $E(X) = \overline{E_{\mathcal{K}}}$ .

**7.2. Claim** For each  $s \in \omega^{<\omega}$ ,  $\overline{E_{\mathcal{K}}} \cap [\{x_s\} \times (0, \varphi(x_s))] = \{x_s\} \times Y_s$ .

*Proof of Claim 7.2.* Let  $s \in \omega^{<\omega}$ . It is clear from the construction that  $\{x_s\} \times Y_s \subseteq \overline{\{\langle x_{s \smallfrown i}, \varphi(x_{s \smallfrown i}) \rangle : i \in \omega\}} \subseteq \overline{E_{\mathcal{K}}}$ .

For the reverse inclusion, let  $s_1, s_2, \dots$  belong to  $\omega^{<\omega}$ , and suppose that  $x_{s_k} \rightarrow x_s$  and  $\varphi(x_{s_k}) \rightarrow y$  as  $k \rightarrow \infty$ . We must prove that  $y \in Y_s \cup \{0, \varphi(x_s)\}$ .

We may assume that  $s_k$  extends  $s$  for all  $k$  and  $s_k(|s|) \rightarrow \infty$  as  $k \rightarrow \infty$ . This implies  $\theta(s_k \upharpoonright |s| + 1) \rightarrow \infty$  as  $k \rightarrow \infty$  by (d). If  $|s_k| = |s| + 1$  for infinitely many  $k$ , then  $y \in Y_s \cup \{\varphi(x_s)\}$  by construction. Assume therefore that (without loss of generality)  $|s_k| \geq |s| + 2$  for all  $k$ .

- For (1), observe that by (e), and (f1),

$$\varphi(x_{s_k}) \leq \varphi(x_{s_k \upharpoonright |s| + 2}) < 2^{-\theta(s_k \upharpoonright |s| + 1)} \rightarrow 0 \text{ as } k \rightarrow \infty.$$

Therefore  $y = 0$ .

- For (2), by (f2) we have  $\varphi(x_{s_k \upharpoonright |s| + 1}) - \varphi(s_k) < 2^{-\theta(s_k \upharpoonright |s| + 1) + 1}$ . Since  $\varphi(x_{s_k \upharpoonright |s| + 1}) \in Y_s \cup \{\varphi(x_s)\}$  for each  $k$  and  $2^{-\theta(s_k \upharpoonright |s| + 1) + 1} \rightarrow 0$  as  $k \rightarrow \infty$ , we have  $y \in Y_s \cup \{\varphi(x_s)\}$ .
- For (3), if for infinitely many  $k$  there exists  $|s| + 2 \leq n \leq |s_k|$  such that  $\varphi(x_{s_k \upharpoonright n}) \in Y_{s_k \upharpoonright n-1}$ , then similarly as with (1) we have that  $\varphi(x_{s_k}) \rightarrow 0$  as  $k \rightarrow \infty$ , so  $y = 0$ . Assume then that (without loss of generality) for all  $k$ , for each  $|s| + 2 \leq n \leq |s_k|$ ,  $\varphi(x_{s_k \upharpoonright n}) = \varphi(x_{s_k \upharpoonright n-1})$ . It follows that  $\varphi(x_{s_k}) = \varphi(x_{s_k \upharpoonright |s| + 1}) \in Y_s \cup \{\varphi(x_s)\}$  for all  $k$ . Thus  $y \in Y_s \cup \{\varphi(x_s)\}$ .

$\square$ (Claim 7.2)

**7.3. Claim** Let  $f \in \omega^{<\omega}$ .

- For (1),  $\varphi(x_f) = 0$ .
- For (2), if  $\varphi(x_f) > 0$  then  $\overline{E_{\mathcal{K}}} \cap [\{x_f\} \times [0, \varphi(x_s)]] = \{\langle x_f, \varphi(x_f) \rangle\}$ .
- For (3), if  $\varphi(x_f) > 0$  then  $\overline{E_{\mathcal{K}}} \cap [\{x_f\} \times [0, \varphi(x_s)]] = \{\langle x_f, 0 \rangle, \langle x_f, \varphi(x_f) \rangle\}$ .

*Proof of Claim 7.3.*

- For (1), observe that for each  $n \geq 1$ , by construction  $\varphi(x_{f \upharpoonright n}) \in Y_{f \upharpoonright n-1} \subset (0, 2^{-\theta(f \upharpoonright n-1)})$ . Hence  $\varphi(x_f) = \lim_{n \rightarrow \infty} \varphi(x_{f \upharpoonright n}) = 0$  because  $\theta(f \upharpoonright n-1) \rightarrow \infty$  as  $n \rightarrow \infty$  by (d).

For (2) and (3), assume now that  $\varphi(x_f) > 0$ . We have already shown above that  $\langle x_f, \varphi(x_f) \rangle$  is in  $\overline{E_{\mathcal{K}}}$ .

- For (3), we also need to show that  $\langle x_f, 0 \rangle \in \overline{E_{\mathcal{K}}}$ . To see this, we construct a sequence as follows: for each  $n \in \omega$ , choose  $i(n) \in \omega$  such that  $\varphi(x_{(f \upharpoonright n) \smallfrown i(n)}) \in Y_{f \upharpoonright n} \subseteq (0, 2^{-\theta(f \upharpoonright n)})$ . Observe that  $x_{(f \upharpoonright n) \smallfrown i(n)} \rightarrow x_f$  as  $n \rightarrow \infty$ . Since  $\theta(f \upharpoonright n) \rightarrow \infty$  as  $n \rightarrow \infty$  by (d), it follows that  $\langle x_f, 0 \rangle \in \overline{\{\langle x_{(f \upharpoonright n) \smallfrown i(n)}, \varphi(x_{(f \upharpoonright n) \smallfrown i(n)}) \rangle : n \in \omega\}} \subseteq \overline{E_{\mathcal{K}}}$ .

For the reverse inclusions, suppose  $s_1, s_2, \dots$  belong to  $\omega^{<\omega}$ , and suppose that  $x_{s_k} \rightarrow x_f$  and  $\varphi(x_{s_k}) \rightarrow y$  as  $k \rightarrow \infty$ . For each  $k$ , let  $n(k) = \max\{n \in \omega : f \upharpoonright n = s_k \upharpoonright n\}$ . Then  $n(k) \rightarrow \infty$  as  $k \rightarrow \infty$ .

- For (2) we must prove that  $y = \varphi(x_f)$ . By (d) and (f2),  $\varphi(x_{f \upharpoonright n(k)}) - \varphi(x_{s_k}) < 2^{-\theta(f \upharpoonright n(k))+1} \rightarrow 0$  as  $k \rightarrow \infty$ . Thus

$$y = \lim_{k \rightarrow \infty} \varphi(x_{s_k}) = \lim_{k \rightarrow \infty} \varphi(x_{f \upharpoonright n(k)}) = \varphi(x_f).$$

- For (3) we must prove that  $y \in \{0, \varphi(x_f)\}$ . First observe that there must exist some  $n_0 \in \omega$  such that  $\varphi(x_{f \upharpoonright n}) = \varphi(x_{f \upharpoonright n_0})$  for all  $n \geq n_0$ , since otherwise  $\varphi(x_f)$  would be 0 as in (1). We may assume without loss of generality that  $n(k) \geq n_0$  for all  $k$ .

If there are infinitely many  $k$  such that  $\varphi(s_k \upharpoonright m(k)) \in Y_{s_k \upharpoonright m(k)-1}$  for some  $n(k) < m(k) \leq |s_k|$ , then for such  $k$ , by (e) and (f3),  $\varphi(x_{s_k}) \leq \varphi(x_{s_k \upharpoonright m(k)}) < 2^{-\theta(s_k \upharpoonright m(k)-1)+1}$ , and it follows from (d) that  $y = 0$  in this case. Otherwise, we have  $\varphi(s_k) = \varphi(x_{f \upharpoonright n(k)}) = \varphi(x_f)$  for all but finitely many  $k$ , hence  $y = \varphi(x_f)$ .

□(Claim 7.3)

This completes the proof that  $X$  is a  $\mathcal{K}$ -fan in all three constructions (1), (2), and (3). Lastly, we identify the spaces  $E(X)$  in each case:

- For (1), it should be clear from the construction that  $E(X) = E_{\mathcal{K}}$  is a countable discrete space.
- For (2), note that  $E(X)$  is completely metrizable and separable. Any neighborhood of an endpoint contains infinitely many endpoints of the form  $\langle x_s, \varphi(x_s) \rangle$  where  $s \in \omega^{<\omega}$ , and in fact will contain the set  $\{x_s\} \times Y_s$  for all but finitely many of them. Since by construction there are sequences of endpoints converging to these sets  $\{x_s\} \times Y_s$ , we conclude that  $E(X)$  is nowhere locally compact. We leave it to the reader to verify that  $E(X)$  is zero-dimensional. Therefore  $E(X)$  is homeomorphic to the irrational numbers.
- For (3), again note that  $E(X)$  is a separable metric space which is not compact. This time  $E(X)$  is locally compact and has no isolated points, since every endpoint has a closed neighborhood homeomorphic to the Cantor set in which all legs have the same height. Therefore  $E(X)$  is homeomorphic to the product of the Cantor set and the natural numbers.

□



Clearly distinct families  $\mathcal{K}$  lead to non-homeomorphic fans in Proposition 7. Hence we conclude the following:

**8. Corollary** There is a family  $\mathcal{F}$  of size  $\mathfrak{c}$  consisting of pairwise non-homeomorphic endpoint rigid smooth fans. Further, we can choose  $\mathcal{F}$  in such a way that  $E(X)$  is homeomorphic to the natural numbers for every  $X \in \mathcal{F}$ , to the irrationals for every  $X \in \mathcal{F}$ , or to the product of the Cantor set with the natural numbers for every  $X \in \mathcal{F}$ .

Constructions (2) and (3) of Proposition 7 yield examples of smooth fans whose endpoint sets have cardinality  $\mathfrak{c}$ . Hence we have the following:

**9. Corollary** There exists a smooth fan with degree of homogeneity  $\mathfrak{c}$ .

A straightforward modification of the constructions from Proposition 7 yields the next two results.

**10. Proposition** Let  $X$  be a smooth fan with vertex  $p$ . Let  $\mathcal{K}$  be a collection as in Definition 5. There exists a  $\mathcal{K}$ -fan  $X'$  such that  $X$  embeds in  $X'$ , and (treating  $X$  as a subset of  $X'$ )  $E(X) \subset E(X')$  and  $X \subset \overline{E(X') \setminus E(X)}$ . Moreover, if  $E_{\mathcal{K}} = \{e_n : n \in \omega\}$  is as in Definition 5, then  $X'$  can be constructed so that either:

- (1)  $E(X') \setminus E(X)$  is homeomorphic to the natural numbers and  $E(X') \setminus E(X) = E_{\mathcal{K}}$ ; or
- (2)  $E(X') \setminus E(X)$  is homeomorphic to the irrationals; or
- (3)  $E(X') \setminus E(X)$  is homeomorphic to the product of the Cantor set with the natural numbers.

**11. Proposition** Let  $X$  be a smooth fan with vertex  $p$ , and assume that  $p \notin \overline{E(X)}$ . Let  $\mathcal{K}$  be a collection as in Definition 5. There exists a  $\mathcal{K}$ -fan  $X'$  such that  $X$  embeds in  $X'$ , and (treating  $X$  as a subset of  $X'$ )  $E(X)$  is a clopen subset of  $E(X')$ . Moreover, if  $E_{\mathcal{K}} = \{e_n : n \in \omega\}$  is as in Definition 5, then  $X'$  can be constructed so that either:

- (1)  $E(X') \setminus E(X)$  is homeomorphic to the natural numbers and  $E(X') \setminus E(X) = E_{\mathcal{K}}$ ; or
- (2)  $E(X') \setminus E(X)$  is homeomorphic to the irrationals; or
- (3)  $E(X') \setminus E(X)$  is homeomorphic to the product of the Cantor set with the natural numbers.

To obtain these last two results, one can start with the fan  $X$ , then add more legs, one for each  $s \in \omega^{<\omega} \setminus \{\emptyset\}$ , starting with a sequence of legs indexed by the sequences in  $\omega^{<\omega}$  of length one, constructed in such a way that they converge to  $X$  (for Proposition 10) or to a subset of  $X$  disjoint from  $E(X)$  and whose intersection with each leg is not order isomorphic to any element of  $\mathcal{K}$  (for Proposition 11), then proceed with the rest of the construction as in Proposition 7.

We next introduce one simple construction on fans which we can use to generate more endpoint rigid fans.

**12. Definition** Given finitely many smooth fans  $X_1, \dots, X_m$  whose vertices are  $p_1, \dots, p_m$ , the *sum* of these fans is

$$\sum_{i=1}^m X_i = \left( \bigsqcup_{i=1}^m X_i \right) / \{p_1, \dots, p_m\}.$$

Given countably infinitely many smooth fans  $X_1, X_2, \dots$  with vertices  $p_1, p_2, \dots$ , consider the one point compactification  $\bigsqcup_{i=1}^{\infty} X_i \cup \{\infty\}$  of  $\bigsqcup_{i=1}^{\infty} X_i$ . The *sum* of these fans is

$$\sum_{i=1}^{\infty} X_i = \left( \bigsqcup_{i=1}^{\infty} X_i \cup \{\infty\} \right) / (\{p_1, p_2, \dots\} \cup \{\infty\}).$$

In both cases, observe that  $\sum_i X_i$  is a smooth fan, and  $E(\sum_i X_i) \approx \bigsqcup_i E(X_i)$ .

**13. Proposition** Suppose that  $\mathcal{K}_1, \mathcal{K}_2, \dots$  is a finite or countably infinite sequence of collections as in Definition 5, and suppose that no two of these collections contain sets which are order isomorphic to each other. For each  $i$ , let  $X_i$  be a  $\mathcal{K}_i$ -fan, and let  $\mathcal{K} = \bigcup_i \mathcal{K}_i$ . Then  $\sum_i X_i$  is a  $\mathcal{K}$ -fan.

#### 4. ENDPOINT SETS IN SMOOTH FANS

In this section we investigate which spaces may be realized as the set of endpoints of a smooth fan, or of an endpoint rigid smooth fan. To start with, it is known from [12, Theorem 7.5] that the set of endpoints of any smooth fan is a Polish space.

*Complete Erdős space* is the subspace

$$\mathfrak{E}_c = \{ \langle x_n \rangle_{n \in \omega} \in \ell^2 : \forall i \in \omega, x_i \in \{0\} \cup \{1/n : n \in \mathbb{N}\} \}$$

of  $\ell^2$ , where  $\ell^2$  is the Hilbert space of square-summable sequences of real numbers. Complete Erdős space was introduced by Erdős in 1940 in [10] as an example of totally disconnected and non-zero-dimensional space. In [11] it was proved that the Lelek fan has its set of endpoints homeomorphic to  $\mathfrak{E}_c$ .

**14. Lemma** For every smooth fan  $X$  the set  $E(X)$  is homeomorphic to a closed subset of  $\mathfrak{E}_c$ .

*Proof.* Let  $X$  be a smooth fan. By Lemma 3,  $E(X) \approx G_0^\varphi$  for some USC function  $\varphi: C \rightarrow [0, 1]$  defined on the Cantor set  $C$ . Thus by [7, Theorem 3.2],  $E(X)$  is homeomorphic to a closed subset of  $\mathfrak{E}_c$ .  $\square$

We are not aware of whether the converse implication of Lemma 14 is true. This would provide a very nice connection between the theory of almost zero-dimensional spaces and the theory of smooth fans.

**15. Question** Let  $E$  be a closed subset of  $\mathfrak{E}_c$ . Is there a smooth fan  $X$  such that  $E(X)$  is homeomorphic to  $E$ ?

We now turn our attention to which spaces may be realized as the endpoint set of an endpoint rigid smooth fan. The results of the previous section produce several examples of such spaces: for example the natural numbers, the irrationals, and the product of the Cantor set with the natural numbers (Propositions 6 and 7). Propositions 10, 11, and 13 (together with Proposition 6) yield many more

examples, such as the disjoint union of any compact zero-dimensional space with any one of the previously mentioned three spaces, disjoint unions of any of these examples, etc.

On the other hand, it is clear that no compact space may be the endpoint set of an endpoint rigid smooth fan. Indeed, if  $X$  is a smooth fan and  $E(X)$  is compact, then  $E(X)$  is zero-dimensional and  $X$  is homeomorphic to the cone over  $E(X)$ , and one can easily find a non-trivial autohomeomorphism of  $E(X)$ , which naturally extends to a non-trivial autohomeomorphism of  $X$ . Theorem 19 of the next section implies that  $\mathfrak{E}_c$  also cannot be realized as the endpoint set of an endpoint rigid smooth fan. We leave the following questions:

**16. Question** Let  $E$  be a non-compact closed subset of  $\mathfrak{E}_c$  that is not homeomorphic to complete Erdős space. Is there an endpoint rigid smooth fan  $X$  such that  $E(X)$  is homeomorphic to  $E$ ?

**17. Question** Let  $E$  be a locally compact, non-compact closed subset of  $\mathfrak{E}_c$ . Is there an endpoint rigid smooth fan  $X$  such that  $E(X)$  is homeomorphic to  $E$ ?

**18. Question** Let  $E$  be a closed subset of  $\mathfrak{E}_c$  such that there is a unique smooth fan  $X$  such that  $E(X)$  is homeomorphic to  $E$ . Is  $E$  either compact or homeomorphic to complete Erdős space?

## 5. CHARACTERIZATION OF THE LELEK FAN

In this section we will prove that the only smooth fan whose set of endpoints is homeomorphic to  $\mathfrak{E}_c$  is the Lelek fan.

**19. Theorem** If  $X$  is a smooth fan and  $E(X)$  is not dense in  $X$ , then  $E(X)$  has a point of zero-dimensionality. In particular,  $E(X)$  is not homeomorphic to  $\mathfrak{E}_c$ .

By Lemma 3, Theorem 19 is implied by the following statement about USC functions, which we prove below. Recall that  $G_0^\varphi$  and  $L_0^\varphi$  are defined in Section 2.2.

**20. Theorem** Let  $\varphi: C \rightarrow [0, \infty)$  be an upper semi-continuous function defined on the Cantor set  $C$ . Suppose there exists a non-empty open subset  $U \subseteq C$  and  $a_0 \in (0, \infty)$  such that  $(U \times \{a_0\}) \cap L_0^\varphi$  is non-empty and disjoint from  $G_0^\varphi$ . Then  $G_0^\varphi$  has a point of zero-dimensionality. In particular,  $G_0^\varphi$  is not homeomorphic to  $\mathfrak{E}_c$ .

*Proof.* Fix a metric  $d$  on  $C$ . We will use the metric  $\rho$  on  $C \times [0, 1]$  defined by

$$\rho(\langle x_1, y_1 \rangle, \langle x_2, y_2 \rangle) = \max\{d(x_1, x_2), |y_1 - y_2|\}.$$

To locate a point in  $G_0^\varphi$  of zero-dimensionality, we will do a recursive construction in  $\omega$  steps. At step  $n$ , we will define a clopen set  $V_n$  in  $C$  and  $a_n, b_n \in (0, \infty]$  with the following properties:

- (1)  $V_{n+1} \subseteq V_n$  for all  $n \in \omega$ ;
- (2)  $a_n \leq a_{n+1} < b_{n+1} \leq b_n$  for all  $n \in \omega$ ;
- (3)  $\emptyset \neq \varphi[V_n] \cap [a_0, \infty) \subset [a_n, b_n)$  for all  $n \in \omega$ ; and
- (4)  $\text{diam}(V_n) < 2^{-n}$  and  $b_n - a_n \leq 2^{-n}$  for all  $n \geq 1$ .

Let us give the construction. In step 0 we start with  $a_0$  as in the hypotheses of the Theorem, let  $V_0$  be a non-empty clopen subset of  $U$  such that  $(V_0 \times \{a_0\}) \cap L_0^\varphi \neq \emptyset$ , and define  $b_0 = \infty$ .

Now let  $n \in \omega$ , and assume that the construction has been carried out for step  $n$ . We now proceed with step  $n + 1$ . Let

$$a_{n+1} = \inf\{y \in (a_0, \infty) : \exists x \in V_n \text{ such that } \varphi(x) = y\}.$$

Note that  $a_{n+1} < b_n$  since by condition (3) there exists a point  $x \in V_n$  with  $\varphi(x) \in [a_n, b_n)$ . Let  $b_{n+1} = \min\{a_{n+1} + 2^{-(n+1)}, b_n\}$ .

Choose a point  $x_{n+1} \in V_n$  such that  $\varphi(x_{n+1}) \in [a_{n+1}, b_{n+1})$ . Let  $V_{n+1}$  be a clopen neighborhood of  $x_{n+1}$  in  $C$  of diameter less than  $2^{-(n+1)}$ , and (using the fact that  $\varphi$  is USC) such that  $\varphi[V_{n+1}] \cap [a_0, \infty) \subset [a_{n+1}, b_{n+1})$ . This completes the recursive construction.

Now, the sets  $V_n \cap \varphi^\leftarrow[[a_0, \infty))$ ,  $n \in \omega$ , are nested non-empty closed subsets of  $C$ , hence there exists a point  $z \in \bigcap_{n \in \omega} (V_n \cap \varphi^\leftarrow[[a_0, \infty)))$ . For each  $n \in \omega$ , let

$$W_n = (V_n \times [a_0, \infty)) \cap G_0^\varphi.$$

Observe that  $\langle z, \varphi(z) \rangle \in W_n$ , and  $W_n$  is clearly closed by definition since  $V_n$  is closed.  $W_n$  is open as well, since  $V_n$  is also open and  $(V_n \times \{a_0\}) \cap G_0^\varphi = \emptyset$ . By condition (3),  $W_n \subset V_n \times [a_n, b_n)$ , hence by condition (4) it has diameter less than  $2^{-n}$ . Thus, the sets  $W_n$ ,  $n \in \omega$ , form a local basis of clopen sets at  $\langle z, \varphi(z) \rangle$ , which is what we wanted to find.  $\square$

From Theorem 19, we obtain a new characterization of the Lelek fan: it is the unique smooth fan whose set of endpoints is homeomorphic to complete Erdős space. As such, we extend a result of W. Charatonik's ([5, Corollary]):

**21. Corollary** The following are equivalent for a smooth fan  $X$  with vertex  $p$ :

- (i)  $X$  is homeomorphic to the Lelek fan;
- (ii)  $E(X)$  is dense in  $X$ ;
- (iii)  $E(X) \cup \{p\}$  is connected;
- (iv) Every non-degenerate confluent image of  $X$  is homeomorphic to it;
- (v) Every non-degenerate monotone image of  $X$  is homeomorphic to it; and
- (vi)  $E(X)$  is homeomorphic to complete Erdős space.

We remark that the analog of Theorem 19 for smooth dendroids does not hold. For a counterexample, consider a smooth dendroid consisting of two copies of the Lelek fan whose tops are joined by an arc. In this dendroid, the set of endpoints is not dense, but it is 1-dimensional at every point.

#### REFERENCES

1. Gerardo Acosta, Logan C. Hoehn, and Yaziel Pacheco Juárez, *Homogeneity degree of fans*, Topology Appl. **231** (2017), 320–328.
2. Gerardo Acosta and Yaziel Pacheco-Juárez,  *$\frac{1}{3}$ -homogeneous dendrites*, Topology Appl. **219** (2017), 55–77.
3. A. V. Arhangel'skii and J. van Mill, *Topological homogeneity*, Recent progress in general topology III., Amsterdam: Atlantis Press, 2014, pp. 1–68.
4. J. J. Charatonik, *On fans*, Diss. Math. (Rozprawy Mat.) **54** (1967), 37 p.
5. W. J. Charatonik, *The Lelek fan is unique*, Houston J. Math. **15** (1989), no. 1, 27–34.
6. H. Cook, *Continua which admit only the identity mapping onto non-degenerate subcontinua*, Fundam. Math. **60** (1967), 241–249.

7. Jan J. Dijkstra and Jan van Mill, *Characterizing complete Erdős space*, Can. J. Math. **61** (2009), no. 1, 124–140.
8. ———, *Erdős space and homeomorphism groups of manifolds*, Mem. Am. Math. Soc., no. 979, Providence, RI: American Mathematical Society (AMS), 2010.
9. Carl Eberhart, *A note on smooth fans*, Colloq. Math. **20** (1969), 89–90.
10. Paul Erdős, *The dimension of the rational points in Hilbert space*, Ann. Math. (2) **41** (1940), 734–736.
11. Kazuhiro Kawamura, Lex G. Oversteegen, and E. D. Tymchatyn, *On homogeneous totally disconnected 1-dimensional spaces*, Fundam. Math. **150** (1996), no. 2, 97–112.
12. A. Lelek, *On plane dendroids and their end points in the classical sense*, Fund. Math. **49** (1961), 301–319.
13. A. J. M. van Engelen, *Homogeneous zero-dimensional absolute Borel sets*, CWI Tracts, 27. Centrum voor Wiskunde en Informatica. Amsterdam: Mathematisch Centrum. III, 133 p., 1986.
14. Fons van Engelen, Arnold W. Miller, and John Steel, *Rigid Borel sets and better quasiorder theory*, Logic and combinatorics, Proc. AMS-IMS-SIAM Conf., Arcata/Calif. 1985, Contemp. Math. 65, 199–222, 1987.

(Hernández-Gutiérrez) DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA METROPOLITANA CAMPUS IZTAPALAPA, AV. SAN RAFAEL ATLIXCO 186, LEYES DE REFORMA 1A SECCIÓN, IZTAPALAPA, 09310, MEXICO CITY, MEXICO

*Email address*, Hernández-Gutiérrez: `rod@xanum.uam.mx`

(Hoehn) NIPISSING UNIVERSITY, DEPARTMENT OF COMPUTER SCIENCE & MATHEMATICS, 100 COLLEGE DRIVE, BOX 5002, NORTH BAY, ONTARIO, CANADA, P1B 8L7

*Email address*, Hoehn: `loganh@nipissingu.ca`