

WEIGHTED BMO ESTIMATES FOR SINGULAR INTEGRALS AND ENDPOINT EXTRAPOLATION IN BANACH FUNCTION SPACES

ZOE NIERAETH AND GUILLERMO REY

ABSTRACT. In this paper we prove sharp weighted BMO estimates for singular integrals, and we show how such estimates can be extrapolated to Banach function spaces.

1. INTRODUCTION

In the 70's Muckenhoupt and Wheeden extended the known $L^\infty \rightarrow \text{BMO}$ estimates for the Hilbert transform to the weighted setting. In particular, they proved that

$$\int_I |Hf - \langle Hf \rangle_I| dx \lesssim_w \|fw\|_{L^\infty} \int_I w^{-1} dx,$$

where $\langle Hf \rangle_I$ is the average of Hf over I , holds if and only if $w^{-1} \in A_\infty \cap B_2$. This is Theorem 1 in [MW76].

The class B_2 , which was introduced in [HMW73], is the collection of all locally integrable weights w satisfying

$$\int_{I^c} \frac{|I|w(x)}{|x - c_I|^2} dx \lesssim \langle w \rangle_I.$$

In the same paper (Theorem 2), Muckenhoupt and Wheeden characterize $w^{-1} \in A_1$ as the class of weights w for which the following weighted BMO estimate holds

$$(1.1) \quad \|w\|_{L^\infty(I)} \frac{1}{|I|} \int_I |Hf - \langle Hf \rangle_I| dx \lesssim_w \|fw\|_{L^\infty}$$

for all intervals I and all functions f .

This result was later used by Harboure, Macías, and Segovia in [HMS88] to prove a BMO-to- L^p extrapolation result, saying that if a sublinear operator T satisfies the same bound as in (1.1) with implicit constant depending only on $[w^{-1}]_{A_1}$, then

$$\|Tf\|_{L^p(v)} \lesssim_v \|f\|_{L^p(v)}$$

for all weights $v \in A_p$ and $1 < p < \infty$.

We should mention the result of [Nie19], where a multilinear extrapolation result is obtained which generalizes and sharpens the one by Harboure, Macías, and Segovia. See also [CPRR19], where sharp linear versions of these two results are obtained.

The purpose of this article is twofold: first, we extend the extrapolation result of Harboure, Macías, and Segovia, to the setting of Banach function spaces. Second, we extend the results of Muckenhoupt and Wheeden to sparse operators as well as to Calderón-Zygmund operators whose kernels satisfy a Dini smoothness condition. We fully recover the results of [CPRR19], and in fact our extrapolation result is sharper, see Remark 3.3.

Let us first state a simplified version of our extrapolation theorem.

2010 *Mathematics Subject Classification.* 42B20, 46E30.

Key words and phrases. Sparse operators, Muckenhoupt weights, Banach function spaces, Calderón-Zygmund operators, BMO, Rubio de Francia extrapolation.

Theorem A. *Let T be a sublinear operator. Suppose there is an increasing function $\phi : [1, \infty) \rightarrow (0, \infty)$ such that for all weights w with $w^{-1} \in A_1$, and all compactly supported functions f with $fw \in L^\infty(\mathbb{R}^d)$ we have*

$$\|w\|_{L^\infty(Q)} \frac{1}{|Q|} \int_Q |Tf - \langle Tf \rangle_Q| dx \leq \phi([w^{-1}]_{A_1}) \|fw\|_{L^\infty(\mathbb{R}^d)}.$$

Then, for all Banach function spaces X over \mathbb{R}^d for which M is bounded on X and its associate space X' , and all compactly supported functions $f \in X$

$$\|Tf\|_X \lesssim_d \|M\|_{X' \rightarrow X'} \phi(2\|M\|_{X \rightarrow X}) \|f\|_X.$$

Our full result is Theorem 3.1 in section 3.

As for our result for Calderón-Zygmund operators, suppose T is an operator represented by

$$Tf(x) = \int_{\mathbb{R}^d} K(x, y) f(y) dy$$

for all x outside of the support of f , and where K satisfies

$$|K(x, y) - K(z, y)| \leq \Omega\left(\frac{|x - z|}{|x - y|}\right) \frac{1}{|x - y|^d}$$

for all x, y, z satisfying $|x - y| > 2|x - z| > 0$, and where $\Omega : [0, \infty) \rightarrow [0, \infty)$ is an increasing subadditive function with $\Omega(0) = 0$.

Similar to the situation in [MW76], it is not sufficient that $w \in A_\infty$ for T to have weighted BMO estimates. What we need is a generalization of the B_2 condition that takes into account the interaction between the weight and the smoothness of the kernel. This is quantified by

$$[w]_{B(\Omega)} = \sup_Q \frac{|Q|}{w(Q)} \int_{Q^c} \frac{w(x)}{|x - c_Q|^d} \Omega\left(\frac{\ell(Q)}{|x - c_Q|}\right) dx.$$

Note that, in dimension one and when $\Omega(t) = t$, $[w]_{B(\Omega)} < \infty$ is the B_2 condition.

With this definition, we have

Theorem B. *Let $w^{-1} \in L^1_{\text{loc}}(\mathbb{R}^d)$ and let T be an operator as above, then*

$$\frac{1}{|Q|} \int_Q |Tf - \langle Tf \rangle_Q| dx \lesssim ([1]_{B(\Omega)} + [w^{-1}]_{B(\Omega)}) [w^{-1}]_{A_\infty} \langle w^{-1} \rangle_Q \|fw\|_{L^\infty(\mathbb{R}^d)}$$

and

$$\|w\|_{L^\infty(Q)} \frac{1}{|Q|} \int_Q |Tf - \langle Tf \rangle_Q| dx \lesssim [w^{-1}]_{A_1}^2 [w^{-1}]_{A_\infty} \|\Omega\|_{\text{Dim}} \|fw\|_{L^\infty(\mathbb{R}^d)}$$

for all compactly supported functions f with $fw \in L^\infty(\mathbb{R}^d)$.

This is Theorem 4.4 in section 4. These estimates are based on sparse domination techniques, so they offer a simplified argument to the results from [MW76] and [MW78].

Notation. We work in the setting of \mathbb{R}^d equipped with the Lebesgue measure dx . For a measurable set $E \subseteq \mathbb{R}^d$ we denote its measure by $|E|$. For a measurable function $f \in L^0(\mathbb{R}^d)$, a measurable set $E \subseteq \mathbb{R}^d$ of finite positive measure, and a $q \in (0, \infty)$ we write

$$\langle f \rangle_{q, E} := \left(\frac{1}{|E|} \int_E |f|^q dx \right)^{\frac{1}{q}},$$

and denote the essential supremum of f in E by $\langle f \rangle_{\infty, E}$. Moreover, we denote the linearized mean of f on E by

$$\langle f \rangle_E := \frac{1}{|E|} \int_E f dx.$$

By a *cube* in \mathbb{R}^d we mean a half-open cube whose sides are parallel to the coordinate axes.

We will write $A \lesssim_{a,b,\dots} B$ to mean that there is a constant C depending only on the parameters a, b, \dots , such that $A \leq CB$. We write $A \approx_{a,b,\dots} B$ when both $A \lesssim_{a,b,\dots} B$ and $B \lesssim_{a,b,\dots} A$.

2. PRELIMINARIES

2.1. Dyadic analysis. In this paper we will be working in \mathbb{R}^d , but our results are also valid in more general spaces of homogeneous type. We will often reduce our arguments to dyadic grids: a dyadic grid is a collection of cubes with the property

$$P \cap Q \neq \emptyset \implies P \subseteq Q \text{ or } Q \subseteq P.$$

For a collection of cubes \mathcal{E} in a dyadic grid \mathcal{D} and a cube $Q_0 \in \mathcal{D}$, we let $\mathcal{E}(Q_0)$ denote the cubes in \mathcal{E} that are contained in Q_0 . We refer the interested reader to the monograph [LN19].

2.2. Muckenhoupt weights. A weight w in \mathbb{R}^d can be associated with the measure through $w(E) := \int_E w \, dx$. A classical result by Muckenhoupt is that the Hardy-Littlewood maximal operator

$$Mf := \sup_Q \langle f \rangle_{1,Q} \mathbb{1}_Q,$$

where the supremum is taken over all cubes Q in \mathbb{R}^d , is bounded $L^p(\mathbb{R}^d, w) \rightarrow L^p(\mathbb{R}^d, w)$ for $p \in (1, \infty)$ precisely when w satisfies the A_p condition

$$[w]_{A_p} := \sup_Q \langle w \rangle_{1,Q} \langle w^{-\frac{1}{p}} \rangle_{p',Q}^p < \infty,$$

and bounded $L^1(\mathbb{R}^d, w) \rightarrow L^{1,\infty}(\mathbb{R}^d, w)$ if and only if w satisfies the A_1 condition

$$[w]_{A_1} := \sup_Q \langle w \rangle_{1,Q} \langle w^{-1} \rangle_{\infty,Q} = \|(Mw)w^{-1}\|_{L^\infty(\mathbb{R}^d)} < \infty.$$

However, in the case $p = \infty$, we have $L^\infty(\mathbb{R}^d, w) = L^\infty(\mathbb{R}^d)$ for all weights w and hence, M is bounded $L^\infty(\mathbb{R}^d, w) \rightarrow L^\infty(\mathbb{R}^d, w)$ for all weights w .

Instead, if we define $L_w^p(\mathbb{R}^d)$ through $\|f\|_{L_w^p(\mathbb{R}^d)} := \|fw\|_{L^p(\mathbb{R}^d)}$, then we do get a meaningful condition for $p = \infty$. Note that for $p \in (1, \infty)$ we have $L_w^p(\mathbb{R}^d) = L^p(\mathbb{R}^d : w^p)$ and hence, $M : L_w^p(\mathbb{R}^d) \rightarrow L_w^p(\mathbb{R}^d)$ is bounded precisely when

$$[w^p]_{A_p}^{\frac{1}{p}} = \sup_Q \langle w \rangle_{p,Q} \langle w^{-1} \rangle_{p',Q} < \infty.$$

For $p = \infty$, this expression yields the condition

$$\sup_Q \langle w \rangle_{\infty,Q} \langle w^{-1} \rangle_{1,Q} = \|M(w^{-1})w\|_{L^\infty(\mathbb{R}^d)} = [w^{-1}]_{A_1}.$$

Indeed, defining $L_w^\infty(\mathbb{R}^d)$ through $\|f\|_{L_w^\infty(\mathbb{R}^d)} = \|fw\|_{L^\infty(\mathbb{R}^d)}$, we have the following result:

Proposition 2.1. *Let w be a weight in \mathbb{R}^d . Then M is bounded $L_w^\infty(\mathbb{R}^d) \rightarrow L_w^\infty(\mathbb{R}^d)$ if and only if $w^{-1} \in A_1$. In this case we have*

$$\|M\|_{L_w^\infty(\mathbb{R}^d) \rightarrow L_w^\infty(\mathbb{R}^d)} = [w^{-1}]_{A_1}.$$

Proof. If M is bounded $L_w^\infty(\mathbb{R}^d) \rightarrow L_w^\infty(\mathbb{R}^d)$, then since $f = w^{-1}$ satisfies $\|f\|_{L_w^\infty(\mathbb{R}^d)} = 1$, we have

$$[w^{-1}]_{A_1} = \|M(w^{-1})\|_{L_w^\infty(\mathbb{R}^d)} \leq \|M\|_{L_w^\infty(\mathbb{R}^d) \rightarrow L_w^\infty(\mathbb{R}^d)}.$$

For the converse, if $w^{-1} \in A_1$, then for all $f \in L_w^\infty(\mathbb{R}^d)$ we have

$$\|Mf\|_{L_w^\infty(\mathbb{R}^d)} \leq \|f\|_{L_w^\infty(\mathbb{R}^d)} \|M(w^{-1})\|_{L_w^\infty(\mathbb{R}^d)} = [w^{-1}]_{A_1} \|f\|_{L_w^\infty(\mathbb{R}^d)}.$$

The assertion follows. \square

If \mathcal{D} is a dyadic grid in \mathbb{R}^d , then we define

$$M^{\mathcal{D}}f := \sup_{Q \in \mathcal{D}} \langle f \rangle_{1,Q} \mathbb{1}_Q$$

and the class $A_1(\mathcal{D})$ through

$$[w]_{A_1(\mathcal{D})} := \sup_{Q \in \mathcal{D}} \langle w \rangle_{1,Q} \langle w^{-1} \rangle_{\infty,Q} = \|(M^{\mathcal{D}}w)w^{-1}\|_{L^\infty(\mathbb{R}^d)}.$$

Then, exactly as in the above result, we have $\|M^{\mathcal{D}}\|_{L_w^\infty(\mathbb{R}^d) \rightarrow L_w^\infty(\mathbb{R}^d)} = [w^{-1}]_{A_1(\mathcal{D})}$.

For a cube $Q \in \mathcal{D}$ we denote by $\mathcal{D}(Q)$ the collection of cubes in \mathcal{D} that are contained in Q . Defining $M^{\mathcal{D}(Q)}$ accordingly, we define the Fujii-Wilson A_∞ constant of a weight through

$$[w]_{A_\infty(\mathcal{D})} := \sup_{Q \in \mathcal{D}} \frac{1}{w(Q)} \int_Q M^{\mathcal{D}(Q)}w \, dx.$$

Finiteness of this constant characterizes the class $\bigcup_{p \geq 1} A_p(\mathcal{D})$. Note that in particular we have

$$[w]_{A_\infty(\mathcal{D})} \leq [w]_{A_1(\mathcal{D})}.$$

In the non-dyadic case, the Fujii-Wilson characteristic is defined as

$$[w]_{A_\infty} := \sup_Q \frac{1}{w(Q)} \int_Q M(\mathbb{1}_Q w) \, dx.$$

where the supremum is taken over all cubes.

We have the following well-known result (which, in fact, is a characterization of $A_\infty(\mathcal{D})$):

Proposition 2.2. *Let $\mathcal{S} \subseteq \mathcal{D}$ be an η -sparse collection of cubes and let $Q_0 \in \mathcal{D}$. Then for a weight w we have*

$$\sum_{\substack{Q \in \mathcal{S} \\ Q \subseteq Q_0}} w(Q) \leq \eta^{-1} [w]_{A_\infty(\mathcal{D})} w(Q_0).$$

Proof. We have

$$\sum_{\substack{Q \in \mathcal{S} \\ Q \subseteq Q_0}} w(Q) \leq \eta^{-1} \sum_{\substack{Q \in \mathcal{S} \\ Q \subseteq Q_0}} \inf_{y \in Q} M^{\mathcal{D}(Q_0)}w(y) |E_Q| \leq \eta^{-1} \int_{Q_0} M^{\mathcal{D}(Q_0)}w \, dx \leq \eta^{-1} [w]_{A_\infty(\mathcal{D})} w(Q_0),$$

as desired. \square

2.3. Weighted BMO spaces. The space $\text{BMO}(\mathbb{R}^d)$ is defined as the space of function $f \in L^0(\mathbb{R}^d)$ for which the sharp maximal function

$$M^\sharp f := \sup_Q \langle f - \langle f \rangle_Q \rangle_{1,Q} \mathbb{1}_Q$$

lies in $L^\infty(\mathbb{R}^d)$. Thus, it is sensible to define a weighted analogue $\text{BMO}_w(\mathbb{R}^d)$ as those $f \in L^0(\mathbb{R}^d)$ for which $M^\sharp f \in L_w^\infty(\mathbb{R}^d)$. This is facilitated through the following definition.

Definition 2.3. For a weight w we define the spaces $\text{BMO}_w^1(\mathbb{R}^d)$ and $\text{BMO}_w^\infty(\mathbb{R}^d)$ through

$$\begin{aligned} \|f\|_{\text{BMO}_w^1(\mathbb{R}^d)} &:= \sup_Q \frac{\langle f - \langle f \rangle_Q \rangle_{1,Q}}{\langle w^{-1} \rangle_{1,Q}} = \sup_Q \frac{1}{w^{-1}(Q)} \int_Q |f - \langle f \rangle_Q| \, dx, \\ \|f\|_{\text{BMO}_w^\infty(\mathbb{R}^d)} &:= \sup_Q \langle w \rangle_{\infty,Q} \langle f - \langle f \rangle_Q \rangle_{1,Q}, \end{aligned}$$

where functions are identified modulo constants.

We also define the weak analogues $\text{BMO}_w^{1,\text{weak}}(\mathbb{R}^d)$, $\text{BMO}_w^{\infty,\text{weak}}(\mathbb{R}^d)$ of these spaces through

$$\begin{aligned}\|f\|_{\text{BMO}_w^{1,\text{weak}}(\mathbb{R}^d)} &:= \sup_Q \inf_{c \in \mathbb{R}} \frac{\|(f - c) \mathbb{1}_Q\|_{L^{1,\infty}(Q)}}{w^{-1}(Q)}, \\ \|f\|_{\text{BMO}_w^{\infty,\text{weak}}(\mathbb{R}^d)} &:= \sup_Q \inf_{c \in \mathbb{R}} \langle w \rangle_{\infty,Q} |Q|^{-1} \|(f - c) \mathbb{1}_Q\|_{L^{1,\infty}(Q)},\end{aligned}$$

where functions are identified modulo constants.

For a dyadic grid \mathcal{D} in \mathbb{R}^d , the spaces $\text{BMO}_w^1(\mathcal{D})$, $\text{BMO}_w^\infty(\mathcal{D})$, $\text{BMO}_w^{1,\text{weak}}(\mathcal{D})$, and $\text{BMO}_w^{\infty,\text{weak}}(\mathcal{D})$ are defined analogously, this time taking the supremum over all cubes in \mathcal{D} , and functions are identified modulo constants on cubes in \mathcal{D} .

Noting that

$$\frac{\langle w \rangle_{\infty,Q}}{|Q|} = \frac{1}{(\text{ess inf}_Q w^{-1})|Q|} \geq \frac{1}{w^{-1}(Q)}$$

for all cubes Q in \mathbb{R}^d , we have

$$\|f\|_{\text{BMO}_w^1(\mathbb{R}^d)} \leq \|f\|_{\text{BMO}_w^\infty(\mathbb{R}^d)}, \quad \|f\|_{\text{BMO}_w^{1,\text{weak}}(\mathbb{R}^d)} \leq \|f\|_{\text{BMO}_w^{\infty,\text{weak}}(\mathbb{R}^d)}$$

and hence, $\text{BMO}_w^\infty(\mathbb{R}^d) \subseteq \text{BMO}_w^1(\mathbb{R}^d)$ and $\text{BMO}_w^{\infty,\text{weak}}(\mathbb{R}^d) \subseteq \text{BMO}_w^{1,\text{weak}}(\mathbb{R}^d)$. Conversely, if $w^{-1} \in A_1$, we have

$$\langle w \rangle_{\infty,Q} \leq \frac{[w^{-1}]_{A_1}}{\langle w^{-1} \rangle_{1,Q}}$$

for all cubes Q in \mathbb{R}^d . Thus, in this case we have

$$(2.1) \quad \|f\|_{\text{BMO}_w^\infty(\mathbb{R}^d)} \leq [w^{-1}]_{A_1} \|f\|_{\text{BMO}_w^1(\mathbb{R}^d)},$$

$$(2.2) \quad \|f\|_{\text{BMO}_w^{\infty,\text{weak}}(\mathbb{R}^d)} \leq [w^{-1}]_{A_1} \|f\|_{\text{BMO}_w^{1,\text{weak}}(\mathbb{R}^d)},$$

and $\text{BMO}_w^\infty(\mathbb{R}^d) = \text{BMO}_w^1(\mathbb{R}^d)$, $\text{BMO}_w^{\infty,\text{weak}}(\mathbb{R}^d) = \text{BMO}_w^{1,\text{weak}}(\mathbb{R}^d)$.

We also note that $L_w^\infty(\mathbb{R}^d) \subseteq \text{BMO}_w^1(\mathbb{R}^d)$ with

$$\|f\|_{\text{BMO}_w^1(\mathbb{R}^d)} \leq 2\|f\|_{L_w^\infty(\mathbb{R}^d)}.$$

Thus, if $w^{-1} \in A_1$, we have $L_w^\infty(\mathbb{R}^d) \subseteq \text{BMO}_w^\infty(\mathbb{R}^d)$ with

$$\|f\|_{\text{BMO}_w^\infty(\mathbb{R}^d)} \leq 2[w^{-1}]_{A_1} \|f\|_{L_w^\infty(\mathbb{R}^d)}.$$

The following result shows that the space $\text{BMO}_w^\infty(\mathbb{R}^d)$ consists of precisely those $f \in L^0(\mathbb{R}^d)$ for which $M^\sharp f \in L_w^\infty(\mathbb{R}^d)$.

Proposition 2.4. *We have*

$$\|f\|_{\text{BMO}_w^\infty(\mathbb{R}^d)} = \|M^\sharp f\|_{L_w^\infty(\mathbb{R}^d)}, \quad \|f\|_{\text{BMO}_w^{\infty,\text{weak}}(\mathbb{R}^d)} = \|M_{\text{weak}}^\sharp f\|_{L_w^\infty(\mathbb{R}^d)},$$

where

$$M_{\text{weak}}^\sharp f := \sup_Q \inf_{c \in \mathbb{R}} |Q|^{-1} \|(f - c) \mathbb{1}_Q\|_{L^{1,\infty}(Q)} \mathbb{1}_Q.$$

For a dyadic grid \mathcal{D} in \mathbb{R}^d , analogously defining the sharp maximal operators $M^{\sharp,\mathcal{D}}$ and $M_{1,\infty}^{\sharp,\mathcal{D}}$ with respect to this grid, we also have

$$\|f\|_{\text{BMO}_w^\infty(\mathcal{D})} = \|M^{\sharp,\mathcal{D}} f\|_{L_w^\infty(\mathcal{D})}, \quad \|f\|_{\text{BMO}_w^{1,\infty}(\mathcal{D})} = \|M_{\text{weak}}^{\sharp,\mathcal{D}} f\|_{L_w^\infty(\mathcal{D})}.$$

Proof. We only prove the first equality, the others being analogous. Since $\mathbb{1}_Q w \leq \langle w \rangle_{\infty,Q}$ for all cubes Q , we have

$$(M^\sharp f)w = \sup_Q (f - \langle f \rangle_Q)_{1,Q} \mathbb{1}_Q w \leq \|f\|_{\text{BMO}_w^\infty(\mathbb{R}^d)}.$$

This proves the inequality $\|(M^\sharp f)w\|_{L^\infty(\mathbb{R}^d)} \leq \|f\|_{\text{BMO}_w^\infty(\mathbb{R}^d)}$. For the converse, fix a cube Q and let $\varepsilon > 0$. Then the set of $x \in Q$ such that $\langle w \rangle_{\infty, Q} \leq (1 + \varepsilon)w(x)$ has positive measure. Hence, there are $x \in Q$ for which

$$\langle w \rangle_{\infty, Q} \langle f - \langle f \rangle_Q \rangle_{1, Q} \leq (1 + \varepsilon) \langle f - \langle f \rangle_Q \rangle_{1, Q} w(x) \mathbb{1}_Q(x) \leq (1 + \varepsilon) \|(M^\sharp f)w\|_{L^\infty(\mathbb{R}^d)}$$

Thus, $\|f\|_{\text{BMO}_w^\infty(\mathbb{R}^d)} \leq (1 + \varepsilon) \|(M^\sharp f)w\|_{L^\infty(\mathbb{R}^d)}$. Letting $\varepsilon \rightarrow 0$, the assertion follows. \square

Proposition 2.5. *Let w be a weight. We have*

$$\begin{aligned} \|f\|_{\text{BMO}_w^1(\mathbb{R}^d)} &\approx \sup_Q \inf_{c \in \mathbb{R}} \frac{\langle f - c \rangle_{1, Q}}{\langle w^{-1} \rangle_{1, Q}}, \\ \|f\|_{\text{BMO}_w^\infty(\mathbb{R}^d)} &\approx \sup_Q \inf_{c \in \mathbb{R}} \langle w \rangle_{\infty, Q} \langle f - c \rangle_{1, Q}. \end{aligned}$$

Thus, in particular,

$$\begin{aligned} \|f\|_{\text{BMO}_w^{1, \text{weak}}(\mathbb{R}^d)} &\lesssim \|f\|_{\text{BMO}_w^1(\mathbb{R}^d)}, \\ \|f\|_{\text{BMO}_w^{\infty, \text{weak}}(\mathbb{R}^d)} &\lesssim \|f\|_{\text{BMO}_w^\infty(\mathbb{R}^d)}. \end{aligned}$$

This similarly holds for $\text{BMO}_w^1(\mathcal{D})$, $\text{BMO}_w^\infty(\mathcal{D})$ for a dyadic grid \mathcal{D} in \mathbb{R}^d .

Proof. We only prove the second equivalence, the first one being analogous. The inequality \geq follows from taking $c = \langle f \rangle_Q$. For the converse inequality, fix a cube Q and note that for any $c \in \mathbb{R}$ we have

$$\langle f - \langle f \rangle_Q \rangle_{1, Q} \leq \langle f - c \rangle_{1, Q} + |\langle f - c \rangle_Q| \leq 2 \langle f - c \rangle_{1, Q}.$$

Thus, $\|f\|_{\text{BMO}_w(\mathbb{R}^d)} \leq 2 \sup_Q \inf_{c \in \mathbb{R}} \langle w \rangle_{\infty, Q} \langle f - c \rangle_{1, Q}$. As the result follows analogously for $\text{BMO}_w(\mathcal{D})$, this proves the assertion. \square

In the unweighted case, it follows from the John-Strömberg characterization of BMO that the weak BMO space is equivalent to the usual one. More precisely, denoting the non-decreasing rearrangement of a function f by f^* , for a cube Q and $\lambda \in (0, 1)$ we define

$$M_\lambda^\sharp f := \sup_Q \omega_\lambda(f; Q) \mathbb{1}_Q,$$

where

$$\omega_\lambda(f; Q) = \inf_{c \in \mathbb{R}} \inf \{t > 0 : |\{x \in Q : |f - c| > t\}| \leq \lambda|Q|\}.$$

Observe that

$$\omega_\lambda(f; Q) \leq \lambda^{-1} |Q|^{-1} \inf_{c \in \mathbb{R}} \|f - c\|_{L^{1, \infty}(Q)},$$

so that $M_\lambda^\sharp f \leq \lambda^{-1} M_{\text{weak}}^\sharp f$.

We can extend John-Strömberg's characterization of BMO through medians (see [Str79]) to the weighted setting using the following pointwise version by A. Lerner [Ler03, Theorem 1.3]:

$$(2.3) \quad M^\sharp f \approx_d M(M_{\frac{1}{2}}^\sharp f).$$

for all $f \in L_{\text{loc}}^1(\mathbb{R}^d)$.

Proposition 2.6. *Let $w^{-1} \in A_1$. Then $\text{BMO}_w^\infty(\mathbb{R}^d) = \text{BMO}_w^{\infty, \text{weak}}(\mathbb{R}^d)$ with*

$$\|f\|_{\text{BMO}_w^\infty(\mathbb{R}^d)} \lesssim_d [w^{-1}]_{A_1} \|f\|_{\text{BMO}_w^{\infty, \text{weak}}(\mathbb{R}^d)}.$$

Proof. By Proposition 2.4, (2.3), and Proposition 2.1, we have

$$\|f\|_{\text{BMO}_w^\infty(\mathbb{R}^d)} \approx_d \|M(M_{\frac{1}{2}}^\sharp f)\|_{L_w^\infty(\mathbb{R}^d)} \leq [w^{-1}]_{A_1} \|M_{\frac{1}{2}}^\sharp f\|_{L_w^\infty(\mathbb{R}^d)}.$$

The result now follows from the fact that $M_{\frac{1}{2}}^\sharp f \lesssim M_{\text{weak}}^\sharp f$. \square

2.4. Banach function spaces. We denote the positive measurable functions on \mathbb{R}^d by $L^0(\mathbb{R}^d)_+$.

Definition 2.7. Suppose $\rho : L^0(\mathbb{R}^d)_+ \rightarrow [0, \infty]$ satisfies

- (i) $\rho(f) = 0$ if and only if $f = 0$ a.e.;
- (ii) $\rho(f+g) \leq \rho(f) + \rho(g)$ and $\rho(\lambda f) = \lambda\rho(f)$ for all $f, g \in L^0(\mathbb{R}^d)_+$ and scalars $\lambda \geq 0$;
- (iii) for every sequence $(f_n)_{n \in \mathbb{N}}$ in $L^0(\mathbb{R}^d)_+$ satisfying $0 \leq f_n \uparrow f$ pointwise a.e. for $f \in L^0(\mathbb{R}^d)_+$, we have $\rho(f_n) \uparrow \rho(f)$.
- (iv) for every measurable $E \subseteq \mathbb{R}^d$ of positive measure there exists a measurable $F \subseteq E$ of positive measure with $\rho(\mathbb{1}_F) < \infty$.

Then we call ρ a function norm. Moreover, we call the space $X \subseteq L^0(\mathbb{R}^d)$ of $f \in L^0(\mathbb{R}^d)$ for which

$$\|f\|_X := \rho(|f|) < \infty,$$

a Banach function space over \mathbb{R}^d .

Given a function norm ρ , we define a new function norm ρ' through

$$\rho'(g) := \sup_{\rho(f)=1} \|fg\|_{L^1(\mathbb{R}^d)}.$$

The K ote dual of X is defined as the Banach function space associated to ρ' , i.e.,

$$X' := \{g \in L^0(\mathbb{R}^d) : fg \in L^1(\mathbb{R}^d) \text{ for all } f \in X\}$$

with

$$\|g\|_{X'} = \sup_{\|f\|_X=1} \|fg\|_{L^1(\mathbb{R}^d)}.$$

Property (iv) is called the saturation property of ρ and is equivalent to the existence of a weak order unit, i.e., an $f > 0$ for which $\rho(f) < \infty$. Property (iv) is also equivalent to the fact that ρ' satisfies (i) and hence, is a function norm. Thus, X' is a Banach function space over \mathbb{R}^d .

The Fatou property (iii) is equivalent to the assertion that $\rho'' = \rho$ and hence, $X'' = X$ isometrically. Thus, in particular we have

$$\|f\|_X = \sup_{\|g\|_{X'}=1} \|fg\|_{L^1(\mathbb{R}^d)}.$$

We say that a Banach function space X over \mathbb{R}^d is order-continuous when for all $(f_n)_{n \in \mathbb{N}}$ in X with $f_n \downarrow 0$ a.e. we have $\|f_n\|_X \rightarrow 0$. We note that X is reflexive if and only if X and X' are order-continuous. For proofs of the above claims we refer the reader to [Zaa67].

By carefully tracking the constants in the proof of [Ler10b, Corollary 4.3], we arrive at the following quantitative version:

Theorem 2.8. *Let X be a Banach function space over \mathbb{R}^d and suppose that $M : X \rightarrow X$ is bounded. Then the following assertions are equivalent:*

- (i) *There is a constant $C_X > 0$ such that*

$$\|f\|_X \leq C_X \|M^\sharp f\|_X$$

for all $f \in X$ with the property that $|\{x \in \mathbb{R}^d : |f(x)| > \alpha\}| < \infty$ for all $\alpha > 0$;

- (ii) *There is a constant $C_{X,\text{weak}} > 0$ such that*

$$\|f\|_X \leq C_{X,\text{weak}} \|M^\sharp_{\text{weak}} f\|_X$$

for all $f \in X$ with the property that $|\{x \in \mathbb{R}^d : |f(x)| > \alpha\}| < \infty$ for all $\alpha > 0$;

- (iii) *$M : X' \rightarrow X'$ is bounded.*

Moreover, if C_X and $C_{X,\text{weak}}$ are the smallest possible constant in (i) and (ii), then

$$(2.4) \quad C_X \leq C_{X,\text{weak}} \lesssim_d \|M\|_{X' \rightarrow X'} \lesssim_d C_X \|M\|_{X \rightarrow X}.$$

Proof. Our strategy will be to prove that (ii) \Rightarrow (i) \Rightarrow (iii) \Rightarrow (ii).

The assertion (ii) \Rightarrow (i) with $C_X \leq C_{X,\text{weak}}$ follows from the fact that $M_{\text{weak}}^\sharp f \leq M^\sharp f$.

For (i) \Rightarrow (iii) we use that it was shown in [Ler10b, Theorem 1.1] that (i) is equivalent to the assertion

(iv) there is a $C > 0$ such that

$$\|fMg\|_{L^1(\mathbb{R}^d)} \leq C\|Mf\|_X\|g\|_{X'}$$

for all $f \in L^1_{\text{loc}}(\mathbb{R}^d)$ and $g \in X'$.

Moreover, if C is the smallest possible constant in this estimate, then

$$C \simeq_d C_X.$$

Thus, we have

$$\|Mg\|_{X'} = \sup_{\|f\|_X=1} \|fMg\|_{L^1(\mathbb{R}^d)} \leq C \sup_{\|f\|_X=1} \|Mf\|_X\|g\|_{X'} = C\|M\|_{X \rightarrow X}\|g\|_{X'}$$

so that $M : X' \rightarrow X'$ with $\|M\|_{X' \rightarrow X'} \lesssim_d C_X\|M\|_{X \rightarrow X}$, as desired

For (iii) \Rightarrow (ii), we use [Ler04, Theorem 1] which states that there is a dimensional constant $\lambda \in (0, 1)$ such that for all f satisfying the property that $|\{x \in \mathbb{R}^d : |f(x)| > \alpha\}| < \infty$ for all $\alpha > 0$ and all $g \in L^1_{\text{loc}}(\mathbb{R}^d)$ we have

$$\|fg\|_{L^1(\mathbb{R}^d)} \lesssim_d \|(M_\lambda^\sharp f)(Mg)\|_{L^1(\mathbb{R}^d)}.$$

Since $M_\lambda^\sharp f \leq \lambda^{-1}M_{\text{weak}}^\sharp f$, it follows that

$$\begin{aligned} \|f\|_X &= \sup_{\|g\|_{X'}=1} \|fg\|_{L^1(\mathbb{R}^d)} \lesssim_d \sup_{\|g\|_{X'}=1} \|(M_{\text{weak}}^\sharp f)(Mg)\|_{L^1(\mathbb{R}^d)} \\ &\leq \sup_{\|g\|_{X'}=1} \|M_{\text{weak}}^\sharp f\|_X\|Mg\|_{X'} = \|M\|_{X' \rightarrow X'}\|M_{\text{weak}}^\sharp f\|_X. \end{aligned}$$

This proves (ii) with $C_{X,\text{weak}} \lesssim_d \|M\|_{X' \rightarrow X'}$ and (2.4). The assertion follows. \square

3. RUBIO DE FRANCIA EXTRAPOLATION FROM WEIGHTED BMO

Theorem 3.1. *Let T be an operator in $L^0(\mathbb{R}^d)$. Suppose there is an increasing function $\phi : [1, \infty) \rightarrow (0, \infty)$ such that for all weights $w^{-1} \in A_1$ and all compactly supported functions $f \in L^\infty_w(\mathbb{R}^d)$ we have*

$$(3.1) \quad \|Tf\|_{\text{BMO}_w^\infty(\mathbb{R}^d)} \leq \phi([w^{-1}]_{A_1})\|f\|_{L^\infty_w(\mathbb{R}^d)}.$$

Then, for all Banach function spaces X over \mathbb{R}^d for which M is bounded on X and X' , and all compactly supported functions $f \in X$ we have

$$\|Tf\|_X \lesssim_d C_X\phi(2\|M\|_{X \rightarrow X})\|f\|_X,$$

where C_X is the constant from Theorem 2.8. If T is a (sub)linear operator and X is order-continuous, then T extends to a bounded operator $X \rightarrow X$ satisfying

$$(3.2) \quad \|T\|_{X \rightarrow X} \lesssim_d C_X\phi(2\|M\|_{X \rightarrow X}),$$

in particular

$$(3.3) \quad \|T\|_{X \rightarrow X} \lesssim_d \|M\|_{X' \rightarrow X'}\phi(2\|M\|_{X \rightarrow X}).$$

An analogous assertion holds when you replace $\text{BMO}_w^\infty(\mathbb{R}^d)$ by $\text{BMO}_w^{\infty,\text{weak}}(\mathbb{R}^d)$ and C_X by $C_{X,\text{weak}}$.

We need the following lemma which is based on the Rubio de Francia algorithm.

Lemma 3.2. *Let X be a Banach function space over \mathbb{R}^d for which $M : X \rightarrow X$. Then for all $f \in X$, $g \in X'$ there exists a weight $w^{-1} \in A_1$ such that*

$$(3.4) \quad [w^{-1}]_{A_1} \leq 2\|M\|_{X \rightarrow X},$$

and $f \in L_w^\infty(\mathbb{R}^d)$, $g \in L_{w^{-1}}^1(\mathbb{R}^d)$ with

$$(3.5) \quad \|f\|_{L_w^\infty(\mathbb{R}^d)} \|g\|_{L_{w^{-1}}^1(\mathbb{R}^d)} \leq 2\|f\|_X \|g\|_{X'}.$$

Proof. We set

$$w^{-1} := \sum_{k=0}^{\infty} \frac{M^k f}{2^k \|M\|_{X \rightarrow X}^k},$$

where we have recursively defined $M^0 f := |f|$, $M^{k+1} f := M(M^k f)$. Then we have

$$M(w^{-1}) \leq \sum_{k=0}^{\infty} \frac{M^{k+1} f}{2^k \|M\|_{X \rightarrow X}^k} \leq 2\|M\|_{X \rightarrow X} w^{-1},$$

proving (3.4).

Since $|f| = M^0 f \leq w^{-1}$, we have $\|f\|_{L_w^\infty(\mathbb{R}^d)} \leq 1$. Combining this with the fact that $\|w^{-1}\|_X \leq 2\|f\|_X$, we have

$$\|f\|_{L_w^\infty(\mathbb{R}^d)} \|g\|_{L_{w^{-1}}^1(\mathbb{R}^d)} \leq \|g w^{-1}\|_{L^1(\mathbb{R}^d)} \leq \|w^{-1}\|_X \|g\|_{X'} \leq 2\|f\|_X \|g\|_{X'}.$$

This proves (3.5), as desired. \square

Proof of Theorem 3.1. Let $f \in X$ have compact support and let $g \in X'$ with $\|g\|_{X'} = 1$. By Lemma 3.2 we can pick a weight $w^{-1} \in A_1$ such that $[w^{-1}]_{A_1} \leq 2\|M\|_{X \rightarrow X}$ and $f \in L_w^\infty(\mathbb{R}^d)$, $g \in L_{w^{-1}}^1(\mathbb{R}^d)$ with

$$\|f\|_{L_w^\infty(\mathbb{R}^d)} \|g\|_{L_{w^{-1}}^1(\mathbb{R}^d)} \leq 2\|f\|_X.$$

Hence, by Proposition 2.4 we have

$$\begin{aligned} \|M^\sharp(Tf)g\|_{L^1(\mathbb{R}^d)} &\leq \|M^\sharp(Tf)\|_{L_w^\infty(\mathbb{R}^d)} \|g\|_{L_{w^{-1}}^1(\mathbb{R}^d)} \leq \phi([w^{-1}]_{A_1}) \|f\|_{L_w^\infty(\mathbb{R}^d)} \|g\|_{L_{w^{-1}}^1(\mathbb{R}^d)} \\ &\leq 2\phi(2\|M\|_{X \rightarrow X}) \|f\|_X. \end{aligned}$$

Hence, since f has compact support and thus $|\{ |f| > \lambda \}| \leq |\text{supp } f| < \infty$ for all $\lambda > 0$, it follows from Theorem 2.8, that

$$\begin{aligned} \|Tf\|_X &\lesssim_d C_X M^\sharp(Tf)\|_X = C_X \sup_{\|g\|_{X'}=1} \|M^\sharp(Tf)g\|_{L^1(\mathbb{R}^d)} \\ &\leq 2C_X \phi(2\|M\|_{X \rightarrow X}) \|f\|_X, \end{aligned}$$

proving the first result.

For the next assertion, since X is order continuous the functions in X of compact support are dense in X . Indeed, given $f \in X$, set $f_n := f \mathbb{1}_{B(0;n)}$. Then $f_n \in X$ has compact support and $|f - f_n| = |f| \mathbb{1}_{\mathbb{R}^d \setminus B(0;n)} \downarrow 0$. Hence, $\|f - f_n\|_X \rightarrow 0$, as desired. Thus, the result follows from the fact that if T is (sub)linear, then it is uniformly continuous and hence, extends to all of X with the same bound.

The assertion for $\text{BMO}_w^\infty(\mathbb{R}^d)$ replaced by $\text{BMO}_w^{\infty, \text{weak}}(\mathbb{R}^d)$ and C_X replaced by $C_{X, \text{weak}}$ is proved analogously. \square

Remark 3.3. It was shown in [CPRR19] that the initial estimate (3.1) implies that for $p \in (1, \infty)$ and $w \in A_p$ we have

$$\|T\|_{L^p(w)} \lesssim_d \|M\|_{L^{p'}(\mathbb{R}^d; w^{1-p'})} \phi(\|M\|_{L^p(\mathbb{R}^d; w)}).$$

We recover this result in (3.3), since if $X = L^p(\mathbb{R}^d; w)$, then $X' = L^{p'}(\mathbb{R}^d; w^{1-p'})$. As a matter of fact, we improve on this estimate in (3.2). Indeed, for $X = L^p(w; \mathbb{R}^d)$ we obtain

$$\|T\|_{L^p(w)} \lesssim_d C_X \phi(\|M\|_{L^p(\mathbb{R}^d; w)}),$$

where C_X is the constant appearing in the Fefferman-Stein inequality

$$\|f\|_{L^p(\mathbb{R}^d;w)} \leq C_X \|M^\sharp f\|_{L^p(\mathbb{R}^d;w)}.$$

While $\|M\|_{L^{p'}(\mathbb{R}^d;w^{1-p'})}$ is bounded if and only if $w \in A_p$, the constant C_X is bounded under much more general conditions. For example, it is bounded when $w \in A_\infty$.

4. WEIGHTED BMO ESTIMATES FOR SINGULAR INTEGRALS

We will begin this section discussing the action of sparse operators on L_w^∞ . These serve as a way to gain intuition for the more complicated Calderón-Zygmund operators. We note, however, that unlike in the classical L^p case, estimates for sparse operators do not immediately imply bounds for Calderón-Zygmund operators because BMO is not a lattice.

4.1. Sparse operators. For $\eta \in (0, 1)$, a collection of sets \mathcal{S} in \mathbb{R}^d is called η -sparse if there exists a pairwise disjoint collection of sets $\{E_S\}_{S \in \mathcal{S}}$ such that $E_S \subseteq S$ and $|E_S| \geq \eta|S|$ for all $S \in \mathcal{S}$. For such a collection we define the *sparse operator* associated to \mathcal{S} as

$$\mathcal{A}_\mathcal{S} f := \sum_{Q \in \mathcal{S}} \langle f \rangle_{1,Q} \mathbb{1}_Q.$$

In general, these operators *do not* map L^∞ to BMO. There are several obstacles that prevent this, which we now describe.

For any family \mathcal{S} of sets in \mathbb{R}^d , let $h_\mathcal{S}$ be its height function:

$$h_\mathcal{S} = \sum_{J \in \mathcal{S}} \mathbb{1}_J.$$

First we note that $\mathcal{A}_\mathcal{S} f$ may not even be locally integrable for $f \in L^\infty(\mathbb{R}^d)$. In fact, the situation is much worse. Indeed, consider the sets $F_n = [0, 1) \cup [n, n+1)$ for $n \geq 1$ and define the family $\mathcal{S} = \{F_n : n \geq 1\}$. For each $n \geq 1$ the subset $E_n = [n, n+1) \subseteq F_n$ has measure $1 = \frac{1}{2}|F_n|$ and the collection $\{E_n\}$ is pairwise disjoint, thus \mathcal{S} is $\frac{1}{2}$ -sparse. However

$$\mathcal{A}_\mathcal{S}(\mathbb{1}_{[0,1)}) = \sum_{n=1}^{\infty} \frac{1}{2} \mathbb{1}_{F_n} = \frac{1}{2} h_\mathcal{S}$$

is identically ∞ on $[0, 1)$.

In general, we need *some* boundedness of the maximal function associated to \mathcal{S} for $\mathcal{A}_\mathcal{S}$ to behave well. In particular, if \mathcal{S} consists of intervals (or generally cubes in any dimension), then $\mathcal{A}_\mathcal{S}$ is bounded on $L^2(\mathbb{R}^d)$, so we have no problems with the local integrability of $\mathcal{A}_\mathcal{S}(f)$. But even this is not enough to reasonably define $\mathcal{A}_\mathcal{S}$ as an operator from $L^\infty(\mathbb{R}^d)$ to BMO(\mathbb{R}^d).

To see this, consider the family $\mathcal{S} = \{[0, 2^n) : n \geq 0\}$. This family is $\frac{1}{2}$ -sparse, however if we define $\mathcal{A}_\mathcal{S}(1)$ with the natural pointwise limit we obtain $\mathcal{A}_\mathcal{S}(1) = \infty$ on $[0, \infty)$.

In general, we need \mathcal{S} to have finite total measure, at least qualitatively. However, this is also not enough because there may still be issues similar to those that appear when comparing BMO and dyadic BMO. Indeed, consider

$$\mathcal{S} = \{[0, 2^{-n}) : n \geq 0\}.$$

This family is $\frac{1}{2}$ -sparse and has finite total measure, thus $\mathcal{A}_\mathcal{S}(f)$ is integrable for all $f \in L^\infty(\mathbb{R}^d)$. However, letting $I_n = [0, 2^{-n})$ and $J_n = [-2^{-n}, 2^{-n})$ for $n \geq 1$, we have

$$\begin{aligned} \frac{1}{|J_n|} \int_{J_n} \mathcal{A}_\mathcal{S}(\mathbb{1}_{[0,1)}) dx &= 2^{n-1} \sum_{m=0}^{\infty} |I_m \cap I_n| \\ &= 2^{n-1} \sum_{m=0}^n |I_n| + 2^{n-1} \sum_{m=n+1}^{\infty} |I_m| \end{aligned}$$

$$\begin{aligned}
&= 2^{n-1}(n+1)2^{-n} + 2^{n-1} \sum_{m=n+1}^{\infty} 2^{-m} \\
&= \frac{n}{2} + 1.
\end{aligned}$$

Thus

$$\begin{aligned}
\frac{1}{|J_n|} \int_{J_n} |\mathcal{A}_S(\mathbb{1}_{[0,1]}) - \langle \mathcal{A}_S(\mathbb{1}_{[0,1]}) \rangle_{J_n}| dx &\geq \frac{1}{|J_n|} \int_{-2^{-n}}^0 \left(\frac{n}{2} + 1\right) dx \\
&\geq \frac{n}{4}
\end{aligned}$$

and hence, $\mathcal{A}_S(\mathbb{1}_{[0,1]})$ is not in $\text{BMO}(\mathbb{R}^d)$.

When the sparse collection consists of cubes belonging to the same dyadic family \mathcal{D} , then we can obtain BMO(\mathcal{D}) estimates.

Theorem 4.1. *Suppose $w^{-1} \in A_1$ and let $\mathcal{S} \subseteq \mathcal{D}$ be an η -sparse collection of cubes. Then*

$$(4.1) \quad \|\mathcal{A}_S f\|_{\text{BMO}_w^1(\mathcal{D})} \lesssim \eta^{-1} [w^{-1}]_{A_\infty(\mathcal{D})} \|f\|_{L_w^\infty(\mathbb{R}^d)},$$

$$(4.2) \quad \|\mathcal{A}_S f\|_{\text{BMO}_w^\infty(\mathcal{D})} \lesssim \eta^{-1} [w^{-1}]_{A_1(\mathcal{D})} [w^{-1}]_{A_\infty(\mathcal{D})} \|f\|_{L_w^\infty(\mathbb{R}^d)}$$

and

$$(4.3) \quad \|\mathcal{A}_S f\|_{\text{BMO}_w^{1,\text{weak}}(\mathcal{D})} \lesssim \eta^{-1} \|f\|_{L_w^\infty(\mathbb{R}^d)},$$

$$(4.4) \quad \|\mathcal{A}_S f\|_{\text{BMO}_w^{\infty,\text{weak}}(\mathcal{D})} \lesssim \eta^{-1} [w^{-1}]_{A_1(\mathcal{D})} \|f\|_{L_w^\infty(\mathbb{R}^d)}.$$

Proof. Fix $Q_0 \in \mathcal{D}$ and let $c := \sum_{\substack{Q \in \mathcal{S} \\ Q \supseteq Q_0}} \langle f \rangle_{1,Q}$. Then, by Proposition 2.2 we have

$$\begin{aligned}
\langle \mathcal{A}_S f - c \rangle_{1,Q_0} &= \frac{1}{|Q_0|} \sum_{\substack{Q \in \mathcal{S} \\ Q \subseteq Q_0}} \int_Q f dx \leq \eta^{-1} \|f\|_{L_w^\infty(\mathbb{R}^d)} \frac{1}{|Q_0|} \sum_{\substack{Q \in \mathcal{S} \\ Q \subseteq Q_0}} w^{-1}(Q) \\
&\leq \eta^{-1} [w^{-1}]_{A_\infty(\mathcal{D})} \|f\|_{L_w^\infty(\mathbb{R}^d)} \langle w^{-1} \rangle_{1,Q_0}.
\end{aligned}$$

Thus, (4.1) follows from Proposition 2.5. The estimate (4.2) then follows from (2.1).

For (4.3) we note that since $\|\mathcal{A}_S(Q_0)\|_{L^1(Q_0) \rightarrow L^{1,\infty}(Q_0)} \lesssim \eta^{-1}$, we have

$$\|\mathcal{A}_S f - c\|_{L^{1,\infty}(Q_0)} = \|\mathcal{A}_S(Q_0) f\|_{L^{1,\infty}(Q_0)} \lesssim \eta^{-1} \|f\|_{L^1(Q_0)} \leq \eta^{-1} \|f\|_{L_w^\infty(\mathbb{R}^d)} w^{-1}(Q_0),$$

as desired. The inequality (4.4) then follows from (2.2). \square

4.2. Calderón-Zygmund operators. We now turn our study to BMO bounds for Calderón-Zygmund operators. In particular, we consider operators T which are weak-type $(1, 1)$ -bounded operator and are given by

$$(4.5) \quad Tf(x) = \int_{\mathbb{R}^d} K(x, y) f(y) dy \quad \text{for all } x \notin \text{supp } f$$

for all f in $L^1(\mathbb{R}^d)$ with compact support.

We will furthermore suppose that K satisfies the following smoothness condition:

$$|K(x, y) - K(z, y)| \leq \Omega\left(\frac{|x-z|}{|x-y|}\right) \frac{1}{|x-y|^d}$$

for all x, y, z satisfying $|x-y| > 2|x-z| > 0$, and where $\Omega : [0, \infty) \rightarrow [0, \infty)$ is an increasing subadditive function with $\Omega(0) = 0$. We can quantify the smoothness of the kernel in terms of the Dini condition of Ω :

$$\|\Omega\|_{\text{Dini}} := \int_0^1 \Omega(t) \frac{dt}{t}.$$

We will be interested in $L_w^\infty \rightarrow BMO_w$ estimates for T , so we should be careful about whether Tf is well-defined. Indeed, if $f \in L_w^\infty(\mathbb{R}^d)$ has compact support, and $w^{-1} \in L_{\text{loc}}^1(\mathbb{R}^d)$, then $f \in L^1(\mathbb{R}^d)$, so we can use the representation (4.5).

To study the $L^\infty \rightarrow BMO$ boundedness of such operators we will employ the following pointwise domination theorem. Originally, a similar theorem was proved by A. Lerner in [Ler10a], but we will use the following version by T. Hytönen:

Theorem 4.2 ([Hyt14, Theorem 2.3]). *For any measurable function f on a cube Q_0 in \mathbb{R}^d there exists a sparse collection $\mathcal{S} \subseteq Q_0$ such that*

$$|f(x) - m_{Q_0}(f)| \leq 2 \sum_{Q \in \mathcal{S}} \omega_\lambda(f; Q) \mathbb{1}_Q(x),$$

where $\lambda = 2^{-d-2}$ and $m_{Q_0}(f)$ is any of the possible medians of f over Q , i.e.: constants m such that

$$|\{x \in Q_0 : f(x) > m\}| \leq \frac{1}{2}|Q| \quad \text{and} \quad |\{x \in Q_0 : f(x) < m\}| \leq \frac{1}{2}|Q|.$$

The idea is to use this theorem, but applied to Tf . One can estimate $\omega_\lambda(Tf; Q)$ in terms of averages of f , in particular we have the following estimate which goes back to [JT85]:

$$(4.6) \quad \omega_\lambda(Tf; Q) \lesssim_d \sum_{m=0}^{\infty} \Omega(2^{-m}) \langle |f| \rangle_{2^m Q}.$$

This, together with Theorem 4.2, yields the following: for every compactly supported function $f \in L^1(\mathbb{R}^d)$ and every cube Q_0 there exists a sparse collection \mathcal{S} of cubes in Q_0 such that

$$(4.7) \quad |Tf(x) - m_{Q_0}(Tf)| \lesssim_d \sum_{m=0}^{\infty} \Omega(2^{-m}) \sum_{Q \in \mathcal{S}} \langle |f| \rangle_{2^m Q} \mathbb{1}_Q(x).$$

We will be able to prove certain weighted estimates for such operators, but unless $w^{-1} \in A_1$ the estimates will depend on the interaction between Ω and the weight. We will quantify this interaction with a variant of the B_2 condition

$$[w]_{B_2} = \sup_I \frac{|I|^2}{w(I)} \int_{I^c} \frac{w(x)}{|x - c_I|^2} dx,$$

where the supremum is taken over all intervals I and c_I is the center of I .

This class was introduced originally in [HMW73], and then in [MW76] where it was part of the characterization of the $L_w^\infty \rightarrow BMO_w^1$ boundedness of the Hilbert transform. In our situation we need to adapt this condition to incorporate the Dini modulus:

$$[w]_{B(\Omega)} = \sup_Q \frac{|Q|}{w(Q)} \int_{Q^c} \frac{w(x)}{|x - c_Q|^d} \Omega\left(\frac{\ell(Q)}{|x - c_Q|}\right) dx$$

where the supremum is taken over all cubes Q . Note that in dimension one and when $\Omega(x) = x$ this reduces to the B_2 condition.

We can prove a result analogous to Lemma 1 from [HMW73] which gives sufficient conditions for a weight to belong to $B(\Omega)$.

Proposition 4.3. *Let $w \in A_p$ and suppose Ω is as above, then*

$$[w]_{B(\Omega)} \lesssim_d [w]_{A_p} \int_0^1 \Omega(t) t^{-d(p-1)} \frac{dt}{t}.$$

In particular, if $w \in A_1$ then

$$[w]_{B(\Omega)} \lesssim_d [w]_{A_1} \|\Omega\|_{\text{Dini}}.$$

Proof. Fix a cube Q , and set $S_m = \{x : 2^{m-1}\ell(Q) \leq |x - c_Q| < 2^m\ell(Q)\}$

$$\begin{aligned}
\int_{Q^c} \frac{w(x)}{|x - c_Q|^d} \Omega\left(\frac{\ell(Q)}{|x - c_Q|}\right) dx &= \sum_{m=1}^{\infty} \int_{S_m} \frac{w(x)}{|x - c_Q|^d} \Omega\left(\frac{\ell(Q)}{|x - c_Q|}\right) dx \\
&\leq \sum_{m=1}^{\infty} \int_{S_m} \frac{w(x)}{(2^{m-1}\ell(Q))^d} \Omega(2^{-(m-1)}) dx \\
&= \sum_{m=0}^{\infty} \frac{\Omega(2^{-m})}{(2^m\ell(Q))^d} \int_{S_{m+1}} w(x) dx \\
&\leq \sum_{m=0}^{\infty} \frac{\Omega(2^{-m})}{(2^m\ell(Q))^d} \int_{2^{m+2}Q} w(x) dx \\
(4.8) \quad &= 2^{2d} \sum_{m=0}^{\infty} \Omega(2^{-m}) \frac{1}{|2^{m+2}Q|} \int_{2^{m+2}Q} w(x) dx.
\end{aligned}$$

Note that by definition

$$\begin{aligned}
\frac{1}{|2^{m+2}Q|} \int_{2^{m+2}Q} w dx &\leq [w]_{A_p} \langle w^{1-p'} \rangle_{2^{m+2}Q}^{1-p} \\
&\leq [w]_{A_p} 2^{(m+2)d(p-1)} \langle w \rangle_Q
\end{aligned}$$

so replacing in (4.8):

$$\begin{aligned}
\frac{|Q|}{w(Q)} \int_{Q^c} \frac{w(x)}{|x - c_Q|^d} \Omega\left(\frac{\ell(Q)}{|x - c_Q|}\right) dx &\lesssim_d [w]_{A_p} \sum_{m=0}^{\infty} \Omega(2^{-m}) 2^{md(p-1)} \\
&\approx_d [w]_{A_p} \int_0^1 \Omega(t) t^{-d(p-1)} \frac{dt}{t}
\end{aligned}$$

□

Remark. When T is the Hilbert transform (or Riesz transforms) then $\Omega(t) = t$ and then the condition becomes

$$[w]_{B(\Omega)} \lesssim_d [w]_{A_p} \int_0^1 t^{-d(p-1)} dt,$$

which is finite when $p < 1 + \frac{1}{d}$. In particular, if $w \in A_{1+\frac{1}{d}}$ then, by self-improvement we can deduce that w is in $B(\Omega)$. This recovers Lemma 1 from [HMW73].

Theorem 4.4. *Let $w^{-1} \in L_{\text{loc}}^1(\mathbb{R}^d)$ and let T be an operator as above, then*

$$(4.9) \quad \|Tf\|_{\text{BMO}_w^1(\mathbb{R}^d)} \lesssim ([1]_{B(\Omega)} + [w^{-1}]_{B(\Omega)}) [w^{-1}]_{A_\infty} \|f\|_{L_w^\infty(\mathbb{R}^d)}$$

$$(4.10) \quad \|Tf\|_{\text{BMO}_w^\infty(\mathbb{R}^d)} \lesssim [w^{-1}]_{A_1}^2 [w^{-1}]_{A_\infty} \|\Omega\|_{\text{Dini}} \|f\|_{L_w^\infty(\mathbb{R}^d)}$$

for all functions $f \in L_w^\infty(\mathbb{R}^d)$ with compact support.

Proof. Take $f \in L^\infty$ with compact support and fix a cube Q_0 . Combining (4.6) and (4.7) we have

$$|Tf(x) - m_{Q_0}(Tf)| \lesssim \sum_{Q \in \mathcal{S}} \sum_{m=0}^{\infty} \Omega(2^{-m}) \langle |f| \rangle_{2^m Q} \mathbb{1}_Q(x)$$

for some sparse collection $\mathcal{S} \subset \mathcal{D}(Q_0)$.

Assume without loss of generality that $fw \leq 1$, then

$$\int_{Q_0} |Tf - m_{Q_0}(Tf)| dx \lesssim \sum_{Q \in \mathcal{S}} \sum_{m=0}^{\infty} \Omega(2^{-m}) \langle w^{-1} \rangle_{2^m Q} |Q|$$

$$= \sum_{Q \in \mathcal{S}} \sum_{m=0}^{\infty} \Omega(2^{-m}) 2^{-md} \int_{2^m Q} w^{-1} dx.$$

We first prove (4.9). Setting $\sigma = w^{-1}$:

$$(4.11) \quad \sum_{m=0}^{\infty} \Omega(2^{-m}) 2^{-md} \int_{2^m Q} \sigma dx = \int_{\mathbb{R}^d} \sigma \sum_{m=0}^{\infty} \Omega(2^{-m}) 2^{-md} \mathbb{1}_{2^m Q} dx.$$

When $x \in Q$ then

$$\sum_{m=0}^{\infty} \Omega(2^{-m}) 2^{-md} \mathbb{1}_{2^m Q}(x) = \mathbb{1}_Q(x) \sum_{m=0}^{\infty} \Omega(2^{-m}) 2^{-md} \lesssim_d [1]_{B(\Omega)} \mathbb{1}_Q(x).$$

When $x \in 2^k Q \setminus 2^{k-1} Q$, for $k \geq 1$:

$$\begin{aligned} \sum_{m=0}^{\infty} \Omega(2^{-m}) 2^{-md} \mathbb{1}_{2^m Q}(x) &= \sum_{m=k}^{\infty} \Omega(2^{-m}) 2^{-md} \\ &\leq \Omega(2^{-k}) \sum_{m=k}^{\infty} 2^{-md} \\ &\lesssim \Omega(2^{-k}) 2^{-kd} \\ &\lesssim \Omega\left(\frac{\ell(Q)}{|x - c_Q|}\right) \frac{|Q|}{|x - c_Q|^d}. \end{aligned}$$

Putting these estimates together, and going back to (4.11):

$$\begin{aligned} \int_{\mathbb{R}^d} \sigma \sum_{m=0}^{\infty} \Omega(2^{-m}) 2^{-md} \mathbb{1}_{2^m Q} dx &\lesssim_d [1]_{B(\Omega)} \int_Q \sigma dx + |Q| \int_{Q^c} \frac{\sigma(x)}{|x - c_Q|^d} \Omega\left(\frac{\ell(Q)}{|x - c_Q|}\right) dx \\ &\approx ([1]_{B(\Omega)} + [\sigma]_{B(\Omega)}) \sigma(Q). \end{aligned}$$

Finally, we can use this estimate to finish with Proposition 2.2

$$\begin{aligned} \sum_{Q \in \mathcal{S}} \sum_{m=0}^{\infty} \Omega(2^{-m}) 2^{-md} \int_{2^m Q} \sigma &\lesssim ([1]_{B(\Omega)} + [\sigma]_{B(\Omega)}) \sum_{Q \in \mathcal{S}} \sigma(Q) \\ &\lesssim ([1]_{B(\Omega)} + [\sigma]_{B(\Omega)}) [\sigma]_{A_\infty} \sigma(Q_0). \end{aligned}$$

Putting everything together, we have shown

$$\frac{1}{w^{-1}(Q_0)} \int_{Q_0} |Tf - m_{Q_0}(Tf)| dx \lesssim ([1]_{B(\Omega)} + [w^{-1}]_{B(\Omega)}) [w^{-1}]_{A_\infty}$$

assuming $|f|w \leq 1$, so (4.9) follows. One can deduce (4.10) by combining (4.9) together with (2.1) and Proposition 4.3. \square

Acknowledgements. Z.N. is supported by the Basque Government through the BERC 2018-2021 program and by the Spanish State Research Agency through BCAM Severo Ochoa excellence accreditation SEV-2017-0718 and through project PID2020-113156GB-I00/AEI/10.13039/501100011033 funded by Agencia Estatal de Investigación and acronym ‘‘HAPDE’’.

G.R. was supported in part by Grant MICIN/AEI/PID2019-105599GB-I00.

REFERENCES

- [CPRR19] Alberto Criado, Carlos Pérez, and Israel P. Rivera-Ríos. Sharp quantitative weighted BMO estimates and a new proof of the Harboure-Macías-Segovia's extrapolation theorem. In *New trends in applied harmonic analysis. Vol. 2—harmonic analysis, geometric measure theory, and applications*, Appl. Numer. Harmon. Anal., pages 241–256. Birkhäuser/Springer, Cham, [2019] ©2019.
- [HMS88] Eleonor Harboure, Roberto A. Macías, and Carlos Segovia. Extrapolation results for classes of weights. *Amer. J. Math.*, 110(3):383–397, 1988.
- [HMW73] Richard Hunt, Benjamin Muckenhoupt, and Richard Wheeden. Weighted norm inequalities for the conjugate function and Hilbert transform. *Trans. Amer. Math. Soc.*, 176:227–251, 1973.
- [Hyt14] Tuomas P. Hytönen. The A_2 theorem: remarks and complements. In *Harmonic analysis and partial differential equations*, volume 612 of *Contemp. Math.*, pages 91–106. Amer. Math. Soc., Providence, RI, 2014.
- [JT85] Björn Jawerth and Alberto Torchinsky. Local sharp maximal functions. *J. Approx. Theory*, 43(3):231–270, 1985.
- [Ler03] Andrei K. Lerner. On the John-Strömberg characterization of BMO for nondoubling measures. *Real Anal. Exchange*, 28(2):649–660, 2002/03.
- [Ler04] Andrei K. Lerner. Weighted norm inequalities for the local sharp maximal function. *J. Fourier Anal. Appl.*, 10(5):465–474, 2004.
- [Ler10a] Andrei K. Lerner. A pointwise estimate for the local sharp maximal function with applications to singular integrals. *Bull. Lond. Math. Soc.*, 42(5):843–856, 2010.
- [Ler10b] Andrei K. Lerner. Some remarks on the Fefferman-Stein inequality. *J. Anal. Math.*, 112:329–349, 2010.
- [LN19] Andrei K. Lerner and Fedor Nazarov. Intuitive dyadic calculus: the basics. *Expo. Math.*, 37(3):225–265, 2019.
- [MW76] Benjamin Muckenhoupt and Richard L. Wheeden. Weighted bounded mean oscillation and the Hilbert transform. *Studia Math.*, 54(3):221–237, 1975/76.
- [MW78] Benjamin Muckenhoupt and Richard L. Wheeden. On the dual of weighted H^1 of the half-space. *Studia Math.*, 63(1):57–79, 1978.
- [Nie19] Zoe Nieraeth. Quantitative estimates and extrapolation for multilinear weight classes. *Math. Ann.*, 375(1-2):453–507, 2019.
- [Str79] Jan-Olov Strömberg. Bounded mean oscillation with Orlicz norms and duality of Hardy spaces. *Indiana Univ. Math. J.*, 28(3):511–544, 1979.
- [Zaa67] Adriaan Cornelis Zaanen. *Integration*. North-Holland Publishing Co., Amsterdam; Interscience Publishers John Wiley & Sons, Inc., New York, 1967. Completely revised edition of An introduction to the theory of integration.

(Zoe Nieraeth) BCAM–BASQUE CENTER FOR APPLIED MATHEMATICS, BILBAO, SPAIN
Email address: zoe.nieraeth@gmail.com

(Guillermo Rey) UNIVERSIDAD AUTÓNOMA DE MADRID
Email address: guillermo.rey@uam.es