# Cosmological Perturbations Engendered by Discrete Relativistic Species

Maksym Brilenkov<sup>a</sup>, Ezgi Canay<sup>b</sup>, Maxim Eingorn<sup>c</sup>

#### Abstract

Within the extension of the  $\Lambda$ CDM model, allowing for the presence of neutrinos or warm dark matter, we develop the analytical cosmological perturbation theory. It covers all spatial scales where the weak gravitational field regime represents a valid approximation. Discrete particles – the sources of the inhomogeneous gravitational field – may be relativistic. Similarly to the previously investigated case of nonrelativistic matter, the Yukawa interaction range is naturally incorporated into the first-order scalar metric corrections.

*Keywords:* inhomogeneous Universe, cosmological perturbations, neutrinos, warm dark matter, cosmic screening

## 1. Introduction

The extension of the conventional  $\Lambda$ CDM cosmological model by means of taking into account relativistic matter represents a very promising research area. Indeed, species with comparatively high peculiar velocities, such as neutrinos and warm dark matter particles, might have an important observable effect on the cosmic microwave background and large-scale structure probed by forth-coming surveys (see [1, 2, 3, 4] and references therein). The study of this effect offers an opportunity to estimate neutrino masses, predict physical properties of warm dark matter candidates and advance in solving of a number of astrophysical and cosmological challenges, such as, for instance, the well-known missing satellite problem (see, in particular, [5] and references therein).

A crucial role in the investigations of the large-scale structure formation belongs to cosmological simulations. Various computer codes have been developed, for example, "GADGET-4" [6] (Newtonian approximation) and "gevolution" [7]

<sup>&</sup>lt;sup>a</sup>Institute of Theoretical Astrophysics, University of Oslo, P.O. Box 1029 Blindern, N-0315 Oslo, Norway <sup>b</sup>Department of Physics, Istanbul Technical University, 34469 Maslak, Istanbul, Turkey <sup>c</sup>Department of Mathematics and Physics, North Carolina Central University, 1801 Fayetteville St., Durham, North Carolina 27707, U.S.A.

Email addresses: maksym.brilenkov@astro.uio.no (Maksym Brilenkov),

ezgicanay@itu.edu.tr (Ezgi Canay), maxim.eingorn@gmail.com (Maxim Eingorn)

(arbitrary spatial scales; ACDM plus neutrinos). Moreover, the analytical perturbation theory covering all scales has been recently formulated in [8, 9, 10] for point-like nonrelativistic masses. Most importantly, it has been demonstrated that the gravitational interaction between these masses obeys the Yukawa law with the finite time-dependent screening length (for the comparison of Yukawa and Newtonian gravitational forces in a cubic simulation box see [11]). The corresponding "screening" computer code [12] runs almost 40% faster than "gevolution" while providing very similar outcomes in the absence of neutrinos.

In this connection, it is absolutely reasonable to generalize the analytical perturbation theory, formulated for the spatially flat Universe containing the standard  $\Lambda$ CDM ingredients, to the case of relativistic particles. This is our main goal here. In Section 2 we present the energy-momentum tensor components and Einstein equations for metric corrections. These equations are then solved exactly in Section 3, and two opposite limiting cases are described in Section 4. Finally, we summarize the results in concluding Section 5.

# 2. Perturbed spacetime

We begin with presenting the perturbed Friedmann-Lemaître-Robertson-Walker metric in the Poisson gauge, that is

$$ds^{2} = a^{2} \left[ (1+2\Phi)d\eta^{2} + 2B_{\alpha}d\eta dx^{\alpha} + (-\delta_{\alpha\beta} + 2\Psi\delta_{\alpha\beta}) dx^{\alpha} dx^{\beta} \right].$$
(1)

In the above expression,  $a(\eta)$  stands for the scale factor, where  $\eta$  is the conformal time, and  $x^{\alpha}$  are the comoving coordinates for  $\alpha = 1, 2, 3$ . The functions  $\Psi$  and  $\Phi$  denote the first-order scalar perturbations whereas  $B_{\alpha}$  is the first-order vector perturbation subject to the gauge condition  $\delta^{\alpha\beta}\partial B_{\alpha}/\partial x^{\beta} = 0$ . Investigation of tensor modes is beyond the scope of our study, therefore the corresponding part in the perturbed metric is disregarded.

Meanwhile, the components of the energy-momentum tensor for a system of point-like particles with comoving radius-vectors  $\mathbf{r}_n$ , momenta  $\mathbf{q}_n$  and masses  $m_n$  are given by (see [13] and particularly Eqs. (3.7), (3.10) and (3.8) in [7])

$$T_0^0 = \frac{c^2}{a^4} \sum_n \sqrt{q_n^2 + a^2 m_n^2} \delta\left(\mathbf{r} - \mathbf{r}_n\right) + \frac{c^2}{a^4} \overline{\sum_n \frac{4q_n^2 + 3a^2 m_n^2}{\sqrt{q_n^2 + a^2 m_n^2}}} \delta\left(\mathbf{r} - \mathbf{r}_n\right) \Psi, \quad (2)$$

$$T_{\alpha}^{0} = -\frac{c^{2}}{a^{4}} \sum_{n} q_{n}^{\alpha} \delta\left(\mathbf{r} - \mathbf{r}_{n}\right) , \qquad (3)$$

$$T_{\beta}^{\alpha} = - \frac{c^{2}}{a^{4}} \sum_{n} \frac{q_{n}^{\alpha} q_{n}^{\beta}}{\sqrt{q_{n}^{2} + a^{2} m_{n}^{2}}} \delta\left(\mathbf{r} - \mathbf{r}_{n}\right) - \frac{c^{2}}{3a^{4}} \overline{\sum_{n} \frac{q_{n}^{2} \left(4q_{n}^{2} + 5a^{2} m_{n}^{2}\right)}{\left(q_{n}^{2} + a^{2} m_{n}^{2}\right)^{3/2}}} \delta\left(\mathbf{r} - \mathbf{r}_{n}\right)} \delta_{\alpha\beta} \Psi, \qquad (4)$$

where  $q_n^2 \equiv \delta_{\alpha\beta} q_n^{\alpha} q_n^{\beta}$ . As we allow for the contribution of relativistic species in our current approach, we do not demand that peculiar velocities of particles are much smaller than the speed of light *c*. Since, in the weak gravitational field regime, the scalar and vector perturbations are small everywhere, the quantities multiplied by them are replaced by their average values, following the well-justified reasoning in [8, 9, 14]. Namely, the multiplications of metric corrections by fluctuations of these quantities are not regarded as significant sources of the inhomogeneous gravitational field. The corresponding nonzero average components

$$\overline{T_0^0} = \frac{c^2}{a^4} \overline{\sum_n \sqrt{q_n^2 + a^2 m_n^2} \delta\left(\mathbf{r} - \mathbf{r}_n\right)} \equiv \overline{\varepsilon}, \qquad (5)$$

$$\overline{T_{\beta}^{\alpha}} = -\frac{c^2}{3a^4} \overline{\sum_{n} \frac{q_n^2}{\sqrt{q_n^2 + a^2 m_n^2}}} \delta\left(\mathbf{r} - \mathbf{r}_n\right) \delta_{\alpha\beta} \equiv -\overline{p} \delta_{\alpha\beta} \tag{6}$$

contribute to the right-hand side of the background Friedmann equations:

$$\frac{3\mathcal{H}^2}{a^2} = \kappa \overline{\varepsilon} + \Lambda \,, \tag{7}$$

$$\frac{2\mathcal{H}' + \mathcal{H}^2}{a^2} = -\kappa \overline{p} + \Lambda \,, \tag{8}$$

where  $\kappa \equiv 8\pi G_N/c^4$  (with  $G_N$  representing the gravitational constant),  $\Lambda$  is the cosmological constant, and  $\mathcal{H}$  denotes the Hubble parameter defined as  $\mathcal{H} \equiv a'/a \equiv (da/d\eta)/a$ .

According to Eq. (1) as well as the energy-momentum tensor components presented above, Einstein equations (linearized with respect to the scalar and vector perturbations) yield

$$\Delta \Psi - 3\mathcal{H} \left( \Psi' + \mathcal{H} \Phi \right) = \frac{\kappa a^2}{2} \left( \frac{c^2}{a^4} \sum_n \sqrt{q_n^2 + a^2 m_n^2} \delta \left( \mathbf{r} - \mathbf{r}_n \right) - \overline{\varepsilon} \right) + \frac{3\kappa a^2}{2} \left( \overline{\varepsilon} + \overline{p} \right) \Psi, \qquad (9)$$

$$\frac{1}{4} \triangle B_{\alpha} + \frac{\partial}{\partial x^{\alpha}} \left( \Psi' + \mathcal{H} \Phi \right) = -\frac{\kappa c^2}{2a^2} \sum_{n} q_n^{\alpha} \delta \left( \mathbf{r} - \mathbf{r}_n \right) \,, \tag{10}$$

$$\frac{\partial^2 \left(\Phi - \Psi\right)}{\partial x^\beta \partial x^\gamma} - \frac{1}{3} \Delta \left(\Phi - \Psi\right) \delta_{\beta\gamma} - \mathcal{H}\left(\frac{\partial B_\gamma}{\partial x^\beta} + \frac{\partial B_\beta}{\partial x^\gamma}\right) - \frac{1}{2} \left(\frac{\partial B_\gamma}{\partial x^\beta} + \frac{\partial B_\beta}{\partial x^\gamma}\right)' \\ = -\frac{\kappa c^2}{a^2} \sum_n \frac{q_n^\beta q_n^\gamma - q_n^2 \delta_{\beta\gamma}/3}{\sqrt{q_n^2 + a^2 m_n^2}} \delta\left(\mathbf{r} - \mathbf{r}_n\right) \,, \tag{11}$$

where the Laplace operator  $\Delta \equiv \delta_{\alpha\beta} \partial^2 / \partial x^{\alpha} \partial x^{\beta}$ .

## 3. Analytical expressions for metric corrections

As it follows from a scalar-vector decomposition of Eq. (11), the difference  $\Phi - \Psi$  satisfies the equation

$$\triangle \triangle (\Phi - \Psi) = -\frac{3\kappa c^2}{2a^2} \sum_{n} \frac{q_n^\beta q_n^\gamma - q_n^2 \delta_{\beta\gamma}/3}{\sqrt{q_n^2 + a^2 m_n^2}} \frac{\partial^2}{\partial x^\beta \partial x^\gamma} \delta\left(\mathbf{r} - \mathbf{r}_n\right) , \qquad (12)$$

whence

$$\Phi = \Psi - \frac{3\kappa c^2}{16\pi a^2} \sum_{n} \frac{q_n^{\beta} q_n^{\gamma} - q_n^2 \delta_{\beta\gamma}/3}{\sqrt{q_n^2 + a^2 m_n^2}} \frac{(x^{\gamma} - x_n^{\gamma}) \left(x^{\beta} - x_n^{\beta}\right)}{\left|\mathbf{r} - \mathbf{r}_n\right|^3} \,.$$
(13)

Similarly, from Eq. (10) we obtain

$$\Psi' + \mathcal{H}\Phi = -\frac{\kappa c^2}{2a}\Xi,\qquad(14)$$

where

$$\Xi = \frac{1}{4\pi a} \sum_{n} \frac{q_n^{\alpha} (x^{\alpha} - x_n^{\alpha})}{|\mathbf{r} - \mathbf{r}_n|^3},$$
(15)

which, when substituted into Eq. (9), yields

$$\Delta \Psi - \frac{a^2}{\lambda^2} \Psi = \frac{\kappa a^2}{2} \left( \frac{c^2}{a^4} \sum_n \sqrt{q_n^2 + a^2 m_n^2} \delta\left(\mathbf{r} - \mathbf{r}_n\right) - \overline{\varepsilon} \right) - \frac{3\mathcal{H}\kappa c^2}{2a} \Xi.$$
(16)

Here we introduce the screening length  $\lambda$  by means of the relationship

$$\frac{a^2}{\lambda^2} = \frac{3\kappa a^2}{2} \left(\overline{\varepsilon} + \overline{p}\right) \,. \tag{17}$$

It is important to emphasize that Eq. (17) can be rewritten with the help of the background Friedmann equations (7) and (8):

$$\lambda = \frac{a}{\sqrt{3\left(\mathcal{H}^2 - \mathcal{H}'\right)}},\tag{18}$$

in complete agreement with Eq. (3.7) in [8]. Earlier, this universal presentation of the screening length  $\lambda$ , predicted in [8], was confirmed for various cosmological models in [15, 16, 17, 18] ( $\Lambda$ CDM plus extra perfect fluids with linear and nonlinear equations of state) and [19] (nonzero spatial curvature). And now we confirm it also for the model which includes relativistic species.

The Helmholtz equation (16) has the analytical solution (see [8] where resembling Eq. (2.27) is analyzed):

$$\Psi = \frac{\kappa\lambda^{2}\overline{\varepsilon}}{2} - \frac{\kappa c^{2}}{8\pi a^{2}} \sum_{n} \frac{\sqrt{q_{n}^{2} + a^{2}m_{n}^{2}}}{|\mathbf{r} - \mathbf{r}_{n}|} \exp\left(-\frac{a|\mathbf{r} - \mathbf{r}_{n}|}{\lambda}\right) + \frac{3\kappa c^{2}\lambda^{2}\mathcal{H}}{8\pi a^{4}} \sum_{n} \frac{q_{n}^{\alpha}\left(x^{\alpha} - x_{n}^{\alpha}\right)}{|\mathbf{r} - \mathbf{r}_{n}|^{3}} \left[1 - \left(1 + \frac{a|\mathbf{r} - \mathbf{r}_{n}|}{\lambda}\right) \exp\left(-\frac{a|\mathbf{r} - \mathbf{r}_{n}|}{\lambda}\right)\right].$$
(19)

For the vector perturbation, we return to Eq. (10) and obtain

$$\frac{1}{4} \triangle B_{\alpha} = -\frac{\kappa c^2}{2a} \left( \frac{1}{a} \sum_{n} q_n^{\alpha} \delta \left( \mathbf{r} - \mathbf{r}_n \right) - \frac{\partial \Xi}{\partial x^{\alpha}} \right), \qquad (20)$$

whence

$$B_{\alpha} = \frac{\kappa c^2}{4\pi a^2} \sum_{n} \left[ \frac{q_n^{\alpha}}{|\mathbf{r} - \mathbf{r}_n|} + \frac{q_n^{\beta} \left(x^{\beta} - x_n^{\beta}\right)}{|\mathbf{r} - \mathbf{r}_n|^3} \left(x^{\alpha} - x_n^{\alpha}\right) \right].$$
(21)

Finally, we write down the equation of motion for the k-th particle of the system as well as the connection between its momentum and peculiar velocity components (see Eq. (3.5) and (3.4) in [7], respectively):

$$(q_k^{\gamma})' = -\sqrt{q_k^2 + a^2 m_k^2} \frac{\partial \Phi}{\partial x^{\gamma}} - \frac{q_k^2}{\sqrt{q_k^2 + a^2 m_k^2}} \frac{\partial \Psi}{\partial x^{\gamma}} - q_k^{\alpha} \frac{\partial B_{\alpha}}{\partial x^{\gamma}}, \qquad (22)$$

$$\tilde{v}_{k}^{\gamma} = \frac{q_{k}^{\gamma}}{\sqrt{q_{k}^{2} + a^{2}m_{k}^{2}}} + \frac{q_{k}^{\gamma}}{\sqrt{q_{k}^{2} + a^{2}m_{k}^{2}}} \left(2 - \frac{q_{k}^{2}}{q_{k}^{2} + a^{2}m_{k}^{2}}\right)\Psi + \frac{q_{k}^{\gamma}}{\sqrt{q_{k}^{2} + a^{2}m_{k}^{2}}}\Phi + B_{\gamma},$$
(23)

where  $\tilde{v}_k^{\gamma} \equiv (x_k^{\gamma})'$ . All metric corrections and their spatial derivatives in Eqs. (22) and (23) are calculated at the point  $\mathbf{r} = \mathbf{r}_k$ , and the summation is over  $n \neq k$ .

### 4. Asymptotic behavior

In the nonrelativistic limit, neglecting all those terms which are quadratic or higher-order in  $q_n^{\alpha}$ , one can reduce Eqs. (13) and (19) to the following one:

$$\Phi = \Psi = \frac{\kappa \lambda^2 \overline{\varepsilon}}{2} + \Phi_{\text{cold}} \,, \tag{24}$$

where

$$\Phi_{\text{cold}} = -\frac{\kappa c^2}{8\pi a} \sum_{n} \frac{m_n}{|\mathbf{r} - \mathbf{r}_n|} \exp\left(-\frac{a|\mathbf{r} - \mathbf{r}_n|}{\lambda}\right) + \frac{3\kappa c^2 \lambda^2 \mathcal{H}}{8\pi a^4} \sum_{n} \frac{q_n^{\alpha} \left(x^{\alpha} - x_n^{\alpha}\right)}{|\mathbf{r} - \mathbf{r}_n|^3} \left[1 - \left(1 + \frac{a|\mathbf{r} - \mathbf{r}_n|}{\lambda}\right) \exp\left(-\frac{a|\mathbf{r} - \mathbf{r}_n|}{\lambda}\right)\right].$$
(25)

At the same time, in Eqs. (22) and (23) one may additionally disregard those terms which contain products of  $q_n^{\alpha}$  and the metric corrections (or their spatial derivatives):

$$(q_k^{\gamma})' = -am_k \frac{\partial \Phi}{\partial x^{\gamma}}, \quad \tilde{v}_k^{\gamma} = \frac{q_k^{\gamma}}{am_k} + B_{\gamma}.$$
 (26)

In the opposite, ultrarelativistic limit, when  $q_n \gg am_n$ , Eqs. (13) and (19) are modified as follows:

$$\Phi = \frac{\kappa \lambda^2 \overline{\varepsilon}}{2} + \Phi_{\text{hot}}, \quad \Psi = \frac{\kappa \lambda^2 \overline{\varepsilon}}{2} + \Psi_{\text{hot}}, \quad (27)$$

where

$$\Phi_{\text{hot}} = \Psi_{\text{hot}} - \frac{3\kappa c^2}{16\pi a^2} \sum_{n} \frac{q_n^{\beta} q_n^{\gamma} - q_n^2 \delta_{\beta\gamma}/3}{q_n} \frac{\left(x^{\gamma} - x_n^{\gamma}\right) \left(x^{\beta} - x_n^{\beta}\right)}{\left|\mathbf{r} - \mathbf{r}_n\right|^3} , \qquad (28)$$

$$\Psi_{\text{hot}} = -\frac{\kappa c^2}{8\pi a^2} \sum_n \frac{q_n}{|\mathbf{r} - \mathbf{r}_n|} \exp\left(-\frac{a|\mathbf{r} - \mathbf{r}_n|}{\lambda}\right) + \frac{3\kappa c^2 \lambda^2 \mathcal{H}}{8\pi a^4} \sum_n \frac{q_n^{\alpha} \left(x^{\alpha} - x_n^{\alpha}\right)}{|\mathbf{r} - \mathbf{r}_n|^3} \left[1 - \left(1 + \frac{a|\mathbf{r} - \mathbf{r}_n|}{\lambda}\right) \exp\left(-\frac{a|\mathbf{r} - \mathbf{r}_n|}{\lambda}\right)\right].$$
(29)

In this case Eqs. (22) and (23) take the form

$$(q_k^{\gamma})' = -q_k \frac{\partial (\Phi + \Psi)}{\partial x^{\gamma}} - q_k^{\alpha} \frac{\partial B_{\alpha}}{\partial x^{\gamma}}, \quad \tilde{v}_k^{\gamma} = \frac{q_k^{\gamma}}{q_k} + \frac{q_k^{\gamma}}{q_k} (\Phi + \Psi) + B_{\gamma}.$$
(30)

Irrespectively of the comparison between  $q_n$  and  $am_n$ , Eq. (21) preserves its form. It is also interesting to note that it is a common situation during the evolution of the Universe when there are simultaneously both types of particles, nonrelativistic and ultrarelativistic, or even three types, with the third one undergoing the transition (or, in simple words, cooling down).

#### 5. Conclusion

Now we summarize our results obtained in the framework of the analyzed cosmological model ( $\Lambda$ CDM plus relativistic species):

• the exact analytical expressions (13), (19) and (21) have been derived for the first-order scalar and vector perturbations generated by discrete particles, and their asymptotic behavior has been studied;

• these expressions are valid at arbitrary scales and represent a direct generalization of their counterparts from [8] to the case of a model which includes neutrinos or warm dark matter (with massless particles also allowed);

• the gravitational interaction between discrete species is characterized by the finite time-dependent screening length, and its universal presentation from [8] has been corroborated. The derived metric corrections along with the corresponding equations of motion are ready to be used in high-precision cosmological simulations and investigations of the role of neutrinos or warm dark matter in the structure formation. Taking into account the indubitable success of the "screening" computer code [12] for nonrelativistic matter, we expect a similar positive outcome of the simulations based on the current analysis. Such promising simulations will be in the focus of our future work.

#### Acknowledgments

The work of Maksym Brilenkov was supported by funding from the European Union's Horizon 2020 research and innovation programme under grant agreement number 772253 (ERC; bits2cosmology).

# References

- J. Lesgourgues, S. Pastor, Massive neutrinos and cosmology. Phys. Rept. 429, 307 (2006), arXiv:astro-ph/0603494
- [2] H.J. de Vega, N.G. Sanchez, Cosmological evolution of warm dark matter fluctuations I: Efficient computational framework with Volterra integral equations. Phys. Rev. D 85, 043516 (2012), arXiv:1111.0290 [astro-ph.CO]
- [3] H. Dupuy, F. Bernardeau, Describing massive neutrinos in cosmology as a collection of independent flows. JCAP 01, 030 (2014), arXiv:1311.5487 [astro-ph.CO]
- [4] Z. Slepian, S.K.N. Portillo, Too hot to handle? Analytic solutions for massive neutrino or warm dark matter cosmologies. MNRAS 478, 516 (2018), arXiv:1710.01785 [astro-ph.CO]
- [5] J. Mastache, A. de la Macorra, Analytic fluid approximation for warm dark matter. arXiv:1909.05132 [astro-ph.CO]
- [6] V. Springel, R. Pakmor, O. Zier, M. Reinecke, Simulating cosmic structure formation with the GADGET-4 code. MNRAS 506, 2871 (2021), arXiv:2010.03567 [astro-ph.IM]
- [7] J. Adamek, D. Daverio, R. Durrer, M. Kunz, gevolution: a cosmological N-body code based on General Relativity. JCAP 07, 053 (2016), arXiv:1604.06065 [astro-ph.CO]
- [8] M. Eingorn, First-order cosmological perturbations engendered by pointlike masses. Astrophys. J. 825, 84 (2016), arXiv:1509.03835 [gr-qc]

- R. Brilenkov, M. Eingorn, Second-order cosmological perturbations engendered by point-like masses. Astrophys. J. 845, 153 (2017), arXiv:1703.10282 [gr-qc]
- [10] E. Canay, M. Eingorn, Duel of cosmological screening lengths. Phys. Dark Univ. 29, 100565 (2020), arXiv:2002.00437 [gr-qc]
- [11] E. Canay, M. Eingorn, Yukawa vs. Newton: gravitational forces in a cubic cosmological simulation box. Eur. Phys. J. Plus 137, 397 (2022), arXiv:2107.01052 [gr-qc]
- [12] M. Eingorn, A.E. Yükselci, A. Zhuk, Screening vs. gevolution: in chase of a perfect cosmological simulation code. Phys. Lett. B 826, 136911 (2022), arXiv:2106.07638 [gr-qc]
- [13] L.D. Landau, E.M. Lifshitz, Course of Theoretical Physics Series, Vol. 2, The Classical Theory of Fields. Oxford: Oxford Pergamon Press (2000)
- [14] N.E. Chisari, M. Zaldarriaga, Connection between Newtonian simulations and general relativity. Phys. Rev. D 83, 123505 (2011), arXiv:1101.3555 [astro-ph.CO]
- [15] M. Eingorn, R. Brilenkov, Perfect fluids with  $\omega = \text{const}$  as sources of scalar cosmological perturbations. Phys. Dark Univ. 17, 63 (2017), arXiv:1509.08181 [gr-qc]
- [16] M. Eingorn, C. Kiefer, A. Zhuk, Scalar and vector perturbations in a universe with discrete and continuous matter sources. JCAP 09, 032 (2016), arXiv:1607.03394 [gr-qc]
- [17] M. Eingorn, C. Kiefer, A. Zhuk, Cosmic screening of the gravitational interaction. Int. J. Mod. Phys. D 26, 1743012 (2017), arXiv:1711.01759 [gr-qc]
- [18] E. Canay, R. Brilenkov, M. Eingorn, A.S. Arapoğlu, A. Zhuk, Scalar and vector perturbations in a universe with nonlinear perfect fluid. EPJC 81, 246 (2021); arXiv:2011.05914 [gr-qc]
- [19] M. Eingorn, A.E. Yükselci, A. Zhuk, Effect of the spatial curvature of the Universe on the form of the gravitational potential. EPJC 79, 655 (2019), arXiv:1905.09502 [gr-qc]