

SELECTION GAMES WITH MINIMAL USCO MAPS

CHRISTOPHER CARUVANA

ABSTRACT. We establish relationships between various topological selection games involving the space of minimal usco maps with various topologies, including the topology of pointwise convergence and the topology of uniform convergence on compact sets, and the underlying domain using full- and limited-information strategies. We also tie these relationships to analogous results related to spaces of continuous functions. The primary games we consider include Rothberger-like games, generalized point-open games, strong fan-tightness games, Tkachuk's closed discrete selection game, and Gruenhage's W -games.

1. INTRODUCTION

Minimal upper-semicontinuous compact-valued functions have a rich history, apparently arising from the study of holomorphic functions and their so-called cluster sets; see [10]. The phrase *minimal usco* was coined by Christensen [7], where a topological game similar to the Banach-Mazur game was considered. In this paper, using some techniques similar to those of Holá and Holý [17], we tie connections between a space X and the space of minimal usco maps with the topology of uniform convergence on certain kinds of subspaces of X similar in spirit to those appearing in [9, 2, 3]; in particular, most of the results come in the form of selection game equivalences or dualities, which rely on a variety of game-related results from [8, 33, 32, 34, 2, 3]. We also tie some of these results to existing results relating topological properties of a space X with the space of continuous real-valued functions on X with various topologies.

Consequences of these results include Corollary 46, which captures [17, Cor. 4.5], that states that X is hemicompact if and only if $MU_k(X)$, the space of minimal usco maps into \mathbb{R} on X with the topology of uniform convergence on compact subsets, is metrizable. Corollary 46 also shows that X is hemicompact if and only if $MU_k(X)$ is not discretely selective. Corollary 51 contains the assertion that X is k -Rothberger if and only if $MU_k(X)$ has strong countable fan-tightness at $\mathbf{0}$, the constant $\{0\}$ function. We end with Corollary 55, which characterizes a space having a countable so-called weak k -covering number in terms of strategic information in selection games.

Date: June 29, 2022.

2010 *Mathematics Subject Classification.* 91A44, 54C60, 54C35, 54D20, 54B20.

Key words and phrases. minimal usco maps, topological selection principles, topology of uniform convergence on compacta.

2. PRELIMINARIES

We use the word *space* to mean *topological space*. When the parent space is understood from context, we use the notation $\text{int}(A)$, $\text{cl}(A)$, and ∂A to denote the interior, closure, and boundary of A , respectively. If we must specify the topological space X , we use $\text{int}_X(A)$, $\text{cl}_X(A)$, and $\partial_X A$.

Given a function $f : X \rightarrow Y$, we denote the graph of f by $\text{gr}(f) = \{\langle x, f(x) \rangle : x \in X\}$. For a set X , we let $\wp(X)$ denote the set of subsets of X and $\wp^+(X) = \wp(X) \setminus \{\emptyset\}$. For sets X and Y , we let

$$\text{Fn}(X, Y) = \bigcup_{A \in \wp^+(X)} Y^A;$$

that is, $\text{Fn}(X, Y)$ is the collection of all Y -valued functions defined on non-empty subsets of X .

When a set X is implicitly serving as the parent space in context, given $A \subseteq X$, we will let $\mathbf{1}_A$ be the indicator function for A ; that is, $\mathbf{1}_A : X \rightarrow \{0, 1\}$ is defined by the rule

$$\mathbf{1}_A(x) = \begin{cases} 1, & x \in A \\ 0, & x \notin A \end{cases}.$$

For any set X , we let $X^{<\omega}$ denote the set of finite sequences of X , $[X]^{<\omega}$ denote the set of finite subsets of X , and, for any cardinal κ , $[X]^\kappa$ denote the set of κ -sized subsets of X .

For a space X , we let $K(X)$ denote the set of all non-empty compact subsets of X . We let $\mathbb{K}(X)$ denote the set $K(X)$ endowed with the Vietoris topology; that is, the topology with basis consisting of sets of the form

$$[U_1, U_2, \dots, U_n] = \left\{ K \in \mathbb{K}(X) : K \subseteq \bigcup_{j=1}^n U_j \wedge K^n \cap \prod_{j=1}^n U_j \neq \emptyset \right\}.$$

For more about this topology, see [24].

Definition 1. For a set X , we say that a family $\mathcal{A} \subseteq \wp^+(X)$ is an **ideal of sets** if

- for $A, B \in \mathcal{A}$, $A \cup B \in \mathcal{A}$, and
- for every $x \in X$, $\{x\} \in \mathcal{A}$.

If X is a space, we say that an ideal of sets \mathcal{A} is an **ideal of closed sets** if \mathcal{A} consists of closed sets

Throughout, we will assume that any ideal of closed sets under consideration doesn't contain the entire space X . Two ideals of closed sets of primary interest are

- the collection of non-empty finite subsets of an infinite space X and
- the collection of non-empty compact subsets of a non-compact space X .

2.1. Selection Games. General topological games have a long history, a lot of which can be gathered from Telgársky's survey [31]. In this paper, we will be dealing only with single-selection games of countable length.

Definition 2. Given sets \mathcal{A} and \mathcal{B} , we define the **single-selection game** $G_1(\mathcal{A}, \mathcal{B})$ as follows.

- For each $n \in \omega$, One chooses $A_n \in \mathcal{A}$ and Two responds with $x_n \in A_n$.
- Two is declared the winner if $\{x_n : n \in \omega\} \in \mathcal{B}$. Otherwise, One wins.

Definition 3. We define strategies of various strength below.

- We use two forms of *full-information* strategies.
 - A **strategy for player One** in $G_1(\mathcal{A}, \mathcal{B})$ is a function $\sigma : (\bigcup \mathcal{A})^{<\omega} \rightarrow \mathcal{A}$. A strategy σ for One is called **winning** if whenever $x_n \in \sigma\langle x_k : k < n \rangle$ for all $n \in \omega$, $\{x_n : n \in \omega\} \notin \mathcal{B}$. If player One has a winning strategy, we write $I \uparrow G_1(\mathcal{A}, \mathcal{B})$.
 - A **strategy for player Two** in $G_1(\mathcal{A}, \mathcal{B})$ is a function $\tau : \mathcal{A}^{<\omega} \rightarrow \bigcup \mathcal{A}$. A strategy τ for Two is **winning** if whenever $A_n \in \mathcal{A}$ for all $n \in \omega$, $\{\tau(A_0, \dots, A_n) : n \in \omega\} \in \mathcal{B}$. If player Two has a winning strategy, we write $II \uparrow G_1(\mathcal{A}, \mathcal{B})$.
- We use two forms of *limited-information* strategies.
 - A **predetermined strategy** for One is a strategy which only considers the current turn number. We call this kind of strategy predetermined because One is not reacting to Two's moves. Formally it is a function $\sigma : \omega \rightarrow \mathcal{A}$. If One has a winning predetermined strategy, we write $I \uparrow_{\text{pre}} G_1(\mathcal{A}, \mathcal{B})$.
 - A **Markov strategy** for Two is a strategy which only considers the most recent move of player One and the current turn number. Formally it is a function $\tau : \mathcal{A} \times \omega \rightarrow \bigcup \mathcal{A}$. If Two has a winning Markov strategy, we write $II \uparrow_{\text{mark}} G_1(\mathcal{A}, \mathcal{B})$.

The reader may be more familiar with selection principles than selection games. For more details on selection principles and relevant references, see [20, 28].

Definition 4. Let \mathcal{A} and \mathcal{B} be collections. The **single-selection principle** $S_1(\mathcal{A}, \mathcal{B})$ for a space X is the following property. Given any $A \in \mathcal{A}^\omega$, there exists $\vec{x} \in \prod_{n \in \omega} A_n$ so that $\{\vec{x}_n : n \in \omega\} \in \mathcal{B}$.

As mentioned in [8, Prop. 15], $S_1(\mathcal{A}, \mathcal{B})$ holds if and only if $I \not\uparrow_{\text{pre}} G_1(\mathcal{A}, \mathcal{B})$. Hence, we may establish equivalences between certain selection principles by addressing the corresponding selection games.

Definition 5. For a space X , an open cover \mathcal{U} of X is said to be **non-trivial** if $\emptyset \notin \mathcal{U}$ and $X \notin \mathcal{U}$.

Definition 6. Let X be a space and \mathcal{A} be a set of closed subsets of X . We say that a non-trivial cover \mathcal{U} of X is an **\mathcal{A} -cover** if, for every $A \in \mathcal{A}$, there exists $U \in \mathcal{U}$ so that $A \subseteq U$.

Definition 7. For a collection \mathcal{A} , we let $\neg\mathcal{A}$ denote the collection of sets which are not in \mathcal{A} . We also define the following classes for a space X and a collection \mathcal{A} of closed subsets of X .

- \mathcal{T}_X is the family of all proper non-empty open subsets of X .
- For $x \in X$, $\mathcal{N}_{X,x} = \{U \in \mathcal{T}_X : x \in U\}$.
- For $A \in \wp^+(X)$, $\mathcal{N}_X(A) = \{U \in \mathcal{T}_X : A \subseteq U\}$.
- $\mathcal{N}_X[\mathcal{A}] = \{\mathcal{N}_X(A) : A \in \mathcal{A}\}$,
- CD_X is the set of all closed discrete subsets of X .
- \mathcal{D}_X is the set of all dense subsets of X .
- For $x \in X$, $\Omega_{X,x} = \{A \subseteq X : x \in \text{cl}(A)\}$.
- For $x \in X$, $\Gamma_{X,x}$ is the set of all sequences of X converging to x .
- \mathcal{O}_X is the set of all non-trivial open covers of X .
- $\mathcal{O}_X(\mathcal{A})$ is the set of all \mathcal{A} -covers.
- $\Lambda_X(\mathcal{A})$ is the set of all \mathcal{A} -covers \mathcal{U} with the property that, for every $A \in \mathcal{A}$, $\{U \in \mathcal{U} : A \subseteq U\}$ is infinite.
- $\Gamma_X(\mathcal{A})$ is the set of all countable \mathcal{A} -covers \mathcal{U} with the property that, for every $A \in \mathcal{A}$, $\{U \in \mathcal{U} : A \subseteq U\}$ is co-finite.

Note that, in our notation, $\mathcal{O}_X([X]^{<\omega})$ is the set of all ω -covers of X , which we will denote by Ω_X , and that $\mathcal{O}_X(K(X))$ is the set of all k -covers of X , which we will denote by \mathcal{K}_X . We also use $\Gamma_\omega(X)$ to denote $\Gamma_X([X]^{<\omega})$ and $\Gamma_k(X)$ to denote $\Gamma_X(K(X))$.

Also note that $\text{S}_1(\mathcal{O}_X, \mathcal{O}_X)$ is the Rothberger property and $\text{G}_1(\mathcal{O}_X, \mathcal{O}_X)$ is the Rothberger game. If we let $\mathbb{P}_X = \{\mathcal{N}_{X,x} : x \in X\}$, then $\text{G}_1(\mathbb{P}_X, \neg\mathcal{O})$ is isomorphic to the point-open game studied by Galvin [12] and Telgársky [30]. The games $\text{G}_1(\mathcal{N}_{X,x}, \neg\Gamma_{X,x})$ and $\text{G}_1(\mathcal{N}_{X,x}, \neg\Omega_{X,x})$ are two variants of Gruenhage's W -game (see [14]). We refer to $\text{G}_1(\mathcal{N}_{X,x}, \neg\Gamma_{X,x})$ as Gruenhage's converging W -game and $\text{G}_1(\mathcal{N}_{X,x}, \neg\Omega_{X,x})$ as Gruenhage's clustering W -game. The games $\text{G}_1(\mathcal{T}_X, \neg\Omega_{X,x})$ and $\text{G}_1(\mathcal{T}_X, \text{CD}_X)$ were introduced by Tkachuk (see [33, 34]) and tied to Gruenhage's W -games in [34, 9]. The strong countable dense fan-tightness game at x is $\text{G}_1(\mathcal{D}_X, \Omega_{X,x})$ and the strong countable fan-tightness game at x is $\text{G}_1(\Omega_{X,x}, \Omega_{X,x})$ (see [1]).

Lemma 8 (See [3, Lemma 4]). For a space X and an ideal of closed sets \mathcal{A} of X , $\mathcal{O}_X(\mathcal{A}) = \Lambda_X(\mathcal{A})$.

In what follows, we say that \mathcal{G} is a *selection game* if there exist classes \mathcal{A}, \mathcal{B} so that $\mathcal{G} = \text{G}_1(\mathcal{A}, \mathcal{B})$.

Since we work with full- and limited-information strategies, we reflect this in our definitions of game equivalence and duality.

Definition 9. We say that two selection games \mathcal{G} and \mathcal{H} are **equivalent**, denoted $\mathcal{G} \equiv \mathcal{H}$, if the following hold:

- $\text{II} \underset{\text{mark}}{\uparrow} \mathcal{G} \iff \text{II} \underset{\text{mark}}{\uparrow} \mathcal{H}$
- $\text{II} \underset{\text{mark}}{\uparrow} \mathcal{G} \iff \text{II} \underset{\text{mark}}{\uparrow} \mathcal{H}$
- $\text{I} \underset{\text{pre}}{\uparrow} \mathcal{G} \iff \text{I} \underset{\text{pre}}{\uparrow} \mathcal{H}$
- $\text{I} \underset{\text{pre}}{\uparrow} \mathcal{G} \iff \text{I} \underset{\text{pre}}{\uparrow} \mathcal{H}$

We also use a preorder on selection games.

Definition 10. Given selection games \mathcal{G} and \mathcal{H} , we say that $\mathcal{G} \leq_{\text{II}} \mathcal{H}$ if the following implications hold:

- $\text{II} \uparrow \mathcal{G} \implies \text{II} \uparrow \mathcal{H}$
- $\text{II} \uparrow_{\text{mark}} \mathcal{G} \implies \text{II} \uparrow_{\text{mark}} \mathcal{H}$
- $\text{I} \uparrow \mathcal{G} \implies \text{I} \uparrow \mathcal{H}$
- $\text{I} \uparrow_{\text{pre}} \mathcal{G} \implies \text{I} \uparrow_{\text{pre}} \mathcal{H}$

Note that \leq_{II} is transitive and that if $\mathcal{G} \leq_{\text{II}} \mathcal{H}$ and $\mathcal{H} \leq_{\text{II}} \mathcal{G}$, then $\mathcal{G} \equiv \mathcal{H}$. We use the subscript of II since each implication in the definition of \leq_{II} is related to a transference of winning plays by Two.

Definition 11. We say that two selection games \mathcal{G} and \mathcal{H} are **dual** if the following hold:

- $\text{I} \uparrow \mathcal{G} \iff \text{II} \uparrow \mathcal{H}$
- $\text{II} \uparrow \mathcal{G} \iff \text{I} \uparrow \mathcal{H}$
- $\text{I} \uparrow_{\text{pre}} \mathcal{G} \iff \text{II} \uparrow_{\text{mark}} \mathcal{H}$
- $\text{II} \uparrow_{\text{mark}} \mathcal{G} \iff \text{I} \uparrow_{\text{pre}} \mathcal{H}$

We note one important way in which equivalence and duality interact.

Lemma 12. Suppose $\mathcal{G}_1, \mathcal{G}_2, \mathcal{H}_1,$ and \mathcal{H}_2 are selection games so that \mathcal{G}_1 is dual to \mathcal{H}_1 and \mathcal{G}_2 is dual to \mathcal{H}_2 . Then, if $\mathcal{G}_1 \leq_{\text{II}} \mathcal{G}_2$, $\mathcal{H}_2 \leq_{\text{II}} \mathcal{H}_1$. Consequently, if $\mathcal{G}_1 \equiv \mathcal{G}_2$, then $\mathcal{H}_1 \equiv \mathcal{H}_2$.

We will use consequences of [8, Cor. 26] to see that a few classes of selection games are dual.

Lemma 13. Let \mathcal{A} be an ideal of closed sets of a space X and \mathcal{B} be a collection.

- (i) By [2, Cor. 3.4] and [8, Thm. 38], $\mathbf{G}_1(\mathcal{O}_X(\mathcal{A}), \mathcal{B})$ and $\mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\mathcal{B})$ are dual. (Note that this is a general form of the duality of the Rothberger game and the point-open game.)
- (ii) By [8, Cor. 33], $\mathbf{G}_1(\mathcal{D}_X, \mathcal{B})$ and $\mathbf{G}_1(\mathcal{T}_X, \neg\mathcal{B})$ are dual.
- (iii) By [8, Cor. 35], for $x \in X$, $\mathbf{G}_1(\Omega_{X,x}, \mathcal{B})$ and $\mathbf{G}_1(\mathcal{N}_{X,x}, \neg\mathcal{B})$ are dual.

We now state the translation theorems we will be using to establish some game equivalences.

Theorem 14 ([3, Thm. 12]). Let $\mathcal{A}, \mathcal{B}, \mathcal{C},$ and \mathcal{D} be collections. Suppose there are functions $\overleftarrow{T}_{\text{I},n} : \mathcal{B} \rightarrow \mathcal{A}$ and $\overrightarrow{T}_{\text{II},n} : (\bigcup \mathcal{A}) \times \mathcal{B} \rightarrow \bigcup \mathcal{B}$ for each $n \in \omega$, so that

- (i) if $x \in \overleftarrow{T}_{\text{I},n}(B)$, then $\overrightarrow{T}_{\text{II},n}(x, B) \in B$, and
- (ii) if $\langle x_n : n \in \omega \rangle \in \prod_{n \in \omega} \overleftarrow{T}_{\text{I},n}(B_n)$ and $\{x_n : n \in \omega\} \in \mathcal{C}$, then $\{\overrightarrow{T}_{\text{II},n}(x_n, B_n) : n \in \omega\} \in \mathcal{D}$.

Then $\mathbf{G}_1(\mathcal{A}, \mathcal{C}) \leq_{\text{II}} \mathbf{G}_1(\mathcal{B}, \mathcal{D})$.

Similar results to Theorem 14 for longer length games and finite-selection games, even with simplified hypotheses, can be found in [3, 4, 5].

We will also need a separation axiom for some results in the sequel.

Definition 15. Let X be a space and \mathcal{A} be an ideal of closed subsets of X . We say that X is **\mathcal{A} -normal** if, given any $A \in \mathcal{A}$ and $U \subseteq X$ open with $A \subseteq U$, there exists an open set V so that $A \subseteq V \subseteq \text{cl}(V) \subseteq U$.

We also say that X is **functionally \mathcal{A} -normal** if, for $A \in \mathcal{A}$ and $U \subseteq X$ open with $A \subseteq U$, there exists a continuous function $f : X \rightarrow \mathbb{R}$ so that $f[A] = \{0\}$ and $f[X \setminus U] = \{1\}$.

Note that, if X is \mathcal{A} -normal, then X is regular. If $\mathcal{A} \subseteq K(X)$ and X is regular, then X is \mathcal{A} -normal. If X is Tychonoff and $\mathcal{A} \subseteq K(X)$, then X is functionally \mathcal{A} -normal.

2.2. Uniform Spaces. To introduce the basics of uniform spaces we'll need in this paper, we mostly follow [18, Chapter 6].

We recall the standard notation involved with uniformities. Let X be a set. The diagonal of X is $\Delta_X = \{\langle x, x \rangle : x \in X\}$. For $E \subseteq X^2$, $E^{-1} = \{\langle y, x \rangle : \langle x, y \rangle \in E\}$. If $E = E^{-1}$, then E is said to be *symmetric*. If $E, F \subseteq X^2$,

$$E \circ F = \{\langle x, z \rangle : (\exists y \in X) \langle x, y \rangle \in F \wedge \langle y, z \rangle \in E\}.$$

For $E \subseteq X^2$, we let $E[x] = \{y \in X : \langle x, y \rangle \in E\}$ and $E[A] = \bigcup_{x \in A} E[x]$.

Definition 16. A **uniformity** on a set X is a set $\mathcal{E} \subseteq \wp^+(X^2)$ which satisfies the following properties:

- For every $E \in \mathcal{E}$, $\Delta_X \subseteq E$.
- For every $E \in \mathcal{E}$, $E^{-1} \in \mathcal{E}$.
- For every $E \in \mathcal{E}$, there exists $F \in \mathcal{E}$ so that $F \circ F \subseteq E$.
- For $E, F \in \mathcal{E}$, $E \cap F \in \mathcal{E}$.
- For $E \in \mathcal{E}$ and $F \subseteq X^2$, if $E \subseteq F$, $F \in \mathcal{E}$.

If, in addition, $\Delta_X = \bigcap \mathcal{E}$, we say that the uniformity \mathcal{E} is **Hausdorff**. By an **entourage** of X , we mean $E \in \mathcal{E}$. The pair (X, \mathcal{E}) is called a **uniform space**.

Definition 17. For a set X , we say that $\mathcal{B} \subseteq \wp^+(X^2)$ is a **base for a uniformity** if

- for every $B \in \mathcal{B}$, $\Delta_X \subseteq B$;
- for every $B \in \mathcal{B}$, there is some $A \in \mathcal{B}$ so that $A \subseteq B^{-1}$;
- for every $B \in \mathcal{B}$, there is some $A \in \mathcal{B}$ so that $A \circ A \subseteq B$; and
- for $A, B \in \mathcal{B}$, there is some $C \in \mathcal{B}$ so that $C \subseteq A \cap B$.

If the uniformity generated by \mathcal{B} is \mathcal{E} , we say that \mathcal{B} is a **base** for \mathcal{E} .

If (X, \mathcal{E}) is a uniform space, then the uniformity \mathcal{E} generates a topology on X in the following way: $U \subseteq X$ is declared to be open provided that, for every $x \in U$, there is some $E \in \mathcal{E}$ so that $E[x] \subseteq U$. An important result about this topology is

Theorem 18 (see [18]). A Hausdorff uniform space (X, \mathcal{E}) is metrizable if and only if \mathcal{E} has a countable base.

With the topology induced by the uniformity, we endow X^2 with the resulting product topology.

Lemma 19 (See [18]). The family of open (or closed) symmetric entourages of a uniformity \mathcal{E} is a base for \mathcal{E} .

For a uniform space (X, \mathcal{E}) , there is a natural way to define a uniformity on $K(X)$ which is directly analogous to the Pompeiu-Hausdorff distance defined in the context of metric spaces.

Definition 20. Let (X, \mathcal{E}) be a uniform space and, for $E \in \mathcal{E}$, define

$$hE = \{\langle K, L \rangle \in K(X)^2 : K \subseteq E[L] \wedge L \subseteq E[K]\}.$$

Just as the Pompeiu-Hausdorff distance on compact subsets generates the Vietoris topology, the analogous uniformity also generates the Vietoris topology.

Theorem 21 (see [6, Chapter 2]). For a uniform space (X, \mathcal{E}) , $\mathcal{B} = \{hE : E \in \mathcal{E}\}$ is a base for a uniformity on $K(X)$; the topology generated by the uniform base \mathcal{B} is the Vietoris topology.

For the set of functions from a space X to a uniform space (Y, \mathcal{E}) , we review the uniformity which generates the topology of uniform convergence on a family of subsets of X . For this review, we mostly follow [18, Chapter 7].

Definition 22. For the set Y^X of functions from a set X to a uniform space (Y, \mathcal{E}) , we define, for $A \in \wp^+(X)$ and $E \in \mathcal{E}$,

$$\mathbf{U}(A, E) = \{\langle f, g \rangle \in (Y^X)^2 : (\forall x \in A) \langle f(x), g(x) \rangle \in E\}.$$

For the set of functions $X \rightarrow \mathbb{K}(Y)$, we let $\mathbf{W}(A, E) = \mathbf{U}(A, hE)$.

If \mathcal{B} is a base for a uniformity on Y and \mathcal{A} is an ideal of subsets of X , then $\{\mathbf{U}(A, B) : A \in \mathcal{A}, B \in \mathcal{B}\}$ forms a base for a uniformity on Y^X . The corresponding topology generated by this base for a uniformity is the topology of uniform convergence on \mathcal{A} . Consequently, $\{\mathbf{W}(A, B) : A \in \mathcal{A}, B \in \mathcal{B}\}$ is a base for a uniformity on $\mathbb{K}(Y)^X$.

2.3. Usco Mappings. In this section, we introduce the basic facts of usco mappings needed for this paper. Of primary use is Theorem 29 which offers a convenient characterization of minimal usco maps.

A *set-valued* function from X to Y is a function $\Phi : X \rightarrow \wp(Y)$. These are sometimes also referred to as *multi-functions*.

Definition 23. A set-valued function $\Phi : X \rightarrow \wp(Y)$ is said to be **upper semicontinuous** if, for every open $V \subseteq Y$,

$$\Phi^{\leftarrow}(V) := \{x \in X : \Phi(x) \subseteq V\}$$

is open in X . An **usco** map from a space X to Y is a set-valued map Φ from X to Y which is upper semicontinuous and whose range contained in $\mathbb{K}(Y)$. An usco map $\Phi : X \rightarrow \mathbb{K}(Y)$ is said to be **minimal** if its graph minimal with respect to the \subseteq relation. Let $MU(X, Y)$ denote the collection of all minimal usco maps $X \rightarrow \mathbb{K}(Y)$.

It is clear that any map $\Phi : X \rightarrow K(Y)$ is usco if and only if Φ is continuous relative to the *upper Vietoris topology* on $K(X)$, which is the topology generated by the

sets $\{K \in K(X) : K \subseteq U\}$ for open $U \subseteq X$. However, inspired by Theorem 21, we maintain that the full Vietoris topology is the desirable topology. At minimum, for most spaces of interest, the full Vietoris topology is Hausdorff. Moreover, the full Vietoris topology on the set of compact subsets has other desirable properties that are intimately related with X , like being metrizable when, and only when, X is metrizable.

As above, it is also clear that any continuous $\Phi : X \rightarrow \mathbb{K}(Y)$ is usco and that there are continuous $\Phi : X \rightarrow \mathbb{K}(Y)$ which are not minimal. As Example 31 will demonstrate, there are minimal usco maps which are not continuous.

Definition 24. Suppose $\Phi : X \rightarrow \wp^+(Y)$. We say that a function $f : X \rightarrow Y$ is a **selection** of Φ if $f(x) \in \Phi(x)$ for every $x \in X$. We let $\text{sel}(\Phi)$ be the set of all selections of Φ .

If $D \subseteq X$ is dense and $f : D \rightarrow Y$ is so that $f(x) \in \Phi(x)$ for each $x \in D$, we say that f is a **densely defined selection** of Φ .

Recall that a point x is said to be an *accumulation point* of a net $\langle x_\lambda : \lambda \in \Lambda \rangle$ if, for every open neighborhood U of x and every $\lambda \in \Lambda$, there exists $\mu \geq \lambda$ so that $x_\mu \in U$.

The notion of subcontinuity was introduced by Fuller [11] which can be extended to so-called densely defined functions in the following way. See also [21].

Definition 25. Suppose $D \subseteq X$ is dense. We say that a function $f : D \rightarrow Y$ is **subcontinuous** if, for every $x \in X$ and every net $\langle x_\lambda : \lambda \in \Lambda \rangle$ in D with $x_\lambda \rightarrow x$, $\langle f(x_\lambda) : \lambda \in \Lambda \rangle$ has an accumulation point.

The notion of semi-open sets was introduced by Levine [22].

Definition 26. For a space X , a set $A \subseteq X$ is said to be **semi-open** if $A \subseteq \text{cl int}(A)$.

The notion of quasicontinuity was introduced by Kempisty [19] and surveyed by Neubrunn [25].

Definition 27. A function $f : X \rightarrow Y$ is said to be **quasicontinuous** if, for each open $V \subseteq Y$, $f^{-1}(V)$ is semi-open in X .

If $D \subseteq X$ is dense and $f : D \rightarrow Y$, we will say that f is quasicontinuous if it is quasicontinuous on D with the subspace topology.

Definition 28. For $f \in \text{Fn}(X, Y)$, define $\bar{f} : X \rightarrow \wp(Y)$ by the rule

$$\bar{f}(x) = \{y \in Y : \langle x, y \rangle \in \text{cl gr}(f)\}.$$

Theorem 29 (Holá, Holý [15]). Suppose Y is regular and that $\Phi : X \rightarrow \wp^+(X)$. Then the following are equivalent:

- (i) Φ is minimal usco.
- (ii) Every selection f of Φ is subcontinuous, quasicontinuous, and $\Phi = \bar{f}$.
- (iii) There exists a selection f of Φ which is subcontinuous, quasicontinuous, and $\Phi = \bar{f}$.
- (iv) There exists a densely defined selection f of Φ which is subcontinuous, quasicontinuous, and $\Phi = \bar{f}$.

A consequence of Theorem 29 we will use is that, for $\Phi \in MU(X, \mathbb{R})$, the function $f : X \rightarrow \mathbb{R}$ defined by $f(x) = \max \Phi(x)$ is a selection of Φ , and hence subcontinuous and quasicontinuous.

We will be using the following to construct certain functions.

Lemma 30. Let $f, g : X \rightarrow Y$ and $U \in \mathcal{T}_X$ and define $h : X \rightarrow Y$ by the rule

$$h(x) = \begin{cases} f(x), & x \in \text{cl}(U); \\ g(x), & x \notin \text{cl}(U). \end{cases}$$

- (i) If f and g are subcontinuous, then h is subcontinuous.
- (ii) If f is constant, and g is quasicontinuous, then h is quasicontinuous.

Consequently, if f is constant and g is both subcontinuous and quasicontinuous, then h is both subcontinuous and quasicontinuous, which implies that \bar{h} is minimal usco.

Proof. (i) Suppose $\langle x_\lambda : \lambda \in \Lambda \rangle$ is so that $x_\lambda \rightarrow x$. If there is a cofinal subnet of $\langle x_\lambda : \lambda \in \Lambda \rangle$ which is contained in $X \setminus \text{cl}(U)$, then we can appeal to the subcontinuity of g to see that $\langle h(x_\lambda) : \lambda \in \Lambda \rangle$ has an accumulation point. Otherwise, we can let $\lambda_0 \in \Lambda$ be so that, for all $\lambda \geq \lambda_0$, $x_\lambda \in \text{cl}(U)$. Then $\langle x_\lambda : \lambda \geq \lambda_0 \rangle$ is a net contained in $\text{cl}(U)$, so we can use the subcontinuity of f to establish that $\langle h(x_\lambda) : \lambda \in \Lambda \rangle$ has an accumulation point.

(ii) Suppose $V \subseteq Y$ is open so that $h^{-1}(V) \neq \emptyset$. Let $x \in h^{-1}(V)$ and suppose W is open with $x \in W$. We proceed by cases.

If $x \in \text{cl}(U)$, then $W \cap U \neq \emptyset$. Notice also that $U \subseteq h^{-1}(V)$ in this case since f is constant. So $W \cap \text{int}(h^{-1}(V)) \neq \emptyset$.

Now, suppose $x \notin \text{cl}(U)$ and observe that $x \in W \setminus \text{cl}(U)$. Since $h(x) \in V$ and $x \notin \text{cl}(U)$, $h(x) = g(x) \in V$. By the quasicontinuity of g , it must be the case that

$$(W \setminus \text{cl}(U)) \cap \text{int}(g^{-1}(V)) \neq \emptyset.$$

Observe that $\text{int}(g^{-1}(V)) \setminus \text{cl}(U) \subseteq h^{-1}(V)$, so $W \cap \text{int}(h^{-1}(V)) \neq \emptyset$.

It follows that, in either case, $h^{-1}(V)$ is semi-open, establishing the quasicontinuity of h .

For the remainder of the proof, combine (i) and (ii) with Theorem 29. \square

Note that, if $F \subseteq X$ is closed and nowhere dense, $\mathbf{1}_F$ is not quasicontinuous since F is not semi-open. Thus, the requirement that we use the closure of an open set in Lemma 30(ii) is, in general, necessary. As a similar example, $\mathbf{1}_{(0,1) \cup (1,2)}$ is not quasicontinuous at 1.

Moreover, consider $U \in \mathcal{T}_X$ with $\partial U \neq \emptyset$ and $\text{cl}(U) \neq X$. Then $f : X \rightarrow \mathbb{R}$ defined by

$$f(x) = \begin{cases} \mathbf{1}_{X \setminus U}(x), & x \in \text{cl}(U); \\ \mathbf{1}_U(x), & x \notin \text{cl}(U). \end{cases}$$

is $\mathbf{1}_{\partial U}$. So the requirement that f be constant (or something stronger than quasicontinuity, at least) in Lemma 30(ii) is, in general, necessary.

Example 31. Consider $MU(\mathbb{R}, \mathbb{R})$. By Lemma 30, $\Phi := \overline{\mathbf{1}}_{[0,1]}$ is minimal usco. However, Φ is not continuous since

$$\{0, 1\} = \{x \in \mathbb{R} : \Phi(x) \in [(-0.5, 0.5), (0.5, 1.5)]\}.$$

Hence, when Y is metrizable, studying the space $MU(X, Y)$ is, in general, different than studying the space of continuous functions into a metrizable space.

We will also be using the following corollary often.

Corollary 32. Suppose $\Phi, \Psi \in MU(X, Y)$ and $U \in \mathcal{T}_X$ are so that there exist $f \in \text{sel}(\Phi)$ and $g \in \text{sel}(\Psi)$ with the property that $f \upharpoonright_U = g \upharpoonright_U$. Then $\Phi \upharpoonright_U = \Psi \upharpoonright_U$.

Proof. We will show that $\Phi(x) \subseteq \Psi(x)$ for each $x \in U$. By symmetry, this will establish that $\Phi(x) = \Psi(x)$ for each $x \in U$.

So let $x \in U$ and $y \in \Phi(x)$ which, by Theorem 29 means that $y \in \overline{f}(x)$. Now, consider any neighborhood $V \times W$ of $\langle x, y \rangle$. Without loss of generality, we may assume that $V \subseteq U$. Then we can find $\langle z, f(z) \rangle \in V \times W$. Since $z \in U$, $f(z) = g(z)$ and, as $V \times W$ was arbitrary, we see that $\langle x, y \rangle \in \text{cl gr}(g)$. Hence, $y \in \overline{g}(x) = \Psi(x)$. \square

Corollary 33. If $A \subseteq X$ is non-empty, $U, V \in \mathcal{T}_X$ are so that $A \subseteq V \subseteq \text{cl}(V) \subseteq U$, $\Phi \in MU(X, Y)$, and $f \in \text{sel}(\Phi)$, then, for $y_0 \in Y$, $g : X \rightarrow Y$ defined by

$$g(x) = \begin{cases} y_0, & x \in \text{cl}(X \setminus \text{cl}(V)); \\ f(x), & \text{otherwise,} \end{cases}$$

has the property that $\Psi := \overline{g} \in MU(X, Y)$, $\langle \Phi, \Psi \rangle \in \mathbf{W}(A, E)$ for any entourage E of Y , and $g[X \setminus U] = \{y_0\}$.

Proof. Note that Theorem 29 and Lemma 30 imply that g is subcontinuous and quasicontinuous. Then $\Psi := \overline{g} \in MU(X, Y)$.

Since V is open, $\text{cl}(X \setminus \text{cl}(V)) \subseteq X \setminus V$. Then $g(x) = f(x)$ for all $x \in V$. By Corollary 32, we see that $\Phi \upharpoonright_A = \Psi \upharpoonright_A$ as $A \subseteq V$. Also, since $X \setminus U \subseteq X \setminus \text{cl}(V) \subseteq \text{cl}(X \setminus \text{cl}(V))$, we see that $g[X \setminus U] = \{y_0\}$. \square

As a final note in this section, we offer the following generalization of [16, Lemma 3.1] to general uniform spaces. Recall that Lemma 19 allows us to restrict our attention to closed entourages.

Corollary 34. Let X be a space and Y be a uniform space. If $\Phi, \Psi \in MU(X, Y)$, a closed entourage E of Y , and a dense $D \subseteq X$ are so that $\langle \Phi(x), \Psi(x) \rangle \in hE$ for all $x \in D$, then $\langle \Phi(x), \Psi(x) \rangle \in hE$ for all $x \in X$.

Proof. Define $F : X \rightarrow \wp(Y)$ by $F(x) = E[\Phi(x)]$.

We first show that the graph of F is closed. Suppose $\langle x, y \rangle \in \text{cl gr}(F)$ and let $\langle \langle x_\lambda, y_\lambda \rangle : \lambda \in \Lambda \rangle$ be a net in $\text{gr}(F)$ so that $\langle x_\lambda, y_\lambda \rangle \rightarrow \langle x, y \rangle$. Since $y_\lambda \in E[\Phi(x_\lambda)]$, we can let $w_\lambda \in \Phi(x_\lambda)$ be so that $y_\lambda \in E[w_\lambda]$. Observe that, since $x_\lambda \rightarrow x$ and $w_\lambda \in \Phi(x_\lambda)$ for each $\lambda \in \Lambda$, by Theorem 29, $\langle w_\lambda : \lambda \in \Lambda \rangle$ has an accumulation point $w \in \Phi(x)$. Since $y_\lambda \rightarrow y$ and w is an accumulation point of $\langle w_\lambda : \lambda \in \Lambda \rangle$, $\langle w, y \rangle$ is an accumulation point of $\langle \langle w_\lambda, y_\lambda \rangle : \lambda \in \Lambda \rangle$. Moreover,

as $\langle w_\lambda, y_\lambda \rangle \in E$ for all $\lambda \in \Lambda$ and E is closed, we see that $\langle w, y \rangle \in E$. Hence, $y \in E[w] \subseteq E[\Phi(x)] = F(x)$. That is, $\langle x, y \rangle \in \text{gr}(F)$ which establishes that $\text{gr}(F)$ is closed.

Now, by Theorem 29, we can let $g : D \rightarrow Y$ be subcontinuous and quasicontinuous so that $g(x) \in \Psi(x)$ for each $x \in D$ and $\Psi = \overline{g}$. Since $\text{gr}(F)$ is closed and $\text{gr}(g) \subseteq \text{gr}(F)$, we see that $\text{cl } \text{gr}(g) \subseteq \text{gr}(F)$. That is, $\Psi(x) \subseteq F(x) = E[\Phi(x)]$ for all $x \in X$.

A symmetric argument shows that $\Phi(x) \subseteq E[\Psi(x)]$ for all $x \in X$, finishing the proof. \square

3. RESULTS

We first note that the clustering version of Gruenhage's W -game is equivalent to an entourage selection game in the realm of topological groups. Such a result holds, for example, for $C_{\mathcal{A}}(X)$ where we define $C_{\mathcal{A}}(X)$ to be the space of continuous real-valued functions on X endowed with the topology of uniform convergence on \mathcal{A} , an ideal of closed subsets of X . This topology is generated by the uniformity outlined in Definition 22.

Recall that a topological group is a (multiplicative) group G with a topology for which the operations $\langle g, h \rangle \mapsto gh$, $G^2 \rightarrow G$, and $g \mapsto g^{-1}$, $G \rightarrow G$, are continuous. Let \mathbf{i} be the identity element of G . Also recall that

$$\{ \{ \langle g, h \rangle \in G^2 : gh^{-1} \in U \} : U \in \mathcal{N}_{G, \mathbf{i}} \}$$

is a basis for a uniformity on G . Let \mathcal{E}_G be the set of all entourages of G with the generated uniformity. Also let

$$\Omega_\Delta = \{ A \subseteq G^2 : \{ gh^{-1} : \langle g, h \rangle \in A \} \in \Omega_{G, \mathbf{i}} \}.$$

Theorem 35. If G is a (multiplicative) topological group and \mathbf{i} is the identity, then, for any $g \in G$,

$$\mathbf{G}_1(\mathcal{N}_{G, g}, \neg\Omega_{G, g}) \equiv \mathbf{G}_1(\mathcal{N}_{G, \mathbf{i}}, \neg\Omega_{G, \mathbf{i}}) \equiv \mathbf{G}_1(\mathcal{E}_G, \neg\Omega_\Delta).$$

Proof. The equivalence

$$\mathbf{G}_1(\mathcal{N}_{G, g}, \neg\Omega_{G, g}) \equiv \mathbf{G}_1(\mathcal{N}_{G, \mathbf{i}}, \neg\Omega_{G, \mathbf{i}})$$

can be seen by using the homeomorphism $x \mapsto gx$, $G \rightarrow G$.

We first show that

$$\mathbf{G}_1(\mathcal{N}_{G, \mathbf{i}}, \neg\Omega_{G, \mathbf{i}}) \leq_{\text{II}} \mathbf{G}_1(\mathcal{E}_G, \neg\Omega_\Delta).$$

Define $\overleftarrow{T}_{\text{I}, n} : \mathcal{E}_G \rightarrow \mathcal{N}_{G, \mathbf{i}}$ so that

$$\{ \langle g, h \rangle \in G^2 : gh^{-1} \in \overleftarrow{T}_{\text{I}, n}(E) \} \subseteq E.$$

We now define $\overrightarrow{T}_{\text{II}, n} : G \times \mathcal{E}_G \rightarrow G^2$ in the following way. For $g \in \overleftarrow{T}_{\text{I}, n}(E)$, let $\overrightarrow{T}_{\text{II}, n}(g, E) = \langle g, \mathbf{i} \rangle$; otherwise, let $\overrightarrow{T}_{\text{II}, n}(g, E) = \langle \mathbf{i}, \mathbf{i} \rangle$. By definition, we see that $\overrightarrow{T}_{\text{II}, n}(g, E) \in E$ when $g \in \overleftarrow{T}_{\text{I}, n}(E)$.

Suppose $\langle g_n : n \in \omega \rangle \in \prod_{n \in \omega} \overleftarrow{T}_{I,n}(E_n)$ for a sequence $\langle E_n : n \in \omega \rangle$ of \mathcal{E}_G so that $\{g_n : n \in \omega\} \notin \Omega_{G,i}$. Then let $U \in \mathcal{N}_{G,i}$ be so that $U \cap \{g_n : n \in \omega\} = \emptyset$. Notice that $\overrightarrow{T}_{II,n}(g_n, E_n) \notin \Omega_\Delta$. Thus, Theorem 14 applies.

We now show that

$$\mathbf{G}_1(\mathcal{E}_G, \neg\Omega_\Delta) \leq_{II} \mathbf{G}_1(\mathcal{N}_{G,i}, \neg\Omega_{G,i}).$$

Define $\overleftarrow{T}_{II,n} : \mathcal{N}_{G,i} \rightarrow \mathcal{E}_G$ by the rule

$$\overleftarrow{T}_{II,n}(U) = \{\langle g, h \rangle \in G^2 : gh^{-1} \in U\}$$

and $\overrightarrow{T}_{II,n} : G^2 \times \mathcal{N}_{G,i} \rightarrow G$ by $\overrightarrow{T}_{II,n}(\langle g, h \rangle, U) = gh^{-1}$. Note that, if $\langle g, h \rangle \in \overleftarrow{T}_{II,n}(U)$, $\overrightarrow{T}_{II,n}(\langle g, h \rangle, U) \in U$. So suppose we have

$$\langle \langle g_n, h_n \rangle : n \in \omega \rangle \in \prod_{n \in \omega} \overleftarrow{T}_{II,n}(U_n)$$

for a sequence $\langle U_n : n \in \omega \rangle$ of $\mathcal{N}_{G,i}$ so that $\{\langle g_n, h_n \rangle : n \in \omega\} \notin \Omega_\Delta$. Evidently, $\overrightarrow{T}_{II,n}(\langle g_n, h_n \rangle, U_n) \notin \Omega_{G,i}$. Again, Theorem 14 applies. \square

For the remainder of the paper, we will be interested only in real set-valued functions; so we will let $MU(X) = MU(X, \mathbb{R})$. We also use, for $\varepsilon > 0$,

$$\Delta_\varepsilon = \{\langle x, y \rangle \in \mathbb{R}^2 : |x - y| < \varepsilon\}.$$

For $A \subseteq X$, we will use $\mathbf{U}(A, \varepsilon) = \mathbf{U}(A, \Delta_\varepsilon)$ and $\mathbf{W}(A, \varepsilon) = \mathbf{W}(A, \Delta_\varepsilon)$. For $Y \subseteq \mathbb{R}$, let $\mathbb{B}(Y, \varepsilon) = \bigcup_{y \in Y} B(y, \varepsilon)$ and note that

$$\begin{aligned} & \mathbf{W}(A, \varepsilon) \\ &= \{ \langle \Phi, \Psi \rangle \in \mathbb{K}(\mathbb{R})^X : (\forall x \in A) [\Phi(x) \subseteq \mathbb{B}(\Psi(x), \varepsilon) \wedge \Psi(x) \subseteq \mathbb{B}(\Phi(x), \varepsilon)] \}. \end{aligned}$$

Then, if \mathcal{A} is an ideal of closed subsets of X , we will use $MU_{\mathcal{A}}(X)$ to denote the set $MU(X)$ with the topology generated by the base for a uniformity $\{\mathbf{W}(A, \varepsilon) : A \in \mathcal{A}, \varepsilon > 0\}$. When $\mathcal{A} = [X]^{<\omega}$, we use $MU_p(X)$ and, when $\mathcal{A} = K(X)$, we use $MU_k(X)$. For $\Phi \in MU(X)$, $A \subseteq X$, and $\varepsilon > 0$, we let $[\Phi; A, \varepsilon] = \mathbf{W}(A, \varepsilon)[\Phi]$. We will use $\mathbf{0}$ to denote the function that is constantly 0 when dealing with real-valued functions and the function that is constantly $\{0\}$ when dealing with usco maps.

Theorem 36. Let X be regular and \mathcal{A} and \mathcal{B} be ideals of closed subsets of X . Then,

- (i) $\mathbf{G}_1(\mathcal{O}_X(\mathcal{A}), \Lambda_X(\mathcal{B})) \leq_{II} \mathbf{G}_1(\Omega_{MU_{\mathcal{A}}(X), \mathbf{0}}, \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}})$,
- (ii) $\mathbf{G}_1(\Omega_{MU_{\mathcal{A}}(X), \mathbf{0}}, \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}) \leq_{II} \mathbf{G}_1(\mathcal{D}_{MU_{\mathcal{A}}(X)}, \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}})$, and
- (iii) if X is \mathcal{A} -normal, $\mathbf{G}_1(\mathcal{D}_{MU_{\mathcal{A}}(X)}, \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}) \leq_{II} \mathbf{G}_1(\mathcal{O}_X(\mathcal{A}), \Lambda_X(\mathcal{B}))$.

Thus, if X is \mathcal{A} -normal, the three games are equivalent.

Proof. We first address (i). Fix some $\mathcal{U}_0 \in \mathcal{O}_X(\mathcal{A})$ and let $W_{\Phi, n} = \Phi^{\leftarrow} [(-2^{-n}, 2^{-n})]$ for $\Phi \in MU(X)$ and $n \in \omega$. Define $\overleftarrow{T}_{I,n} : \Omega_{MU_{\mathcal{A}}(X), \mathbf{0}} \rightarrow \mathcal{O}_X(\mathcal{A})$ by the rule

$$\overleftarrow{T}_{I,n}(\mathcal{F}) = \begin{cases} \{W_{\Phi, n} : \Phi \in \mathcal{F}\}, & (\forall \Phi \in \mathcal{F}) W_{\Phi, n} \neq X; \\ \mathcal{U}_0, & \text{otherwise.} \end{cases}$$

To see that $\overleftarrow{T}_{I,n}$ is defined, let $\mathcal{F} \in \Omega_{MU_{\mathcal{A}}(X), \mathbf{0}}$ be so that $W_{\Phi,n} \neq X$ for every $\Phi \in \mathcal{F}$. Let $A \in \mathcal{A}$ be arbitrary and choose $\Phi \in [\mathbf{0}; A, 2^{-n}] \cap \mathcal{F}$. It follows that $A \subseteq W_{\Phi,n}$. Hence, $\overleftarrow{T}_{I,n}(\mathcal{F}) \in \mathcal{O}_X(\mathcal{A})$.

We now define

$$\overrightarrow{T}_{II,n} : \mathcal{T}_X \times \Omega_{MU_{\mathcal{A}}(X), \mathbf{0}} \rightarrow MU(X)$$

in the following way. Let

$$\mathfrak{T}_n = \{\mathcal{F} \in \Omega_{MU_{\mathcal{A}}(X), \mathbf{0}} : (\exists \Phi \in \mathcal{F}) W_{\Phi,n} = X\}$$

and $\mathfrak{T}_n^* = \Omega_{MU_{\mathcal{A}}(X), \mathbf{0}} \setminus \mathfrak{T}_n$. For each $\langle U, \mathcal{F} \rangle \in \mathcal{T}_X \times \mathfrak{T}_n$, let $\overrightarrow{T}_{II,n}(U, \mathcal{F}) \in \mathcal{F}$ be so that $W_{\overrightarrow{T}_{II,n}(U, \mathcal{F}), n} = X$. For $\langle U, \mathcal{F} \rangle \in \mathcal{T}_X \times \mathfrak{T}_n^*$ so that $U \in \overleftarrow{T}_{I,n}(\mathcal{F})$, let $\overrightarrow{T}_{II,n}(U, \mathcal{F}) \in \mathcal{F}$ be so that $U = W_{\overrightarrow{T}_{II,n}(U, \mathcal{F}), n}$. For $\langle U, \mathcal{F} \rangle \in \mathcal{T}_X \times \mathfrak{T}_n^*$ so that $U \notin \overleftarrow{T}_{I,n}(\mathcal{F})$, let $\overrightarrow{T}_{II,n}(U, \mathcal{F}) = \mathbf{0}$. By construction, if $U \in \overleftarrow{T}_{I,n}(\mathcal{F})$, then $\overrightarrow{T}_{II,n}(U, \mathcal{F}) \in \mathcal{F}$.

To finish this application of Theorem 14, assume that we have

$$\langle U_n : n \in \omega \rangle \in \prod_{n \in \omega} \overleftarrow{T}_{I,n}(\mathcal{F}_n)$$

for some sequence $\langle \mathcal{F}_n : n \in \omega \rangle$ of $\Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}$ so that $\{U_n : n \in \omega\} \in \Lambda_X(\mathcal{B})$. For each $n \in \omega$, let $\Phi_n = \overrightarrow{T}_{II,n}(U_n, \mathcal{F}_n)$. Now, let $B \in \mathcal{B}$ and $\varepsilon > 0$ be arbitrary. Choose $n \in \omega$ so that $2^{-n} < \varepsilon$ and $B \subseteq U_n$. If $\mathcal{F}_n \in \mathfrak{T}_n$, then Φ_n has the property that $X = \Phi_n^{\leftarrow} [(-2^{-n}, 2^{-n})]$; hence, $\Phi_n \in [\mathbf{0}; B, \varepsilon]$. Otherwise, $B \subseteq U_n = \Phi_n^{\leftarrow} [(-2^{-n}, 2^{-n})]$ which also implies that $\Phi_n \in [\mathbf{0}; B, \varepsilon]$. Thus, $\{\Phi_n : n \in \omega\} \in \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}$.

(ii) holds since $\mathcal{D}_{MU_{\mathcal{A}}(X)} \subseteq \Omega_{MU_{\mathcal{A}}(X), \mathbf{0}}$.

Lastly, we address (iii). We define $\overleftarrow{T}_{I,n} : \mathcal{O}_X(\mathcal{A}) \rightarrow \mathcal{D}_{MU_{\mathcal{A}}(X)}$ by the rule

$$\overleftarrow{T}_{I,n}(\mathcal{U}) = \{\Phi \in MU(X) : (\exists U \in \mathcal{U})(\exists f \in \text{sel}(\Phi)) f[X \setminus U] = \{1\}\}.$$

To see that $\overleftarrow{T}_{I,n}$ is defined, let $\mathcal{U} \in \mathcal{O}_X(\mathcal{A})$ and consider a basic open set $[\Phi; A, \varepsilon]$. Then let $U \in \mathcal{U}$ be so that $A \subseteq U$ and, by \mathcal{A} -normality, let V be open so that $A \subseteq V \subseteq \text{cl}(V) \subseteq U$. Define $f : X \rightarrow \mathbb{R}$ by the rule

$$f(x) = \begin{cases} 1, & x \in \text{cl}(X \setminus \text{cl}(V)); \\ \max \Phi(x), & \text{otherwise.} \end{cases}$$

By Corollary 33, $\overline{f} \in [\Phi; A, \varepsilon] \cap \overleftarrow{T}_{I,n}(\mathcal{U})$. Hence, $\overleftarrow{T}_{I,n}(\mathcal{U}) \in \mathcal{D}_{MU_{\mathcal{A}}(X)}$.

We define $\overrightarrow{T}_{II,n} : MU(X) \times \mathcal{O}_X(\mathcal{A}) \rightarrow \mathcal{T}_X$ in the following way. Fix some $U_0 \in \mathcal{T}_X$. For $\langle \Phi, \mathcal{U} \rangle \in MU(X) \times \mathcal{O}_X(\mathcal{A})$, if

$$\{U \in \mathcal{U} : (\exists f \in \text{sel}(\Phi)) f[X \setminus U] = \{1\}\} \neq \emptyset,$$

let $\vec{T}_{\text{II},n}(\Phi, \mathcal{U}) \in \mathcal{U}$ be so that there exists $f \in \text{sel}(\Phi)$ with the property that $f[X \setminus \vec{T}_{\text{II},n}(\Phi, \mathcal{U})] = \{1\}$; otherwise, let $\vec{T}_{\text{II},n}(\Phi, \mathcal{U}) = U_0$. By construction, if $\Phi \in \overleftarrow{T}_{\text{I},n}(\mathcal{U})$, then $\vec{T}_{\text{II},n}(\Phi, \mathcal{U}) \in \mathcal{U}$.

Suppose we have

$$\langle \Phi_n : n \in \omega \rangle \in \prod_{n \in \omega} \overleftarrow{T}_{\text{I},n}(\mathcal{U}_n)$$

for a sequence $\langle \mathcal{U}_n : n \in \omega \rangle$ of $\mathcal{O}_X(\mathcal{A})$ with the property that $\{\Phi_n : n \in \omega\} \in \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}$. For each $n \in \omega$, let $U_n = \vec{T}_{\text{II},n}(\Phi_n, \mathcal{U}_n)$. Since \mathcal{B} is an ideal of sets, we need only show that $\langle U_n : n \in \omega \rangle$ is a \mathcal{B} -cover. So let $B \in \mathcal{B}$ be arbitrary and let $n \in \omega$ be so that $\Phi_n \in [\mathbf{0}; B, 1]$. Then we can let $f \in \text{sel}(\Phi_n)$ be so that $f[X \setminus U_n] = \{1\}$. Since $\Phi_n \in [\mathbf{0}; B, 1]$, we see that, for each $x \in B$, $f(x) \in \Phi_n(x) \subseteq (-1, 1)$. Hence, $B \cap (X \setminus U_n) = \emptyset$, which is to say that $B \subseteq U_n$. So Theorem 14 applies. \square

We now establish some relationships between these games and games on the space of continuous real-valued functions.

Corollary 37. Let \mathcal{A} and \mathcal{B} be ideals of closed subsets of X and suppose that X is \mathcal{A} -normal. Then

$$\begin{aligned} \mathcal{G} := \mathbf{G}_1(\mathcal{O}_X(\mathcal{A}), \mathcal{O}_X(\mathcal{B})) &\equiv \mathbf{G}_1(\Omega_{MU_{\mathcal{A}}(X), \mathbf{0}}, \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}) \\ &\equiv \mathbf{G}_1(\mathcal{D}_{MU_{\mathcal{A}}(X)}, \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}), \end{aligned}$$

$$\begin{aligned} \mathcal{H} := \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg \mathcal{O}_X(\mathcal{B})) &\equiv \mathbf{G}_1(\mathcal{N}_{MU_{\mathcal{A}}(X), \mathbf{0}}, \neg \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}) \\ &\equiv \mathbf{G}_1(\mathcal{T}_{MU_{\mathcal{A}}(X)}, \neg \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}), \end{aligned}$$

and \mathcal{G} is dual to \mathcal{H} . If X is functionally \mathcal{A} -normal, then

$$\begin{aligned} \mathbf{G}_1(\mathcal{O}_X(\mathcal{A}), \mathcal{O}_X(\mathcal{B})) &\equiv \mathbf{G}_1(\Omega_{MU_{\mathcal{A}}(X), \mathbf{0}}, \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}) \\ &\equiv \mathbf{G}_1(\Omega_{C_{\mathcal{A}}(X), \mathbf{0}}, \Omega_{C_{\mathcal{B}}(X), \mathbf{0}}) \\ &\equiv \mathbf{G}_1(\mathcal{D}_{MU_{\mathcal{A}}(X)}, \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}) \\ &\equiv \mathbf{G}_1(\mathcal{D}_{C_{\mathcal{A}}(X)}, \Omega_{C_{\mathcal{B}}(X), \mathbf{0}}) \end{aligned}$$

and

$$\begin{aligned} \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg \mathcal{O}_X(\mathcal{B})) &\equiv \mathbf{G}_1(\mathcal{N}_{MU_{\mathcal{A}}(X), \mathbf{0}}, \neg \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}) \\ &\equiv \mathbf{G}_1(\mathcal{N}_{C_{\mathcal{A}}(X), \mathbf{0}}, \neg \Omega_{C_{\mathcal{B}}(X), \mathbf{0}}) \\ &\equiv \mathbf{G}_1(\mathcal{T}_{MU_{\mathcal{A}}(X)}, \neg \Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}) \\ &\equiv \mathbf{G}_1(\mathcal{T}_{C_{\mathcal{A}}(X)}, \neg \Omega_{C_{\mathcal{B}}(X), \mathbf{0}}). \end{aligned}$$

Proof. Apply Theorem 36, Lemmas 8, 12, and 13, and [3, Cor. 14]. \square

In [4, Thm. 31], inspired by Li [23], the game $\mathbf{G}_1(\mathcal{O}_X(\mathcal{A}), \mathcal{O}_X(\mathcal{B}))$ is shown to be equivalent to the selective separability game on certain hyperspaces of X , which we only note in passing here for the interested reader.

Recall that a subset A of a topological space is *sequentially compact* if every sequence in A has a subsequence which converges to a point of A .

Lemma 38. Suppose $\Phi \in MU(X)$ and that $A \subseteq X$ is sequentially compact. Then $\Phi[A]$ is bounded.

Proof. Suppose $\Phi : X \rightarrow \mathbb{K}(\mathbb{R})$ is unbounded on $A \subseteq X$ which is sequentially compact. For each $n \in \omega$, let $x_n \in A$ be so that there is some $y \in \Phi(x_n)$ with $|y| \geq n$. Then let $y_n \in \Phi(x_n)$ be so that $|y_n| \geq n$ and define $f : X \rightarrow \mathbb{R}$ to be a selection of Φ so that $f(x_n) = y_n$ for $n \in \omega$. Since A is sequentially compact, we can find $x \in A$ and a subsequence $\langle x_{n_k} : k \in \omega \rangle$ so that $x_{n_k} \rightarrow x$. Notice that $\langle f(x_{n_k}) : k \in \omega \rangle$ does not have an accumulation point. Therefore f is not subcontinuous and, by Theorem 29, Φ is not a minimal usco map. \square

Theorem 39. Let \mathcal{A} and \mathcal{B} be ideals of closed subsets of X . If X is \mathcal{A} -normal and \mathcal{B} consists of sequentially compact sets, then

$$\mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\Lambda_X(\mathcal{B})) \leq_{\text{II}} \mathbf{G}_1(\mathcal{T}_{MU_{\mathcal{A}}(X)}, \text{CD}_{MU_{\mathcal{B}}(X)}).$$

Consequently,

$$\begin{aligned} \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\mathcal{O}_X(\mathcal{B})) &\equiv \mathbf{G}_1(\mathcal{N}_{MU_{\mathcal{A}}(X)}, \neg\Omega_{MU_{\mathcal{B}}(X)}, \mathbf{0}) \\ &\equiv \mathbf{G}_1(\mathcal{T}_{MU_{\mathcal{A}}(X)}, \neg\Omega_{MU_{\mathcal{B}}(X)}, \mathbf{0}) \\ &\equiv \mathbf{G}_1(\mathcal{T}_{MU_{\mathcal{A}}(X)}, \text{CD}_{MU_{\mathcal{B}}(X)}). \end{aligned}$$

Proof. Let $\pi_1 : MU(X) \times \mathcal{A} \times \mathbb{R} \rightarrow MU(X)$, $\pi_2 : MU(X) \times \mathcal{A} \times \mathbb{R} \rightarrow \mathcal{A}$, and $\pi_3 : MU(X) \times \mathcal{A} \times \mathbb{R} \rightarrow \mathbb{R}$ be the standard coordinate projection maps. Define a choice function $\gamma : \mathcal{T}_{MU_{\mathcal{A}}(X)} \rightarrow MU(X) \times \mathcal{A} \times \mathbb{R}$ so that

$$[\pi_1(\gamma(W)); \pi_2(\gamma(W)), \pi_3(\gamma(W))] \subseteq W.$$

Let $\Psi_W = \pi_1(\gamma(W))$ and $A_W = \pi_2(\gamma(W))$. Now we define $\overleftarrow{T}_{I,n} : \mathcal{T}_{MU_{\mathcal{A}}(X)} \rightarrow \mathcal{N}_X[\mathcal{A}]$ by $\overleftarrow{T}_{I,n}(W) = \mathcal{N}_X(A_W)$.

We now define $\overrightarrow{T}_{II,n} : \mathcal{T}_X \times \mathcal{T}_{MU_{\mathcal{A}}(X)} \rightarrow MU(X)$ in the following way. For $A \in \mathcal{A}$ and $U \in \mathcal{N}_X(A)$, let $V_{A,U}$ be open so that

$$A \subseteq V_{A,U} \subseteq \text{cl}(V_{A,U}) \subseteq U.$$

For $W \in \mathcal{T}_{MU_{\mathcal{A}}(X)}$ and $U \in \overleftarrow{T}_{I,n}(W)$, define $f_{W,U,n} : X \rightarrow \mathbb{R}$ by the rule

$$f_{W,U,n}(x) = \begin{cases} n, & x \in \text{cl}(X \setminus \text{cl}(V_{A_W,U})); \\ \max \Psi_W(x), & \text{otherwise.} \end{cases}$$

Then we set

$$\overrightarrow{T}_{II,n}(U, W) = \begin{cases} \overline{f}_{W,U,n}, & U \in \overleftarrow{T}_{I,n}(W); \\ \mathbf{0}, & \text{otherwise.} \end{cases}$$

By Corollary 33, $\overrightarrow{T}_{II,n}(U, W) \in MU(X)$ and, if $U \in \overleftarrow{T}_{I,n}(W)$,

$$\overrightarrow{T}_{II,n}(U, W) \in [\Psi_W; A_W, \pi_3(\gamma(W))] \subseteq W.$$

Suppose we have a sequence

$$\langle U_n : n \in \omega \rangle \in \prod_{n \in \omega} \overleftarrow{T}_{I,n}(W_n)$$

for a sequence $\langle W_n : n \in \omega \rangle$ of $\mathcal{T}_{MU_{\mathcal{A}}(X)}$ so that $\{U_n : n \in \omega\} \notin \Lambda_X(\mathcal{B})$. Let $\Phi_n = \vec{T}_{\text{II},n}(U_n, W_n)$ for each $n \in \omega$. We can find $N \in \omega$ and $B \in \mathcal{B}$ so that, for every $n \geq N$, $B \not\subseteq U_n$. Now, suppose $\Phi \in MU(X) \setminus \{\Phi_n : n \in \omega\}$ is arbitrary. By Lemma 38, $\Phi[B]$ is bounded, so let $M > \sup |\Phi[B]|$ and $n \geq \max\{N, M + 1\}$. Now, for $x \in B \setminus U_n$, note that $n \in \Phi_n(x)$ and that, for $y \in \Phi(x)$,

$$y \leq \sup |\Phi[B]| < M \leq n - 1 \implies y - n < -1 \implies |y - n| > 1.$$

In particular, $\Phi_n(x) \not\subseteq \mathbb{B}(\Phi(x), 1)$ which establishes that $\Phi_n \notin [\Phi; B, 1]$. Hence, $\{\Phi_n : n \in \omega\}$ is closed and discrete and Theorem 14 applies.

For what remains, observe that

$$\mathbf{G}_1(\mathcal{T}_{MU_{\mathcal{A}}(X)}, \text{CD}_{MU_{\mathcal{B}}(X)}) \leq_{\text{II}} \mathbf{G}_1(\mathcal{T}_{MU_{\mathcal{A}}(X)}, \neg\Omega_{MU_{\mathcal{B}}(X), \mathbf{0}})$$

since, if Two can produce a closed discrete set, then Two can avoid clustering around $\mathbf{0}$. Hence, by Corollary 37 we obtain that

$$\begin{aligned} \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\mathcal{O}_X(\mathcal{B})) &= \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\Lambda_X(\mathcal{B})) \\ &\leq_{\text{II}} \mathbf{G}_1(\mathcal{T}_{MU_{\mathcal{A}}(X)}, \text{CD}_{MU_{\mathcal{B}}(X)}) \\ &\leq_{\text{II}} \mathbf{G}_1(\mathcal{T}_{MU_{\mathcal{A}}(X)}, \neg\Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}) \\ &\equiv \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\mathcal{O}_X(\mathcal{B})). \end{aligned}$$

This complete the proof. \square

Corollary 40. Let \mathcal{A} and \mathcal{B} be ideals of closed subsets of X . If X is functionally \mathcal{A} -normal and \mathcal{B} consists of sequentially compact sets, then

$$\begin{aligned} \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\mathcal{O}_X(\mathcal{B})) &\equiv \mathbf{G}_1(\mathcal{N}_{MU_{\mathcal{A}}(X), \mathbf{0}}, \neg\Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}) \\ &\equiv \mathbf{G}_1(\mathcal{N}_{C_{\mathcal{A}}(X), \mathbf{0}}, \neg\Omega_{C_{\mathcal{B}}(X), \mathbf{0}}) \\ &\equiv \mathbf{G}_1(\mathcal{T}_{MU_{\mathcal{A}}(X)}, \neg\Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}) \\ &\equiv \mathbf{G}_1(\mathcal{T}_{C_{\mathcal{A}}(X)}, \neg\Omega_{C_{\mathcal{B}}(X), \mathbf{0}}) \\ &\equiv \mathbf{G}_1(\mathcal{T}_{MU_{\mathcal{A}}(X)}, \text{CD}_{MU_{\mathcal{B}}(X)}) \\ &\equiv \mathbf{G}_1(\mathcal{T}_{C_{\mathcal{A}}(X)}, \text{CD}_{C_{\mathcal{B}}(X)}). \end{aligned}$$

Proof. Apply Theorem 39 and [3, Cor. 15]. \square

We now offer some relationships related to Gruenhage's W -games.

Proposition 41. Let \mathcal{A} and \mathcal{B} be ideals of closed subsets of X . Then

- (i) $\mathbf{G}_1(\mathcal{N}_{MU_{\mathcal{A}}(X), \mathbf{0}}, \neg\Omega_{MU_{\mathcal{B}}(X), \mathbf{0}}) \leq_{\text{II}} \mathbf{G}_1(\mathcal{N}_{MU_{\mathcal{A}}(X), \mathbf{0}}, \neg\Gamma_{MU_{\mathcal{B}}(X), \mathbf{0}})$ and
- (ii) $\mathbf{G}_1(\mathcal{N}_{MU_{\mathcal{A}}(X), \mathbf{0}}, \neg\Gamma_{MU_{\mathcal{B}}(X), \mathbf{0}}) \leq_{\text{II}} \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\Gamma_X(\mathcal{B}))$.

Proof. (i) is evident since, if Two can avoid clustering at $\mathbf{0}$, they can surely avoid converging to $\mathbf{0}$.

(ii) Fix $U_0 \in \mathcal{T}_X$ and define $\overleftarrow{T}_{\text{I},n} : \mathcal{N}_X[\mathcal{A}] \rightarrow \mathcal{N}_{MU_{\mathcal{A}}(X), \mathbf{0}}$ by $\overleftarrow{T}_{\text{I},n}(\mathcal{N}_X(A)) = [\mathbf{0}; A, 2^{-n}]$. Then define $\vec{T}_{\text{II},n} : MU(X) \times \mathcal{N}_X[\mathcal{A}] \rightarrow \mathcal{T}_X$ by $\vec{T}_{\text{II},n}(\Phi, \mathcal{N}_X(A)) = \Phi^{\leftarrow} [(-2^{-n}, 2^{-n})]$. Note that, if $\Phi \in [\mathbf{0}; A, 2^{-n}] = \overleftarrow{T}_{\text{I},n}(\mathcal{N}_X(A))$, then $A \subseteq \Phi^{\leftarrow} [(-2^{-n}, 2^{-n})]$, which establishes that $\vec{T}_{\text{II},n}(\Phi, \mathcal{N}_X(A)) \in \mathcal{N}_X(A)$.

Suppose we have

$$\langle \Phi_n : n \in \omega \rangle \in \prod_{n \in \omega} \overleftarrow{T}_{I,n}(\mathcal{N}_X(A_n))$$

for a sequence $\langle A_n : n \in \omega \rangle$ of \mathcal{A} so that $\langle \Phi_n : n \in \omega \rangle \notin \Gamma_{MU_{\mathcal{B}}(X), \mathbf{0}}$. Then we can find $B \in \mathcal{B}$, $\varepsilon > 0$, and $N \in \omega$ so that $2^{-N} < \varepsilon$ and, for all $n \geq N$, $\Phi_n \notin [\mathbf{0}; B, \varepsilon]$.

To finish this application of Theorem 14, we need to show that

$$B \not\subseteq \overrightarrow{T}_{II,n}(\Phi_n, \mathcal{N}_X(A_n))$$

for all $n \geq N$. So let $n \geq N$ and note that, since $\Phi_n \notin [\mathbf{0}; B, \varepsilon]$, there is some $x \in B$ and $y \in \Phi_n(x)$ so that $|y| \geq \varepsilon > 2^{-N} \geq 2^{-n}$. That is, $\Phi_n(x) \not\subseteq (-2^{-n}, 2^{-n})$ and so $x \notin \overrightarrow{T}_{II,n}(\Phi_n, \mathcal{N}_X(A_n))$. This finishes the proof. \square

Though particular applications of Corollaries 37 and 40 abound, we record a few that capture the general spirit using ideals of usual interest after recalling some other facts and some names for particular selection principles. We suppress the relationships with $C_{\mathcal{A}}(X)$ in the following applications in the interest of space.

Definition 42. We identify some particular selection principles by name.

- $S_1(\Omega_{X,x}, \Omega_{X,x})$ is known as the **strong countable fan-tightness property for X at x** .
- $S_1(\mathcal{O}_X, \Omega_{X,x})$ is known as the **strong countable dense fan-tightness property for X at x** .
- $S_1(\mathcal{T}_X, CD_X)$ is known as the **discretely selective property for X** .
- We refer to $S_1(\Omega_X, \Omega_X)$ as the **ω -Rothberger property** and $S_1(\mathcal{K}_X, \mathcal{K}_X)$ as the **k -Rothberger property**.

Definition 43. For a partially ordered set (\mathbb{P}, \leq) and collections $\mathcal{A}, \mathcal{B} \subseteq \mathbb{P}$ so that, for every $B \in \mathcal{B}$, there exists some $A \in \mathcal{A}$ with $B \subseteq A$, we define the **cofinality of \mathcal{A} relative to \mathcal{B}** by

$$\text{cof}(\mathcal{A}; \mathcal{B}, \leq) = \min\{\kappa \in \text{CARD} : (\exists \mathcal{F} \in [\mathcal{A}]^\kappa)(\forall B \in \mathcal{B})(\exists A \in \mathcal{F}) B \subseteq A\}$$

where CARD is the class of cardinals.

Lemma 44. Let $\mathcal{A}, \mathcal{B} \subseteq \wp^+(X)$ for a space X .

As long as X is T_1 ,

$$I \uparrow_{\text{pre}} G_1(\mathcal{N}_X[\mathcal{A}], \neg \mathcal{O}_X(\mathcal{B})) \iff \text{cof}(\mathcal{A}; \mathcal{B}, \subseteq) \leq \omega.$$

(See [13, 32], and [3, Lemma 23].)

If \mathcal{A} consists of G_δ sets,

$$\begin{aligned} I \uparrow G_1(\mathcal{N}_X[\mathcal{A}], \neg \mathcal{O}_X(\mathcal{B})) &\iff I \uparrow_{\text{pre}} G_1(\mathcal{N}_X[\mathcal{A}], \neg \mathcal{O}_X(\mathcal{B})) \\ &\iff \text{cof}(\mathcal{A}; \mathcal{B}, \subseteq) \leq \omega. \end{aligned}$$

(See [12, 30] and [3, Lemma 24].)

Observe that Lemma 44 informs us that, for a T_1 space X ,

- $\text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathbb{P}_X, \neg\mathcal{O}_X)$ if and only if X is countable,
- $\text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_X[K(X)], \neg\mathcal{O}_X)$ if and only if X is σ -compact, and
- $\text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_X[K(X)], \neg\mathcal{K}_X)$ if and only if X is hemicompact.

Corollary 45. For an ideal \mathcal{A} of closed subsets of a T_1 space X , $\text{cof}(\mathcal{A}; \mathcal{A}, \subseteq) \leq \omega$ if and only if $MU_{\mathcal{A}}(X)$ is metrizable.

Proof. If $\{A_n : n \in \omega\} \subseteq \mathcal{A}$ is so that, for every $A \in \mathcal{A}$, there is an $n \in \omega$ with $A \subseteq A_n$, notice that the family $\{\mathbf{W}(A_n, 2^{-m}) : n, m \in \omega\}$ is a countable base for the uniformity on $MU_{\mathcal{A}}(X)$; so Theorem 18 demonstrates that $MU_{\mathcal{A}}(X)$ is metrizable.

Now, suppose $MU_{\mathcal{A}}(X)$ is metrizable, which implies that $MU_{\mathcal{A}}(X)$ is first-countable. Using a descending countable basis at $\mathbf{0}$, we see that

$$\text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_{MU_{\mathcal{A}}(X), \mathbf{0}}, \neg\Gamma_{MU_{\mathcal{A}}(X), \mathbf{0}}),$$

and, in particular,

$$\text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_{MU_{\mathcal{A}}(X), \mathbf{0}}, \neg\Omega_{MU_{\mathcal{A}}(X), \mathbf{0}}).$$

By Corollary 40, we see that

$$\text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\mathcal{O}_X(\mathcal{A})).$$

So, by Lemma 44, $\text{cof}(\mathcal{A}; \mathcal{A}, \subseteq) \leq \omega$. □

As a particular consequence of this, we see that

Corollary 46. For any regular space X , the following are equivalent.

- (i) X is countable.
- (ii) $MU_p(X)$ is metrizable.
- (iii) $MU_p(X)$ is not discretely selective.
- (iv) $\text{II} \uparrow_{\text{mark}} \mathbf{G}_1(\Omega_X, \Omega_X)$.
- (v) $\text{II} \uparrow_{\text{mark}} \mathbf{G}_1(\Omega_{MU_p(X), \mathbf{0}}, \Omega_{MU_p(X), \mathbf{0}})$.
- (vi) $\text{II} \uparrow_{\text{mark}} \mathbf{G}_1(\mathcal{D}_{MU_p(X)}, \Omega_{MU_p(X), \mathbf{0}})$.

Also, the following are equivalent.

- (i) X is hemicompact.
- (ii) $\mathbb{K}(X)$ is hemicompact. (See [2, Thm. 3.22].)
- (iii) $MU_k(X)$ is metrizable. (See [17, Cor. 4.5].)
- (iv) $MU_k(X)$ is not discretely selective.
- (v) $\text{II} \uparrow_{\text{mark}} \mathbf{G}_1(\mathcal{K}_X, \mathcal{K}_X)$.
- (vi) $\text{II} \uparrow_{\text{mark}} \mathbf{G}_1(\Omega_{MU_k(X), \mathbf{0}}, \Omega_{MU_k(X), \mathbf{0}})$.
- (vii) $\text{II} \uparrow_{\text{mark}} \mathbf{G}_1(\mathcal{D}_{MU_k(X)}, \Omega_{MU_k(X), \mathbf{0}})$.

Before the next corollary, we recall Tkachuk's strategy strengthening for the generalized point-open games.

Theorem 47 (See [3, Cor. 11] and [32]). Let \mathcal{A} and \mathcal{B} be ideals of closed subsets of X . Then

$$I \uparrow G_1(\mathcal{N}_X[\mathcal{A}], \neg\mathcal{O}_X(\mathcal{B})) \iff I \uparrow G_1(\mathcal{N}_X[\mathcal{A}], \neg\Gamma_X(\mathcal{B}))$$

and

$$I \uparrow_{\text{pre}} G_1(\mathcal{N}_X[\mathcal{A}], \neg\mathcal{O}_X(\mathcal{B})) \iff I \uparrow_{\text{pre}} G_1(\mathcal{N}_X[\mathcal{A}], \neg\Gamma_X(\mathcal{B}))$$

Corollary 48. For any regular space X , the following are equivalent.

- (i) $\text{II} \uparrow G_1(\Omega_X, \Omega_X)$.
- (ii) $\text{II} \uparrow G_1(\Omega_{MU_p(X)}, \mathbf{0}, \Omega_{MU_p(X)}, \mathbf{0})$.
- (iii) $\text{II} \uparrow G_1(\mathcal{D}_{MU_p(X)}, \Omega_{MU_p(X)}, \mathbf{0})$.
- (iv) $I \uparrow G_1(\mathcal{T}_{MU_p(X)}, \text{CD}_{MU_p(X)})$.
- (v) $I \uparrow G_1(\mathcal{N}_X[[X]^{<\omega}], \neg\Omega_X)$.
- (vi) $I \uparrow G_1(\mathcal{N}_X[[X]^{<\omega}], \neg\Gamma_\omega(X))$.

Also, the following are equivalent.

- (i) $\text{II} \uparrow G_1(\mathcal{K}_X, \mathcal{K}_X)$.
- (ii) $\text{II} \uparrow G_1(\Omega_{MU_k(X)}, \mathbf{0}, \Omega_{MU_k(X)}, \mathbf{0})$.
- (iii) $\text{II} \uparrow G_1(\mathcal{D}_{MU_k(X)}, \Omega_{MU_k(X)}, \mathbf{0})$.
- (iv) $I \uparrow G_1(\mathcal{T}_{MU_k(X)}, \text{CD}_{MU_k(X)})$.
- (v) $I \uparrow G_1(\mathcal{N}_X[K(X)], \neg\mathcal{K}_X)$.
- (vi) $I \uparrow G_1(\mathcal{N}_X[K(X)], \neg\Gamma_k(X))$.

In general, Corollaries 46 and 48 are strictly separate, as the following example demonstrates.

Example 49. Let X be the one-point Lindelöfication of ω_1 with the discrete topology. In [2, Ex. 3.24], it is shown that X has the property that $\text{II} \uparrow G_1(\mathcal{K}_X, \mathcal{K}_X)$, but $\text{II} \not\uparrow_{\text{mark}} G_1(\mathcal{K}_X, \mathcal{K}_X)$.

However, according to Theorem 50, if Two can win against predetermined strategies in some Rothberger-like games, Two can actually win against full-information strategies in those games.

Theorem 50. Let X be any space.

- (i) By Pawlikowski [26],

$$I \uparrow_{\text{pre}} G_1(\mathcal{O}_X, \mathcal{O}_X) \iff I \uparrow G_1(\mathcal{O}_X, \mathcal{O}_X).$$

- (ii) By Scheepers [29] (see also [5, Cor. 4.12]),

$$I \uparrow_{\text{pre}} G_1(\Omega_X, \Omega_X) \iff I \uparrow G_1(\Omega_X, \Omega_X).$$

- (iii) By [5, Thm. 4.21],

$$I \uparrow_{\text{pre}} G_1(\mathcal{K}_X, \mathcal{K}_X) \iff I \uparrow G_1(\mathcal{K}_X, \mathcal{K}_X).$$

Corollary 51. For any regular space X , the following are equivalent.

- (i) X is ω -Rothberger.
- (ii) $X^{<\omega}$ is Rothberger, where $X^{<\omega}$ is the disjoint union of X^n for all $n \geq 1$. (See [27] and [5, Cor. 3.11].)
- (iii) $\mathcal{P}_{\text{fin}}(X)$ is Rothberger, where $\mathcal{P}_{\text{fin}}(X)$ is the set $[X]^{<\omega}$ with the subspace topology inherited from $\mathbb{K}(X)$. (See [5, Cor. 4.11].)
- (iv) I $\nabla \mathbf{G}_1(\Omega_X, \Omega_X)$.
- (v) $MU_p(X)$ has strong countable fan-tightness at $\mathbf{0}$.
- (vi) $MU_p(X)$ has strong countable dense fan-tightness at $\mathbf{0}$.
- (vii) II $\nabla \mathbf{G}_1(\mathcal{N}_X[[X]^{<\omega}], \neg\Omega_X)$.
- (viii) II $\overset{\text{mark}}{\nabla} \mathbf{G}_1(\mathcal{N}_X[[X]^{<\omega}], \neg\Omega_X)$.
- (ix) II $\nabla \mathbf{G}_1(\mathcal{T}_{MU_p(X)}, \text{CD}_{MU_p(X)})$.
- (x) II $\overset{\text{mark}}{\nabla} \mathbf{G}_1(\mathcal{T}_{MU_p(X)}, \text{CD}_{MU_p(X)})$.
- (xi) II $\nabla \mathbf{G}_1(\mathcal{N}_{MU_p(X), \mathbf{0}}, \neg\Omega_{MU_p(X), \mathbf{0}})$.
- (xii) II $\overset{\text{mark}}{\nabla} \mathbf{G}_1(\mathcal{N}_{MU_p(X), \mathbf{0}}, \neg\Omega_{MU_p(X), \mathbf{0}})$.

Also, the following are equivalent.

- (i) X is k -Rothberger.
- (ii) I $\nabla \mathbf{G}_1(\mathcal{K}_X, \mathcal{K}_X)$.
- (iii) $MU_k(X)$ has strong countable fan-tightness at $\mathbf{0}$.
- (iv) $MU_k(X)$ has strong countable dense fan-tightness at $\mathbf{0}$.
- (v) II $\nabla \mathbf{G}_1(\mathcal{N}_X[K(X)], \neg\mathcal{K}_X)$.
- (vi) II $\overset{\text{mark}}{\nabla} \mathbf{G}_1(\mathcal{N}_X[K(X)], \neg\mathcal{K}_X)$.
- (vii) II $\nabla \mathbf{G}_1(\mathcal{T}_{MU_k(X)}, \text{CD}_{MU_k(X)})$.
- (viii) II $\overset{\text{mark}}{\nabla} \mathbf{G}_1(\mathcal{T}_{MU_k(X)}, \text{CD}_{MU_k(X)})$.
- (ix) II $\nabla \mathbf{G}_1(\mathcal{N}_{MU_k(X), \mathbf{0}}, \neg\Omega_{MU_k(X), \mathbf{0}})$.
- (x) II $\overset{\text{mark}}{\nabla} \mathbf{G}_1(\mathcal{N}_{MU_k(X), \mathbf{0}}, \neg\Omega_{MU_k(X), \mathbf{0}})$.

Using the techniques of Gerlits and Nagy [13], Galvin [12], and Telgársky [30], we offer an analog of Lemma 44 related to the so-called *weak k -covering number* in [17].

Definition 52. For a collection \mathcal{A} of closed subsets of X , define the **weak covering number** of \mathcal{A} to be

$$wkc(\mathcal{A}) = \min \left\{ \kappa \in \text{CARD} : (\exists \mathcal{F} \in [\mathcal{A}]^\kappa) X = \text{cl} \left(\bigcup \mathcal{F} \right) \right\}.$$

Definition 53. For a space X , let DO_X be the set of all $\mathcal{U} \subseteq \mathcal{T}_X$ so that $X = \text{cl}(\bigcup \mathcal{U})$.

Lemma 54. Let $\mathcal{A} \subseteq \wp^+(X)$ where X is a regular space. Then

$$\text{I } \overset{\text{pre}}{\nabla} \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\text{DO}_X) \iff wkc(\mathcal{A}) \leq \omega.$$

If X is metrizable and $\mathcal{A} \subseteq K(X)$, then

$$\begin{aligned} \text{I} \uparrow \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\text{DO}_X) &\iff \text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\text{DO}_X) \\ &\iff \text{wkc}(\mathcal{A}) \leq \omega. \end{aligned}$$

Proof. The implications

$$\text{wkc}(\mathcal{A}) \leq \omega \implies \text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\text{DO}_X)$$

and

$$\text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\text{DO}_X) \implies \text{I} \uparrow \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\text{DO}_X)$$

are evident and hold with no assumptions on X .

Suppose X is regular and that $\text{wkc}(\mathcal{A}) > \omega$. Any predetermined strategy for One in $\mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\text{DO}_X)$ corresponds to a sequence $\langle A_n : n \in \omega \rangle \in \mathcal{A}^\omega$. We show that no such strategy can be winning for One. Since $\bigcup\{A_n : n \in \omega\}$ is not dense in X , we can find $U \in \mathcal{T}_X$ so that $U \cap \bigcup\{A_n : n \in \omega\} = \emptyset$. As X is assumed to be regular, without loss of generality, we can assume that $\text{cl}(U) \cap \bigcup\{A_n : n \in \omega\} = \emptyset$. It follows that $A_n \subseteq X \setminus \text{cl}(U) =: V$ for all $n \in \omega$. Hence, $\langle \mathcal{N}_X(A_n) : n \in \omega \rangle$ is not a winning strategy for One. That is, $\text{I} \not\uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\text{DO}_X)$.

Now suppose that X is metrizable, $\mathcal{A} \subseteq K(X)$, and $\text{I} \uparrow \mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\text{DO}_X)$. Let σ be a winning strategy for One which is coded by elements of \mathcal{A} . Let $\mathbb{T}_0 = \langle \rangle$ and, for $n \in \omega$,

$$\mathbb{T}_{n+1} = \left\{ w \widehat{\ } \langle \sigma(w), \mathbb{B}(\sigma(w), 2^{-\ell}) \rangle : w \in \mathbb{T}_n \wedge \ell \in \omega \right\}.$$

Notice that $\mathcal{F} = \bigcup_{n \in \omega} \{\sigma(w) : w \in \mathbb{T}_n\}$ is a countable subset of \mathcal{A} .

We show that $X = \text{cl}(\bigcup \mathcal{F})$ by way of contradiction. Suppose $X \setminus \text{cl}(\bigcup \mathcal{F}) \neq \emptyset$ and let $V \in \mathcal{T}_X$ be open so that $\text{cl}(V) \cap \text{cl}(\bigcup \mathcal{F}) = \emptyset$. For $A_0 := \sigma(\emptyset)$, let $\ell_0 \in \omega$ be so that $\text{cl}(V) \cap \mathbb{B}(A_0, 2^{-\ell_0}) = \emptyset$ and set $w_0 = \langle A_0, \mathbb{B}(A_0, 2^{-\ell_0}) \rangle \in \mathbb{T}_1$.

For $n \in \omega$, suppose we have $w_n \in \mathbb{T}_{n+1}$ defined. Now, for $A_{n+1} := \sigma(w_n)$, let $\ell_{n+1} \in \omega$ be so that $\text{cl}(V) \cap \mathbb{B}(A_{n+1}, 2^{-\ell_{n+1}}) = \emptyset$. Then define $w_{n+1} = w_n \widehat{\ } \langle A_{n+1}, \mathbb{B}(A_{n+1}, 2^{-\ell_{n+1}}) \rangle \in \mathbb{T}_{n+2}$.

Now, we have a run of the game according to σ with the property that

$$\text{cl}(V) \cap \bigcup_{n \in \omega} \mathbb{B}(A_n, 2^{-\ell_n}) = \emptyset.$$

It follows that

$$V \cap \text{cl} \left(\bigcup_{n \in \omega} \mathbb{B}(A_n, 2^{-\ell_n}) \right) = \emptyset,$$

which contradicts the assumption that σ is a winning strategy. \square

Consequently, for a regular space X ,

- $\text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_X[[X]^{<\omega}], \neg\text{DO}_X)$ if and only if X is separable and

- $\text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_X[K(X)], \neg\text{DO}_X)$ if and only if X admits a countable collection of compact subsets whose union is dense.

Also, by Lemma 13, $\mathbf{G}_1(\mathcal{N}_X[\mathcal{A}], \neg\text{DO}_X)$ is dual to $\mathbf{G}_1(\mathcal{O}_X(\mathcal{A}), \text{DO}_X)$. Hence, we obtain

Corollary 55. If X is a regular space, then

$$\begin{aligned} X \text{ is separable} &\iff \text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_X[[X]^{<\omega}], \neg\text{DO}_X) \\ &\iff \text{II} \uparrow_{\text{mark}} \mathbf{G}_1(\Omega_X, \text{DO}_X) \end{aligned}$$

and

$$\begin{aligned} wkc(K(X)) \leq \omega &\iff \text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_X[K(X)], \neg\text{DO}_X) \\ &\iff \text{II} \uparrow_{\text{mark}} \mathbf{G}_1(\mathcal{K}_X, \text{DO}_X). \end{aligned}$$

If X is, in addition, metrizable, then

$$\begin{aligned} X \text{ is separable} &\iff \text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_X[[X]^{<\omega}], \neg\text{DO}_X) \\ &\iff \text{I} \uparrow \mathbf{G}_1(\mathcal{N}_X[[X]^{<\omega}], \neg\text{DO}_X) \\ &\iff \text{II} \uparrow_{\text{mark}} \mathbf{G}_1(\Omega_X, \text{DO}_X) \\ &\iff \text{II} \uparrow \mathbf{G}_1(\Omega_X, \text{DO}_X) \end{aligned}$$

and

$$\begin{aligned} wkc(K(X)) \leq \omega &\iff \text{I} \uparrow_{\text{pre}} \mathbf{G}_1(\mathcal{N}_X[K(X)], \neg\text{DO}_X) \\ &\iff \text{I} \uparrow \mathbf{G}_1(\mathcal{N}_X[K(X)], \neg\text{DO}_X) \\ &\iff \text{II} \uparrow_{\text{mark}} \mathbf{G}_1(\mathcal{K}_X, \text{DO}_X) \\ &\iff \text{II} \uparrow \mathbf{G}_1(\mathcal{K}_X, \text{DO}_X). \end{aligned}$$

4. QUESTIONS

It is tempting to conjecture that an analogous result to Theorem 35 holds for the space of minimal usco maps, except that the natural candidate of pointwise set difference between two minimal usco maps may not generally produce a minimal usco map. For $A, B \subseteq \mathbb{R}$, let $A - B = \{x - y : x \in A, y \in B\}$ and consider $\Phi := \bar{\mathbf{I}}_{[0,1]}$. Note that the pointwise difference $\Phi - \Phi : X \rightarrow \mathbb{R}$ defined by $\Phi(x) - \Phi(x)$ has the property that

$$\Phi(x) - \Phi(x) = \begin{cases} \{0\}, & x \notin \{0, 1\}; \\ \{-1, 0, 1\}, & x \in \{0, 1\}. \end{cases}$$

Since $\text{gr}(\mathbf{0}) \subseteq \text{gr}(\Phi - \Phi)$, we see that $\Phi - \Phi$ is not minimal.

We also show that the pointwise Pompeiu-Hausdorff distance between two minimal usco maps need not be quasicontinuous. For compact $A, B \subseteq \mathbb{R}$, let

$$H_d(A, B) = \inf\{\varepsilon > 0 : A \subseteq \mathbb{B}(B, \varepsilon) \wedge B \subseteq \mathbb{B}(A, \varepsilon)\}.$$

We can define $f : X \rightarrow \mathbb{R}$ by $f(x) = H_d(\overline{\mathbf{1}}_{[0,1]}(x), \overline{\mathbf{1}}_{\mathbb{R} \setminus [0,1]}(x))$ and observe that

$$f(x) = \begin{cases} 1, & x \notin \{0, 1\}; \\ 0, & x \in \{0, 1\}. \end{cases}$$

So f is not quasicontinuous. However, this doesn't mean that the pointwise Pompeiu-Hausdorff distance cannot be used to establish an analogous result to Theorem 35 for $MU_{\mathcal{A}}(X)$.

We end with a few questions.

Question 1. Is there a result similar to Theorem 35 for general uniform spaces? Along these lines, are there selection games that characterize the diagonal degree and the uniformity degree of a uniform space?

Question 2. Can results similar to Theorems 36 and 39 be established relative to $\Omega_{MU_{\mathcal{A}}(X), \Phi}$ for any $\Phi \in MU(X)$?

Question 3. How many of the equivalences and dualities of this paper can be established for games of longer length and for finite-selection games?

Question 4. How much of this theory can be recovered when we study $MU(X, Y)$ for $Y \neq \mathbb{R}$, for example, when Y is $[0, 1]$, any metrizable space, any topological group, or any uniform space?

REFERENCES

- [1] Doyel Barman and Alan Dow, *Selective separability and SS^+* , Topol. Proc. **37** (2011), 181–204.
- [2] Christopher Caruvana and Jared Holshouser, *Closed discrete selection in the compact open topology*, Topology Proc. **56** (2020), 25–55.
- [3] ———, *Selection games on continuous functions*, Topology Appl. **279** (2020), 107253.
- [4] ———, *Selection games on hyperspaces*, Topology Appl. **300** (2021), 107771.
- [5] ———, *Selection games and the Vietoris space*, Topology Appl. **307** (2022), 107772.
- [6] Charles Castaing and Michel Valadier, *Convex Analysis and Measurable Multifunctions*, Lecture Notes in Mathematics, Springer Berlin, Heidelberg, 1977.
- [7] Jens Peter Reus Christensen, *Theorems of Namioka and B. E. Johnson type for upper semicontinuous and compact valued set-valued mappings*, Proc. Am. Math. Soc. **86** (1982), 649–655.
- [8] Steven Clontz, *Dual selection games*, Topology Appl. **272** (2020), 107056.
- [9] Steven Clontz and Jared Holshouser, *Limited information strategies and discrete selectivity*, Topology Appl. **265** (2019), 106815.
- [10] E. F. Collingwood and A. J. Lohwater, *The theory of cluster sets*, Camb. Tracts Math. Math. Phys., vol. 56, Cambridge University Press, Cambridge, 1966.
- [11] R. V. Fuller, *Relations among continuous and various non-continuous functions*, Pac. J. Math. **25** (1968), 495–509.
- [12] Fred Galvin, *Indeterminacy of point-open games*, Bull. Acad. Pol. Sci., Sér. Sci. Math. Astron. Phys. **26** (1978), 445–449.
- [13] J. Gerlits and Zs. Nagy, *Some properties of $C(X)$. I*, Topology Appl. **14** (1982), 151–161.
- [14] Gary Gruenhage, *Infinite games and generalizations of first-countable spaces*, General Topology Appl. **6** (1976), 339–352.

- [15] Ľubica Holá and Dušan Holý, *New characterizations of minimalusco maps*, Rocky Mt. J. Math. **44** (2014), no. 6, 1851–1866.
- [16] ———, *Minimalusco and minimalusco maps and compactness*, J. Math. Anal. Appl. **439** (2016), no. 2, 737–744.
- [17] ———, *Minimalusco maps and cardinal invariants of the topology of uniform convergence on compacta*, RACSAM **116** (2022), 27.
- [18] John L. Kelley, *General Topology*, Graduate Texts in Mathematics, Springer New York, 1975.
- [19] Stefan Kempisty, *Sur les fonctions quasicontinues*, Fundam. Math. **19** (1932), no. 1, 184–197.
- [20] Lj.D.R. Kočinac, *Selected results on selection principles*, Proceedings of the Third Seminar on Geometry and Topology (Tabriz, Iran), July 15–17, 2004, pp. 71–104.
- [21] Alojzy Lechicki and Sandro Levi, *Extensions of semicontinuous multifunctions.*, Forum mathematicum **2** (1990), no. 4, 341–360.
- [22] Norman Levine, *Semi-open sets and semi-continuity in topological spaces*, Am. Math. Mon. **70** (1963), 36–41.
- [23] Zuquan Li, *Selection principles of the Fell topology and the Vietoris topology*, Topology Appl. **212** (2016), 90–104.
- [24] Ernest Michael, *Topologies on spaces of subsets*, Trans. Am. Math. Soc. **71** (1951), 152–182.
- [25] T. Neubrunn, *Quasi-continuity*, Real Anal. Exch. **14** (1989), no. 2, 259–306.
- [26] Janusz Pawlikowski, *Undetermined sets of point-open games*, Fundam. Math. **144** (1994), no. 3, 279–285.
- [27] Masami Sakai, *Property C''' and function spaces*, Proc. Am. Math. Soc. **104** (1988), no. 3, 917–919.
- [28] Marion Scheepers, *Combinatorics of open covers. I: Ramsey theory*, Topology Appl. **69** (1996), no. 1, 31–62.
- [29] ———, *Combinatorics of open covers. III: Games, $C_p(X)$* , Fundam. Math. **152** (1997), no. 3, 231–254.
- [30] Rastislav Telgársky, *Spaces defined by topological games*, Fundam. Math. **88** (1975), 193–223.
- [31] ———, *Topological games: on the 50th anniversary of the Banach-Mazur game*, Rocky Mt. J. Math. **17** (1987), 227–276.
- [32] V. V. Tkachuk, *A C_p -theory problem book. Functional equivalencies*, Probl. Books Math., Cham: Springer, 2016.
- [33] ———, *Closed discrete selections for sequences of open sets in function spaces*, Acta Math. Hung. **154** (2018), no. 1, 56–68.
- [34] ———, *Two point-picking games derived from a property of function spaces*, Quaest. Math. **41** (2018), no. 6, 729–743.

SCHOOL OF SCIENCES, INDIANA UNIVERSITY KOKOMO, KOKOMO, IN, 46902, UNITED STATES OF AMERICA

Email address: chcaru@iu.edu

URL: <https://chcaru.pages.iu.edu/>