A pilot study on canonical gravity with mass dimension one fermions

R. de C. Lima¹,* T. M. Guimarães²,[†] and S. H. Pereira^{3‡} ¹Instituto Federal de Educação,

Ciência e Tecnologia do Norte de Minas Gerais (IFNMG), Januária, MG, 39480-000, Brazil. ²Instituto Federal de Educação, Ciência e Tecnologia do Paraná (IFPR), Ivaiporã, PR, 86870-000, Brazil. and ³Universidade Estadual Paulista (UNESP) Faculdade de Engenharia de Guaratinguetá, Departamento de Física Guaratinguetá, SP, 12516-410, Brazil.

Abstract

The mass dimension one (MDO) fermionic field is built on a complete set of dual-helicity self-spinors of the charge conjugation operator, which obeys the statistic of Fermi-Dirac. These spinors are a potential candidate for the description of dark matter. The redefinition of the dual structure of this object holds a local adjacent theory and satisfies complete Lorentz invariance. In this work, we investigate important aspects of the interaction of this fermion with gravity in the light of a canonical formulation in ADM formalism. We construct an action via tetrad fields using a manifold on a family of space-like surfaces Σ_t that carries the MDO matter field, and additionally we propose a condition for the action to have a term associated with dark energy. We found the Hamiltonian and diffeomorphism constraints at the classical level for gravitational dynamics with the immersion of this material content in space-time, which leads us to the interpretation of the contribution of dark matter energy density, parallel to the lapse function of foliation and its directional flux of energy density in the hypersurface of manifold.

PACS numbers: 11.10.-z, 04.20.Fy, 95.35.+d

I. INTRODUCTION

Proposed in its first version in 2005, the fermionic field of spin-1/2 with mass dimension one, abbreviated to MDO, is built on a complete set of eigenspinors of the charge conjugation operator C, the so called $Elko^1$ spinors [1]. The construction of the field is characterized by the presence of spinors belonging to the $(0, 1/2) \oplus (1/2, 0)$ family of the Weyl representation space. In its original formulation, these fields were quantum objects based on a representation of the subgroups HOM(2) and SIM(2) of the Lorentz group, whose algebra is based on Very Special Relativity (VSR) [2]. A few years ago, a modification of the dual of the field - taking advantage of the fact that only bilinears are observable in nature - endowed the field with complete Lorentz (Poincaré) symmetry [3]. Thus, the old non-local field turns into a local one after a redefinition of phase factors in the right and left-hand components of the field. These recent developments put the theory on solid grounds and stimulated several works in broad areas of Physics, with numerous results explored in the book [4], in Particle Phenomenology [5] and Cosmology [6], as brief examples. More recently, Ref. [7] presents a detailed discussion of duals and adjoints of the field, and the fermionic self-interaction and interactions with a real scalar field at one-loop shows that the earlier problem of unitarity violation are absent. Also, a quantum field theoretic calculation establishes the Newtonian gravitational interaction for a mass dimension

^{*}Electronic address: rodrigo.lima@ifnmg.edu.br

[†]Electronic address: thiago.moreira@ifpr.edu.br

[‡]Electronic address: s.pereira@unesp.br

¹ Elko is a german acronym for *Eigenspinoren des Ladungskonjugationsoperators*

one dark matter candidate. The partition function and main thermodynamic properties were studied and a review on the localization of higher-dimensional ELKOs on flat and bent branes were discussed.

The crucial difference between Dirac and MDO spinors is related to parity. For Dirac spinors, the parity is intrinsic to the theory and, as a direct consequence, Dirac dynamics is achieved. For the MDO spinors this does not happen, as well analyzed in [8]. However, since its construction is relativistic, the Klein-Gordon dynamics is assured, thus, its quantum field must inherit this dynamic. The MDO field is electrically neutral because it is eigenspinor of the charge-conjugation operator and since it presents a null or almost null interaction with photons, it has become a prominent theoretical dark matter candidate.

Some works that focused on understanding the interaction of the MDO fermion with gravity have been studied and deserve to be highlighted, such as in quantum field theory in curved spaces [9] and in the covariant formulation, via gravitons interactions [10]. Now we propose to investigate the possible interaction between MDO fermions and gravity, through the canonical (Hamiltonian) formulation. Our motivation is the fact that the canonical approach was born as an attempt to build a quantum theory in which the metric functions are exempt from background perturbation, unlike the covariant perturbative formulation.

The paper is organized as follow. In the Section (II), we extend the gravitational action developed by Palatini-Holst, incorporating fermionic matter with mass dimension one and spin-1/2, through the tetrad-ADM formalism. In Section (III), we investigate the classical hamiltonian and diffeomorphism constraints arising from this dark matter field with gravity. As a consequence, an energy density and associated directional energy flux density emerges in conjunction with the MDO-gravity coupling, which furnish a term similar to the cosmological constant. Conclusions are in Section IV.

II. MASS DIMENSION ONE FERMIONS ACTION COUPLED TO CANONICAL GRAVITY

A. Notation and preliminaries

For the purpose of studying the connection of matter to space-time through the canonical treatment of gravity, it is necessary to deal with certain preliminary notations.

The Dirac gamma matrices, γ^{μ} , in the Weyl basis are expressed by [11]:

$$\gamma^{0} = \begin{pmatrix} 0_{2\times2} & 1_{2\times2} \\ 1_{2\times2} & 0_{2\times2} \end{pmatrix}, \quad \gamma^{i} = \begin{pmatrix} 0_{2\times2} & -\sigma^{i} \\ \sigma^{i} & 0_{2\times2} \end{pmatrix}, \tag{1}$$

where σ^i are the Pauli matrices:

$$\sigma^{1} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^{2} = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^{3} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$
 (2)

The tetrad vector fields $e^a_{\mu}(x)$ are defined by $e^a_{\mu}(x)e^b_{\nu}(x)\eta_{ab} = g_{\mu\nu}(x)$, being $g_{\mu\nu}$ the metric of spacetime and $\eta_{ab} = (1, -1, -1, -1)$ the metric of a local minkowskian manifold. The matrices- γ in the arbitrary spacetime are given by $\gamma^{\mu} = e^a_{\mu}\gamma^a$ and obey Clifford's algebra:

$$\{\gamma^{\mu}, \gamma^{\nu}\} = \gamma^{\mu}\gamma^{\nu} + \gamma^{\nu}\gamma^{\mu} = 2g^{\mu\nu} \tag{3}$$

and

$$\gamma^{5} = i\gamma^{0}\gamma^{1}\gamma^{2}\gamma^{3} = \begin{pmatrix} 1_{2\times 2} & 0_{2\times 2} \\ 0_{2\times 2} & -1_{2\times 2} \end{pmatrix}.$$
 (4)

In Quantum Field Theory (QFT) is used, in general, the metric notation (+, -, -, -). On the other hand, in General Relativity (GR) and Loop Quantum Gravity (LQG) the usual notation is (-, +, +, +). To transcribe the fermionic fields in QFT to GR/LQG, we just multiply all the Dirac matrices (1) by $i \in Im(\mathbb{C})$ [12]. In this paper we will make use of this last notation. Note that in fact Clifford algebra does not change,

$$\{\gamma^{\mu}, \gamma^{\nu}\}_{(QFT)} = 2g^{\mu\nu} = 2(+, -, -, -) \longmapsto \{\gamma^{\mu}, \gamma^{\nu}\}_{(GR)} = \{i\gamma^{\mu}, i\gamma^{\nu}\} = i^{2}\{\gamma^{\mu}, \gamma^{\nu}\}_{(QFT)} = i^{2}2g^{\mu\nu} \Leftrightarrow \{\gamma^{\mu}, \gamma^{\nu}\}_{(GR)} = -2g^{\mu\nu} = 2(-, +, +, +).$$

In curved spaces, the covariant derivatives acting on the usual and dual MDO fermionic fields are defined by [11]:

$$\nabla_{\mu}\lambda \equiv \partial_{\mu}\lambda - \Gamma_{\mu}\lambda, \quad \nabla_{\mu}\lambda \equiv \partial_{\mu}\lambda + \lambda\Gamma_{\mu}, \tag{5}$$

where λ is the usual MDO spinor field, $\overline{\lambda}$ is the dual MDO spinor field and the spin connection Γ_{μ} as well as the switch of two covariant derivatives acting on the spinor are expressed as (see Appendix (A)):

$$\Gamma_{\mu} = \frac{i}{4} \omega_{\mu}^{\ ab} \sigma_{ab}, \quad [\nabla_{\mu}, \nabla_{\nu}] \lambda = -\frac{i}{4} F_{\mu\nu}^{\ ab} \sigma_{ab} \lambda, \tag{6}$$

where $\omega_{\mu}^{\ ab}\sigma_{ab} = e_{\nu}^{a}\partial_{\mu}e^{\nu b} + e_{\nu}^{a}e^{\rho b}\Gamma_{\ \mu\rho}^{\nu}$ is the connection written via tetrad fields and the Levi-Civita connection, under indices of the spacetime (external indices μ) and of Lorentz (internal indices a). $F_{\mu\nu}^{\ ab} = \partial_{\mu}\omega_{\nu}^{\ ab} - \partial_{\nu}\omega_{\mu}^{\ ab} + \omega_{\mu}^{\ ac}\omega_{\nu c}^{\ b} - \omega_{\nu}^{\ ac}\omega_{\mu c}^{\ b}$ is the "curvature" due to the spin connection on the spinor λ , with similar function to the Riemann curvature, coming from the related connection, for tensors in curved spaces, and $\sigma_{ab} = i/2[\gamma_a, \gamma_b]$.

The set of the MDO spinor field is represented by eigenstate of the charge conjugation operator C for fields with spin 1/2 with two possible eigenvalues (±1). Thus, $\lambda = (\lambda^S, \lambda^A)$ are the eigenstates self-conjugated and anti self-conjugated associated with eigenvalues (+1, -1), respectively. In addition, each of these objects has a dual helicity $\alpha = \{\pm, \mp\}$ that emerges from its structures, as demonstrated in [1], and therefore may be expressed as four fields with notation $\lambda = (\lambda^S_{\{\pm, \mp\}}, \lambda^A_{\{\pm, \mp\}})$.

Recently, Ahluwalia demonstrated that the dual spinor fields associated with the mass dimension one fermion guarantee local symmetry of Lorentz, [3], allowing to obtain the self-conjugated and anti-self-conjugated duals as

$$\vec{\lambda}_{\alpha}^{S} = \tilde{\lambda}_{\alpha}^{S} \mathcal{A}, \quad \vec{\lambda}_{\alpha}^{A} = \tilde{\lambda}_{\alpha}^{A} \mathcal{B}, \tag{7}$$

with $\widetilde{\lambda}_{\alpha}(p^{\mu}) \stackrel{def}{=} [\Xi(p^{\mu})\lambda_{\alpha}(p^{\mu})]^{\dagger}\gamma^{0}$, where the "Dirac-type" operator is denoted as $\Xi(p^{\mu}) = (\mathcal{G}(p^{\mu})/m)\gamma_{\mu}p^{\mu}$, in which the matrix \mathcal{G} is explicitly given by

$$G(p^{\mu}) \stackrel{def}{=} \begin{pmatrix} 0 & 0 & 0 & -ie^{-i\varphi} \\ 0 & 0 & ie^{i\varphi} & 0 \\ 0 & -ie^{-i\varphi} & 0 & 0 \\ ie^{i\varphi} & 0 & 0 & 0 \end{pmatrix}.$$
 (8)

Assuming that such objects were constructed via Weyl's left hand spinors with the momentum in spherical coordinates: $p^{\mu} = (E, p \sin \theta \cos \varphi, p \sin \theta \sin \varphi, p \cos \theta)$ and $p = |\vec{p}|$ [1, 3]. The operators \mathcal{A} and \mathcal{B} are conditions to guarantee the locality of the theory when the sum of spin is extracted $\sum_{\alpha} \lambda_{\alpha}^{S/A} \overline{\lambda}_{\alpha}^{S/A}$ [3], using the so-called *tau-deformation*.

B. Building the MDO fermion action in canonical gravity

The Einstein-Hilbert action [13] with the MDO fermion action (dark matter) in curved space [9, 11, 14] is given by:

$$\mathcal{S} = \mathcal{S}_{EH} + \mathcal{S}_{MDO} = \frac{1}{16\pi G} \int_{\mathcal{M}} d^4 x \sqrt{-g} R + \frac{1}{2} \int_{\mathcal{M}} d^4 x \sqrt{-g} [g^{\mu\nu} (\nabla_\mu \vec{\lambda} \nabla_\nu \lambda) - m^2 \vec{\lambda} \lambda - \xi R \vec{\lambda} \lambda], \tag{9}$$

where g is the determinant of metric $g_{\mu\nu}$, R is the Ricci scalar curvature, $\nabla_{\mu} \lambda$ and $\nabla_{\nu} \lambda$ are the covariant derivatives acting on the dual and usual MDO fields, respectively, m is the mass of the field λ , characterizing the term of selfinteraction $\lambda \lambda$, and ξ a constant of coupling between the MDO spinor field and the gravitational field. The boundary term $\partial \mathcal{M}$ of the manifold \mathcal{M} is not being considered. Explicitly, the quadratic kinetic term of MDO spinor is described by:

$$\nabla_{\mu} \dot{\lambda} \nabla_{\nu} \lambda = (\partial_{\mu} \dot{\lambda} + \dot{\lambda} \Gamma_{\mu}) (\partial_{\nu} \lambda - \Gamma_{\nu} \lambda) = \partial_{\mu} \dot{\lambda} \partial_{\nu} \lambda - (\partial_{\mu} \dot{\lambda}) \Gamma_{\nu} \lambda + \dot{\lambda} \Gamma_{\mu} (\partial_{\nu} \lambda) - \dot{\lambda} \Gamma_{\mu} \Gamma_{\nu} \lambda$$
$$= \partial_{\mu} \vec{\lambda} \partial_{\nu} \lambda - (\partial_{\mu} \vec{\lambda}) \left[\frac{i}{4} \omega_{\nu}^{cd} \sigma_{cd} \lambda \right] + \vec{\lambda} \left[\frac{i}{4} \omega_{\mu}^{ab} \sigma_{ab} \right] (\partial_{\nu} \lambda) - \vec{\lambda} \left[\frac{i}{4} \frac{i}{4} \omega_{\mu}^{ab} \omega_{\nu}^{cd} \sigma_{ab} \sigma_{cd} \lambda \right].$$
(10)

Rewriting the action (9) in terms of the tetrad fields, assuming the Einstein-Cartan formalism, considering the gravitational action S_{EH} equivalent to the action of Palatini-Holst [15] and using the spin connection (5), we have

(see Appendix (B)):

$$S = \frac{1}{16\pi G} \int_{\mathcal{M}} d^{4}x (ee_{I}^{\mu} e_{J}^{\nu} P^{IJ}{}_{KL} F_{\mu\nu}{}^{KL}(\omega)) (1 - 8\pi G\xi \,\overline{\lambda} \,\lambda) + \frac{1}{2} \int_{\mathcal{M}} d^{4}x (e) [(e_{I}^{\mu} e_{J}^{\nu} \eta^{IJ} \partial_{\mu} \,\overline{\lambda} \,\partial_{\nu} \lambda) - m^{2} \,\overline{\lambda} \,\lambda] + \frac{i}{8} \int_{\mathcal{M}} d^{4}x (ee_{I}^{\mu} e_{J}^{\nu} \eta^{IJ}) [\overline{\lambda} \,\omega_{\mu}{}^{IJ} \sigma_{IJ} (\partial_{\nu} \lambda) - (\partial_{\mu} \,\overline{\lambda}) \omega_{\nu}{}^{MN} \sigma_{MN} \lambda - i/4 \,\overline{\lambda} \,(\omega_{\mu}{}^{IJ} \omega_{\nu}{}^{MN} \sigma_{IJ} \sigma_{MN}) \lambda],$$
(11)

where the curvature $F_{\mu\nu}{}^{KL}(\omega) = 2\partial_{[\mu}\omega_{\nu]}{}^{IJ} + [\omega_{\mu}, \omega_{\nu}]$ is described in terms of the Lorentz connection $\omega_{\mu}{}^{IJ}$. The term $P^{IJ}{}_{KL}$ of Holst action, as well as its inverse, in the compact form [12], is written as:

$$P^{IJ}_{\ \ KL} = \delta^{[I}_{K} \delta^{J]}_{L} - \frac{1}{\gamma} \frac{\epsilon^{IJ}_{\ \ KL}}{2}, \quad P^{-1}_{IJ}^{\ \ KL} = \frac{\gamma^{2}}{\gamma^{2} + 1} \left(\delta^{[K}_{I} \delta^{L]}_{J} + \frac{1}{\gamma} \frac{\epsilon_{IJ}^{\ \ KL}}{2} \right), \tag{12}$$

where γ (it can not to be confused with Dirac matrices γ^{μ}) is the parameter of Barbero-Immirzi and ϵ^{IJ}_{KL} is a tensor completely antisymmetric. It is important to clarify that the coupling term between the gravitational field and the fermionic field were merged in the Palatini-Holst action in order to propose $\xi = 2G^{-1}\rho_{vac}/(R \lambda \lambda)$, being dependent on purely scalar factors and, therefore, invariant. Such term assumes a similar function to the cosmological constant Λ , in the Einstein field equations in cosmological scenarios, for models of the type $\Lambda(t)$ [16–18], once $\Lambda = 8\pi\rho_{vac}$ and $\rho_{vac} = 5,96 \times 10^{-27} kg/m^3$ is the vacuum density [19]. It is also valid for scalar-tensor theories à la Brans-Dick [20–22]. This proposal is constructed for reference frames in which the Holst term recovers the exact scalar Ricci curvature, R, originally present in the Einstein-Hilbert action. Furthermore, it is noted that the matter part, in our case, is written as a sum of two parts in the action of the MDO field: the first part resembles the dynamics of a massive scalar field and the second part carries the quadratic information of the pure fermionic portion of the MDO field with spin connection (the last term integrated over the \mathcal{M} manifold of action (11)). This is due to the Klein-Gordon dynamic nature of energy conservation and the structure of this fermionic field, respectively.

A canonical (Hamiltonian) formalism is built from a Legendre transformation in action. For this, it will be necessary to transcribe the action (11) from manifold \mathcal{M} , provided with a metric, to a family of constant foliations Σ_t , and determined by the function of time t, which are embedded in the manifold $(\mathcal{M}, g_{ab}) \mapsto (\Sigma_t, t) \equiv \Sigma_t \times \mathbb{R} \subseteq \mathcal{M}$, as originally prescribed by [23], known as ADM (Arnowitt-Deser-Misner) formalism. Instead of working with spacetime tensors, we will use the description of how a vector field t^{μ} evolves from a hypersurface Σ_t to another, $\Sigma_{t+\delta t}$, in direction to future, since it is using a Lorentzian metric signature. Following the same prescription used in [12], the four-vector, in spacetime, t^{μ} is

$$t^{\mu} = N n^{\mu} + N^{\mu}, \tag{13}$$

where n^{μ} is the vector normal to the hypersurface Σ_t , N^{μ} is the *shift* vector tangent to the surface Σ_t and N is the *lapse* function which dictates the temporal transition from one hypersurface to another, therefore, $N^{\mu}n_{\mu} = 0$. The spatial evolution of t^a must obey $t^a \nabla_a t = 1$, being a, b, c,... spatial tensor indices. The metric $g_{\mu\nu}$ is described, therefore, as

$$g_{\mu\nu} = q_{\mu\nu} - n_{\mu}n_{\nu}.$$
 (14)

Since we are using the tetrad formalism in addition to the ADM formalism, it is necessary to perform a gauge fixation in internal vector fields of the tetrad, thus it can be decomposed into internal time units, as a vector and a *triad* (spatial part of the tetrad). Setting the internal vector field as a constant $n_I = -\delta_I^0$, where $n^I n_I = 1$, we get:

$$n^a = n^I e^a_I, \quad e^a_I = \varepsilon^a_I - n^a n_I, \tag{15}$$

where n^a is the unit normal to foliation, $\varepsilon_I^a n_a = \varepsilon_I^a n^I = 0$, with $\varepsilon_I^a := triad$. From equation (13), one has

$$n^a = N^{-1}(t^a - N^a), (16)$$

for the (14) projection of the normal and tangential fields in Σ_t .

Thus, using ADM formalism decomposition and tetrad fields, the action of MDO matter with the Palatini-Holst

gravitation is described as (see Appendix (C)):

$$S[\varepsilon, \omega, \lambda] = S_{PH,\xi} + S_{\lambda, \overline{\lambda}_{S}} + S_{\lambda, \overline{\lambda}_{F}}$$

$$= \frac{1}{16\pi G} \int_{\mathbb{R}} dt \int_{\Sigma} d^{3}x N \sqrt{q} \Omega_{IJ}^{ab} P^{IJ}{}_{KL} F_{ab} {}^{KL}(\omega) (1 - 8\pi G\xi \,\overline{\lambda} \,\lambda)$$

$$+ \frac{1}{2} \int_{\mathbb{R}} dt \int_{\Sigma} d^{3}x N \sqrt{q} [\Omega_{IJ}^{ab} \eta^{IJ} (\partial_{a} \,\overline{\lambda} \,\partial_{b} \lambda) - m^{2} \,\overline{\lambda} \,\lambda]$$

$$- \frac{1}{8} \int_{\mathbb{R}} dt \int_{\Sigma} d^{3}x N \sqrt{q} \Omega_{IJ}^{ab} \eta^{IJ} \{i/2(\overline{\lambda} \,\omega_{a}^{IJ} [\gamma_{I}, \gamma_{J}] (\partial_{b} \lambda) - (\partial_{a} \,\overline{\lambda}) \omega_{b}^{MN} [\gamma_{M}, \gamma_{N}] \lambda) + i/16 \,\overline{\lambda} \,(\omega_{a}^{IJ} \omega_{b}^{MN} [\gamma_{I}, \gamma_{J}] [\gamma_{M}, \gamma_{N}] \lambda\}, \qquad (17)$$

where $\Omega_{IJ}^{ab} = (\varepsilon_I^a \varepsilon_J^b - 2N^{-1}n_I t^a \varepsilon_J^b + 2N^{-1}N^a n_I \varepsilon_J^b)$. The subscripts in the action of matter, $S_{\lambda,\overline{\lambda}S}$ and $S_{\lambda,\overline{\lambda}F}$, are to emphasize the difference between two terms: the Klein Gordon (scalar) and the pure quadratic Dirac (fermionic), respectively, due to the aforementioned nature of the spinor field.

The equation (17), for now, presents the Lagrangian density of a dark fermionic matter coupled to gravity with the contribution of dark energy for models $\Lambda(t)$ or Brans-Dick theory, admitting $\xi = 2G^{-1}\rho_{vac}/(R \lambda \lambda)$.

III. HAMILTONIAN CONSTRAINT BETWEEN MDO AND GRAVITY

The action (17), and its Lagrangian density, is the most complete way of treating the Hamiltonian formalism in terms of the matter field of interest. In the scope of this work, the scenario without torsion and with the Palatini-Holst dual term recovers the usual Ricci scalar structure. Under a family of foliations Σ in the manifold and with the aid of the expression (9), one has

$$S = \frac{1}{16\pi G} \int_{\mathbb{R}} dt \int_{\Sigma} d^3 x N (G^{abcd} K_{ab} K_{cd} + \sqrt{q}^{(3)} R) [1 - 8\pi G \xi \,\vec{\lambda} \,\lambda] + \frac{1}{2} \int_{\mathbb{R}} dt \int_{\Sigma} d^3 x N \sqrt{q} [(q^{\mu\nu} + n^{\mu} n^{\nu}) (q^{\alpha}_{\mu} D_{\alpha} \,\vec{\lambda} \,q^{\beta}_{\nu} D_{\beta} \lambda) - m^2 \,\vec{\lambda} \,\lambda],$$
(18)

where $D_{\alpha} = q_{\alpha}^{\mu} \nabla_{\mu}$ is the projection of the covariant derivative operator on \mathcal{M} onto the hypersurface Σ_t and the pair (G^{abcd}, G_{abcd}) is the DeWitt metric, $K_{ab} = \Gamma^{\mu}_{ab} n_{\mu} = -\Gamma^{0}_{ab} / \sqrt{-g^{00}}$ is a symmetrical "velocity" associated with the three-metric q_{ab} [27], expressed in terms of the spatial components of the equation (14). Thus, both are written respectively as:

$$G^{abcd} = \frac{\sqrt{q}}{2} (q^{ac} q^{bd} + q^{ad} q^{bc} - 2q^{ab} q^{cd}), \qquad G_{abcd} = \frac{1}{2\sqrt{q}} (q_{ac} q_{bd} + q_{ad} q_{bc} - 2q_{ab} q_{cd})$$
(19)

and

$$K_{\mu\nu} \equiv q_{\mu}^{\ \rho} q_{\nu}^{\ \sigma} \nabla_{\rho} n_{\sigma} = q_{\mu}^{\ \rho} \nabla_{\rho} n_{\nu} = K_{\nu\mu}, \tag{20}$$

which characterizes a purely special form, as well as $q_{\mu\nu}$, furthermore $K_{\mu\nu}$ is orthogonal to n^{μ} ($K_{\mu\nu}n^{\mu} = K_{\mu\nu}n^{\nu} = 0$).

In order to seek a canonical approach that couples this dark matter candidate with gravity, we will start by choosing a configuration variable and defining the momentum for the MDO ($'p = \partial L/\partial \dot{q}'$), in order to ensure the Legendre transformation:

$$H_{\lambda,\vec{\lambda}} = p_{\lambda}\dot{\lambda} + p_{\vec{\lambda}}\dot{\vec{\lambda}} - \mathcal{L}_{\lambda,\vec{\lambda}}.$$
 (21)

Starting from the explicit Lagrangian density of the MDO fermion,

$$\mathcal{L}_{\lambda,\vec{\lambda}} = \frac{1}{2} \sqrt{-g} [\partial^{\nu} \vec{\lambda} \partial_{\nu} \lambda - \partial^{\nu} \vec{\lambda} \Gamma_{\nu} \lambda + \vec{\lambda} \Gamma^{\nu} \partial_{\nu} \lambda - \vec{\lambda} \Gamma^{\nu} \Gamma_{\nu} \lambda - m^{2} \vec{\lambda} \lambda], \qquad (22)$$

we can get the following operators

$$\frac{\partial \mathcal{L}_{\lambda,\bar{\lambda}}}{\partial(\partial_{\alpha}\lambda)} = \frac{\sqrt{-g}}{2} (\partial^{\nu} \,\bar{\lambda} \,\delta^{\alpha}_{\nu} + \bar{\lambda} \,\Gamma^{\nu}\delta^{\alpha}_{\nu}) \quad \text{and} \quad \frac{\partial \mathcal{L}_{\lambda,\bar{\lambda}}}{\partial(\partial_{\alpha} \,\bar{\lambda})} = \frac{\sqrt{-g}}{2} (\delta^{\alpha}_{\beta}\partial^{\beta}\lambda - \delta^{\alpha}_{\beta}\Gamma^{\beta}\lambda) \tag{23}$$

allowing us to construct their conjugate momentum as

$$p_{\lambda} = \frac{\partial \mathcal{L}_{\lambda,\vec{\lambda}}}{\partial(\partial_{0}\lambda)} = \frac{\sqrt{-g}}{2} (\partial^{0} \vec{\lambda} + \vec{\lambda} \Gamma^{0}) \quad \text{and} \quad p_{\vec{\lambda}} = \frac{\partial \mathcal{L}_{\lambda,\vec{\lambda}}}{\partial(\partial_{0} \vec{\lambda})} = \frac{\sqrt{-g}}{2} (\partial^{0}\lambda - \Gamma^{0}\lambda). \tag{24}$$

Then, the respective "velocity" of the field in terms of momentum is determined, with the help of (24), as

$$\dot{\lambda} = \nabla_0 \lambda = (\partial_0 \lambda - \Gamma_0 \lambda) = \frac{2}{\sqrt{-g}} g_{00} p_{\vec{\lambda}} \quad \text{and} \quad \dot{\vec{\lambda}} = \nabla_0 \ \vec{\lambda} = (\partial_0 \ \vec{\lambda} + \vec{\lambda} \ \Gamma_0) = \frac{2}{\sqrt{-g}} g_{00} p_{\lambda}. \tag{25}$$

We also note that the Lagrangian density (22) can be written more explicitly as

$$\mathcal{L}_{\lambda,\vec{\lambda}} = \frac{1}{2}\sqrt{-g} \underbrace{\left[\partial^{0} \vec{\lambda} \partial_{0}\lambda - \partial^{0} \vec{\lambda} \Gamma_{0}\lambda + \vec{\lambda} \Gamma^{0} \partial_{0}\lambda - \vec{\lambda} \Gamma^{0} \Gamma_{0}\lambda\right]}_{\mathbb{A}} - \frac{1}{2}\sqrt{-g} \left[\partial^{i} \vec{\lambda} \partial_{i}\lambda + m^{2} \vec{\lambda} \lambda\right] - \frac{1}{2}\sqrt{-g} \left[\vec{\lambda} \Gamma^{i} \partial_{i}\lambda - \partial^{i} \vec{\lambda} \Gamma_{i}\lambda - \vec{\lambda} \Gamma^{i} \Gamma_{i}\lambda\right],$$
(26)

where

$$\mathbb{A} = (\partial^0 \vec{\lambda} + \vec{\lambda} \Gamma^0)(\partial_0 \lambda - \Gamma_0 \lambda) = \frac{2}{\sqrt{-g}} \frac{2g_{00}}{\sqrt{-g}} (p_\lambda p_{\vec{\lambda}}).$$
(27)

Thus, one has

$$\mathcal{L}_{\lambda,\vec{\lambda}} = \frac{2}{\sqrt{-g}} g_{00}(p_{\lambda}p_{\vec{\lambda}}) - \frac{1}{2}\sqrt{-g} [\partial^{i}\vec{\lambda}\partial_{i}\lambda + m^{2}\vec{\lambda}\lambda] - \frac{1}{2}\sqrt{-g} [\vec{\lambda}\Gamma^{i}\partial_{i}\lambda - \partial^{i}\vec{\lambda}\Gamma_{i}\lambda - \vec{\lambda}\Gamma^{i}\Gamma_{i}\lambda].$$
(28)

With the Legendre transformation performed on $\mathcal{L}_{\lambda,\overline{\lambda}}$, in the expression (28), we develop the Hamiltonian density of the dark fermion in curved space,

$$\mathcal{H}_{\lambda,\vec{\lambda}} = p_{\lambda}\dot{\lambda} + p_{\vec{\lambda}}\dot{\vec{\lambda}} - \mathcal{L}_{\lambda,\vec{\lambda}} = \frac{2}{\sqrt{-g}}g_{00}(p_{\vec{\lambda}}p_{\lambda}) + \frac{1}{2}\sqrt{-g}(\partial^{i}\vec{\lambda}\partial_{i}\lambda + m^{2}\vec{\lambda}\lambda) + \frac{1}{2}\sqrt{-g}[\vec{\lambda}\Gamma^{i}\partial_{i}\lambda - \partial^{i}\vec{\lambda}\Gamma_{i}\lambda - \vec{\lambda}\Gamma^{i}\Gamma_{i}\lambda)]$$
(29)

which in turn allows us to finally describe the Hamiltonian of the MDO fermion in a foliation scenario, Σ_t , à lá ADM, in the form:

$$\begin{aligned}
H_{\lambda,\vec{\lambda}} &= \int_{\Sigma} d^{3}x \mathcal{H}_{\lambda,\vec{\lambda}} \\
&= \int_{\Sigma} d^{3}x N \left(\frac{2p_{\lambda}p_{\lambda}}{\sqrt{q}} + \frac{\sqrt{q}}{2} (q^{ab} \partial_{a} \vec{\lambda} \partial_{b} \lambda + m^{2} \vec{\lambda} \lambda) \right) \\
&+ \int_{\Sigma} d^{3}x N^{a} \left(\frac{1}{2} q^{b}_{a} [\vec{\lambda} \Gamma_{b} \partial_{c} \lambda - \partial_{b} \vec{\lambda} \Gamma_{c} \lambda - \vec{\lambda} \Gamma_{b} \Gamma_{c} \lambda] n^{c} \right) \\
&\equiv \int_{\Sigma} d^{3}x (N \mathcal{H}_{\perp,\lambda,\vec{\lambda}} + N^{a} \mathcal{H}_{a,\lambda,\vec{\lambda}}),
\end{aligned}$$
(30)

noting that the transformations $\sqrt{-g} \mapsto N\sqrt{q}$ and $\sqrt{-g}g^{ij}A_jB_i \mapsto N^a q^b_a(A_bB_c)n^c$ were performed to map $\mathcal{M} \longrightarrow \mathbb{R} \times \Sigma$.

The equation (30) allows us to visualize $\mathcal{H}_{\perp,\lambda,\overline{\lambda}}$ as the energy density of the MDO fermionic field $(T_{00} = \rho_{\lambda,\overline{\lambda}})$, which "flows" through the *lapse* function, and consequently orthogonal to Σ , at a given fixed instant of time t. The term $\mathcal{H}_{a,\lambda,\overline{\lambda}}$ can be interpreted as a "flux" $(J_{a,\lambda,\overline{\lambda}} \equiv q_a^{\ b} T_{bc} n^c)$ associated with the MDO fermion, which moves through the shift vector N^a tangential to the hypersurface Σ_t . It is notable that this directional energy density flux depends exclusively on the spatial derivative and the spin connection of the mass-dimension-one spinor field.

We can clarify the physical meaning of the Hamiltonian and diffeomorphism constraints for gravity in this scenery. It is important to note that in the absence of matter, the structures of gravitational constraints, denoted in the expression below by $\mathcal{H}_{\perp, g}$ (Hamiltonian) and $\mathcal{H}_{a, g}$ (diffeomorphism) respectively, are well established in the literature, according to [23, 27]. Now, we obtained a result with geometry of space-time and a peculiar material content, i.e., gravity and the MDO fermion, as developed from the action (18). Therefore, one has:

$$\mathcal{H}_{\perp} = \mathcal{H}_{\perp, g} + \mathcal{H}_{\perp, \lambda, \overline{\lambda}} = \left(16\pi G G_{abcd} p^{ab} p^{cd} - \frac{\sqrt{q}}{16\pi G} {}^{(3)} R\right) \left[1 - 8\pi G \xi \ \overline{\lambda} \ \lambda\right] + \rho_{\lambda, \overline{\lambda}} \approx 0 \quad \text{and} \tag{31}$$

$$\mathcal{H}_{a} = \mathcal{H}_{a, g} + \mathcal{H}_{a, \lambda, \overline{\lambda}} = -2D_{b}p_{a}^{\ b}[1 - 8\pi G\xi \ \overline{\lambda} \ \lambda] + J_{a, \lambda, \overline{\lambda}} \approx 0, \tag{32}$$

or even more explicitly, using the equation (29),

$$\mathcal{H}_{\perp} = \left(16\pi G G_{abcd} p^{ab} p^{cd} - \frac{\sqrt{q}}{16\pi G} R\right) \left[1 - 8\pi G \xi \,\vec{\lambda} \,\lambda\right] + \left(\frac{2p_{\vec{\lambda}} p_{\lambda}}{\sqrt{q}} + \frac{\sqrt{q}}{2} (q^{ab} \partial_a \,\vec{\lambda} \,\partial_b \lambda + m^2 \,\vec{\lambda} \,\lambda)\right) \approx 0 \quad \text{and} \quad (33)$$

$$\mathcal{H}_{a} = -2D_{b}p_{a}^{\ b}[1 - 8\pi G\xi \ \vec{\lambda} \ \lambda] + \left(\frac{1}{2}q_{a}^{b}[\vec{\lambda} \ \Gamma_{b}\partial_{c}\lambda - \partial_{b} \ \vec{\lambda} \ \Gamma_{c}\lambda - \vec{\lambda} \ \Gamma_{b}\Gamma_{c}\lambda]n^{c}\right) \approx 0, \tag{34}$$

remembering that the term $-8\pi G\xi \[\lambda] \lambda$ was joined to the Ricci scalar, in the geometric portion of the action, in order to guarantee a similar behavior to the cosmological constant in $\Lambda(t)$ models, from the coupling constant ξ between MDO and gravity, and from the orthonormal invariance $\[\lambda] (x)\lambda(x') = \pm 2m\delta(x-x')$ for the same helicities of the fermion, in its self-conjugated (S) and anti-self-conjugated (A) forms, respectively.

Just as a comparative analysis, when incorporating a scalar Lagrangian density field $\mathcal{L}_{\phi} = \sqrt{-g}(-1/2g^{\mu\nu}\nabla_{\mu}\phi\nabla_{\nu}\phi - 1/2m^{2}\phi^{2})$ to same gravitational scenario [27], the constraints of scalar field ϕ , $\mathcal{H}_{\perp,\phi}$ and $\mathcal{H}_{a,\phi}$, in curved space-time are given by

$$\mathcal{H}_{\perp,\phi} = \rho_{\phi} = \left(\frac{p_{\phi}^2}{2\sqrt{q}} + \frac{\sqrt{q}}{2}(q^{ab}\partial_a\phi\partial_b\phi + m^2\phi^2)\right) \quad e \quad \mathcal{H}_{a,\phi} = J_{a,\phi} = p_{\phi}\partial_a\phi. \tag{35}$$

It is possible to notice, in (33)-(34) and (35), that the energy density of the MDO fermionic field, $(\rho_{\lambda,\overline{\lambda}})$, closely resembles the scalar field (ρ_{ϕ}) , since the dynamic equation of both fields follows the Klein Gordon equation in the same way. However, while the "current" - directional density of energy flux - in the hypersurface Σ_t of the MDO field $(J_{a,\overline{\lambda},\lambda})$ is a function of the spatial derivatives and spin connections, the scalar field current is also a function of the spatial derivative, but it has no spin connection term, as expected. This difference is clarified due to the spinor structure of the fermion in curved space.

IV. CONCLUSION

In this work we present a proposal for the incorporation of the mass-dimension-one fermion with gravity, under a canonical approach. We start the first section with the Hamiltonian formulation of General Relativity using the ADM formalism [23], which foliate a pseudo-Riemannian manifold, equipped with a physical metric $(\mathcal{M}, g_{\mu\nu})$, for a globally hyperbolic spacetime, on Cauchy hypersurfaces Σ_t for each fixed $t \in \mathbb{R}$, being $\mathcal{M} \simeq \mathbb{R} \times \Sigma$. In this formulation, the existence of two fundamental forms, represented by the three-metric decomposition and the extrinsic curvature $(q_{\mu\nu}, K_{\mu\nu})$, in conjunction with the projection of the covariant derivative D_{μ} in the hypersurface, are responsible for characterizing the so-called ADM action, whose Legendre transformation on its Lagrangian density results in a geometric Hamiltonian and diffeomorphism constraints associated with gravity and which establish the so-called "dynamics of the theory in the absence of matter". In the subsequent section, we effectively incorporate the MDO fermion into the Einstein-Hilbert action. Aiming on a greater completeness via fermionic and gravitational interaction, the geometric part of the Einstein-Hilbert action was rewritten in terms of tetrads into an extension of Palatini [15], which holds a dual field for the original gravitational theory. The portion associated with the coupling constant between fermion and gravity was rewritten in order to reproduce the behavior of a cosmological constant term. Performing the same foliation process, in the light of the ADM formalism, we build an action that combines dark matter MDO and gravity via Palatini-Holst in ADM-tetrad (17), comprising the most complete form for a Hamiltonian analysis in terms of the field of interest.

Noticing that the dual term of Palatini-Holst, when it is null in eq (17), recovers the original form for the Ricci scalar, we can investigate the Hamiltonian formulation developed according to the ADM formalism. For our context, we will obtain the constraints between the gravitational hamiltonian density and the MDO fermion. After the proper Legendre transformation for the fermion Lagrangian density (22), in curved space, we were able to write the Hamiltonian in a

foliation scenario Σ_t by means of ADM formalism, which explained the Hamiltonian component $\mathcal{H}_{\perp,\lambda,\vec{\lambda}}$ as the energy density of the MDO fermionic field $(T_{00} = \rho_{\lambda,\vec{\lambda}})$, which "flows" through the lapse function, orthogonal to Σ at a given instant of fixed time t. Moreover, $\mathcal{H}_{a,\lambda,\vec{\lambda}}$ can be decomposed as a directional energy flux density, which is a function of spatial derivatives and spin connections of the MDO field under the representation $(J_{a,\lambda,\vec{\lambda}} \equiv q_a^{\ b} T_{bc} n^c)$, which moves through the shift vector N^a tangential to the hypersurface Σ_t .

In this way, we obtained the Hamiltonian and diffeomorphic constraints due to the incorporation of the MDO matter to the action, via ADM formalism, expressed by the equations (34)-(35), which elucidate three ingredients for canonical dynamics, namely: gravity, dark matter and dark energy. The latter one can be interpreted through the varying cosmological constant like term in $\Lambda(t)$ models, now encoded in the term $-8\pi G\xi \ \lambda \ \lambda$, where $\ \lambda \ \lambda$ is an orthonormal invariant for each and every observable, $\ \lambda \ (x)\lambda(x') = \pm 2m\delta(x-x')$, dictated for the same helicities of the fermion in its self-conjugated (S) and anti-self-conjugated (A) form, respectively.

The role of this coupling due to ξ can also be interpreted as a contributing part in Brans-Dicke [28] scalar-tensor theories. We also notice that the energy density $\rho_{\lambda,\overline{\lambda}}$ coming from $\mathcal{H}_{\perp,\lambda,\overline{\lambda}}$ has a similar form to that expected for a scalar field (ρ_{ϕ}) in the same scenario, unlike the structure observed for directional flow of energy between both matter fields, J_a . It should be noted that this construction related to the constraints was carried out on a general metric $g_{\mu\nu} = q_{\mu\nu} - n_{\mu}n_{\nu}$, thus being able to be applied to different scenarios of cosmological interests such as FLRW, black holes, wormholes among others.

Furthermore, a possible hypothesis raised for future investigations would be to resort to Ashtekar's new canonical variables $(E_i^a(x), A_b^j(x'))$ [29], which presently support the bases of the modern canonical quantization theory of gravity, Loop Quantum Gravity (LQG). LQG is a quantum theory of gravitation based on a geometric formulation, whose intention is to unify Quantum Mechanics and General Relativity, incorporating the matter of the Standard Model to the established framework for the case of pure quantum gravity, and use them directly on the action (17) in search of new MDO-gravity bonds that are potential for quantization.

V. ACKNOWLEDGEMENT

RdCL thank to Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001. SHP acknowledges financial support from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (No. 303583/2018-5 and 308469/2021-6).

Appendix A: Covariant derivatives commutation in MDO spinor

Using the definition of the spin connection (6) in the covariant derivative in equation (5), and performing the commutation of two covariant derivatives over MDO spinor λ , has been:

$$\begin{split} [\nabla_{\mu}, \nabla_{\nu}]\lambda &= \nabla_{\mu}\nabla_{\nu}\lambda - \nabla_{\nu}\nabla_{\mu}\lambda = \nabla_{\mu}\left(\partial_{\nu} - \frac{i}{4}\omega_{\nu}^{ab}\sigma_{ab}\right)\lambda - \nabla_{\nu}\left(\partial_{\mu} - \frac{i}{4}\omega_{\mu}^{ab}\sigma_{ab}\right)\lambda \\ &= \left[\partial_{\mu}\left(\partial_{\nu} - \frac{i}{4}\omega_{\nu}^{ab}\sigma_{ab}\right)\lambda - \Gamma_{\mu\nu}^{\rho}\left(\partial_{\rho} - \frac{i}{4}\omega_{\rho}^{ab}\sigma_{ab}\right)\lambda - \frac{i}{4}\omega_{\mu}^{ab}\sigma_{ab}(\partial_{\nu}\lambda) + \frac{i}{4}\omega_{\mu}^{ab}\sigma_{ab}\frac{i}{4}\omega_{\nu}^{cd}\sigma_{cd}\lambda\right] \\ &- \left[\partial_{\nu}\left(\partial_{\mu} - \frac{i}{4}\omega_{\mu}^{ab}\sigma_{ab}\right)\lambda - \Gamma_{\nu\mu}^{\rho}\left(\partial_{\rho} - \frac{i}{4}\omega_{\rho}^{ab}\sigma_{ab}\right)\lambda - \frac{i}{4}\omega_{\nu}^{ab}\sigma_{ab}(\partial_{\mu}\lambda) + \frac{i}{4}\omega_{\nu}^{cd}\sigma_{cd}\frac{i}{4}\omega_{\mu}^{ab}\sigma_{ab}\lambda\right], \quad (A1) \end{split}$$

being $(\partial_{\mu}\partial_{\nu} - \partial_{\nu}\partial_{\mu})\lambda = 0$ and assuming a torsion free scenario $(\Gamma^{\rho}_{\mu\nu} - \Gamma^{\rho}_{\nu\mu}) = 0$, we have

$$\begin{split} [\nabla_{\mu}, \nabla_{\nu}]\lambda &= -\frac{i}{4}\partial_{\mu}(\omega_{\nu}^{ab}\sigma_{ab}\lambda) - \frac{i}{4}\omega_{\mu}^{ab}\sigma_{ab}(\partial_{\nu}\lambda) + \frac{i}{4}\partial_{\nu}(\omega_{\mu}^{ab}\sigma_{ab}\lambda) + \frac{i}{4}\omega_{\nu}^{ab}\sigma_{ab}(\partial_{\mu}\lambda) \\ &+ \frac{i}{4}\frac{i}{4}(\omega_{\mu}^{ab}\sigma_{ab}\omega_{\nu}^{cd}\sigma_{cd} - \omega_{\nu}^{cd}\sigma_{cd}\omega_{\mu}^{ab}\sigma_{ab})\lambda \\ &= -\frac{i}{4}(\partial_{\mu}\omega_{\nu}^{ab})\sigma_{ab}\lambda - \frac{i}{4}\omega_{\nu}^{ab}\sigma_{ab}(\partial_{\mu}\lambda) - \frac{i}{4}\omega_{\mu}^{ab}\sigma_{ab}(\partial_{\nu}\lambda) + \frac{i}{4}(\partial_{\nu}\omega_{\mu}^{ab})\sigma_{ab}\lambda \\ &+ \frac{i}{4}\omega_{\mu}^{ab}\sigma_{ab}(\partial_{\nu}\lambda) + \frac{i}{4}\omega_{\nu}^{ab}\sigma_{ab}(\partial_{\mu}\lambda) + \frac{i}{4}\frac{i}{4}(\omega_{\mu}^{ab}\sigma_{ab}\omega_{\nu}^{cd}\sigma_{cd} - \omega_{\nu}^{cd}\sigma_{cd}\omega_{\mu}^{ab}\sigma_{ab})\lambda \\ &= -\frac{i}{4}(\partial_{\mu}\omega_{\nu}^{ab} - \partial_{\nu}\omega_{\mu}^{ab})\sigma_{ab}\lambda + \frac{i}{4}\frac{i}{4}\omega_{\mu}^{ab}\omega_{\nu}^{cd}(\sigma_{ab}\sigma_{cd} - \sigma_{cd}\sigma_{ab})\lambda. \end{split}$$
(A2)

Once $\sigma^{ab} = i/2[\gamma^a, \gamma^b]$ and $\{\gamma^a, \gamma^b\} = 2\eta^{ab}$, immediately we have $\gamma^b \gamma^a = 2\eta^{ab} - \gamma^a \gamma^b$ and $\sigma^{ab} = i/2[\gamma^a, \gamma^b] = i/2(\gamma^a \gamma^b - 2\eta^{ab} + \gamma^a \gamma^b) = i(\gamma^a \gamma^b - \eta^{ab})$. Using σ^{ab} in the last term of expression (A2);

$$[\nabla_{\mu}, \nabla_{\nu}]\lambda = -\frac{i}{4} (\partial_{\mu}\omega_{\nu}^{ab} - \partial_{\nu}\omega_{\mu}^{ab})\sigma_{ab}\lambda - \frac{i}{4} (\omega_{\mu}^{ac}\omega_{\nu c}{}^{b} - \omega_{\nu}^{ac}\omega_{\mu c}{}^{b})\sigma_{ab}\lambda$$
$$= -\frac{i}{4} [(\partial_{\mu}\omega_{\nu}^{ab} - \partial_{\nu}\omega_{\mu}^{ab}) + (\omega_{\mu}^{ac}\omega_{\nu c}{}^{b} - \omega_{\nu}^{ac}\omega_{\mu c}{}^{b})]\sigma_{ab}\lambda.$$
(A3)

Therefore, from equation (A3), it is possible to describe the commutation between the covariant derivatives on the spinor λ as

$$[\nabla_{\mu}, \nabla_{\nu}]\lambda = -\frac{i}{4}F_{\mu\nu}{}^{ab}\sigma_{ab}\lambda, \tag{A4}$$

where $F_{\mu\nu}{}^{ab} = \partial_{\mu}\omega_{\nu}{}^{ab} - \partial_{\nu}\omega_{\mu}{}^{ab} + \omega_{\mu}{}^{ac}\omega_{\nu c}{}^{b} - \omega_{\nu}{}^{ac}\omega_{\mu c}{}^{b}$.

Appendix B: From Einstein-Hilbert action to Palatini-Hilbert action with the term of Holst

It is known, from the General Relativity, that the Riemann tensor can be obtained using the commutation of two covariant derivatives acting on a vector, it means that

$$[\nabla_{\mu}, \nabla_{\nu}]V_{\rho} = R^{\alpha}_{\nu \ \mu\rho}V_{\alpha}, \quad R^{\alpha}_{\nu \ \mu\rho} = \partial_{\nu}\Gamma^{\alpha}_{\mu\rho} - \partial_{\mu}\Gamma^{\alpha}_{\nu\rho} + \Gamma^{\sigma}_{\rho\mu}\Gamma^{\alpha}_{\nu\sigma} - \Gamma^{\sigma}_{\rho\nu}\Gamma^{\alpha}_{\mu\sigma}. \tag{B1}$$

The relation between the curvature tensor $R_{\nu \ \mu\rho}^{\ \alpha}$ and $F_{\mu\nu}^{\ ab}$ originates from the commutation between the covariant derivatives under an internal index object (tetrad), $[\nabla_{\mu}, \nabla_{\nu}]S_a$:

$$[\nabla_{\mu}, \nabla_{\nu}]S_{a} = \nabla_{\mu}\nabla_{\nu}S_{a} - \nabla_{\nu}\nabla_{\mu}S_{a} = [\partial_{\mu}(\partial_{\nu}S_{a} + \omega_{\nu a}{}^{b}S_{b}) - \Gamma^{\alpha}_{\mu\nu}(\partial_{\alpha}S_{a} + \omega_{\alpha a}{}^{b}S_{b}) + \omega_{\mu a}{}^{c}(\partial_{\nu}S_{c} + \omega_{\nu c}{}^{b}S_{b})] - [\partial_{\nu}(\partial_{\mu}S_{a} + \omega_{\mu a}{}^{b}S_{b}) - \Gamma^{\alpha}_{\nu\mu}(\partial_{\alpha}S_{a} + \omega_{\alpha a}{}^{b}S_{b}) + \omega_{\nu a}{}^{c}(\partial_{\mu}S_{c} + \omega_{\mu c}{}^{b}S_{b})],$$
(B2)

again, because of the torsion free scenario $\Gamma^{\alpha}_{\mu\nu} - \Gamma^{\alpha}_{\nu\mu} = 0$, we get

$$[\nabla_{\mu}, \nabla_{\nu}]S_{a} = \partial_{\mu}\partial_{\nu}S_{a} + (\partial_{\mu}\omega_{\nu a}{}^{b})S_{b} + \omega_{\nu a}{}^{b}(\partial_{\mu}S_{b}) + \omega_{\mu a}{}^{c}(\partial_{\nu}S_{c}) + \omega_{\mu a}{}^{c}\omega_{\nu c}{}^{b}S_{b} - \partial_{\nu}\partial_{\mu}S_{a} - (\partial_{\nu}\omega_{\mu a}{}^{b})S_{b} - \omega_{\mu a}{}^{b}(\partial_{\nu}S_{b}) - \omega_{\nu a}{}^{c}(\partial_{\mu}S_{c}) - \omega_{\nu a}{}^{c}\omega_{\mu c}{}^{b}S_{b} = (\partial_{\mu}\omega_{\nu a}{}^{b})S_{b} - (\partial_{\nu}\omega_{\mu a}{}^{b})S_{b} + \omega_{\mu a}{}^{c}\omega_{\nu c}{}^{b}S_{b} - \omega_{\nu a}{}^{c}\omega_{\mu c}{}^{b}S_{b},$$
(B3)

and it allows us to concluded that

$$[\nabla_{\mu}, \nabla_{\nu}]S^a = F_{\mu\nu}{}^{ab}S_b, \tag{B4}$$

where $F_{\mu\nu}{}^{ab} = \partial_{\mu}\omega_{\nu}{}^{ab} - \partial_{\nu}\omega_{\mu}{}^{ab} + \omega_{\mu}{}^{ac}\omega_{\nu c}{}^{b} - \omega_{\nu}{}^{ac}\omega_{\mu c}{}^{b}$. Now, with the vector V_{α} in terms of tetrads and using the equation (B4), the expression (B1) can be rewriting as

$$R^{\,\alpha}_{\nu\ \mu\rho}V_{\alpha} = [\nabla_{\mu}, \nabla_{\nu}]V_{\rho} = [\nabla_{\mu}, \nabla_{\nu}]e^{a}_{\rho}V_{a} = e^{a}_{\rho}[\nabla_{\mu}, \nabla_{\nu}]V_{a} = e^{a}_{\rho}F_{\mu\nu a}^{\ b}V_{b} = e^{a}_{\rho}F_{\mu\nu a}^{\ b}e^{a}_{b}V_{\alpha}, \tag{B5}$$

then the relation between tensor curvature and $F_{\mu\nu}^{\ \ ab}$ takes the form:

$$R^{\ \alpha}_{\nu\ \mu\rho} = e^a_\rho F^{\ b}_{\mu\nu a} e^{\alpha}_b. \tag{B6}$$

From equation (B6):

$$R_{\nu\sigma\mu\rho} = g_{\sigma\alpha}R_{\nu\ \mu\rho}^{\ \alpha} = g_{\sigma\alpha}e_{\rho}^{a}F_{\mu\nu a}^{\ b}e_{b}^{\alpha} = \eta_{cd}e_{\sigma}^{c}e_{\alpha}^{d}e_{\rho}^{a}F_{\mu\nu a}^{\ b}e_{b}^{\alpha} = \eta_{cd}e_{\sigma}^{c}e_{\rho}^{a}F_{\mu\nu a}^{\ b}\delta_{b}^{d} = \eta_{cd}e_{\sigma}^{c}e_{\rho}^{a}F_{\mu\nu a}^{\ b}d = R_{\mu\rho\nu\sigma}.$$
 (B7)

Once the Ricci's scalar is $R = g^{\mu\nu}R_{\mu\nu} = R_{\mu\rho\nu\sigma}g^{\mu\nu}g^{\rho\sigma}$, from the equation (B7) it follows that

$$R = \eta_{cd} e^c_{\sigma} e^a_{\rho} F_{\mu\nu a}{}^d \eta^{gh} e^a_g e^\nu_h \eta^{jk} e^\rho_j e^\sigma_k = \eta_{cd} e^c_{\sigma} e^a_{\rho} F_{\mu\nu a}{}^d \eta^{ch} e^a_c e^\rho_h e^\sigma_a e^\sigma_k e^\sigma_k e^\sigma_k e^\sigma_k e^\sigma_k e^\sigma_h e$$

Making the change $k \to a$ and $d \to b$, the scalar of curvature, via tetrads and connection (e, ω) , is

$$R = F_{\mu\nu}{}^{ab}e^{\mu}_{a}e^{\nu}_{b}.\tag{B9}$$

The Jacobian term for volume correction in curved spaces, $\sqrt{-g}$, in terms of tetrads is characterized by

$$\sqrt{-g} = \left[-\det(\eta_{ab}e^a_{\mu}e^b_{\nu})\right]^{1/2} = \left[-\det(\eta_{ab})\det(e^a_{\mu})\det(e^b_{\nu})\right]^{1/2} = \left[\det(e^a_{\mu})\det(e^b_{\nu})\right]^{1/2} = (e^2)^{1/2} = e.$$
(B10)

Thus, Einstein-Hilbert action can be written with the Riemann-Cartan formalism, via equations (B9) and (B10), in the so-called Palatini-Hilbert action [24]:

$$\mathcal{S}_{EH} = \frac{1}{16\pi G} \int_{\mathcal{M}} d^4 x \sqrt{-g} R \longmapsto \mathcal{S}_{PH} = \frac{1}{16\pi G} \int_{\mathcal{M}} d^4 x(e) F_{\mu\nu}{}^{ab} e^{\mu}_a e^{\nu}_b. \tag{B11}$$

Almost 77 years after the Palatini's work, Holst obtained a dual term associated with curvature $F_{\mu\nu}{}^{ab}$ in the Hamiltonian formalism for gravity, now called the term of Holst (self-dual of Palatini action) [15]:

$$-\frac{1}{2\gamma}ee^{\mu}_{I}e^{\nu}_{J}F_{\mu\nu}{}^{IJ}(\omega), \tag{B12}$$

where (I, J) are the spatial components of Lorentz internal indices (a, b) and γ is the Barberp-Immirzi parameter, proposed by both authors in refs. [25, 26], in which the self-dual formulation corresponds to the choice of $\gamma = -i$. This Holst term allows the Palatini-Hilbert action to be generalized to

$$S_{PH} = \frac{1}{16\pi G} \int_{\mathcal{M}} d^4 x (e e_I^{\mu} e_J^{\nu} P^{IJ}{}_{KL} F_{\mu\nu}{}^{KL}(\omega)), \tag{B13}$$

being that, this value $\gamma = \pm i$ can not be performed to deal with its inverse:

$$P^{IJ}{}_{KL} = \delta^{[I}_{K} \delta^{J]}_{L} - \frac{1}{\gamma} \frac{\epsilon^{IJ}{}_{KL}}{2}, \quad P^{-1}{}_{IJ}{}^{KL} = \frac{\gamma^{2}}{\gamma^{2} + 1} \left(\delta^{[K}_{I} \delta^{L]}_{J} + \frac{1}{\gamma} \frac{\epsilon_{IJ}{}^{KL}}{2} \right). \tag{B14}$$

Appendix C: Palatini-Hilbert-Holst coupling MDO action in ADM and tetrad formalism

Knowing that $|det(e_a^i)| = \sqrt{q}$ and $|e| = |\sqrt{-g}| = N\sqrt{q}$, and using the decomposition (15), the gravitational Lagrangian density of action (11) is expressed as $ee_I^{\mu}e_J^{\nu}P^{IJ}{}_{KL}F_{\mu\nu}{}^{KL}(\omega) \mapsto N\sqrt{q}e_I^ae_J^bP^{IJ}{}_{KL}F_{ab}{}^{KL}(\omega)$:

$$N\sqrt{q}e_{I}^{a}e_{J}^{b}P^{IJ}{}_{KL}F_{ab}{}^{KL}(\omega) = N\sqrt{q}(\varepsilon_{I}^{a} - n^{a}n_{I})(\varepsilon_{J}^{b} - n^{b}n_{J})P^{IJ}{}_{KL}F_{ab}{}^{KL}(\omega)$$
$$= N\sqrt{q}(\varepsilon_{I}^{a}\varepsilon_{J}^{b} - \varepsilon_{I}^{a}n^{b}n_{J} - n^{a}n_{I}\varepsilon_{J}^{b} + n^{a}n_{I}n^{b}n_{J})P^{IJ}{}_{KL}F_{ab}{}^{KL}(\omega),$$
(C1)

where the last three terms inside parentheses are, from equation (16),

$$n^{a}n_{I}n^{b}n_{J} - \varepsilon_{I}^{a}n^{b}n_{J} - n^{a}n_{I}\varepsilon_{J}^{b} = (N^{-1}(t^{a} - N^{a}))n_{I}(N^{-1}(t^{b} - N^{b}))n_{J} - \varepsilon_{I}^{a}(N^{-1}(t^{b} - N^{b}))n_{J} - (N^{-1}(t^{a} - N^{a}))n_{I}\varepsilon_{J}^{b}$$

$$= N^{-2}(t^{a} - N^{a})(t^{b} - N^{b})(\delta_{I}^{0}\delta_{J}^{0}) - N^{-1}\varepsilon_{I}^{a}t^{b}n_{J} + N^{-1}\varepsilon_{I}^{a}N^{b}n_{J} - N^{-1}t^{a}n_{I}\varepsilon_{J}^{b} + N^{-1}N^{a}n_{I}\varepsilon_{J}^{b}$$

$$= -N^{-1}[\varepsilon_{I}^{a}t^{b}n_{J} + \varepsilon_{J}^{b}t^{a}n_{I}] + N^{-1}[N^{a}n_{I}\varepsilon_{J}^{b} + N^{b}n_{J}\varepsilon_{I}^{a}]$$

$$= -2N^{-1}n_{I}t^{a}\varepsilon_{J}^{b} + 2N^{-1}N^{a}n_{I}\varepsilon_{J}^{b}, \qquad (C2)$$

so that, using the equations (C2) in (C1), the gravitational Lagrangian density becomes

$$ee_I^{\mu}e_J^{\nu}P_{\ KL}^{IJ}F_{\mu\nu}^{\ KL}(\omega) \mapsto N\sqrt{q}(\varepsilon_I^a\varepsilon_J^b - 2N^{-1}n_It^a\varepsilon_J^b + 2N^{-1}N^an_I\varepsilon_J^b)P_{\ KL}^{IJ}F_{ab}^{\ KL}(\omega). \tag{C3}$$

Following the same prescription, the plots from the part to the "scalar" and fermionic of the MDO spinor field in action (11) gives

$$e[(e_I^{\mu}e_J^{\nu}\eta^{IJ}\partial_{\mu}\vec{\lambda}\partial_{\nu}\lambda) - m^2\vec{\lambda}\lambda] \mapsto N\sqrt{q}[(\varepsilon_I^a\varepsilon_J^b - 2N^{-1}n_It^a\varepsilon_J^b + 2N^{-1}N^an_I\varepsilon_J^b)\eta^{IJ}(\partial_a\vec{\lambda}\partial_b\lambda) - m^2\vec{\lambda}\lambda]$$
(C4)

and

$$ee_{I}^{\mu}e_{J}^{\nu}\eta^{IJ}[\vec{\lambda}\,\omega_{\mu}^{IJ}\sigma_{IJ}(\partial_{\nu}\lambda) - (\partial_{\mu}\,\vec{\lambda})\omega_{\nu}^{MN}\sigma_{MN}\lambda - i/4\,\vec{\lambda}\,(\omega_{\mu}^{IJ}\omega_{\nu}^{MN}\sigma_{IJ}\sigma_{MN})\lambda] \mapsto \\ N\sqrt{q}[\varepsilon_{I}^{a}\varepsilon_{J}^{b} - 2N^{-1}n_{I}t^{a}\varepsilon_{J}^{b} + 2N^{-1}N^{a}n_{I}\varepsilon_{J}^{b}]\eta^{IJ}\{i/2(\vec{\lambda}\,\omega_{a}^{IJ}[\gamma_{I}\gamma_{J} - \gamma_{J}\gamma_{I}](\partial_{b}\lambda) - (\partial_{a}\,\vec{\lambda})\omega_{b}^{MN}[\gamma_{M}\gamma_{N} - \gamma_{N}\gamma_{M}]\lambda) \\ + i/16\,\vec{\lambda}\,(\omega_{a}^{IJ}\omega_{b}^{MN}[\gamma_{I}\gamma_{J} - \gamma_{J}\gamma_{I}][\gamma_{M}\gamma_{N} - \gamma_{N}\gamma_{M}])\lambda\}.$$
(C5)

Thus, from the expressions (C3), (C4) and (C5), the action of MDO matter with the Palatini-Holst gravitation (11) is described in ADM formalism and tetrad field as

$$\begin{split} \mathcal{S}[\varepsilon,\omega,\lambda] &= \mathcal{S}_{PH,\xi} + \mathcal{S}_{\lambda,\vec{\lambda}_{S}} + \mathcal{S}_{\lambda,\vec{\lambda}_{F}} \\ &= \frac{1}{16\pi G} \int_{\mathbb{R}} dt \int_{\Sigma} d^{3}x N \sqrt{q} \Omega_{IJ}^{ab} P^{IJ}{}_{KL} F_{ab} {}^{KL}(\omega) (1 - 8\pi G\xi \,\vec{\lambda} \,\lambda) \\ &\quad + \frac{1}{2} \int_{\mathbb{R}} dt \int_{\Sigma} d^{3}x N \sqrt{q} [\Omega_{IJ}^{ab} \eta^{IJ} (\partial_{a} \,\vec{\lambda} \,\partial_{b} \lambda) - m^{2} \,\vec{\lambda} \,\lambda] \\ &\quad - \frac{1}{8} \int_{\mathbb{R}} dt \int_{\Sigma} d^{3}x N \sqrt{q} \Omega_{IJ}^{ab} \eta^{IJ} \{i/2(\vec{\lambda} \,\omega_{a}^{IJ}[\gamma_{I},\gamma_{J}](\partial_{b} \lambda) \\ &\quad - (\partial_{a} \,\vec{\lambda}) \omega_{b}^{MN}[\gamma_{M},\gamma_{N}] \lambda) + i/16 \,\vec{\lambda} \,(\omega_{a}^{IJ} \omega_{b}^{MN}[\gamma_{I},\gamma_{J}][\gamma_{M},\gamma_{N}]) \lambda \}, \end{split}$$
(C6)

where $\Omega_{IJ}^{ab} = (\varepsilon_I^a \varepsilon_J^b - 2N^{-1}n_I t^a \varepsilon_J^b + 2N^{-1}N^a n_I \varepsilon_J^b)$. Being $S_{\lambda, \overline{\lambda}_S}$ and $S_{\lambda, \overline{\lambda}_F}$ the scalar and fermionic parts of the action, respectively.

- [1] D. V. Ahluwalia and D. Grumiller, JCAP **0507**, 012 (2005) doi:10.1088/1475-7516/2005/07/012.
- [2] A. G. Cohen and S. L. Glashow, Phys. Rev. Lett. 97 (2006), 021601 doi:10.1103/PhysRevLett.97.021601.
- [3] D. V. Ahluwalia, Adv. Appl. Clifford Algebras 27, no. 3, 2247 (2017) doi:10.1007/s00006-017-0775-1.
- [4] D. Ahluwalia, Eur. Phys. J. ST 229 (2020), 1997-2001 doi:10.1017/9781316145593
- [5] L. Duarte, M. Dias and F. de Campos, Eur. Phys. J. ST 229 (2020) no.11, 2133-2146 doi:10.1140/epjst/e2020-900267-5
- [6] S. H. Pereira, R. d. Lima, M. E. S. Alves, T. M. Guimarães, J. F. Jesus and A. P. S. Souza, Eur. Phys. J. ST 229 (2020) no.11, 2079-2116 doi:10.1140/epjst/e2020-900271-x
- [7] D. V. Ahluwalia, J. M. H. da Silva, C. Y. Lee, Y. X. Liu, S. H. Pereira and M. M. Sorkhi, Phys. Rept. 967 (2022), 1-43 doi:10.1016/j.physrep.2022.04.003
- [8] C. H. Coronado Villalobos and R. J. Bueno Rogerio, EPL 116 (2016) no.6, 60007 doi:10.1209/0295-5075/116/60007
- [9] S. H. Pereira and R. C. Lima, Int. J. Mod. Phys. D 26, no. 12, 1730028 (2017) doi:10.1142/S0218271817300282.
- [10] R. J. Bueno Rogerio, R. de C. Lima, L. Duarte, J. M. Hoff da Silva, M. Dias and C. R. Senise, EPL 128 (2019) no.2, 20004 doi:10.1209/0295-5075/128/20004
- [11] C. G. Boehmer, J. Burnett, D. F. Mota and D. J. Shaw, JHEP 1007, 053 (2010) doi:10.1007/JHEP07(2010)053.
- [12] M. Bojowald and R. Das, Phys. Rev. D 78, 064009 (2008) doi:10.1103/PhysRevD.78.064009.
- [13] Feynman, Richard P. (1995). Feynman Lectures on Gravitation. Addison-Wesley. p. 136, eq. (10.1.2).
- [14] J. M. Hoff da Silva and S. H. Pereira, JCAP 1403, 009 (2014) doi:10.1088/1475-7516/2014/03/009.
- [15] S. Holst, Phys. Rev. D 53, 5966 (1996) doi:10.1103/PhysRevD.53.5966.
- [16] P. J. E. Peebles and B. Ratra, Astrophys. J. Lett. **325** (1988), L17 doi:10.1086/185100.
- [17] L. Xu, J. Lu and W. Li, Phys. Lett. B 690 (2010), 333-336 doi:10.1016/j.physletb.2010.05.058
- [18] A. H. A. Alfedeel and A. Abebe, Int. J. Geom. Meth. Mod. Phys. 17 (2020) no.6, 2050076 doi:10.1142/S0219887820500760
- [19] P. A. R. Ade et al. [Planck Collaboration], Astron. Astrophys. 594, A13 (2016) doi:10.1051/0004-6361/201525830.
- [20] C. Brans and R. H. Dicke, Mach's principle and a relativistic theory of gravitation, Physical review, 124, 3, APS, 1961.
- [21] P. A. P. Molinari, F. P. Devecchi and M. O. Ribas, Phys. Rev. D 105 (2022) no.4, 043527 doi:10.1103/PhysRevD.105.043527
- [22] V. S. M. and S. Y., New Astron. 95 (2022), 101823 doi:10.1016/j.newast.2022.101823
- [23] R. L. Arnowitt, S. Deser and C. W. Misner, Phys. Rev. 116, 1322 (1959), doi:10.1103/PhysRev.116.1322.
- [24] A. Palatini (1919) Deduzione invariantiva delle equazioni gravitazionali dal principio di Hamilton, Rend. Circ. Mat. Palermo 43, 203-212 [English translation by R.Hojman and C. Mukku in P.G. Bergmann and V. De Sabbata (eds.) Cosmology and Gravitation, Plenum Press, New York (1980)].
- [25] J. F. Barbero G., Phys. Rev. D 51, 5507 (1995) doi:10.1103/PhysRevD.51.5507.
- [26] G. Immirzi, Class. Quant. Grav. 14, L177 (1997) doi:10.1088/0264-9381/14/10/002.
- [27] C. Kiefer, Quantum gravity, vol. 136 of International Series of Monographs on Physics (2007).
- [28] S. J. Gabriele Gionti and S. J, Phys. Rev. D 103 (2021) no.2, 024022 doi:10.1103/PhysRevD.103.024022.
- [29] A. Ashtekar, Phys. Rev. D 36 (1987), 1587-1602 doi:10.1103/PhysRevD.36.1587.