

Memory effects from holonomies

Ali Seraj and Turmoli Neogi

Centre for Gravitational Waves, Université Libre de Bruxelles,
International Solvay Institutes, CP 231, B-1050 Brussels, Belgium.

Abstract. We show that known electromagnetic and gravitational memory effects can be formulated in terms of certain holonomies over suitable paths on a holographic screen in asymptotically flat spacetimes. In particular, the leading and subleading displacement memory effects correspond to ‘translational’ holonomies, while the gyroscopic and spin memory effects are derived from ‘rotational’ holonomies. Moreover, electromagnetic and gravitational kick memory effects correspond to spatial derivatives of the translational holonomy. The gravitational holonomy solves the parallel or ‘affine’ transport equations, depending on an invariant of the Poincaré representation.

Contents

1	Introduction and summary of results	1
2	EM Memory Effect	3
2.1	Holonomies	5
3	Gravitational Memory effects	6
3.1	Displacement and kick memory	7
3.2	Spin memory	8
3.3	Gyroscopic memory	8
4	Gravitational memory from holonomy	9
4.1	Formal development	9
4.2	Asymptotic frame	11
4.3	Translational holonomy	12
4.4	Rotational holonomy	14
5	Discussion and outlook	15
	References	16

1 Introduction and summary of results

Gravitational wave (GW) memory refers to permanent effects in the configuration of an experimental setup after the passage of GWs. Its classical example is the change in the

physical distance between free test masses initially at rest which arises due to a permanent change in transverse traceless components of the metric after the passage of the GW burst. This is the well known ‘displacement memory’ whose linear form dates back to the 70’s [1,2]. A more rigorous analysis of this effect in the 90’s by Christodoulou [3] and Blanchet and Damour [4] showed that there is a nonlinear contribution to the memory. The latter can be understood as the linear memory caused by emitted gravitons [5], *i.e.* the ability of gravitons to gravitate [6]. While the linear effect is dominant in scattering processes, nonlinear memory has the major contribution in the displacement memory from bound binary systems. In the perturbative coalescence regime, the amplitude of the memory is about 20 percent of the amplitude of the peak of the oscillatory waveform [7]. However, the low frequency nature of the memory, makes it hard to detect in GW experiments. Prospects for detection of GW memory is discussed in [6, 8–11]. Thanks to the BMS flux-balance equations [12], memory effects can be implemented in numerical waveforms, implying that the effect can be enhanced in the merger phase [13–15].

In recent years, new types of permanent GW effects have been discovered that are not simply related to the net change in the metric components, but are rather sourced by certain time integrals of the waveform. This includes the spin and center of mass memory effects [16–18], and more recently the gyroscopic memory effect [19,20]. Indeed, the set of positive integer modes in the Mellin transform of the Bondi news lead to a tower of subleading memory effects [21,22], which can be detected by studying curve deviation [23]. While all these effects are subdominant with respect to the standard displacement memory effect, and thereby hard to detect in the near future, they are still very interesting due to their relation to the symmetry structures of asymptotically flat spacetimes [16, 24–28], leading to the interpretation of memory effects as a ‘vacuum transition’ [24, 29, 30].

Memory effects show up also in gauge theories [31–33], though to a more limited extent. The passage of electromagnetic radiation leads to a net shift in transverse components of the gauge field A_μ . Interestingly, the physical effect of this shift on a test charges is not a displacement, but a kick. Our treatment of EM memory effect in terms of Wilson loops sheds some light on why this should be the case. Finally, it should be mentioned that there are novel memory effects in modified theories of gravity [34–36] with interesting symmetry interpretation [37].

The memory effect is a nonlocal effect in both space and time. Therefore, one might expect that nonlocal quantities like holonomies and Wilson loops might be good candidates to quantify memory effects. Progress along this direction was made in [38,39] where a ‘generalized’ gravitational holonomy was introduced through a modification of the parallel transport equation called the ‘affine’ transport equation. This holonomy computed over a specific closed path was shown to encode various gravitational memory effects.

In this paper, we develop further this interesting idea. We make the relationship between memory and holonomy more direct by introducing different paths, each of which giving a certain type of memory. Moreover, we find that things are more transparent in the first order formulation of gravity. In particular, we show that the affine transport equation corresponds to gauging the Poincaré algebra, while the parallel transport equation corresponds to gauging the Lorentz algebra. In the former case, the holonomy naturally splits into a ‘translational’ part and a ‘rotational’ part. We will show that the translational holonomy reveals the leading and subleading displacement memory ef-

fects, while the rotational holonomy reproduces gyroscopic and spin memory effects. Finally, electromagnetic and gravitational kick memory effects are obtained by deriving the holonomy with respect to the spatial position of the loop.

In AdS/CFT, Wilson loops in the field theory side correspond to the area of certain minimal surfaces in the bulk, and the phase transition occurs when the minimal surface starts to meet the black hole horizon [40]. It is then natural to consider Wilson loops/holonomies in asymptotically flat spacetimes. Motivated by this, we consider holonomies that lie on an asymptotic hypersurface of constant radius, viewed as a holographic screen. It would be interesting to see if there is an alternative way to compute these holonomies as surface integrals in the bulk. The nonabelian Stokes theorem [41] which generalizes the usual Stokes theorem to Lie algebra valued differential one-forms may be useful in this regard.

This paper is organized as follows. In section 2, we study electromagnetic kick memory effect and show that it can be derived from a natural Wilson loop/holonomy. Moreover, we argue that there is another natural Wilson loop that captures a subleading memory, which corresponds to the Aharonov-Bohm phase on the same loop. Then we turn to general relativity in section 3, and review known gravitational memory effects. In section 4, after formal development of the gravitational holonomy and the corresponding transport equations in section 4.1, we introduce a suitable frame to compute gravitational holonomies in section 4.2. We then compute gravitational holonomies and show that they encode memory effects discussed in section 3. In particular, the leading and subleading displacement memory effects are derived from the translational holonomy in section 4.3, while the gyroscopic and spin memory effects are obtained from the rotational holonomy in section 4.4, either on a timelike geodesic or on a helical path in spacetime. We end with discussion and outlook in section 5.

2 EM Memory Effect

Consider a test particle with mass m and charge q , in Minkowski space, subject to electromagnetic radiation produced by some source J^μ at a large distance r . We assume that the source is centered around the origin $r = 0$ of a retarded coordinate system (u, r, θ^A) in which the metric reads

$$ds^2 = -du^2 - 2dudr + r^2\gamma_{AB}d\theta^Ad\theta^B. \quad (2.1)$$

In these coordinates, Maxwell equations $\partial_\mu F^{\mu\nu} = J^\nu$ imply that the gauge field behaves at large distance as

$$A = \frac{\bar{A}_u}{r}du + \frac{\bar{A}_r}{r^2}dr + \bar{A}_Ad\theta^A + \text{subleading}. \quad (2.2)$$

Note that coordinate basis vectors ∂_A are not normalized, since $\partial_A \cdot \partial_B = r^2\gamma_{AB}$. From the viewpoint of an experimenter, it is more natural to use instead an orthonormal tetrad, *i.e.* a set of four basis vectors $e_{\hat{\mu}} = e_{\hat{\mu}}^\mu \partial_\mu$ such that $e_{\hat{\mu}} \cdot e_{\hat{\nu}} = \eta_{\hat{\mu}\hat{\nu}}$ where $\eta_{\hat{\mu}\hat{\nu}} = \text{diag}(-1, 1, 1, 1)$. The dual basis one forms are given by $e^{\hat{\mu}} = e^{\hat{\mu}}_\mu dx^\mu$. In this frame, components of any tensor is given by contraction with suitable number of the tetrad $e_{\hat{\mu}}^\mu$ or its inverse $e^{\hat{\mu}}_\mu$. For example, $A_{\hat{\mu}} = A_\mu e_{\hat{\mu}}^\mu$ or $F^{\hat{\mu}\hat{\nu}} = e^{\hat{\mu}}_\mu e^{\hat{\nu}}_\nu F^{\mu\nu}$. A suitable frame in Minkowski spacetime

is

$$\mathbf{e}_{\hat{0}} = \partial_u, \quad \mathbf{e}_{\hat{r}} = \partial_r - \partial_u, \quad \mathbf{e}_{\hat{A}} = \frac{1}{r} E_{\hat{A}}^A \partial_A, \quad (2.3)$$

$$\mathbf{e}^{\hat{0}} = du + dr, \quad \mathbf{e}^{\hat{r}} = dr, \quad \mathbf{e}^{\hat{A}} = r E^{\hat{A}}_A d\theta^A. \quad (2.4)$$

where $\mathbf{e}_{\hat{0}}$ is adapted to observers at rest, $\mathbf{e}_{\hat{r}}$ is along the radial direction specified by outgoing rays, and $E_{\hat{A}}^A$ form a dyad on the round sphere $\gamma_{AB} E_{\hat{A}}^A E_{\hat{B}}^B = \delta_{\hat{A}\hat{B}}$. In this basis, we have

$$A = \frac{\bar{A}_{\hat{0}}}{r} \mathbf{e}^{\hat{0}} + \frac{\bar{A}_{\hat{r}}}{r^2} \mathbf{e}^{\hat{r}} + \frac{\bar{A}_{\hat{A}}}{r} \mathbf{e}^{\hat{A}} + \text{subleading}. \quad (2.5)$$

where the leading components relate to those in the coordinate basis as

$$\bar{A}_{\hat{0}} = \bar{A}_u = -\bar{A}_{\hat{r}}, \quad \bar{A}_{\hat{A}} = E_{\hat{A}}^A \bar{A}_A \quad (2.6)$$

Also, up to subleading corrections, components of the field strength are given by

$$F_{\hat{0}\hat{A}} = -F_{\hat{r}\hat{A}} = \frac{1}{r} E_{\hat{A}}^A \dot{\bar{A}}_A \quad (2.7)$$

$$F_{\hat{0}\hat{r}} = \frac{1}{r^2} (\bar{A}_0 + \dot{\bar{A}}_r), \quad F_{\hat{A}\hat{B}} = \frac{1}{r^2} E_{\hat{A}}^A E_{\hat{B}}^B (\partial_A \bar{A}_B - \partial_B \bar{A}_A) \quad (2.8)$$

Note that here $\bar{A}_A(u, x^B)$ is a tensor living on the unit sphere and therefore it is contracted with the dyad on the sphere $E_{\hat{A}}^A$. This will be frequently used in the paper, following the Bondi framework in which one performs an asymptotic expansion in $1/r$ to reduce dynamical fields to covariant tensors living on the unit sphere, called the ‘celestial’ sphere.

Consider a test charged particle far from the source. Its dynamics is given by

$$\frac{d}{d\tau} V^{\hat{\mu}} = \frac{q}{m} F^{\hat{\mu}}_{\hat{\nu}} V^{\hat{\nu}} \quad (2.9)$$

where τ is the proper time on the particle’s worldline and $V^{\hat{\mu}}$ is the four velocity in the local frame. To solve this, we perform an asymptotic expansion for the velocity as $V^{\hat{\mu}} = \bar{V}^{\hat{\mu}} + \frac{v^{\hat{\mu}}}{r} + \dots$. Inserting this expansion in (2.9), we find an equation at each power of $1/r$. The first equation implies that $\bar{V}^{\hat{\mu}} = \text{const}$. Let us further assume that the charge has zero initial velocity in the local frame, *i.e.* $\bar{\mathbf{V}} = \mathbf{e}_{\hat{0}}$, which is thus preserved by the evolution. Now radiative effects appear at $O(1/r)$, where we find that the temporal and radial components of $v^{\hat{\mu}}$ are vanishing, while

$$\dot{v}_{\hat{A}} = -\frac{q}{m} E_{\hat{A}}^A \dot{\bar{A}}_A \quad (2.10)$$

which gives the electromagnetic kick memory upon integration over time:

$$\Delta V_{\hat{A}} = -\frac{q}{mr} E_{\hat{A}}^A \Delta \bar{A}_A \quad (2.11)$$

The kick memory is given by the nonlocal (in time) quantity $\int du \dot{A}_A = \Delta \bar{A}_A$, is the soft photon current [42], which is the canonical pair of large gauge transformations. These two properties underlie the correspondence between memory effects, soft theorems and asymptotic symmetries [42].




Figure 1: Two different loops used for computing holonomies. The first one Γ_{\square} is a spacetime rectangular loop. The second one Γ_{\circ} is a spatial loop in transverse directions, *i.e.* in the plane tangent to the asymptotic sphere.

2.1 Holonomies

We will show that various memory effects can be derived from another nonlocal quantity, namely *holonomy*, see *e.g.* [43] for a nice exposition. Consider the parallel transport equation over a path Γ parameterized by $\tau \in [\tau_0, \tau_f]$:

$$\frac{D}{D\tau}W \equiv \frac{dx^\mu}{d\tau}D_\mu W = 0, \quad D_\mu = \partial_\mu + qA_\mu \quad (2.12)$$

where the connection $A_\mu = A_\mu^a T_a$ gauges the Lie algebra represented by generators T_a , and q is the charge of the test particle coupled to the gauge field. For a given path Γ (not necessarily closed), the holonomy $W(\Gamma)$, as an element of the gauge group, is a solution to (2.12) with initial condition $W[x(\tau_0)] = I$:

$$W[\Gamma] = \mathcal{P} \exp \left(-q \int_{\Gamma} dx^\mu A_\mu \right) \quad (2.13)$$

where \mathcal{P} is the path ordering operator. Note that we use natural units where $\hbar = c = G = 1$. The trace of this operator is called a Wilson line, or a Wilson loop if the path is closed. In Abelian gauge theories like Maxwell, the trace is trivial and thus the two notions coincide.

The fact that the radial component of the gauge field is subleading at large distance, see Eq.(2.5), implies that nontrivial holonomies at leading order are computed on a $r = \text{const}$ hypersurface. We can therefore consider two natural loops that are depicted in figure (1). The first loop Γ_{\square} , which we call ‘spacetime loop’, consists of a spatial segment of length L along a transverse vector $\vec{m} = m^{\hat{A}} e_{\hat{A}}$ and a temporal segment between u_0 to u_f . The other path Γ_{\circ} which we call the “spatial loop” is a closed path in the transverse directions (tangent to the sphere). In both cases, we assume that the spatial extent L of the loops are finite which means that $L/r \ll 1$.

EM Kick memory from the spacetime loop. The path ordered exponential (2.13) along the spacetime loop can be Taylor expanded. Since we are interested here in the leading $1/r$ effect, we can truncate quadratic and higher terms. Moreover, the leading

contribution of the two integrals along time direction are canceled and they lead to $O(1/r^2)$ effect. Therefore, the loop integral reduces to the difference of two spacelike line integrals $\int_0^x dx^A A_A(u)$ at $u = u_0$ and $u = u_f$, leading to

$$W = 1 - \frac{q}{r} \int_0^x dx^A \Delta \bar{A}_A \quad (2.14)$$

Therefore, the kick memory is given by

$$\boxed{\Delta P_{\hat{A}} = \partial_{\hat{A}} W} \quad (2.15)$$

where $P_{\hat{\mu}} = mV_{\hat{\mu}}$ is the momentum of the particle and $\partial_{\hat{A}} \equiv E_{\hat{A}}^A \partial_A$ following the convention explained below Eq.(2.7). By restoring units we get $\Delta P_{\hat{A}} = -\hbar \partial_{\hat{A}} W$, in which the operator $\hbar \partial_{\hat{A}}$ is a representation of momentum, and thus it is natural that the memory effect is a kick in electrodynamics, in contrast to gravity, where there is a displacement memory. This is because here the holonomy has no index, and to obtain a vectorial object, one needs to take a derivative.

Aharanov-Bohm phase from spatial loop. Now we argue that the Wilson loop over the other path $\Gamma_{\circlearrowleft}$ reveals a well known measurable quantity, namely the Aharanov-Bohm (AB) effect

$$W = P \exp \left(- \int_{\Gamma_{\circlearrowleft}} dx^A \bar{A}_A \right) \quad (2.16)$$

This quantity gives the relative phase caused by EM field on two similar particles starting at the same point, follow different paths and interfere at another point, thus forming a closed loop as in figure (1). This shows that the AB effect can be thought of as another EM memory effect. See Ref. [44] for further insight on this point.

3 Gravitational Memory effects

One can consistently incorporate gravitational effects, including gravitational radiation, in the Minkowski metric (2.1). This is conveniently done in Bondi gauge [45, 46]

$$g_{rr} = g_{rA} = 0, \quad \partial_r \det(r^{-2} g_{AB}) = 0, \quad (3.1)$$

in which any metric can be written as [47, 48]

$$ds^2 = -e^{2\beta}(F du^2 + 2du dr) + r^2 h_{AB} \left(d\theta^A - \frac{U^A}{r^2} du \right) \left(d\theta^B - \frac{U^B}{r^2} du \right) \quad (3.2)$$

where F, β, U^A, h_{AB} are functions of all coordinates. To understand the structure of radiative spacetimes, one solves Einstein equations far from the source, treating $1/r$ as a small parameter and imposing as a boundary condition that the metric tends to the Minkowski metric in the limit $r \rightarrow \infty$, *i.e.* $\lim_{r \rightarrow \infty} h_{AB} = \gamma_{AB}$, where γ_{AB} is the round metric on the unit sphere, usually called the ‘celestial sphere’. At leading orders, one finds the following asymptotic expansion

$$ds^2 = - \left(1 - \frac{2m}{r} \right) du^2 - 2du dr + \left(r^2 \gamma_{AB} + r C_{AB} \right) d\theta^A d\theta^B + D^B C_{AB} du d\theta^A. \quad (3.3)$$

The symmetric trace-free tensor $C_{AB}(u, \theta^A)$, called the Bondi shear, is the shear of the congruence of outgoing null rays and encodes the gravitational waveform received at retarded time u and angles θ^A on the celestial sphere. On the other hand, $m(u, \theta^A)$ is the Bondi mass aspect whose flux is fixed by the radiation

$$\partial_u m = \frac{1}{4} D_A D_B \dot{C}^{AB} - \frac{1}{8} \dot{C}_{AB} \dot{C}^{AB} \quad (3.4)$$

Note that γ_{AB} and its inverse γ^{AB} are used to lower or raise indices of tensors living on the celestial sphere.

3.1 Displacement and kick memory

The geodesic deviation equation, describing the relative separation X^μ of two nearby freely falling test masses, moving on a geodesic congruence with velocity V^μ in gravitational field reads

$$\ddot{X}^\mu = V^\alpha V^\beta R^\mu{}_{\alpha\beta\nu} X^\nu. \quad (3.5)$$

If one test mass is considered at the origin of the asymptotically normalized frame (2.3), then $X^{\hat{\mu}}$ describes the position of the second test mass. It turns out that the most dominant component of the Riemann tensor for the metric (3.3) is $R_{0\hat{A}0\hat{B}} = -\frac{1}{2r} \ddot{C}_{\hat{A}\hat{B}}$, where $C_{\hat{A}\hat{B}} = E_{\hat{A}}{}^A E_{\hat{B}}{}^B C_{AB}$ which implies that the leading geodesic deviation effect occurs in the transverse plane and is given by

$$\ddot{X}_{\hat{A}} = \frac{1}{2r} \ddot{C}_{\hat{A}\hat{B}} X^{\hat{B}}. \quad (3.6)$$

Upon integration over time twice, we find

$$X_{\hat{A}} = X_{\hat{A}}(0) + V_{\hat{A}}(0)(u - u_0) + \frac{1}{2r} \int^u \int^{u'} du'' \ddot{C}_{\hat{A}\hat{B}}(u'', X^A(u'')) X^{\hat{B}}(u''). \quad (3.7)$$

Let us assume that the spacetime is radiative in the interval $u \in (u_0, u_f)$ and consider a free test mass with initial position $X_0^{\hat{A}} \equiv X^{\hat{A}}(u_0)$ and velocity $V_0^{\hat{A}} \equiv V^{\hat{A}}(u_0)$ in the asymptotic frame (2.3). In the absence of radiation, $\dot{C}_{AB} = 0$ and accordingly the test mass follows a straight line up to subleading corrections at $O(1/r^2)$ which includes the Newtonian acceleration. In particular, the final motion at $u \geq u_f$, can be parameterized as

$$X^{\hat{A}}(u) = \tilde{X}^{\hat{A}}(u_0) + (u - u_0) \tilde{V}^{\hat{A}}(u_0) \quad u \geq u_f, \quad (3.8)$$

where the parameters

$$\tilde{X}^{\hat{A}}(u_0) = X_0^{\hat{A}} + d^{\hat{A}}/r, \quad \tilde{V}^{\hat{A}}(u_0) = V_0^{\hat{A}} + v^{\hat{A}}/r, \quad (3.9)$$

differ from the initial position and velocity by a displacement memory $d_{\hat{A}}/r$ and a kick memory $v_{\hat{A}}/r$, given by [34, 37, 39]¹

$$d_{\hat{A}} = \frac{1}{2} X_0^{\hat{B}} \Delta C_{\hat{A}\hat{B}} - V_0^{\hat{B}} \int_{u_0}^{u_f} du (C_{\hat{A}\hat{B}}(u) - C_{\hat{A}\hat{B}}(u_f)), \quad (3.10a)$$

$$v_{\hat{A}} = -\frac{1}{2} V_0^{\hat{B}} \Delta C_{\hat{A}\hat{B}}. \quad (3.10b)$$

¹Note that $d_{\hat{A}}$ has dimension length squared, so that $\frac{d_{\hat{A}}}{r}$ is a displacement. We factored out $1/r$ in (3.9) to stress the size of memory effects. Similar comment applies to $v_{\hat{A}}$.

The first term in Eq.(3.10a) is the standard result of Christodoulou [3], while the second term is a ‘subleading’ correction in the presence of initial relative velocity. The standard displacement effect, *i.e.* the first term in (3.10a) is sourced by $\Delta C_{\hat{A}\hat{B}}$, which is the soft graviton current, while the subleading displacement memory is sourced by $\int_{u_0}^{u_f} du (C_{\hat{A}\hat{B}}(u) - C_{\hat{A}\hat{B}}(u_f))$, which is the subleading soft graviton current [49] (up to a corner contribution). Finally Eq. (3.10b) is a kick memory analogous to that in Maxwell theory (2.11) and is again sourced by the leading soft graviton current.

3.2 Spin memory

The spin memory effect was introduced in [16]. It is indeed the effect of gravitational waves on a Sagnac interferometer, consisting of two light beams counter orbiting a closed loop Γ , constructed by optical fiber or mirrors. Taking the loop to lie on the sphere as in Γ_{\circ} in figure (1), the spin memory is given by the difference ΔT in the travel time T of the two light beams. In the approximation where T is much smaller than the period of the gravitational wave, and assuming the validity of geometric optics, the spin memory is given by [16]

$$\Delta T = \int_{\Gamma_{\circ}} d\theta^A D^B C_{AB} = \int_{D(\Gamma_{\circ})} d^2\theta \sqrt{\gamma} D_A D_B \tilde{C}^{AB}, \quad (3.11)$$

where $D(\Gamma_{\circ})$ is the area enclosed by the loop Γ_{\circ} and the coefficient $\tilde{C}_{AB} \equiv \epsilon_{AC} C_B^C$ is the dual shear [50].

3.3 Gyroscopic memory

The effect of gravitational field on gyroscopes, *i.e.* objects carrying spin, has been of interest for the whole history of general relativity. The famous Lens-Thirring effect explains how a rotating star/black hole affects the spin of the gyroscope, and this has been tested by the Gravity Probe B experiment [51]. The effect of gravitational waves on gyroscopes has been discussed in [52–54], and more thoroughly recently in [19, 20] for freely falling gyroscopes.

The evolution of a freely falling small gyroscope with velocity V^μ and spin S^μ is given by the parallel transport equation $V^\alpha \nabla_\alpha S^\mu = 0$. With respect to a local frame which is comoving with the gyroscope, *i.e.* when $e_{\hat{0}}^\mu = V^\mu$, the components of the gyroscope’s spin are purely spatial given by $S^{\hat{i}} = S^\mu e_{\hat{i}\mu}$, while $S^{\hat{0}} = -S^\mu u_\mu = 0$. The parallel transport equation then reads

$$\frac{dS^{\hat{i}}}{d\tau} = \Omega^{\hat{i}\hat{j}} S^{\hat{j}}, \quad \Omega^{\hat{i}\hat{j}} \equiv -V^\alpha \omega^{\hat{i}\hat{j}}{}_\alpha, \quad (3.12)$$

where $\omega^{\hat{\mu}\hat{\nu}}{}_\alpha = e^{\hat{\mu}}{}_\beta \nabla_\alpha e^{\hat{\nu}\beta}$ is the spin connection of the local frame with respect to which the orientation of the spin is computed. It was shown in [19, 20] that the effect of GWs on a gyroscope with initial spin $S^{\hat{A}}(u_0)$ is a net change of orientation, given by a rotation $\Delta\Phi$ in the transverse plane orthogonal to the direction of propagation of GWs²

$$\Delta S^{\hat{A}} = \epsilon^{\hat{A}\hat{B}} S_{\hat{B}}(u_0) \Delta\Phi, \quad \Delta\Phi = -\frac{1}{r^2} \int du \left(\frac{1}{4} D_A D_B \tilde{C}^{AB} - \frac{1}{8} N_{AB} \tilde{C}^{AB} \right) \quad (3.13)$$

²Our convention in defining the dual shear is opposite of that of [19, 20] and matches with [50, 55], which causes a relative minus sign in the angle $\Delta\Phi$.

The first term in the integrand also appears in the spin memory Eq.(3.11), which shows the close connection between the two effects. However, there is an additional nonlinear contribution to the gyroscopic memory, which is related to the charge of gravitational electric-magnetic duality, as shown in [19, 20].

4 Gravitational memory from holonomy

4.1 Formal development

In this section, we derive the above gravitational memory effects from certain gravitational holonomies. To this end, it is convenient to think of general relativity as a gauge theory of the Poincaré group (see [56] and references therein). Let us define the ‘Poincaré gauge field’ as

$$A_\mu = e^{\hat{\mu}}{}_\nu P_{\hat{\mu}} + \frac{1}{2} \omega^{\hat{\mu}\hat{\nu}}{}_\mu M_{\hat{\mu}\hat{\nu}} \quad (4.1)$$

where $P_{\hat{\mu}}, M_{\hat{\mu}\hat{\nu}}$ respectively denote the generators of Poincaré translations and Lorentz transformations, which are respectively gauged by the vielbein $e^{\hat{\mu}}{}_\mu$ and the spin connection $\omega^{\hat{\mu}\hat{\nu}}{}_\mu$. The explicit form of $P_{\hat{\mu}}, M_{\hat{\mu}\hat{\nu}}$ depends on the representation, *i.e.* the probe which is coupled to gravity. From (4.1), we define the ‘Poincaré holonomy’ as

$$W[\Gamma] = \mathcal{P} \exp \left(- \int_\Gamma dx^\mu A_\mu \right) = \mathcal{P} \exp \left(W^{\hat{\mu}} P_{\hat{\mu}} + \frac{1}{2} W^{\hat{\mu}\hat{\nu}} M_{\hat{\mu}\hat{\nu}} \right) \quad (4.2)$$

where

$$W^{\hat{\mu}} = - \int_\Gamma dx^\mu e^{\hat{\mu}}{}_\mu = \frac{1}{r} \bar{W}^{\hat{\mu}} + \frac{1}{r^2} W_{(2)}^{\hat{\mu}} + \mathcal{O}(1/r^3), \quad (4.3)$$

$$W^{\hat{\mu}\hat{\nu}} = - \int_\Gamma dx^\mu \omega^{\hat{\mu}\hat{\nu}}{}_\mu = \frac{1}{r^2} \bar{W}^{\hat{\mu}\hat{\nu}} + \mathcal{O}(1/r^3) \quad (4.4)$$

The last equalities above come from an asymptotic expansion of fields, as we will see explicitly in section 4.2. Equation (4.2) can be simplified using the Zassenhaus formula, stating that for two operators X, Y

$$e^{X+Y} = e^X e^Y e^{-\frac{1}{2}[X,Y]} e^{\frac{1}{6}(2[Y,[X,Y]]+[X,[X,Y]])} \dots \quad (4.5)$$

where \dots refer to exponentials involving more commutators. Taking $X = W^{\hat{\mu}} P_{\hat{\mu}}, Y = \frac{1}{2} W^{\hat{\mu}\hat{\nu}} M_{\hat{\mu}\hat{\nu}}$, and given the asymptotic form of $W^{\hat{\mu}}, W^{\hat{\mu}\hat{\nu}}$, we find that

$$W = \mathcal{P} \left(\exp(W^{\hat{\mu}} P_{\hat{\mu}}) \exp\left(\frac{1}{2} W^{\hat{\mu}\hat{\nu}} M_{\hat{\mu}\hat{\nu}}\right) \right) + \mathcal{O}(1/r^3) \quad (4.6)$$

$$= 1 + W^{\hat{\mu}} P_{\hat{\mu}} + \frac{1}{2} W^{\hat{\mu}\hat{\nu}} M_{\hat{\mu}\hat{\nu}} + \frac{1}{2} \mathcal{P}(W^{\hat{\mu}} W^{\hat{\nu}}) P_{\hat{\mu}} P_{\hat{\nu}} + \mathcal{O}(1/r^3) \quad (4.7)$$

Finite dimensional representation. We can explicitly compute the holonomy once we have a specific representation of the Poincaré algebra. In a d dimensional spacetime, a finite dimensional representation is given by a $d + 1$ dimensional vector Ψ^a , which transforms under Poincare group as

$$\Psi^a \equiv \begin{pmatrix} \psi^{\hat{\mu}} \\ c \end{pmatrix} \mapsto \begin{pmatrix} \Lambda^{\hat{\mu}}{}_{\hat{\nu}} & T^{\hat{\mu}} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \psi^{\hat{\mu}} \\ c \end{pmatrix} \quad (4.8)$$

where $c \in \mathbb{R}$ is an arbitrary real number. Note that c is invariant under the symmetry transformation and therefore the representation is decomposed into various orbits labeled by c , while $\psi^{\hat{\mu}}$ represents the physical state of the system. Depending on the value of c , $\psi^{\hat{\mu}}$ represent different physical systems. In particular, on the orbits $c = 1, c = 0$, the state transforms as

$$X^{\hat{\mu}} \equiv \psi^{\hat{\mu}}[c = 1], \quad X^{\hat{\mu}} \mapsto \Lambda^{\hat{\mu}}_{\hat{\nu}} X^{\hat{\nu}} + T^{\hat{\mu}}, \quad (4.9)$$

$$S^{\hat{\mu}} \equiv \psi^{\hat{\mu}}[c = 0], \quad S^{\hat{\mu}} \mapsto \Lambda^{\hat{\mu}}_{\hat{\nu}} \psi^{\hat{\nu}}. \quad (4.10)$$

$X^{\hat{\mu}}$ can represent the position of a particle in the local frame, while $S^{\hat{\mu}}$ can represent the spin of a gyroscope.

By linearizing (4.8) around the identity, we can find the matrix representation of the Poincaré algebra

$$(M_{\hat{\mu}\hat{\nu}})^a_b = \delta_{\hat{\mu}}^a \eta_{\hat{\nu}b} - \delta_{\hat{\nu}}^a \eta_{\hat{\mu}b} \quad (P_{\hat{\mu}})^a_b = \delta_{\hat{\mu}}^a \delta_b^{d+1} \quad (4.11)$$

In particular, translation $P_{\hat{\mu}}$ is nilpotent, *i.e.* $P^n = 0$ for $n \geq 2$ and therefore, the holonomy (4.7) reduces to

$$W = 1 + \frac{1}{r} \bar{W}^{\hat{\mu}} P_{\hat{\mu}} + \frac{1}{r^2} \left(\frac{1}{2} \bar{W}^{\hat{\mu}\hat{\nu}} M_{\hat{\mu}\hat{\nu}} + W_{(2)}^{\hat{\mu}} P_{\hat{\mu}} \right) + \mathcal{O}(1/r^3) \quad (4.12)$$

Therefore, at leading order in $1/r$ expansion, the holonomy is purely translational given by $\bar{W}^{\hat{\mu}}$, while it consists of a rotational (Lorentz) part $\bar{W}^{\hat{\mu}\hat{\nu}}$ as well as a subleading translational part $W_{(2)}^{\hat{\mu}}$ at $\mathcal{O}(1/r^2)$. In fact, it turns out that the only nontrivial components at leading order are $\bar{W}^{\hat{A}}, \bar{W}^{\hat{A}\hat{B}}$, *i.e.* when they involve transverse directions. We show that $\bar{W}^{\hat{A}}$ encodes the (leading and subleading) displacement memory, while the rotational holonomy $\bar{W}^{\hat{A}\hat{B}}$ reproduces the gyroscopic and spin memory once integrated along appropriate paths.

Poincaré holonomy and affine transport equation. A natural question is: which transport equation does the Poincaré holonomy solve? To answer this question, we consider again the finite dimensional representation discussed above. Assume that the initial state of the probe system is given by Ψ_0^a . Now the evolution of the initial state by the holonomy (4.2) is given by $\Psi^a(\tau) = (W_{\tau})^a_b \Psi_0^b$, where W_{τ} is the holonomy along an open path with endpoint $x^{\mu}(\tau)$ and with tangent vector $k^{\mu} = dx^{\mu}/d\tau$. It satisfies the differential equation

$$\frac{d}{d\tau} \Psi^a = -k^{\mu} (A_{\mu})^a_b \Psi^b \quad (4.13)$$

Using the decomposition $\Psi^a = (\psi^{\hat{\mu}}, c)$, and equations (4.1),(4.11), we find that the evolution equation, depending on the value of c , is given by

$$\frac{d}{d\tau} \psi^{\hat{\mu}} = -c k^{\hat{\mu}} - k^{\mu} \omega_{\mu}^{\hat{\mu}}{}_{\hat{\nu}} \psi^{\hat{\nu}} \quad (4.14)$$

Taking the last term to the left hand side, and multiplying the equation with inverse tetrad to go back to the coordinate basis reveals

$$k^{\mu} \nabla_{\mu} \psi^{\nu} = -c k^{\nu} \quad \psi^{\mu} \equiv e_{\hat{\mu}}^{\mu} \psi^{\hat{\mu}}. \quad (4.15)$$

Therefore, on the orbit $c = 0$, the evolution reduces to the parallel transport equation, while for $c = 1$, we recover the ‘affine’ transport equation of Refs. [38, 39]. The latter case ($c = 1$) corresponds to the evolution of the position of a test mass in the local frame, or equivalently, the physical distance between two nearby test masses, while the former case ($c = 0$) describes the evolution of the spin of a gyroscope.

4.2 Asymptotic frame

To compute the holonomy (4.12), we need to pick a particular frame. In this paper, we use the ‘source oriented frame’ constructed in [19, 20, 30], which is orthonormal everywhere in spacetime and therefore provides an extension of the asymptotic frame (2.3) inside the bulk. It is constructed such that the timelike basis vector $e_{\hat{0}}$ coincides with a geodesic congruence that is at rest at some initial time u_0 . Moreover, the radial basis vector $e_{\hat{r}}$ is tangent to the spatial path followed by rays arriving from the source. However, as we will discuss later, the essence of our result, namely the fact that holonomies encode memory effects is independent of the choice of frame. To support this, we study the gauge transformation properties of the translational holonomy $\bar{W}^{\hat{A}}$.

At large distance, the dual basis one-forms of the source oriented frame are given up to subleading corrections by

$$e^{\hat{0}} = (1 - m/r)du + (1 + m_0/r)dr - \left(\frac{1}{r} \int_{u_0}^u du' \partial_{AM} \right) d\theta^A \quad (4.16)$$

$$e^{\hat{r}} = -\frac{m - m_0}{r}du + (1 + m_0/r)dr - \left(\frac{1}{r} \int_{u_0}^u du' \partial_{AM} \right) d\theta^A \quad (4.17)$$

$$e^{\hat{A}} = E^{\hat{A}}{}_{\hat{B}} \left(\frac{1}{r} D_C C^{BC} du + (r\delta^B{}_A + \frac{1}{2} C^B{}_A) d\theta^A \right) \quad (4.18)$$

where $E^{\hat{A}}{}_{\hat{A}}(\theta^A)$ is a time-independent dyad on the sphere such that $\gamma^{AB} E^{\hat{A}}{}_A E^{\hat{B}}{}_B = \delta^{\hat{A}\hat{B}}$. The associated spin connections are given by $\omega = \bar{\omega} + \omega'$ where

$$\bar{\omega}_{\hat{r}\hat{A}} = -E_{\hat{A}\hat{A}} d\theta^A, \quad \bar{\omega}_{\hat{A}\hat{B}} = E_{\hat{A}}{}^B D_B E_{\hat{B}C} d\theta^C, \quad (4.19)$$

and

$$\omega'_{\hat{r}\hat{A}} = E_{\hat{A}}{}^A \left(\frac{1}{4r^2} N_{AB} D_C C^{BC} du + \frac{1}{2r^2} D^B C_{AB} dr + \frac{1}{2} N_{AB} d\theta^B \right), \quad (4.20)$$

$$\omega'_{\hat{A}\hat{B}} = \epsilon_{\hat{A}\hat{B}} \left[-\frac{1}{4r^2} (D_A D_B \tilde{C}^{AB} - \frac{1}{2} N_{AB} \tilde{C}^{AB}) du + \frac{1}{2r} D^A \tilde{C}_{AB} d\theta^B \right] + \mathcal{O}(r^{-3}) dr \quad (4.21)$$

In the above result, the $\bar{\omega}$ part of the spin connection is nonzero even in Minkowski background, while the genuine radiation effects are encoded in ω' . It is possible to define a rotated frame for which the connection is solely given by ω' . It was shown in [19, 20] that the latter corresponds to a frame whose spatial axes are tied to distant stars rather than to the ray arriving from the source.

In this section we compute gravitational holonomies over the paths depicted in Figure 2 and show that they reproduce gravitational memory effects discussed in section 3.




Figure 2: Paths implemented to compute gravitational holonomies. The paths are non-dynamical and specified by the initial data, while the dynamics is encoded in the holonomy. Figure (2a) is a trapezoidal spacetime loop whose spatial component is tangent the sphere and reveals leading and subleading displacement memories. Figure (2b) is a geodesic, which appears as a straight line in the comoving frame (4.16). Finally, figure (2c) is a helical path whose projection on transverse directions forms a loop. The last two figures give the gyroscopic memory and also the spin memory after an assumption on GW frequency.

Remark. Paths depicted in (2) do not have a dynamical content themselves, *i.e.* they can be thought of as spacetime paths in the Minkowski background. They are specified by the given data of the problem, including the initial data of the system and the time interval of the process. The dynamical content appears when one computes the holonomy along these paths.

4.3 Translational holonomy

Consider two test masses with initial relative physical distance and velocity $X_0 \equiv X(u_0)$ and $V_0 \equiv V(u_0)$, which can be thought of as the position and velocity of the second test mass in the local inertial frame (4.16) in which the first mass is at the origin. Also assume that the GW is being emitted during the time interval (u_0, u_f) . From this data, we construct a trapezoid depicted in figure (2a), and compute the translational holonomy (4.3):

$$\begin{aligned}
 W^{\hat{A}} = & - \int_0^{X_0} d\theta^A e^{\hat{A}}_{\ A}(u_0) + \int_0^{X(u)} d\theta^A e^{\hat{A}}_{\ A}(u_f) \\
 & + \int_{u_0}^u du e^{\hat{A}}_{\ u}(x_0) - \left(\int_{u_0}^u du e^{\hat{A}}_{\ u}(x) - \int_{X_0}^{X(u)} d\theta^A e^{\hat{A}}_{\ A} \right), \quad (4.22)
 \end{aligned}$$

with $u \geq u_f$. The first (second) line contains the contributions of the spatial (timelike) segments. After plugging $e^{\hat{A}}_{\ u}$ and $e^{\hat{A}}_{\ A}$ from (4.16), we find at leading order

$$\boxed{\bar{W}^{\hat{A}} = \frac{1}{2} \left[X_0^{\hat{B}} \Delta C^{\hat{A}}_{\ \hat{B}} - \beta^{\hat{B}} \int_{u_0}^{u_f} du (C^{\hat{A}}_{\ \hat{B}}(u) - C^{\hat{A}}_{\ \hat{B}}(u_f)) \right]} \quad (4.23)$$

where $\beta^{\hat{A}}$ is the slope of the trapezoid. Comparing this with (3.10a), we see that this holonomy reproduces the leading displacement memory effect, and the subleading cor-

rection once we take $\beta^{\hat{A}} = 2V^{\hat{A}}(u_0)$. As mentioned before, while the loop should be determined by the initial data, it does not represent the real path of the test masses³.

Recall an important property of holonomies. For two paths Γ_1, Γ_2 , the composition $\Gamma_1 \circ \Gamma_2$ is another loop in which overlapping segments with reverse arrows cancel out. The composition rule states that

$$W_{\Gamma_1 \circ \Gamma_2}(A) = W_{\Gamma_2}(A) \cdot W_{\Gamma_1}(A), \quad \forall A, \quad (4.24)$$

The trapezoid in figure (2a) can be decomposed into a rectangular loop as in Fig.(1a), and a triangular loop. The former gives the first term in (4.23), *i.e.* the leading displacement memory, while the triangular loop gives the second term, *i.e.* the subleading displacement memory.

Gauge transformation of the holonomy. We computed the holonomy above in a specific frame given by (4.16). However, the choice of frame is arbitrary and parametrized by internal Lorentz symmetries. In contrast to the Wilson loop, holonomy is not gauge invariant, but transforms under a gauge transformation $A \rightarrow g \cdot A = g^{-1}Ag + g^{-1}dg$ as

$$W[\Gamma, g \cdot A] = g^{-1}[\tau_f] W[\Gamma, A] g[\tau_0] \quad (4.25)$$

where τ_0, τ_f refer to the starting and end points of the path Γ . Now let us see how our result above transforms at leading order in $1/r$ under an infinitesimal Local Lorentz transformation $g = I + T^{\hat{A}}P_{\hat{A}} + \lambda^{\hat{A}\hat{B}}M_{\hat{A}\hat{B}}$. At leading order, $W[\Gamma, A] = 1 + \frac{1}{r}\bar{W}^{\hat{A}}P_{\hat{A}} + O(1/r^2)$, and therefore the gauge transformation induces

$$\bar{W}^{\hat{A}}P_{\hat{A}} \rightarrow \bar{W}^{\hat{A}} \left((\delta_{\hat{A}}^{\hat{B}} + \lambda_{\hat{A}}^{\hat{B}})P_{\hat{B}} + \lambda^{\hat{0}\hat{A}}P_{\hat{0}} \right), \quad (4.26)$$

implying that the translational holonomy $W_{(1)}^{\hat{A}}$ is a Lorentz vector, rotated by $\lambda^{\hat{A}\hat{B}}$ and boosted by $\lambda^{\hat{0}\hat{A}}$. This is expected since ‘displacement’ memory also transforms as a vector.

Kick memory. Now to reproduce the kick memory Eq.(3.10b), we need a structure like that of Maxwell, Eq.(2.15), *i.e.* an additional derivative acting on the holonomy. In this case, the index structure forces us to take

$$\boxed{v^{\hat{A}} = -V_0^{\hat{B}}\partial_{\hat{B}}\bar{W}^{\hat{A}}} \quad (4.27)$$

where $V_0^{\hat{B}}$ is the initial relative velocity and $\partial_{\hat{B}} = \partial/\partial X^{\hat{A}}(u_0)$ is derivative with respect to the initial position of the test mass. Using (4.23), we see that the above equation reproduces (3.10b). This explains why in gravity, the kick effect happens only in the presence of initial relative velocity, as opposed to EM. We saw that in Maxwell, the holonomy is a scalar, and the kick memory is obtained by taking a spatial derivative.

³This factor of 2 is potentially related to the fact that in the gravitoelectromagnetism (GEM) formulation of general relativity, gravitomagnetic charge of a test mass m is twice its gravitoelectric charge $q_B = 2q_E = -2m$ [57]. In this setup, motion of the test mass is given in a Fermi normal coordinate by the force law $\mathbf{F} = q_E\mathbf{E} + q_B\mathbf{v} \times \mathbf{B}$, where \mathbf{E}, \mathbf{B} are the gravitoelectric and gravitomagnetic fields. From this equation, it is clear that velocity effects are sourced through the gravitomagnetic charge and this leads to a relative factor of two between the first and second terms in the displacement memory effects Eq.(4.23).

However, in gravity, the translational holonomy already has an index, and if we take a derivative, we need to contract with another vector, given by the initial data. Dimensional analysis implies that the contracting vector needs to be a velocity, as in (4.27), which explains the absence of the kick memory in gravity for two test masses initially at rest.

Time translation. Intuitively, we may expect that $W^{\hat{0}}$, *i.e.* the time-translational holonomy gives the so called “relative proper time memory” [24,39]. However, computing this quantity, we find that

$$W^{\hat{0}} = \mathcal{O}(1/r^3). \quad (4.28)$$

Given that our frame (4.16) is adapted to freely falling observers, this result matches that of Strominger and Zhiboedov [24]. Computing the first nontrivial term in the expansion of $W^{\hat{0}}$, requires a more accurate expansion of the frame (4.16), which goes beyond the scope of this paper. However, it would be nice to compute this and compare with Eq.(2.6) of [39].

4.4 Rotational holonomy

In this section, we study the rotational holonomy (4.4) and show that for suitable choice of the path, it reproduces the gyroscopic and spin memory effects. Consider a generic path at large distance. Using (4.20), we find that the effect of radiation on the rotational holonomy is given by

$$W^{\hat{A}\hat{B}} = -\frac{1}{2r^2}\epsilon^{\hat{A}\hat{B}} \left[\int du (D_A D_B \tilde{C}^{AB} - \frac{1}{2} N_{AB} \tilde{C}^{AB}) - 2 \int r d\theta^A D^B \tilde{C}_{AB} \right] + \text{subleading}. \quad (4.29)$$

If the path is a timelike geodesic, as in figure (2b), which has zero velocity at some initial time, the angular velocity remains at $\mathcal{O}(1/r^2)$ at later times [19,20]. Therefore, the last term in (4.29) is subleading and we find

$$\boxed{\bar{W}^{\hat{A}\hat{B}} = -\frac{1}{2r^2}\epsilon^{\hat{A}\hat{B}} \left[\int du (D_A D_B \tilde{C}^{AB} - \frac{1}{2} N_{AB} \tilde{C}^{AB}) \right]}, \quad (4.30)$$

which coincides with the gyroscopic memory effect (3.13). This result is expected, because the evolution of a freely falling gyroscope is given by the parallel transport equation (3.12), which is actually solved by the holonomy

$$S^{\hat{i}} = \mathcal{P} \exp \left(- \int_{\Gamma} dx^{\mu} \omega_{\mu}^{\hat{i}\hat{j}} \right) S_0^{\hat{j}} \quad (4.31)$$

over a geodesic Γ followed by the gyroscope. The net change in the spin is given by $S^{\hat{j}}(u_f) - S_0^{\hat{j}}$ which is at leading order given by minus the rotational holonomy $W^{\hat{i}\hat{j}}$ defined in (4.4). In particular, at $\mathcal{O}(1/r^2)$, the only effect appears in $\hat{i}\hat{j} = \hat{A}\hat{B}$ components, which gives the gyroscopic memory, *i.e.* a rotation in the transverse plane, given by (3.13).

Another interesting situation is when the path is not a geodesic, but instead its spatial projection forms a loop in the transverse plane, as depicted in figure (2c). In this case,

the angular integral in (4.29) can also be written as a surface integral

$$\bar{W}^{\hat{A}\hat{B}} = -\frac{1}{2r^2}\epsilon^{\hat{A}\hat{B}} \left[\int du (D_A D_B \tilde{C}^{AB} - \frac{1}{2} N_{AB} \tilde{C}^{AB}) - r \int_{D(\Gamma)} d^2\theta \sqrt{q} D_A D_B C^{AB} \right]. \quad (4.32)$$

Since the size of the path, denoted by L , is by assumption much smaller than the distance r to the source, the angular integral is subleading by a factor L/r with respect to the first integral (since $d\theta \sim L/r$), and we again reproduce the gyroscopic memory (4.30). This second path is more closely related to the setup of the Sagnac effect, where light beams counter-orbit a closed spatial path. It is not surprising that we get the same gyroscopic memory from the latter, since Sagnac interferometer is nothing but an optical gyroscope.

To reproduce the spin memory [16], we further assume that L is much smaller than the typical wavelength λ of the gravitational wave. Then the second term in (4.30) is smaller than the first term by a factor L/λ , and Eq.(4.30) simplifies to

$$\bar{W}^{\hat{A}\hat{B}} = -\frac{\Delta u}{2r^2}\epsilon^{\hat{A}\hat{B}} D_A D_B \tilde{C}^{AB}(u_0). \quad (4.33)$$

To relate this to the spin memory, we Taylor expand (3.11) around the experiment space-time event (u_0, θ_0) , and we find $\Delta T = \frac{\text{Area}}{r^2} D_A D_B \tilde{C}^{AB}(u_0, \theta_0)$. Therefore the simplified rotational holonomy is related to the spin memory (3.11) as

$$\bar{W}^{\hat{A}\hat{B}} = -\frac{\Delta T}{2L}\epsilon^{\hat{A}\hat{B}}. \quad (4.34)$$

5 Discussion and outlook

In this paper, we showed that known types of memory effects can be derived from certain Wilson loops/ holonomies on a holographic screen, *i.e.* a hypersurface of constant radius at large distance in asymptotically flat spacetimes. Using the Nonabelian Stokes theorem [41] which generalizes the usual Stokes theorem to Lie algebra valued differential one-forms, one may be able to write equivalent expressions for these holonomies as surface integrals. Inspired by results in AdS/CFT, such surface integrals may be considered as holographic dual expressions for the holonomies. From the boundary point of view, it is believed that gravity in asymptotically flat spacetimes may be holographically described by a Carrollian theory [58–64]. If such a correspondence exists, holonomies discussed here may be related to ‘out of time ordered correlators’ (OTOC) on the boundary. Such observables have already received much attention in the context of AdS/CFT [65]. Further investigations along these lines will be of great interest.

Another point that we plan to further investigate in a future work is the investigation of memory effects from the viewpoint of gravito-electromagnetism for a better understanding of these effects and the peculiar feature that we encountered in relating the subleading displacement memory to the corresponding holonomy. Finally, it would be interesting to study more subleading memory effects and their relationship to holonomies, in particular, the relative proper time memory mentioned around Eq.(4.28).

Acknowledgments

A. Seraj is grateful to Aldo Riello for proposing the possible relationship between Wilson loops and memory effects. We also thank H. Afshar, W. Merbis, B. Oblak, J. Salzer and M.M. Sheikh Jabbari for useful discussions. This research is supported by the ERC Advanced Grant “High-Spin-Grav”, and the ERC Starting Grant “Holography for realistic black holes”.

References

- [1] Y. B. Zel’dovich and A. G. Polnarev, *Radiation of gravitational waves by a cluster of superdense stars*, Sov. Astron. **18** (1974) 17.
- [2] V. B. Braginsky and L. P. Grishchuk, *Kinematic Resonance and Memory Effect in Free Mass Gravitational Antennas*, Sov. Phys. JETP **62** (1985) 427–430.
- [3] D. Christodoulou, *Nonlinear nature of gravitation and gravitational wave experiments*, Phys. Rev. Lett. **67** (1991) 1486–1489.
- [4] L. Blanchet and T. Damour, *Hereditary effects in gravitational radiation*, Phys. Rev. D **46** (1992) 4304–4319.
- [5] K. S. Thorne, *Gravitational-wave bursts with memory: The Christodoulou effect*, Phys. Rev. D **45** (1992), no. 2 520–524.
- [6] M. Favata, *The gravitational-wave memory effect*, Class. Quant. Grav. **27** (2010) 084036 [[1003.3486](#)].
- [7] A. G. Wiseman and C. M. Will, *Christodoulou’s nonlinear gravitational wave memory: Evaluation in the quadrupole approximation*, Phys. Rev. D **44** (1991), no. 10 R2945–R2949.
- [8] P. D. Lasky, E. Thrane, Y. Levin, J. Blackman and Y. Chen, *Detecting gravitational-wave memory with LIGO: implications of GW150914*, Phys. Rev. Lett. **117** (2016), no. 6 061102 [[1605.01415](#)].
- [9] M. Hübner, C. Talbot, P. D. Lasky and E. Thrane, *Measuring gravitational-wave memory in the first LIGO/Virgo gravitational-wave transient catalog*, Phys. Rev. D **101** (2020), no. 2 023011 [[1911.12496](#)].
- [10] O. M. Boersma, D. A. Nichols and P. Schmidt, *Forecasts for detecting the gravitational-wave memory effect with Advanced LIGO and Virgo*, Phys. Rev. D **101** (2020), no. 8 083026 [[2002.01821](#)].
- [11] M. Hübner, P. Lasky and E. Thrane, *Memory remains undetected: Updates from the second LIGO/Virgo gravitational-wave transient catalog*, Phys. Rev. D **104** (2021), no. 2 023004 [[2105.02879](#)].
- [12] G. Compère, R. Oliveri and A. Seraj, *The Poincaré and BMS flux-balance laws with application to binary systems*, JHEP **10** (2020) 116 [[1912.03164](#)].
- [13] K. Mitman *et. al.*, *Adding gravitational memory to waveform catalogs using BMS balance laws*, Phys. Rev. D **103** (2021), no. 2 024031 [[2011.01309](#)].
- [14] K. Mitman, J. Moxon, M. A. Scheel, S. A. Teukolsky, M. Boyle, N. Deppe, L. E. Kidder and W. Throwe, *Computation of displacement and spin gravitational memory in numerical relativity*, Phys. Rev. D **102** (2020), no. 10 104007 [[2007.11562](#)].
- [15] K. Mitman *et. al.*, *Fixing the BMS frame of numerical relativity waveforms*, Phys. Rev. D **104** (2021), no. 2 024051 [[2105.02300](#)].

- [16] S. Pasterski, A. Strominger and A. Zhiboedov, *New Gravitational Memories*, JHEP **12** (2016) 053 [[1502.06120](#)].
- [17] D. A. Nichols, *Spin memory effect for compact binaries in the post-Newtonian approximation*, Phys. Rev. D **95** (2017), no. 8 084048 [[1702.03300](#)].
- [18] D. A. Nichols, *Center-of-mass angular momentum and memory effect in asymptotically flat spacetimes*, Phys. Rev. D **98** (2018), no. 6 064032 [[1807.08767](#)].
- [19] A. Seraj and B. Oblak, *Gyroscopic Gravitational Memory*, [2112.04535](#).
- [20] A. Seraj and B. Oblak, *The Precession Caused by Gravitational Waves*, [2203.16216](#).
- [21] L. Freidel, D. Pranzetti and A.-M. Raclariu, *Higher spin dynamics in gravity and $w_{1+\infty}$ celestial symmetries*, [2112.15573](#).
- [22] G. Compère, R. Oliveri and A. Seraj, *Metric reconstruction from celestial multipoles*, [2206.12597](#).
- [23] A. M. Grant and D. A. Nichols, *Persistent gravitational wave observables: Curve deviation in asymptotically flat spacetimes*, Phys. Rev. D **105** (2022), no. 2 024056 [[2109.03832](#)].
- [24] A. Strominger and A. Zhiboedov, *Gravitational Memory, BMS Supertranslations and Soft Theorems*, JHEP **01** (2016) 086 [[1411.5745](#)].
- [25] A. Strominger, *$w(1+\text{infinity})$ and the Celestial Sphere*, [2105.14346](#).
- [26] A. Strominger, *$w_{1+\infty}$ Algebra and the Celestial Sphere: Infinite Towers of Soft Graviton, Photon, and Gluon Symmetries*, Phys. Rev. Lett. **127** (2021), no. 22 221601.
- [27] L. Freidel, D. Pranzetti and A.-M. Raclariu, *Sub-subleading soft graviton theorem from asymptotic Einstein's equations*, JHEP **05** (2022) 186 [[2111.15607](#)].
- [28] M. Geiller and C. Zwikel, *The partial Bondi gauge: Further enlarging the asymptotic structure of gravity*, [2205.11401](#).
- [29] E. Himwich, Z. Mirzaiyan and S. Pasterski, *A Note on the Subleading Soft Graviton*, JHEP **04** (2021) 172 [[1902.01840](#)].
- [30] M. Godazgar, G. Long and A. Seraj, *Gravitational memory effects and higher derivative actions*, [2206.12339](#).
- [31] L. Bieri and D. Garfinkle, *Perturbative and gauge invariant treatment of gravitational wave memory*, Phys. Rev. D **89** (2014), no. 8 084039 [[1312.6871](#)].
- [32] L. Susskind, *Electromagnetic Memory*, [1507.02584](#).
- [33] S. Pasterski, *Asymptotic Symmetries and Electromagnetic Memory*, JHEP **09** (2017) 154 [[1505.00716](#)].
- [34] S. Tahura, D. A. Nichols, A. Saffer, L. C. Stein and K. Yagi, *Brans-Dicke theory in Bondi-Sachs form: Asymptotically flat solutions, asymptotic symmetries and gravitational-wave memory effects*, Phys. Rev. D **103** (2021), no. 10 104026 [[2007.13799](#)].
- [35] S. Tahura, D. A. Nichols and K. Yagi, *Gravitational-wave memory effects in Brans-Dicke theory: Waveforms and effects in the post-Newtonian approximation*, Phys. Rev. D **104** (2021), no. 10 104010 [[2107.02208](#)].
- [36] S. Hou and Z.-H. Zhu, *Gravitational memory effects and Bondi-Metzner-Sachs symmetries in scalar-tensor theories*, JHEP **01** (2021) 083 [[2005.01310](#)].
- [37] A. Seraj, *Gravitational breathing memory and dual symmetries*, JHEP **05** (2021) 283 [[2103.12185](#)].

- [38] E. E. Flanagan and D. A. Nichols, *Observer dependence of angular momentum in general relativity and its relationship to the gravitational-wave memory effect*, Phys. Rev. D **92** (2015), no. 8 084057 [[1411.4599](#)]. [Erratum: Phys.Rev.D 93, 049905 (2016)].
- [39] E. E. Flanagan, A. M. Grant, A. I. Harte and D. A. Nichols, *Persistent gravitational wave observables: general framework*, Phys. Rev. D **99** (2019), no. 8 084044 [[1901.00021](#)].
- [40] M. Ammon and J. Erdmenger, *Gauge/gravity duality: Foundations and applications*. Cambridge University Press, Cambridge, 4, 2015.
- [41] U. Schreiber and K. Waldorf, *Smooth functors vs. differential forms*, Homology, Homotopy and Applications **13** (2011), no. 1 143–203.
- [42] A. Strominger, *Lectures on the Infrared Structure of Gravity and Gauge Theory*, [1703.05448](#).
- [43] R. Loll, *Gauge theory and gravity in the loop formulation*, Lect. Notes Phys. **434** (1994) 254–288.
- [44] P. Mao, H. Ouyang, J. B. Wu and X. Wu, *New electromagnetic memories and soft photon theorems*, Phys. Rev. D **95**, no.12, 125011 (2017) [[1703.06588](#)]
- [45] H. Bondi, M. G. J. van der Burg and A. W. K. Metzner, *Gravitational waves in general relativity. 7. Waves from axisymmetric isolated systems*, Proc. Roy. Soc. Lond. A **269** (1962) 21–52.
- [46] R. K. Sachs, *Gravitational waves in general relativity. 8. Waves in asymptotically flat space-times*, Proc. Roy. Soc. Lond. A **270** (1962) 103–126.
- [47] G. Barnich and C. Troessaert, *Symmetries of asymptotically flat 4 dimensional space-times at null infinity revisited*, Phys. Rev. Lett. **105** (2010) 111103 [[0909.2617](#)].
- [48] G. Barnich and C. Troessaert, *Aspects of the BMS/CFT correspondence*, JHEP **05** (2010) 062 [[1001.1541](#)].
- [49] M. Campiglia and J. Peraza, *Generalized BMS charge algebra*, Phys. Rev. D **101** (2020), no. 10 104039 [[2002.06691](#)].
- [50] H. Godazgar, M. Godazgar and C. N. Pope, *Tower of subleading dual BMS charges*, JHEP **03** (2019) 057 [[1812.06935](#)].
- [51] C. W. F. Everitt *et. al.*, *Gravity Probe B: Final Results of a Space Experiment to Test General Relativity*, Phys. Rev. Lett. **106** (2011) 221101 [[1105.3456](#)].
- [52] L. Herrera and J. L. Hernandez Pastora, *On the influence of gravitational radiation on a gyroscope*, Class. Quant. Grav. **17** (2000) 3617–3626 [[gr-qc/0006080](#)].
- [53] D. Bini and F. de Felice, *Gyroscopes and gravitational waves*, Class. Quant. Grav. **17** (2000) 4627–4635.
- [54] F. Sorge, D. Bini and F. de Felice, *Gravitational waves, gyroscopes and frame dragging*, Class. Quant. Grav. **18** (2001) 2945–2958.
- [55] L. Freidel and D. Pranzetti, *Gravity from symmetry: Duality and impulsive waves*, [2109.06342](#).
- [56] D. Ivanenko and G. Sardanashvily, *The Gauge Treatment of Gravity*, Phys. Rept. **94** (1983) 1–45.
- [57] B. Mashhoon, *Gravitoelectromagnetism: A Brief review*, [gr-qc/0311030](#).
- [58] J.-M. Lévy-Leblond, *Une nouvelle limite non-relativiste du groupe de poincaré*, in *Annales de l’IHP Physique théorique*, vol. 3, pp. 1–12, 1965.

- [59] C. Duval and P. A. Horvathy, *Non-relativistic conformal symmetries and Newton-Cartan structures*, J. Phys. A **42** (2009) 465206 [[0904.0531](#)].
- [60] C. Duval, G. W. Gibbons and P. A. Horvathy, *Conformal Carroll groups and BMS symmetry*, Class. Quant. Grav. **31** (2014) 092001 [[1402.5894](#)].
- [61] L. Ciambelli, C. Marteau, A. C. Petkou, P. M. Petropoulos and K. Siampos, *Flat holography and Carrollian fluids*, JHEP **07** (2018) 165 [[1802.06809](#)].
- [62] A. Bagchi, A. Mehra and P. Nandi, *Field Theories with Conformal Carrollian Symmetry*, JHEP **05** (2019) 108 [[1901.10147](#)].
- [63] Y. Herfray, *Carrollian manifolds and null infinity: A view from Cartan geometry*, [2112.09048](#).
- [64] L. Donnay, A. Fiorucci, Y. Herfray and R. Ruzziconi, *A Carrollian Perspective on Celestial Holography*, [2202.04702](#).
- [65] J. Maldacena, S. H. Shenker and D. Stanford, *A bound on chaos*, JHEP **08** (2016) 106 [[1503.01409](#)].