

Local well-posedness of the 1d compressible Navier–Stokes system with rough data

Ke Chen,^{*} Ruilin Hu,[†] Quoc-Hung Nguyen[‡]

June 29, 2022

Abstract

This paper presents a new approach to the local well-posedness of the 1d compressible Navier–Stokes systems with rough initial data. Our approach is based on establishing some smoothing and Lipschitz-type estimates for the 1d parabolic equation with piecewise continuous coefficients.

1 Introduction

In this paper, we study the compressible Navier–Stokes equations in Lagrangian coordinates, which can be written as (see [16])

$$\begin{cases} v_t - u_x = 0, \\ u_t + p_x = \left(\frac{\mu u_x}{v} \right)_x, \\ \left(e + \frac{1}{2} u^2 \right)_t + (pu)_x = \left(\frac{\kappa}{v} \theta_x + \frac{\mu}{v} u u_x \right)_x. \end{cases} \quad (1.1)$$

Here we denote v the specific volume, u the velocity, p the pressure, e the specific internal energy, θ the absolute temperature. And $\mu, \kappa > 0$ are viscosity and heat conductivity coefficients. The above equations describe the conservation of mass, momentum and energy, respectively.

We will consider two systems. The first system is the so-called p -system (see [16]), which is a general model of isentropic gas dynamics (when $p(v) = Av^{-\gamma}$). In the isentropic case, the temperature is held constant hence energy must be added to the system, whence the conservation of energy is absent. The p -system can be written as

$$\begin{cases} v_t - u_x = 0, \\ u_t + (p(v))_x = \left(\frac{\mu u_x}{v} \right)_x. \end{cases} \quad (1.2)$$

In this paper we consider a more general system which only requires $p \in W^{2,\infty}$.

Another system studied in this paper is (1.1) for the polytropic ideal gas which has the constitutive relations

$$p(v, \theta) = \frac{K\theta}{v}, \quad e = c\theta,$$

^{*}E-mail address: kchen18@fudan.edu.cn, Fudan University, 220 Handan Road, Yangpu, Shanghai, 200433, China.

[†]E-mail address: huruilin16@mails.ucas.ac.cn, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing, 100190, China.

[‡]E-mail address: qhnguyen@amss.ac.cn, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing, 100190, China.

where K and heat capacity $\mathbf{c}$ are both positive constants. The system can be written as

$$\begin{cases} v_t - u_x = 0, \\ u_t + (p(v, \theta))_x = \left(\frac{\mu u_x}{v}\right)_x, \\ \theta_t + \frac{p(v, \theta)}{\mathbf{c}} u_x - \frac{\mu}{\mathbf{c}v} (u_x)^2 = \left(\frac{\kappa}{\mathbf{c}v} \theta_x\right)_x. \end{cases} \quad (1.3)$$

Let us give a review of classical results. The study about compressible Navier-Stokes equations in gas dynamics started with Nash in [15], where he considered general elliptic and parabolic equations. Then Itaya [5, 6] solved the compressible Navier-Stokes equations for initial data in Hölder space. Kazhikhov [9] established a priori estimates and proved the existence of weak and classical solutions. For the equation of 3-d ideal gas, Nishida and Matsumura in [12] applied the energy method to give the existence of a global solution with data in Hölder space. The result of large-time behavior of global solution was provided by Kawashima [8]. Using the energy method, Hoff [4] proved the existence of a global solution for discontinuous data. The results in the book of Lions [10] and the book of Feireisl [2] are applicable to a larger space of initial data.

Recently, Liu and Yu [11] studied the system (1.2) with $BV \cap L^1$ data, under the assumption $p'(v) < 0$. Based on exact analysis of the associated linear equations and their Green's functions, they initiated a new method, which converts the differential equations into integral equations, and proved that for initial data v_0, u_0 satisfying

$$\|v_0 - 1\|_{L^1}, \|v_0\|_{BV}, \|u_0\|_{L^1}, \|u_0\|_{BV} < \delta \ll 1,$$

there exist $t_\sharp > 0$ and $C_\sharp > 0$ such that the system (1.2) admits a unique weak solution (v, u) for $t \in (0, t_\sharp)$, and

$$\|v(t, \cdot) - 1\|_{L^1} + \|v(t, \cdot)\|_{BV} + \|\sqrt{t}u_x(t, \cdot)\|_{L^\infty} \leq 2C_\sharp \delta, \quad \forall t \in (0, t_\sharp).$$

Moreover, they also established global result for polytropic gases $p(v) = Av^{-\gamma}$ with $1 \leq \gamma < e$. Their result was extended to the full compressible Navier-Stokes system (1.3) by Wang, Yu, and Zhang [18].

In this paper, we are interested in the local well-posedness of (1.2) and (1.3) with rough data, which allows v_0 to have a finite jump. The idea to study the elliptic problem with jump coefficients comes from [13, 14]. More precisely, without loss of generality, let $x = 0$ be the jump point. Suppose there exists $\varepsilon > 0$ such that

$$|v_0(x) - v_0(y)| \leq \delta, \quad \text{if } |x - y| \leq \varepsilon \text{ and } xy > 0, \quad (1.4)$$

for some $\delta > 0$ that will be fixed later.

Throughout the paper, we fix two constants $0 < \alpha < \gamma \ll 1$. Define the norms

$$\begin{aligned} \|f\|_{X_T^{\sigma,p}(E)} &= \sup_{0 < s < T} s^\sigma \|f(s)\|_{L^p(E)} + \sup_{0 < s < t < T} s^{\sigma+\alpha} \frac{\|f(t) - f(s)\|_{L^p(E)}}{(t-s)^\alpha}, \\ \|f\|_{Y_T(E)} &= \sup_{0 < s < T} \|f(s)\|_{L^\infty(E)} + \sup_{0 < s < t < T} s^\alpha \frac{\|f(t) - f(s)\|_{L^\infty(E)}}{(t-s)^\alpha}, \\ \|f\|_{L_T^p(E)} &= \|f\|_{L_T^p(E)} = \left(\int_0^T \int_E |f(t, x)|^p dx dt \right)^{\frac{1}{p}}, \quad \text{for any measurable set } E \subseteq \mathbb{R}, \end{aligned}$$

and we omit the domain if $E = \mathbb{R}$. We write several norms together to denote the sum of them. Moreover, we denote $A \lesssim B$ if there exists a universal constant C such that $A \leq CB$. We will solve the system (1.2) and (1.3) by the fixed point theorem in the Banach space equipped with the norms above. We note that the norms $\|\cdot\|_{X_T^{\sigma,p}}$ and $\|\cdot\|_{Y_T}$ contain Hölder derivatives in time.

For the system (1.2), we impose the following conditions for initial data

$$v_0 \in L^\infty, \inf_x v_0 \geq \lambda_0 > 0, u_0 = \partial_x \bar{u}_0 \text{ with } \bar{u}_0 \in \dot{C}^{2\gamma}. \quad (1.5)$$

Simplified versions of our main results are stated in the following theorems.

Theorem 1.1 *Assume $p \in W^{2,\infty}(\mathbb{R})$. There exists $\delta_0 > 0$ such that if initial data (v_0, u_0) satisfies the conditions (1.4) and (1.5) with $\delta = \delta_0$, then the system (1.2) admits a unique local solution (v, u) in $[0, T]$ satisfying*

$$\inf_{t \in [0, T]} \inf_x v(t, x) \geq \frac{\lambda_0}{2}, \quad \|v\|_{Y_T} \leq 2\|v_0\|_{L^\infty}, \quad \|\partial_x u\|_{X_T^{1-\gamma, \infty}} \leq M,$$

for some $T, M > 0$.

For the full N-S system (1.3), we use L^p setting, and set the following norm and corresponding Banach space

$$\|(w, \vartheta)\|_{Z_T} := \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2}, 2}, X_T^{\frac{3}{4}, \infty}\}} \|\partial_x w\|_\star + \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2}, 2}\}} \|\vartheta\|_\star + \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6}, \frac{6}{5}}\}} \|\partial_x \vartheta\|_\star.$$

The above norms have the same scaling in the parabolic setting. The norms $\|\partial_x w\|_{L_T^2}$, $\|\vartheta\|_{L_T^2}$ and $\|\partial_x \vartheta\|_{L_T^{\frac{6}{5}}}$ are imposed in view of the parabolic structure of the equation of (u, θ) . Other norms $\|\partial_x w\|_{X_T^{\frac{1}{2}, 2}}$, $\|\partial_x w\|_{X_T^{\frac{3}{4}, \infty}}$, $\|\vartheta\|_{X_T^{\frac{1}{2}, 2}}$, $\|\partial_x \vartheta\|_{X_T^{\frac{5}{6}, \frac{6}{5}}}$ are necessary to obtain local estimates of v .

We assume the initial data satisfies that

$$\inf_x v_0(x) \geq \lambda_0 > 0, \quad \|v_0\|_{L^\infty} < \infty, \quad \|u_0\|_{L^2} < \infty, \quad \|\theta_0\|_{\dot{W}^{-\frac{2}{3}, \frac{6}{5}}} < \infty. \quad (1.6)$$

The following result is the local well-posedness of system (1.3).

Theorem 1.2 *There exists $\delta_0 > 0$ such that if initial data (v_0, u_0, θ_0) satisfies (1.4) and (1.6) with $\delta = \delta_0$, then the system (1.3) has a unique local solution (v, u) in $[0, T]$ satisfying*

$$\inf_{t \in [0, T]} \inf_x v(t, x) \geq \frac{\lambda_0}{2}, \quad \|v\|_{Y_T} \leq 2\|v_0\|_{L^\infty}, \quad \|(u, \theta)\|_{Z_T} \leq B.$$

for some $T, B > 0$.

Note that we do not need u_0, θ_0 decay at infinity. Hence the condition (1.6) can be relaxed to local L^p norms (see Theorem 5.2):

$$\inf_x v_0(x) \geq \lambda_0 > 0, \quad \|v_0\|_{L^\infty} < \infty, \quad \sup_{z \in \mathbb{R}} \|u_0 \chi_z\|_{L^2} < \infty, \quad \sup_{z \in \mathbb{R}} \|\theta_0 \chi_z\|_{\dot{W}^{-\frac{2}{3}, \frac{6}{5}}} < \infty. \quad (1.7)$$

Here χ_z is a smooth cutoff function satisfying $\mathbf{1}_{[z-1, z+1]} \leq \chi_z \leq \mathbf{1}_{[z-2, z+2]}$, where $\mathbf{1}$ is the indicator function.

Our method is based on analysis of heat equation with jump coefficient $\phi(x)$ satisfying (3.2),

$$\begin{aligned} \partial_t f(t, x) - \partial_x(\phi(x) \partial_x f(t, x)) &= \partial_x F(t, x) + R(t, x), \\ f(0, x) &= f_0(x). \end{aligned}$$

We convert the differential equations into integral equations and give two different formulas of solution. One formula (3.9) is suitable for estimates near jump points, and another one (3.37) behaves well for estimates far from jump points. Our main results rely on Lemma 3.1.

We make some remarks about the choice of initial data. For the isentropic system (1.2), we requires $u_0 \in \dot{B}_{\infty, \infty}^{-1+2\gamma}$. Generally in Schauder theory of parabolic/elliptic equations, to control the Lipschitz norm of solution, it requires the coefficient to be at least Dini (see [7]). Due to

the roughness of coefficient, the space $\dot{B}_{\infty,\infty}^{-1+2\gamma}$ is optimal in the sense that we can take $\gamma > 0$ arbitrarily close to 0. For the full N-S system (1.2), we require $u_0 \in L^2$ and $\theta_0 \in \dot{W}^{-\frac{2}{3}, \frac{6}{5}}$. Taking a glance at the equation of θ , we need $(u_x)^2, \theta u_x \in L_T^1$ to make θ well-defined. By classical theory of heat equation, $u_x \in L_T^2$ requires $u_0 \in L^2$, and $\theta \in L_T^2$ requires $\theta_0 \in \dot{H}^{-1}$. Taking force terms into consideration, we finally choose $\theta_0 \in \dot{W}^{-\frac{2}{3}, \frac{6}{5}} \subset \dot{H}^{-1}$.

The rest of this paper is organized as follows. We introduce some primary properties for heat equations and heat kernel in Section 2. We establish the main lemma 3.1 in Section 3. In Section 4 we apply Lemma 3.1 to obtain local well-posedness for system (1.2) and prove Theorem 1.1. Section 5 is devoted to prove local wellposedness for system (1.3) and prove Theorem 1.2.

2 Preliminaries

We introduce the following estimates for the standard heat kernel $\mathbf{K}(t, x) = (4\pi t)^{-\frac{1}{2}} e^{-\frac{x^2}{4t}}$ in $\mathbb{R}$ which satisfies

$$\begin{aligned}\partial_t \mathbf{K}(t, x) - \partial_{xx} \mathbf{K}(t, x) &= 0, \\ \lim_{t \rightarrow 0^+} \mathbf{K}(t, x) &= \text{Dirac}(x).\end{aligned}$$

The following estimates for heat kernel will be used frequently in our proof.

Lemma 2.1 *There holds*

$$\begin{aligned}|\partial_t^j \partial_x^m \mathbf{K}(t, x)| &\lesssim \frac{1}{(t^{\frac{1}{2}} + |x|)^{1+2j+m}}, \\ \left(\int_0^\infty |\partial_x^m \mathbf{K}(t, x)|^p dt \right)^{\frac{1}{p}} &\lesssim |x|^{\frac{2}{p}-1-m}, \quad \left(\int_{\mathbb{R}} |\partial_x^m \mathbf{K}(t, x)|^p dx \right)^{\frac{1}{p}} \lesssim t^{\frac{1}{2}(\frac{1}{p}-1-m)}, \\ \left(\int_{\mathbb{R}} |\partial_t^j \partial_x^m \mathbf{K}(t, x)|^p |x|^{\sigma p} dx \right)^{\frac{1}{p}} &\lesssim \frac{1}{t^{j+\frac{1}{2}(m-\frac{1}{p}+1-\sigma)}}, \\ \left(\int_{\mathbb{R}} |\partial_t^j \partial_x^m \mathbf{K}(t+a, x) - \partial_t^j \partial_x^m \mathbf{K}(t, x)|^p |x|^{\sigma p} dx \right)^{\frac{1}{p}} &\lesssim \frac{1}{t^{j+\frac{1}{2}(m-\frac{1}{p}+1-\sigma)}} \min\{1, \frac{a}{t}\},\end{aligned}$$

for any $m, j = 0, 1, 2$, any $\sigma, a, t \geq 0$ and $p \in [1, +\infty]$.

It is easy to check the above estimates by the definition of heat kernel and the fact that $b^m e^{-b} \lesssim_m 1$, $\forall b > 0, m \in \mathbb{N}^+$. We omit details here.

Lemma 2.2 *Let $h \in \dot{W}^{\sigma,p}(\mathbb{R})$ for some $\sigma \in (0, 1), p \in [1, \infty)$. Define $\mathbf{h}(x) = h(x)\mathbf{1}_{x \geq 0} + h(-x)\mathbf{1}_{x < 0}$ and $\bar{\mathbf{h}}(x) = (h(x) - h(-x))\mathbf{1}_{x \leq 0}$. Then we have*

$$\int_{\mathbb{R}} \int_0^\infty |\mathbf{h}(x) - \mathbf{h}(x-z)|^p \frac{dx dz}{|z|^{1+\sigma p}} + \int_{\mathbb{R}} \int_0^\infty |\bar{\mathbf{h}}(x) - \bar{\mathbf{h}}(x-z)|^p \frac{dx dz}{|z|^{1+\sigma p}} \lesssim \|h\|_{\dot{W}^{\sigma,p}}^p.$$

Proof. By definition, one can check that

$$\begin{aligned}\frac{|\mathbf{h}(x) - \mathbf{h}(x-z)|^p}{|z|^{1+\sigma p}} &= \mathbf{1}_{x \geq z} \frac{|h(x) - h(x-z)|^p}{|z|^{1+\sigma p}} + \mathbf{1}_{x < z} \frac{|h(x) - h(z-x)|^p}{|z|^{1+\sigma p}} \\ &\leq \frac{|h(x) - h(x-z)|^p}{|z|^{1+\sigma p}} + \frac{|h(x) - h(z-x)|^p}{|2x-z|^{1+\sigma p}}, \\ \frac{|\bar{\mathbf{h}}(x) - \bar{\mathbf{h}}(x-z)|^p}{|z|^{1+\sigma p}} &= \mathbf{1}_{x < z} \frac{|h(x-z) - h(z-x)|^p}{|z|^{1+\sigma p}} \\ &\leq \frac{|h(x) - h(x-z)|^p}{|z|^{1+\sigma p}} + \frac{|h(x) - h(z-x)|^p}{|2x-z|^{1+\sigma p}}.\end{aligned}$$

The result is standard in view of the fact that

$$\|h\|_{\dot{W}^{\sigma,p}}^p = \iint_{\mathbb{R}^2} \frac{|h(x) - h(y)|^p}{|x - y|^{1+\sigma p}} dx dy.$$

This completes the proof. ■

We introduce the following Schauder type lemma, the idea can be found in [1].

Lemma 2.3 *Let $\mathbf{K}$ be the heat kernel, and let*

$$g(t, x) = \int_0^t \int_{\mathbb{R}} \partial_t \mathbf{K}(t - \tau, x - y) f(\tau, y) dy d\tau.$$

Then we have

$$\|g\|_{X_T^{\sigma,p}} \lesssim \|f\|_{X_T^{\sigma,p}}, \quad \forall T > 0, \quad \sigma \in (0, 1 - \alpha), \quad p \in [1, \infty]. \quad (2.1)$$

Proof. One has

$$\begin{aligned} g(t, x) &= \int_0^t \int_{\mathbb{R}} \partial_t \mathbf{K}(t - \tau, x - y) (f(\tau, y) - f(t, y)) dy d\tau + \int_0^t \int_{\mathbb{R}} \partial_t \mathbf{K}(t - \tau, x - y) f(t, y) dy d\tau \\ &:= g_1(t, x) + g_2(t, x). \end{aligned}$$

By Lemma 2.1 we get

$$\|g_1(t)\|_{L^p} \lesssim \int_0^t \|\partial_t \mathbf{K}(t - \tau)\|_{L^1} \|f(\tau) - f(t)\|_{L^p} d\tau \lesssim \int_0^t (t - \tau)^{\alpha-1} \tau^{-\sigma-\alpha} d\tau \|f\|_{X_T^{\sigma,p}} \lesssim t^{-\sigma} \|f\|_{X_T^{\sigma,p}}.$$

Moreover, observe that $\int_0^t \partial_t \mathbf{K}(t - \tau, x - y) d\tau = K(t, x - y) - \mathbf{Dirac}(x - y)$, hence

$$g_2(t, x) = \int_{\mathbb{R}} \mathbf{K}(t, x - y) f(t, y) dy - f(t, x).$$

Then

$$\|g_2(t)\|_{L^p} \lesssim (1 + \|\mathbf{K}(t)\|_{L^1}) \|f(t)\|_{L^p} \lesssim \|f(t)\|_{L^p} \lesssim t^{-\sigma} \|f\|_{X_T^{\sigma,p}}.$$

Thus, we obtain that

$$\sup_{s \in [0, T]} s^\sigma \|g(s)\|_{L^p} \lesssim \|f\|_{X_T^{\sigma,p}}. \quad (2.2)$$

In the following we will denote $a = t - s > 0$ for convenience, and we denote $\delta_\beta f(s, x) = f(s + \beta, x) - f(s, x)$. We write

$$\begin{aligned} g(t) - g(s) &= \int_0^s \delta_a \partial_t \mathbf{K}(s - \tau) * f(\tau) d\tau + \int_s^t \partial_t \mathbf{K}(t - \tau) * f(\tau) d\tau \\ &= \int_0^s \delta_a \partial_t \mathbf{K}(s - \tau) * (f(\tau) - f(s)) d\tau + \int_s^t \partial_t \mathbf{K}(t - \tau) * (f(\tau) - f(t)) d\tau \\ &\quad + \left(\int_0^s \delta_a \partial_t \mathbf{K}(s - \tau) * f(s) d\tau + \int_s^t \partial_t \mathbf{K}(t - \tau) * f(t) d\tau \right) \\ &:= I_1 + I_2 + I_3. \end{aligned}$$

Applying Lemma 2.1, we have

$$\begin{aligned} \|I_1\|_{L_x^p} &\lesssim \int_0^s \min\left\{1, \frac{a}{s - \tau}\right\} \frac{1}{(s - \tau)^{1-\alpha} \tau^{\sigma+\alpha}} d\tau \|f\|_{X_T^{\sigma,p}} \\ &\lesssim a^\alpha s^{-\sigma-\alpha} \|f\|_{X_T^{\sigma,p}}, \end{aligned}$$

where we use the fact that

$$\begin{aligned} & \int_0^s \min\{1, \frac{a}{s-\tau}\} \frac{1}{(s-\tau)^{1-\alpha} \tau^{\sigma+\alpha}} d\tau \\ & \lesssim a^\alpha \int_0^{\frac{s}{2}} \frac{1}{(s-\tau)^{\sigma+\alpha}} d\tau + s^{-\sigma-\alpha} \int_{\frac{s}{2}}^s \min\{1, \frac{a}{s-\tau}\} \frac{1}{(s-\tau)^{1-\alpha}} d\tau \lesssim a^\alpha s^{-\sigma-\alpha}. \end{aligned} \quad (2.3)$$

For I_2 , we have

$$\|I_2\|_{L^p} \lesssim \int_s^t \frac{1}{(t-\tau)^{1-\alpha}} \frac{1}{\tau^{\sigma+\alpha}} d\tau \|f\|_{X_T^{\sigma,p}} \lesssim a^\alpha s^{-\sigma-\alpha} \|f\|_{X_T^{\sigma,p}}.$$

For I_3 , we have

$$\begin{aligned} \|I_3\|_{L^p} & \lesssim \|f(s) * \mathbf{K}(t-s) - f(s) * \mathbf{K}(t) - f(s) + f(s) * \mathbf{K}(s) + f(t) - f(t) * \mathbf{K}(t-s)\|_{L^p} \\ & \lesssim \|f(t) - f(s)\|_{L^p} + \|f(s) * (\mathbf{K}(t) - \mathbf{K}(s))\|_{L^p} \lesssim s^{-\sigma-\alpha} (t-s)^\alpha \|f\|_{X_T^{\sigma,p}}, \end{aligned}$$

where we applied Lemma 2.1 to get $\|\mathbf{K}(t) - \mathbf{K}(s)\|_{L^1} \lesssim \frac{(t-s)^\alpha}{s^\alpha}$ in the last inequality. Then we obtain that

$$\sup_{0 < s < t < T} s^{1-\gamma+\alpha} \frac{\|g(t) - g(s)\|_{L^p}}{(t-s)^\alpha} \lesssim \|f\|_{X_T^{\sigma,p}}. \quad (2.4)$$

By (2.2) and (2.4), we get (2.1). The proof is complete. $\blacksquare$

Remark 2.4 The result in Lemma 2.3 is still true for $g(t, x) = \int_0^t \int_{\mathbb{R}} \partial_t \mathbf{G}(t-\tau, x-y, x) f(\tau, y) dy$, where the kernel $\mathbf{G}(t, z, x)$ satisfies

$$\begin{aligned} \sup_z |\partial_t \mathbf{G}(t, z, x)| & \lesssim |\partial_t \mathbf{K}(\zeta t, x)|, \\ \sup_z |\partial_t \mathbf{G}(t, z, x) - \partial_t \mathbf{G}(s, z, x)| & \lesssim |\partial_t \mathbf{K}(\zeta s, x) - \partial_t \mathbf{K}(\zeta t, x)|, \end{aligned}$$

for a universal constant $\zeta > 0$.

Lemma 2.5 Let $0 \leq \beta < 1 - \alpha$; and $\tilde{\sigma} < 1$, $1 \leq q \leq p \leq \infty$, $1 + \frac{1}{p} = \frac{1}{r} + \frac{1}{q}$, $1 + \sigma - \tilde{\sigma} - \beta \geq 0$. Suppose that $G(t, y, x)$ satisfies

$$\sup_x \|G(t, \cdot, x)\|_{L^r} \lesssim t^{-\beta}, \quad \sup_x \|G(t, \cdot, x) - G(s, \cdot, x)\|_{L^r} \lesssim s^{-\beta} \min\{1, \frac{t-s}{s}\}, \quad (2.5)$$

for any $0 < s < t < T$. Then for $g(t, x) = \int_0^t \int G(t-\tau, x-y, x) f(\tau, y) dy d\tau$, we have

$$\|g\|_{X_T^{\sigma,p}} \lesssim T^{1+\sigma-\tilde{\sigma}-\beta} \|f\|_{X_T^{\tilde{\sigma},q}}.$$

Proof. By Young's inequality,

$$\begin{aligned} \|g(t)\|_{L^p} & \lesssim \int_0^t \sup_x \|G(t-\tau, \cdot, x)\|_{L^r} \|f(\tau)\|_{L^q} d\tau \\ & \lesssim \int_0^t \frac{1}{(t-\tau)^\beta} \frac{1}{\tau^{\tilde{\sigma}}} d\tau \|f\|_{X_T^{\tilde{\sigma},q}} \lesssim t^{1-\beta-\tilde{\sigma}} \|f\|_{X_T^{\tilde{\sigma},q}}. \end{aligned}$$

Hence

$$\sup_{t \in [0, T]} t^\sigma \|g(t)\|_{L^p} \lesssim T^{1-\beta+\sigma-\tilde{\sigma}} \|f\|_{X_T^{\tilde{\sigma},q}}. \quad (2.6)$$

Moreover, for any $0 < s < t < T$, denote $a = t - s$,

$$g(t) - g(s) = \int_0^s \int_{\mathbb{R}} \delta_a G(s - \tau, x - y, x) f(\tau, y) dy d\tau + \int_s^t \int_{\mathbb{R}} G(t - \tau, x - y, x) f(\tau, y) dy d\tau.$$

By (2.5) and Young's inequality, one has

$$\begin{aligned} \|g(t) - g(s)\|_{L^p} &\lesssim \|f\|_{X_T^{\tilde{\sigma}, q}} \int_0^s \frac{\min\{1, \frac{a}{s-\tau}\}}{(s-\tau)^\beta} \frac{1}{\tau^{\tilde{\sigma}}} d\tau + \|f\|_{X_T^{\tilde{\sigma}, q}} \int_s^t \frac{1}{(t-\tau)^\beta} \frac{1}{\tau^{\tilde{\sigma}}} d\tau \\ &\stackrel{(2.3)}{\lesssim} a^\alpha s^{-\sigma-\alpha} T^{1-\beta+\sigma-\tilde{\sigma}} \|f\|_{X_T^{\tilde{\sigma}, q}}. \end{aligned}$$

Combining this with (2.6), we obtain the result. ■

Lemma 2.6 *Suppose*

$$g(t, x) = \int_0^t \int_{\mathbb{R}} \mathbf{K}(t - \tau, x - y) R(\tau, y) dy d\tau.$$

Let $l = 0, 1$. Then for any $\frac{1}{1+2\alpha} < p < \frac{1}{2\alpha}$ and $\sigma \geq \frac{1+l}{2} - \frac{1}{2p}$, there holds

$$\|\partial_x^l g\|_{X_T^{\sigma, p}} \lesssim T^{\sigma - \frac{1+l}{2} + \frac{1}{2p}} (\|R\|_{L_T^1} + \|R\|_{X_T^{1,1}}).$$

Proof. Note that

$$\partial_x^l g(t, x) = \int_0^t \int_{\mathbb{R}} \partial_x^l \mathbf{K}(t - \tau, x - y) R(\tau, y) dy d\tau.$$

By Young's inequality and Lemma 2.1,

$$\|\partial_x^l g(t)\|_{L^p} \lesssim \int_0^t \|\partial_x^l \mathbf{K}(t - \tau)\|_{L^p} \|R(\tau)\|_{L^1} d\tau \lesssim \int_0^t (t - \tau)^{\frac{1}{2p} - \frac{1+l}{2}} \|R(\tau)\|_{L^1} d\tau.$$

We divide the integral into two parts. When $\tau \in [0, t/2]$, we have $(t - \tau)^{\frac{1}{2p} - \frac{1+l}{2}} \sim t^{\frac{1}{2p} - \frac{1+l}{2}}$. And when $\tau \in [t/2, t]$, one has $\|R(\tau)\|_{L^1} \lesssim t^{-1} \|R\|_{X_T^{1,1}}$. This yields that

$$\|\partial_x^l g(t)\|_{L^p} \lesssim t^{\frac{1}{2p} - \frac{1+l}{2}} (\|R\|_{L_T^1} + \|R\|_{X_T^{1,1}}). \quad (2.7)$$

It remains to check the Hölder norms. Let $0 < s < t < T$ and $a = t - s$, one has

$$\begin{aligned} &\partial_x^l g(t, x) - \partial_x^l g(s, x) \\ &= \int_0^s \int_{\mathbb{R}} \delta_a \partial_x^l \mathbf{K}(s - \tau, x - y) R(\tau, y) dy d\tau + \int_s^t \int_{\mathbb{R}} \partial_x^l \mathbf{K}(t - \tau, x - y) R(\tau, y) dy d\tau := \mathcal{R}_1 + \mathcal{R}_2. \end{aligned}$$

By Young's inequality and Lemma 2.1,

$$\|\mathcal{R}_1\|_{L^p} \lesssim \int_0^s \|\delta_a \partial_x^l \mathbf{K}(s - \tau)\|_{L^p} \|R(\tau)\|_{L^1} d\tau \lesssim \int_0^s (s - \tau)^{\frac{1}{2p} - \frac{1+l}{2}} \min\{1, \frac{a}{s - \tau}\} \|R(\tau)\|_{L^1} d\tau.$$

We discuss $\tau \in [0, s/2]$ and $\tau \in [s/2, s]$ separately and get that

$$\|\mathcal{R}_1\|_{L^p} \lesssim a^\alpha s^{\frac{1}{2p} - \frac{1+l}{2} - \alpha} \int_0^{s/2} \|R(\tau)\|_{L^1} d\tau + s^{-1} \|R\|_{L_T^1} \int_{s/2}^s (s - \tau)^{\frac{1}{2p} - \frac{1+l}{2}} \min\{1, \frac{a}{s - \tau}\} d\tau.$$

By a change of variable, we can write

$$\begin{aligned} &\int_{s/2}^s (s - \tau)^{\frac{1}{2p} - \frac{1+l}{2}} \min\{1, \frac{a}{s - \tau}\} d\tau = \int_0^{s/2} \tau^{\frac{1}{2p} - \frac{1+l}{2}} \min\{1, \frac{a}{\tau}\} d\tau \\ &\lesssim \mathbf{1}_{a \leq s/2} \left(\int_0^a \tau^{\frac{1}{2p} - \frac{1+l}{2}} d\tau + a \int_a^{s/2} \tau^{\frac{1}{2p} - \frac{1+l}{2} - 1} d\tau \right) + \mathbf{1}_{a \geq s/2} \int_0^s \tau^{\frac{1}{2p} - \frac{1+l}{2}} d\tau \\ &\lesssim \mathbf{1}_{a \leq s/2} a^{\frac{1}{2p} + \frac{1-l}{2}} + \mathbf{1}_{a \geq s/2} s^{\frac{1}{2p} + \frac{1-l}{2}} \lesssim a^\alpha s^{\frac{1}{2p} + \frac{1-l}{2} - \alpha}, \end{aligned}$$

provided $\frac{1}{2p} + \frac{1+l}{2} - \alpha \geq 0$. Hence

$$\|\mathcal{R}_1\|_{L^p} \lesssim a^\alpha s^{\frac{1}{2p} - \frac{1+l}{2} - \alpha} (\|R\|_{L_T^1} + \|R\|_{X_T^{1,1}}). \quad (2.8)$$

Then we deal with $\mathcal{R}_2$. By Young's inequality,

$$\|\mathcal{R}_2\|_{L^p} \lesssim \int_s^t \|\partial_x^l \mathbf{K}(t-\tau)\|_{L^p} \|R(\tau)\|_{L^1} d\tau \lesssim \int_s^t (t-\tau)^{\frac{1}{2p} - \frac{1+l}{2}} \|R(\tau)\|_{L^1} d\tau.$$

We consider two cases. If $s < t/2$, then one has $a \sim t$. Hence

$$\int_s^t (t-\tau)^{\frac{1}{2p} - \frac{1+l}{2}} \|R(\tau)\|_{L^1} d\tau \lesssim t^{\frac{1}{2p} - \frac{1+l}{2}} (\|R\|_{L_T^1} + \|R\|_{X_T^{1,1}}) \lesssim a^\alpha s^{\frac{1}{2p} - \frac{1+l}{2} - \alpha} (\|R\|_{L_T^1} + \|R\|_{X_T^{1,1}}),$$

provided $\frac{1}{2p} - \frac{1+l}{2} - \alpha < 0$.

If $s \geq t/2$, we can write

$$\begin{aligned} \int_s^t (t-\tau)^{\frac{1}{2p} - \frac{1+l}{2}} \|R(\tau)\|_{L^1} d\tau &\lesssim \int_s^t (t-\tau)^{\frac{1}{2p} - \frac{1+l}{2}} d\tau t^{-1} \|R\|_{X_T^{1,1}} \lesssim a^{\frac{1}{2p} + \frac{1-l}{2}} t^{-1} \|R\|_{X_T^{1,1}} \\ &\lesssim a^\alpha s^{\frac{1}{2p} - \frac{1+l}{2} - \alpha} \|R\|_{X_T^{1,1}}. \end{aligned}$$

We conclude that

$$\|\mathcal{R}_2\|_{L^p} \lesssim a^\alpha s^{\frac{1}{2p} - \frac{1+l}{2} - \alpha} (\|R\|_{L_T^1} + \|R\|_{X_T^{1,1}}).$$

Combining this with (2.8) yields that

$$\|\partial_x^l g(t) - \partial_x^l g(s)\|_{L^p} \lesssim (t-s)^\alpha s^{\frac{1}{2p} - \frac{1+l}{2} - \alpha} (\|R\|_{L_T^1} + \|R\|_{X_T^{1,1}}). \quad (2.9)$$

Then we complete the proof in view of (2.7) and (2.9). $\blacksquare$

Remark 2.7 The result in Lemma 2.6 is still true for $g(t, x) = \int_0^t \int \mathbf{G}(t-\tau, x, x-y) R(\tau, y) dy d\tau$, where $\mathbf{G}(t, x, y)$ satisfies that

$$\begin{aligned} \sup_z \|\partial_2^l \mathbf{G}(t, z, \cdot)\|_{L^p} &\lesssim t^{-\frac{1+l}{2} + \frac{1}{2p}}, \\ \sup_z \|\partial_2^l \mathbf{G}(t, z, \cdot) - \partial_2^l \mathbf{G}(s, z, \cdot)\|_{L^p} &\lesssim s^{-\frac{1+l}{2} + \frac{1}{2p}} \min\{1, \frac{t-s}{s}\}, \text{ for } l = 0, 1, p \in [1, \infty]. \end{aligned}$$

3 Main lemma and its proof

We consider the system (1.2) and the system (1.3) with jump initial data v_0 . Observe that the equation for v is hyperbolic, $v(t)$ will have jump at some points for any $t > 0$. To deal with u in (1.2) and (u, θ) in (1.3), we need to consider the following parabolic equation with jump coefficient.

$$\begin{aligned} \partial_t f(t, x) - \partial_x(\phi(x) \partial_x f(t, x)) &= \partial_x F(t, x) + R(t, x), \\ f(0, x) &= f_0(x). \end{aligned} \quad (3.1)$$

Here the coefficient function satisfies $0 < c_0 \leq \phi(x) \leq c_0^{-1}, \forall x \in \mathbb{R}$, and there exist $\varepsilon > 0$, $\{a_n\}_{n=1}^N \subset \mathbb{R}$ satisfying

$$\begin{aligned} \min_{i \neq j} |a_i - a_j| &\geq 10\varepsilon, \quad \|\phi'\|_{L^\infty(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} \leq C_\phi, \\ \phi(x) &= \begin{cases} c_n^+, & x \in [a_n, a_n + \varepsilon], \\ c_n^-, & x \in [a_n - \varepsilon, a_n], \end{cases} \quad n = 1, \dots, N, \end{aligned} \quad (3.2)$$

for some $\{c_n^\pm\}_{n=1}^N \subset \mathbb{R}^+$. Here we denote $\mathbb{I}_\varepsilon = \cup_{n=1}^N [a_n - \varepsilon, a_n + \varepsilon]$.

Lemma 3.1 *Let f be a solution to (3.1) with initial data $f_0 = \partial_x \bar{f}_0$. There exists $T^* = T^*(c_0, C_\phi, \varepsilon) > 0$ such that for any $0 < T < T^*$,*

$$\sum_{\star \in \{L_T^2, X_T^{1/2,2}\}} \|f\|_\star + \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6}, \frac{6}{5}}\}} \|\partial_x f\|_\star \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}} + \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6}, \frac{6}{5}}\}} \|F\|_\star + T^{\frac{1}{4}} \sum_{\star \in \{L_T^1, X_T^{1,1}\}} \|R\|_\star.$$

Moreover, if $R = 0$, we have

$$\begin{aligned} \|\partial_x f\|_{X_T^{1-\gamma, \infty}} &\lesssim \|\bar{f}_0\|_{\dot{C}^{2\gamma}} + \|F\|_{X_T^{1-\gamma, \infty}}, \\ \|\partial_x f\|_{L_T^2} + \|\partial_x f\|_{X_T^{\frac{1}{2}, 2}} + \|\partial_x f\|_{X_T^{\frac{3}{4}, \infty}} &\lesssim \|f_0\|_{L^2} + \|F\|_{L_T^2} + \|F\|_{X_T^{\frac{1}{2}, 2}} + \|F\|_{X_T^{\frac{3}{4}, \infty}}. \end{aligned}$$

Proof. Applying Lemma 3.7-Lemma 3.10 to obtain that there exists $T_0 = T_0(C_\phi, \varepsilon) > 0$ such that for any $0 < T \leq T_0$,

$$\begin{aligned} &\sum_{\star \in \{L_T^2, X_T^{\frac{1}{2}, 2}\}} \|f\|_\star + \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6}, \frac{6}{5}}\}} \|\partial_x f\|_\star \\ &\leq \mathbf{C} \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}} + \mathbf{C} \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6}, \frac{6}{5}}\}} \|F\|_\star + \mathbf{C} T^{\frac{1}{4}} \sum_{\star \in \{L_T^1, X_T^{1,1}\}} \|R\|_\star + \frac{1}{5} (\|f\|_{L_T^2} + \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6}, \frac{6}{5}}\}} \|\partial_x f\|_\star), \end{aligned}$$

and, if $R = 0$,

$$\begin{aligned} \|\partial_x f\|_{X_T^{1-\gamma, \infty}} &\leq \mathbf{C} (\|\bar{f}_0\|_{\dot{C}^{2\gamma}} + \|F\|_{X_T^{1-\gamma, \infty}} + T^{\frac{1}{3}} \|\partial_x f\|_{X_T^{1-\gamma, \infty}}), \\ \|f\|_{L_T^6} + \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2}, 2}, X_T^{\frac{3}{4}, \infty}\}} \|\partial_x f\|_\star &\leq \mathbf{C} \|f_0\|_{L^2} + \mathbf{C} \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2}, 2}, X_T^{\frac{3}{4}, \infty}\}} \|F\|_\star \\ &\quad + \frac{1}{10} (\|f\|_{L_T^6} + \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2}, 2}, X_T^{\frac{3}{4}, \infty}\}} \|\partial_x f\|_\star). \end{aligned}$$

where a constant $\mathbf{C} > 0$ depends only on c_0 , C_ϕ and ε . This implies the results by taking $0 < T^* = \min\{\frac{1}{(\mathbf{C}+1)^{10}}, T_0\}$. Then we complete the proof. $\blacksquare$

We first introduce some elementary lemmas that will be used frequently in our proof.

Lemma 3.2 *Let $g(t, x) = \int_{\mathbb{R}} \partial_x \mathbf{K}(t, x-y) \bar{g}_0(y) dy$. Then for any $T > 0$,*

$$\begin{aligned} \|\partial_x g\|_{X_T^{1-\gamma, \infty}} &\lesssim \|\bar{g}_0\|_{\dot{C}^{2\gamma}}, \\ \|\partial_x g\|_{X_T^{\frac{1}{2}, 2}} + \|\partial_x g\|_{X_T^{\frac{3}{4}, \infty}} &\lesssim \|\partial_x \bar{g}_0\|_{L^2}, \\ \|g\|_{X_T^{\frac{1}{2}, 2}} &\lesssim \|\bar{g}_0\|_{L^2}, \quad \|\partial_x g\|_{X_T^{\frac{5}{6}, \frac{6}{5}}} \lesssim \|\bar{g}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}}. \end{aligned}$$

Proof. Observe that $\int_{\mathbb{R}} \partial_x^2 \mathbf{K}(t, x) dx = 0$. Hence

$$\partial_x g(t, x) = \int_{\mathbb{R}} \partial_x^2 \mathbf{K}(t, x-y) (\bar{g}_0(y) - \bar{g}_0(x)) dy = - \int_{\mathbb{R}} \partial_x^2 \mathbf{K}(t, z) \Delta_z \bar{g}_0(x) dz,$$

where we denote $\Delta_z \bar{g}_0(x) = \bar{g}_0(x) - \bar{g}_0(x-z)$. Note that $|\Delta_z \bar{g}_0(x)| \leq |z|^{2\gamma} \|\bar{g}_0\|_{\dot{C}^{2\gamma}}$. Hence, by Lemma 2.1, we have

$$\|\partial_x g(t)\|_{L^\infty} \lesssim \int_{\mathbb{R}} |\partial_x^2 \mathbf{K}(t, x)| |x|^{2\gamma} dx \|\bar{g}_0\|_{\dot{C}^{2\gamma}} \lesssim t^{\gamma-1} \|\bar{g}_0\|_{\dot{C}^{2\gamma}}.$$

Moreover, for any $0 < s < t < T$,

$$\begin{aligned} \|\partial_x g(t) - \partial_x g(s)\|_{L^\infty} &\lesssim \int_{\mathbb{R}} |\partial_x^2 \mathbf{K}(t, x) - \partial_x^2 \mathbf{K}(s, x)| |x|^{2\gamma} dx \|\bar{g}_0\|_{\dot{C}^{2\gamma}} \\ &\lesssim s^{\gamma-1} \min\{1, \frac{t-s}{s}\} \|\bar{g}_0\|_{\dot{C}^{2\gamma}} \lesssim (t-s)^\alpha s^{\gamma-1-\alpha} \|\bar{g}_0\|_{\dot{C}^{2\gamma}}. \end{aligned}$$

Hence

$$\|\partial_x g\|_{X_T^{1-\gamma,\infty}} \lesssim \|\bar{g}_0\|_{\dot{C}^{2\gamma}}. \quad (3.3)$$

By Lemma 2.1 and Young's inequality, we have

$$\begin{aligned} t^{\frac{1}{2}} \|\partial_x g(t)\|_{L^2} + t^{\frac{3}{4}} \|\partial_x g\|_{L^\infty} &\lesssim (t^{\frac{1}{2}} \|\partial_x \mathbf{K}(t)\|_{L^1} + t^{\frac{3}{4}} \|\partial_x \mathbf{K}(t)\|_{L^2}) \|\partial_x \bar{g}_0\|_{L^2} \lesssim \|\partial_x \bar{g}_0\|_{L^2}, \\ t^{\frac{1}{2}} \|g(t)\|_{L^2} &\lesssim t^{\frac{1}{2}} \|\partial_x \mathbf{K}(t)\|_{L^1} \|\bar{g}_0\|_{L^2} \lesssim \|\bar{g}_0\|_{L^2}, \end{aligned} \quad (3.4)$$

and

$$\begin{aligned} s^{\frac{1}{2}+\alpha} \|\partial_x g(t) - \partial_x g(s)\|_{L^2} + s^{\frac{3}{4}+\alpha} \|\partial_x g(t) - \partial_x g(s)\|_{L^\infty} \\ \lesssim (s^{\frac{1}{2}+\alpha} \|\partial_x K(t) - \partial_x K(s)\|_{L^1} + s^{\frac{3}{4}+\alpha} \|\partial_x K(t) - \partial_x K(s)\|_{L^2}) \|\partial_x \bar{g}_0\|_{L^2} \\ \lesssim (t-s)^\alpha \|\partial_x \bar{g}_0\|_{L^2}. \end{aligned} \quad (3.5)$$

Similarly, it is easy to check that

$$s^{\frac{1}{2}+\alpha} \|g(t) - g(s)\|_{L^2} \lesssim (t-s)^\alpha \|\bar{g}_0\|_{L^2}. \quad (3.6)$$

By (3.3)-(3.6), one has

$$\|\partial_x g\|_{X_T^{1-\gamma,\infty}} \lesssim \|\bar{g}_0\|_{\dot{C}^{2\gamma}}, \quad \|\partial_x g\|_{X_T^{\frac{1}{2},2}} + \|\partial_x g\|_{X_T^{\frac{3}{4},\infty}} \lesssim \|\partial_x \bar{g}_0\|_{L^2}, \quad \|g\|_{X_T^{\frac{1}{2},2}} \lesssim \|\bar{g}_0\|_{L^2}. \quad (3.7)$$

It remains to estimate $\|\partial_x g\|_{X_T^{\frac{5}{6},\frac{6}{5}}}$. By Hölder's inequality and Lemma 2.1,

$$\begin{aligned} \|\partial_x g(t)\|_{L^{\frac{6}{5}}} &\lesssim \int_{\mathbb{R}} |\partial_x^2 \mathbf{K}(t, z)| \|\Delta_z \bar{f}_0\|_{L^{\frac{6}{5}}} dz \lesssim \left(\int_{\mathbb{R}} |\partial_x^2 \mathbf{K}(t, z)|^6 |z|^7 dz \right)^{\frac{1}{6}} \left(\int_{\mathbb{R}} \frac{\|\Delta_z \bar{f}_0\|_{L^{\frac{6}{5}}}^{6/5}}{|z|^{7/5}} dz \right)^{5/6} \\ &\lesssim t^{-\frac{5}{6}} \|\bar{g}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}}. \end{aligned}$$

Moreover, for any $0 < s < t < T$,

$$\begin{aligned} \|\partial_x g(t) - \partial_x g(s)\|_{L^{\frac{6}{5}}} &\lesssim \int_{\mathbb{R}} |\partial_x^2 \mathbf{K}(t, z) - \partial_x^2 \mathbf{K}(s, z)| \|\Delta_z \bar{f}_0\|_{L^{\frac{6}{5}}} dz \\ &\lesssim \left(\int_{\mathbb{R}} |\partial_x^2 \mathbf{K}(t, z) - \partial_x^2 \mathbf{K}(s, z)|^6 |z|^7 dz \right)^{\frac{1}{6}} \left(\int_{\mathbb{R}} \frac{\|\Delta_z \bar{f}_0\|_{L^{\frac{6}{5}}}^{6/5}}{|z|^{7/5}} dz \right)^{5/6} \\ &\lesssim s^{-\frac{5}{6}-\alpha} (t-s)^\alpha \|\bar{g}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}}. \end{aligned}$$

Then we get

$$\|\partial_x g\|_{X_T^{\frac{5}{6},\frac{6}{5}}} \lesssim \|\bar{g}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}}.$$

Combining this with (3.7), we complete the proof. $\blacksquare$

Remark 3.3 The result in Lemma 3.2 is still true for $g(t, x) = \int_{\mathbb{R}} \partial_y \mathbf{G}(t, x - y, x) \bar{g}_0(y) dy$, where $\mathbf{G}(t, x, z)$ satisfies that $\int \partial_x^2 \mathbf{G}(t, x, z) dx = 0$, and

$$\begin{aligned} \max_{l=0,1} \sup_z |\partial_x^m \partial_z^l \mathbf{G}(t, z, x)| &\lesssim |\partial_x^m \mathbf{K}(\zeta t, x)|, \\ \max_{l=0,1} \sup_z |\partial_x^m \partial_z^l \mathbf{G}(t, z, x) - \partial_x^m \partial_z^l \mathbf{G}(s, z, x)| &\lesssim |\partial_x^m \mathbf{K}(\zeta s, x) - \partial_x^m \mathbf{K}(\zeta t, x)|, \quad m = 1, 2, \end{aligned}$$

for a universal constant $\zeta > 0$.

The following is the Hardy's inequality.

Lemma 3.4 For any non-negative measurable function $h : \mathbb{R} \rightarrow \mathbb{R}^+$, there holds for any $p \in (1, \infty)$,

$$\int_0^\infty \left(\int_0^\infty h(x) \frac{dx}{x+y} \right)^p dy \lesssim \int_0^\infty h(x)^p dx.$$

Proof. By Hardy's inequality [3],

$$\begin{aligned} \int_0^\infty \left(\int_0^\infty h(x) \frac{dx}{x+y} \right)^p dy &\leq \int_0^\infty \left(\int_0^y h(x) dx \right)^p \frac{dy}{y^p} + \int_0^\infty \left(\int_y^\infty h(x) \frac{dx}{x} \right)^p dy \\ &\leq \left(\frac{p^p}{(p-1)^p} + p^p \right) \int_0^\infty h(x)^p dx. \end{aligned}$$

This completes the proof. ■

The following lemma helps us to control boundary terms in (3.10).

Lemma 3.5 *Let $r \in [1, +\infty)$, $p, q \in (1, +\infty)$ be such that $\frac{1}{p} + 1 = \frac{1}{q} + \frac{1}{r}$. Suppose that $\tilde{G}(t, x)$ satisfies*

$$\int_0^\infty |\tilde{G}(t, x)|^r dt \lesssim \frac{1}{|x|}.$$

Then for

$$g(t, x) = \mathbf{1}_{x \geq 0} \int_0^t \int_0^{+\infty} \tilde{G}(t - \tau, x + y) f(\tau, y) dy d\tau,$$

there holds

$$\|g\|_{L_T^p} \lesssim \|f\|_{L_T^q}.$$

Proof. By Young's inequality,

$$\begin{aligned} \|g\|_{L_T^p} &\lesssim \left\| \int_0^\infty \left(\int_0^t |\tilde{G}(t, x + y)|^r dt \right)^{\frac{1}{r}} \|f(y)\|_{L_t^q} dy \right\|_{L_x^p(\mathbb{R}^+)} \\ &\lesssim \left\| \int_0^\infty \|f(y)\|_{L_t^q} \frac{dy}{|x + y|^{\frac{1}{r}}} \right\|_{L_x^p(\mathbb{R}^+)}. \end{aligned}$$

When $r > 1$, then by Young's inequality, we obtain that

$$\left\| \int_0^\infty \|f(y)\|_{L_{t,T}^q} \frac{dy}{|x + y|^{\frac{1}{r}}} \right\|_{L_x^p(\mathbb{R}^+)} \lesssim \|f\|_{L_T^q} \| |x|^{-\frac{1}{r}} \|_{L_w^r} \lesssim \|f\|_{L_T^q}.$$

When $r = 1$, we obtain the result by Lemma 3.4. This completes the proof. ■

3.1 Estimates near jump points

We first study estimates near jump points. Let $\mathbb{I}_\varepsilon^n = [a_n - \varepsilon, a_n + \varepsilon]$. Without loss of generality, we assume $a_n = 0$. Then we can write equation (3.1) as

$$\begin{aligned} \partial_t f - \partial_x(\bar{\phi} f_x) &= \partial_x \tilde{F} + R, \\ f(0, x) &= f_0(x), \end{aligned} \tag{3.8}$$

where we denote $\tilde{F} = F + (\phi - \bar{\phi})f_x$ with $\bar{\phi} = \mathbf{1}_{x \leq 0} c_n^- + \mathbf{1}_{x > 0} c_n^+$. For simplicity, we omit the subscript n in our proof and denote $c^\pm = c_n^\pm$.

By Lemma 6.2, the solution of (3.8) has formula $f = f^+ \mathbf{1}_{x \geq 0} + f^- \mathbf{1}_{x < 0}$, where

$$f^\pm(t, x) = f_L^\pm + f_N^\pm + f_R^\pm. \tag{3.9}$$

Here $f_L^\pm = f_1^\pm + f_{1,B}^\pm$, $f_N^\pm = f_2^\pm + f_{2,B}^\pm$, $f_R^\pm = f_3^\pm + f_{3,B}^\pm$, with

$$\begin{aligned} f_1^\pm(t, x) &= \int_{\mathbb{R}^\pm} (\mathbf{K}(c_\pm t, x - y) - \mathbf{K}(c_\pm t, x + y)) f_0(y) dy, \\ f_2^\pm(t, x) &= \int_0^t \int_{\mathbb{R}^\pm} (\mathbf{K}(c_\pm(t - \tau), x - y) - \mathbf{K}(c_\pm(t - \tau), x + y)) \partial_x \tilde{F}(\tau, y) dy d\tau, \\ f_3^\pm(t, x) &= \int_0^t \int_{\mathbb{R}^\pm} (\mathbf{K}(c_\pm(t - \tau), x - y) - \mathbf{K}(c_\pm(t - \tau), x + y)) R(\tau, y) dy d\tau, \\ f_{m,B}^\pm &= \frac{-2\sqrt{c_\pm}}{\sqrt{c_+} + \sqrt{c_-}} \int_0^t \mathbf{K}(c_\pm(t - \tau), x) (c_+ \partial_x f_m^+(\tau, 0) - c_- \partial_x f_m^-(\tau, 0)) d\tau, \quad m = 1, 2, 3. \end{aligned}$$

Here $f_{m,B}$ means these are boundary terms. Denote $\beta_m(\tau, y) = c_+ f_m^+(\tau, y) + c_- f_m^-(\tau, -y)$, then

$$\begin{aligned} f_{m,B}^+ &= C_+ \int_0^t \int_0^\infty \partial_y (\mathbf{K}(c_+(t - \tau), x + y) \partial_x \beta_m(\tau, y)) dy d\tau \\ &= -C_+ \int_0^t \int_0^\infty \partial_x^2 \mathbf{K}(c_+(t - \tau), x + y) \beta_m(\tau, y) dy d\tau \\ &\quad + C_+ \int_0^t \int_0^\infty \mathbf{K}(c_+(t - \tau), x + y) \partial_x^2 \beta_m(\tau, y) dy d\tau, \end{aligned}$$

where we denote $C_\pm = \frac{-2\sqrt{c_\pm}}{\sqrt{c_+} + \sqrt{c_-}}$. Note that

$$\begin{aligned} \partial_x^2 \beta_1(\tau, y) &= \partial_t f_1^+(\tau, y) + \partial_t f_1^-(\tau, -y), \\ \partial_x^2 \beta_2(\tau, y) &= \partial_t f_2^+(\tau, y) - \partial_x \tilde{F}(\tau, y) + \partial_t f_2^-(\tau, -y) - \partial_x \tilde{F}(\tau, -y), \\ \partial_x^2 \beta_3(\tau, y) &= \partial_t f_3^+(\tau, y) - R(\tau, y) + \partial_t f_3^-(\tau, -y) - R(\tau, -y). \end{aligned}$$

Hence,

$$\begin{aligned} f_{1,B}^+(t, x) &= C_+(c_+ - c_-) \int_0^t \int_0^\infty \partial_x^2 \mathbf{K}(c_+(t - \tau), x + y) f_1^-(\tau, -y) dy d\tau \\ &\quad - C_+ \int_0^\infty \mathbf{K}(c_+ t, x + y) (f_0(y) + f_0(-y)) dy, \\ f_{2,B}^+(t, x) &= C_+(c_+ - c_-) \int_0^t \int_0^\infty \partial_x^2 \mathbf{K}(c_+(t - \tau), x + y) f_2^-(\tau, -y) dy d\tau \\ &\quad - C_+ \int_0^t \int_0^\infty \mathbf{K}(c_+(t - \tau), x + y) ((\partial_x \tilde{F})(\tau, y) + (\partial_x \tilde{F})(\tau, -y)) dy d\tau, \\ f_{3,B}^+(t, x) &= C_+(c_+ - c_-) \int_0^t \int_0^\infty \partial_x^2 \mathbf{K}(c_+(t - \tau), x + y) f_3^-(\tau, -y) dy d\tau \\ &\quad - C_+ \int_0^t \int_0^\infty \mathbf{K}(c_+(t - \tau), x + y) (R(\tau, y) + R(\tau, -y)) dy. \end{aligned} \tag{3.10}$$

Remark 3.6 Note that we can recover the standard heat kernel from our formula when $c_+ = c_- = c$. More precisely, we have

$$\begin{aligned} f_L(t, x) &= \int_{\mathbb{R}} \mathbf{K}(ct, x - y) f_0(y) dy, \quad f_N(t, x) = \int_0^t \int_{\mathbb{R}} \mathbf{K}(c(t - \tau), x - y) \partial_x F(\tau, y) dy, \\ f_R(t, x) &= \int_0^t \int_{\mathbb{R}} \mathbf{K}(c(t - \tau), x - y) R(\tau, y) dy. \end{aligned}$$

The following lemma will help us to analyze the regularity of the solution near the jump points.

Lemma 3.7 *Let f be a solution to (3.1) with initial data $f_0 = \partial_x \bar{f}_0$. Then there exists $T > 0$ such that*

$$\|f\|_{X_T^{\frac{1}{2},2}(\mathbb{I}_\varepsilon)} + \|\partial_x f\|_{X_T^{\frac{5}{6},\frac{6}{5}}(\mathbb{I}_\varepsilon)} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}} + \|F\|_{X_T^{\frac{5}{6},\frac{6}{5}}} + T^{\frac{1}{3}}\|f_x\|_{X_T^{\frac{5}{6},\frac{6}{5}}} + T^{\frac{1}{4}}(\|R\|_{L_T^1} + \|R\|_{X_T^{1,1}}).$$

Moreover, if $R = 0$,

$$\begin{aligned} \|\partial_x f\|_{X_T^{1-\gamma,\infty}(\mathbb{I}_\varepsilon)} &\lesssim \|\bar{f}_0\|_{\dot{C}^{2\gamma}} + \|F\|_{X_T^{1-\gamma,\infty}} + T^{\frac{1}{3}}\|\partial_x f\|_{X_T^{1-\gamma,\infty}}, \\ \|\partial_x f\|_{X_T^{\frac{1}{2},2}(\mathbb{I}_\varepsilon)} + \|\partial_x f\|_{X_T^{\frac{3}{4},\infty}(\mathbb{I}_\varepsilon)} &\lesssim \|\partial_x \bar{f}_0\|_{L^2} + \|F\|_{X_T^{\frac{3}{4},\infty}} + T^{\frac{1}{3}}(\|\partial_x f\|_{X_T^{\frac{1}{2},2}} + \|\partial_x f\|_{X_T^{\frac{3}{4},\infty}}). \end{aligned}$$

Proof. Recall that $\mathbb{I}_\varepsilon = \cup_{n=1}^N \mathbb{I}_\varepsilon^n$. It suffices to consider one interval $\mathbb{I}_\varepsilon^n = [a_n - \varepsilon, a_n + \varepsilon]$. Without loss of generality, we assume $a_n = 0$. The solution has formula (3.9). Using integration by parts, one has

$$\begin{aligned} f_1^+(t, x) &= \int_{\mathbb{R}^+} (\partial_x \mathbf{K}(c_+ t, x - y) + \partial_x \mathbf{K}(c_+ t, x + y)) \bar{f}_0(y) dy \\ &= \int_{\mathbb{R}} \partial_x \mathbf{K}(c_+ t, x - y) \mathbf{f}_0(y) dy, \end{aligned}$$

where $\mathbf{f}_0(y)$ is the even extension of $\bar{f}_0 \mathbf{1}_{x \geq 0}$

$$\mathbf{f}_0(y) = \bar{f}_0(y) \mathbf{1}_{y \geq 0} + \bar{f}_0(-y) \mathbf{1}_{y < 0}.$$

It is easy to check that

$$\|\mathbf{f}_0\|_{\dot{C}^{2\gamma}} \leq \|\bar{f}_0\|_{\dot{C}^{2\gamma}}.$$

Applying Lemma 3.2 and Lemma 2.2 to get

$$\begin{aligned} \|\partial_x f_1^+\|_{X_T^{1-\gamma,\infty}(\mathbb{R}^+)} &\lesssim \|\mathbf{f}_0\|_{\dot{C}^{2\gamma}} \lesssim \|\bar{f}_0\|_{\dot{C}^{2\gamma}}, \\ \|\partial_x f_1^+\|_{X_T^{\frac{1}{2},2}(\mathbb{R}^+)} + \|\partial_x f_1^+\|_{X_T^{\frac{3}{4},\infty}(\mathbb{R}^+)} &\lesssim \|\partial_x \bar{f}_0\|_{L^2}, \quad \|f_1^+\|_{X_T^{\frac{1}{2},2}(\mathbb{R}^+)} \lesssim \|\bar{f}_0\|_{L^2}, \\ \|\partial_x f_1^+\|_{X_T^{\frac{5}{6},\frac{6}{5}}(\mathbb{R}^+)} &\lesssim \left(\int_{\mathbb{R}} \int_0^\infty |\mathbf{f}_0(x) - \mathbf{f}_0(x-z)|^{\frac{6}{5}} \frac{dx dz}{|z|^{\frac{7}{5}}} \right)^{5/6} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}}. \end{aligned} \tag{3.11}$$

Recall the formula of boundary term $f_{1,B}^+$ in (3.10), using integration by parts one has

$$\begin{aligned} f_{1,B}^+ &= C_+(c_+ - c_-) \int_0^t \int_0^\infty \partial_x^2 \mathbf{K}(c_+(t-\tau), x+y) f_1^-(\tau, -y) dy d\tau \\ &\quad - C_+ \int_0^\infty \partial_x \mathbf{K}(c_+ t, x+y) (\bar{f}_0(y) - \bar{f}_0(-y)) dy, \\ &:= I_1 + I_2. \end{aligned}$$

Applying Lemma 2.3 to obtain that for $(\sigma, p) \in \{(1-\gamma, \infty), (\frac{1}{2}, 2), (\frac{3}{4}, \infty), (\frac{5}{6}, \frac{6}{5})\}$,

$$\|\partial_x^l I_1\|_{X_T^{\sigma,p}} \lesssim \|\partial_x^l f_1\|_{X_T^{\sigma,p}}, \quad l = 1, 2.$$

Moreover, observe that

$$I_2 = -C_+ \int_{\mathbb{R}} \partial_x \mathbf{K}(c_+ t, x-y) \bar{\mathbf{f}}_0(y) dy,$$

where $\bar{\mathbf{f}}_0(y) = (\bar{f}_0(-y) - \bar{f}_0(y)) \mathbf{1}_{y \leq 0}$. Following the proof of Lemma 3.2 to yield

$$\begin{aligned} \|\partial_x I_2\|_{X_T^{1-\gamma,\infty}} &\lesssim \|\bar{\mathbf{f}}_0\|_{\dot{C}^{2\gamma}} \lesssim \|\bar{f}_0\|_{\dot{C}^{2\gamma}}, \\ \|\partial_x I_2\|_{X_T^{\frac{1}{2},2}} + \|\partial_x I_2\|_{X_T^{\frac{3}{4},\infty}} &\lesssim \|\partial_x (\bar{f}_0(-y) - \bar{f}_0(y)) \mathbf{1}_{y \leq 0}\|_{L^2} \lesssim \|\partial_x \bar{f}_0\|_{L^2}, \\ \|I_2\|_{X_T^{\frac{1}{2},2}} &\lesssim \|\bar{f}_0\|_{L^2} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}}, \end{aligned}$$

and

$$\|\partial_x I_2\|_{X_T^{\frac{5}{6}, \frac{6}{5}}(\mathbb{R}^+)} \lesssim \left(\int_{\mathbb{R}} \int_{\mathbb{R}^+} |\bar{f}_0(x) - \bar{f}_0(x-z)|^{\frac{6}{5}} \frac{dx dz}{|z|^{7/5}} \right)^{5/6} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{5}{6}}},$$

where we applied Lemma 2.2 in the last inequality. Similar arguments hold for $\partial_x f_{1,B}^-$. Hence

$$\begin{aligned} \|\partial_x f_{1,B}\|_{X_T^{1-\gamma, \infty}} &\lesssim \|\partial_x f_1\|_{X_T^{1-\gamma, \infty}} + \|\bar{f}_0\|_{\dot{C}^{2\gamma}}, \\ \|\partial_x f_{1,B}\|_{X_T^{\frac{1}{2}, 2}} + \|\partial_x f_{1,B}\|_{X_T^{\frac{3}{4}, \infty}} &\lesssim \|\partial_x f_1\|_{X_T^{\frac{1}{2}, 2}} + \|\partial_x f_1\|_{X_T^{\frac{3}{4}, \infty}} + \|\partial_x \bar{f}_0\|_{L^2}, \\ \|f_{1,B}\|_{X_T^{\frac{1}{2}, 2}} + \|\partial_x f_{1,B}\|_{X_T^{\frac{5}{6}, \frac{6}{5}}} &\lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}} + \|f_1\|_{X_T^{\frac{1}{2}, 2}} + \|\partial_x f_1\|_{X_T^{\frac{5}{6}, \frac{6}{5}}}. \end{aligned}$$

Combining this with (3.11), we obtain that

$$\begin{aligned} \|\partial_x f_L\|_{X_T^{1-\gamma, \infty}} &\lesssim \|\bar{f}_0\|_{\dot{C}^{2\gamma}}, \quad \|\partial_x f_L\|_{X_T^{\frac{1}{2}, 2}} + \|\partial_x f_L\|_{X_T^{\frac{3}{4}, \infty}} \lesssim \|\partial_x \bar{f}_0\|_{L^2}, \\ \|f_L\|_{X_T^{\frac{1}{2}, 2}} + \|\partial_x f_L\|_{X_T^{\frac{5}{6}, \frac{6}{5}}} &\lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}}. \end{aligned} \quad (3.12)$$

Then we estimate nonlinear term $f_N = f_2 + f_{2,B}$. We first estimate f_2 . By integration by parts,

$$\begin{aligned} f_2^\pm(t, x) &= \int_0^t \int_{\mathbb{R}^\pm} (\partial_x \mathbf{K}(c_\pm(t-\tau), x-y) + \partial_x \mathbf{K}(c_\pm(t-\tau), x+y)) F(\tau, y) dy d\tau \\ &\quad + \int_0^t \int_{\mathbb{R}^\pm} (\partial_x \mathbf{K}(c_\pm(t-\tau), x-y) + \partial_x \mathbf{K}(c_\pm(t-\tau), x+y)) (\phi - \bar{\phi})(y) f_x(\tau, y) dy d\tau \\ &= f_{2,1}^\pm(t, x) + f_{2,2}^\pm(t, x). \end{aligned} \quad (3.13)$$

By Lemma 2.3, we obtain

$$\|\partial_x f_{2,1}\|_{X_T^{\sigma, p}} \lesssim \|F\|_{X_T^{\sigma, p}}, \quad \|\partial_x f_{2,2}\|_{X_T^{\sigma, p}} \lesssim \|\partial_x f\|_{X_T^{\sigma, p}}. \quad (3.14)$$

And Lemma 2.5 implies that

$$\|f_{2,1}\|_{X_T^{\frac{1}{2}, 2}} \lesssim \|F\|_{X_T^{\frac{5}{6}, \frac{6}{5}}}. \quad (3.15)$$

Observe that $\phi(y) - \bar{\phi}(y) = 0$ for $y \in [-\varepsilon, \varepsilon]$. And

$$|\phi(y) - \bar{\phi}(y)| = |\phi(y) - \phi(x)| \lesssim C_\phi |x - y|, \quad (3.16)$$

for $x \in (0, \varepsilon), y \in (0, 4\varepsilon)$ or $x \in (-\varepsilon, 0), y \in (-4\varepsilon, 0)$. This implies a better estimate for $f_{2,2}$ when $x \in (-\varepsilon, \varepsilon)$. For simplicity, we only consider $f_{2,2}^+(t, x)$ with $x \in (0, \varepsilon)$, while $f_{2,2}^-(t, x)$ with $x \in (-\varepsilon, 0)$ can be done similarly. We can write

$$\partial_x f_{2,2}^+(t, x) = \int_0^t \int_{\mathbb{R}^+} G(t-\tau, x, y) f_x(\tau, y) dy d\tau,$$

where

$$G(t, x, y) = (\partial_x^2 \mathbf{K}(c_+ t, x-y) + \partial_x^2 \mathbf{K}(c_+ t, x+y)) (\phi - \bar{\phi})(y) \mathbf{1}_{x, y \in \mathbb{R}^+}.$$

For $x \in (0, \varepsilon)$, $0 < s < t < T$,

$$\begin{aligned} \int_{\mathbb{R}} |G(t, x, y)| dy &\lesssim C_\phi \int_{|y| \leq 4\varepsilon} |\partial_x^2 \mathbf{K}(c_+ t, y)| |y| dy + \int_{|y| \geq 4\varepsilon} |\partial_x^2 \mathbf{K}(c_+ t, y)| dy, \\ \int_{\mathbb{R}} |G(t, x, y) - G(s, x, y)| dy &\lesssim C_\phi \int_{|y| \leq 4\varepsilon} |\partial_x^2 \mathbf{K}(c_+ t, y) - \partial_x^2 \mathbf{K}(c_+ s, y)| |y| dy \\ &\quad + \int_{|y| \geq 4\varepsilon} |\partial_x^2 \mathbf{K}(c_+ t, y) - \partial_x^2 \mathbf{K}(c_+ s, y)| dy. \end{aligned}$$

By Lemma 2.1 we obtain that

$$\begin{aligned} \int_{|y| \leq 4\varepsilon} |\partial_x^2 \mathbf{K}(c_+(t, y))| |y| dy &\lesssim t^{-\frac{1}{2}}, \quad \int_{|y| \geq 4\varepsilon} |\partial_x^2 \mathbf{K}(c_+(t, y))| dy \lesssim \int_{|y| \geq 4\varepsilon} \frac{1}{(|y| + t^{\frac{1}{2}})^3} dy \lesssim \varepsilon^{-1} t^{-\frac{1}{2}}, \\ \int_{|y| \leq 4\varepsilon} |\partial_x^2 \mathbf{K}(c_+(t, y)) - \partial_x^2 \mathbf{K}(c_+(s, y))| |y| dy &\lesssim t^{-\frac{1}{2}} \min\{1, \frac{t-s}{s}\}, \\ \int_{|y| \geq 4\varepsilon} |\partial_x^2 \mathbf{K}(c_+(t, y)) - \partial_x^2 \mathbf{K}(c_+(s, y))| dy &\lesssim \varepsilon^{-1} t^{-\frac{1}{2}} \min\{1, \frac{t-s}{s}\}. \end{aligned}$$

Hence

$$\begin{aligned} \sup_{x \in (0, \varepsilon)} \int_{\mathbb{R}} |G(t, x, y)| dy &\lesssim (C_\phi + \varepsilon^{-1}) t^{-\frac{1}{2}}, \\ \sup_{x \in (0, \varepsilon)} \int_{\mathbb{R}} |G(t, x, y) - G(s, x, y)| dy &\lesssim (C_\phi + \varepsilon^{-1}) t^{-\frac{1}{2}} \min\{1, \frac{t-s}{s}\}. \end{aligned}$$

Then Lemma 2.5 implies that

$$\begin{aligned} \|\partial_x f_{2,2}\|_{X_T^{\sigma,p}([- \varepsilon, \varepsilon])} &\lesssim (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{X_T^{\sigma,p}}, \\ \|f_{2,2}\|_{X_T^{\frac{1}{2},2}([- \varepsilon, \varepsilon])} &\lesssim (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{X_T^{\frac{5}{6}, \frac{6}{5}}}. \end{aligned}$$

Combining this with (3.14), (3.15) to yield

$$\begin{aligned} \|\partial_x f_2\|_{X_T^{\sigma,p}([- \varepsilon, \varepsilon])} &\lesssim \|F\|_{X_T^{\sigma,p}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{X_T^{\sigma,p}}, \\ \|f_2\|_{X_T^{\frac{1}{2},2}([- \varepsilon, \varepsilon])} &\lesssim \|F\|_{X_T^{\frac{5}{6}, \frac{6}{5}}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{X_T^{\frac{5}{6}, \frac{6}{5}}}. \end{aligned} \tag{3.17}$$

For boundary terms, we can write

$$\begin{aligned} \partial_x^l f_{2,B}^+ &= C_+(c_- - c_+) \int_0^t \int_0^\infty \partial_x^2 \mathbf{K}(c_+(t-\tau), x+y) (\partial_x^l f_2^-)(\tau, -y) \mathbf{1}_{[0, \varepsilon]} dy d\tau \\ &\quad + C_+(c_- - c_+) \int_0^t \int_0^\infty \partial_x^2 \mathbf{K}(c_+(t-\tau), x+y) (\partial_x^l f_2^-)(\tau, -y) \mathbf{1}_{(\varepsilon, \infty)} dy d\tau \\ &\quad + C_+ \int_0^t \int_0^\infty \partial_x^{1+l} \mathbf{K}(c_+(t-\tau), x+y) (\tilde{F}(\tau, y) - \tilde{F}(\tau, -y)) dy d\tau \\ &:= II_{1,l} + II_{2,l} + II_{3,l}. \end{aligned} \tag{3.18}$$

Applying Lemma 2.3 to get

$$\|II_{1,l}\|_{X_T^{\sigma,p}} \lesssim \|\partial_x^l f_2\|_{X_T^{\sigma,p}([- \varepsilon, \varepsilon])}. \tag{3.19}$$

One can check from Lemma 2.1 that

$$\sup_{x \in \mathbb{R}^+} \int \partial_x^2 \mathbf{K}(c_+ t, x+y) \mathbf{1}_{[\varepsilon, \infty)} dy \lesssim \int_0^\infty \frac{1}{(t^{\frac{1}{2}} + y)^3} dy \lesssim \varepsilon^{-1} t^{-\frac{1}{2}}. \tag{3.20}$$

Then Lemma 2.5 implies that

$$\begin{aligned} \|II_{2,1}\|_{X_T^{\sigma,p}} &\lesssim \varepsilon^{-1} T^{\frac{1}{2}} \|\partial_x f_2\|_{X_T^{\sigma,p}} \stackrel{(3.14)}{\lesssim} \varepsilon^{-1} T^{\frac{1}{2}} (\|F\|_{X_T^{\sigma,p}} + \|\partial_x f\|_{X_T^{\sigma,p}}), \\ \|II_{2,0}\|_{X_T^{\frac{1}{2},2}} &\lesssim \varepsilon^{-1} T^{\frac{1}{2}} \|f_2\|_{X_T^{\frac{1}{2},2}} \stackrel{(3.14)}{\lesssim} \varepsilon^{-1} T^{\frac{1}{2}} (\|F\|_{X_T^{\frac{5}{6}, \frac{6}{5}}} + \|\partial_x f\|_{X_T^{\frac{5}{6}, \frac{6}{5}}}). \end{aligned} \tag{3.21}$$

Finally, following the estimate for f_2 , we can prove that

$$\begin{aligned} \|II_{3,0}\|_{X_T^{\frac{1}{2},2}([0, \varepsilon])} &\lesssim \|F\|_{X_T^{\frac{5}{6}, \frac{6}{5}}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{X_T^{\frac{5}{6}, \frac{6}{5}}}, \\ \|II_{3,1}\|_{X_T^{\sigma,p}([0, \varepsilon])} &\lesssim \|F\|_{X_T^{\sigma,p}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{X_T^{\sigma,p}}. \end{aligned} \tag{3.22}$$

One can estimate $f_{2,B}^-$ similarly. By (3.19), (3.21) and (3.22), we obtain that

$$\begin{aligned}\|\partial_x f_{2,B}\|_{X_T^{\sigma,p}([- \varepsilon, \varepsilon])} &\lesssim \|\partial_x f_2\|_{X_T^{\sigma,p}([- \varepsilon, \varepsilon])} + (1 + \varepsilon^{-1} T^{\frac{1}{2}}) \|F\|_{X_T^{\sigma,p}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|\partial_x f\|_{X_T^{\sigma,p}}, \\ \|f_{2,B}\|_{X_T^{\frac{1}{2},2}([- \varepsilon, \varepsilon])} &\lesssim \|\partial_x f_2\|_{X_T^{\frac{5}{6},\frac{6}{5}}([- \varepsilon, \varepsilon])} + (1 + \varepsilon^{-1} T^{\frac{1}{2}}) \|F\|_{X_T^{\frac{5}{6},\frac{6}{5}}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|\partial_x f\|_{X_T^{\frac{5}{6},\frac{6}{5}}}.\end{aligned}$$

Combining this with (3.17), we obtain that

$$\begin{aligned}\|\partial_x f_N\|_{X_T^{\sigma,p}([- \varepsilon, \varepsilon])} &\lesssim (1 + \varepsilon^{-1} T^{\frac{1}{2}}) \|F\|_{X_T^{\sigma,p}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{X_T^{\sigma,p}}, \\ \|f_N\|_{X_T^{\frac{1}{2},2}([- \varepsilon, \varepsilon])} &\lesssim (1 + \varepsilon^{-1} T^{\frac{1}{2}}) \|F\|_{X_T^{\frac{5}{6},\frac{6}{5}}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{X_T^{\frac{5}{6},\frac{6}{5}}}.\end{aligned}\tag{3.23}$$

Finally, we estimate f_R . For f_3 , we can write

$$f_3^\pm(t, x) = \int_0^t \int_{\mathbb{R}} \mathbf{K}(c_\pm(t - \tau), x - y) R^\pm(\tau, y) dy d\tau,$$

where $R^\pm$ are odd extensions of R :

$$R^+(t, x) = R(t, x) \mathbf{1}_{x \geq 0} - R(-x) \mathbf{1}_{x < 0}, \quad R^-(t, x) = R(t, x) \mathbf{1}_{x < 0} - R(-x) \mathbf{1}_{x \geq 0}.$$

From Lemma 2.6, we obtain

$$\|f_3\|_{X_T^{\frac{1}{2},2}} + \|\partial_x f_3\|_{X_T^{\frac{5}{6},\frac{6}{5}}} \lesssim T^{\frac{1}{4}} (\|R\|_{L_T^1} + \|R\|_{X_T^{1,1}}).$$

The proof for boundary terms $f_{3,B}^\pm$ are similar as previous arguments. We conclude that

$$\|f_R\|_{X_T^{\frac{1}{2},2}} + \|\partial_x f_R\|_{X_T^{\frac{5}{6},\frac{6}{5}}} \lesssim T^{\frac{1}{4}} (\|R\|_{L_T^1} + \|R\|_{X_T^{1,1}}).\tag{3.24}$$

We conclude from (3.12), (3.23) and (3.24) that

$$\begin{aligned}\|f\|_{X_T^{\frac{1}{2},2}([- \varepsilon, \varepsilon])} + \|\partial_x f\|_{X_T^{\frac{5}{6},\frac{6}{5}}([- \varepsilon, \varepsilon])} &\lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}} + (1 + \varepsilon^{-1} T^{\frac{1}{2}}) \|F\|_{X_T^{\frac{5}{6},\frac{6}{5}}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{X_T^{\frac{5}{6},\frac{6}{5}}} \\ &\quad + T^{\frac{1}{4}} (\|R\|_{L_T^1} + \|R\|_{X_T^{1,1}}),\end{aligned}$$

and, when $R = 0$,

$$\begin{aligned}\|\partial_x f\|_{X_T^{1-\gamma,\infty}([- \varepsilon, \varepsilon])} &\lesssim \|\bar{f}_0\|_{\dot{C}^{2\gamma}} + (1 + \varepsilon^{-1} T^{\frac{1}{2}}) \|F\|_{X_T^{1-\gamma,\infty}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|\partial_x f\|_{X_T^{1-\gamma,\infty}}, \\ \|\partial_x f\|_{X_T^{\frac{1}{2},2}([- \varepsilon, \varepsilon])} + \|\partial_x f\|_{X_T^{\frac{3}{4},\infty}([- \varepsilon, \varepsilon])} &\lesssim \|\partial_x \bar{f}_0\|_{L^2} + (1 + \varepsilon^{-1} T^{\frac{1}{2}}) (\|F\|_{X_T^{\frac{1}{2},2}} + \|F\|_{X_T^{\frac{3}{4},\infty}}) \\ &\quad + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} (\|f_x\|_{X_T^{\frac{1}{2},2}} + \|f_x\|_{X_T^{\frac{3}{4},\infty}}).\end{aligned}$$

Fix $T = \frac{1}{(1+C_\phi+\varepsilon^{-1})^{10}}$, we obtain the results. This completes the proof. $\blacksquare$

The following lemma gives estimates of L^p norms near jump points.

Lemma 3.8 *Suppose $f_0 = \partial_x \bar{f}_0$. There exists $T > 0$ such that*

$$\|f\|_{L_T^2(\mathbb{I}_\varepsilon)} + \|\partial_x f\|_{L_T^{\frac{6}{5}}(\mathbb{I}_\varepsilon)} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}} + \|F\|_{L_T^{\frac{6}{5}}} + T^{\frac{1}{3}} \|\partial_x f\|_{L_T^{\frac{6}{5}}} + T^{\frac{1}{4}} \|R\|_{L_T^1}.$$

Moreover, if $R = 0$, there holds

$$\|f\|_{L_T^6(\mathbb{I}_\varepsilon)} + \|\partial_x f\|_{L_T^2(\mathbb{I}_\varepsilon)} \lesssim \|f_0\|_{L^2} + \|F\|_{L_T^2} + T^{\frac{1}{3}} \|\partial_x f\|_{L_T^2}.$$

Proof.

1. *Estimate f_L .*

For simplicity, we only consider $f_1^+ \mathbf{1}_{x \geq 0}$, while $f_1^- \mathbf{1}_{x < 0}$ can be done similarly. Integrate by parts we obtain

$$f_1^+(t, x) = \int_{\mathbb{R}} \partial_x \mathbf{K}(b_+ t, x - y) \mathbf{f}_0(y) dy, \quad \text{where } \mathbf{f}_0(y) = \bar{f}_0(y) \mathbf{1}_{y \geq 0} + \bar{f}_0(-y) \mathbf{1}_{y < 0}.$$

By Parseval's identity (see (6.2)), we have for any $T > 0$,

$$\begin{aligned} \|f_1^+ \mathbf{1}_{x \geq 0}\|_{L_T^2} &\sim \|\mathbf{f}_0\|_{L^2} \lesssim \|\bar{f}_0\|_{L^2} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}}, \\ \|\partial_x f_1^+ \mathbf{1}_{x \geq 0}\|_{L_T^2} &\sim \|\partial_x \mathbf{f}_0\|_{L^2} \lesssim \|f_0\|_{L^2}. \end{aligned} \quad (3.25)$$

Then following (6.3) in Lemma 6.1, we obtain from Lemma 2.2 that

$$\begin{aligned} \|\mathbf{1}_{x \geq 0} \partial_x f_1^+\|_{L_T^{6/5}}^{6/5} &\lesssim \int_{\mathbb{R}} \frac{\|\mathbf{f}_0(\cdot) - \mathbf{f}_0(\cdot - z)\|_{L^{6/5}(\mathbb{R}^+)}^{6/5}}{|z|^{\frac{7}{5}}} dz \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}}^{6/5}, \\ \|\mathbf{1}_{x \geq 0} f_1^+\|_{L_T^6}^6 &\lesssim \int_{\mathbb{R}} \frac{\|\mathbf{f}_0(\cdot) - \mathbf{f}_0(\cdot - z)\|_{L^6(\mathbb{R}^+)}^6}{|z|^5} dz \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{2}{3}, 6}}^6 \lesssim \|\bar{f}_0\|_{\dot{W}^{1, 2}}^6 = \|f_0\|_{L^2}^6. \end{aligned} \quad (3.26)$$

By (3.25) and (3.26) we obtain that

$$\|f_1\|_{L_T^2} + \|\partial_x f_1\|_{L_T^{\frac{6}{5}}} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}}, \quad \|f_1\|_{L_T^6} + \|\partial_x f_1\|_{L_T^2} \lesssim \|f_0\|_{L^2}. \quad (3.27)$$

Then we estimate $f_{1,B}$. Recall the formula

$$\begin{aligned} f_{1,B}^+(t, x) &= C_+(c_+ - c_-) \int_0^t \int_0^\infty \partial_x^2 \mathbf{K}(c_+(t - \tau), x + y) f_1^-(\tau, -y) dy d\tau \\ &\quad - C_+ \int_{\mathbb{R}} \partial_x \mathbf{K}(c_+ t, x - y) \bar{\mathbf{f}}_0(y) dy \\ &:= I_1(t, x) + I_2(t, x). \end{aligned}$$

From Lemma 2.1 one has $(\int_0^\infty |\partial_x \mathbf{K}(c_+ \tau, x + y)|^2 d\tau)^{\frac{1}{2}} \lesssim \frac{1}{x+y}$. Applying Lemma 3.5 yields that

$$\|\partial_x^l I_1\|_{L_T^p} \lesssim \|\partial_x^l f_1\|_{L_T^p}, \quad l = 0, 1, \quad p > 1. \quad (3.28)$$

Moreover, recall that $\bar{\mathbf{f}}_0(y) = (\bar{f}_0(-y) - \bar{f}_0(y)) \mathbf{1}_{y \leq 0}$. We can write

$$\begin{aligned} \|I_2\|_{L_T^2}^2 &\lesssim \int_0^\infty \left(\int_0^\infty \|\partial_x \mathbf{K}(c_+ \tau, x + y)\|_{L^2} (\bar{f}_0(y) - \bar{f}_0(-y)) dy \right)^2 dx \\ &\lesssim \int_0^\infty \left(\int_0^\infty (\bar{f}_0(y) - \bar{f}_0(-y)) \frac{dy}{x + y} \right)^2 dx \lesssim \|\bar{f}_0\|_{L^2}^2, \end{aligned} \quad (3.29)$$

where we apply Lemma 3.4 in the last inequality. Similarly, we obtain

$$\|\partial_x I_2\|_{L_T^2} \lesssim \|f_0\|_{L^2}. \quad (3.30)$$

It remains to estimate $\|\partial_x I_2\|_{L_T^{\frac{6}{5}}}$ and $\|I_2\|_{L_T^6}$. Following (6.3) in Lemma 6.1, one gets

$$\|\partial_x^l I_2\|_{L_T^p(\mathbb{R}^+)} \lesssim \int_{\mathbb{R}} \int_0^\infty \frac{|\bar{\mathbf{f}}_0(x) - \bar{\mathbf{f}}_0(x - z)|^p}{|z|^{(1+l)p-1}} dx dz.$$

Applying Lemma 2.2 to obtain that

$$\|\partial_x I_2\|_{L_T^{\frac{6}{5}}} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}}, \quad \|I_2\|_{L_T^6} \lesssim \|f_0\|_{L^2}.$$

Similar argument holds for $f_{1,B}^-$. Combining this with (3.28), (3.29) and (3.30), we obtain that

$$\|f_{1,B}\|_{L_T^2} + \|\partial_x f_{1,B}\|_{L_T^{\frac{6}{5}}} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}}, \quad \|f_{1,B}\|_{L_T^6} + \|\partial_x f_{1,B}\|_{L_T^2} \lesssim \|f_0\|_{L^2}. \quad (3.31)$$

By (3.27) and (3.31) we obtain that

$$\|f_L\|_{L_T^2} + \|\partial_x f_L\|_{L_T^{\frac{6}{5}}} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}}, \quad \|f_L\|_{L_T^6} + \|\partial_x f_L\|_{L_T^2} \lesssim \|f_0\|_{L^2}. \quad (3.32)$$

2. *Estimate f_N .*

We first estimate $f_2^\pm = f_{2,1}^\pm + f_{2,2}^\pm$ as defined in (3.13). For simplicity, we only consider $\partial_x^l f_2^+$, and the other part $\partial_x^l f_2^-$ can be done parallelly. Note that

$$\begin{aligned} f_{2,1}^+(t, x) &= \int_0^t \int_{\mathbb{R}^+} (\partial_x \mathbf{K}(c_+(t-\tau), x-y) + \partial_x \mathbf{K}(c_+(t-\tau), x+y)) F(\tau, y) dy d\tau \\ &= \int_0^t \int_{\mathbb{R}} \partial_x \mathbf{K}(c_+(t-\tau), x-y) (F(\tau, y) \mathbf{1}_{y \geq 0} + F(\tau, -y) \mathbf{1}_{y < 0}) dy d\tau. \end{aligned}$$

Following the proof of Lemma 6.1, we obtain that

$$\|\partial_x^l f_{2,1}\|_{L_T^p} \lesssim \|F\|_{L_T^{\frac{3p}{3+(1-l)p}}}, \quad \text{for } l = 0, 1, \quad p > 1. \quad (3.33)$$

Then we consider $f_{2,2}$. Recall that

$$\partial_x^l f_{2,2}^+(t, x) = \int_0^t \int_0^\infty \partial_x^l G(t-\tau, x, y) f_x(\tau, y) dy d\tau, \quad l = 0, 1,$$

where

$$G(t, x, y) = (\partial_x \mathbf{K}(c_\pm t, x-y) + \partial_x \mathbf{K}(c_\pm t, x+y))(\phi - \bar{\phi})(y).$$

Then Lemma 6.1 yields that

$$\|\partial_x^l f_{2,1}\|_{L_T^p} \lesssim \|f_x\|_{L_T^{\frac{3p}{3+(1-l)p}}}, \quad \text{for } l = 0, 1, \quad p > 1. \quad (3.34)$$

By (3.16) and Lemma 2.1 we can check that

$$\sup_{x \in [0, \varepsilon]} \left(\int_0^T \int_0^\infty |\partial_x^l G(t, x, y)|^{\frac{3}{2+l}} dy dt \right)^{\frac{2+l}{3}} \lesssim (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}}.$$

Applying Young's inequality we obtain that

$$\begin{aligned} \|\partial_x^l f_{2,2}\|_{L_T^p([0, \varepsilon])} &\lesssim \sup_{x \in [0, \varepsilon]} \left(\int_0^T \int_0^\infty |\partial_x^l G(t, x, y)|^{\frac{3}{2+l}} dy dt \right)^{\frac{2+l}{3}} \|f_x\|_{L_T^{\frac{3p}{3+(1-l)p}}} \\ &\lesssim (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{L_T^{\frac{3p}{3+(1-l)p}}}. \end{aligned}$$

Similarly, we have

$$\|\partial_x^l f_{2,2}\|_{L_T^p(\mathbb{I}_\varepsilon)} \lesssim (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{L_T^{\frac{3p}{3+(1-l)p}}}.$$

Combining this with (3.33), we obtain that

$$\|\partial_x^l f_2\|_{L_T^p(\mathbb{I}_\varepsilon)} \lesssim \|F\|_{L_T^{\frac{3p}{3+(1-l)p}}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{L_T^{\frac{3p}{3+(1-l)p}}}. \quad (3.35)$$

Then we consider the boundary term $f_{2,B}$. Recall the formula $\partial_x^l f_{2,B}^+ = II_{1,l} + II_{2,l} + II_{3,l}$ in 3.18. We apply Lemma 3.5 to obtain that

$$\|II_{1,l}\|_{L_T^p} \lesssim \|\partial_x^l f_2\|_{L_T^p([0,\varepsilon])}.$$

For $II_{2,l}$, the kernel is not singular in view of the fact that $x + y \geq \varepsilon$. Applying Young's inequality to obtain

$$\|II_{2,l}\|_{L_T^p} \lesssim \|\partial_x^l f_2\|_{L_T^p} \int_0^t \int_\varepsilon^\infty |\partial_t \mathbf{K}(c_+ \tau, y)| dy d\tau \stackrel{(3.20)}{\lesssim} \varepsilon^{-1} T^{\frac{1}{2}} \|\partial_x^l f_2\|_{L_T^p}.$$

One can estimate $II_{3,l}$ similarly as $\partial_x^l f_2$. We conclude that

$$\begin{aligned} & \|\partial_x^l f_{2,B}^+\|_{L_T^p([0,\varepsilon])} \\ & \lesssim \|\partial_x^l f_2\|_{L_T^p([0,\varepsilon])} + \varepsilon^{-1} T^{\frac{1}{2}} \|\partial_x^l f_2\|_{L_T^p} + \|F\|_{L_T^{\frac{3p}{3+(1-l)p}}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{L_T^{\frac{3p}{3+(1-l)p}}} \\ & \stackrel{(3.34)}{\lesssim} \|\partial_x^l f_2\|_{L_T^p([0,\varepsilon])} + (1 + \varepsilon^{-1} T^{\frac{1}{2}}) \|F\|_{L_T^{\frac{3p}{3+(1-l)p}}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{L_T^{\frac{3p}{3+(1-l)p}}}. \end{aligned}$$

Combining this with (3.35), we get

$$\|\partial_x^l f_N\|_{L_T^p(\mathbb{I}_\varepsilon)} \lesssim (1 + \varepsilon^{-1} T^{\frac{1}{2}}) \|F\|_{L_T^{\frac{3p}{3+(1-l)p}}} + (C_\phi + \varepsilon^{-1}) T^{\frac{1}{2}} \|f_x\|_{L_T^{\frac{3p}{3+(1-l)p}}}. \quad (3.36)$$

3. Estimate f_R .

Recall the formula

$$\partial_x^l f_3^\pm(t, x) = \int_0^t \int_{\mathbb{R}^\pm} (\partial_x^l \mathbf{K}(c_\pm(t-\tau), x-y) - \partial_x^l \mathbf{K}(c_\pm(t-\tau), x+y)) R(\tau, y) dy d\tau,$$

and

$$\begin{aligned} \partial_x^l f_{3,B}^+(t, x) &= C_+(c_+ - c_-) \int_0^t \int_0^\infty \partial_x^2 \mathbf{K}(c_+(t-\tau), x+y) \partial_x^l f_3^-(\tau, -y) dy d\tau \\ &\quad - C_+ \int_0^t \int_0^\infty \partial_x^l \mathbf{K}(c_+(t-\tau), x+y) (R(\tau, y) + R(\tau, -y)) dy. \end{aligned}$$

It follows from Young's inequality and Lemma 3.5 that

$$\|f_R\|_{L_T^2} + \|\partial_x f_R\|_{L_T^{\frac{6}{5}}} \lesssim \|\mathbf{K}\|_{L_T^2} \|R\|_{L_T^1} + \|\partial_x \mathbf{K}\|_{L_T^{\frac{6}{5}}} \|R\|_{L_T^1} \lesssim T^{\frac{1}{4}} \|R\|_{L_T^1}.$$

Combining this with (3.32) and (3.36), we obtain the result by taking $T = \frac{1}{(10+C_\phi+\varepsilon^{-1})^{10}}$. This completes the proof. $\blacksquare$

3.2 Interior estimates

In this part we estimate the solution f on domain $\mathbb{R} \setminus \mathbb{I}_\varepsilon$. The solution to (3.1) satisfies (see Lemma 6.3)

$$\begin{aligned} f(t, x) &= \mathbf{H}(t, \cdot, x) * f_0(\cdot) + \int_0^t \partial_1 \mathbf{H}(t-\tau, \cdot, x) * F(\tau, \cdot) d\tau + \int_0^t \mathbf{H}(t-\tau, \cdot, x) * R(\tau, \cdot) d\tau \\ &\quad - \int_0^t \int \partial_1 \mathbf{H}(t-\tau, x-y, x) (\phi(x) - \phi(y)) \partial_x f(\tau, y) dy d\tau \\ &:= f_L(t, x) + f_N(t, x) + f_R(t, x) + f_E(t, x), \end{aligned} \quad (3.37)$$

where we denote $\mathbf{H}(t, x, y) = \mathbf{K}(\phi(y)t, x)$, $\partial_1 \mathbf{H}(t, x, y) = \partial_x \mathbf{H}(t, x, y)$ and $\partial_2 \mathbf{H}(t, x, y) = \partial_y \mathbf{H}(t, x, y)$. We approximate the kernel of equation (3.1) by $\mathbf{H}(t, x, y)$ and f_E is the error term.

Lemma 3.9 *Let f satisfy (3.37) with data $f_0 = \partial_x \bar{f}_0$. There exists $T > 0$ such that*

$$\|f\|_{X_T^{\frac{1}{2},2}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} + \|\partial_x f\|_{X_T^{\frac{5}{6},\frac{6}{5}}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}} + \|F\|_{X_T^{\frac{5}{6},\frac{6}{5}}} + T^{\frac{1}{4}}(\|R\|_{X_T^{1,1}} + \|R\|_{L_T^1}) + T^{\frac{1}{3}}\|\partial_x f\|_{X_T^{\frac{5}{6},\frac{6}{5}}}.$$

Moreover, if $R = 0$, then

$$\begin{aligned} \|\partial_x f\|_{X_T^{\frac{1}{2},2}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} + \|\partial_x f\|_{X_T^{\frac{3}{4},\infty}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} &\lesssim \|f_0\|_{L^2} + \|F\|_{X_T^{\frac{1}{2},2}} + \|F\|_{X_T^{\frac{3}{4},\infty}} + T^{\frac{1}{3}}(\|\partial_x f\|_{X_T^{\frac{1}{2},2}} + \|\partial_x f\|_{X_T^{\frac{3}{4},\infty}}), \\ \|\partial_x f\|_{X_T^{1-\gamma,\infty}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} &\lesssim \|\bar{f}_0\|_{\dot{C}^{2\gamma}} + \|F\|_{X_T^{1-\gamma,\infty}} + T^{\frac{1}{3}}\|\partial_x f\|_{X_T^{1-\gamma,\infty}}. \end{aligned}$$

Proof. We first estimate

$$f_L(t, x) = \int_{\mathbb{R}} \mathbf{H}(t, x - y, x) f_0(y) dy = \int_{\mathbb{R}} \partial_1 \mathbf{H}(t, x - y, x) \bar{f}_0(y) dy.$$

One has

$$\partial_x f_L(t, x) = \int_{\mathbb{R}} (\partial_1^2 \mathbf{H}(t, x - y, x) + \partial_1 \partial_2 \mathbf{H}(t, x - y, x)) \bar{f}_0(y) dy.$$

Note that $\mathbf{H}(t, x, y)$ is a time rescaling of heat kernel. We can write

$$\partial_1^2 \mathbf{H}(t, z, x) + \partial_1 \partial_2 \mathbf{H}(t, z, x) = \partial_x^2 \mathbf{K}(\phi(x)t, z) + t\phi'(x)\partial_t \partial_x \mathbf{K}(\phi(x)t, z) := \bar{G}(t, z, x).$$

By Lemma 2.1, it is easy to check that $\bar{G}(t, z, x)$ satisfies the conditions in Remark 3.3. Then Lemma 3.2 and Remark 3.3 imply that

$$\begin{aligned} \|f_L\|_{X_T^{\frac{1}{2},2}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} + \|\partial_x f_L\|_{X_T^{\frac{5}{6},\frac{6}{5}}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} &\lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}}, \quad \|\partial_x f_L\|_{X_T^{1-\gamma,\infty}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} \lesssim \|\bar{f}_0\|_{\dot{C}^{2\gamma}}, \\ \|\partial_x f_L\|_{X_T^{\frac{1}{2},2}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} + \|\partial_x f_L\|_{X_T^{\frac{3}{4},\infty}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} &\lesssim \|\partial_x \bar{f}_0\|_{L^2}. \end{aligned} \tag{3.38}$$

Then we estimate the nonlinear term

$$f_N(t, x) = \int_0^t \int_{\mathbb{R}} \partial_1 \mathbf{H}(t - \tau, x - y, x) F(\tau, y) dy d\tau.$$

We have

$$\partial_x f_N(t, x) = \int_0^t \int_{\mathbb{R}} \bar{G}(t - \tau, x - y, x) F(\tau, y) dy d\tau.$$

By Remark 2.4 we have

$$\|\partial_x f_N\|_{X_T^{\sigma,p}} \lesssim \|F\|_{X_T^{\sigma,p}}, \quad \forall \sigma \in (0, 1 - \alpha), \quad p \in [1, \infty]. \tag{3.39}$$

Since

$$\sup_{z \in \mathbb{R} \setminus \mathbb{I}_\varepsilon} \|\partial_1 \mathbf{H}(t, \cdot, z)\|_{L^{\frac{3}{2}}} \lesssim t^{-\frac{2}{3}}.$$

Then Lemma 2.5 implies that

$$\left\| \int_0^t \int_{\mathbb{R}} \partial_1 \mathbf{H}(t - \tau, x - y, x) F(\tau, y) dy d\tau \right\|_{X_T^{\frac{1}{2},2}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} \lesssim \|F\|_{X_T^{\frac{5}{6},\frac{6}{5}}}.$$

Combining this with (3.39) to obtain that

$$\begin{aligned} \|f_N\|_{X_T^{\frac{1}{2},2}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} + \|\partial_x f_N\|_{X_T^{\frac{5}{6},\frac{6}{5}}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} &\lesssim \|F\|_{X_T^{\frac{5}{6},\frac{6}{5}}}, \\ \|\partial_x f_N\|_{X_T^{\frac{1}{2},2}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} + \|\partial_x f_N\|_{X_T^{\frac{3}{4},\infty}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} &\lesssim \|F\|_{X_T^{\frac{1}{2},2}} + \|F\|_{X_T^{\frac{3}{4},\infty}}, \\ \|\partial_x f_N\|_{X_T^{1-\gamma,\infty}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} &\lesssim \|F\|_{X_T^{1-\gamma,\infty}}. \end{aligned} \tag{3.40}$$

Then we estimate

$$f_R = \int_0^t \int_{\mathbb{R}} \mathbf{H}(t - \tau, x - y, x) R(\tau, y) dy d\tau.$$

By Lemma 2.6, we obtain that

$$\|f_R\|_{X_T^{\frac{1}{2}, 2}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} + \|\partial_x f_R\|_{X_T^{\frac{5}{6}, \frac{6}{5}}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} \lesssim T^{\frac{1}{4}} (\|R\|_{X_T^{1,1}} + \|R\|_{L_T^1}). \quad (3.41)$$

Finally, we estimate f_E . One has

$$f_E(t, x) = - \int_0^t \int_{\mathbb{R}} G_1(t - \tau, x, y) \partial_x f(\tau, y) dy d\tau,$$

where

$$G_1(t - \tau, x, y) = \partial_1 \mathbf{H}(t - \tau, x - y, x) (\phi(x) - \phi(y)).$$

Note that

$$|\phi(x) - \phi(y)| \lesssim C_\phi |x - y|, \quad \forall x \in \mathbb{R} \setminus \mathbb{I}_\varepsilon, \quad |y - x| \leq \varepsilon.$$

Hence for any $0 < s < t < T$,

$$\begin{aligned} \sup_{x \in \mathbb{R} \setminus \mathbb{I}_\varepsilon} \int_{\mathbb{R}} |\partial_x G_1(t, x, y)| dy &\lesssim (\varepsilon^{-1} + C_\phi) t^{-\frac{1}{2}}, \\ \sup_{x \in \mathbb{R} \setminus \mathbb{I}_\varepsilon} \int_{\mathbb{R}} |\partial_x G_1(t, x, y) - \partial_x G_1(s, x, y)| dy &\lesssim (\varepsilon^{-1} + C_\phi) s^{-\frac{1}{2}} \min\{1, \frac{t-s}{s}\}, \\ \sup_{x \in \mathbb{R} \setminus \mathbb{I}_\varepsilon} \|G_1(t, x, \cdot)\|_{L^{\frac{3}{2}}} &\lesssim C_\phi t^{-\frac{1}{6}}, \\ \sup_{x \in \mathbb{R} \setminus \mathbb{I}_\varepsilon} \|G_1(t, x, \cdot) - G_1(s, x, \cdot)\|_{L^{\frac{3}{2}}} &\lesssim C_\phi t^{-\frac{1}{6}} \min\{1, \frac{t-s}{s}\}. \end{aligned}$$

Combining this with Lemma 2.5 to obtain that for $(\sigma, p) \in \{(1 - \gamma, \infty), (\frac{1}{2}, 2), (\frac{3}{4}, \infty), (\frac{5}{6}, \frac{6}{5})\}$,

$$\|\partial_x f_E\|_{X_T^{\sigma, p}} \lesssim T^{\frac{1}{2}} (C_\phi + \varepsilon^{-1}) \|\partial_x f\|_{X_T^{\sigma, p}}, \quad (3.42)$$

and

$$\|f_E\|_{X_T^{\frac{1}{2}, 2}} \lesssim T^{\frac{1}{2}} C_\phi \|\partial_x f\|_{X_T^{\frac{5}{6}, \frac{6}{5}}}. \quad (3.43)$$

Take $T = \frac{1}{(1 + C_\phi + \varepsilon^{-1})^{10}}$. Then the result follows from (3.38), (3.40), (3.41), (3.42) and (3.43). ■

Lemma 3.10 *Suppose $f_0 = \partial_x \bar{f}_0$. There exists $T > 0$ such that*

$$\begin{aligned} \|f\|_{L_T^2(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} + \|\partial_x f\|_{L_T^{\frac{6}{5}}(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} &\leq C_1 (\|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}} + \|F\|_{L_T^{\frac{6}{5}}} + T^{\frac{1}{4}} \|R\|_{L_T^1}) \\ &\quad + \frac{1}{10} (\|f\|_{L_T^2} + \|\partial_x f\|_{L_T^{\frac{6}{5}}}). \end{aligned}$$

Moreover, if $R = 0$, there holds

$$\|f\|_{L_T^6(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} + \|\partial_x f\|_{L_T^2(\mathbb{R} \setminus \mathbb{I}_\varepsilon)} \leq C_1 (\|f_0\|_{L^2} + \|F\|_{L_T^2}) + \frac{1}{10} (\|f\|_{L_T^6} + \|\partial_x f\|_{L_T^2}).$$

Proof. There exists a sequence $\{\chi_n\}_{n \in \mathbb{Z}} \subset \mathbb{R} \setminus \mathbb{I}_\varepsilon$ such that $\sum_n \chi_n^\ell(x) = 1$, $\forall x \in \mathbb{R} \setminus \mathbb{I}_\varepsilon$ and $\text{supp}(\chi_j^\ell) \cap \text{supp}(\chi_k^\ell) = \emptyset$ if $|j - k| \geq 2$, where $\chi_n^\ell : \mathbb{R} \rightarrow [0, 1]$ is a smooth cutoff function satisfying $\mathbf{1}_{[z_n - \ell/2, z_n + \ell/2]} \leq \chi_n^\ell \leq \mathbf{1}_{[z_n - \ell, z_n + \ell]}$ with $0 < \ell < \varepsilon$. We rewrite the equation (3.1) as

$$\begin{aligned} \partial_t(f \chi_n^\ell) - \phi(z_n) \partial_x^2(f \chi_n^\ell) &= \partial_x F_1 + R_1, \\ (f \chi_n^\ell)_{t=0} &= \partial_x(\bar{f}_0 \chi_n^\ell) - \bar{f}_0 \partial_x \chi_n^\ell, \end{aligned}$$

where

$$F_1 = F\chi_n^\ell + (\phi(x) - \phi(z_n))\partial_x f\chi_n^\ell + \phi(z_n)f\partial_x \chi_n^\ell, \quad R_1 = \phi(x)\partial_x f\partial_x \chi_n^\ell + R\chi_n^\ell.$$

Applying Lemma 6.1 to obtain that

$$\|f\chi_n^\ell\|_{L_T^2} + \|\partial_x(f\chi_n^\ell)\|_{L_T^{\frac{6}{5}}} \lesssim \|\bar{f}_0\chi_n^\ell\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}} + T^{\frac{1}{3}}\|\bar{f}_0\partial_x \chi_n^\ell\|_{L_T^{\frac{6}{5}}} + \|F_1\|_{L_T^{\frac{6}{5}}} + T^{\frac{1}{4}}\|R_1\|_{L_T^1},$$

and

$$\|f\chi_n^\ell\|_{L_T^6} + \|\partial_x(f\chi_n^\ell)\|_{L_T^2} \lesssim \|f_0\chi_n^\ell\|_{L^2} + \|F_1\|_{L_T^2} + T^{\frac{1}{4}}\ell^{-1}\|R_1\|_{L_T^2}, \quad \text{if } R = 0.$$

Note that

$$\begin{aligned} \|F_1\|_{L_T^{\frac{6}{5}}} &\lesssim \|F\chi_n^\ell\|_{L_T^{\frac{6}{5}}} + \|\phi(x) - \phi(z_n)\|_{L^\infty(B_\ell)}\|\partial_x f\|_{L_T^{\frac{6}{5}}(B_\ell)} + \|f\|_{L_{t,x,T}^2(B_\ell)}\|\partial_x \chi_n^\ell\|_{L_T^3} \\ &\lesssim \|F\|_{L_T^{\frac{6}{5}}(B_\ell)} + \ell C_\phi\|\partial_x f\|_{L_T^{\frac{6}{5}}(B_\ell)} + \ell^{-\frac{2}{3}}T^{\frac{1}{3}}\|f\|_{L_T^2(B_\ell)}, \\ \|R_1\|_{L_T^1} &\lesssim \|\partial_x f\|_{L_T^{\frac{6}{5}}(B_\ell)}\|\partial_x \chi_n^\ell\|_{L_T^4} + \|R\|_{L_T^1(B_\ell)} \\ &\lesssim T^{\frac{1}{4}}\ell^{-\frac{3}{4}}\|\partial_x f\|_{L_T^{\frac{6}{5}}(B_\ell)} + \|R\|_{L_T^1(B_\ell)}, \end{aligned}$$

where we denote $B_\ell = [z_n - \ell, z_n + \ell]$. Moreover, we can check that

$$\sum_n \|\bar{f}_0\chi_n^\ell\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}} + \ell^{-1}\|\bar{f}_0\|_{L^2} \lesssim (1 + \ell^{-1})\|\bar{f}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}}.$$

Taking sum of n , we obtain that

$$\begin{aligned} \|f\|_{L_T^2(\mathbb{R}\setminus\mathbb{I}_\varepsilon)} + \|\partial_x f\|_{L_T^{\frac{6}{5}}(\mathbb{R}\setminus\mathbb{I}_\varepsilon)} &\leq \tilde{C}_1(1 + \ell^{-1})\|\bar{f}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}} + \tilde{C}_1\|F\|_{L_T^{\frac{6}{5}}} + \tilde{C}_1T^{\frac{1}{4}}\|R\|_{L_T^1} \\ &\quad + \tilde{C}_1(\ell C_\phi + T^{\frac{1}{3}}\ell^{-\frac{3}{4}})(\|\partial_x f\|_{L_T^{\frac{6}{5}}} + \|f\|_{L_T^2}). \end{aligned}$$

Similarly, when $R = 0$, we have

$$\begin{aligned} \|f\|_{L_T^6(\mathbb{R}\setminus\mathbb{I}_\varepsilon)} + \|\partial_x f\|_{L_T^2(\mathbb{R}\setminus\mathbb{I}_\varepsilon)} &\leq \tilde{C}_1(1 + \ell^{-1})\|f_0\|_{L^2} + \tilde{C}_1\|F\|_{L_T^2} \\ &\quad + \tilde{C}_1(\ell C_\phi + \ell^{-1}T^{\frac{1}{4}})(\|\partial_x f\|_{L_T^2} + \|f\|_{L_T^6}). \end{aligned}$$

Then we obtain the result by taking $\ell = (10 + \tilde{C}_1 + C_\phi)^{-10}$ and $T = \ell^{10}$. This completes the proof. $\blacksquare$

Remark 3.11 We have two ways to prove Lemma 3.10. One is to approximate the kernel (see (3.37)), and the other is to cut off in space. We choose the latter one for simplicity. It is equivalent to approximate the kernel piece wisely, namely, approximate the kernel by $\sum_n \mathbf{H}(t, x, z_n)\chi_n^l(x)$.

4 Well-posedness for the isentropic Navier–Stokes equations

In this section, we consider the Cauchy problem of (1.2) with initial data $(v(0, x), u(0, x)) = (v_0, u_0)$ satisfying

$$v_0 \in L^\infty, \quad \inf_x v_0 \geq \lambda_0 > 0, \quad u_0 = \partial_x \bar{u}_0 \text{ with } \bar{u}_0 \in \dot{C}^{2\gamma}. \quad (4.1)$$

The initial data v_0 is allowed to have finite jump. More precisely, we assume there exists a sequence $\{a_n\}_{n=1}^N$ satisfying

$$\min_{j \neq k} |a_j - a_k| \geq \sigma > 0, \quad \|v_0 - b_{\varepsilon, \eta}\|_{L^\infty} \leq \varepsilon_0, \quad (4.2)$$

for some constants $0 < \varepsilon, \eta \ll \sigma$. Here

$$b_{\varepsilon, \eta}(x) = \sum_{n=1}^{N+1} (v_0 * \rho_\eta \mathbf{1}_{(a_{n-1}+\varepsilon, a_n-\varepsilon)} + (v_0 * \rho_\eta)(a_n + \varepsilon) \mathbf{1}_{[a_n, a_n+\varepsilon]} + v_0 * \rho_\eta(a_n - \varepsilon) \mathbf{1}_{[a_n-\varepsilon, a_n]}) , \quad (4.3)$$

where we take $a_0 = -\infty$ and $a_{N+1} = +\infty$.

The main result is the following

Theorem 4.1 *There exists $\varepsilon_0 > 0$ such that if initial data (v_0, u_0) satisfies (4.1) and (4.2), then the system (1.3) admits a unique local solution (v, u) in $[0, T]$ satisfying*

$$\inf_{t \in [0, T]} \inf_x v(t, x) \geq \frac{\lambda_0}{2}, \quad \|v\|_{Y_T} \leq 2\|v_0\|_{L^\infty}, \quad \|\partial_x u\|_{X_T^{1-\gamma, \infty}} \leq M.$$

for some $T, M > 0$.

Clearly, the theorem (1.1) is as a consequence of Theorem 4.1.

Define the space

$$\mathcal{E}_{T, M} := \{w : w(0, x) = u_0(x), \|\partial_x w\|_{X_T^{1-\gamma, \infty}} \leq M\}.$$

Consider $w \in \mathcal{E}_{T, M}$, we define a map $\mathcal{T}w = u$, where u is a solution to the equation

$$\partial_t u - \mu \left(\frac{\partial_x u}{b_{\varepsilon, \eta}} \right)_x = -(p(v))_x + \mu \left(\left(\frac{1}{v} - \frac{1}{b_{\varepsilon, \eta}} \right) \partial_x w \right)_x =: \tilde{F}(v, w)_x, \quad (4.4)$$

where v is defined by

$$v(t, x) = v_0(x) + \int_0^t \partial_x w(s, x) ds. \quad (4.5)$$

The following is a key lemma to prove Theorem 4.1.

Lemma 4.1 *For any initial data*

$$v_0 \in L^\infty, \quad \inf_x v_0 \geq \lambda_0 > 0, \quad u_0 = \partial_x \bar{u}_0 \text{ with } \bar{u}_0 \in \dot{C}^{2\gamma},$$

there exists $\varepsilon_0 > 0$ such that if (4.2) holds, then

$$\begin{aligned} \|\partial_x(\mathcal{T}w)\|_{X_T^{1-\gamma, \infty}} &\leq M, \quad \forall w \in \mathcal{E}_{T, M}, \\ \|\partial_x(\mathcal{T}w_1 - \mathcal{T}w_2)\|_{X_T^{1-\gamma, \infty}} &\leq \frac{1}{2} \|\partial_x(w_1 - w_2)\|_{X_T^{1-\gamma, \infty}}, \quad \forall w_1, w_2 \in \mathcal{E}_{T, M}, \end{aligned}$$

for some $T, M > 0$ depending on $\varepsilon_0, \lambda_0, \|v_0\|_{L^\infty}$ and $\|\bar{u}_0\|_{\dot{C}^{2\gamma}}$.

Proof. We first show the estimates of v . By definition (4.5), it is easy to check that

$$\begin{aligned} \sup_{s \in [0, T]} \|v - b_{\varepsilon, \eta}\|_{L^\infty} &\lesssim \|v_0 - b_{\varepsilon, \eta}\|_{L^\infty} + \int_0^T \|w_x(\tau)\|_{L^\infty} d\tau \lesssim \|v_0 - b_{\varepsilon, \eta}\|_{L^\infty} + T^\gamma \|\partial_x w\|_{X_T^{1-\gamma, \infty}}, \\ \sup_{0 < s < t < T} s^\alpha \frac{\|v(t) - v(s)\|_{L^\infty}}{(t-s)^\alpha} &\lesssim \sup_{0 < s < t < T} s^\alpha \frac{\int_s^t \|\partial_x w\|_{L^\infty} d\tau}{(t-s)^\alpha} \\ &\lesssim \|\partial_x w\|_{X_T^{1-\gamma, \infty}} \sup_{s < t < T} s^\alpha \frac{t^\gamma - s^\gamma}{(t-s)^\alpha} \lesssim T^\gamma \|\partial_x w\|_{X_T^{1-\gamma, \infty}}. \end{aligned}$$

Hence,

$$\|v - b_{\varepsilon, \eta}\|_{Y_T} \leq \|v_0 - b_{\varepsilon, \eta}\|_{L^\infty} + T^\gamma \|\partial_x w\|_{X_T^{1-\gamma, \infty}} \stackrel{(4.2)}{\leq} \varepsilon_0 + T^\gamma M.$$

Moreover, we have

$$\inf_{s \in [0, T]} \inf_x v(s, x) \geq \inf_x v_0(x) - \sup_{s \in [0, T]} \|v(s) - v_0\|_{L^\infty} \geq \lambda_0 - T^\gamma \|\partial_x w\|_{X_T^{1-\gamma, \infty}} \geq \lambda_0 - T^\gamma M.$$

We can take T small enough such that $T^\gamma M \leq \frac{1}{10} \min\{\varepsilon_0, \lambda_0\}$, then

$$\|v - b_{\varepsilon, \eta}\|_{Y_T} \leq 2\varepsilon_0, \quad \inf_{s \in [0, T]} \inf_x v(s, x) \geq \frac{\lambda_0}{2}. \quad (4.6)$$

Applying Lemma 3.1 to (4.4), there exists $T_1 > 0$ such that

$$\|\partial_x(\mathcal{T}w)\|_{X_T^{1-\gamma, \infty}} \lesssim \|\bar{u}_0\|_{\dot{C}^{2\gamma}} + \|\tilde{F}(v, w)\|_{X_T^{1-\gamma, \infty}}, \quad \forall 0 < T < T_1. \quad (4.7)$$

We have

$$\|\tilde{F}(v, w)\|_{X_T^{1-\gamma, \infty}} \lesssim \|p(v)\|_{X_T^{1-\gamma, \infty}} + \left\| \left(\frac{1}{v} - \frac{1}{b_{\varepsilon, \eta}} \right) \partial_x w \right\|_{X_T^{1-\gamma, \infty}}.$$

Note that $p \in W^{2, \infty}$, hence

$$\begin{aligned} \sup_{t \in [0, T]} t^{1-\gamma} \|p(v)(t)\|_{L^\infty} &\lesssim T^{1-\gamma}, \\ \frac{\|p(v)(t) - p(v)(s)\|_{L^\infty}}{(t-s)^\alpha} &\lesssim \frac{\|v(t) - v(s)\|_{L^\infty}}{(t-s)^\alpha} \lesssim s^{-\alpha} \|v - b_\varepsilon\|_{Y_T}, \quad \forall 0 < s < t < T. \end{aligned}$$

Hence,

$$\|p(v)\|_{X_T^{1-\gamma, \infty}} \lesssim T^{1-\gamma} (1 + \|v - b_\varepsilon\|_{Y_T}). \quad (4.8)$$

Denote $q(t, x) = (\frac{1}{v(t, x)} - \frac{1}{b_{\varepsilon, \eta}(x)}) \partial_x w(t, x)$, we have

$$\|q(t)\|_{L^\infty} \lesssim \|v(t) - b_{\varepsilon, \eta}\|_{L^\infty} \|\partial_x w(t)\|_{L^\infty} \lesssim t^{\gamma-1} \|v - b_{\varepsilon, \eta}\|_{Y_T} \|\partial_x w\|_{X_T^{1-\gamma, \infty}}. \quad (4.9)$$

Moreover, for $0 < s < t < T$, we have

$$q(t, x) - q(s, x) = \left(\frac{1}{v(t)} - \frac{1}{v(s)} \right) \partial_x w(t) + \left(\frac{1}{v(s)} - \frac{1}{b_{\varepsilon, \eta}} \right) (\partial_x w(t) - \partial_x w(s)).$$

Note that

$$\begin{aligned} \left\| \left(\frac{1}{v(t)} - \frac{1}{v(s)} \right) \partial_x w(t) \right\|_{L^\infty} &\lesssim \|v(t) - v(s)\|_{L^\infty} \|\partial_x w(t)\|_{L^\infty} \\ &\lesssim (t-s)^\alpha s^{\gamma-1-\alpha} \|v - b_{\varepsilon, \eta}\|_{Y_T} \|\partial_x w\|_{X_T^{1-\gamma, \infty}}, \\ \left\| \left(\frac{1}{v(s)} - \frac{1}{b_{\varepsilon, \eta}} \right) (\partial_x w(t) - \partial_x w(s)) \right\|_{L^\infty} &\lesssim \|v(s) - b_{\varepsilon, \eta}\|_{L^\infty} \|w(t) - w(s)\|_{L^\infty} \\ &\lesssim (t-s)^\alpha s^{\gamma-1-\alpha} \|v - b_{\varepsilon, \eta}\|_{Y_T} \|\partial_x w\|_{X_T^{1-\gamma, \infty}}. \end{aligned}$$

Then one gets

$$\sup_{0 < s < t < T} s^{1-\gamma+\alpha} \frac{\|q(t) - q(s)\|_{L^\infty}}{(t-s)^\alpha} \lesssim \|v - b_{\varepsilon, \eta}\|_{Y_T} \|\partial_x w\|_{X_T^{1-\gamma, \infty}}. \quad (4.10)$$

Hence we get from (4.9) and (4.10) that

$$\|q\|_{X_T^{1-\gamma, \infty}} \lesssim \|v - b_{\varepsilon, \eta}\|_{Y_T} \|\partial_x w\|_{X_T^{1-\gamma, \infty}}.$$

Combining this with (4.8) to obtain that

$$\|\tilde{F}(v, w)\|_{X_T^{1-\gamma, \infty}} \leq C_1 T^{1-\gamma} (1 + \|v - b_\varepsilon\|_{Y_T}) + C_1 \|v - b_{\varepsilon, \eta}\|_{Y_T} \|\partial_x w\|_{X_T^{1-\gamma, \infty}}. \quad (4.11)$$

We conclude from (4.6), (4.7) and (4.11) that

$$\|\partial_x(\mathcal{T}w)\|_{X_T^{1-\gamma, \infty}} \leq C_1 \|\bar{u}_0\|_{\dot{C}^{2\gamma}} + C_1 T^{1-\gamma} (1 + 2\varepsilon_0) + 2C_1 \varepsilon_0 M, \quad \forall w \in \mathcal{E}_{T, M}.$$

Then we obtain $\|\partial_x(\mathcal{T}w)\|_{X_T^{1-\gamma, \infty}} \leq M$ by taking $M = 2C_1 \|\bar{u}_0\|_{\dot{C}^{2\gamma}} + 1$, $\varepsilon_0 < \frac{1}{(1+C_1)^{10}}$ and $T < \min\{\frac{1}{(1+C_1)^{10}}, T_1\}$. This implies that $\mathcal{T}$ maps $\mathcal{E}_{T, M}$ to itself.

It remains to prove that $\mathcal{T}$ is contraction. Consider $w_1, w_2 \in \mathcal{E}_{T, M}$, let $v_m(t, x) = v_0(x) + \int_0^t \partial_x w_m(s, x) ds$, $m = 1, 2$. Then it is easy to check that both v_1 and v_2 satisfy (4.6), and

$$\|v_1 - v_2\|_{Y_T} \lesssim T^\gamma \|\partial_x(w_1 - w_2)\|_{X_T^{1-\gamma, \infty}}. \quad (4.12)$$

Thanks to Lemma 3.1, there exists $T_2 > 0$ such that

$$\|\partial_x(\mathcal{T}w_1 - \mathcal{T}w_2)\|_{X_T^{1-\gamma, \infty}} \lesssim \|\tilde{F}(v_1, w_1) - \tilde{F}(v_2, w_2)\|_{X_T^{1-\gamma, \infty}}, \quad \forall 0 < T < T_2. \quad (4.13)$$

Note that

$$\tilde{F}(v_1, w_1) - \tilde{F}(v_2, w_2) = p(v_2) - p(v_1) + \mu \left(\left(\frac{1}{v_1} - \frac{1}{b_{\varepsilon, \eta}} \right) \partial_x w_1 - \left(\frac{1}{v_2} - \frac{1}{b_{\varepsilon, \eta}} \right) \partial_x w_2 \right).$$

We first estimate $p(v_2) - p(v_1)$. Note that $p \in W^{2, \infty}$, hence

$$\|p(v_2(t)) - p(v_1(t))\|_{L^\infty} \lesssim \|v_1 - v_2\|_{L^\infty} \lesssim \|v_1 - v_2\|_{Y_T},$$

and

$$\begin{aligned} & \|p(v_2(t)) - p(v_1(t)) - (p(v_2(s)) - p(v_1(s)))\|_{L^\infty} \\ & \lesssim \|(v_2 - v_1)(t) - (v_2 - v_1)(s)\|_{L^\infty} + \|v_2(t) - v_1(t)\|_{L^\infty} (\|v_1(t) - v_1(s)\|_{L^\infty} + \|v_2(t) - v_2(s)\|_{L^\infty}) \\ & \lesssim (t - s)^\alpha s^{-\alpha} \|v_1 - v_2\|_{Y_T} (1 + \|v_1 - b_{\varepsilon, \eta}\|_{Y_T} + \|v_2 - b_{\varepsilon, \eta}\|_{Y_T}). \end{aligned}$$

Hence

$$\begin{aligned} \|p(v_2) - p(v_1)\|_{X_T^{1-\gamma, \infty}} & \lesssim T^{1-\gamma} \|v_1 - v_2\|_{Y_T} (1 + \|v_1 - b_{\varepsilon, \eta}\|_{Y_T} + \|v_2 - b_{\varepsilon, \eta}\|_{Y_T}) \\ & \stackrel{(4.12)}{\lesssim} T \|\partial_x(w_1 - w_2)\|_{X_T^{1-\gamma, \infty}}. \end{aligned} \quad (4.14)$$

Then we estimate $\mathcal{Q} = \left(\frac{1}{v_1} - \frac{1}{b_{\varepsilon, \eta}} \right) \partial_x w_1 - \left(\frac{1}{v_2} - \frac{1}{b_{\varepsilon, \eta}} \right) \partial_x w_2$. We have

$$\mathcal{Q} = \left(\frac{1}{v_1} - \frac{1}{v_2} \right) \partial_x w_1 + \left(\frac{1}{v_2} - \frac{1}{b_{\varepsilon, \eta}} \right) \partial_x(w_1 - w_2).$$

For any $0 < t < T$, one has

$$\|\mathcal{Q}(t)\|_{L^\infty} \lesssim \|(v_1 - v_2)(t)\|_{L^\infty} \|\partial_x w_1(t)\|_{L^\infty} \lesssim t^{\gamma-1} \|v_1 - v_2\|_{Y_T} \|\partial_x w_1\|_{X_T^{1-\gamma, \infty}}. \quad (4.15)$$

For any $0 < s < t < T$,

$$\begin{aligned} & \|\mathcal{Q}(t) - \mathcal{Q}(s)\|_{L^\infty} \\ & \lesssim \|(v_1 - v_2)(t) - (v_1 - v_2)(s)\|_{L^\infty} \|\partial_x w_1(t)\|_{L^\infty} \\ & \quad + \|(v_1 - v_2)(t)\|_{L^\infty} (\|v_1(t) - v_1(s)\|_{L^\infty} + \|v_2(t) - v_2(s)\|_{L^\infty}) \|\partial_x w_1(t)\|_{L^\infty} \\ & \quad + \|(v_1 - v_2)(s)\|_{L^\infty} \|\partial_x(w_1(t) - w_1(s))\|_{L^\infty} + \|v_2(t) - v_2(s)\|_{L^\infty} \|\partial_x(w_1 - w_2)(t)\|_{L^\infty} \\ & \quad + \|v_2(s) - b_{\varepsilon, \eta}\|_{L^\infty} \|\partial_x(w_1 - w_2)(t) - \partial_x(w_1 - w_2)(s)\|_{L^\infty} \\ & \lesssim (t - s)^\alpha s^{\gamma-1-\alpha} \|v_1 - v_2\|_{Y_T} \|\partial_x w_1\|_{X_T^{1-\gamma, \infty}} (1 + \|v_1 - b_{\varepsilon, \eta}\|_{Y_T} + \|v_2 - b_{\varepsilon, \eta}\|_{Y_T}) \\ & \quad + (t - s)^\alpha s^{\gamma-1-\alpha} \|v_2 - b_{\varepsilon, \eta}\|_{Y_T} \|\partial_x(w_1 - w_2)\|_{X_T^{1-\gamma, \infty}}. \end{aligned} \quad (4.16)$$

Combining (4.12), (4.15), (4.16) with the fact that $\|v_1 - b_{\varepsilon,\eta}\|_{Y_T} + \|v_2 - b_{\varepsilon,\eta}\|_{Y_T} \lesssim \varepsilon_0$ and $\|\partial_x w_1\|_{X_T^{1-\gamma,\infty}} \leq M$, we obtain

$$\|\mathcal{Q}\|_{X_T^{1-\gamma,\infty}} \lesssim MT^\gamma \|\partial_x(w_1 - w_2)\|_{X_T^{1-\gamma,\infty}}. \quad (4.17)$$

We conclude from (4.13), (4.14) and (4.17) that

$$\|\partial_x(\mathcal{T}w_1 - \mathcal{T}w_2)\|_{X_T^{1-\gamma,\infty}} \leq C_2(MT^\gamma + T) \|\partial_x(w_1 - w_2)\|_{X_T^{1-\gamma,\infty}} \leq \frac{1}{2} \|\partial_x(w_1 - w_2)\|_{X_T^{1-\gamma,\infty}}.$$

Here we fix $T = \min \left\{ \frac{1}{(1+C_1+C_2)10}, T_1, T_2 \right\}$. This completes the proof. $\blacksquare$

5 Well-posedness for the full Navier–Stokes system

In this section, we consider the Cauchy problem of (1.3) with initial data $(v, u, \theta)(0, x) = (v_0, u_0, \theta_0)(x)$ satisfying

$$\inf_x v_0(x) \geq \lambda_0 > 0, \quad \|v_0\|_{L^\infty} < \infty, \quad \|u_0\|_{L^2} < \infty, \quad \|\theta_0\|_{\dot{W}^{-\frac{2}{3}, \frac{6}{5}}} < \infty. \quad (5.1)$$

Moreover, we suppose that there exist $0 < \varepsilon, \eta \ll 1$ such that v_0 satisfies the condition (4.2) for some $\varepsilon_0 > 0$ that will be fixed later in our proof.

Theorem 5.1 *There exists $\varepsilon_0 > 0$ such that if initial data (v_0, u_0, θ_0) satisfies (5.1) and (4.2), then the system (1.3) admits a unique local solution in $[0, T]$ satisfying*

$$\inf_{t \in [0, T]} \inf_x v(t, x) \geq \frac{\lambda_0}{2}, \quad \|v\|_{Y_T} \leq 2\|v_0\|_{L^\infty}, \quad \|(u, \theta)\|_{Z_T} \leq B.$$

for some $T, B > 0$.

Clearly, the theorem (1.2) is as a consequence of Theorem 5.1.

Define the space

$$\mathcal{X}_{T,B} = \{(w, \vartheta) : (w, \vartheta)|_{t=0} = (u_0, \theta_0), \|(w, \vartheta)\|_{Z_T} \leq B\}.$$

Let $(w, \vartheta) \in \mathcal{X}_{T,B}$, we define a map $\mathcal{M}(w, \vartheta) = (u, \theta)$, where u, θ solve the equations

$$\begin{aligned} u_t - \mu \left(\frac{u_x}{b_{\varepsilon,\eta}} \right)_x + (p(v, \theta))_x &= \mu \left(\left(\frac{1}{v} - \frac{1}{b_{\varepsilon,\eta}} \right) \partial_x w \right)_x, \quad u(0, x) = u_0(x), \\ \theta_t - \frac{\kappa}{\mathbf{c}} \left(\frac{\theta_x}{b_{\varepsilon,\eta}} \right)_x &= -\frac{p(v, \vartheta)}{\mathbf{c}} w_x + \frac{\mu}{\mathbf{c}v} (w_x)^2 + \frac{\kappa}{\mathbf{c}} \left(\left(\frac{1}{v} - \frac{1}{b_{\varepsilon,\eta}} \right) \partial_x \vartheta \right)_x, \quad \theta(0, x) = \theta_0(x), \end{aligned}$$

where v is fixed by

$$v(t, x) = v_0(x) + \int_0^t \partial_x w(s, x) ds,$$

and the function $b_{\varepsilon,\eta}$ is defined in (4.3).

The following is the key lemma to prove Theorem 5.1.

Lemma 5.1 *There exists $\varepsilon_0 > 0$ such that if initial data (v_0, u_0, θ_0) satisfies (5.1) and (4.2), then we have*

$$\begin{aligned} \|\mathcal{M}(w, \vartheta)\|_{Z_T} &\leq B, \quad \forall (w, \vartheta) \in \mathcal{X}_{T,B}, \\ \|\mathcal{M}(w_1, \vartheta_1) - \mathcal{M}(w_2, \vartheta_2)\|_{Z_T} &\leq \frac{1}{2} \|(w_1 - w_2, \vartheta_1 - \vartheta_2)\|_{Z_T}, \quad \forall (w_k, \vartheta_k) \in \mathcal{X}_{T,B}, \quad k = 1, 2. \end{aligned}$$

for some $T, B > 0$.

Proof.

1. *Estimate v*

We have

$$\begin{aligned} \|v(t) - b_{\varepsilon, \eta}\|_{L^\infty} &\lesssim \|v_0 - b_{\varepsilon, \eta}\|_{L^\infty} + \int_0^t \|w_x(\tau)\|_{L^\infty} d\tau, \\ \inf_{t \in [0, T]} \inf_x v(t, x) &\geq \inf_x v_0(x) - \int_0^t \|w_x(\tau)\|_{L^\infty} d\tau. \end{aligned}$$

By Hölder's inequality,

$$\int_0^T \|w_x(\tau)\|_{L^\infty} d\tau \leq CT^{\frac{1}{4}} \|w_x\|_{X_T^{\frac{3}{4}, \infty}} \leq CT^{\frac{1}{4}} B.$$

We take T small enough such that $CT^{\frac{1}{4}} B \leq \frac{1}{100} \min\{\lambda_0, \varepsilon_0\}$, and

$$\|v - b_{\varepsilon, \eta}\|_{Y_T} \leq 2\varepsilon_0, \quad \inf_{s \in [0, T]} \inf_x v(s, x) \geq \frac{\lambda_0}{2}. \quad (5.2)$$

2. *Estimate θ*

$$\theta_t - \frac{\kappa}{\mathbf{c}} \left(\frac{\theta_x}{b_{\varepsilon, \eta}} \right)_x = -\frac{p(v, \vartheta)}{\mathbf{c}} w_x + \frac{\mu}{\mathbf{c}v} (w_x)^2 + \frac{\kappa}{\mathbf{c}} \left(\left(\frac{1}{v} - \frac{1}{b_{\varepsilon, \eta}} \right) \partial_x \vartheta \right)_x := R(v, w, \vartheta) + \partial_x \tilde{F}(v, \vartheta).$$

Applying Lemma 3.1 with the equation above for θ to obtain

$$\sum_{\star \in \{L_T^2, X_T^{\frac{1}{2}, 2}\}} \|\theta\|_{\star} + \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6}, \frac{6}{5}}\}} \|\theta_x\|_{\star} \lesssim \|\bar{\theta}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}} + \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6}, \frac{6}{5}}\}} \|\tilde{F}\|_{\star} + T^{\frac{1}{4}} \sum_{\star \in \{L_T^1, X_T^{1,1}\}} \|R\|_{\star}.$$

It is easy to check that

$$\begin{aligned} \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6}, \frac{6}{5}}\}} \|\tilde{F}\|_{\star} &\lesssim \|v - b_{\varepsilon, \eta}\|_{L_T^\infty} (\|\partial_x \vartheta\|_{L_T^{\frac{6}{5}}} + \|\partial_x \vartheta\|_{X_T^{\frac{5}{6}, \frac{6}{5}}}) \lesssim \varepsilon_0 B, \\ \sum_{\star \in \{L_T^1, X_T^{1,1}\}} \|R\|_{\star} &\lesssim \|w_x\|_{L_T^2}^2 + \|\vartheta\|_{L_T^2}^2 + \|w_x\|_{X_T^{\frac{1}{2}, 2}}^2 + \|\vartheta\|_{X_T^{\frac{1}{2}, 2}}^2 \lesssim B^2. \end{aligned}$$

Hence we obtain

$$\begin{aligned} \|\theta\|_{L_T^2} + \|\theta_x\|_{L_T^{\frac{6}{5}}} + \|\theta\|_{X_T^{\frac{1}{2}, 2}} + \|\theta_x\|_{X_T^{\frac{5}{6}, \frac{6}{5}}} \\ \leq \tilde{C}_0 \|\bar{\theta}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}} + \tilde{C}_0 \varepsilon_0 B + \tilde{C}_0 T^{\frac{1}{4}} B^2 \leq \frac{B}{3}, \end{aligned} \quad (5.3)$$

provided $\|\bar{\theta}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}} \leq \frac{B}{\tilde{C}_0}$, $\varepsilon_0 \leq \frac{1}{10(1+\tilde{C}_0)^{10}}$ and $T \leq \frac{1}{(1+\tilde{C}_0+B)^{10}}$.

3. *Estimate u*

Recall the equation

$$u_t - \mu \left(\frac{u_x}{b_\varepsilon} \right)_x = -(p(v, \theta))_x + \mu \left(\left(\frac{1}{v} - \frac{1}{b_\varepsilon} \right) w_x \right)_x =: \partial_x F(v, w, \theta).$$

Applying Lemma 3.1 with the equation above for u , we obtain that

$$\|u_x\|_{L_T^2} + \|u_x\|_{X_T^{\frac{1}{2}, 2}} + \|u_x\|_{X_T^{\frac{3}{4}, \infty}} \lesssim \|u_0\|_{L^2} + \|F\|_{L_T^2} + \|F\|_{X_T^{\frac{1}{2}, 2}} + \|F\|_{X_T^{\frac{3}{4}, \infty}}.$$

It is easy to check that

$$\begin{aligned} \|F\|_{L_T^2} &\lesssim \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2},2}, X_T^{\frac{1}{2},2}\}} \|\theta\|_{\star} + \|v - b_{\varepsilon}\|_{L^\infty} \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2},2}\}} \|w_x\|_{\star}, \\ \|F\|_{X_T^{\frac{3}{4},\infty}} &\lesssim \|\theta\|_{X_T^{\frac{3}{4},\infty}} + \|v - b_{\varepsilon}\|_{L^\infty} \|w_x\|_{X_T^{\frac{3}{4},\infty}}. \end{aligned}$$

By the interpolation inequality, we have $\|\theta\|_{X_T^{\frac{3}{4},\infty}} \lesssim \|\theta\|_{X_T^{\frac{1}{2},2}} + \|\theta_x\|_{X_T^{\frac{5}{6},\frac{6}{5}}}$. Hence

$$\begin{aligned} &\sum_{\star \in \{L_T^2, X_T^{\frac{1}{2},2}, X_T^{\frac{3}{4},\infty}\}} \|u_x\|_{\star} \\ &\leq \tilde{C}_1(\|u_0\|_{L^2} + \|\theta\|_{L_T^2} + \|\theta\|_{X_T^{\frac{1}{2},2}} + \|\theta_x\|_{X_T^{\frac{5}{6},\frac{6}{5}}} + \|v - b_{\varepsilon,\eta}\|_{L^\infty}(\|w_x\|_{L_T^2} + \|w_x\|_{X_T^{\frac{1}{2},2}})) \\ &\stackrel{(5.2),(5.3)}{\leq} \tilde{C}_1(\|u_0\|_{L^2} + \tilde{C}_0\|\bar{\theta}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}} + \tilde{C}_0\varepsilon_0 B + \tilde{C}_0 T^{\frac{1}{4}} B^2 + \varepsilon_0 B) \\ &\leq \frac{B}{3}, \end{aligned}$$

provided $B \geq (10 + \tilde{C}_0 + \tilde{C}_1 + \tilde{C}_2)(\|u_0\|_{L^2} + \|\bar{\theta}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}})$, $\varepsilon_0 \leq \frac{1}{100(1 + \tilde{C}_0 + \tilde{C}_1 + \tilde{C}_2)^{10}}$ and $T \leq \varepsilon_0^{10} B^{-10}$.

Combining this with (5.3), we obtain

$$\|(u, \theta)\|_{Z_T} \leq B.$$

This implies that $\mathcal{M}$ maps $\mathcal{X}_{T,B}$ to itself. It remains to prove that $\mathcal{M}$ is contraction. Consider $(w_1, \vartheta_1), (w_2, \vartheta_2) \in \mathcal{X}_{T,B}$, let $v_m(t, x) = v_0(x) + \int_0^t \partial_x w_m(s, x) ds$, $m = 1, 2$. Then we have

$$\begin{aligned} \|v_m - b_{\varepsilon,\eta}\|_{L_T^\infty} &\leq 2\varepsilon_0, \quad \inf_{s \in [0, T]} \inf_x v_m(s, x) \geq \frac{\lambda_0}{2}, \\ \|v_1 - v_2\|_{L_T^\infty} &\leq \int_0^T \|\partial_x(w_1 - w_2)(s)\|_{L^\infty} ds \lesssim T^{\frac{1}{4}} \|\partial_x(w_1 - w_2)\|_{X_T^{\frac{3}{4},\infty}}. \end{aligned} \tag{5.4}$$

Denote $(u_m, \theta_m) = \mathcal{M}(w_m, \vartheta_m)$, $m = 1, 2$. We have

$$\begin{aligned} \partial_t(u_1 - u_2) - \kappa \left(\frac{\partial_x(u_1 - u_2)}{b_{\varepsilon,\eta}} \right)_x &= \partial_x(F(v_1, w_1, \theta_1) - F(v_2, w_2, \theta_2)), \\ \partial_t(\theta_1 - \theta_2) - \frac{\kappa}{c} \left(\frac{\partial_x(\theta_1 - \theta_2)}{b_{\varepsilon,\eta}} \right)_x &= R(v_1, w_1, \vartheta_1) - R(v_2, w_2, \vartheta_2) + \partial_x(\tilde{F}(v_1, \vartheta_1) - \tilde{F}(v_2, \vartheta_2)). \end{aligned}$$

Applying Lemma 3.1 with $(f, \phi, C_\phi) = (\theta_1 - \theta_2, b_{\varepsilon,\eta}^{-1}, \eta^{-1}\|v_0\|_{L^\infty})$ to get

$$\begin{aligned} &\sum_{\star \in \{L_T^2, X_T^{\frac{1}{2},2}\}} \|\theta_1 - \theta_2\|_{\star} + \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6},\frac{6}{5}}\}} \|\partial_x(\theta_1 - \theta_2)\|_{\star} \\ &\lesssim \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6},\frac{6}{5}}\}} \|\tilde{F}(v_1, \vartheta_1) - \tilde{F}(v_2, \vartheta_2)\|_{\star} + T^{\frac{1}{4}} \sum_{\star \in \{L_T^1, X_T^{1,1}\}} \|R(v_1, w_1, \vartheta_1) - R(v_2, w_2, \vartheta_2)\|_{\star}. \end{aligned} \tag{5.5}$$

Then (5.4) yields

$$\begin{aligned} &\sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6},\frac{6}{5}}\}} \|(\tilde{F}(v_1, \vartheta_1) - \tilde{F}(v_2, \vartheta_2))\|_{\star} \\ &\lesssim \|v_1 - v_2\|_{L^\infty} \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6},\frac{6}{5}}\}} \|\partial_x \vartheta_1\|_{\star} + \|v_1 - b_{\varepsilon,\eta}\|_{L^\infty} \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6},\frac{6}{5}}\}} \|\partial_x(\vartheta_1 - \vartheta_2)\|_{\star} \\ &\lesssim (T^{\frac{1}{5}} B + \varepsilon_0) \|(w_1, \vartheta_1) - (w_2, \vartheta_2)\|_{Z_T}. \end{aligned} \tag{5.6}$$

Moreover, we have

$$|R(v_1, w_1, \vartheta_1) - R(v_2, w_2, \vartheta_2)| \lesssim |\vartheta_1 - \vartheta_2| |\partial_x w_1| + |\vartheta_2| |\partial_x(w_1 - w_2)| + |\vartheta_2| |\partial_x w_2| |v_1 - v_2| \\ + |\partial_x(w_1 - w_2)| (|\partial_x w_1| + |\partial_x w_2|) + |\partial_x w_2|^2 |v_1 - v_2|.$$

By Hölder's inequality, it is easy to check that

$$\begin{aligned} & \|R(v_1, w_1, \vartheta_1) - R(v_2, w_2, \vartheta_2)\|_{L_T^1} \\ & \lesssim \|\vartheta_1 - \vartheta_2\|_{L_T^2} \|\partial_x w_1\|_{L_T^2} + (\|\vartheta_2\|_{L_T^2} + \|\partial_x w_1\|_{L_T^2} + \|\partial_x w_2\|_{L_T^2}) \|\partial_x(w_1 - w_2)\|_{L_T^2} \\ & \quad + (\|\vartheta_2\|_{L_T^2}^2 + \|\partial_x w_2\|_{L_T^2}^2) \|v_1 - v_2\|_{L_T^\infty} \\ & \lesssim (B + B^2 T^{\frac{1}{5}}) \|(w_1, \vartheta_1) - (w_2, \vartheta_2)\|_{Z_T}. \end{aligned} \quad (5.7)$$

Similarly, one gets

$$\sum_{\star \in \{L_T^1, X_T^{1,1}\}} \|R(v_1, w_1, \vartheta_1) - R(v_2, w_2, \vartheta_2)\|_{\star} \lesssim (B + B^2 T^{\frac{1}{5}}) \|(w_1, \vartheta_1) - (w_2, \vartheta_2)\|_{Z_T}.$$

Combining this with (5.5), (5.6) and (5.7) to obtain

$$\begin{aligned} & \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2},2}\}} \|\theta_1 - \theta_2\|_{\star} + \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{6},\frac{6}{5}}\}} \|\partial_x(\theta_1 - \theta_2)\|_{\star} \\ & \leq \tilde{C}_3 (T^{\frac{1}{5}} B + \varepsilon_0 + T^{\frac{1}{4}} (B + B^2 T^{\frac{1}{5}})) \|(w_1, \vartheta_1) - (w_2, \vartheta_2)\|_{Z_T} \\ & \leq 2\tilde{C}_3 \varepsilon_0 \|(w_1, \vartheta_1) - (w_2, \vartheta_2)\|_{Z_T}. \end{aligned} \quad (5.8)$$

Then we estimate $u_1 - u_2$. By Lemma 3.1, we get

$$\sum_{\star \in \{L_T^2, X_T^{\frac{1}{2},2}, X_T^{\frac{3}{4},\infty}\}} \|\partial_x(u_1 - u_2)\|_{\star} \lesssim \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2},2}, X_T^{\frac{3}{4},\infty}\}} \|F(v_1, w_1, \theta_1) - F(v_2, w_2, \theta_2)\|_{\star}.$$

Note that

$$|F(v_1, w_1, \theta_1) - F(v_2, w_2, \theta_2)| \lesssim |\theta_1 - \theta_2| + |\theta_2| |v_1 - v_2| + |v_1 - b_{\varepsilon,\eta}| |\partial_x(w_1 - w_2)| + |v_1 - v_2| |\partial_x w_2|.$$

Using the interpolation $\|f\|_{X_T^{\frac{3}{4},\infty}} \lesssim \|f\|_{X_T^{\frac{1}{2},2}} \|\partial_x f\|_{X_T^{\frac{5}{6},\frac{6}{5}}}$, we get

$$\begin{aligned} & \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2},2}, X_T^{\frac{3}{4},\infty}\}} \|F(v_1, w_1, \theta_1) - F(v_2, w_2, \theta_2)\|_{\star} \\ & \lesssim \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2},2}\}} \|\theta_1 - \theta_2\|_{\star} + \|\partial_x(\theta_1 - \theta_2)\|_{X_T^{\frac{5}{6},\frac{6}{5}}} + \|v_1 - b_{\varepsilon,\eta}\|_{L^\infty} \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2},2}, X_T^{\frac{3}{4},\infty}\}} \|\partial_x(w_1 - w_2)\|_{\star} \\ & \quad + \left(\sum_{\star \in \{L_T^2, X_T^{\frac{1}{2},2}\}} \|\theta_2\|_{\star} + \|\partial_x \theta_2\|_{X_T^{\frac{5}{6},\frac{6}{5}}} + \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2},2}, X_T^{\frac{3}{4},\infty}\}} \|\partial_x w_2\|_{\star} \right) \|v_1 - v_2\|_{L^\infty} \\ & \lesssim (\tilde{C}_3 \varepsilon_0 + B T^{\frac{1}{5}}) \|(w_1, \vartheta_1) - (w_2, \vartheta_2)\|_{Z_T}. \end{aligned}$$

Then we get

$$\begin{aligned} & \sum_{\star \in \{L_T^2, X_T^{\frac{1}{2},2}, X_T^{\frac{3}{4},\infty}\}} \|\partial_x(u_1 - u_2)\|_{\star} \\ & \leq \tilde{C}_4 (\tilde{C}_3 \varepsilon_0 + B T^{\frac{1}{5}}) \|(w_1, \vartheta_1) - (w_2, \vartheta_2)\|_{Z_T} \leq \frac{1}{5} \|(w_1, \vartheta_1) - (w_2, \vartheta_2)\|_{Z_T}, \end{aligned}$$

by taking $B = (10 + \sum_{m=1}^4 \tilde{C}_m)(\|u_0\|_{L^2} + \|\bar{\theta}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}})$, $\varepsilon_0 = 100(1 + \sum_{m=1}^4 \tilde{C}_m)^{-10}$ and $T = \varepsilon_0^{10} B^{-10}$. Combining this with (5.8) yields

$$\|\mathcal{M}(w_1, \vartheta_1) - \mathcal{M}(w_1, \vartheta_2)\|_{Z_T} \leq \frac{1}{2} \|(w_1 - w_2, \vartheta_1 - \vartheta_2)\|_{Z_T}.$$

This completes the proof. $\blacksquare$

In the rest part of this section, we give a local version of Theorem 5.1. Define the local norms

$$\|h\|_{\tilde{L}^p} := \sup_z \|h\|_{L^p([z, z+1])}, \quad \|h\|_{\dot{W}^{s,p}} := \sup_z \|h\chi_z\|_{\dot{W}^{s,p}}, \quad \|f\|_{\tilde{X}_T^{\sigma,p}} = \sup_z \|f\chi_z\|_{\tilde{X}_T^{\sigma,p}},$$

for $h : \mathbb{R} \rightarrow \mathbb{R}$, $f : \mathbb{R}^+ \times \mathbb{R} \rightarrow \mathbb{R}$. Here χ_z is a smooth cutoff function satisfying $\mathbf{1}_{[z-1, z+1]} \leq \chi_z \leq \mathbf{1}_{[z-2, z+2]}$. Moreover, we denote $\|f\|_{\tilde{L}_T^p} = \|f\|_{L_t^p \tilde{L}_x^p}$. We introduce the following lemma.

Lemma 5.2 *Let*

$$g(t, x) = \int_0^t \int_{|y| \geq \rho} \partial_t K(t - \tau, y) F(\tau, x - y) dy d\tau,$$

for some $\rho \in (0, 1)$. Then for any $0 < T < 1$, $p \geq 1$, $\sigma \in (0, 1 - \alpha)$,

$$\|g\|_{\tilde{L}_T^p} \lesssim_\rho \|F\|_{\tilde{L}_T^p}, \quad \|g\|_{\tilde{X}_T^{\sigma,p}} \lesssim_\rho \|F\|_{\tilde{X}_T^{\sigma,p}}.$$

Proof. By Lemma 2.1 and Hölder's inequality,

$$\begin{aligned} |g(t, x)| &\lesssim \int_0^t \int_{|y| \geq \rho} \frac{1}{((t - \tau)^{\frac{1}{2}} + |y|)^3} |F(\tau, x - y)| dy d\tau \\ &\lesssim \sum_{n=0}^{\infty} \int_0^t \int_{|y| \in [\rho+n, \rho+n+1]} \frac{1}{((t - \tau)^{\frac{1}{2}} + |y|)^3} |F(\tau, x - y)| dy d\tau \\ &\lesssim \sum_{n=0}^{\infty} \frac{1}{(\rho + n)^3} \min\{\|F\|_{\tilde{L}_T^p}, \|F\|_{\tilde{X}_T^{\sigma,p}}\} \lesssim \rho^{-3} \min\{\|F\|_{\tilde{L}_T^p}, \|F\|_{\tilde{X}_T^{\sigma,p}}\}, \end{aligned}$$

for any $x \in \mathbb{R}$, $t \in (0, T)$.

Hence for any $z \in \mathbb{R}$,

$$\|g\|_{L_T^p([z, z+1])} + \|g\|_{X_T^{\sigma,p}([z, z+1])} \lesssim \|g\|_{L_T^\infty([z, z+1])} \lesssim \min\{\|F\|_{\tilde{L}_T^p}, \|F\|_{\tilde{X}_T^{\sigma,p}}\}.$$

Similarly, one can check that $\sup_{0 < s < t < T} s^{\sigma+\alpha} \frac{\|g(t) - g(s)\|_{\tilde{L}^p}}{(t-s)^\alpha} \lesssim \|F\|_{\tilde{X}_T^{\sigma,p}}$. This completes the proof. $\blacksquare$

The following is a local version of Lemma 3.1.

Lemma 5.3 *Let f be a solution to (3.1) with initial data $f_0 = \partial_x \bar{f}_0$. There exists $T > 0$ such that*

$$\sum_{\star \in \{\tilde{L}_T^2, \tilde{X}_T^{\frac{1}{2}, 2}\}} \|f\|_{\star} + \sum_{\star \in \{\tilde{L}_T^{\frac{6}{5}}, \tilde{X}_T^{\frac{5}{5}, \frac{6}{5}}\}} \|\partial_x f\|_{\star} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}} + \sum_{\star \in \{\tilde{L}_T^{\frac{6}{5}}, \tilde{X}_T^{\frac{5}{5}, \frac{6}{5}}\}} \|F\|_{\star} + T^{\frac{1}{4}} \sum_{\star \in \{\tilde{L}_T^1, \tilde{X}_T^{1,1}\}} \|R\|_{\star}.$$

Moreover, if $R = 0$, we have

$$\sum_{\star \in \{\tilde{L}_T^2, \tilde{X}_T^{\frac{1}{2}, 2}, \tilde{X}_T^{\frac{3}{4}, \infty}\}} \|\partial_x f\|_{\star} \lesssim \|f_0\|_{\tilde{L}^2} + \sum_{\star \in \{\tilde{L}_T^2, \tilde{X}_T^{\frac{1}{2}, 2}, \tilde{X}_T^{\frac{3}{4}, \infty}\}} \|F\|_{\star}.$$

Proof. Denote $\mathbb{I}_\varepsilon = \cup_{n=1}^\infty [a_n - \varepsilon, a_n + \varepsilon]$. Fix $r > 100$ such that $\mathbb{I}_\varepsilon \subset B_{r/10} = [-r/10, r/10]$. We consider a compact interval $\mathbb{K}$, which is either $B_{r/2}$ or any $[z - \frac{1}{2}, z + \frac{1}{2}] \subset \mathbb{R} \setminus B_{r/2}$. We divide the solution into two parts $f = f_1 + f_2$, such that

$$\begin{cases} \partial_x f_1 - \partial_x(\phi(x)\partial_x f_1) = \partial_x(F\chi) + R\chi, \\ f_1(0, x) = \partial_x(\bar{f}_0(x)\chi(x)), \end{cases} \quad \begin{cases} \partial_x f_2 - \partial_x(\phi(x)\partial_x f_2) = \partial_x(F(1 - \chi)) + R(1 - \chi), \\ f_2(0, x) = \partial_x(\bar{f}_0(x)(1 - \chi)(x)). \end{cases}$$

Here χ is a smooth cutoff function satisfying

$$\begin{cases} \mathbf{1}_{B_r} \leq \chi \leq \mathbf{1}_{B_{2r}}, & \text{if } \mathbb{K} = B_{r/2}, \\ \mathbf{1}_{[z-1, z+1]} \leq \chi \leq \mathbf{1}_{[z-2, z+2]}, & \text{if } \mathbb{K} = [z - \frac{1}{2}, z + \frac{1}{2}] \subset \mathbb{R} \setminus B_{r/2}. \end{cases}$$

We first consider the main term f_1 . Applying Lemma 3.1 to obtain that

$$\sum_{\star \in \{L_T^2, X_T^{\frac{1}{2}, 2}\}} \|f_1\|_\star + \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{5}, \frac{6}{5}}\}} \|\partial_x f_1\|_\star \lesssim \|\bar{f}_0\chi\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}} + \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{5}, \frac{6}{5}}\}} \|F\chi\|_\star + T^{\frac{1}{4}} \sum_{\star \in \{L_T^1, X_T^{1, 1}\}} \|R\chi\|_\star.$$

Note that χ is compactly supported. Hence $\|g\chi\|_{L^p} \lesssim \|g\|_{\tilde{L}^p}$. Then we obtain

$$\sum_{\star \in \{L_T^2, X_T^{\frac{1}{2}, 2}\}} \|f_1\|_\star + \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{5}, \frac{6}{5}}\}} \|\partial_x f_1\|_\star \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}} + \sum_{\star \in \{L_T^{\frac{6}{5}}, X_T^{\frac{5}{5}, \frac{6}{5}}\}} \|F\|_\star + T^{\frac{1}{4}} \sum_{\star \in \{\tilde{L}_T^1, \tilde{X}_T^{1, 1}\}} \|R\|_\star. \quad (5.9)$$

Similarly, when $R = 0$ we have

$$\sum_{\star \in \{\tilde{L}_T^2, \tilde{X}_T^{\frac{1}{2}, 2}, \tilde{X}_T^{\frac{3}{4}, \infty}\}} \|\partial_x f_1\|_\star \lesssim \|\bar{f}_0\|_{\tilde{L}^2} + \sum_{\star \in \{\tilde{L}_T^2, \tilde{X}_T^{\frac{1}{2}, 2}, \tilde{X}_T^{\frac{3}{4}, \infty}\}} \|F\|_\star.$$

Then we estimate the perturbation part f_2 . Note that for any $y \in \text{supp}(1 - \chi)$ and any $x \in \mathbb{K}$, there holds $|x - y| \geq \frac{1}{2}$, which removes the singularity of kernel. The good decay property of the kernel helps us to control local norms (see Lemma 5.2). We consider $\partial_x f_{2,N} = \int_0^t \partial_1^2 \mathbf{H}(t - \tau, x - y, x) F(\tau, y) (1 - \chi(y)) dy d\tau$ for an example. Other terms can be done similarly. Applying Lemma 5.2 we obtain that

$$\sum_{\star \in \{\tilde{L}_T^p(\mathbb{K}), \tilde{X}_T^{\sigma, p}\}} \|\partial_x f_{2,N}\|_\star \lesssim \sum_{\star \in \{\tilde{L}_T^p, \tilde{X}_T^{\sigma, p}\}} \|F\|_\star, \quad p \geq 1.$$

Similarly, we obtain that

$$\sum_{\star \in \{\tilde{L}_T^2, \tilde{X}_T^{\frac{1}{2}, 2}\}} \|f_2\|_\star + \sum_{\star \in \{\tilde{L}_T^{\frac{6}{5}}, \tilde{X}_T^{\frac{5}{5}, \frac{6}{5}}\}} \|\partial_x f_2\|_\star \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}} + \sum_{\star \in \{\tilde{L}_T^{\frac{6}{5}}, \tilde{X}_T^{\frac{5}{5}, \frac{6}{5}}\}} \|F\|_\star + T^{\frac{1}{4}} \sum_{\star \in \{\tilde{L}_T^1, \tilde{X}_T^{1, 1}\}} \|R\|_\star,$$

and

$$\sum_{\star \in \{\tilde{L}_T^2, \tilde{X}_T^{\frac{1}{2}, 2}, \tilde{X}_T^{\frac{3}{4}, \infty}\}} \|\partial_x f_2\|_\star \lesssim \|\bar{f}_0\|_{\tilde{L}^2} + \sum_{\star \in \{\tilde{L}_T^2, \tilde{X}_T^{\frac{1}{2}, 2}, \tilde{X}_T^{\frac{3}{4}, \infty}\}} \|F\|_\star, \quad \text{if } R = 0.$$

Combining this with (5.9), we obtain the result. $\blacksquare$

Denote

$$\|(w, \vartheta)\|_{\tilde{Z}_T} := \sum_{\star \in \{\tilde{L}_T^2, \tilde{X}_T^{\frac{1}{2}, 2}, \tilde{X}_T^{\frac{3}{4}, \infty}\}} \|\partial_x w\|_\star + \sum_{\star \in \{\tilde{L}_T^2, \tilde{X}_T^{\frac{1}{2}, 2}\}} \|\vartheta\|_\star + \sum_{\star \in \{\tilde{L}_T^{\frac{6}{5}}, \tilde{X}_T^{\frac{5}{5}, \frac{6}{5}}\}} \|\partial_x \vartheta\|_\star.$$

Following the proof of Theorem 5.1, we obtain the following result.

Theorem 5.2 *There exists $\varepsilon_0 > 0$ such that if initial data (v_0, u_0, θ_0) satisfies (1.7) and (4.2), then the system (1.3) admits a unique local solution (v, u, θ) in $[0, T]$ satisfying*

$$\inf_{t \in [0, T]} \inf_x v(t, x) \geq \frac{\lambda_0}{2}, \quad \|v\|_{Y_T} \leq 2\|v_0\|_{L^\infty}, \quad \|(u, \theta)\|_{\tilde{Z}_T} \leq B.$$

for some $T, B > 0$.

6 Appendix

Consider the heat equation

$$\partial_t f - \partial_x^2 f = \partial_x F + R, \quad f(0, x) = \partial_x \bar{f}_0(x) + \tilde{f}_0(x). \quad (6.1)$$

We introduce the following lemma.

Lemma 6.1 *Let f be a solution to (6.1), for any $T \in (0, 1)$, there holds*

$$\begin{aligned} \|f\|_{L_T^2} + \|\partial_x f\|_{L_T^{\frac{6}{5}}} &\lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}} + T^{\frac{1}{3}} \|\tilde{f}_0\|_{L^{\frac{6}{5}}} + \|F\|_{L_T^{\frac{6}{5}}} + T^{\frac{1}{4}} \|R\|_{L_T^1}, \\ \|f\|_{L_T^6} + \|\partial_x f\|_{L_T^2} &\lesssim \|\partial_x \bar{f}_0\|_{L^2} + \|\tilde{f}_0\|_{L^2} + \|F\|_{L_T^2} + T^{\frac{1}{2}} \|R\|_{L_T^2}. \end{aligned}$$

Proof. The solution has formula

$$\begin{aligned} f(t, x) &= \int_{\mathbb{R}} \partial_x \mathbf{K}(t, x - y) \bar{f}_0(y) dy + \int_{\mathbb{R}} \mathbf{K}(t, x - y) \tilde{f}_0(y) dy \\ &\quad + \int_0^t \int_{\mathbb{R}} \partial_x \mathbf{K}(t - \tau, x - y) F(\tau, y) dy d\tau + \int_0^t \int_{\mathbb{R}} \mathbf{K}(t - \tau, x - y) R(\tau, y) dy d\tau \\ &:= f_{L1}(t, x) + f_{L2}(t, x) + f_N(t, x) + f_R(t, x). \end{aligned}$$

We first estimate f_{L1} . By Parseval's identity and Sobolev inequality, it is easy to check that

$$\begin{aligned} \|f_{L1}\|_{L_\infty^2}^2 &= \int_0^\infty \int_{\mathbb{R}} |\xi|^2 |e^{-|\xi|^2 \tau} \mathcal{F}(\bar{f}_0)(\xi)|^2 d\xi d\tau = \frac{1}{2} \int_{\mathbb{R}} |\mathcal{F}(\bar{f}_0)(\xi)|^2 d\xi = \frac{1}{2} \|\bar{f}_0\|_{L^2}^2 \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3}, \frac{6}{5}}}^2, \\ \|\partial_x f_{L1}\|_{L_\infty^2}^2 &= \int_0^\infty \int_{\mathbb{R}} |\xi|^4 |e^{-|\xi|^2 \tau} \mathcal{F}(\bar{f}_0)(\xi)|^2 d\xi d\tau = \frac{1}{2} \int_{\mathbb{R}} |\xi|^2 |\mathcal{F}(\bar{f}_0)(\xi)|^2 d\xi = \frac{1}{2} \|\partial_x \bar{f}_0\|_{L^2}^2. \end{aligned} \quad (6.2)$$

Let $(l, p) = (1, \frac{6}{5})$ or $(0, 6)$. Note that

$$\partial_x^l f_{L1}(t, x) = \int_{\mathbb{R}} \partial_x^{1+l} \mathbf{K}(t, x - y) (\bar{f}_0(y) - \bar{f}_0(x)) dy.$$

Set $K_0(t, x) = |\partial_x^{1+l} \mathbf{K}(t, x)|$. Denote $\Delta_z g(x) = g(x) - g(x - z)$. Then by a change of variable,

$$|\partial_x^l f_{L1}(t, x)| \lesssim \int_{\mathbb{R}} K_0(t, z) |\Delta_z \bar{f}_0(x)| dz.$$

Then we have

$$\begin{aligned} \|\partial_x^l f_{L1}\|_{L_T^p}^p &= \int_0^\infty \int_{\mathbb{R}} \left(\int_{\mathbb{R}} K_0(t, z) |\Delta_z \bar{f}_0(x)| dz \right)^p dx dt \\ &\lesssim \int_0^\infty \left(\int_{\mathbb{R}} K_0(t, z) \|\Delta_z \bar{f}_0\|_{L_x^p} dz \right)^p dt \\ &\lesssim \int_0^\infty \left(\int_{\mathbb{R}} K_0(t, z)^{\frac{p}{p-1}} (1 + |z|^2 t^{-1})^{\frac{\gamma p}{p-1}} dz \right)^{p-1} \left(\int_{\mathbb{R}} \frac{\|\Delta_z \bar{f}_0\|_{L_x^p}^p}{(1 + |z|^2 t^{-1})^{\gamma p}} dz \right) dt, \end{aligned}$$

where we applied Minkowski's inequality in the first inequality, and Hölder's inequality in the second inequality. It is easy to check that

$$\left(\int_{\mathbb{R}} K_0(t, z)^{\frac{p}{p-1}} (1 + |z|^2 t^{-1})^{\frac{\gamma p}{p-1}} dz \right)^{p-1} \lesssim t^{\frac{-(l+1)p-1}{2}},$$

where we fix $\gamma = \frac{1+l}{2}$. Hence,

$$\begin{aligned} \|\partial_x^l f_{L1}(t, x)\|_{L_T^p}^p &\lesssim \int_0^\infty t^{\frac{-(l+1)p-1}{2}} \left(\int_{\mathbb{R}} \frac{\|\Delta_z \bar{f}_0(x)\|_{L_x^p}^p}{(1 + |z|^2 t^{-1})^{\gamma p}} dz \right) dt \\ &\lesssim \int_{\mathbb{R}} \frac{\|\Delta_z \bar{f}_0(x)\|_{L_x^p}^p}{|z|^{(1+l)p-1}} dz = c \|\bar{f}_0\|_{\dot{W}^{\frac{p-2}{p}+l, p}}^p. \end{aligned} \quad (6.3)$$

Hence we obtain that

$$\|f_{L1}\|_{L_T^6} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{2}{3},6}} \lesssim \|\partial_x \bar{f}_0\|_{L^2}, \quad \|\partial_x f_{L1}\|_{L_\infty^{\frac{6}{5}}} \lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}}. \quad (6.4)$$

Combining this with (6.2) and (6.4), one has

$$\begin{aligned} \|f_{L1}\|_{L_T^2} + \|\partial_x f_{L1}\|_{L_T^{\frac{6}{5}}} &\lesssim \|\bar{f}_0\|_{\dot{W}^{\frac{1}{3},\frac{6}{5}}}, \\ \|f_{L1}\|_{L_T^6} + \|\partial_x f_{L1}\|_{L_T^2} &\lesssim \|\partial_x \bar{f}_0\|_{L^2}. \end{aligned} \quad (6.5)$$

Then we estimate f_{L2} . By Parseval's identity, we obtain that

$$\|\partial_x f_{L2}\|_{L_T^2} \lesssim \int_0^\infty \int_{\mathbb{R}} |\xi|^2 |e^{-|\xi|^2 \tau} \mathcal{F}(\tilde{f}_0)(\xi)|^2 d\xi d\tau = \frac{1}{2} \int_{\mathbb{R}} |\mathcal{F}(\tilde{f}_0)(\xi)|^2 d\xi = \frac{1}{2} \|\tilde{f}_0\|_{L^2}^2.$$

Moreover, by Hölder's inequality and Young's inequality,

$$\begin{aligned} \|f_{L2}\|_{L_T^2} &\lesssim \|\mathbf{K}\|_{L_{t,T}^2 L_x^{\frac{3}{2}}} \|\tilde{f}_0\|_{L^{\frac{6}{5}}} \lesssim T^{\frac{1}{3}} \|\tilde{f}_0\|_{L^{\frac{6}{5}}}, \\ \|f_{L2}\|_{L_T^6} &\lesssim \left\| \int_{\mathbb{R}} \|\mathbf{K}(x-y)\|_{L_{t,T}^6} |\tilde{f}_0(y)| dy \right\|_{L_x^6} \lesssim \left\| \int_{\mathbb{R}} |\tilde{f}_0(y)| \frac{dy}{|x-y|^{\frac{2}{3}}} \right\|_{L_x^6} \lesssim \|\tilde{f}_0\|_{L^2}, \\ \|\partial_x f_{L2}\|_{L_T^{\frac{6}{5}}} &\lesssim \|\partial_x \mathbf{K}\|_{L_{t,T}^{\frac{6}{5}} L_x^1} \|\tilde{f}_0\|_{L^{\frac{6}{5}}} \lesssim T^{\frac{1}{3}} \|\tilde{f}_0\|_{L^{\frac{6}{5}}}. \end{aligned}$$

Hence we obtain that

$$\begin{aligned} \|f_{L2}\|_{L_T^2} + \|\partial_x f_{L2}\|_{L_T^{\frac{6}{5}}} &\lesssim T^{\frac{1}{3}} \|\tilde{f}_0\|_{L^{\frac{6}{5}}}, \\ \|f_{L2}\|_{L_T^6} + \|\partial_x f_{L2}\|_{L_T^2} &\lesssim \|\tilde{f}_0\|_{L^2}. \end{aligned} \quad (6.6)$$

Then we estimate the forced term f_N . We can write

$$\partial_t f_N - \partial_x^2 f_N = \partial_x F_0, \quad \text{in } \mathbb{R} \times \mathbb{R}, \quad \text{where } F_0 = F \mathbf{1}_{t>0}.$$

We take Fourier transform in (t, x) and obtain that

$$f_N = \mathcal{F}_{t,x}^{-1} \left(\frac{\xi \mathcal{F}_T(F_0)(\tau, \xi)}{i\tau + \xi^2} \right),$$

where we denote $\mathcal{F}_{t,x}$ the Fourier transform with respect to (t, x) , and $\mathcal{F}_T^{-1}$ its inverse. It follows from the parabolic Calderon-Zygmund theory (see [17]) that

$$\|\partial_x f_N\|_{L_T^p} \lesssim \|F_0\|_{L_T^p} \lesssim \|F\|_{L_T^p}, \quad p \in (1, \infty).$$

On the other hand, we have $|\partial_x \mathbf{K}(t, x)| \lesssim (t^{\frac{1}{2}} + |x|)^{-2}$. Then by Young's inequality,

$$\begin{aligned} &\left\| \int_0^t \int_{\mathbb{R}} \partial_x \mathbf{K}(t-\tau, x-y) F(\tau, y) dy d\tau \right\|_{L_T^p} \\ &\lesssim \left\| \int_{\mathbb{R}} \|\partial_x \mathbf{K}(\cdot, x-y)\|_{L_{t,T}^{\frac{3}{2}}} \|F(y)\|_{L_{t,T}^{\frac{3p}{3+p}}} dy \right\|_{L_x^p} \lesssim \left\| \int_{\mathbb{R}} \|F(y)\|_{L_{t,T}^{\frac{3p}{3+p}}} \frac{dy}{|x-y|^{\frac{2}{3}}} \right\|_{L_x^p} \\ &\lesssim \|F\|_{L_T^{\frac{3p}{3+p}}}. \end{aligned}$$

Hence we obtain that for $l = 0, 1$,

$$\|\partial_x^l f_N\|_{L_T^p} = \left\| \int_0^t \int_{\mathbb{R}} \partial_x^{1+l} \mathbf{K}(t-\tau, x-y) F(\tau, y) dy d\tau \right\|_{L_T^p} \lesssim \|F\|_{L_T^{\frac{3p}{3+(1-l)p}}}. \quad (6.7)$$

Finally, we estimate f_R . By Hölder's inequality we have

$$\|\partial_x^l f_R\|_{L_T^p} \lesssim \|\partial_x^l \mathbf{K}\|_{L_T^p} \|R\|_{L_T^1} \lesssim T^{\frac{1}{4}} \|R\|_{L_T^1}, \quad (l, p) \in \{(0, 2), (1, 6/5)\},$$

and

$$\|\partial_x^l f_R\|_{L_T^p} \lesssim \|\partial_x^l \mathbf{K}\|_{L_T^{\frac{2p}{2+p}}} \|R\|_{L_T^1} \lesssim T^{\frac{1}{2}} \|R\|_{L_T^2}, \quad (l, p) \in \{(0, 6), (1, 2)\}.$$

Combining this with (6.5), (6.6) and (6.7), we finish the proof. $\blacksquare$

Consider the parabolic equation with jump coefficients,

$$\begin{aligned} \partial_t f^\pm - c_\pm f_{xx}^\pm &= F, \quad \text{in } (0, T) \times \mathbb{R}^\pm, \\ f^+(t, 0) &= f^-(t, 0), \quad c_+ \partial_x f^+(t, 0) - c_- \partial_x f^-(t, 0) = 0, \quad t \in (0, T), \\ f(0, x) &= f_0(x), \quad x \in \mathbb{R}. \end{aligned} \quad (6.8)$$

Lemma 6.2 *Solution to (6.8) has formula $f = f^+ \mathbf{1}_{x \geq 0} + f^- \mathbf{1}_{x < 0}$, where*

$$f^\pm(t, x) = f_M^\pm(t, x) + f_B^\pm(t, x).$$

with

$$\begin{aligned} f_M^\pm(t, x) &= \int_{\mathbb{R}^\pm} (\mathbf{K}(c_\pm t, x - y) - \mathbf{K}(c_\pm t, x + y)) f_0(y) dy \\ &\quad + \int_0^t \int_{\mathbb{R}^\pm} (\mathbf{K}(c_\pm(t - \tau), x - y) - \mathbf{K}(c_\pm(t - \tau), x + y)) F(\tau, y) dy d\tau, \end{aligned} \quad (6.9)$$

$$f_B^\pm(t, x) = \frac{-2\sqrt{c_\pm}}{\sqrt{c_+} + \sqrt{c_-}} \int_0^t \mathbf{K}(c_\pm(t - \tau), x) (c_+ \partial_x f_M^+(\tau, 0) - c_- \partial_x f_M^-(\tau, 0)) d\tau. \quad (6.10)$$

Proof. By superposition principle, we have $f^\pm(t, x) = f_M^\pm(t, x) + f_B^\pm(t, x)$, where $f_M^\pm$ solves the following equation on half space

$$\begin{cases} \partial_t f_M^+ - c_+ \partial_x^2 f_M^+ = F, & \text{in } (0, T) \times \mathbb{R}^+, \\ f_M^+(0, x) = f_0(x), & x \in \mathbb{R}^+, \\ f_M^+(t, 0) = 0, & t \in (0, T). \end{cases} \quad \begin{cases} \partial_t f_M^- - c_- \partial_x^2 f_M^- = F, & \text{in } (0, T) \times \mathbb{R}^-, \\ f_M^-(0, x) = f_0(x), & x \in \mathbb{R}^-, \\ f_M^-(t, 0) = 0, & t \in (0, T). \end{cases}$$

And $f_B^\pm$ solves the system

$$\begin{aligned} \partial_t f_B^\pm - c_\pm \partial_x^2 f_B^\pm &= 0, \quad \text{in } (0, T) \times \mathbb{R}^\pm, \\ f_B^+(0, x) &= 0 \text{ in } \mathbb{R}^+, \quad f_B^-(0, x) = 0 \text{ in } \mathbb{R}^-, \\ f_B^+(t, 0) &= f_B^-(t, 0), \quad t \in (0, T), \\ c_+ \partial_x f_B^+(t, 0) - c_- \partial_x f_B^-(t, 0) &= -(c_+ \partial_x f_M^+(t, 0) - c_- \partial_x f_M^-(t, 0)), \quad t \in (0, T). \end{aligned} \quad (6.11)$$

Indeed, integrate by parts we have

$$\begin{aligned} &\int_0^t \int_{\mathbb{R}^+} (\mathbf{K}(c_+(t - \tau), x - y) - \mathbf{K}(c_+(t - \tau), x + y)) F(\tau, y) dy d\tau \\ &= \int_0^t \int_{\mathbb{R}^+} (\mathbf{K}(c_+(t - \tau), x - y) - \mathbf{K}(c_+(t - \tau), x + y)) (\partial_t f_M^+ - c_+ \partial_x^2 f_M^+)(\tau, y) dy d\tau \\ &= f_M^+(t, x) - \int_{\mathbb{R}^+} (\mathbf{K}(c_+ t, x - y) - \mathbf{K}(c_+ t, x + y)) f_0(y) dy \\ &\quad + \int_0^t \int_{\mathbb{R}^+} \partial_t (\mathbf{K}(c_+(t - \tau), x - y) - \mathbf{K}(c_+(t - \tau), x + y)) f_M^+(\tau, y) dy d\tau \\ &\quad - c_+ \int_0^t \int_{\mathbb{R}^+} \partial_x^2 (\mathbf{K}(c_+(t - \tau), x - y) - \mathbf{K}(c_+(t - \tau), x + y)) f_M^+(\tau, y) dy d\tau \\ &= f_M^+(t, x) - \int_{\mathbb{R}^+} (\mathbf{K}(c_+ t, x - y) - \mathbf{K}(c_+ t, x + y)) f_0(y) dy, \end{aligned}$$

where we use the fact that

$$\partial_t \mathbf{K}(c_+ t, x) - c_+ \partial_x^2 \mathbf{K}(c_+ t, x) = 0, \quad \forall t > 0, x \in \mathbb{R}.$$

We also get similar estimate for f_M^- . Hence we obtain formula (6.9).

It remains to check that formula (6.10) satisfies equation (6.11). For simplicity, we only check the Neumann boundary condition

$$c_+ \partial_x f_B^+(t, 0) - c_- \partial_x f_B^-(t, 0) = -(c_+ \partial_x f_M^+(t, 0) - c_- \partial_x f_M^-(t, 0)) := h(t), \quad t \in (0, T). \quad (6.12)$$

And other conditions are trivial. Note that

$$\begin{aligned} c_+ \partial_x f_B^+(t, 0) - c_- \partial_x f_B^-(t, 0) &= \frac{2c_+^{\frac{3}{2}}}{\sqrt{c_+} + \sqrt{c_-}} \lim_{x \rightarrow 0^+} \int_0^t \partial_x \mathbf{K}(c_+(t - \tau), x) h(\tau) d\tau \\ &\quad - \frac{2c_-^{\frac{3}{2}}}{\sqrt{c_+} + \sqrt{c_-}} \lim_{x \rightarrow 0^-} \int_0^t \partial_x \mathbf{K}(c_-(t - \tau), x) h(\tau) d\tau. \end{aligned} \quad (6.13)$$

Note that the heat kernel has scaling $\mathbf{K}(t, x) = t^{-\frac{1}{2}} \mathbf{K}(1, \frac{x}{\sqrt{t}})$.

$$\begin{aligned} \int_0^t \partial_x \mathbf{K}(c_+(t - \tau), x) h(\tau) d\tau &= - \int_0^t \frac{x}{2c_+ \tau} K(c_+ \tau, x) h(t - \tau) d\tau \\ &= - \int_0^t \frac{x}{2(c_+ \tau)^{\frac{3}{2}}} K(1, \frac{x}{\sqrt{c_+ \tau}}) h(t - \tau) d\tau. \end{aligned}$$

Let $\tilde{\tau} = \frac{x}{\sqrt{c_+ \tau}}$, we obtain that

$$\int_0^t \partial_x \mathbf{K}(c_+(t - \tau), x) h(\tau) d\tau = c_+^{-1} \int_{\frac{x}{\sqrt{c_+ t}}}^{\infty} \mathbf{K}(1, \tilde{\tau}) h(t - \frac{x^2}{c_+ \tilde{\tau}^2}) d\tilde{\tau}.$$

We can write

$$\int_{\tilde{t}}^{\infty} \mathbf{K}(1, \tilde{\tau}) h(t - \frac{x^2}{c_+ \tilde{\tau}^2}) d\tilde{\tau} = \left(\int_{\tilde{t}^{1/2}}^{\infty} + \int_{\tilde{t}}^{\tilde{t}^{1/2}} \right) \mathbf{K}(1, \tilde{\tau}) h(t - \frac{x^2}{c_+ \tilde{\tau}^2}) d\tilde{\tau},$$

where we denote $\tilde{t} = \frac{x}{\sqrt{c_+ t}}$. We have $\tilde{t} \rightarrow 0$ when $x \rightarrow 0$, and the integral on $[\tilde{t}, \tilde{t}^{1/2}]$ tends to 0. Hence,

$$\lim_{x \rightarrow 0^+} \int_0^t \partial_x \mathbf{K}(c_+(t - \tau), x) h(\tau) d\tau = c_+^{-1} \int_0^{\infty} \mathbf{K}(1, \tilde{\tau}) d\tilde{\tau} h(t) = \frac{1}{2c_+} h(t).$$

Similarly, one has

$$\lim_{x \rightarrow 0^-} \int_0^t \partial_x \mathbf{K}(c_-(t - \tau), x) h(\tau) d\tau = -\frac{1}{2c_-} h(t). \quad (6.14)$$

Then we obtain (6.12) from (6.13)-(6.14). This completes the proof. $\blacksquare$

Consider the following parabolic equation with variable coefficient,

$$\begin{aligned} \partial_t f - \partial_x(\phi(x) f_x) &= F, \quad (t, x) \in (0, T) \times \mathbb{R}, \\ f(0, x) &= f_0(x). \end{aligned} \quad (6.15)$$

Lemma 6.3 *the Solution f to (6.15) satisfies*

$$\begin{aligned} f(t, x) &= \int_{\mathbb{R}} \mathbf{H}(t, x - y, x) f_0(y) dy + \int_0^t \int_{\mathbb{R}} \mathbf{H}(t - \tau, x - y, x) F(\tau, y) dy d\tau \\ &\quad - \int_0^t \int_{\mathbb{R}} \partial_1 \mathbf{H}(t - \tau, x - y, x) (\phi(x) - \phi(y)) f_x(\tau, y) dy d\tau, \end{aligned}$$

where we denote $\mathbf{H}(t, x, y) = \mathbf{K}(\phi(y)t, x)$, $\partial_1 \mathbf{H}(t, x, y) = \partial_x \mathbf{H}(t, x, y)$.

Proof. Integrate by parts we obtain

$$\begin{aligned} &\int_0^t \int_{\mathbb{R}} \mathbf{H}(t - \tau, x - y, x) F(\tau, y) dy d\tau \\ &= \int_0^t \int_{\mathbb{R}} \mathbf{H}(t - \tau, x - y, x) (\partial_t f - \partial_x(\phi f_x))(\tau, y) dy d\tau \\ &= f(t, x) - \int_{\mathbb{R}} \mathbf{H}(t, x - y, x) f_0(y) dy + \int_0^t \int_{\mathbb{R}} \partial_t \mathbf{H}(t - \tau, x - y, x) f(\tau, y) dy d\tau \\ &\quad + \int_0^t \int_{\mathbb{R}} \partial_1 \mathbf{H}(t - \tau, x - y, x) (\phi(x) - \phi(y)) f_x(\tau, y) dy d\tau \\ &\quad - \phi(x) \int_0^t \int_{\mathbb{R}} \partial_1^2 \mathbf{H}(t - \tau, x - y, x) f(\tau, y) dy d\tau. \end{aligned}$$

It is easy to check that

$$\partial_t \mathbf{H}(t - \tau, x - y, x) - \phi(x) \partial_1^2 \mathbf{H}(t - \tau, x - y, x) = 0.$$

Then we obtain the result. ■

Acknowledgements: Q.H.N. is supported by the Academy of Mathematics and Systems Science, Chinese Academy of Sciences startup fund, and the National Natural Science Foundation of China (No. 12050410257 and No. 12288201) and the National Key R&D Program of China under grant 2021YFA1000800.

References

- [1] Ke Chen, Q. H. Nguyen. *The Peskin problem with $\dot{B}_{\infty, \infty}^{-1}$ initial data*. arXiv:2107.13854.
- [2] E. Feireisl. *Dynamics of viscous compressible fluids*. Oxford Lecture Series in Mathematics and Its Applications. Oxford University Press, Oxford, 2004.
- [3] G. H. Hardy, J. E. Littlewood and G. Pólya. *Inequalities* (Second ed.). Cambridge, 1952.
- [4] D. Hoff. *Global well-posedness of the Cauchy problem for the Navier-Stokes equations of nonisentropic flow with discontinuous initial data*. Journal of Differential Equations, 95(1):33-74, 1992.
- [5] N. Itaya. *On the Cauchy problem for the system of fundamental equations describing the movement of compressible viscous fluid*. Kōdai Mathematical Seminar Reports, 23(1):60-120, 1971.
- [6] N. Itaya. *On the initial value problem of the motion of compressible viscous fluid, especially on the problem of uniqueness*. J. Math. Kyoto Univ, 16(2): 413-427, 1976.
- [7] T. Jin, V. Maz'ya and J. V. Schaftingen. *Pathological solutions to elliptic problems in divergence form with continuous coefficients*. Comptes Rendus Mathématique, 347(13-14): 773-778, 2009.

- [8] S. Kawashima. *Large-time behaviour of solutions to hyperbolic-parabolic systems of conservation laws and applications*. Proceedings of the Royal Society of Edinburgh Section A Mathematics, 106(1-2):169-194, 1987.
- [9] A. V. Kazhikhov, V. V. Shelukhin. *Unique global solution with respect to time of initial-boundary value problems for one-dimensional equations of a viscous gas*. Journal of Applied Mathematics and Mechanics, 41(2):282-291, 1977.
- [10] P. L. Lions. *Mathematical topics in fluid mechanics*. Vol. 2. Compressible models. Oxford Lecture Series in Mathematics and Its Applications. Oxford University Press, New York, 1998.
- [11] T. P. Liu and S. H. Yu. *Navier-Stokes equations in gas dynamics: Green's function, singularity, and well-posedness*. Communications on Pure and Applied Mathematics, 75:223-348, 2022.
- [12] A. Matsumura, T. Nishida. *The initial value problem for the equations of motion of viscous and heat-conductive gases*. Journal of Mathematics of Kyoto University, 20(1):67-104, 1980.
- [13] H. M. Nguyen and Q. H. Nguyen. *Discreteness of interior transmission eigenvalues revisited*. Calculus of Variations and Partial Differential Equations, 56(2):51, 2017.
- [14] H. M. Nguyen, and Q. H. Nguyen. *The Weyl law of transmission eigenvalues and the completeness of generalized transmission eigenfunctions*. Journal of Functional Analysis. 281.8:109-146, 2021.
- [15] J. Nash. *Continuity of solutions of parabolic and elliptic equations*. Amer. J. Math. 80 (1958), 931-954. doi:10.2307/2372841.
- [16] J. Smoller. *Shock waves and Reaction-Diffusion Equations*. Springer. 1994.
- [17] E. M. Stein. *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integral*, 43. Princeton University Press, Princeton. 1993.
- [18] H. Wang, S. H. Yu, and X. Zhang. *Global well-posedness of compressible Navier-Stokes equation with $BV \cap L^1$ initial data*. Arch. Rational Mech. Anal. 2022. <https://doi.org/10.1007/s00205-022-01787-z>.