

# Regularity and drift by Osgood vector fields

Joonhyun La

June 30, 2022

## Abstract

We consider the problem of loss and propagation of regularity of transport equation with Osgood vector field. As an application, we obtain a quantitative stability estimate for 2D incompressible Euler equation with generalized Yudovich initial data.

## 1 Introduction

We consider the transport equation

$$\begin{cases} \partial_t \theta + u \cdot \nabla_x \theta = 0, \\ \theta(0, \cdot) = \theta_0, \end{cases} \quad (1)$$

where  $u$  is a divergence-free field. It is well-known that if  $u$  has an Osgood modulus of continuity, that is,

$$|u(x, t) - u(y, t)| \leq C\varphi(|x - y|), \quad (2)$$

for some increasing function  $\varphi : (0, m) \rightarrow (0, \infty)$  satisfying

$$\begin{aligned} \lim_{r \rightarrow 0^+} \varphi(r) &= 0, \\ \int_0^m \frac{dr}{\varphi(r)} &= \infty, \end{aligned} \quad (3)$$

then the associated ODE for the flow

$$\begin{cases} \frac{d}{dt} \phi(x, t) = u(\phi(x, t), t), \\ \phi(x, 0) = x \end{cases} \quad (4)$$

has the unique solution (e.g. [4], [11]) and there is unique weak integrable solution for (1), given by

$$\theta(x, t) = \theta_0(\phi^{-1}(x, t)) \quad (5)$$

([3], [6]) In that regard, in terms of existence and uniqueness of solution all Osgood modulus of continuities are equally good: they generate unique flows.

What about the propagation of regularity? Suppose that  $\theta_0 \in \dot{H}^\sigma(\mathbb{R}^d)$  for some  $\sigma > 0$ . Is  $\theta(t) \in \dot{H}^\sigma(\mathbb{R}^d)$ ? If  $u$  is not Lipschitz, the answer is negative in general and there is loss of regularity. However, the degree of loss differs depending on the regularity of  $u$ . If  $u$  is log-Lipschitz or better, we have losing estimates, that is,  $\theta(t) \in \dot{H}^{\sigma(t)}(\mathbb{R}^d)$ , for  $\sigma(t) > 0$  (e.g. [4]). On the other hand, if  $u \in \dot{W}^{1,p}$ ,  $p < \infty$ , one can immediately lose all Sobolev regularities: in [2], it has been shown that there is a divergence-free field  $u \in \cap_{1 \leq p < \infty} \dot{W}^{1,p}$  and  $\theta_0 \in H^\sigma(\mathbb{R}^d)$

such that  $\theta(t) \notin \dot{H}^s(\mathbb{R}^d)$  for every  $s > 0$ . On the other hand, logarithm of a derivative can be preserved (e.g. [8], [5], [13])

Osgood vector fields  $u$  which are worse than log-Lipschitz lie between these two well known cases, and one may naturally ask the following questions. First, does Sobolev regularity propagate, even in a losing manner? Second, is there any kind of regularity, better than logarithm of a derivative, which is propagated? In this paper, we attempt to answer these questions.

First, we show that Sobolev regularity will not be propagated in general.

**Theorem 1.** *Let  $d \geq 2$  and  $0 < \sigma < \frac{d}{2}$ . For each admissible growth function  $\Theta$  (see definition 1), there exists a divergence-free vector field  $u$  with a modulus of continuity  $\varphi_\Theta$  (see definition 2) and  $\theta_0 \in H^\sigma(\mathbb{R}^d)$  such that  $\theta(t) \notin H^\sigma(\mathbb{R}^d)$  for every  $t > 0, s > 0$ .*

As  $\varphi_\Theta$  is an Osgood modulus of continuity (Lemma 1), we see that all Sobolev regularity may loss immediately for Osgood velocity fields. Also, one may find  $\theta_0 \in H^\sigma(\mathbb{R}^d)$  for any  $\sigma > 0$ : see Remark 3.

Second, we show that certain regularities propagate, which depend on the modulus of continuity of  $u$ .

**Theorem 2.** *Suppose that  $u$  is a divergence-free vector field with an Osgood modulus of continuity  $\varphi$ , and let  $\mu_0$  be a modulus of continuity.*

1. *Suppose that  $\theta_0$  has modulus of continuity  $\mu_0$ , and  $\theta(t)$  be the solution of (1). Then  $\theta(t)$  has a modulus of continuity  $\mu_0 \circ \mu_{[u]_\varphi, t}$  for  $t \geq 0$  (see (29) for the definition of  $\mu_{[u]_\varphi, t}$ ).*
2. *Let  $T > 0$  and suppose that  $A(u) := \int_0^1 \frac{\mu_0 \circ \mu_{[u]_\varphi, T}(r)}{r} dr < \infty$ . Suppose that there exists an  $g \in L^2(\mathbb{R}^d)$  such that (30) holds. Then for  $t \in [0, T]$ ,*

$$\int_{B_1(0)} \int_{\mathbb{R}^d} \frac{|\theta(x, t) - \theta(x + h, t)|^2}{|h|^d} \mu_0 \circ \mu_{[u]_\varphi, t}(|h|)^{-1} dx dh \leq C \|g\|_{L^2(\mathbb{R}^d)}^2 A(u). \quad (6)$$

As an application, we obtain a quantitative stability result for 2D Euler's equation in generalized Yudovich class with additional regularity for initial data, which improves that of [12]. Let  $n \in \mathbb{Z}_{\geq 1}$ , and  $\Theta_n : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$  with

$$\Theta_n(z) = \log z \log_2 z \cdots \log_{n-1} z$$

for sufficiently large  $z > 1$ . Let  $\omega_{0,1}, \omega_{0,2} \in Y_{\Theta_n}(\mathbb{R}^2)$  with  $u_{0,i} = \nabla^\perp \Delta^{-1} \omega_i \in L^2(\mathbb{R}^2)$  for  $i = 1, 2$ , where

$$Y_{\Theta_n}(\mathbb{R}^2) = \left\{ f \in \cap_{1 \leq p < \infty} L^p(\mathbb{R}^2) \mid \|f\|_{L^p(\mathbb{R}^2)} \leq \Theta_n(p) \right\}, \quad \|f\|_{Y_{\Theta_n}} = \sup_p \frac{\|f\|_{L^p}}{\Theta_n(p)}. \quad (7)$$

Furthermore, suppose that  $\omega_{0,i} \in H^s(\mathbb{R}^2)$  for some  $0 < s \leq 1, i = 1, 2$ . Let  $\omega_i, i = 1, 2$  be solutions of the initial-value problem

$$\begin{cases} \partial_t \omega_i + u_i \cdot \nabla_x \omega_i = 0, \\ u_i = \nabla^\perp \Delta^{-1} \omega_i, \\ \omega_i(0, x) = \omega_{0,i}(x). \end{cases} \quad (8)$$

**Theorem 3.** *If  $\|u_{0,1} - u_{0,2}\|_{L^2}$  is sufficiently small, then there exist constants  $C, C_1, C_2, M > 0, \gamma < 0$  such that*

$$\|\omega_1(t) - \omega_2(t)\|_{L^2}^2 \leq C \left( \mu_{\omega, n, t} \left( \left( 2 + \frac{C \max_i \|\omega_{0,i}\|_{L^2}^2}{\mu_{\omega, n, t}(\|u_{0,1} - u_{0,2}\|_{L^2}^2)} \right)^\gamma \right) \right)^s \quad (9)$$

where  $\mu_{\omega,n,t}$  is defined by

$$\mu_{\omega,n,t}(r) := \frac{C_1}{e_{n-1} \left( (\log_{n-1}(C_2/r))^{1/\exp(tM \max_i \|\omega_{0,i}\|_{Y_{\Theta_n}})} \right)}. \quad (10)$$

**Remark 1.** Theorem (3) recovers the rate obtained in Proposition 18 of [12] for Yudovich case ( $n = 1$ ) and improves the rate for localized Yudovich case. In the case  $n = 1$ , which is Yudovich case  $\omega \in L^\infty$ ,  $\mu_{\omega,n,t}^i(r) = r^{C(t)}$ , by Remark 4. Thus,  $\|\omega_1(t) - \omega_2(t)\|_{L^2}^2 \leq C\|u_{0,1} - u_{0,2}\|_{L^2}^{C(t)}$ , which is the algebraic rate obtained in [12] (see also [7]). On the other hand, when  $n > 1$ , we note that the result in [12] gives the rate of  $1/\log(\mu_{\omega,n,t}^2(\|u_{0,1} - u_{0,2}\|_{L^2}^2))$ . Since  $\mu_{\omega,n,t}^1$  vanishes faster than logarithmic function by Lemma 6, the estimate (9) gives a smaller upper bound.

**Acknowledgement** The author greatly thanks Theo Drivas and In-Jee Jeong for stimulating discussion. Also, the author is deeply grateful to Gianluca Crippa and Anna Mazzucato for encouraging and helpful comments.

## 2 Immediate loss of Sobolev regularity

### 2.1 Growth functions and Osgood modulus of continuity

**Definition 1.** A function  $\Theta : [1, \infty) \rightarrow (0, \infty)$  is called an admissible growth function if  $\Theta$  is a differentiable function which is increasing with  $\lim_{x \rightarrow \infty} \Theta(x) = \infty$ , and there is a constant  $M > 1$ ,  $C > 0$  such that

1.  $\Theta(xy) \leq \Theta(x) + \Theta(y) + C$  for all  $x, y \geq M$ , and
2.  $x\Theta'(x) \leq C\Theta(x)$  for all  $x > M$ .

**Remark 2.** The last two conditions are technical: they are satisfied for many slowly growing functions, and they are useful in obtaining certain bounds, but it would be interesting to obtain the same result with relaxed conditions.

An admissible growth function can grow slowly.

**Proposition 1.**  $\Theta(x) = \log_m x$ ,  $x > e_m(2)$ , which is defined recursively by  $\log_{m+1} x = \log_m(\log x)$ , with  $\log_1 x = \log x$ , and  $e_m = \log_m^{-1}$  for  $m \in \mathbb{N}$  is an admissible growth function ( $\Theta(x)$  for  $x \leq e_m(2)$  can be chosen appropriately.)

*Proof.* It suffices to show the last two conditions. The last condition holds immediately: for  $x > e_m(2)$ ,  $\Theta'(x) = \prod_{j=1}^{m-1} \frac{1}{\log_j(x)} \frac{1}{x}$ , so  $\Theta'(x)x \leq C < C\Theta(x)$ . For the condition  $\Theta(xy) \leq \Theta(x) + \Theta(y) + C$ , we use induction on  $m$ : for  $m = 1$ ,  $\Theta(x) = \log x$  and the condition obviously holds. Suppose that For  $\log_m x$  the condition hold, with  $M = M_m, C = C_m$ . Then for  $m + 1$ , for  $x, y \geq e^{M_m} = M_{m+1}$ , we have

$$\begin{aligned} \Theta(xy) &= \log_m(\log x + \log y) \leq \log_m((M_m + 2)\max(x, y)) \\ &\leq C_m + \log_m(M_m + 2) + \log_{m+1} \max(x, y) \\ &\leq (C_m + \log_m(M_m + 2)) + \log_{m+1} x + \log_{m+1} y = \Theta(x) + \Theta(y) + C_{m+1}. \end{aligned} \quad (11)$$

□

Also, an admissible growth function cannot grow too fast.

**Proposition 2.** *There exists a constant  $C' > 0$  such that  $\Theta(x) \leq C'(\log x + 1)$  for all  $x \geq 1$ .*

*Proof.* This is immediate from the last property.  $\square$

As a consequence,  $\Theta$  is associated with an Osgood modulus of continuity. We use the following modulus of continuity, following [9].

**Lemma 1.** *Let  $\Theta$  be an admissible growth function. The function  $\varphi_\Theta : [0, \infty) \rightarrow [0, \infty)$  defined by*

$$\varphi_\Theta(r) = \begin{cases} r \log(e/r) \Theta(\log(e/r)), & 0 < r < e^{-2}, \\ e^{-2} 3 \Theta(3), & r > e^{-2} \end{cases} \quad (12)$$

*is an Osgood modulus of continuity.*

*Proof.*

$$\int_0^{e^{-2}} \frac{dr}{\varphi_\Theta(r)} = \int_3^\infty \frac{dp}{p \Theta(p)} \geq \int_3^\infty \frac{dp}{C' p (\log p + 1)} = \infty. \quad (13)$$

$\square$

**Definition 2.** *For an admissible growth function  $\Theta$ ,  $\varphi_\Theta$  defined in lemma 1 is called the Osgood modulus of continuity associated with  $\Theta$ .*

As the growth of  $\Theta$  becomes slower,  $\varphi_\Theta$  gets closer to log-Lipschitz. Since the class of admissible growth functions contains plenty of slowly growing functions (Proposition 1), one can find  $\varphi_\Theta$  which is close to log-Lipschitz.

Next, we show that if  $L^p$  norm of  $\nabla u$  grows mildly, then  $u$  is Osgood.

**Lemma 2** ([9]). *Suppose that  $\Theta(p)$  is an admissible growth function and there exists a constant  $C > 0$  such that  $\|\nabla u\|_{L^p(\mathbb{R}^d)} \leq Cp \Theta(p)$  for all  $p \geq 1$ . Then  $u$  has a modulus of continuity  $\varphi_\Theta$ .*

*Proof.* Let  $p > 2d$ , and we apply Morrey's inequality:

$$|u(x) - u(y)| \leq C_d \|\nabla u\|_{L^p(\mathbb{R}^d)} |x - y|^{1 - \frac{d}{p}}, \quad (14)$$

where  $C_d$  depends only on the dimension  $d$ . For  $0 < |x - y| \leq e^{-2}$ , take  $p = 1 - \log |x - y| \geq 3$ . Then one immediately can see that

$$\begin{aligned} |u(x) - u(y)| &\leq C_d \log \left( \frac{e}{|x - y|} \right) \Theta \left( \log \left( \frac{e}{|x - y|} \right) \right) |x - y|^{1 - \frac{d}{\log e/|x - y|}} \\ &\leq C \log \left( \frac{e}{|x - y|} \right) \Theta \left( \log \left( \frac{e}{|x - y|} \right) \right) |x - y| = C \varphi_\Theta(|x - y|), \end{aligned} \quad (15)$$

since  $\lim_{r \rightarrow 0^+} r^{-d/\log(e/r)} = e^d$ .  $\square$

## 2.2 Alberti-Crippa-Mazzucato construction

Our proof crucially relies on the construction made in [2], so we briefly summarize it.

**Proposition 3** ([1]). *Let  $d \geq 2$ ,  $Q = [-\frac{1}{2}, \frac{1}{2}]^d$ . There exist a divergence-free velocity field  $v \in C^\infty(\mathbb{R}_{\geq 0} \times \mathbb{R}^d)$  and a corresponding non-trivial solution  $\rho \in C^\infty(\mathbb{R}_{\geq 0} \times \mathbb{R}^d)$  of the continuity equation (1) such that*

1. *for every  $t \geq 0$ ,  $v(t, \cdot)$  and  $\rho(t, \cdot)$  are bounded and supported in  $Q$ ,*

2. there is a constant  $B > 0$ , independent of  $p$  such that  $\|\nabla v(t, \cdot)\|_{L^p(\mathbb{R}^d)} \leq B$  for all  $1 \leq p \leq \infty$ , and
3. for every  $0 < s < 2$  there exist constants  $C_s, c > 0$  such that for every  $t \geq 0$ ,  $\|\rho(t, \cdot)\|_{\dot{H}^s(\mathbb{R}^d)} \geq C_s e^{sct}$ .

Next, we consider a sequence of cubes  $Q_n$  in  $\mathbb{R}^n$  of side-length  $\lambda_n \searrow 0$  and center  $q_n \in \mathbb{R}^d$  which are disjoint to each other, contained in a compact set, and converging to the origin. Next, we set a sequence  $\tau_n \searrow 0$  and in each cube  $Q_n$  define

$$u_n(t, x) = \frac{\lambda_n}{\tau_n} v\left(\frac{t}{\tau_n}, \frac{x - q_n}{\lambda_n}\right), \theta_n(t, x) = \rho\left(\frac{t}{\tau_n}, \frac{x - q_n}{\lambda_n}\right). \quad (16)$$

We note the following: for detailed computations one may refer to [2].

1.  $\theta_n$  solves  $\partial_t \theta_n + u_n \cdot \nabla_x \theta_n = 0$ .
2. The distance between support of  $\theta_n$  and  $Q_n^c$  is at least  $\lambda_n$ ,
3.  $\|u_n\|_{\dot{W}^{1,p}(\mathbb{R}^d)} \leq \frac{\lambda_n^{\frac{d}{p}}}{\tau_n}$ ,  $\|\theta_n(0)\|_{\dot{H}^\sigma(\mathbb{R}^d)} \leq \lambda_n^{\frac{d}{2}-\sigma}$ , and  $\|\theta_n(t)\|_{\dot{H}^s(\mathbb{R}^d)}^2 \geq \lambda_n^{d-2s} \left(C_s^2 \exp\left(\frac{2sct}{\tau_n}\right) - \frac{C}{s}\right)$  for some constant  $C$ , for every  $0 < s < 1$ .

The key lemma is the following:

**Lemma 3.** *There exists  $\lambda_n \searrow 0$ ,  $\tau_n \searrow 0$  such that*

1.  $\sum_n \lambda_n < \infty$ , and thus  $Q_n$  can be chosen to be disjoint, contained in a compact set and converging to the origin,
2. there exists a constant  $C > 0$  such that  $\sum_n \frac{\lambda_n^{\frac{d}{p}}}{\tau_n} \leq Cp\Theta(p)$  for all  $1 \leq p < \infty$ , so that  $u = \sum_n u_n$  is in  $\dot{W}^{1,p}$  with  $\|u\|_{\dot{W}^{1,p}} \leq Cp\Theta(p)$ ,
3. there exists  $C > 0$  such that  $\frac{\lambda_n}{\tau_n} \leq C$  for every  $n$ , so that  $u(t, \cdot) \in L^\infty(\mathbb{R}^d)$  uniformly in  $t$ ,
4.  $\sum_n \lambda_n^{\frac{d}{2}-\sigma} < \infty$ , so that  $\theta_0 = \sum_n \theta_n(0, \cdot) \in \dot{H}^\sigma(\mathbb{R}^d)$ ,
5.  $\sum_n \lambda_n^{d-2s} \exp\left(\frac{2sct}{\tau_n}\right) = \infty$  for every  $0 < s < 1$  and  $t > 0$ , so that  $\theta(t, \cdot) = \sum_n \theta_n(t, \cdot) \notin \dot{H}^s(\mathbb{R}^d)$ .

Note that  $\theta(t, \cdot) \in L^\infty(\mathbb{R}^d)$  uniformly in  $t$ . Also, since each  $Q_n$  are disjoint to each other, and  $\cup_n Q_n$  is contained in a compact set  $K$ ,  $\sum_{k=1}^n u_k \rightarrow u$ ,  $\sum_{k=1}^n \theta_k \rightarrow \theta$  strongly in  $L^2([0, T] \times K)$  for every  $T > 0$ . Thus, for any  $\psi \in C_0^\infty((0, \infty) \times \mathbb{R}^d)$ ,  $0 = \int \partial_t \psi \sum_{k=1}^n \theta_k + \nabla_x \psi \cdot (\sum_{k=1}^n u_k) \sum_{k=1}^n \theta_k \rightarrow \int \partial_t \psi \theta + \nabla_x \psi \cdot u \theta$ , that is,  $\theta$  is a weak solution of (1) with  $u$ . Moreover, as  $u$  is Osgood by Lemma 1,  $\theta$  is the unique integrable weak solution. Since  $\theta(t, \cdot) \notin \dot{H}^s(\mathbb{R}^d)$  for any  $t > 0$  and any  $0 < s < 1$ , while  $\theta(t, \cdot) \in L^2(\mathbb{R}^2)$  uniformly in time, this proves the Theorem 1.

**Remark 3.** *One can extend Theorem 1 to  $\sigma \geq \frac{d}{2}$  immediately: one may introduce another factor  $\gamma_n \searrow 0$  and set  $\theta = \sum_n \gamma_n \theta_n$ . In fact, along with  $\lambda_n, \tau_n$ , setting  $\gamma_n = \lambda_n^\sigma$  works.*

## 2.3 Proof of Lemma 3

Let  $\Theta$  be an admissible growth function. We set

$$\lambda_n = e^{-e^n}, \tau_n^{-1} = \log(1/\lambda_n)\Theta(\log(1/\lambda_n)) = e^n \Theta(e^n). \quad (17)$$

Obviously, condition 1, 3, and 4 of Lemma 3 are satisfied. For condition 5, we see that

$$\begin{aligned} \sum_n \lambda^{d-2s} \exp\left(\frac{2sct}{\tau_n}\right) &= \sum_n \exp((2s-d)\log(1/\lambda_n) + \log(1/\lambda_n)\Theta(\log(1/\lambda_n))) \\ &\geq C + \sum_{n \geq N} \exp(C \log(1/\lambda_n)\Theta(\log(1/\lambda_n))) = \infty. \end{aligned} \quad (18)$$

It remains to verify condition 2. Since

$$\sum_n \frac{\lambda_n^{\frac{d}{p}}}{\tau_n} = \sum_{n=1}^{\infty} e^{-\frac{d}{p}e^n} e^n \Theta(e^n) = \sum_{n=1}^{\infty} F(e^n), \quad (19)$$

where  $F(x) = x\Theta(x)e^{-\frac{d}{p}x}$ . Note that  $\lim_{x \rightarrow \infty} F(e^x) = 0$ , and  $F$  is bounded; moreover, either there is  $\bar{x}_p \geq 1$  such that  $F(e^x)$  attains maximum at  $\bar{x}_p$ ,  $F(e^x)$  is increasing on  $[1, \bar{x}_p]$ , and decreasing on  $(\bar{x}_p, \infty)$  or  $F$  is decreasing on  $\mathbb{R}_{\geq 1}$ . Therefore, comparing the series with integrals, we have

$$\sum_{n=1}^{\infty} F(e^n) \leq \int_1^{\infty} F(e^x) dx + 2F(e^{\bar{x}_p}), \quad (20)$$

where the last term of (20) (if exists) comes from that for  $n = 1, \dots, [\bar{x}_p] - 1$ , one has  $F(e^n) \leq \int_n^{n+1} F(e^x) dx$ , and for  $n = [\bar{x}_p] + 2, \dots$  one has  $F(e^n) \leq \int_{n-1}^n F(e^x) dx$ , while at  $n = [\bar{x}_p], [\bar{x}_p] + 1$ , one has  $F(e^n) \leq F(e^{\bar{x}_p})$ . We estimate each term of (20).

**Estimate of  $F(e^{\bar{x}_p})$ .** Note that  $F(e^{\bar{x}_p}) = \|F\|_{L^\infty}$ , and it is attained either at the unique point  $y$  such that  $F' = 0$  or  $y = 1$ . Since  $F'(y) = (\Theta(y) + y\Theta'(y) - \frac{d}{p}y\Theta(y))e^{-\frac{d}{p}y}$ , we have

$$\Theta(y) + y\Theta'(y) = \frac{d}{p}y\Theta(y). \quad (21)$$

Since  $\Theta$  is admissible, we have  $(C+1)\Theta(y) \geq \frac{d}{p}y\Theta(y)$ , that is,  $y \leq \frac{(C+1)}{d}p$ . Therefore,

$$\begin{aligned} \|F\|_{L^\infty} = F(y) &\leq \frac{C+1}{d}p\Theta\left(\frac{C+1}{d}p\right) \leq C'p\Theta\left(\left(M + \frac{C+1}{d}\right)p\right) \\ &\leq C'p\left(\Theta(p) + \Theta\left(\left(M + \frac{C+1}{d}\right)p\right) + C\right) \leq C''p\Theta(p) \end{aligned} \quad (22)$$

for large enough  $p$ .

**Estimate of  $\int_1^\infty F(e^x) dx$**  We note that, by change of variable  $z = e^x$ ,

$$\int_1^\infty F(e^x) dx = \int_1^\infty \Theta(e^x) e^{-\frac{d}{p}e^x} e^x dx = \int_e^\infty \Theta(z) e^{-\frac{d}{p}z} dz. \quad (23)$$

By putting  $y = \frac{d}{p}z$ , we have

$$\int_e^\infty \Theta(z) e^{-\frac{d}{p}z} dz = \frac{p}{d} \int_{\frac{de}{p}}^\infty \Theta\left(\frac{p}{d}y\right) e^{-y} dy. \quad (24)$$

Note that for  $p > M$ ,  $\Theta(\frac{p}{d}y) \leq \Theta(pM/d) \leq \Theta(pM) \leq \Theta(p) + \Theta(M) + C$  if  $y \leq M$ , and if  $y > M$   $\Theta(\frac{p}{d}y) \leq \Theta(py) \leq \Theta(p) + \Theta(y) + C$ . In any case, we have a constant  $C' > 0$  satisfying

$$\Theta\left(\frac{p}{d}y\right) \leq \Theta(p) + \Theta(y) + C', y \geq 1. \quad (25)$$

Also, if  $\frac{de}{p} \leq y \leq 1$ , we have  $\Theta\left(\frac{p}{d}y\right) \leq \Theta(p)$ . Thus, we have

$$\begin{aligned} \frac{p}{d} \int_{\frac{de}{p}}^{\infty} \Theta\left(\frac{p}{d}y\right) e^{-y} dy &\leq \frac{p}{d} \int_{\frac{de}{p}}^1 \Theta(p) e^{-y} dy + \frac{p}{d} \int_1^{\infty} (\Theta(p) + \Theta(y) + C') e^{-y} dy \\ &\leq C'' p \Theta(p) \end{aligned} \quad (26)$$

for some constant  $C'' > 0$ , due to the integrability of  $\Theta(y)e^{-y}$ .

### 3 Propagation of regularity

Still stronger regularity is propagated than  $\nabla u \in L^p$  case: in the latter case, only logarithm of derivatives are propagated, provided that the initial data of (1)  $\theta_0$  has one derivative (see, for example, [5].)

#### 3.1 Propagation of modulus for $\theta$

In this subsection, we establish propagation of modulus of continuity result, based on observations made in [10]. We start with a result proved in [10].

**Lemma 4** ([10]). *Suppose that  $\varphi$  is an Osgood modulus of continuity for  $u$ , and define*

$$\mathcal{M}(z) := \int_z^m \frac{dr}{\varphi(r)}, \quad R(z) := \exp(-\mathcal{M}(z)), \quad (27)$$

and

$$[u(t)]_{\varphi} := \sup_{x \neq y} \frac{|u(x, t) - u(y, t)|}{\varphi(|x - y|)}. \quad (28)$$

Then  $R$  is monotonely increasing,  $\lim_{z \rightarrow 0^+} R(z) = 0$ , and

$$|\phi^{-1}(x, t) - \phi^{-1}(y, t)| \leq R^{-1} \left( e^{\int_0^t [u(s)]_{\varphi} ds} R(|x - y|) \right) =: \mu_{[u]_{\varphi}, t}(|x - y|), \quad (29)$$

where  $\phi(x, t)$  is the unique flow generated by  $u$ : i.e. the unique solution of (4) and  $\phi^{-1}(x, t)$  is its back-to-label map.

*Proof.* This is an immediate consequence of Osgood's lemma and time-reversal.  $\square$

An immediate consequence is the following.

**Lemma 5.** *Let  $\mu_0$  be a modulus of continuity, and let  $g$  be a measurable function satisfying*

$$|\theta_0(x) - \theta_0(y)| \leq (g(x) + g(y))\mu_0(|x - y|). \quad (30)$$

Then

$$|\theta(x, t) - \theta(y, t)| \leq (g_t(x) + g_t(y))\mu_0(\mu_{[u]_{\varphi}, t}(|x - y|)) \quad (31)$$

for  $g_t = g \circ \phi_t^{-1}$ .

*Proof.* We may simply recall that  $\phi^{-1}$  is volume preserving and

$$\begin{aligned} |\theta(x, t) - \theta(y, t)| &= |\theta_0(\phi^{-1}(x, t)) - \theta_0(\phi^{-1}(y, t))| \\ &\leq (g(\phi^{-1}(x, t)) + g(\phi^{-1}(y, t)))\mu_0(|\phi^{-1}(x, t) - \phi^{-1}(y, t)|) \\ &\leq (g(\phi^{-1}(x, t)) + g(\phi^{-1}(y, t)))\mu_0(\mu_{[u]_{\varphi}, t}(|x - y|)). \end{aligned} \quad (32)$$

$\square$

This immediately gives the first part of Theorem 2.

**Remark 4.** It is worth noting that the modulus of continuity  $\mu_{0,\varphi} = R$  plays a special role: if  $\theta_0$  has a modulus of continuity  $R$ , then  $\theta(t)$  still has the same modulus of continuity  $R$  since  $\mu_{0,\varphi} \circ \mu_{[u]_{\varphi},t}(r) = e^{\int_0^t [u(s)]_{\varphi} ds} R(r) = e^{\int_0^t [u(s)]_{\varphi} ds} \mu_{0,\varphi}(r)$ .

**Remark 5.** The following are explicit calculation of  $\mu_{[u]_{\varphi},t}(r)$  for some  $\varphi$ .

- If  $u$  is Lipschitz, that is,  $\varphi(z) = z$ , then  $\mathcal{M}(z) = \log(1/z)$  for sufficiently small  $z$ ,  $R(z) = z = R^{-1}(z)$ , and thus  $\mu_{[u]_{\varphi},t}(r) = e^{\int_0^t [u(s)]_{\varphi} ds} r$ . Thus, if  $\theta_0$  is Lipschitz or Holder, then  $\theta(t)$  retains the same modulus of continuity.
- If  $u$  is log-Lipschitz, that is,  $\varphi(z) = z \log(1/z)$ , then  $\mathcal{M}(z) = \log \log(1/z)$  for sufficiently small  $z$ ,  $R(z) = \frac{1}{\log(1/z)}$ ,  $R^{-1}(z) = e^{-1/z}$ , and thus  $\mu_{[u]_{\varphi},t}(r) = r^{\frac{1}{\exp(\int_0^t [u(s)]_{\varphi} ds)}}$ . Thus, if  $\theta_0$  is Lipschitz or Holder, then  $\theta(t)$  remains Holder, with its Holder exponent decreasing over time.
- If  $\varphi(z) = z \log(1/z) \log_2(1/z) \cdots \log_n(1/z)$  for sufficiently small  $z$ , then  $\mathcal{M}(z) = \log_{n+1}(1/z)$ ,  $R(z) = \frac{1}{\log_n(1/z)}$ ,  $R^{-1}(z) = \frac{1}{e_n(1/z)}$  where  $e_n$  is the inverse of  $\log_n$ . Therefore,  $\mu_{[u]_{\varphi},t}(r) = \frac{1}{e_{n-1} \left( (\log_{n-1}(1/r))^{1/\exp(\int_0^t [u(s)]_{\varphi} ds)} \right)}$ .
- If  $\varphi(z) = (1-\alpha)z^\alpha$ ,  $\alpha \in (0,1)$ , then  $\mathcal{M}(z) = 1 - z^{1-\alpha}$ ,  $R(z) = e^{-1}e^{z^{1-\alpha}}$ ,  $R^{-1}(z) = \frac{1}{1-\alpha} \log(ez)$ , and  $\mu_{[u]_{\varphi},t}(r) = \frac{1}{1-\alpha} \left( \int_0^t [u]_{\varphi,t}(s) ds + r^{1-\alpha} \right)$ . In particular,  $\mu_{[u]_{\varphi},t}$  does not vanish as  $r \rightarrow 0$ , and therefore  $\mu_0 \circ \mu_{[u]_{\varphi},t}$  is not a modulus of continuity in general.

Regarding the vanishing rate of  $\mu_{[u]_{\varphi},t}(r)$  for  $\varphi(z) = z \log(1/z) \log_2(1/z) \cdots \log_n(1/z)$ , we have the following result.

**Lemma 6.** Let  $\varphi_n(z) = z \log(1/z) \log_2(1/z) \cdots \log_n(1/z)$  and define

$$\mu_{n,t}(r) := \frac{1}{e_{n-1} \left( (\log_{n-1}(1/r))^{1/\exp(\int_0^t [u(s)]_{\varphi} ds)} \right)}. \quad (33)$$

- $\mu_{n,t}$  modulus of continuity is worse than any Holder modulus of continuity:  $\lim_{r \rightarrow 0^+} \frac{\mu_{n,t}(r)}{r^\alpha} = \infty$  for any  $\alpha \in (0,1]$  for  $n \geq 2$ .
- $\mu_{n,t}$  modulus of continuity is better than any logarithmic modulus:  $\lim_{r \rightarrow 0^+} \frac{\mu_{n,t}(r)}{1/(\log(1/r))^a} = 0$  for any  $a \geq 1$ .

*Proof.* For the first claim, it suffices to show that  $\lim_{r \rightarrow 0^+} \frac{\log_{n-1} 1/r^\alpha}{(\log_{n-1} 1/r)^\beta} = \infty$  for  $\alpha > 0$  sufficiently small and  $\beta \in (0,1)$ . For  $n = 2$ , it obviously holds. For  $n \geq 3$ , as  $\log_{n-2} x$  is an admissible growth function,  $\log_{n-1}(1/r) \leq \log_{n-1}(1/r^\alpha) + \log_{n-2} 1/\alpha + C$  for sufficiently small  $\alpha > 0$  and a constant  $C > 0$ . Thus,  $\frac{\log_{n-1} 1/r^\alpha}{(\log_{n-1} 1/r)^\beta} \geq \frac{\log_{n-1} 1/r^\alpha}{(\log_{n-1}(1/r^\alpha) + C)^\beta} \rightarrow \infty$  as  $r \rightarrow 0^+$ .

For the second claim, it suffices to show that  $\lim_{r \rightarrow 0^+} \frac{\log_{n-1}(\log(1/r))^a}{(\log_{n-1}(1/r))^\beta} = 0$  for any  $\beta \in (0,1)$  and  $a \geq 1$  sufficiently large, and again it is obvious for  $n = 2$ . For  $n \geq 3$ , since  $\log_{n-2}$  is an admissible growth function, we have  $\log_{n-1}(\log(1/r))^a \leq \log_{n-2} a + \log_n(1/r) + C$  for sufficiently large  $a \geq 1$  and a constant  $C > 0$ , and the desired estimate follows.  $\square$



## 3.2 Propagation of Sobolev-type functionals

Finally, we prove the second part of Theorem 2. Our argument is inspired by that of [5]. We use the layer-cake representation, (31), and monotonicity of  $\mu_{[u]_\varphi, t}$  on  $t$ , and volume-preserving property of the flow  $\phi_t^{-1}$  to obtain

$$\begin{aligned}
\int_{\mathbb{R}^d} |\theta(x, t) - \theta(x + h, t)|^2 dx &= 2 \int_0^\infty r |\{x : |\theta(x, t) - \theta(x + h, t)| > r\}| dr \\
&\leq 2 \int_0^\infty r |\{x : g_t(x) + g_t(x + h) > r/\mu_0 \circ \mu_{[u]_\varphi, T}(|h|)\}| dr \\
&= 2(\mu_0 \circ \mu_{[u]_\varphi, T}(|h|))^2 \int_0^\infty r |\{x : g_t(x) + g_t(x + h) > r\}| dr \\
&\leq 2(\mu_0 \circ \mu_{[u]_\varphi, T}(|h|))^2 \int_0^\infty r |\{x : 2g_t(x) > r\}| dr \leq C(\mu_0 \circ \mu_{[u]_\varphi, T}(|h|))^2 \|g\|_{L^2(\mathbb{R}^d)}^2.
\end{aligned} \tag{34}$$

Therefore, we have

$$\begin{aligned}
&\int_{B_1(0)} \int_{\mathbb{R}^d} \frac{|\theta(x, t) - \theta(x + h, t)|^2}{|h|^d} \mu_0 \circ \mu_{[u]_\varphi, t}(|h|)^{-1} dx dh \\
&\leq C \|g\|_{L^2(\mathbb{R}^d)}^2 \int_{B_1(0)} \frac{\mu_0 \circ \mu_{[u]_\varphi, T}(|h|)}{|h|^d} dh \leq C \|g\|_{L^2(\mathbb{R}^d)}^2 \int_0^1 \frac{\mu_0 \circ \mu_{[u]_\varphi, T}(r)}{r} dr = CA(u) \|g\|_{L^2(\mathbb{R}^d)}^2.
\end{aligned} \tag{35}$$

**Remark 6.** In the case of  $\varphi(z) = z \log(1/z) \log_2(1/z) \cdots \log_n(1/z)$ , by Lemma 6,  $\mu_{[u]_\varphi, t} = o(1/\log(1/r)^a)$  for any  $a \geq 1$ : therefore we have

$$\int_{B_1(0)} \int_{\mathbb{R}^d} \frac{|\theta(x, t) - \theta(x + h, t)|^2}{|h|^d} \mu_{[u]_\varphi, t}(|h|)^{-1} dx dh \geq C \int_{B_1(0)} \int_{\mathbb{R}^d} \frac{|\theta(x, t) - \theta(x + h, t)|^2}{|h|^d (\log(1/|h|))^{1-p}} dx dh \tag{36}$$

for some  $C > 0$  and any  $p > 1$ . In this sense, transport equation with Osgood velocity field  $u$  with modulus of continuity  $\varphi$  propagates more regularity than logarithm of a derivative.

## 4 Stability result for 2D Euler equation

### 4.1 An interpolation lemma

We start with an interpolation lemma, which is analogous to Proposition 3.5 of [5].

**Lemma 7.** Let  $\varepsilon \in (0, 1)$ , and let  $\mu : (0, 1) \rightarrow \mathbb{R}_{>0}$  be a continuous increasing function. Then there exists a constant  $C > 0$  such that the following holds.

$$\begin{aligned}
C^{-1} \|f\|_{L^2(\mathbb{R}^2)}^2 &\leq \mu(\varepsilon) \int_{B_1(0)} \int_{\mathbb{R}^d} \frac{|f(x + h) - f(x)|^2}{|h|^d} \mu(|h|)^{-1} dx dh \\
&\quad + |\log \varepsilon| \frac{\|f\|_{L^2(\mathbb{R}^2)}^2}{\log \left( 2 + \frac{\|f\|_{L^2(\mathbb{R}^d)}^2}{\|f\|_{H^{-1}(\mathbb{R}^d)}^2} \right)}.
\end{aligned} \tag{37}$$

*Proof.* We closely follow the proof of Proposition 3.5 of [5]. We first fix  $\delta \in (0, 1)$ ,  $\psi \in C_c^\infty(\mathbb{R}^d)$ , a radial function, with  $\psi \equiv 1$  in  $B_{2/3} \setminus B_{1/2}$ ,  $\psi \equiv 0$  in  $(B_{5/6} \setminus B_{1/3})^c$ , and  $\int \psi_{\mathbb{R}^d} = 1$ , and write  $\psi_\delta(x) = \delta^{-d} \psi(x/\delta)$ . We decompose  $f$  by

$$f = f * \psi_\delta + (f - f * \psi_\delta) \tag{38}$$

and treat each term separately. We have

$$\begin{aligned}
\|f - f * \psi_\delta\|_{L^2}^2 &= \int_{\mathbb{R}^d} \left| \int_{\mathbb{R}^d} (f(x+h) - f(x)) \psi_\delta(|h|) dh \right|^2 dx \\
&\leq \int_{\mathbb{R}^d} \int_{\delta/3 \leq |h| \leq \delta} \frac{|f(x+h) - f(x)|^2}{\delta^d} dh \|\psi\|_{L^2}^2 dx \\
&\leq C \int_{\delta/3 \leq |h| \leq \delta} \int_{\mathbb{R}^d} \frac{|f(x+h) - f(x)|^2}{|h|^d} dx dh, \\
\|f * \psi_\delta\|_{L^2}^2 &= \|\hat{\psi}_\delta \hat{f}\|_{L^2}^2 \leq \sup_\xi |\log(2 + |\xi|)| |\hat{\psi}(\delta\xi)|^2 \int_{\mathbb{R}^d} \frac{|\hat{f}(\xi)|^2}{\log(2 + |\xi|)} d\xi \\
&\leq \max(\sup_{|\xi| \leq 2} 2 \log 2 |\hat{\psi}(\delta\xi)|^2, \sup_{|\xi| > 2} 2 \log |\xi| |\hat{\psi}(\delta\xi)|^2) \int_{\mathbb{R}^d} \frac{|\hat{f}(\xi)|^2}{\log(2 + |\xi|)} d\xi \\
&\leq C \log(\max(1/2, 1/\delta)) \int_{\mathbb{R}^d} \frac{|\hat{f}(\xi)|^2}{\log(2 + |\xi|)} d\xi.
\end{aligned} \tag{39}$$

For the treatment of  $\int_{\mathbb{R}^d} \frac{|\hat{f}(\xi)|^2}{\log(2 + |\xi|)} d\xi$ , we split the integral by low and high frequencies and optimize:

$$\begin{aligned}
\int_{\mathbb{R}^d} \frac{|\hat{f}(\xi)|^2}{\log(2 + |\xi|)} d\xi &= \int_{|\xi| \leq \nu} \frac{|\xi|^2}{\log(2 + |\xi|)} |\xi|^{-2} |\hat{f}(\xi)|^2 d\xi + \int_{|\xi| > \nu} \frac{|\hat{f}(\xi)|^2}{\log(2 + |\xi|)} d\xi \\
&\leq \frac{\nu^2}{\log(2 + \nu)} \int_{|\xi| \leq \nu} |\xi|^{-2} |\hat{f}(\xi)|^2 d\xi + \frac{1}{\log(2 + \nu)} \int_{|\xi| \geq \nu} |\hat{f}(\xi)|^2 d\xi \\
&\leq C \frac{\nu^2}{\log(2 + \nu^2)} \|f\|_{\dot{H}^{-1}}^2 + \frac{C}{\log(2 + \nu^2)} \|f\|_{L^2}^2,
\end{aligned} \tag{40}$$

and taking  $\nu = \|f\|_{L^2} / \|f\|_{\dot{H}^{-1}}$  gives

$$\int_{\mathbb{R}^d} \frac{|\hat{f}(\xi)|^2}{\log(2 + |\xi|)} d\xi \leq \frac{C \|f\|_{L^2}^2}{\log\left(2 + \frac{\|f\|_{L^2(\mathbb{R}^d)}^2}{\|f\|_{\dot{H}^{-1}(\mathbb{R}^d)}^2}\right)}. \tag{41}$$

Thus, we have

$$\begin{aligned}
\|f\|_{L^2}^2 &\leq C \int_{\delta/3 \leq |h| \leq \delta} \int_{\mathbb{R}^d} \frac{|f(x+h) - f(x)|^2}{|h|^d} dx dh \\
&\quad + C \log(\max(1/2, 1/\delta)) \frac{\|f\|_{L^2}^2}{\log\left(2 + \frac{\|f\|_{L^2(\mathbb{R}^d)}^2}{\|f\|_{\dot{H}^{-1}(\mathbb{R}^d)}^2}\right)}.
\end{aligned} \tag{42}$$

Integrating  $\delta$  over  $(\varepsilon, 1)$  with weight  $\frac{\mu(\delta)^{-1}}{\delta}$  gives us the following:

$$\begin{aligned}
\|f\|_{L^2}^2 &\leq C \frac{1}{\int_\varepsilon^1 \frac{\mu(\delta)^{-1}}{\delta} d\delta} \int_\varepsilon^1 \int_{\delta/3 \leq |h| \leq \delta} \int_{\mathbb{R}^d} \frac{|f(x+h) - f(x)|^2}{|h|^d} dx dh \frac{\mu(\delta)^{-1}}{\delta} d\delta \\
&\quad + C \frac{1}{\int_\varepsilon^1 \frac{\mu(\delta)^{-1}}{\delta} d\delta} \int_\varepsilon^1 \log(\max(1/2, 1/\delta)) \frac{\|f\|_{L^2}^2}{\log\left(2 + \frac{\|f\|_{L^2(\mathbb{R}^d)}^2}{\|f\|_{\dot{H}^{-1}(\mathbb{R}^d)}^2}\right)} \frac{\mu(\delta)^{-1}}{\delta} d\delta \\
&\leq C \frac{\int_\varepsilon^1 \frac{d\delta}{\delta}}{\int_\varepsilon^1 \frac{\mu(\delta)^{-1}}{\delta} d\delta} \int_{B_1(0)} \int_{\mathbb{R}^d} \frac{|f(x+h) - f(x)|^2}{|h|^d} dx dh + C |\log \varepsilon| \frac{\|f\|_{L^2}^2}{\log\left(2 + \frac{\|f\|_{L^2(\mathbb{R}^d)}^2}{\|f\|_{\dot{H}^{-1}(\mathbb{R}^d)}^2}\right)}.
\end{aligned} \tag{43}$$

Since  $\mu$  is increasing, we have  $\frac{\int_{\varepsilon}^1 \frac{d\delta}{\delta}}{\int_{\varepsilon}^1 \frac{\mu(\delta)^{-1}}{\delta} d\delta} \leq \mu(\varepsilon)$ , and (37) follows.  $\square$

## 4.2 Proof of Theorem 3

From Lemma 1 and Lemma 2, we see that  $u_1, u_2$  have modulus of continuity  $\varphi_{\Theta_n}$  as in (12), with

$$[u_i]_{\varphi_{\Theta_n}} \leq M \|\omega_{i,0}\|_{Y_{\Theta_n}}, i = 1, 2. \quad (44)$$

Therefore, there is  $C_1, C_2 > 0$  such that

$$\mu[u_i]_{\varphi_{\Theta_n}, t}(r) \leq \mu_{\omega, n, t}(r), i = 1, 2. \quad (45)$$

Next, we have the following stability estimate of velocity for Euler equation:

**Proposition 4** ([12]). *For some constant  $C_1, C_2 > 0$ , we have*

$$\|u_1(t) - u_2(t)\|_{L^2}^2 \leq \mu_{\omega, n, t}(\|u_{0,1} - u_{0,2}\|_{L^2}^2), \quad (46)$$

One can adjust  $C_1, C_2 > 0$  appropriately (if necessary) so that (45) and (46) are simultaneously satisfied. Next, by the standard Lusin estimate, we have

$$|\omega_{0,i}(x) - \omega_{0,i}(y)| \leq C|x - y|^s (D_s \omega_{0,i}(x) + D_s \omega_{0,i}(y)), i = 1, 2, \quad (47)$$

where

$$D_s f(x) := \left( \int_{\mathbb{R}^d} \frac{|f(x+h) - f(x)|^2}{|h|^{d+2s}} dh \right)^{\frac{1}{2}} \in L^2. \quad (48)$$

Therefore, by the same argument as in Theorem 2, we have

$$\int_{B_1(0)} \int_{\mathbb{R}^d} \frac{|\omega_i(x, t) - \omega_i(x+h, t)|^2}{|h|^d} (\mu_{\omega, n, t}(|h|))^{-s} dx dh \leq C \|D_s \omega_i\|_{L^2}^2 \int_0^1 \frac{(\mu_{\omega, n, t}(r))^s}{r} dr < \infty, \quad (49)$$

since  $\mu_{\omega, n, t}$  vanishes faster than any logarithm by Lemma 6. Therefore, applying (37) with  $\mu(r) = (\mu_{\omega, n, t}(r))^s$  and  $f = \omega_1 - \omega_2$  gives that for sufficiently small  $\varepsilon > 0$ , we have

$$\begin{aligned} C^{-1} \|\omega_1(t) - \omega_2(t)\|_{L^2}^2 &\leq (\mu_{\omega, n, t}(\varepsilon))^s + |\log \varepsilon| \frac{\|\omega_1(t) - \omega_2(t)\|_{L^2}^2}{\log(2 + \|\omega_1(t) - \omega_2(t)\|_{L^2}^2 / \|u_1(t) - u_2(t)\|_{L^2}^2)} \\ &\leq (\mu_{\omega, n, t}(\varepsilon))^s + |\log \varepsilon| \frac{\|\omega_1(t) - \omega_2(t)\|_{L^2}^2}{\log\left(2 + \frac{\|\omega_1(t) - \omega_2(t)\|_{L^2}^2}{\mu_{\omega, n, t}(\|u_{0,1} - u_{0,2}\|_{L^2}^2)}\right)}. \end{aligned} \quad (50)$$

Now take

$$\varepsilon = \left( 2 + \frac{\|\omega_1(t) - \omega_2(t)\|_{L^2}^2}{\mu_{\omega, n, t}(\|u_{0,1} - u_{0,2}\|_{L^2}^2)} \right)^{\gamma} \quad (51)$$

for some  $\gamma < 0$  so that

$$\frac{|\log \varepsilon|}{\log\left(2 + \frac{\|\omega_1(t) - \omega_2(t)\|_{L^2}^2}{\mu_{\omega, n, t}(\|u_{0,1} - u_{0,2}\|_{L^2}^2)}\right)} < C^{-1}/2,$$

and then we have

$$\|\omega_1(t) - \omega_2(t)\|_{L^2}^2 \leq C \mu_{\omega, n, t} \left( \left( 2 + \frac{\|\omega_1(t) - \omega_2(t)\|_{L^2}^2}{\mu_{\omega, n, t}(\|u_{0,1} - u_{0,2}\|_{L^2}^2)} \right)^{\gamma} \right). \quad (52)$$

Replacing  $\|\omega_1(t) - \omega_2(t)\|_{L^2}^2$  on the right-hand side by  $2 \max_i \|\omega_{0,i}\|_{L^2}$  gives (9).

## References

- [1] G. Alberti, G. Crippa, and A. L. Mazzucato. Exponential self-similar mixing by incompressible flows. *J. Amer. Math. Soc.*, 32(2):445–490, 2019.
- [2] G. Alberti, G. Crippa, and A. L. Mazzucato. Loss of regularity for the continuity equation with non-Lipschitz velocity field. *Ann. PDE*, 5(1):Paper No. 9, 19, 2019.
- [3] L. Ambrosio and P. Bernard. Uniqueness of signed measures solving the continuity equation for Osgood vector fields. *Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl.*, 19(3):237–245, 2008.
- [4] H. Bahouri, J.-Y. Chemin, and R. Danchin. *Fourier analysis and nonlinear partial differential equations*, volume 343 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer, Heidelberg, 2011.
- [5] E. Brué and Q.-H. Nguyen. Sharp regularity estimates for solutions of the continuity equation drifted by Sobolev vector fields. *Anal. PDE*, 14(8):2539–2559, 2021.
- [6] L. Caravenna and G. Crippa. A directional Lipschitz extension lemma, with applications to uniqueness and Lagrangianity for the continuity equation. *Comm. Partial Differential Equations*, 46(8):1488–1520, 2021.
- [7] P. Constantin, T. D. Drivas, and T. M. Elgindi. Inviscid limit of vorticity distributions in the Yudovich class. *Comm. Pure Appl. Math.*, 75(1):60–82, 2022.
- [8] G. Crippa and C. De Lellis. Estimates and regularity results for the DiPerna-Lions flow. *J. Reine Angew. Math.*, 616:15–46, 2008.
- [9] G. Crippa and G. Stefani. An elementary proof of existence and uniqueness for the euler flow in localized yudovich space. *arxiv preprint*.
- [10] T. D. Drivas, T. Elgindi, and J. La. Propagation of singularities by osgood vector fields and for 2d inviscid incompressible fluids. *arxiv preprint*.
- [11] P. Hartman. *Ordinary differential equations*, volume 38 of *Classics in Applied Mathematics*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2002. Corrected reprint of the second (1982) edition [Birkhäuser, Boston, MA; MR0658490 (83e:34002)], With a foreword by Peter Bates.
- [12] C. Kim and J. M. La. Vorticity convergence from boltzmann to 2d incompressible euler equations below yudovich class. *arxiv preprint*.
- [13] D. Meyer and C. Seis. Propagation of regularity for transport equations. a littlewood-paley approach. *arxiv preprint*.