

MITTAG-LEFFLER MODULES AND DEFINABLE SUBCATEGORIES. II

PHILIPP ROTHMALER

This note on countably generated relative Mittag-Leffler modules is a continuation of [R3], henceforth referred to as Part I. Its terminology, preliminaries, and results are freely used throughout. (In particular, modules are left R -modules unless stated otherwise.) All unspecified citations are to that paper. Theorem 7.1(1) states:

The countably generated \mathcal{K} -Mittag-Leffler modules are precisely the uniform \mathcal{L} -pure images of \mathcal{L} - ω -limits.

Recall from Def. 7.4, an \mathcal{L} - ω -limit is an ω -limit of finitely presented modules that is \mathcal{L} -pure on images. Here, and throughout, \mathcal{K} and \mathcal{L} are definably dual classes of right, resp., left R -modules in the sense of Def. 2.5, i.e., they generate mutually elementarily dual definable subcategories, cf. Conv. 2.7. By the main theorem of [R1] (or Part I), the \mathcal{K} -Mittag-Leffler modules are precisely the \mathcal{L} -atomic modules. One could therefore replace \mathcal{K} -Mittag-Leffler by \mathcal{L} -atomic everywhere, as was done in [P3].

In Theorem 7.1(2) ‘uniform \mathcal{L} -pure images’ were incorrectly omitted. The resulting discrepancy in the classical case was detected and communicated to me by Jan Trlifaj, for which I am very grateful.

After taking the opportunity to correct the statement of Thm. 7.1(2), I continue the study of uniform purity of epimorphisms in order to derive the main result, Thm. 3.1(2), which states that—provided $R_R \in \langle \mathcal{K} \rangle$ (equivalently, $R_{\#} \subseteq \langle \mathcal{L} \rangle$), the definable subcategory generated by \mathcal{L} —every countably generated \mathcal{K} -Mittag-Leffler module in $\langle \mathcal{L} \rangle$ is a direct summand of a $\langle \mathcal{L} \rangle$ -preenvelope of a union of an \mathcal{L} -pure ω -chain of finitely presented modules. In conclusion I present a number of examples that starts with and grew out of the study of \mathcal{L} -purity (of monomorphisms in $\mathbb{Z}\text{-Mod}$) for $\mathcal{L} = \text{Div}$, the definable subcategory of divisible abelian groups.

1. THE CORRECTED THEOREM 7.1(2)

If $R_R \in \langle \mathcal{K} \rangle$, the countably generated \mathcal{K} -Mittag-Leffler modules are precisely the uniform \mathcal{L} -pure images of unions of \mathcal{L} -pure ω -chains of finitely presented modules.

Deleting the incorrect application of Lemma 5.11—which simply does not apply to ‘right pure’ maps of §5.4 (but only to ‘left pure’ maps of §§5.1+2)—the proof given for Thm. 7.1(2) (in Part I) yields just that.

Date: June 30, 2022.

2. UNIFORM PURITY

This entire section goes back to [R1, Lemma 3.9], where it was proved model-theoretically that countably generated relative Mittag-Leffler modules are relatively pure projective. Here I reiterate this material in terms introduced in Part I, especially some splitting behavior of relative Mittag-Leffler modules—the terms were coined recently, the proofs are old.

In [R2, Prop. 2.5] it was proved that every pure epimorphism from a module in \mathcal{L} onto a countably generated \mathcal{L} -atomic module splits. As pointed out there, this result is just a variant of [R1, Lemma 3.9]. Here I present another variant. To make the arguments more transparent, I distinguish three kinds of right \mathcal{L} -purity, i.e., purity for epimorphisms, and I break up the proof accordingly.

- Definition 2.1.** (1) [R4, Def.1.1(2)]. The map $g : B \rightarrow C$ is an \mathcal{L} -pure epimorphism if for every tuple \bar{c} in C and every pp formula φ it satisfies, there is a g -preimage \bar{b} in B which satisfies a pp formula $\psi \leq_{\mathcal{L}} \varphi$.
- (2) [R3, Def. 5.12]. The map g is a *uniform \mathcal{L} -pure epimorphism* if every tuple in $\bar{c} \in C$ has a g -preimage \bar{b} such that $\text{pp}_B(\bar{b}) \sim_{\mathcal{L}} \text{pp}_C(\bar{c})$.
- (3) The map g is a *strict uniform \mathcal{L} -pure epimorphism* if for all tuples $\bar{b} \in B$ and $\bar{c} \in C$ with $g(\bar{b}) = \bar{c}$ and $\text{pp}_B(\bar{b}) \sim_{\mathcal{L}} \text{pp}_C(\bar{c})$, every element $c \in C$ has a g -preimage b such that $\text{pp}_B(\bar{b}, b) \sim_{\mathcal{L}} \text{pp}_C(\bar{c}, c)$.
- (4) The prefix \mathcal{L} is omitted when it is all of $R\text{-Mod}$.
- (5) Note that always $\text{pp}_B(\bar{b}) \subseteq \text{pp}_C(\bar{c})$ and that the properties above are an invariant of the definable subcategory generated by \mathcal{L} .
- (6) In particular, w.l.o.g., $\mathcal{L} = \langle \mathcal{L} \rangle$ everywhere in these concepts.

Upon iterating the strict condition one easily sees that *strict uniform \mathcal{L} -pure epimorphisms* are *uniform \mathcal{L} -pure epimorphisms*. In turn, by [R3, Rem. 2.1], *uniform \mathcal{L} -pure epimorphisms* are \mathcal{L} -pure epimorphisms.

- Remark 2.2.** (1) If $B \in \mathcal{L}$ in (1), then \bar{b} in B satisfies the formula φ itself. Hence every \mathcal{L} -pure epimorphism $g : B \rightarrow C$ is a pure epimorphism (i.e., $R\text{-Mod}$ -pure epic).
- (2) If $B \in \mathcal{L}$ in (2), then \bar{b} in B realizes the entire type $\text{pp}_C(\bar{c})$, hence $\text{pp}_B(\bar{b}) = \text{pp}_C(\bar{c})$. Thus every uniform \mathcal{L} -pure epimorphism $g : B \rightarrow C$ is a uniform pure epimorphism (i.e., uniform $R\text{-Mod}$ -pure epic).
- (3) Similarly, if $B \in \mathcal{L}$, then every strict uniform \mathcal{L} -pure epimorphism $g : B \rightarrow C$ is a strict uniform pure epimorphism (i.e., strictly uniform $R\text{-Mod}$ -pure epic).

Arguments like the following are standard in model theory. This particular one is a major ingredient in the proof of aforementioned [R1, Lemma 3.9]. Note, $\mathcal{L} = R\text{-Mod}$ here.

Lemma 2.3. *Strict uniform epimorphisms onto countably generated modules split.*

Proof. Suppose $C = \langle c_0, c_1, c_2, \dots \rangle$ and let $g : B \rightarrow C$ be a strict uniform epimorphism. Starting from the empty tuple, choose a preimage $b_0 \in B$ of c_0 of same pp type, and continue and successively choose preimages $b_1, b_2, \dots, b_i, \dots$ of $c_1, c_2, \dots, c_i, \dots$, respectively, such that $\text{pp}_B(b_0, b_1, \dots, b_i) = \text{pp}_C(c_0, c_1, \dots, c_i)$ for all i . Then the assignment $h(c_i) = b_i$ defines a map, since if $\sum_{i < n} c_i = 0$ in C then $\sum_{i < n} x_i \doteq 0 \in \text{pp}_C(c_1, c_2, \dots, c_i) = \text{pp}_B(b_1, b_2, \dots, b_i)$, hence $\sum_{i < n} b_i = 0$ in B . Applying the same argument to formulas expressing linear dependence shows, h is a homomorphism. Consequently, h is a section of g , this proving that g splits. \square

Lemma 2.4. (\mathcal{L} -) pure epimorphisms from \mathcal{L} onto \mathcal{L} -atomic modules are uniform.

Proof. Suppose $B \in \mathcal{L}$ and $g : B \rightarrow C$ is a pure epimorphism onto the \mathcal{L} -atomic module C . To prove g is uniform, let $\bar{c} \in C$ and choose an \mathcal{L} -generator $\varphi \in q$. By purity, \bar{c} has a g -preimage \bar{b} satisfying φ in B . As $B \in \mathcal{L}$, its pp-type, $p := \text{pp}_B(\bar{b})$, contains all of q . The reverse inclusion follows from $g(\bar{b}) = \bar{c}$. Thus $p = q$, which proves, g is uniform. \square

Proposition 2.5. Suppose, $B \in \mathcal{L}$ and C is \mathcal{L} -atomic.

Then every \mathcal{L} -pure epimorphism $B \rightarrow C$ is a strict uniform epimorphism (i.e., strictly uniform R -Mod-pure epic) and $C \in \langle \mathcal{L} \rangle$.

Proof. Let $g : B \rightarrow C$ be \mathcal{L} -pure epic. By the first part of the hypothesis and Rem. 2.2(1), g is a pure epimorphism. Together with the second part of the hypothesis and Lemma 2.4, this implies, g is a uniform pure epimorphism. To prove it is strictly so, let $\bar{b} \in B$ and $\bar{c} \in C$ with $g(\bar{b}) = \bar{c}$ and $\text{pp}_B(\bar{b}) = \text{pp}_C(\bar{c})$ and pick an element $c \in C$. We have to find a g -preimage b of c with $\text{pp}_B(\bar{b}, b) = \text{pp}_C(\bar{c}, c)$.

Set $p := \text{pp}_B(\bar{b})$, $q = \text{pp}_C(\bar{c})$, $q' := \text{pp}_C(\bar{c}, c)$, and choose \mathcal{L} -generators $\varphi = \varphi(\bar{x}) \in q$ and $\varphi' = \varphi'(\bar{x}, x) \in q'$.

(One could carry out the argument below entirely on the \mathcal{L} level to prove $\text{pp}_B(\bar{b}, b) \sim_{\mathcal{L}} \text{pp}_C(\bar{c}, c)$ and then apply Rem. 2.2(3) to get equality. However, there is no gain, and we argue simply on the R -Mod level.)

Note, $\exists x \varphi' \in q$, hence $\varphi \leq_{\mathcal{L}} \exists x \varphi'$. (Even $\varphi \sim_{\mathcal{L}} \exists x \varphi'$, for, as $q \subseteq q'$, we have $\varphi' = \varphi'(\bar{x}, x) \leq_{\mathcal{L}} \varphi(\bar{x})$, hence also $\exists x \varphi' \leq_{\mathcal{L}} \varphi$.) Accordingly, choose $b' \in B$ with $B \models \varphi'(\bar{b}, b')$. Then $C \models \varphi'(\bar{c}, g(b'))$, hence $C \models \varphi'(\bar{0}, g(b') - c)$.

Set $c'' := g(b') - c$. By uniformity, c'' has a g -preimage $b'' \in B$ of same pp type: $\text{pp}_B(b'') = \text{pp}_C(c'')$. This implies $B \models \varphi'(\bar{0}, b'')$, which, together with $B \models \varphi'(\bar{b}, b')$, yields $B \models \varphi'(\bar{b}, b' - b'')$.

The final claim is that $b := b' - b''$ is the sought-after element in B . Clearly, $g(b) = g(b' - b'') = g(b') - c'' = c$. To see that $p' := \text{pp}_B(\bar{b}, b) = q'$, note that $p' \subseteq q'$ and that both types contain the latter's \mathcal{L} -generator. Invoking $B \in \mathcal{L}$ once again, this entails $q' \subseteq p'$, which shows, g is strict uniform.

Finally, as definable subcategories are preserved under pure epic images, this implies $C \in \langle \mathcal{L} \rangle$. \square

Invoking Lemma 2.3, this yields the following splitting result.

Corollary 2.6. Suppose, C is a countably generated \mathcal{L} -atomic module.

If C is an \mathcal{L} -pure epimorphic image of a module $B \in \mathcal{L}$, then C is a direct summand of B (and a member of $\langle \mathcal{L} \rangle$). \square

Beware, in the \mathcal{L} -pure chains considered below, \mathcal{L} -purity is a left purity, i.e., a purity for monomorphisms (as opposed to epimorphisms, as considered previously). Here, as usual, $N \subseteq M$ is an \mathcal{L} -pure embedding if for every tuple \bar{n} in N , the pp types of \bar{n} in N and of \bar{n} in M are \mathcal{L} -equivalent.

Remark 2.7. Over any ring, if $M \subseteq N$, then M is \mathcal{L} -pure in N iff the following holds for every tuple \bar{a} in M : if \bar{a} satisfies a pp formula φ in N , then it also satisfies some pp formula $\psi \leq_{\mathcal{L}} \varphi$ in M , see (the proof of) [R3, Lemma 5.3].

Remark 2.8. (1) By Theorem 7.1(1) (of Part I), every countably generated \mathcal{L} -atomic module is a uniform \mathcal{L} -pure image of an \mathcal{L} - ω -limit. If \mathcal{L} is large enough

as to contain every such \mathcal{L} - ω -limit, then, by the corollary, every countably generated \mathcal{L} -atomic module is a direct summand of an \mathcal{L} - ω -limit.

- (2) Similarly, Theorem 7.1(2), see above, can be improved on as follows in case \mathcal{L} contains all unions of \mathcal{L} -pure chains of finitely presented modules and all absolutely pure modules (the latter is equivalent to $R_R \in \langle \mathcal{K} \rangle$).

Then the countably generated \mathcal{K} -Mittag-Leffler modules are precisely the direct summands of unions of \mathcal{L} -pure chains of finitely presented modules.

- (3) This is the case if $\mathcal{L} = R\text{-Mod}$. As then the unions in question are direct sums of finitely presented modules, we get back the classical result from [RG, p. 74, 2.2.2] that countably generated Mittag-Leffler-modules are pure-projective.

This suggests the question posed in Section 4. But before asking more questions I combine these splitting facts with what is known about preenvelopes in definable subcategories.

3. ENTER PREENVELOPES

Next we prove that every countably generated \mathcal{K} -Mittag-Leffler module N in \mathcal{L} is a direct summand of a $\langle \mathcal{L} \rangle$ -preenvelope of some \mathcal{L} - ω -limit. We know from Theorem 7.1(1) that there is an \mathcal{L} - ω -limit M and a uniform \mathcal{L} -pure epimorphism $h : M \twoheadrightarrow N$. Let $\mathcal{D} = \langle \mathcal{L} \rangle$ be the definable subcategory generated by \mathcal{L} . By [RS, Corollary 3.5(c)], every module M has a \mathcal{D} -preenvelope, i.e., a map $\varepsilon_M : M \rightarrow D_M$, with $D_M \in \mathcal{D}$, through which every other map from M to a member of \mathcal{D} factors. Applying this to the \mathcal{L} - ω -limit M , we see that h factors through ε_M , which gives us $h_D : D_M \twoheadrightarrow N$ such that $h = h_D \varepsilon_M$. The same simple argument that shows h_D is surjective also proves that it is, as a matter of fact, a uniform \mathcal{L} -pure epimorphism (remember, morphisms preserve pp formulas (and types)!). Thus Cor.2.6 implies that h_D splits, this making N a direct summand of D_M , which proves (1) below. (Note for the converse that definable subcategories are closed under direct summands).

Theorem 3.1. *Let $\mathcal{D} = \langle \mathcal{L} \rangle$, the definable subcategory generated by \mathcal{L} .*

Suppose N is a countably generated \mathcal{K} -Mittag-Leffler (= \mathcal{L} -atomic) module.

There is a \mathcal{D} -preenvelope D of an \mathcal{L} - ω -limit, M , and a uniform \mathcal{L} -pure epimorphism $h_D : D \twoheadrightarrow N$. For any such D and h_D the following holds.

- (1) *$N \in \mathcal{D}$ if and only if (h_D splits and) N is a direct summand of D .*
- (2) *If $R_R \in \langle \mathcal{K} \rangle$ (equivalently, $R_{\#} \subseteq \mathcal{D}$), then every countably generated \mathcal{K} -Mittag-Leffler module in \mathcal{D} is a direct summand of a \mathcal{D} -preenvelope of a union of an \mathcal{L} -pure ω -chain of finitely presented modules.*
- (3) *If \mathcal{D} contains $R_{\#}$ and an \mathcal{L} -pure ω -chain of finitely presented modules whose union is M , then $M \in \mathcal{D}$ is its own \mathcal{D} -preenvelope and, if N is in \mathcal{D} as well, N is a direct summand of M .*
- (4) *If $\mathcal{D} = R\text{-Mod}$, then M is a direct sum of finitely presented modules, hence N is pure-projective. (This is, once again, the aforementioned classical result from [RG].)*

Proof. (1) was proved above. For (2), just put (1) together with Theorem 7.1(2) above. For (3), as definable subcategories are closed under limits, $M \in \mathcal{D}$, and it remains to apply (1). Finally, (4) is a special case of (3). \square

Remark 3.2. There is another interesting module in \mathcal{D} , namely, the direct limit D_{∞} of \mathcal{D} -preenvelopes of the corresponding finitely presented modules constituting

the \mathcal{L} -chain of the \mathcal{L} - ω -limit M . (Note, by [P3], these preenvelopes can be taken strict \mathcal{D} -atomic (= \mathcal{K} -Mittag-Leffler), which does not (seem to) make D_∞ , however, \mathcal{D} -atomic, at least not automatically.) By properties of limits, there is an epimorphism $h_\infty : M \rightarrow D$. As $D_\infty \in \mathcal{D}$, h_∞ factors through ε_M . But it's not clear (to me) if $h : M \twoheadrightarrow N$ factors through D_∞ as well.

4. WHEN ARE \mathcal{L} - ω -LIMITS IN \mathcal{L} ?

I have no conclusive answer. All I have is a simple criterion for any direct limit to be in $\langle \mathcal{L} \rangle$. We need it only for \mathcal{L} - ω -limits, so, for simplicity, it is formulated here only for direct systems of partial order type ω , i.e., for chains.

Remark 4.1. (1) Let for simplicity (and w.l.o.g.) $\mathcal{L} = \langle \mathcal{L} \rangle$, axiomatized by a collection, Ψ , of pp implications—written as pp pairs φ/ψ , i.e., it is assumed that $\psi \leq \varphi$, while the inverse implication, $\varphi \leq \psi$, (stating the closing of the pair) is an axiom of \mathcal{L} . Cf. [P2].

Consider an ω -system $A_0 \xrightarrow{f_0} A_1 \xrightarrow{f_1} A_2 \xrightarrow{f_2} \dots$. Given $i \leq j$, let $f_i^j : A_i \rightarrow A_j$ be the corresponding composition of maps. Suppose A is the limit of this chain.

Then $A \in \mathcal{L}$ if and only if the following holds for every $\varphi/\psi \in \Psi$ and $i < \omega$.

If $a_i \in \varphi(A_i)$, then $f_i^j(a_i) \in \psi(A_j)$, for some $j \geq i$.

(2) Mutatis mutandis, the same holds for directed systems of any shape.

When constructing the \mathcal{L} - ω -limit M_Φ (in [R3, Not.7.9]) for a given \mathcal{L} -atomic module N , what condition on the ω -system associated with M_Φ ensures its membership in $\langle \mathcal{L} \rangle$?

5. WHEN ALL EMBEDDINGS ARE \mathcal{L} -PURE

Example 5.1 (Divisible abelian groups, i.e., $R = \mathbb{Z}$). Let $\mathcal{L} = \text{Div}$, the class of divisible abelian groups, which constitutes the subcategory defined by the closing of the pp pairs of the form $(x \doteq x)/(r|x)$, where r runs over all nonzero integers.

- (1) Every embedding of abelian groups is Div-pure: \mathbb{Z} being an RD ring, every pp formula φ is equivalent to a finite conjunction of basic RD-formulas, $r|\mathfrak{b}\bar{x}$, where \mathfrak{b} is a row vector of integers. If one of those conjuncts has $r = 0$, then that conjunct is quantifier-free and is thus true of \bar{n} in M iff it is true of it in N . So, w.l.o.g., all $r \neq 0$ in φ . In that case, those conjuncts define everything in any divisible abelian group. Consequently, $\varphi(D) = D^n$ (where n is the number of free variables in φ) in every $D \in \text{Div}$. This means, $\bar{x} \doteq \bar{x} \leq_{\text{Div}} \varphi$, and therefore the formula ψ in Rem. 2.7 can be taken to be $\bar{x} \doteq \bar{x}$, which is always true—this showing that every embedding is Div-pure, as desired.
- (2) Thus every chain of finitely generated (= finitely presented) abelian groups is a Div-chain, hence every union of such is a Div- ω -limit, and hence every countable (= countably generated) abelian group is a Div- ω -limit.
- (3) Consequently, every countable abelian group is Div-atomic (and itself a Div- ω -limit—no uniform epis needed).
- (4) The Prüfer groups are examples of members of Div that are Div- ω -limits of finite groups (which are obviously not in Div).

- (5) Note, \mathcal{K} (the elementary dual of $\mathcal{L} = \text{Div}$) is the class of torsionfree (= flat) abelian groups. So the Div-atomic modules are exactly the $\mathfrak{b}_{\mathbb{Z}}$ -Mittag-Leffler, or simply \mathbb{Z} -Mittag-Leffler, abelian groups.
- (6) Consequently, every abelian group is \mathbb{Z} -Mittag-Leffler, which is a special case of Goodearl's result that every left module over a left noetherian ring R is R -Mittag-Leffler (and conversely), see [Goo, p.1] (or [R3, Cor.4.3(2)]) and Cor. 5.8 below.

Example 5.2 (Divisible modules over RD domains, special case). *If R is a countable and left noetherian RD domain, the situation is the exact same as over \mathbb{Z} , since 'finitely generated' = 'finitely presented' and 'countable' = 'countably generated'.*

Recall, a submodule N of a module M is said to be RD-pure or relatively divisible if $rM \cap N = rN$ for every $r \in R$. (This is purity for the pp formulas of the form $r|x$ with $r \in R$.) An RD ring is a ring over which RD-purity (relative divisibility) is (full) purity, or, equivalently, every finitely presented module is a direct summand of a direct sum of cyclically presented modules. A ring is RD iff every pp formula is equivalent to a finite conjunction of basic RD-formulas, i.e., formulas of the form $r|\mathfrak{b}x$ with $r \in R$ and \mathfrak{b} a row vector over R iff every unary pp formula is equivalent to a finite conjunction of unary basic RD-formulas $r|sx$ ($r, s \in R$). This property is left-right symmetric. See [P2, §2.4.2] or [PPR] for all this.

The argument in (1) above can be employed to yield a much farther reaching statement.

Proposition 5.3. *Suppose, all embeddings of (left) R -modules are \mathcal{L} -pure.*

- (1) *Then all countably presented modules are \mathcal{L} -atomic (= \mathcal{K} -Mittag-Leffler).*
- (2) *If R is, in addition, left noetherian, all modules are \mathcal{L} -atomic (= \mathcal{K} -Mittag-Leffler).*

Proof. (1). By hypothesis, every chain of morphisms of (left R -) modules is pure on images. Hence, every ω -limit of finitely presented modules, in particular, every countably presented module, is an \mathcal{L} - ω -limit and therefore \mathcal{L} -atomic (= \mathcal{K} -Mittag-Leffler). (See, for example, [GT, Lemma 2.11 (or §13.2.2)] for the fact that countably presented modules are ω -limits of finitely presented modules.)

(2). Every countably generated module is the union of an ascending chain of finitely generated submodules. Any chain is, by the first hypothesis, an \mathcal{L} -pure chain. On the other hand, as R is noetherian, finitely generated is the same as finitely presented. Consequently, every countably generated module is an \mathcal{L} - ω -limit (even with all maps monic) and thus \mathcal{L} -atomic. Then every module is a directed union of an ω -closed (i.e., closed under unions of countable subchains) system of \mathcal{L} -atomic submodules. By a result of Herbera–Trlifaj, see [R3, Cor. 11], this suffices to make *all* modules \mathcal{L} -atomic. \square

One obtains a classical instance with $\mathcal{L} = R\text{-Mod}$ (and $\mathcal{K} = \text{Mod-}R$), for which all embeddings of (left) modules are \mathcal{L} -pure (= pure) if and only if R is von Neumann regular if and only all embeddings of right modules are \mathcal{K} -pure (= pure). As a von Neumann regular ring is one-sided noetherian if and only if it is semisimple artinian, (2) below should come to no surprise. For (1), recall from [RG] (or Thm. 3.1(4) above) that countably generated Mittag-Leffler modules are pure projective, hence projective in case they are flat.

Corollary 5.4. *Let R be von Neumann regular.*

- (1) *All countably presented (left or right) R -modules are Mittag-Leffler, hence projective.*
- (2) *If R is, in addition, noetherian, all (left or right) R -modules are Mittag-Leffler.*

Next we apply the proposition in order to extend Example 5.2 to arbitrary RD domains. Recall first that over *any* RD ring, on either side, the divisible modules are exactly the absolutely pure modules and the torsion-free modules are exactly the flat modules ([PPR, Lemma 2.16] or [P2, Prop.2.4.16]), so $\mathcal{L} = {}_R\text{Div} = {}_R\sharp$ and $\mathcal{K} = \text{Tf}_R = \flat_R$ (with Tf the class of torsion-free modules). In particular, \mathcal{K} -Mittag-Leffler is the same as R -Mittag-Leffler. If R is a domain (RD or not), on either side, Div is a definable subcategory for the same simple reason as over \mathbb{Z} , see above; similarly Tf is one too.

Remember, (3) below is a special case of Goodearl's result mentioned in Example 5.1(6).

Corollary 5.5. *Let R be an RD domain and $\mathcal{L} = {}_R\text{Div} = {}_R\sharp$.*

- (1) *Then all countably presented left R -modules are \mathcal{L} -atomic (= R -Mittag-Leffler).*
- (2) *Similarly, for right modules.*
- (3) *If R is, in addition, left noetherian, all modules are \mathcal{L} -atomic (= R -Mittag-Leffler).*

Proof. The proof in Example 5.1(1) above that all embeddings are \mathcal{L} -pure, did in fact use only the fact that the ring $R = \mathbb{Z}$ was an RD domain. \square

Note that by [PPR, Rem.5.8], not every left noetherian RD domain is right noetherian, so in contrast to (1) and (2), statement (3) is not left-right symmetric.

We now turn to a broader reason for this behavior of RD domains.

Lemma 5.6. *If R is left coherent, all embeddings in $R\text{-Mod}$ are ${}_R\sharp$ -pure.*

Proof. By [P1, Thm.15.41], every injective (left) R -module has complete elimination of quantifiers if and only if R is left coherent. (Taking an injective model of the largest theory of injectives shows that that this is equivalent to complete quantifier elimination *universally* for all injectives.)

Let $M \subseteq N$ be any inclusion in $R\text{-Mod}$. To show it is ${}_R\sharp$ -pure, let \bar{a} , an arbitrary tuple in M , satisfy a certain pp formula φ in N . We must find a formula $\alpha \leq_{{}_R\sharp} \varphi$ that \bar{a} satisfies in M . Being existential, φ holds of \bar{a} also in an injective envelope $E(N)$ of N . But there it is ${}_R\sharp$ -equivalent to a quantifier-free formula α . Then α holds of \bar{a} in $E(N)$, hence, being quantifier-free, also in M . \square

Remark 5.7. This phenomenon is a consequence of the classical result of Eklof and Sabbagh that over a left coherent ring the theory of all (left) modules has a model-completion (and conversely). One would expect it to take place, mutatis mutandis, in other classes of modules having a model-completion, cf. [P1, §15.3].

- Corollary 5.8.** (1) *If R is left coherent, all countably presented left R -modules are \mathcal{L} -atomic (= R -Mittag-Leffler).*
 (2) (Goodearl) *If R is left noetherian, all modules are \mathcal{L} -atomic (= R -Mittag-Leffler).*

Question 5.9 (Converses). *The converse of (2) is part of Goodearl's classical result, [Goo, p.1] (see [R1, Cor.2.7] or [R3, Cor.4.3(2)] for a model-theoretic proof).*

Is the converse of (1) also true? How about the converse of the lemma? Are all inclusions \mathcal{L} -pure if all (countably presented) modules are \mathcal{L} -atomic (cf. Prop. 5.3)? At least for $\mathcal{L} = {}_R\sharp$?

For a final application of the proposition I first introduce some terminology.

- Definition 5.10.** (1) [R5]. A **high formula** is a unary pp formula γ that defines the entire module in every injective (equivalently, in every absolutely pure) module, i.e., $\gamma(E) = E$ for every injective E . The collection of all high formulas is denoted by Γ , with a left or right subscript for the ring, if necessary.
- (2) A **left high ring** is a ring over which every unary left pp formula is equivalent to a finite conjunction of high formulas and quantifier-free formulas.

- Remark 5.11.** (1) Clearly, Γ is closed under finite conjunction (and so is the set of qf formulas). Hence over a high ring, every unary pp formula is equivalent to the conjunction of a high formula and a quantifier-free formula.
- (2) High rings figured (without a specific name) in [R5, Cor. 6.24], where they were shown to admit a decomposition theorem for pure injectives into an ‘Ulm length 0 module’ and a ‘reduced module,’ see there for terminology.
- (3) [R5, Rem. 6.25]. *RD-domains* are two-sided high, as every basic RD-formula is either high or quantifier-free, see [R5, Cor. 2.12(1)].

- Question 5.12.** (1) *Is highness of rings left-right symmetric? Presumably, not.*
- (2) *Find examples of high rings other than RD domains.*

Remark 5.13. The known arguments for (1) (and Question 6.4(1) below) to have affirmative answers when high formulas are replaced by RD-formulas do not seem to work here, since highness of a formula $\mathfrak{A}|bx$ is not given by the matrix \mathfrak{A} alone, but by the condition $l(\mathfrak{A}) \subseteq l(\mathfrak{b})$, [R5, Prop. 2.8(2)].

Cor. 5.5 can be generalized to high Warfield rings. Recall from [Pu] (or [PPR] and [P2]) that a ring is called **left Warfield** if every finitely presented left module is a direct summand of a direct sum of cyclic finitely presented modules. This is equivalent to saying that every left pp formula (in the variables \bar{x}) is equivalent to a finite conjunction of formulas of the form $\mathfrak{a}|\mathfrak{b}\bar{x}$, where \mathfrak{a} and \mathfrak{b} are row vectors. (This is a special case of the much more general [PPR, Thm. 2.5] or [P2, Cor. 2.4.3].) Clearly, RD rings are left and right Warfield (and, by a result of Puninski, conversely, every two-sided Warfield ring is RD, see the above sources).

Corollary 5.14. *Let R be a left high, left Warfield ring and $\mathcal{L} = {}_R\text{Div} = {}_R\sharp$.*

- (1) *Then all countably presented left R -modules are \mathcal{L} -atomic (= R -Mittag-Leffler).*
- (2) *If R is, in addition, left noetherian, all modules are \mathcal{L} -atomic (= R -Mittag-Leffler).*

Proof. We are going to verify that every embedding is \mathcal{L} -pure. To this end, consider modules $M \subseteq N$ and a tuple \bar{a} in M satisfying a certain pp formula φ in N . All we have to do is find a pp formula $\psi \leq_{\mathcal{L}} \varphi$ that \bar{a} satisfies in M . As R is Warfield, φ is equivalent to a conjunction of formulas φ_i ($i < m$) of the form $\mathfrak{a}_i|\mathfrak{b}_i\bar{x}$ with \mathfrak{a}_i and \mathfrak{b}_i row vectors. Let θ_i be the (unary) formula $\mathfrak{a}_i|x$, $i < m$. By highness, every θ_i is equivalent to a conjunction $\gamma_i \wedge \alpha_i$ with $\gamma_i \in \Gamma$ and α_i quantifier-free. Note, $\varphi \sim \bigwedge_{i < m} \theta_i(\mathfrak{b}_i\bar{x})$. Let ψ be the conjunction of all the $\alpha_i(\mathfrak{b}_i\bar{x})$. Clearly \bar{a} satisfies ψ in N . As ψ is quantifier-free, it does so also in M . It remains to verify $\psi \leq_{\mathcal{L}} \varphi$. To this end let $L \in \mathcal{L} = {}_R\sharp$ and $\bar{c} \in \psi(L)$, i.e., $\mathfrak{b}_i\bar{c} \in \alpha_i(L)$ for all i . By highness of γ_i ,

we have also $\mathfrak{b}_i\bar{c} \in L = \gamma_i(L)$, hence $\mathfrak{b}_i\bar{c} \in \theta_i(L)$ for all i . Consequently, $\bar{c} \in \varphi(L)$, as desired. \square

6. CONCLUDING REMARKS: HIGH PURITY

One may call submodule M of a module N **high-pure** if $\gamma(N) \cap M = \gamma(M)$ for every $\gamma \in \Gamma$.

Lemma 6.1. *Over domains, high purity implies RD-purity.*

Consequently, over RD-domains (full) purity, high purity and RD-purity are all the same.

Proof. By [R5, Cor. 2.11(5)], the formula $r|x$ is high iff r is not a right zero divisor. Over a domain this is true iff $r \neq 0$. For $r = 0$, the condition $rM \cap N = rN$ is trivial. For $r \neq 0$, we infer it from high purity. \square

Question 6.2. *What can be said about RD-rings in general?*

Remark 6.3. As quantifier-free formulas always pass down, over left high rings, high purity is purity.

Question 6.4. (1) *Is a ring high if high purity is purity?*

(2) *For full purity, RD-purity, and high purity, respectively, it suffices to inspect unary pp formulas. What about \sharp -purity? For what \mathcal{L} would the same be true?*

REFERENCES

- [GT] Göbel, R. and Trlifaj, J., *Approximations and Endomorphism Algebras of Modules*, vol.1, 2nd ed., de Gruyter 2012.
- [Goo] Goodearl, K. R., Distributing tensor product over direct product, *Pacific J. Math.* **43** (1972) 107-110.
- [P1] Prest, M., *Model Theory and Modules*, London Math. Soc. Lecture Note Series, vol. 130, Cambridge University Press 1987.
- [P2] Prest, M., *Purity, Spectra and Localisation*, Encyclopedia of Math. and its Applications, vol. 121, Cambridge University Press 2009.
- [P3] Prest, M., Strictly atomic modules in definable categories, arXiv:2109.13750 [math.RT], 2021, 30 pp.
- [Pu] Puninski, G., *Warfield rings*, Algebra and Logic **33**, (1994) 147–159.
- [PPR] Puninski, G., Prest, M., and Rothmaler, Ph., Rings described by various purities, *Comm. in Algebra* **27**(5) (1999) 2127-2162.
- [RS] Rada, J. and Saorin, M., Rings characterized by (pre)envelopes and (pre)covers of their modules, *Comm. Algebra* **26**(3) (1998) 899–912.
- [RG] Raynaud, M. and Gruson, L. : Critères de platitude et de projectivité, *Seconde partie*, *Invent. Math.* **13** (1971) 52-89.
- [R1] Rothmaler, Ph., *Mittag-Leffler modules and positive atomicity*, Habilitationsschrift, Christian-Albrechts-Universität zu Kiel 1994.
- [R2] Rothmaler, Ph., Torsion-free, Divisible, and Mittag-Leffler Modules, *Comm. Algebra* **43**:8 (2015) 3342-3364.
- [R3] Rothmaler, Mittag-Leffler modules and definable subcategories, in ‘Model Theory of Modules, Algebras and Categories,’ eds. A. Facchini, L. Gregory, S. L’Innocente, M. Tressl, *Contemporary Mathematics* **730** (2019) 171–196.
- [R4] Rothmaler, Ph., Strict Mittag-Leffler modules and purely generated classes, in ‘Categorical, Homological and Combinatorial Methods in Algebra,’ eds. A. K. Srivastava, A. Leroy, I. Herzog, P. A. Guil Asensio, *Contemporary Mathematics* **751** (2020) 303-327.
- [R5] Rothmaler, Ph., High and low formulas in modules, arXiv:2111.01986 [math.RA], 2021, 47pp.