

Search for Close Stellar Encounters with the Solar System Based on Data from the Gaia DR3 Catalogue

V. V. Bobylev¹, A. T. Bajkova

*Pulkovo Astronomical Observatory, Russian Academy of Sciences,
Pulkovskoe sh. 65, St. Petersburg, 196140 Russia*

Abstract—A search for close encounters of stars with the Solar System was performed using data from the Gaia DR3 catalog. We considered 31 stars with the approach parameter $d_{\min} < 1$ pc. Among them, 15 stars are appearing as candidates for close encounters for the first time. The status of the stars GJ 710 and HD 7977 has been confirmed as candidates for deep penetration into the inner region of the Oort cloud. In particular, for GJ 710 and HD 7977, respectively, the following estimates of approach parameters are obtained: $t_{\min} = 1.324 \pm 0.026$ Myr and $d_{\min} = 0.052 \pm 0.002$ pc, $t_{\min} = -2.830 \pm 0.025$ Myr and $d_{\min} = 0.071 \pm 0.027$ pc. Among the newly identified candidates, the most interesting is the white dwarf WD 0810-353, for which the following approach parameters were found: $t_{\min} = 0.029 \pm 0.001$ Myr and $d_{\min} = 0.150 \pm 0.003$ pc.

INTRODUCTION

Close (less than ~ 1 pc) encounters of field stars with the Solar System can lead to perturbation of the cometary Oort Cloud (Oort, 1950), which can provoke the emergence of the so-called comet shower from the outer boundaries of the Oort Cloud (~ 0.5 pc) to the inner region of the Solar System, and towards the Earth in particular. In addition to stellar flybys, the cometary Oort Cloud is subject to perturbations from giant molecular clouds, and is also affected by the Galactic tide.

The early history of searches for close encounters of stars with the Solar System using various catalogs is reflected in the works of Revina (1988), Matthews (1994), Mülläri, Orlov (1996), Garcia-Sánchez et al. (1999; 2001), Bobylev (2010a; 2010b), Anderson, Francis (2012), Dybczyński, Berski (2015), Bailer-Jones (2015), Feng, Bailer-Jones (2015). In particular, it was found that one of the record holders, a candidate for penetration into the Oort Cloud, is the star GJ 710 (Garcia-Sanchez et al., 2001; Bobylev, 2010a; Berski and Dybczyński, 2016).

The Gaia space experiment (Prusti et al. 2016) is dedicated to the determination of high-precision trigonometric parallaxes, proper motions, and a number of photometric characteristics of more than 1.5 billion stars. Using kinematic data on stars from various versions of this project, a large number of works have been performed on the search for close encounters of stars with the Solar System (Berski, Dybczyński, 2016; Bobylev and Bajkova, 2017;

¹vbobylev@gaoran.ru

Fuente Markos and Fuente Markos, 2018; Bailer-Jones et al., 2018; Darma et al., 2019; Torres et al., 2019; Wysoczańska et al., 2020; Bobylev and Bajkova, 2020). Thus, the analysis of data from the Gaia TGAS catalog (Tycho-Gaia Astrometric Solution, Lindegren et al. 2016) led to the discovery of several new candidates for a very close flyby, namely, for getting inside the Oort cloud to distances less than 0.5 pc. Based on the data from the Gaia DR2 catalog (Brown et al. 2018), it was found that approaches to the Solar System to distances less than 5 pc over the time interval ± 5 Ma may have about 3000 stars, less than 1 pc — about 30 stars and 5–6 stars can have approaches to distances less than 0.25 pc.

In the Gaia EDR3 version (Gaia Early Data Release 3, Brown et al. 2021), the values of trigonometric parallaxes and stellar proper motions have been significantly refined compared to the previous version. In this case, the radial velocities of the stars were copied from the Gaia DR2 catalog. Works of Bobylev and Bajkova (2021) and Dybczyński et al. (2022) are devoted to the analysis of these data in order to search for close encounters of stars with the Solar System. In particular, Dybczyński et al. (2022) created an updated StePPeD electronic database (Stellar Potential Perturbers Database¹), which contains a large amount of information about all currently known candidate stars for close encounters with the Solar System.

In the recently published version of Gaia DR3 (Vallenari et al. 2022), the radial velocities of stars are significantly improved — previously measured values are refined and new ones are determined for a large number of stars. The values of trigonometric parallaxes and stellar proper motions were copied from an earlier version of the catalog, Gaia EDR3.

The purpose of this work is to search for close encounters of stars with the solar system using the latest data from the latest version of the Gaia — Gaia DR3 catalog.

METHODS

Linear method

From observations, the radial velocity V_r and two projections of the tangential velocity $\mu_\alpha \cos \delta$ and μ_δ are known. The heliocentric distance to the star d is calculated in terms of its trigonometric parallax π , $d = 1/\pi$. The parallax π is expressed in milliseconds of arc (mas), and the proper motion components $\mu_\alpha \cos \delta$ and μ_δ are expressed in milliseconds of arc per year (mas/yr). In the present work, very close stars with relative parallax errors much smaller than 10% are used, so there is no need to take into account any corrections like the Lutz-Kelker effect (1973).

The minimum distance between the trajectories of the star and the Sun d_{\min} at the moment of approach t_{\min} is calculated based on the following relations (Matthews, 1994):

$$\begin{aligned} d_{\min} &= d / \sqrt{1 + (V_r/V_t)^2}, \\ t_{\min} &= -d \cdot V_r / (V_t^2 + V_r^2), \end{aligned} \tag{1}$$

where d is a heliocentric distance to the star in pc, V_r is the star velocity along the line of sight and $V_t = 4.74d\sqrt{\mu_\alpha^2 \cos^2 \delta + \mu_\delta^2}$ is the star velocity perpendicular to the line of sight. Since d is expressed here in pc, and the velocities V_r and V_t are in km/s, then d_{\min} will be expressed in pc, and t_{\min} in million years.

¹<https://pad2.astro.amu.edu.pl/stars/>

Construction of galactic orbits of stars

To construct the galactic orbits of stars, we use the axisymmetric gravitational potential of the Galaxy Φ , which is represented as the sum of three components — the central spherical bulge Φ_b , the disk Φ_d , and the massive spherical dark matter halo Φ_h :

$$\Phi = \Phi_b + \Phi_d + \Phi_h. \quad (2)$$

The Φ_b bulge and Φ_d disk potentials are presented in the form proposed by Miyamoto and Nagai (1975), and the halo component is presented according to Navarro et al. (1997). The specific values of the parameters of the potential model used can be found in the works of Bajkova and Bobylev (2016; 2017), where it is designated as model III.

The galactic orbits of stars are constructed in a coordinate system related to the Local Standard of Rest. Therefore, the peculiar velocity of the Sun with the values $(U, V, W)_\odot = (11.0, 12.0, 7.2)$ km/s from Schönrich et al. (2010) is excluded from the initial velocities. The elevation of the Sun above the galactic plane $h_\odot = 16$ pc is also taken into account (Bobylev and Bajkova 2016). The gravitational interaction between the star and the Sun is not taken into account.

In the work of Bobylev and Bajkova (2021), when analyzing data from the Gaia EDR3 catalog, the parameters of the approach of stars were calculated using the linear method, the method of integrating orbits in an axisymmetric potential, and also integrating the orbits in the potential, taking into account the spiral density wave. It was concluded that the results obtained by the second and third methods are practically the same. Therefore, in this paper, we integrate the orbits of stars without taking into account the spiral density wave.

Estimating parameter errors

The errors in determining Δd_{\min} and t_{\min} are estimated using the Monte Carlo method. It is assumed that the errors in the kinematic parameters of the stars are distributed according to the normal law with a root-mean-square deviation σ .

When using the linear method, measurement errors are added to the proper motion components $\mu_\alpha \cos \delta$ and μ_δ , parallax π , and radial velocity of the star V_r . In the method of constructing galactic orbits in the potential of the Galaxy, measurement errors are added to the rectangular coordinates X, Y, Z of stars and their spatial velocities U, V, W .

DATA

First, from the catalog Gaia DR3, placed in the Strasbourg database under the number I/355, we selected stars using the following restrictions:

$$\pi > 7 \text{ mas}, \quad \sigma_\pi/\pi < 15\%, \quad \sigma_{V_r} < 15 \text{ km/s}. \quad (3)$$

There were about 350 000 stars in this preliminary sample. Then we examine this sample for close approaches by two methods. Using the recommendations made in Bobylev and Bajkova (2021), we first select candidates based on a simpler linear method, then we apply a more complex second method for estimating the approach parameters using the gravitational potential of the Galaxy to the selected stars.

Table 1: Initial data on stars

Gaia EDR3	π , mas	$\mu_\alpha \cos \delta$, mas/yr	μ_δ , mas/yr	V_r , km/s
4270814637616488064	$52.39 \pm .02$	$-0.414 \pm .019$	$-0.108 \pm .017$	-14.42 ± 0.26
510911618569239040	$13.21 \pm .03$	$0.144 \pm .024$	$0.010 \pm .034$	26.76 ± 0.21
5544743925212648320	$89.50 \pm .02$	$-65.479 \pm .016$	$-29.204 \pm .018$	-373.7 ± 8.2
5571232118090082816	$10.23 \pm .01$	$0.094 \pm .010$	$0.461 \pm .011$	82.48 ± 0.42
3372104035275483392	$11.97 \pm .03$	$0.557 \pm .043$	$0.438 \pm .031$	-47.45 ± 3.74
929788371508812288	$13.69 \pm .06$	$-2.043 \pm .053$	$3.232 \pm .042$	195.7 ± 6.6
3118526069444386944	$7.60 \pm .05$	$0.252 \pm .049$	$0.045 \pm .049$	40.07 ± 0.55
4116451378388951424	$11.86 \pm .33$	$-2.102 \pm .226$	$3.655 \pm .184$	-264.4 ± 5.5
1952802469918554368	$141.89 \pm .02$	$161.454 \pm .017$	$-119.74 \pm .019$	-82.91 ± 0.18
3295253979286613376	$9.33 \pm .02$	$0.261 \pm .026$	$-0.622 \pm .020$	-61.63 ± 1.89
5261593808165974784	$15.35 \pm .01$	$-0.091 \pm .012$	$-2.212 \pm .015$	71.08 ± 0.58
3054509410098672000	$24.96 \pm .02$	$0.630 \pm .023$	$5.493 \pm .020$	70.35 ± 0.35
3600338081985998080	$25.90 \pm .09$	$-3.175 \pm .111$	$-0.317 \pm .054$	-37.17 ± 1.95
398496965625177216	$13.72 \pm .04$	$-2.817 \pm .046$	$-3.043 \pm .034$	-151.4 ± 7.9
4536673181955253504	$10.19 \pm .25$	$0.739 \pm .186$	$-0.729 \pm .261$	-71.67 ± 0.86
2926732831673735168	$8.85 \pm .01$	$-0.737 \pm .009$	$0.533 \pm .013$	66.46 ± 0.15
3260079227925564160	$32.11 \pm .03$	$-3.615 \pm .030$	$-4.959 \pm .020$	-32.71 ± 0.37
3621143693841328896	$7.52 \pm .05$	$0.860 \pm .054$	$-0.148 \pm .046$	85.2 ± 9.0
3320184202856435840	$123.19 \pm .02$	$-444.69 \pm .018$	$-925.14 \pm .015$	-414.0 ± 10.4
5473864079915092736	$9.98 \pm .05$	$8.481 \pm .047$	$9.750 \pm .051$	712.5 ± 14.0
5553958176239495040	$40.27 \pm .02$	$-24.002 \pm .031$	$-0.309 \pm .030$	78.61 ± 3.17
1791617849154434688	$11.38 \pm .02$	$-0.386 \pm .014$	$-1.172 \pm .013$	55.82 ± 0.25
5346007675222666752	$44.99 \pm .02$	$-18.348 \pm .017$	$-21.173 \pm .016$	76.29 ± 0.41
6396469681261213568	$9.80 \pm .02$	$0.490 \pm .014$	$0.188 \pm .016$	52.22 ± 0.19
6224087389269263488	$31.43 \pm .15$	$-3.235 \pm .175$	$-0.553 \pm .157$	18.03 ± 0.45
3972130276695660288	$59.92 \pm .03$	$-20.813 \pm .030$	$6.629 \pm .020$	31.14 ± 0.17
1926461164913660160	$316.48 \pm .04$	$112.527 \pm .036$	$-1591.6 \pm .027$	-77.29 ± 0.19
5853498713190525696	$768.07 \pm .05$	$-3781.7 \pm .03$	$769.465 \pm .051$	-21.94 ± 0.22
4155835025908320640	$10.50 \pm .04$	$5.892 \pm .038$	$0.062 \pm .030$	-283.0 ± 10.7
2933503521200215424	$7.43 \pm .02$	$0.248 \pm .013$	$-0.020 \pm .016$	32.66 ± 0.73
4306481867124380672	$9.75 \pm .04$	$-0.523 \pm .024$	$0.145 \pm .022$	-33.21 ± 0.19

Table 2: Additional information on stars

Gaia EDR3	alternative designation	StePPeD	Mass, M_{\odot}
4270814637616488064	GJ 710	P0107	0.65
510911618569239040	HD 7977	P0230	1.10
5544743925212648320	WD 0810-353		
5571232118090082816	UCAC4 237-008148, AB	P0506	0.77
3372104035275483392			
929788371508812288			
3118526069444386944		P0533	0.87
4116451378388951424			
1952802469918554368	LSPM J2146+3813	P0416	0.20
3295253979286613376			
5261593808165974784	UCAC4 076-006432	P0522	0.55
3054509410098672000			
3600338081985998080			
398496965625177216			
4536673181955253504	UCAC4 574-064555	P0567	0.60
2926732831673735168	BD-21 1529	P0287	1.02
3260079227925564160	UCAC4 464-006057	P0526	0.45
3621143693841328896			
3320184202856435840			
5473864079915092736			
5553958176239495040	UCAC4 213-008644		
1791617849154434688	TYC 1662-1962-1	P0189	0.71
5346007675222666752			
6396469681261213568	TYC 9327-264-1	P0382	0.89
6224087389269263488	TYC 6760-1510-1		
3972130276695660288	GJ 3649	P0178	0.55
1926461164913660160	GJ 905	P0413	0.15
5853498713190525696	Proxima Cen	P1037-C	
4155835025908320640			
2933503521200215424	HD 49995	P0286	1.32
4306481867124380672	HD 179939	P0111	1.69

Table 3: Parameters for the approach of stars to the Solar System

Gaia EDR3	$t_{\min},$ Myr	$d_{\min},$ pc	$t_{\min},$ Myr	$d_{\min},$ pc	$\sigma_t,$ Myr	$\sigma_d,$ pc
	(1)		(2)			
4270814637616488064	1.325	0.051	1.324	0.052	0.026	0.002
510911618569239040	−2.826	0.146	−2.830	0.071	0.025	0.027
5544743925212648320	0.030	0.113	0.030	0.164	0.001	0.003
5571232118090082816	−1.185	0.258	−1.185	0.203	0.007	0.006
3372104035275483392	1.759	0.493	1.762	0.476	0.156	0.062
929788371508812288	−0.373	0.494	−0.374	0.513	0.013	0.018
3118526069444386944	−3.281	0.524	−3.288	0.529	0.054	0.108
4116451378388951424	0.319	0.537	0.320	0.540	0.012	0.047
1952802469918554368	0.084	0.569	0.085	0.552	0.001	0.001
3295253979286613376	1.739	0.596	1.741	0.556	0.058	0.022
5261593808165974784	−0.916	0.626	−0.917	0.652	0.008	0.009
3054509410098672000	−0.569	0.598	−0.570	0.658	0.003	0.003
3600338081985998080	1.038	0.606	1.037	0.670	0.060	0.039
398496965625177216	0.481	0.689	0.482	0.730	0.028	0.038
4536673181955253504	1.368	0.660	1.368	0.735	0.041	0.150
2926732831673735168	−1.700	0.828	−1.701	0.786	0.007	0.006
3260079227925564160	0.951	0.862	0.953	0.800	0.008	0.015
3621143693841328896	−1.561	0.858	−1.558	0.809	0.156	0.062
3320184202856435840	0.019	0.771	0.020	0.842	0.001	0.021
5473864079915092736	−0.141	0.863	−0.142	0.857	0.054	0.108
5553958176239495040	−0.315	0.892	−0.317	0.859	0.058	0.022
1791617849154434688	−1.574	0.809	−1.574	0.860	0.026	0.002
5346007675222666752	−0.291	0.859	−0.292	0.861	0.013	0.018
6396469681261213568	−1.953	0.495	−1.949	0.872	0.008	0.015
6224087389269263488	−1.763	0.873	−1.761	0.901	0.001	0.001
3972130276695660288	−0.534	0.925	−0.534	0.912	0.060	0.039
1926461164913660160	0.037	0.933	0.038	0.917	0.008	0.009
5853498713190525696	0.027	0.958	0.028	0.939	0.158	0.191
4155835025908320640	0.337	0.895	0.337	0.940	0.003	0.003
2933503521200215424	−4.118	0.653	−4.113	0.980	0.098	0.038
4306481867124380672	3.086	0.814	3.086	0.980	0.001	0.003

Note. (1) — linear method, (2) — axisymmetric potential.

RESULTS

Tables 1–3 show the results of the search for candidate stars for close encounters with the Solar System. Table 1 gives the initial data on stars extracted from the Gaia DR3 catalog.

Table 2 gives the numbers of selected stars from the Gaia EDR3 catalog and their alternative designations, as well as numbers from the StePPeD database (Dybczyński et al., 2022) and mass estimates taken from the StePPeD database. Note that for many stars in Table 2 there are no alternative designations. Moreover, most of these stars are missing from the SIMBAD ² search database. That is, the complete set of kinematic parameters ($\pi, \mu_\alpha \cos \delta, \mu_\delta$, and V_r) necessary for analysis for such stars, appeared for the first time.

Table 3 gives the parameters of the approach of stars to the Solar System t_{\min} and d_{\min} calculated by two methods. In tables 1–3 the stars are sorted by the parameter d_{\min} , found when constructing orbits using the axisymmetric potential (method (2) in Table reftab-3). The Tables give 31 stars with $d_{\min} < 1$ pc.

DISCUSSION

As can be seen from Table 2, the masses of candidates are small. The magnitude of the perturbation that it can exert on the cometary cloud strongly depends on the mass of a passing star. The most massive star of our list occupies the last line in the tables. The outer boundary of the cometary Oort cloud is not exactly known. It is assumed that it is about 0.48 pc. In our opinion, 5-6 stars, in which the value of the parameter d_{\min} is inside the outer boundary of the Oort cloud, may be of the greatest interest. We will discuss them in more detail.

GJ 710 and HD 7977. The first two stars in Tables 1–3 and earlier, when using data from the Gaia EDR3 catalog, occupied the first two lines of candidates for the closest encounters (Bobylev and Bajkova, 2021). We note that the random errors in the estimates of the parameters t_{\min} and d_{\min} , have approximately halved, which is due to improved radial velocities for these stars in the Gaia DR3 catalog.

Fuente Marcos and Fuente Marcos (2022) estimated the parameters of the approach of the star GJ 710 using data from the Gaia DR3 catalog. These authors have shown that the distribution of the distances of the minimum approach to the Solar System has a median value of 0.052 pc, and with a probability of 90% is in the range of 0.048–0.056 pc; the corresponding perihelion passage time lies in the range of 1.26 and 1.33 million years with a confidence of 90%, with the most likely value of 1.29 million years. The values of these parameters found by us are in good agreement with the estimates of Fuente Marcus and Fuente Marcus.

WD 0810-353 ranks third in the tables 1–3. This star appears for the first time in the list of candidates for the closest approaches to the solar system. According to the SIMBAD database, this is a white dwarf with large proper motions, therefore it is classified as HPMS (High Proper Motion Star). According to Bagnulo and Landstreet (2020), this white dwarf has a mass of $0.63M_\odot$ and an age of 2.7 Gyr.

This star also has a huge radial velocity $V_r = -373.7 \pm 8.2$ km/s. For white dwarfs, it is important to take into account the gravitational redshift of spectral lines when determining the radial velocities. Such an effect is a consequence of a strong gravitational field on

²<https://simbad.u-strasbg.fr/simbad/>

the surface of white dwarfs, whose influence is equivalent to velocities of several tens of km/s (see, for example, Greenstein and Trimble (1967)). We performed new calculations with the radial velocity value $V_r = -423.7 \pm 8.2$ km/s, adding -50 km/s to the measured value, and found the following approach parameters: $t_{\min} = 0.029 \pm 0.001$ million years and $d_{\min} = 0.150 \pm 0.003$ pc. In this case, taking into account the correction for the gravitational redshift does not have a critical effect on the estimation of the rendezvous parameters.

In general, we can conclude that in about 30 thousand years the star will rapidly sweep past the solar system. Its influence on the objects of the Oort cloud will be very short.

UCAC4 237-008148. This low-mass binary star is number four in the tables 1–3. For example, in the work of Dybczyński et al. (2022), when analyzing data from the Gaia EDR3 catalog, the following approach parameters were found for it: $t_{\min} = -1.084 \pm 0.004$ Myr and $d_{\min} = 0.199 \pm 0.008$ pc. We can see that there is good agreement with the data in Table 3, especially good agreement is in the estimation of parameter errors.

Gaia EDR3 3372104035275483392 and Gaia EDR3 929788371508812288. These two stars rank fifth and sixth in our tables. They appear for the first time in the list of candidates for the closest approaches to the solar system. Unfortunately, they are not in the SIMBAD database, so little is known about these stars.

In general, we can conclude that due to the use of selection criteria (3), the approach parameters of all stars are determined reliably, since the errors in determining these parameters are small (see the last two columns of Table 3) .

CONCLUSION

A search for close encounters of stars with the Solar System was carried out using data from the latest Gaia DR3 catalog. The paper considers 31 stars with the approach parameters $d_{\min} < 1$ pc. Among them, 15 stars for the first time act as candidates for close encounters with the solar system. These 15 stars were identified mainly due to the appearance of measurements of their radial velocities in the Gaia DR3 catalog.

For the stars GJ 710 and HD 7977, their status as candidates for very close encounters, candidates for deep penetration into the inner region of the Oort cloud, has been confirmed.

Among the newly identified candidates, the most interesting is the white dwarf WD 0810-353. For this star, the following approach parameters were found, taking into account the model correction for the effect of gravitational redshift in spectral lines when measuring its radial velocity: $t_{\min} = 0.029 \pm 0.001$ Myr and $d_{\min} = 0.150 \pm 0.003$ pc.

REFERENCES

1. E. Anderson, et al., *Astron. Lett.* **38**, 331 (2012).
2. S. Bagnulo and J.D. Landstreet, *Astron. Astrophys.* **643**, 134 (2020).
3. C.A.L. Bailer-Jones, *Astron. Astrophys.* **575**, 35 (2015).
4. C.A.L. Bailer-Jones, J. Rybizki, R. Andrae, and M. Fouesneau, *Astron. Astrophys.* **616**, 37 (2018).
5. A.T. Bajkova and V.V. Bobylev, *Astron. Lett.* **42**, 567 (2016).
6. A.T. Bajkova and V.V. Bobylev, *Open Astron.* **26**, 72 (2017).

7. F. Berski and P.A. Dybczyński, *Astron. Astrophys.* **595**, L10 (2016).
8. V.V. Bobylev, *Astron. Lett.* **36**, 220 (2010a).
9. V.V. Bobylev, *Astron. Lett.* **36**, 816 (2010b).
10. V.V. Bobylev and A.T. Bajkova, *Astron. Lett.* **42**, 1 (2016).
11. V.V. Bobylev and A.T. Bajkova, *Astron. Lett.* **43**, 559 (2017).
12. V.V. Bobylev and A.T. Bajkova, *Astron. Lett.* **46**, 245 (2020).
13. V.V. Bobylev and A.T. Bajkova, *Astron. Lett.* **47**, 180 (2021).
14. Gaia Collaboration, A.G.A. Brown, A. Vallenari, T. Prusti, et al., *Astron. Astrophys.* **616**, 1 (2018).
15. Gaia Collaboration, A.G.A. Brown, A. Vallenari, T. Prusti, et al., *Astron. Astrophys.* **649**, 1 (2021).
16. R. Darma, W. Hidayat, and M.I. Arifyanto, *J. Phys.: Conf. Ser.* **1245**, 012028 (2019).
17. P.A. Dybczyński and F. Berski, *MNRAS* **449**, 2459 (2015).
18. P.A. Dybczyński, F. Berski, J. Tokarek, et al., arXiv: 2206.1104 (2022).
19. F. Feng and C.A.L. Bailer-Jones, *MNRAS* **454**, 3267 (2015).
20. R. de la Fuente Marcos and C. de la Fuente Marcos, *Res. Not. Am. Astron. Soc.* **2**, 30 (2018).
21. R. de la Fuente Marcos and C. de la Fuente Marcos, *Res. Not. Am. Astron. Soc.* **6**, 136 (2022).
22. J. Garcia-Sánchez, R.A. Preston, D.L. Jones, et al., *Astron. J.* **117**, 1042 (1999).
23. J. Garcia-Sánchez, P.R. Weissman, R.A. Preston, et al., *Astron. Astrophys.* **379**, 634 (2001).
24. J.L. Greenstein and V.L. Trimble, *Astrophys. J.* **149**, 283 (1967).
25. Gaia Collaboration, L. Lindegren, U. Lammers, U. Bastian, et al., *Astron. Astrophys.* **595**, A4 (2016).
26. T.E. Lutz and D.H. Kelker, *PASP* **85**, 573 (1973).
27. M. Miyamoto and R. Nagai, *PASP* **27**, 533 (1975).
28. R.A.J. Matthews, *Royal Astron. Soc. Quart. Journ.* **35**, 1 (1994).
29. A.A. Mülläri and V.V. Orlov, *Earth, Moon, and Planets* (Kluwer, Netherlands, **72**, p. 19, 1996).
30. J.F. Navarro, C.S. Frenk and S.D.M. White, *Astrophys. J.* **490**, 493 (1997).
31. J.H. Oort, *Bull. Astron. Inst. Netherland* **11**, No 408, 91 (1950).
32. Gaia Collaboration, T. Prusti, J.H.J. de Bruijne, A.G.A. Brown, et al., *Astron. Astrophys.* **595**, 1 (2016).

- 33. I.A. Revina, *Analysis of motion of celestial bodies and estimation of accuracy of their observations* (Latvian University, Riga, p. 121, 1988).
- 34. S. Torres, M.X. Cai, A.G.A. Brown, and S. Portegies Zwart, *Astron. Astrophys.* **629**, 139 (2019).
- 35. R. Schönrich, J. Binney and W. Dehnen, *MNRAS* **403**, 1829 (2010).
- 36. Gaia Collaboration, A. Vallenari, A.G.A. Brown, T. Prusti, et al., in press (2022).
- 37. R. Wysoczańska, P.A. Dybczyński, and M. Polińska, *Astron. Astrophys.* **640**, 129 (2020).