A stochastic model solvable without integrability

F Mathieu¹ and E. Ragoucy²

Laboratoire d'Annecy-le-vieux de Physique Théorique LAPTh CNRS and Université Savoie Mont Blanc. BP 110, F-74941 Annecy-le-Vieux Cedex, France.

Abstract

We introduce a model with diffusive and evaporation/condensation processes, depending on 3 parameters obeying some inequalities. The model can be solved in the sense that all correlation functions can be computed exactly without the use of integrability.

We show that the mean field approximation is not exact in general. This can be shown by looking at the analytical expression of the two-point correlation functions, that we provide. We confirm our analysis by numerics based on direct diagonalisation of the Markov matrix (for small values of the number of sites) and also by Monte-Carlo simulations (for a higher number of sites).

Although the model is symmetric in its diffusive rates, it exhibits a left / right asymmetry driven by the evaporation/condensation processes. We also argue that the model can be taken as a one-dimensional model for catalysis or fracturing processes.

 $^{^{1}} fabien.mathieu@lapth.cnrs.fr\\$

²eric.ragoucy@lapth.cnrs.fr

1 Introduction

One-dimensional Markovian models of hard core interacting particles hopping on a lattice, such as the Asymmetric Simple Exclusion Process (ASEP), are commonly studied as solvable non-equilibrium statistical physics problems [6,19]. The ASEP model is known to be integrable, which provides powerful tools to study it [2]. Some of its variations keep the integrability property [5], but this property is rather fragile: it does generally not survive modifications. The matrix ansatz (or DEHP method, named from its authors) is another way to get insight on some models [6,17], but outside the integrability framework, the elaboration of a matrix ansatz as hard as a direct diagonalization of the Markov process [16], see also [1] for a recent review on the subject. However, asking for the stationarity of the densities (one-point correlation functions) is in general enough to get their analytical expressions. The same is true for the current, but starting with two-point correlation functions, the calculation becomes in general laborious [7]. Few examples are known for such calculations, but they are often done for fermions on graphs, see e.g. [8, 10, 15], or use the mean field approximation [18]

In the present paper, we study a model that allows a decoupling of the *n*-point correlation functions from the higher ones. We take advantage of this hierarchy to compute the correlation functions step by step. We provide explicit formulae for the densities, the currents and the 2-points correlation functions and observe that the correlation functions differ from the mean field approximation. We also provide the general recursion relations that allow to compute the *n*-point correlation functions.

The model has some characteristics of the Symmetric Simple Exclusion Process (SSEP) but also allows evaporation and condensation of particles on the lattice. The condensation and evaporation rates are free parameters of the model, obeying some inequalities to ensure the Markovian property. This model possesses two distinguished features:

- It catches and releases single particles and pairs of particles, the rate of the later being different from the rate of the former. This allows to see it as a one-dimensional Markovian model for fracturing or catalysis of molecules.
- Although the model is symmetric in its diffusive rates, the evaporation/condensation processes induce a right/left asymmetry. As a consequence, the model has some properties characteristic of a SSEP model, but also properties closer to an ASEP one.

This model is a priori not integrable, in the sense that it is not connected to the R-matrices that one can find in the literature. We come back on this point in the conclusion.

After presenting the general features of the model in section 2, we derive formulae for the one-site observables in section 3 and explicit the catalysis by comparing the evaporation currents of single particles against the evaporation current of pairs of particles. We show what combination of the parameters determines whether the lattice bounds particles together or splits them apart. In section 4, we discuss n-point correlation functions, giving an explicit formula for 2-point correlation functions and presenting the general recursion relations for any n-points correlation function. We investigate in section 5 a special case where the model is integrable and can be solved completely. Finally, in section 6, we present numerical results and plot observables.

2 The model

We study a SSEP-like model on a one-dimensional lattice where in addition to the usual symmetric interaction of particles on the lattice, evaporation from and condensation on the lattice can occur. These processes can apply to particles, and also to 'molecules' constituted of two particles sitting on two adjacent sites. The lattice is coupled to two reservoirs, that can exchange particles in the usual way. The rates can be divided in two parts: one corresponding to the DiSSEP model [2], where only 'molecules' can condensate or evaporate, plus a second additional part where single particles can also condensate or evaporate. This allows us to interpret the model as a fracturing process, with 'molecules' constituted of two 'particles' on adjacent sites, condensing or evaporating from the lattice at rates which vary from the ones of single 'particles'. Hence, the one-dimensional lattice can be thought of as a catalyst immersed in a gaz of molecules, fracturing them into particles, see also discussion in section 3.3.

The values of the different rates are shown in figures 1 and 2.




Figure 1: Dynamical rules of the model (DiSSEP part).

It leads to the following local Markov matrices. In the bulk:

$$M = \begin{pmatrix} -c_0 - c_1 - c_2 & 1 + c_2 - c_0 & c_1 + c_0 - 1 & c_0 \\ c_2 & -1 - c_1 - c_2 & 1 & c_1 \\ c_1 & 1 & -1 - c_1 - c_2 & c_2 \\ c_0 & c_1 + c_0 - 1 & 1 + c_2 - c_0 & -c_0 - c_1 - c_2 \end{pmatrix}.$$
 (2.1)

The off-diagonal terms of M are positive when

$$0 \le c_k, \ k = 0, 1, 2 \quad \text{and} \quad 1 - c_1 \le c_0 \le 1 + c_2.$$
 (2.2)

Note that these constraints are automatically satisfied if one chooses $c_1 = c_0 = c_2 \ge \frac{1}{2}$. It leads to a one-parameter model that could be worth studying.




Figure 2: Dynamical rules of the model (additional transitions for single particles).

On the left boundary, we have

$$B = \begin{pmatrix} -\alpha & \gamma \\ \alpha & -\gamma \end{pmatrix}.$$
 (2.3)

On the right boundary, we have

$$\bar{B} = \begin{pmatrix} -\delta & \beta \\ \delta & -\beta \end{pmatrix}.$$
(2.4)

The full Markovian matrix takes the form

$$\mathbb{M} = B_1 + \sum_{j=1}^{L-1} M_{j,j+1} + \bar{B}_L.$$
(2.5)

2.1 General properties

Particle/hole symmetry

The exchange of particles and holes corresponds to a conjugation

$$\begin{cases} M_{j,j+1} \to V \otimes V M_{j,j+1} V \otimes V = M_{j,j+1} \\ B \to V B V \quad ; \quad \bar{B} \to V \bar{B} V \end{cases} \quad \text{with} \quad V = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}. \tag{2.6}$$

Remark that, in the bulk, the model is invariant under this transformation: only the boundaries break it. It corresponds to the exchange $\alpha \leftrightarrow \gamma$ and $\beta \leftrightarrow \delta$.

Transposition in the bulk

The transposition of the matrix M is still a Markovian matrix. To connect the matrix M^T with a matrix of the form (2.1), one needs to reverse the orientation of the lattice, which amounts to conjugate the matrix by the permutation matrix

$$P = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Then, the matrix $M' = P M^T P$ is of the form (2.1) with parameters

$$c'_0 = c_0, \quad c'_1 = 1 + c_2 - c_0 \quad \text{and} \quad c'_2 = c_1 + c_0 - 1,$$
 (2.7)

obeying $c'_{j} \ge 0$ and $1 - c'_{1} \le c'_{0} \le 1 + c'_{2}$.

When $c_0 = 1$, M is self-transposed and P is not needed, since it commutes with M. In that case, the model is simpler and the mean field approximation becomes exact, see section 5.

Comparison with DiSSEP

A SSEP model with dissipation, called DiSSEP, has already been studied in [2,5]. Its local Markov matrix reads:

$$W = \begin{pmatrix} -\kappa & 0 & 0 & \kappa \\ 0 & -1 & 1 & 0 \\ 0 & 1 & -1 & 0 \\ \kappa & 0 & 0 & -\kappa \end{pmatrix}.$$
 (2.8)

The matrices W and M coincide only when the parameters take the values $\kappa = c_0 = 1$ and $c_1 = c_2 = 0$, which corresponds to the free fermions point, see a detailed study in [5]. We will show that for the present model, $c_0 = 1$ (with c_1 and c_2 kept free) is enough to simplify the presentation, see section 5.

Note also that we checked that for generic values of their parameters it is <u>not</u> possible to relate the two matrices M and W by transformations of the form

$$M = U \otimes U W U^{-1} \otimes U^{-1} + T \otimes \mathbb{I}_2 - \mathbb{I}_2 \otimes T + \lambda \mathbb{I}_4, \qquad (2.9)$$

where U is an arbitrary invertible 2×2 matrix, T an arbitrary 2×2 matrix corresponding to telescoping terms, and λ a free parameter.

Case of periodic lattice

In the case of a periodic lattice, where the site L + 1 is identified with the first site, the model is at equilibrium. Indeed, the stationary state can be computed, it takes the form

$$|\mathcal{S}_0\rangle = \begin{pmatrix} 1\\1 \end{pmatrix} \otimes \ldots \otimes \begin{pmatrix} 1\\1 \end{pmatrix}.$$
 (2.10)

The densities are constant, $\langle n_j \rangle = \frac{1}{2}$, $\forall j$, and there is no current. This shows the similarity between the present model and the SSEP model: it is the presence of reservoirs that brings the system out of equilibrium.

3 Currents and densities

As we shall see, when asking for the stationarity of the densities and currents, we get recursion relations involving densities only. There is a decoupling from the higher correlation functions, such that the recursion can be solved. This property extends to higher correlation functions, see section 4.

To simplify the presentation, we introduce the ratios:

$$\Lambda = \frac{\gamma - \alpha}{2c_0 + 2c_1 + \gamma + \alpha}, \quad \widehat{\Lambda} = \frac{\beta - \delta}{2 + 2c_2 + \beta + \delta} \quad \text{and} \quad \Gamma = \frac{1 - c_0}{1 + c_0 + c_1 + c_2}. \tag{3.1}$$

The parameter Λ (resp. $\widehat{\Lambda}$) encodes the dependence in the right (resp. left) boundary, while Γ corresponds to the site (i.e. the position) dependence for densities and currents (see below). When $\Gamma = 0$ or 1, there is no dependence in the position (except for the first and the last sites).

Remark that Γ is a decreasing function of $c_0 \geq 0$ with positive parameters c_1 and c_2 . Without any constraint, the extreme values of Γ are obtained for $c_0 \to \infty$ and c_1 , c_2 finite (which gives $\Gamma = -1$) and for $c_0 = 0$ with $c_1 = c_2 = 0$ (which provides $\Gamma = 1$). However, due to the constraints (2.2), when $c_0 \to \infty$, we must take $c_2 \to \infty$, while to get $c_0 = 0$ we must choose $c_1 = 1$. This leads to

$$-\frac{1}{2} \le \Gamma \le \frac{1}{2}.$$

One can define two sub-classes of models that lead to the extremal values for Γ :

• One can take $c_2 = c_0 - 1$. The inequalities (2.2) are then satisfied for $c_0 \ge 1$. Then, one gets $\Gamma = \frac{1-c_0}{2c_0+c_1}$, and the limit $c_0 \to \infty$ (keeping c_1 finite) gives $\Gamma = -\frac{1}{2}$. Up to a scaling factor c_0 , the limit provides a model with no free parameter and no diffusion. Its local Markov matrix reads

$$\lim_{c_0 \to \infty} \left(\frac{1}{c_0} M_{c_2 = c_0 - 1} \right) = \begin{pmatrix} -2 & 0 & 1 & 1\\ 1 & -1 & 0 & 0\\ 0 & 0 & -1 & 1\\ 1 & 1 & 0 & -2 \end{pmatrix} .$$
(3.2)

• One can take $c_1 = 1 - c_0$ and $c_2 = \mathfrak{c}_2 c_0$. The inequalities (2.2) are then satisfied for $c_0 \leq 1$. Then, one gets $\Gamma = \frac{1-c_0}{2+c_0\mathfrak{c}_2}$, and the value $c_0 = 0$ (keeping \mathfrak{c}_2 finite) gives $\Gamma = \frac{1}{2}$. Its local Markov matrix at $c_0 = 0$ has no free parameter and reads:

$$M' = \begin{pmatrix} -1 & 1 & 0 & 0 \\ 0 & -2 & 1 & 1 \\ 1 & 1 & -2 & 0 \\ 0 & 0 & 1 & -1 \end{pmatrix}.$$
 (3.3)

3.1 Densities

The time evolution of the densities read

$$\frac{d}{dt} \langle n_1 \rangle = \alpha + c_0 + c_1 - (2c_0 + 2c_1 + \alpha + \gamma) \langle n_1 \rangle,
\frac{d}{dt} \langle n_j \rangle = 2c_0 + c_1 + c_2 - (2 + 2c_0 + 2c_1 + 2c_2) \langle n_j \rangle + (2 - 2c_0) \langle n_{j-1} \rangle, \qquad 1 < j < L, \quad (3.4)
\frac{d}{dt} \langle n_L \rangle = \delta + c_0 + c_2 - (2 + 2c_2 + \delta + \beta) \langle n_L \rangle + (2 - 2c_0) \langle n_{L-1} \rangle.$$

Asking for stationarity, we get

$$\langle n_j \rangle = \frac{1}{2} \left(1 - \Lambda \Gamma^{j-1} \right), \qquad 1 \le j < L,$$

$$\langle n_L \rangle = \frac{1}{2} \left(1 - \widehat{\Lambda} \right) - \frac{1 - c_0}{2 + 2c_2 + \delta + \beta} \Lambda \Gamma^{L-2}.$$

$$(3.5)$$

Performing a direct diagonalisation of the Markov matrix, we checked that the above formulas correspond to the densities when L = 2, 3, 4.

Remark that if one chooses the parameters c_k in such a way that $\Gamma < 0$, the densities follow a damped oscillating form. We illustrate it in figure 3.




Figure 3: Oscillation of the densities close to the left boundary

Note that, as in the SSEP case, the differential equation for the one-point correlation function decouples from the higher correlations, despite there is evaporation and condensation. This confirms that the model with symmetric evaporation–condensation process is very similar to the SSEP.

However, although we are dealing with a SSEP-like model, which is symmetric in the bulk, the left/right symmetry is broken here by the evaporation/condensation processes. Indeed, the densities depend essentially on the left boundary, and mildly on the right one. We checked that there is no possibility to connect the model to an ASEP model through transformations of the form (2.9).

3.2 Currents

From the expression of the one-point correlation function, we can deduce the currents. There are two types of currents: the lattice current, corresponding to particles moving in the lattice, and the evaporation current corresponding to evaporation/condensation of particles and 'molecules' on the lattice.

3.2.1 Lattice current and reservoir densities

The current in the lattice reads

$$J_{0,1}^{\text{latt}} = \alpha \langle 1 - n_1 \rangle - \gamma \langle n_1 \rangle,$$

$$J_{j,j+1}^{\text{latt}} = \langle n_j (1 - n_{j+1}) \rangle - \langle (1 - n_j) n_{j+1} \rangle = \langle n_j \rangle - \langle n_{j+1} \rangle, \qquad 1 \le j \le L - 1, \qquad (3.6)$$

$$J_{L,L+1}^{\text{latt}} = \beta \langle n_L \rangle - \delta \langle 1 - n_L \rangle,$$

where by convention j = 0 and j = L + 1 stand for the left and the right reservoir respectively. Plugging the values (3.5), we get:

$$J_{0,1}^{\text{latt}} = -(c_0 + c_1)\Lambda,$$

$$J_{j,j+1}^{\text{latt}} = \frac{1}{2}\Lambda(\Gamma^j - \Gamma^{j-1}) = -\frac{2c_0 + c_1 + c_2}{2(1 + c_0 + c_1 + c_2)}\Lambda\Gamma^{j-1}, \quad 1 \le j \le L - 2,$$

$$J_{L-1,L}^{\text{latt}} = \frac{1}{2}\widehat{\Lambda} - \frac{\beta + \delta + 2c_0 + 2c_2}{2(2 + 2c_2 + \delta + \beta)}\Lambda\Gamma^{L-2},$$

$$J_{L,L+1}^{\text{latt}} = (1 + c_2)\widehat{\Lambda} - \frac{(\beta + \delta)(1 - c_0)}{2 + 2c_2 + \delta + \beta}\Lambda\Gamma^{L-2}.$$
(3.7)

Remark that generically, the currents depend only on the sum c_1+c_2 , while it depends separately on c_1 and c_2 for the first and the two last sites.

Extending the relation $J_{j,j+1}^{latt} = \langle n_j \rangle - \langle n_{j+1} \rangle$ to j = 1 and j = L+1 allows us to define the

reservoir densities:

$$\langle n_0 \rangle = \frac{1}{2} \left(1 - (1 + 2c_0 + 2c_1)\Lambda \right),$$

$$\langle n_{L+1} \rangle = \frac{\delta + (1 - \beta + \delta)(1 + c_2)}{2 + 2c_2 + \delta + \beta} - \frac{(1 - \beta - \delta)(1 - c_0)}{2 + 2c_2 + \delta + \beta} \Lambda \Gamma^{L-2}.$$
(3.8)

3.2.2 Evaporation current

To compute the evaporation current, one needs to consider three adjacent sites (because of the condensation of 'molecules'). For $2 \le j \le L - 1$, it is given by

$$J_{j}^{ev} = (1 + c_{2} - c_{0}) \left\langle (1 - n_{j-1})n_{j} - n_{j-1}(1 - n_{j}) \right\rangle + (c_{2} + c_{0}) \left\langle n_{j-1}n_{j} - (1 - n_{j-1})(1 - n_{j}) \right\rangle + (c_{1} + c_{0} - 1) \left\langle n_{j}(1 - n_{j+1}) - (1 - n_{j})n_{j+1} \right\rangle + (c_{1} + c_{0}) \left\langle n_{j}n_{j+1} - (1 - n_{j})(1 - n_{j+1}) \right\rangle = (2c_{0} - 1) \left\langle n_{j-1} \right\rangle + 2(c_{2} + c_{1} + c_{0}) \left\langle n_{j} \right\rangle + \left\langle n_{j+1} \right\rangle - (c_{1} + c_{2} + 2c_{0}).$$

$$(3.9)$$

The same calculation for the first and last sites leads to

$$J_1^{ev} = (2c_1 + 2c_0 - 1)\langle n_1 \rangle + \langle n_2 \rangle - (c_0 + c_1), J_L^{ev} = (2c_0 - 1)\langle n_{L-1} \rangle + (2c_2 + 1)\langle n_L \rangle - (c_2 + c_0).$$
(3.10)

Plugging formulas (3.5) in the above expressions, we get

$$J_{1}^{\text{ev}} = -\frac{1}{2}\Lambda \left(2c_{0} + 2c_{1} - 1 + \Gamma\right) ,$$

$$J_{j}^{\text{ev}} = -\frac{1}{2}\Lambda \left(1 - \Gamma\right)^{2}\Gamma^{j-2} ,$$

$$J_{L-1}^{\text{ev}} = -\frac{1}{2}\widehat{\Lambda} - \left(\frac{1 - c_{0}}{2 + 2c_{2} + \delta + \beta} + c_{0} + c_{1} + c_{2}\right)\Lambda\Gamma^{L-2} + \frac{1 - 2c_{0}}{2}\Lambda\Gamma^{L-3} ,$$

$$J_{L}^{\text{ev}} = -\frac{1}{2}(1 + 2c_{2})\widehat{\Lambda} + \frac{(\delta + \beta)(1 - 2c_{0}) - 2(c_{0} + c_{2})}{2(2 + 2c_{2} + \delta + \beta)}\Lambda\Gamma^{L-2} .$$

(3.11)

Using the expressions (3.7) and (3.11), it is easy to check the current conservation on each site of the lattice:

$$J_{j-1,j}^{latt} = J_{j,j+1}^{latt} + J_j^{ev}, \qquad 1 \le j \le L.$$

We plot below (figures 4 and 5) the currents for a system of L = 50 sites: numerical results obtained from a Monte-Carlo simulation coincide with the values obtained from the analytical results.




Figure 4: Lattice current




Figure 5: Evaporation current

3.3 Catalysis vs fracturing processes

We want to compare evaporation of molecules with respect to evaporation of single particles. Through this comparison, we can understand whether the lattice (understood as immersed in a volume, while connected to the two reservoirs) acts as a catalyst in the coagulation (resp. fracturing) reaction of particles (resp. molecules).

The current corresponding to molecule evaporation is given by

$$J_{j,j+1}^{mol} = c_0 \left\langle n_j n_{j+1} - (1 - n_j)(1 - n_{j+1}) \right\rangle = c_0 \left(\left\langle n_j \right\rangle + \left\langle n_{j+1} \right\rangle - 1 \right), \quad 1 \le j \le L - 1.$$
(3.12)

It leads to

$$J_{j,j+1}^{mol} = -\frac{1}{2}c_0\Lambda(1+\Gamma)\Gamma^{j-1}, \quad 1 \le j \le L-2$$

$$J_{L-1,L}^{mol} = -\frac{1}{2}c_0\widehat{\Lambda} - c_0\frac{\delta+\beta+4+2c_0+2c_2}{2(2+2c_2+\delta+\beta)}\Lambda\Gamma^{L-2}$$
(3.13)

When $J_{j-1,j}^{mol}$ is positive, molecules evaporate from the lattice more than they condensate, whereas when $J_{j-1,j}^{mol}$ is negative, molecules mostly condensate onto the lattice. When $\Gamma < 0$, the sign of $J_{j-1,j}^{mol}$ alternates with respect to the parity of j.

To compare the evaporation of molecule with that of single particles, we need to consider $J_{j-1}^{ev} + J_j^{ev}$, which contains both molecules and particles evaporation (on 2 sites), and remove the part coming from molecules. On one site, the current evaporation without molecule evaporation reads:-

$$J_{1}^{\text{part}} = (2c_{1} + c_{0} - 1)\langle n_{1} \rangle + (1 - c_{0})\langle n_{2} \rangle - c_{1},$$

$$J_{j}^{\text{part}} = (c_{1} + c_{2})(2\langle n_{j} \rangle - 1) + (1 - c_{0})(\langle n_{j+1} \rangle - \langle n_{j-1} \rangle), \quad 2 \leq j \leq L - 1,$$

$$J_{L}^{\text{part}} = (1 - c_{0} + 2c_{2})\langle n_{L} \rangle - (1 - c_{0})\langle n_{L-1} \rangle - c_{2}.$$

(3.14)

For $2 \leq j \leq L - 3$, it leads to

$$J_{j,j+1}^{\text{part}} = J_j^{\text{part}} + J_{j+1}^{\text{part}} = \Lambda \Gamma^{j-1} (1+\Gamma) \left(\frac{1-c_0}{2} \left(\frac{1}{\Gamma} - \Gamma \right) - (c_1 + c_2) \right)$$
(3.15)

We show in figure 6 the two parts of the evaporation current.

Remark that the ratio $\mathbf{r}_{j,j+1} = J_{j-1,j}^{\text{part}}/J_{j-1,j}^{\text{mol}}$ is constant all over the lattice except when $j \in \{1, L-2, L-1\}$:

$$\mathfrak{r}_{j,j+1} = \frac{(1-c_0)(\Gamma - \frac{1}{\Gamma}) + 2(c_1 + c_2)}{c_0} = \frac{(c_1 + c_2)^2 - 4c_0}{c_0(1+c_0 + c_1 + c_2)}, \quad 1 < j < L-2.$$
(3.16)

It is also independent of L, despite the total evaporation current J_j^{ev} tends to zero in the bulk when $L \to \infty$.

We define

$$J_{tot}^{ev} = \sum_{j=1}^{L} J_j^{ev}, \qquad J_{tot}^{mol} = \sum_{j=1}^{L-1} J_{j,j+1}^{mol}, \qquad J_{tot}^{part} = \sum_{j=1}^{L} J_j^{part}$$

In the limit $L \to \infty$, the total evaporation current simplifies, which allows us to determine the propensity of the lattice to bound particles into molecules or to split them apart.

$$J_{\infty}^{ev} = \lim_{L \to \infty} J_{tot}^{ev} = -(c_0 + c_1)\Lambda - (1 + c_2)\widehat{\Lambda},$$

$$J_{\infty}^{mol} = \lim_{L \to \infty} J_{tot}^{mol} = -\frac{c_0}{2} \left(\frac{1 + \Gamma}{1 - \Gamma}\Lambda + \widehat{\Lambda}\right),$$

$$J_{\infty}^{part} = \lim_{L \to \infty} J_{tot}^{part} = \left(\frac{2c_0\Gamma}{1 - \Gamma} - c_1\right)\Lambda + (c_0 - 1 - c_2)\widehat{\Lambda}.$$
(3.17)

Now, the quantity

$$2J_{\infty}^{mol} - J_{\infty}^{part} = \left(c_1 - c_0 \frac{1+3\Gamma}{1-\Gamma}\right)\Lambda + (1+c_2 - 2c_0)\widehat{\Lambda}$$

is positive if molecules are created by the lattice, and negative otherwise. The factor 2 before J_{tot}^{mol} is necessary to compare the flow of 2-particles molecules with the flow of single atoms. Remark that this quantity depends on the left boundary, but not on the right one. We show in figure 7 how the local quantity $2J_j^{mol} - J_j^{part}$ evolves with respect to the position of the site.

3.4 Left-right asymmetry

As already noticed, the currents and densities depends heavily on the left boundary parameters, while right boundary has only little impact. This left-right asymmetry may sound surprising since the model is of SSEP-type for its diffusive aspects.

Looking at the densities (3.5), it is clear that the asymmetry is due to the presence of $\Gamma \neq 0$. Indeed, when $c_0 = 1$, the symmetry between the left and right boundary is restored, see section 5. Then, this asymmetry may be linked to the fact that the transposition of local Markov matrix (2.1) is still a Markov matrix, but of a different kind. Indeed, when $c_0 = 1$, the Markov matrix is self-transposed. This suggests that the asymmetry may be induced by the evaporation/condensation processes, but a physical explanation is lacking at this point.




Figure 6: Evaporation currents (for molecules and particules separately)




Figure 7: Comparison of the molecule and particle flows

4 *n*-point correlation functions

The stationarity of the *n*-point correlation function leads to a recursion relation involving ℓ -point correlation function with $\ell \leq n$ only. In other words, we have a decoupling of the *n*-point correlation functions from the higher ones. Thus, they can be computed exactly. We illustrate it for n = 2 in section 4.1, and prove the decoupling in the general case in section 4.2. Let us stress that although we get an analytical expression for the correlation functions, they differ from the mean field: we illustrate it in section 6 through numerical calculations.

To simplify the presentation, we use the notations (3.1) and

$$\Theta = \frac{1 - c_0}{1 + c_0 + 2c_1 + 2c_2}, \qquad \Xi = \frac{1 - c_0}{1 + c_0 + c_1 + c_2 + \frac{\alpha + \gamma + \beta + \delta}{2}}, \qquad (4.1)$$

$$\Upsilon = \frac{1 - c_0}{1 + 2c_0 + 2c_1 + c_2 + \frac{\alpha + \gamma}{2}}, \qquad \Delta = \frac{1 - c_0}{2 + c_0 + c_1 + 2c_2 + \frac{\beta + \delta}{2}}.$$

4.1 Two-point correlation function

Considering the stationarity of the two-point correlation functions, we get the following recursion relations for 1 < i < j - 1 < L - 1:

$$\langle n_i n_j \rangle - \frac{\Gamma}{2} \Big(\langle n_{i-1} n_j \rangle + \langle n_i n_{j-1} \rangle \Big) = \frac{1 - \Gamma}{4} \Big(\langle n_i \rangle + \langle n_j \rangle \Big).$$
 (4.2)

For 1 < i < L - 1, we have

$$\langle n_i n_{i+1} \rangle - \Theta \langle n_{i-1} n_{i+1} \rangle = \frac{(-1 + c_0 + c_1 + c_2) \langle n_{i+1} \rangle + (1 - c_0 + c_1 + c_2) \langle n_i \rangle + c_0}{2 + 2c_0 + 4c_1 + 4c_2}, \quad (4.3)$$

$$\langle n_1 n_{i+1} \rangle - \Upsilon \langle n_1 n_i \rangle = \frac{(2c_0 + c_1 + c_2)\langle n_1 \rangle + (\alpha + c_0 + c_1)\langle n_{i+1} \rangle}{\alpha + \gamma + 2 + 4c_0 + 4c_1 + 2c_2}, \qquad (4.4)$$

$$\langle n_{i}n_{L}\rangle - \Delta\left(\langle n_{i}n_{L-1}\rangle + \langle n_{i-1}n_{L}\rangle\right) = \frac{(\delta + c_{0} + c_{2})\langle n_{i}\rangle + (2c_{0} + c_{1} + c_{2})\langle n_{L}\rangle}{\beta + \delta + 4 + 2c_{0} + 2c_{1} + 4c_{2}}, \quad (4.5)$$

together with the boundary cases

$$\langle n_1 n_2 \rangle = \frac{(1 - c_0 + c_1 + c_2) \langle n_1 \rangle + (\alpha - 1 + c_1) \langle n_2 \rangle + c_0}{\alpha + \gamma + 2c_0 + 4c_1 + 2c_2}, \qquad (4.6)$$

$$\langle n_1 n_L \rangle - \Xi \langle n_1 n_{L-1} \rangle = \frac{(\delta + c_0 + c_2) \langle n_1 \rangle + (\alpha + c_0 + c_1) \langle n_L \rangle}{\alpha + \gamma + \beta + \delta + 2 + 2c_0 + 2c_1 + 2c_2}, \qquad (4.7)$$

$$\langle n_{L-1}n_L \rangle - \frac{2(1-c_0)\langle n_{L-2}n_L \rangle}{\beta+\delta+2+2c_1+4c_2} = \frac{(-1+c_0+c_1+c_2)\langle n_L \rangle + (\delta+1-2c_0+c_2)\langle n_{L-1} \rangle + c_0}{\beta+\delta+2+2c_1+4c_2}.$$
(4.8)

Notice that the correlation functions $\langle n_i n_j \rangle$ with j < L decouple from the correlation functions $\langle n_i n_L \rangle$, $\forall i$. Then, we can solve the former independently from the later. It leads to the following expressions for the two-point correlation functions.

$$\langle n_{i}n_{j} \rangle = \frac{1}{4} \left(1 - (\Gamma^{i-1} + \Gamma^{j-1})\Lambda + \Gamma^{i+j-2} \left(\frac{j-i-1}{\Theta} - \frac{j-i-2}{\Gamma} \right) \omega \right) + \frac{1}{4} \sum_{k=3}^{j} \left(\frac{\Gamma}{2} \right)^{i+j+1-k} a_{k} \left\{ \binom{i+j-k-1}{i-1} - \binom{i+j-k-1}{j-2} + \sum_{s=1}^{i+2-k} \left[\binom{i+j-k-1-s}{j-3} - \binom{i+j-k-1-s}{j-2} \right] \left(\frac{2\Theta}{\Gamma} \right)^{s} \right\}$$

$$(4.9)$$

for $1 \leq i < j \leq L - 1$, where

$$a_{k} = \left(\lambda_{1} k + \lambda_{0}\right) \Gamma^{k-3} - \frac{2 + 2c_{2} - \alpha - \gamma}{2(1 - c_{0})} (\lambda_{0} + \lambda_{1}) \Upsilon^{k-2}, \qquad 3 \le k \le L - 1, \qquad (4.10)$$

with

$$\lambda_{1} = -2 \frac{c_{1} + c_{2}}{1 - c_{0}} \omega , \qquad \lambda_{0} = -2 \frac{1 + c_{0}}{1 - c_{0}} \omega + 2 \frac{(\alpha - \gamma)^{2}}{(2c_{0} + 2c_{1} + \alpha + \gamma)^{2}},$$

$$\omega = 4 \langle n_{1}n_{2} \rangle - \frac{2(c_{0} + c_{1})(1 + c_{0} + c_{1} + c_{2}) + (\alpha + \gamma)c_{0} + 2\alpha(c_{1} + c_{2}) + 3\alpha - \gamma}{(1 + c_{0} + c_{1} + c_{2})(2c_{0} + 2c_{1} + \alpha + \gamma)}.$$
(4.11)

The expression (4.9) is obtained as follows. The first line is a particular solution of the recurrence relations (4.2) and (4.3), whatever the value of ω . The two last lines, whatever the value of the coefficients a_k , solve them with vanishing r.h.s. The coefficients a_k are fixed in such a way that the last relation (4.4) is also obeyed and ω is given by the initial value (4.6). This allows to get all two-point functions $\langle n_i n_j \rangle$ with $1 \leq i < j < L$.

The remaining two-point functions are more involved. We get

$$\langle n_1 n_L \rangle = \frac{\alpha + c_0 + c_1}{(2c_0 + 2c_1 + \alpha + \gamma)(2 + 2c_2 + \beta + \delta)} \left(1 + c_2 + \delta - (1 - c_0) \Lambda \Gamma^{L-2} \right) - \frac{1}{8} (\lambda_0 + \lambda_1) \Gamma \Xi \Upsilon^{L-3},$$
(4.12)

and for $2 \le i \le L - 2$,

$$\langle n_i n_L \rangle = \Delta^{i-1} \langle n_1 n_L \rangle + \sum_{k=1}^{i-1} \Delta^{i-k} \langle n_{k+1} n_{L-1} \rangle - \frac{1}{4} \frac{\delta + c_0 + c_2}{1 - c_0} \Lambda \frac{\Delta \Gamma^i - \Gamma \Delta^i}{\Gamma - \Delta} + \frac{\Delta - \Delta^i}{\Delta - 1} \left(\frac{\delta + c_0 + c_2}{4(1 - c_0)} + \frac{2c_0 + c_1 + c_2}{2(1 - c_0)} (1 - \widehat{\Lambda}) - \frac{1 - c_0}{\delta + \beta + 2 + 2c_2} \Lambda \Gamma^{L-2} \right).$$

$$(4.13)$$

Finally $\langle n_{L-1}n_L \rangle$ is given by formula (4.8).

4.2 Hierarchical relations for *n*-point correlation function

As for the two-point correlation functions, the n-point correlation functions decouple from the higher ones, and thus can be exactly computed recursively. To show it, we introduce the following row vectors

$$\mathbf{u} = (1,1), \quad \mathbf{f} = (0,1), \quad \mathbf{v}_n(i_1,\ldots,i_n) = \mathbf{u} \otimes \cdots \otimes \mathbf{u} \otimes \underbrace{\mathbf{f}}_{i_1} \otimes \mathbf{u} \ldots \mathbf{u} \otimes \underbrace{\mathbf{f}}_{i_n} \otimes \mathbf{u} \ldots \mathbf{u} \quad (4.14)$$

so that the *n*-point correlation functions read $\langle n_{i_1} \dots n_{i_n} \rangle = \mathbf{v}_n(i_1, \dots, i_n) | \mathcal{S} \rangle$.

The time evolution of this correlation function is ruled by the action of the Markov matrix \mathbb{M} on $\mathbf{v}_n(i_1,\ldots,i_n)$. Thus, the recursion relations for the correlation functions are obtained from the identities

$$\begin{aligned} \mathbf{u} \otimes \mathbf{u} M &= 0 \quad ; \quad \mathbf{u} B = 0 \quad ; \quad \mathbf{u} \overline{B} = 0 \\ \mathbf{u} \otimes \mathbf{f} M &= (c_0 + c_2) \mathbf{u} \otimes \mathbf{u} - 2(1 + c_2) \mathbf{u} \otimes \mathbf{f} + 2(1 - c_0) \mathbf{f} \otimes \mathbf{u} \\ \mathbf{f} \otimes \mathbf{u} M &= (c_0 + c_1) \left(\mathbf{u} \otimes \mathbf{u} - 2 \mathbf{f} \otimes \mathbf{u} \right) \\ \mathbf{f} \otimes \mathbf{f} M &= c_0 \mathbf{u} \otimes \mathbf{u} + (c_1 - 1) \mathbf{u} \otimes \mathbf{f} + (1 - 2c_0 + c_2) \mathbf{f} \otimes \mathbf{u} - 2(c_1 + c_2) \mathbf{f} \otimes \mathbf{f} \\ \mathbf{f} B &= \alpha \mathbf{u} - (\alpha + \gamma) \mathbf{f} \\ \mathbf{f} \overline{B} &= \delta \mathbf{u} - (\delta + \beta) \mathbf{f} \end{aligned}$$
(4.15)

Since $\mathbf{f} \otimes \mathbf{f}$ occurs only in the action of M on $\mathbf{f} \otimes \mathbf{f}$ (and because $\mathbf{u} \otimes \mathbf{u} M = 0$) this proves that the recursion relation for the *n*-point functions involves only correlations of *n*-point at most.

4.3 *n*-point correlations recursion relations

The aim of this section is to compute explicitly the recursion relations obeyed by the *n*-point correlation fonctions. Let $T = \{i_1 \cdots i_n\}$ the set of sites where we want to calculate the correlation, i.e.

$$\langle n_T \rangle = \langle n_{i_1} \cdots n_{i_n} \rangle. \tag{4.16}$$

We decompose T as a union of disjoint segments:

$$T = \bigcup_{a=1}^{N} [i_a, j_a] \quad \text{with} \quad j_{a-1} + 1 < i_a \le j_a < i_{a+1} - 1, \ a = 1..., N,$$
(4.17)

where by convention $j_0 = -1$ and $i_{N+1} = L + 2$.

The recursion relations are obtained through the action of the Markov matrix

$$0 = \mathbb{M} \langle n_T \rangle = \sum_{a=1}^N M_{[i_a, j_a]} \langle n_T \rangle + \delta_{i_1, 1} B_1 \langle n_T \rangle + \delta_{j_N, L} \bar{B}_L \langle n_T \rangle, \qquad (4.18)$$

where $M_{[i_a, j_a]} = \bigoplus_{k=i_a}^{j_a-1} M_{k,k+1}$. It is clear that there is a decoupling of the different segments $[i_a, j_a]$ and that the contributions of each segment will add up with no interference. Then, it

is sufficient to consider the sets T = [i, j] which appear as building blocks for the calculation of the recursion relations.

We obtain, for T = [i, j] with $1 \le i \le j \le L$:

$$\begin{pmatrix} 2(j+1-i)(c_{1}+c_{2})+2(1+c_{0})+\delta_{i,1}\left(\alpha+\gamma-2-2c_{2}\right)+\delta_{j,L}\left(\beta+\delta-2c_{0}-2c_{1}\right)\right)\langle n_{[i,j]}\rangle \\ -2(1-c_{0})(1-\delta_{i,1})\langle n_{i-1}n_{[i+1,j]}\rangle \\ = \\ \begin{pmatrix} c_{0}+c_{2}+(1-\delta_{i,j})\left(c_{1}-1\right)+\delta_{i,1}\left(\alpha-c_{0}-c_{2}\right)\right)\langle n_{[i+1,j]}\rangle \\ +\left(c_{0}+c_{1}+(1-\delta_{i,j})\left(1-2c_{0}+c_{2}\right)+\delta_{j,L}\left(\delta-c_{1}-c_{0}\right)\right)\langle n_{[i,j-1]}\rangle \\ +c_{0}\sum_{k=i}^{j-1}\langle n_{[i,k-1]}n_{[k+2,j]}\rangle +\left(c_{1}+c_{2}-2c_{0}\right)\sum_{k=i+1}^{j-1}\langle n_{[i,k-1]}n_{[k+1,j]}\rangle,$$

$$(4.19)$$

(4.19) where we have used the conventions that $n_{[i,i]} = n_i$, $n_{[k,\ell]} = 1$ when $k > \ell$, and $\sum_{k=a}^{b} (\cdots) = 0$ if a > b.

In (4.19), the left-hand-side corresponds to (j + 1 - i)-point correlation functions, while the right-hand-side involves only *m*-point correlation functions with $m \leq j - i$. Since for i = 1, the l.h.s. contains only the correlation $\langle n_{[1,j]} \rangle$, it can be solved by recursion hypothesis. Then, the correlation $\langle n_{[2,j+1]} \rangle$ can be computed once one knows $\langle n_{[1,j]} \rangle$ and the correlations with a lower number of points, and so on. It is the left-right asymmetry of the model that allows the calculation through this (double) recursion.

When $T = \bigcup_{a=1}^{N} [i_a, j_a]$, the calculation is more involved, but follows also a recursion: one first needs to compute all recursions $\langle n_{[1,k]} \rangle$ with $k \leq |T|$ following the method given above. Then, similarly, one can compute $\langle n_{[1,k]}n_{\ell} \rangle$ starting from $\ell = k+2$, and so on.

5 The special case $c_0 = 1$

As already mentioned, the comparison with the DISSEP model is obtained for $c_0 = 1$ and $c_1 = c_2 = 0$ and corresponds to the free fermions point. Here, we consider a more general case by setting $c_0 = 1$ but keeping c_1 and c_2 free. Then we have $\Gamma = 0$ and expressions simplify. The bulk Markov matrix reads

$$M\big|_{c_0=1} = \begin{pmatrix} -1 - c_1 - c_1 & c_2 & c_1 & 1\\ c_2 & -1 - c_1 - c_2 & 1 & c_1\\ c_1 & 1 & -1 - c_1 - c_2 & c_2\\ 1 & c_1 & c_2 & -1 - c_1 - c_2 \end{pmatrix}.$$
 (5.1)

The densities take the form

$$\langle n_0 \rangle = \frac{\alpha + (1 + \alpha - \gamma)(1 + c_1)}{2 + 2c_1 + \alpha + \gamma} \quad ; \quad \langle n_1 \rangle = \frac{\alpha + 1 + c_1}{2 + 2c_1 + \alpha + \gamma};$$

$$\langle n_j \rangle = \frac{1}{2}, \quad 2 \le j \le L - 1;$$

$$\langle n_L \rangle = \frac{\delta + 1 + c_2}{2 + 2c_2 + \beta + \delta} \quad ; \quad \langle n_{L+1} \rangle = \frac{\delta + (1 + \delta - \beta)(1 + c_2)}{2 + 2c_2 + \delta + \beta}.$$
 (5.2)

For the current in the lattice, we have

$$J_{01}^{latt} = \frac{(\alpha - \gamma)(1 + c_1)}{2 + 2c_1 + \alpha + \gamma} \qquad ; \qquad J_{12}^{latt} = \frac{\alpha - \gamma}{2(2 + 2c_1 + \alpha + \gamma)} ;$$

$$J_{j,j+1}^{latt} = 0 , \quad 2 \le j \le L - 2 ; \qquad (5.3)$$

$$J_{L-1,L}^{latt} = \frac{\beta - \delta}{2(2 + 2c_2 + \beta + \delta)} \qquad ; \qquad J_{L,L+1}^{latt} = \frac{(\beta - \delta)(1 + c_2)}{2 + 2c_2 + \beta + \delta} .$$

Finally, the evaporation current reads

$$J_{1}^{ev} = \frac{(\alpha - \gamma)(1 + 2c_{1})}{2(2 + 2c_{1} + \alpha + \gamma)} \quad ; \qquad J_{2}^{ev} = \frac{\alpha - \gamma}{2(2 + 2c_{1} + \alpha + \gamma)} ;$$

$$J_{j}^{ev} = 0 , \quad 3 \le j \le L - 2 ; \qquad (5.4)$$

$$J_{L-1}^{ev} = \frac{\delta - \beta}{2(2 + 2c_{2} + \beta + \delta)} \quad ; \qquad J_{L}^{ev} = \frac{(\delta - \beta)(1 + 2c_{2})}{2(2 + 2c_{2} + \beta + \delta)} .$$

When $c_1 = c_2 = 0$, these values agree with the ones computed in [5].

Remark that we also have

$$J_{j}^{part} = -2(c_{1} + c_{2})(\langle n_{j} \rangle + \langle n_{j+1} \rangle - 1) = 0 \quad \text{and} \quad J_{j,j+1}^{mol} = \langle n_{j} \rangle + \langle n_{j+1} \rangle - 1 = 0, \quad (5.5)$$

which shows that in the limit $c_0 \rightarrow 1$, there is an opposite flux of condensation/evaporation for particules and for molecules, the ratio being constant and equal to $-2(c_1+c_2)$. The competition between the two processes is parametrized by $c_1 + c_2$, which can be adjusted to favor one or the other.

In fact, when $c_0 = 1$, the bulk Markov matrix is diagonalizable:

$$(U \otimes U) M \big|_{c_0 = 1} (U^{-1} \otimes U^{-1}) = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & -2(1 + c_2) & 0 & 0 \\ 0 & 0 & -2(1 + c_1) & 0 \\ 0 & 0 & 0 & -2(c_1 + c_2) \end{pmatrix},$$
(5.6)

where

$$U = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}.$$

Performing this transformation for the full Markov matrix (2.5)

$$\mathbb{M}\big|_{c_0=1} \to U \otimes U \otimes \cdots \otimes U \mathbb{M}\big|_{c_0=1} U^{-1} \otimes U^{-1} \otimes \cdots \otimes U^{-1},$$

it is easy to see that for the transformed Markov matrix, the eigenstate with zero eigenvalue reads

$$|0\rangle = \begin{pmatrix} 1\\\Lambda \end{pmatrix} \otimes \begin{pmatrix} 1\\0 \end{pmatrix} \otimes \cdots \otimes \begin{pmatrix} 1\\0 \end{pmatrix} \otimes \begin{pmatrix} 1\\\widehat{\Lambda} \end{pmatrix}, \qquad (5.7)$$

where Λ and $\widehat{\Lambda}$ are given in (3.1) (with $c_0 = 1$). This allows us to compute the stationary state of the Markov matrix \mathbb{M} :

$$|\mathcal{S}\rangle = U^{-1} \otimes U^{-1} \otimes \cdots \otimes U^{-1} |0\rangle = \begin{pmatrix} 1+\Lambda\\ 1-\Lambda \end{pmatrix} \otimes \begin{pmatrix} 1\\ 1 \end{pmatrix} \otimes \cdots \otimes \begin{pmatrix} 1\\ 1 \end{pmatrix} \otimes \begin{pmatrix} 1+\widehat{\Lambda}\\ 1-\widehat{\Lambda} \end{pmatrix}.$$
 (5.8)

Remark that the symmetry between the left and the right boundaries has been re-established.

Then, the calculation of the correlation functions becomes easy, and one can show that we get the results given above. For the two-point functions, one can check directly that the mean field approximation becomes exact at $c_0 = 1$:

$$\langle n_i n_j \rangle = \langle n_i \rangle \langle n_j \rangle, \quad 1 \le i \ne j \le L.$$

As we show in section 6 below, it is not true for generic values of c_0 .

6 Numerical calculations

To confirm our analysis, we have performed numerical calculations from two different methods. The first method consists in solving numerically the linear system which must satisfy the stationary state, *i.e.* finding the normalized vector spanning the kernel of the Markov matrix. We used this method to get really accurate results up to L = 14 sites. For larger values of L, we used a Monte-Carlo approach.

We have computed numerically (figures 8, 9, 10, and 11) the connected two-point correlation $\langle n_j n_k \rangle_c = \langle n_j n_k \rangle - \langle n_j \rangle \langle n_k \rangle$ up to L = 14 sites by solving numerically the linear system (first method), and compared it against our analytical results.




Figure 8: Connected 2-point correlation $\langle n_i n_{i+1} \rangle_c$




Figure 9: Connected 2-point correlation $\langle n_i n_{i+2} \rangle_c$




Figure 10: Connected 2-point correlation $\langle n_1 n_i \rangle_c$




Figure 11: Connected 2-points correlation $\langle n_i n_L \rangle_c$

7 Conclusion

The model we have studied possesses several features that are worth mentioning. The left/right asymmetry, in particular, is a striking point. Although it is clear that this asymmetry roots in the evaporation/condensation part of the model, a deeper understanding of this property needs to be developed. Although the thermodynamical limit $L \to \infty$ seems to be trivial (most densities reach $\frac{1}{2}$ and most of the currents are zero), the model being symmetric in its diffusive part, it is tempting to look for a WASEP-like model in the thermodynamical limit. We hope to come back on this point in the next future. A generalization of the model to the case of asymmetric diffusion rates is also a line that we wish to develop.

We have already mentioned that such hierarchical relations were studied also in the literature, see e.g. [8, 10, 15], but the models where based on fermions, and/or dealt with discrete time or periodic cases. The present model does not fall in these classes.

As already mentioned, the model can be solved at the level of correlation functions, but there is no known integrable structure. Nevertheless, since all correlation functions can be computed analytically, a more detailed search of its possible integrability is worth doing. We have checked that for generic values of the parameters, the matrix M does not obey a braid relation. This prevents it from being part of a Hecke, a Temperley-Lieb or a BMW algebra that would lead to a baxterisation procedure [11, 13, 14]. However, there are several types of new algebras that can be used for a baxterisation, see for instance the new algebras introduced in [3, 4]. It would be interesting to look for a four dimensional representation of them that could lead to the Markov matrix of the present paper.

The search of a matrix ansatz is also a direction that one could look at. It seems however that the associated algebra, if it exists, is not a 'diffusion algebra', as they were studied in [9,12]. Obviously, if the model appears to be integrable, a search for a matrix ansatz would be greatly simplified, following the lines given in [2].

Finally, the fracturing / catalysis approach is also a point that needs to be explored. A qualitative comparison with a true catalysis reaction in one dimension would be a first step to see whether the model can be used as a toy model for such processes.

Acknowledgements

We are grateful to E. Bertin, N. Crampé and V. Lecomte for fruitful discussions and comments during the course of this work.

References

 R.A. Blythe and M.R. Evans, Nonequilibrium Steady States of Matrix Product Form: A Solver's Guide, J. Phys. A40 (2007) R333-R441 and arXiv:0706.1678.

- [2] N. Crampe, E. Ragoucy and M. Vanicat, Integrable approach to simple exclusion processes with boundaries. Review and progress, J. Stat. Mech. (2014) P11032 and arXiv:1408.5357.
- [3] N. Crampe, L. Frappat, E. Ragoucy, M. Vanicat, A new braid-like algebra for Baxterisation, Comm. Math. Phys. 349 (2017) 271–283 and arXiv:1509.05516.
- [4] N. Crampe, E. Ragoucy, M. Vanicat, Back to Baxterisation, Comm. Math. Phys. 365 (2019) 1079–1090 and arXiv:1708.02754.
- [5] N. Crampe, E. Ragoucy, V. Rittenberg and M. Vanicat, Integrable dissipative exclusion process: Correlation functions and physical properties, Phys. Rev. E94 (2016) 032102 and arXiv:1603.06796.
- [6] B. Derrida, M. R Evans, V. Hakim, and V. Pasquier, Exact solution of a 1d asymmetric exclusion model using a matrix formulation, J. Phys. A26 (1993) 1493.
- [7] B. Derrida, Non-equilibrium steady states: fluctuations and large deviations of the density and of the current, JSTAT (2007) 07023.
- [8] V. Eisler, Crossover between ballistic and diffusive transport: The Quantum Exclusion Process, arXiv:1104.4050.
- [9] F. Essler, V. Rittenberg, Representations of the quadratic Algebra and Partially Asymmetric Diffusion with Open Boundaries, cond-mat/950611.
- [10] C. Giardina, J. Kurchan, F. Redig, Duality and exact correlations for a model of heat conduction, J. Math. Phys. 48(2007) 033301.
- [11] A.P. Isaev, Quantum groups and Yang-Baxter equations, Max-Planck Institut f
 ür Mathematik (2004).
- [12] A.P. Isaev, P.N. Pyatov and V.Rittenberg, Diffusion algebras, J. Phys. A34 (2001) 5815 and arXiv:cond-mat/0103603.
- [13] M. Jimbo, A q-difference analogue of U (gl (n + 1)), Hecke algebra and the Yang-Baxter equation, Lett. Math. Phys. **11** (1986) 247.
- [14] V.F.R. Jones, Baxterisation, Int. J. Mod. Phys. B 4 (1990) 701. In: Proceedings of "Yang-Baxter equations, conformal invariance and integrability in statistical mechanics and field theory", Canberra (1989).
- [15] I. Klich, Closed hierarchies and non-equilibrium steady states of driven systems, Annals of Physics 404 (2019) 66-80 and arXiv:1508.05957.

- [16] K. Krebs and S. Sandow, Matrix Product Eigenstates for One-Dimensional Stochastic Models and Quantum Spin Chains, J. Phys. A30 (1997) 3165 and arXiv:cond-mat/9610029.
- [17] S. Sandow, Partially Asymmetric Exclusion Process with Open Boundaries, Phys. Rev. E50 (1994) 2660 and arXiv:cond-mat/9405073.
- [18] T. Sasamoto, S. Mori, M. Wadati, One-dimensional asymmetric exclusion model with open boundaries, J. Phys. Soc. Japan 65 (1996) 2000.
- [19] C. A Tracy and H. Widom, Asymptotics in ASEP with step initial condition, Comm. Math. Phys. 290 (2009) 129–154 and arXiv:0807.1713.