

Proof of Weierstrass gap theorem

Gollakota V V Hemasundar

Department of Mathematics,

SIWS College, Mumbai-400031, INDIA

E-mail: gvvhemasundar@yahoo.co.in

Abstract

In this expository note we give proof of the Weierstrass gap theorem in Cohomology terminology . We analyze gap sequence for finding possible gaps and non-gaps on X .

Keywords: Weierstrass gap theorem, Meromorphic functions, Compact Riemann surfaces

AMS: 30F10

1 Introduction:

Let X be a compact Riemann surface of genus g . One of the important questions is the existence of meromorphic functions having pole at a single point p on X . By the Riemann Roch theorem, we can show that there always exists a non-constant meromorphic function $f \in \mathcal{M}(X)$ which has a pole of order $\leq g + 1$ at p and is holomorphic in $X \setminus \{p\}$.

One of the basic results in this topic is Weierstrass gap theorem, which states that

Theorem 1 *For a surface of genus $g \geq 1$ there are precisely g integers*

$$(1) \quad 1 = n_1 < n_2 < \cdots < n_g < 2g$$

such that there does not exist a meromorphic function on X with a pole of order n_k at p .

The numbers n_j , for $j = 1, \dots, g$ are called “gaps” at p and their complement in \mathbb{N} are called “non-gaps”. Further, the sequence is uniquely determined by the point p .

The case $g = 0$ is trivial as there is always a function on the sphere with a simple pole. The case $g = 1$ is clear since there is no meromorphic function with a single simple pole.

In this note we furnish the proof of Theorem 1 in the language of sheaf cohomology, in the spirit of proof given in Springer [4]. The proof of Noether's theorem was given in Farkas and Kra [1] from which the proof of Weierstrass gap theorem was deduced as a special case. The proofs are generally given as an application of Riemann-Roch Theorem. In section 2, we mention some of the consequences of Riemann-Roch and Serre Duality theorems. The proofs may be found in [2]. The proof of Weierstrass gap theorem is given in Section 3. In Section 4, we form a combinatorial problem which appears to be a by product from the statement of Weierstrass gap theorem and analyze gaps and non-gaps.

2 Some Consequences of Riemann-Roch and Serre Duality Theorems

Let D be a divisor on X .

If $\deg D < 0$ then $H^0(X, \mathcal{O}_D) = 0$.

We can use Serre-Duality theorem to obtain equality of dimensions:

$$(2) \quad \dim H^0(X, \Omega_{-D}) = \dim H^1(X, \mathcal{O}_D)$$

For $D = 0$ we obtain

$$(3) \quad \dim H^0(X, \Omega) = g = \dim H^1(X, \mathcal{O})$$

Here $H^0(X, \Omega) = \Omega(X)$ which denotes the sheaf of holomorphic 1-forms on X .

The following equality can be obtained as an application of Serre Duality Theorem:

$$\dim H^1(X, \Omega) = \dim H^0(X, \mathcal{O}) = 1$$

Note 2 *This is a known fact again saying there are no non-constant holomorphic maps on a compact Riemann surface.*

Suppose X is a compact Riemann surface of genus g . Let K be the canonical divisor on X . Then $\deg K = 2g - 2$.

3 Proof of Theorem 1 (Weierstrass Gap Theorem)

Proof. Suppose $P \in X$. If D is a zero divisor, then $\dim H^1(X, \mathcal{O}) = g$ and $\deg(D) = 0$.

By the Riemann Roch theorem $\dim H^0(X, \mathcal{O}) = 1$. Therefore, there are no non-constant holomorphic function on X .

Define the divisor D_P such that

$$D_P(P) = \begin{cases} 1 & \text{if } x = P \\ 0 & \text{if } x \neq P \end{cases}$$

$\deg(D_P) = 1$. Once again by the Riemann -Roch theorem,

$$\dim H^0(X, \mathcal{O}_{D_P}) = 2 - g + \dim H^0(X, \Omega_{-D_P})$$

If $\dim H^0(X, \Omega_{-D_P}) = g$, then $\dim H^0(X, \mathcal{O}_{D_P}) = 2$, hence there exists $f \in \mathcal{M}(X)$ which has a simple pole at P and is holomorphic in $X \setminus \{P\}$.

If $\dim H^0(X, \Omega_{-D_P}) = g - 1$, then $\dim H^0(X, \mathcal{O}_{D_P}) = 1$, hence there is no meromorphic function which has a simple pole at P and is holomorphic in $X \setminus \{P\}$.

Now we want to see the effect of changing $D_P = (n-1)D_P$ to $D_P = nD_P$. By the Riemann - Roch Theorem

$$\dim H^0(X, \mathcal{O}_{-D_{(n-1)P}}) = n - g + \dim H^0(X, \Omega_{-D_{(n-1)P}})$$

and

$$\dim H^0(X, \mathcal{O}_{-D_{nP}}) = n + 1 - g + \dim H^0(X, \Omega_{-D_{nP}})$$

If

$$\dim H^0(X, \Omega_{-D_{(n-1)P}}) = \dim H^0(X, \Omega_{-D_{nP}})$$

then

$$\dim H^0(X, \mathcal{O}_{-D_{nP}}) = \dim H^0(X, \mathcal{O}_{-D_{(n-1)P}}) + 1.$$

So there exists a meromorphic function $f \in \mathcal{M}(X)$ with a pole of order n at P and is holomorphic in $X \setminus \{P\}$.

If

$$\dim H^0(X, \Omega_{-D_{(n-1)P}}) = \dim H^0(X, \Omega_{-D_{nP}}) - 1$$

then

$$\dim H^0(X, \mathcal{O}_{-D_{nP}}) = \dim H^0(X, \mathcal{O}_{-D_{(n-1)P}}).$$

So there will not exist a function with a pole of order n at P and holomorphic in $X \setminus \{P\}$.

So if $\dim H^0(X, \Omega_{-D_{nP}})$ remains the same as n increases by 1, a new linearly independent function is added in going from the sheaf $\mathcal{O}_{D_{nP}}$ to $\mathcal{O}_{D_{(n+1)P}}$.

From the Eq. 3 we have $\dim H^0(X, \Omega) = g$. We have already seen that if D is the divisor of a non-vanishing meromorphic 1-form on a compact Riemann surface of genus g , then $\deg(\omega) = 2g - 2$.

Let K be its canonical divisor. Then we can deduce that

$$\dim H^0(X, \Omega_{-D_{(2g-1)P}}) = 0$$

Therefore, the number of times $\dim H^0(X, \Omega_{-D_{nP}})$ does not remain the same must be g times and at each change it decreases by 1.

It completes the proof. ■

4 Analyzing gaps and non-gaps

Suppose $p \in X$. If f has a pole of order s at p , and g has a pole of order t at p , then fg has a pole of order $s + t$ at p . Therefore, the set of non-gaps forms an additive sub-semi group of \mathbb{N} .

Let d be the least non-gap value at the point p , if $n > d$ is a gap then $n - d$ is again a gap value. Therefore, all the gaps occur in finite arithmetical sequences of the form $j, j + d, j + 2d, \dots, j + \lambda_j d$, where $j = 1, 2, \dots, d - 1$ and $\lambda_j = 0, 1, 2, \dots$. See pp. 124 [3]. A point p is called a hyperelliptic Weierstrass point if the non-gap sequence starts with 2 and the hyperelliptic Riemann surfaces are characterized by the gap sequence:

$$(4) \quad P = \{1, 3, \dots, 2g - 1\}$$

hence the non-gaps are $Q = \{2, 4, \dots, 2g\}$

For the Exceptional Riemann surfaces the gap sequence P and the non-gap sequence Q are given by

$$(5) \quad P = \{1, 2, 3, \dots, g - 1, g + 1\} \text{ and } Q = \{g, g + 2, \dots, 2g - 1\}.$$

For example, if $g = 3$ we see the possible gap sequences are

$\{1, 3, 5\}$, $\{1, 2, 3\}$, $\{1, 2, 4\}$, $\{1, 2, 5\}$

and corresponding non-gaps are:

$\{2, 4, 6\}$, $\{4, 5, 6\}$, $\{3, 5, 6\}$, $\{3, 4, 6\}$.

The following problem may be of some interest to see possible gaps and non-gaps at a given point.

Question 1:

Write the numbers 2 to $2g-1$ in to two (disjoint) parts $P = \{n_1, n_2, \dots, n_{g-1}\}$ and $Q = \{m_1, m_2, \dots, m_{g-1}\}$ such that no number in P is a sum of any combination of numbers in Q . How many pairs of such P and Q exist?

Clearly then $\{1\} \cup P$ gives possible gaps and $Q \cup \{2g\}$ gives possible non-gaps.

References

- [1] H.M. Farkas and I. Kra, Riemann surfaces, GTM, Springer, Second Edition, 1991.
- [2] O. Forster, Lectures on Riemann surfaces, GTM 81, Springer-Verlag, New York, 1993.
- [3] R.C. Gunning, Lectures on Riemann surfaces, Princeton University Press, second printing 1968.
- [4] George Springer, Introduction to Riemann surfaces, Addison-Wesley Publishing Company, Inc, 1957.