

GOODNESS IN n -ANGULATED CATEGORIES

Sebastian H. Martensen

Abstract

We generalise the notions of good, middling good, and Verdier good morphisms of distinguished triangles in triangulated categories, first introduced by Neeman, to the setting of n -angulated categories, introduced in Geiss, Keller, and Oppermann. We then prove that all morphisms of n -angles in an $(n - 2)$ -cluster tilting n -angulated category are middling good for $n > 3$, and we provide an example of a morphism in exotic n -angulated categories which is not middling good.

1 Introduction

Triangulated categories were introduced independently by Puppe and Verdier in 1962 and 1963 and, today, they turn up in the settings of representation theory, algebraic geometry, and algebraic topology to name a few. However, even in a model-free setting, the triangulation of a category is a rich structure. The Adams spectral sequence, which is classically defined over the stable homotopy category and calculates stable homotopy groups from mod p cohomology operations, extends to general triangulated categories that are equipped with a *projective* or *injective class* (see [5]). This more general Adams spectral sequence is dependent on the chosen class and, given an appropriate class, it will for example specialise to the Adams–Novikov spectral sequence in the category of spectra and, in the derived category of a differential graded algebra, it can specialise to a spectral sequence that converges to sets of morphisms.

Christensen and Frankland found (see [7]) that, by choosing suitable fill-in morphisms between quotients in Adams towers, one might be able to generalize the Moss pairing to the general Adams spectral sequence in triangulated categories. Of interest in this context are fill-ins which are both *good* and *Verdier good* (see Definitions 3.1 and 3.4 below).

In short, a morphism of distinguished triangle is called *good* if its mapping cone (see Definition 2.7 below) is a distinguished triangle, it is called *Verdier good* if the third morphism between objects factors through two specific octahedra, and furthermore we call a morphism *middling good* if it extends to a (4×4) -diagram wherein the first three rows and columns are all distinguished triangles. Starting from a commutative square, it turns out that all three conditions are almost entirely dependent on the choice of fill-in map between the cone objects. In [6] and [10], certain relationships between these concepts are found and certain classes of morphisms of distinguished triangles are found to be good if and only if they are Verdier good. It is certain, however, that both goodness and Verdier goodness are stronger conditions than middling goodness. Something to take note of, is that even with all the extra structure inherent to

Date: June 30, 2022

2020 Mathematics Subject Classification. Primary 18G80.

Key words and phrases. good morphism, Verdier good morphism, middling good morphism, n -angulated category.

the derived category $D(\mathbb{Z})$, which is an *algebraic triangulated category*, there are simple examples of morphisms which are not middling good; take for instance the following diagram arising from a mapping of the embedding of the exact sequence

$$0 \longrightarrow \mathbb{Z} \xrightarrow{n} \mathbb{Z} \xrightarrow{q_n} \mathbb{Z}/n \longrightarrow 0$$

in degree 0 to its left rotation:

$$\begin{array}{ccccccc} \mathbb{Z}[0] & \xrightarrow{n} & \mathbb{Z}[0] & \xrightarrow{q_n} & \mathbb{Z}/n[0] & \xrightarrow{\varepsilon_n} & \mathbb{Z}[1] \\ \downarrow 0 & & \downarrow 0 & & \downarrow \varepsilon_n & & \downarrow 0 \\ \mathbb{Z}[0] & \xrightarrow{q_n} & \mathbb{Z}/n[0] & \xrightarrow{\varepsilon_n} & \mathbb{Z}[1] & \xrightarrow{-n} & \mathbb{Z}[1]. \end{array}$$

One can show explicitly that this morphism does not extend to a (4×4) -diagram or one can appeal to [6, Proposition 7.13].

In the setting of n -angulated categories, as introduced by Geiss, Keller, and Oppermann in [8], even simple questions regarding the relationship of these three goodness criteria are difficult to answer; the extra commutativity relations and the distance between objects in an n -angle become obstructions to the tools and methods readily available in the triangulated setting. So far, the most concrete examples of n -angulated categories are those arising from $(n-2)$ -cluster tilting subcategories of triangulated categories described in [8] and the so-called exotic n -angulated categories described in [3].

What we shall see in this paper, is that in $(n-2)$ -cluster tilting n -angulated categories, not only do all commutative squares extend to an $(n+1) \times (n+1)$ -diagram wherein the first n rows and columns are n -angles, in fact, all morphisms of n -angles extend to such a diagram for $n > 3$ (Theorem 4.7). Since $(n-2)$ -cluster tilting n -angulated categories are algebraic n -angulated categories in the sense of [9], it is somewhat surprising that something similar does not hold for, say, $D(\mathbb{Z})$. Naturally one would want to know whether this holds true in general for n -angulated categories with $n > 3$. We provide a counterexample, i.e., for all $n > 3$ we find an n -angulated category and a morphism of n -angles within which is not middling good (Example 4.12).

We begin in Section 2 by recalling the standard definitions related to n -angulated categories. In Section 3.1 we recall the definitions and some of the results about the various goodness criteria, and in Section 3.2 we generalise them to the n -angulated setting. In Section 4.1 we recall the definition and n -angulation of $(n-2)$ -cluster tilting categories and prove our main result, and in Section 4.2 we provide an example of a morphism in exotic n -angulated categories which is not middling good.

Acknowledgements

I would like thank Marius Thauale for the many discussions that resulted in this project and for all of the advice he has imparted along the way. I would also like to thank Petter Bergh and Martin Frankland for their helpful comments and interest in the project.

2 Preliminary theory

Let \mathcal{C} be an additive category equipped with an automorphism $\Sigma: \mathcal{C} \rightarrow \mathcal{C}$ and let n an integer greater than or equal to three.

Definition 2.1 A diagram in \mathcal{C} of the form

$$A_1 \xrightarrow{\alpha_1} A_2 \xrightarrow{\alpha_2} A_3 \xrightarrow{\alpha_3} \cdots \xrightarrow{\alpha_{n-1}} A_n \xrightarrow{\alpha_n} \Sigma A_1$$

is called an n - Σ -sequence. We may denote such a sequence by A_\bullet and refer to the maps α_i as the *structure maps* of the sequence. In particular, we may say that α_1 is the *base* of the n - Σ -sequence and that the α_i are the *cones of α_1* for $i \in \{2, \dots, n\}$.

Definition 2.2 The *left* and *right rotations* of A_\bullet are the n - Σ -sequences

$$A_2 \xrightarrow{\alpha_2} A_3 \xrightarrow{\alpha_3} \cdots \xrightarrow{\alpha_{n-1}} A_n \xrightarrow{\alpha_n} \Sigma A_1 \xrightarrow{(-1)^n \Sigma \alpha_1} \Sigma A_2$$

and

$$\Sigma^{-1} A_n \xrightarrow{(-1)^n \Sigma^{-1} \alpha_n} A_1 \xrightarrow{\alpha_1} A_2 \xrightarrow{\alpha_2} A_3 \xrightarrow{\alpha_3} \cdots \xrightarrow{\alpha_{n-1}} A_n,$$

respectively. We will borrow the notation of the shift functor from cochain complexes and thus denote by $A_\bullet[1]$ the left rotation of A_\bullet and by $A_\bullet[-1]$ the right rotation. For an object A of \mathcal{C} , a sequence of the form

$$A \xrightarrow{1} A \longrightarrow 0 \longrightarrow \cdots \longrightarrow 0 \longrightarrow \Sigma A \quad (2.1)$$

and any of its rotations are called *trivial n - Σ -sequence*. The n - Σ -sequence (2.1), in particular, we denote by $(\Gamma A)_\bullet$.

An n - Σ -sequence A_\bullet is *exact* if the induced sequence

$$\cdots \longrightarrow \mathcal{C}(-, A_1) \longrightarrow \mathcal{C}(-, A_2) \longrightarrow \cdots \longrightarrow \mathcal{C}(-, A_n) \longrightarrow \mathcal{C}(-, \Sigma A_1) \longrightarrow \cdots$$

is exact as a cochain complex.

Definition 2.3 A *morphism* of n - Σ -sequences $\varphi_\bullet: A_\bullet \rightarrow B_\bullet$ is an n -tuple of morphisms $\varphi_\bullet = (\varphi_1, \varphi_2, \dots, \varphi_n)$ constituting a commutative diagram

$$\begin{array}{ccccccccccc} A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\ \downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & & & \downarrow \varphi_n & & \downarrow \Sigma \varphi_1 \\ B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma B_1. \end{array}$$

The morphism φ_\bullet is an *isomorphism* if φ_i is an isomorphism for each $i \in \{1, \dots, n\}$; if for some $1 \leq i \leq n$ the maps φ_i and φ_{i+1} are isomorphisms (identifying $\varphi_{n+1} := \Sigma \varphi_1$) we say that φ_\bullet is a *weak isomorphism*.

Remark 2.4. The collection of n - Σ -sequences over \mathcal{C} and the collection of morphisms between them constitutes a category. We may thus consider Γ a functor from \mathcal{C} to n - Σ -sequences taking an object to the trivial n - Σ -sequence (2.1) and extending a morphism of objects to a morphism of n - Σ -sequences in the obvious way as illustrated below.

$$\begin{array}{ccccccccccc}
A & & A & \xrightarrow{1} & A & \longrightarrow & 0 & \longrightarrow & \cdots & \longrightarrow & 0 & \longrightarrow & \Sigma A \\
\downarrow f & \dashv \Gamma & \downarrow f & & \downarrow f & & \downarrow & & & & \downarrow & & \downarrow \Sigma f \\
B & & B & \xrightarrow{1} & B & \longrightarrow & 0 & \longrightarrow & \cdots & \longrightarrow & 0 & \longrightarrow & \Sigma B.
\end{array} \quad \blacklozenge$$

Definition 2.5 Let \mathcal{N} denote a collection of n - Σ -sequences in \mathcal{C} . The triple $(\mathcal{C}, \Sigma, \mathcal{N})$ is an n -angulated category if it satisfies the following four axioms:

- (N1) (a) \mathcal{N} is closed under direct sums, direct summands, and isomorphisms of n - Σ -sequences.
(b) For any object A in \mathcal{C} , the trivial n - Σ -sequence $(\Gamma A)_\bullet$ is in \mathcal{N} .
(c) For each morphism $\alpha_1: A_1 \rightarrow A_2$ in \mathcal{C} there exists an n - Σ -sequence in \mathcal{N} with base α_1 .
- (N2) An n - Σ -sequence belongs to \mathcal{N} if and only if its left rotation belongs to \mathcal{N} .
- (N3) Given that the solid part of the diagram

$$\begin{array}{ccccccccccc}
A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\
\downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & & & \downarrow \varphi_n & & \downarrow \Sigma \varphi_1 \\
B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma B_1,
\end{array}$$

commutes and has rows in \mathcal{N} , the dotted morphisms exist such that $(\varphi_1, \varphi_2, \varphi_3, \dots, \varphi_n)$ is a morphism of n - Σ -sequence.

- (N4) Given that the solid part of Figure 1 commutes and has rows in \mathcal{N} , the

$$\begin{array}{ccccccccccc}
A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & A_4 & \xrightarrow{\alpha_4} & \cdots & \xrightarrow{\alpha_{n-2}} & A_{n-1} & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\
\parallel & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & \downarrow \varphi_4 & & & & \downarrow \varphi_{n-1} & & \downarrow \varphi_n & & \parallel \\
A_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & B_4 & \xrightarrow{\beta_4} & \cdots & \xrightarrow{\beta_{n-2}} & B_{n-1} & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma A_1 \\
\downarrow \alpha_1 & & \parallel & & \downarrow \psi_3 & & \downarrow \psi_4 & & & & \downarrow \psi_{n-1} & & \downarrow \psi_n & & \downarrow \Sigma \alpha_1 \\
A_2 & \xrightarrow{\varphi_2} & B_2 & \xrightarrow{\gamma_2} & C_3 & \xrightarrow{\gamma_3} & C_4 & \xrightarrow{\gamma_4} & \cdots & \xrightarrow{\gamma_{n-2}} & C_{n-1} & \xrightarrow{\gamma_{n-1}} & C_n & \xrightarrow{\gamma_n} & \Sigma A_2
\end{array}$$

Figure 1: An octahedron in an n -angulated category.

dotted morphisms exist such that each square commutes and such that the n - Σ -sequence of Figure 2 belongs to \mathcal{N} .

The collection \mathcal{N} is called an n -angulation of the category \mathcal{C} with respect to the autoequivalence Σ , and the elements of \mathcal{N} are called n -angles. A category \mathcal{T} is triangulated if and only if it is 3-angulated, so one may refer to the distinguished triangles of \mathcal{T} as 3-angles should one choose.

$$\begin{array}{c}
A_3 \xrightarrow{\begin{pmatrix} \alpha_3 \\ \varphi_3 \end{pmatrix}} A_4 \oplus B_3 \xrightarrow{\begin{pmatrix} -\alpha_4 & 0 \\ \varphi_4 & -\beta_3 \\ \lambda_4 & \psi_3 \end{pmatrix}} A_5 \oplus B_4 \oplus C_3 \xrightarrow{\mu_3} A_6 \oplus B_5 \oplus C_4 \\
\curvearrowright \mu_4 \curvearrowleft \\
\cdots \xrightarrow{\mu_{n-3}} A_n \oplus B_{n-1} \oplus C_{n-2} \xrightarrow{\eta} B_n \oplus C_{n-1} \xrightarrow{\begin{pmatrix} \psi_n & \gamma_{n-1} \end{pmatrix}} C_n \xrightarrow{(\Sigma\alpha_2)\gamma_n} \Sigma A_3
\end{array}$$

with $\mu_i = \begin{pmatrix} -\alpha_{i+2} & 0 & 0 \\ (-1)^i \varphi_{i+2} & -\beta_{i+1} & 0 \\ \lambda_{i+2} & \psi_{i+1} & \gamma_i \end{pmatrix}$ and $\eta = \begin{pmatrix} (-1)^n \varphi_n & -\beta_{n-1} & 0 \\ \lambda_n & \psi_{n-1} & \gamma_{n-2} \end{pmatrix}$.

Figure 2: The n -angle associated to the octahedron.

We will often refer to the diagram of Figure 1 as an *octahedron* and to the n -angle of Figure 2 as the *associated n -angle*. We will also refer to the axioms by their name (N1)–(N4), and in the case of triangulated categories we may at times refer to the relevant axioms as (TR1)–(TR4).

Remark 2.6. Similarly to the triangulated setting, (N1)(ii), (N2), and (N3) ensures that for an n -angle A_\bullet with structure morphisms α_\bullet , the map α_i is both a weak cokernel of α_{i-1} and a weak kernel of α_{i+1} . \blacklozenge

The category of n - Σ -sequences, in, for example, an n -angulated category, shares some similarities with categories of cochain complexes over abelian categories and, as such, we can borrow some concepts from this theory. As an example, one could bring the concept of chain homotopies to the setting of n - Σ -sequences as in [3] and thus we can think of an n -angle as being *nullhomotopic* (or *contractible*) in the sense of regular chain complexes.

Of course, when we have a notion of homotopy, the next thing we want is a notion of mapping cone, ideally one that is homotopy invariant. This already exists in categories of cochain complexes, and the idea directly translates to the setting of n - Σ -sequences.

Definition 2.7 Let $\varphi_\bullet: A_\bullet \rightarrow B_\bullet$ be a morphism of n - Σ -sequences

$$\begin{array}{ccccccccccc}
A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\
\downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & & & \downarrow \varphi_n & & \downarrow \Sigma\varphi_1 \\
B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma B_1
\end{array}$$

Then the mapping cone of φ_\bullet is the n - Σ -sequence

$$\begin{array}{c}
A_2 \oplus B_1 \xrightarrow{\begin{pmatrix} -\alpha_2 & 0 \\ \varphi_2 & \beta_1 \end{pmatrix}} A_3 \oplus B_2 \xrightarrow{\begin{pmatrix} -\alpha_3 & 0 \\ \varphi_3 & \beta_2 \end{pmatrix}} A_4 \oplus B_3 \\
\curvearrowright \begin{pmatrix} -\alpha_4 & 0 \\ \varphi_4 & \beta_3 \end{pmatrix} \curvearrowleft \\
\cdots \xrightarrow{\begin{pmatrix} -\alpha_n & 0 \\ \varphi_n & \beta_{n-1} \end{pmatrix}} \Sigma A_1 \oplus B_n \xrightarrow{\begin{pmatrix} -\Sigma\alpha_1 & 0 \\ \Sigma\varphi_1 & \beta_n \end{pmatrix}} \Sigma A_2 \oplus \Sigma B_1.
\end{array}$$

However, a mapping cone should be the same type of object as the source and target of the morphism from which it was constructed and, while the mapping cone is an n - Σ -sequence by construction, if we start from two n -angles

in an n -angulated category, the resulting mapping cone may not be an n -angle. This inspires the following definition.

Definition 2.8 In a n -angulated category, a morphism of n -angles

$$\varphi_{\bullet}: A_{\bullet} \rightarrow B_{\bullet}$$

is a *good morphism* if the mapping cone is an n -angle.

Remark 2.9. The ‘‘octahedral axiom’’ (N4) has a few variants. The original definition of n -angulated categories stated in [8] says the following:

For any two n -angles A_{\bullet} and B_{\bullet} and a solid commutative square such as the one below, there exist fill-in morphisms $\varphi_3, \dots, \varphi_n$ such that the full diagram commutes and such that $(\varphi_1, \varphi_2, \varphi_3, \dots, \varphi_n)$ is a good morphism of n -angles.

$$\begin{array}{ccccccccccc} A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\ \downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & & & \downarrow \varphi_n & & \downarrow \Sigma \varphi_1 \\ B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma B_1 \end{array}$$

It was proven in [4] that, fixing the axioms (N1)–(N3), this axiom is equivalent to an axiom defined in said article, called (N4*), and it was later proven in [1] that (N4*) is equivalent to the axiom (N4) stated in Definition 2.5. \blacklozenge

3 Goodness criteria

In this section we recall the various goodness criteria in the original triangulated setting and generalise their respective definitions to the n -angulated setting.

3.1 Triangulated categories

Definition 2.8 is a generalisation of the following definition, originally introduced by Neeman in [10].

Definition 3.1 In a triangulated category, a morphism of distinguished triangles

$$\begin{array}{ccccccc} A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \Sigma A_1 \\ \downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & \downarrow \Sigma \varphi_1 \\ B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \Sigma B_1 \end{array} \quad (3.1)$$

is a *good morphism* if the mapping cone

$$A_2 \oplus B_1 \xrightarrow{\begin{pmatrix} -\alpha_2 & 0 \\ \varphi_2 & \beta_1 \end{pmatrix}} A_3 \oplus B_2 \xrightarrow{\begin{pmatrix} -\alpha_3 & 0 \\ \varphi_3 & \beta_2 \end{pmatrix}} \Sigma A_1 \oplus B_3 \xrightarrow{\begin{pmatrix} -\Sigma \alpha_1 & 0 \\ \Sigma \varphi_1 & \beta_3 \end{pmatrix}} \Sigma A_2 \oplus \Sigma B_1$$

is a distinguished triangle.

As was discussed in the case of n -angulated categories in Remark 2.9, the octahedral axiom may be rephrased in terms of good morphisms. This is originally due to Neeman, the proof split between [10, Theorem 1.8] and [11, Proposition 1.4.6].

Theorem 3.2 (A. Neeman). *Suppose \mathcal{C} is a pretriangulated category. The category \mathcal{C} is triangulated if and only if for each diagram as the one below with both rows consisting of distinguished triangles, a map $\varphi_3: A_3 \rightarrow B_3$ exists making the diagram a good morphism. We may refer to φ_3 as a good fill-in.*

$$\begin{array}{ccccccc}
A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \Sigma A_1 \\
\downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & \downarrow \Sigma \varphi_1 \\
B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \Sigma B_1
\end{array} \tag{3.2}$$

Definition 3.3 A map of distinguished triangles (3.1) is *middling good* if it extends to a diagram

$$\begin{array}{ccccccc}
A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \Sigma A_1 \\
\downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & \downarrow \Sigma \varphi_1 \\
B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \Sigma B_1 \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
C_1 & \dashrightarrow & C_2 & \dashrightarrow & C_3 & \dashrightarrow & \Sigma C_1 \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
\Sigma A_1 & \xrightarrow{\Sigma \alpha_1} & \Sigma A_2 & \xrightarrow{\Sigma \alpha_2} & \Sigma A_3 & \xrightarrow{\Sigma \alpha_3} & \Sigma^2 A_1
\end{array}$$

Ω

such that all squares commute, except for the square Ω which anticommutes, and such that the first three rows and columns are all distinguished triangles. We may refer to φ_3 as a *middling good fill-in*.

Definition 3.4 A map of distinguished triangles (3.1) is *Verdier good* if φ_3 factors through an object S as $\nu_2 \mu_1$ where μ_1 arises from the octahedron

$$\begin{array}{ccccccc}
A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \Sigma A_1 \\
\parallel & & \downarrow \varphi_2 & & \downarrow \mu_1 & & \parallel \\
A_1 & \xrightarrow{\varphi_2 \alpha_1} & B_2 & \xrightarrow{\sigma_2} & S & \xrightarrow{\sigma_3} & \Sigma A_1 \\
\downarrow \alpha_1 & & \parallel & & \downarrow \mu_2 & & \downarrow \Sigma \alpha_1 \\
A_2 & \xrightarrow{\varphi_2} & B_2 & \xrightarrow{\tau_2} & T & \xrightarrow{\tau_3} & \Sigma A_2 \\
& & & & \downarrow \mu_3 & \swarrow \Sigma \alpha_2 & \\
& & & & \Sigma A_3 & &
\end{array} \tag{3.3}$$

and ν_2 arises from the octahedron

$$\begin{array}{ccccccc}
A_1 & \xrightarrow{\varphi_1} & B_1 & \xrightarrow{\rho_2} & R & \xrightarrow{\rho_3} & \Sigma A_1 \\
\parallel & & \downarrow \beta_1 & & \downarrow \nu_1 & & \parallel \\
A_1 & \xrightarrow{\beta_1 \varphi_1} & B_2 & \xrightarrow{\sigma_2} & S & \xrightarrow{\sigma_3} & \Sigma A_1 \\
\downarrow \varphi_1 & & \parallel & & \downarrow \nu_2 & & \downarrow \Sigma \varphi_1 \\
B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \Sigma B_1 \\
& & & & \downarrow \nu_3 & \swarrow \Sigma \rho_2 & \\
& & & & \Sigma R & &
\end{array} \tag{3.4}$$

A fill-in that factors in this way is called a *Verdier good fill-in*.

Note that $\beta_1 \varphi_1 = \varphi_2 \alpha_1$, so we can, in fact, use the same object S in both octahedra. Since the two octahedral diagrams share the middle row, pictorially the two octahedra will share a face and the two arising distinguished triangles form a “butterfly” with S at its center. Thus φ_3 should arise as the composition of maps going from the lower lefthand corner to the upper righthand corner in Figure 3. Conversely, extending φ_1 and φ_2 to distinguished triangles with




Figure 3: Two octahedra glued together by a face. In blue: the “butterfly.”

cones R and T respectively, the other composition through S , namely $\mu_2 \nu_1$, is a Verdier good fill-in for a morphism between these two new distinguished triangles.

Remark 3.5. Given the solid part of the diagram (3.2), grafting together (3.3) and (3.4) via their shared row and composing the result produces a Verdier good morphism from (TR4) and (TR1)(c). Thus a Verdier good fill-in always exists.

This notion of goodness was originally applied by Verdier in his proof of the so-called (4×4) -lemma which is described in [2] in order to prove that a middling good fill-in always exists, and said proof entails that Verdier good morphisms are always middling good. Furthermore, it is proven in [10, Theorem 2.3] that good morphisms are always middling good, so middling goodness is

the weakest goodness constraint we can apply to a morphism. It is also known that middling good morphisms are not necessarily good nor Verdier good, however, we can say for certain when a middling good morphism is Verdier good, as Verdier goodness is equivalent to being middling good and satisfying a specific extra condition as stated in [6, Lemma 3.4]. \blacklozenge

3.2 Higher triangulated categories

In this section we introduce the generalised notions of Verdier good and middling good morphisms in n -angulated categories.

Definition 3.6 In an n -angulated category, a morphism of n -angles

$$\begin{array}{ccccccccccc}
A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\
\downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & & & \downarrow \varphi_n & & \downarrow \Sigma \varphi_1 \\
B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma B_1
\end{array} \quad (3.5)$$

is *Verdier good* if for $3 \leq i \leq n$ the morphism φ_i factors as $\nu_{2i}\mu_{1i}$ through an object S_i , where the μ_{1i} arise from an octahedron

$$\begin{array}{ccccccccccccccc}
A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & A_4 & \xrightarrow{\alpha_4} & \cdots & \xrightarrow{\alpha_{n-2}} & A_{n-1} & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma A_1 \\
\parallel & & \downarrow \varphi_2 & & \downarrow \mu_{13} & \swarrow \lambda_4 & \downarrow \mu_{14} & & & & \downarrow \mu_{1,n-1} & \swarrow \lambda_n & \downarrow \mu_{1n} & & \parallel \\
A_1 & \xrightarrow{\varphi_2 \alpha_1} & B_2 & \xrightarrow{\sigma_2} & S_3 & \xrightarrow{\sigma_3} & S_4 & \xrightarrow{\sigma_4} & \cdots & \xrightarrow{\sigma_{n-2}} & S_{n-1} & \xrightarrow{\sigma_{n-1}} & S_n & \xrightarrow{\sigma_n} & \Sigma A_1 \\
\downarrow \alpha_1 & & \parallel & & \downarrow \mu_{23} & \swarrow \tau_3 & \downarrow \mu_{24} & & & & \downarrow \mu_{2,n-1} & \swarrow \tau_{n-1} & \downarrow \mu_{2n} & & \downarrow \Sigma \alpha_1 \\
A_2 & \xrightarrow{\varphi_2} & B_2 & \xrightarrow{\tau_2} & T_3 & \xrightarrow{\tau_3} & T_4 & \xrightarrow{\tau_4} & \cdots & \xrightarrow{\tau_{n-2}} & T_{n-1} & \xrightarrow{\tau_{n-1}} & T_n & \xrightarrow{\tau_n} & \Sigma T_2
\end{array} \quad (3.6)$$

and the ν_{2i} arise from an octahedron

$$\begin{array}{ccccccccccccccc}
A_1 & \xrightarrow{\varphi_1} & B_1 & \xrightarrow{\rho_2} & R_3 & \xrightarrow{\rho_3} & R_4 & \xrightarrow{\rho_4} & \cdots & \xrightarrow{\rho_{n-2}} & R_{n-1} & \xrightarrow{\rho_{n-1}} & R_n & \xrightarrow{\rho_n} & \Sigma A_1 \\
\parallel & & \downarrow \beta_1 & & \downarrow \nu_{13} & \swarrow \gamma_4 & \downarrow \nu_{14} & & & & \downarrow \nu_{1,n-1} & \swarrow \gamma_n & \downarrow \nu_{1n} & & \parallel \\
A_1 & \xrightarrow{\beta_1 \varphi_1} & B_2 & \xrightarrow{\sigma_2} & S_3 & \xrightarrow{\sigma_3} & S_4 & \xrightarrow{\sigma_4} & \cdots & \xrightarrow{\sigma_{n-2}} & S_{n-1} & \xrightarrow{\sigma_{n-1}} & S_n & \xrightarrow{\sigma_n} & \Sigma A_1 \\
\downarrow \varphi_1 & & \parallel & & \downarrow \nu_{23} & \swarrow \beta_3 & \downarrow \nu_{24} & & & & \downarrow \nu_{2,n-1} & \swarrow \beta_{n-1} & \downarrow \nu_{2n} & & \downarrow \Sigma \varphi_1 \\
B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & B_4 & \xrightarrow{\beta_4} & \cdots & \xrightarrow{\beta_{n-2}} & B_{n-1} & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma B_2
\end{array} \quad (3.7)$$

Remark 3.7. A Verdier good fill-in always exists by a similar argument to that of triangulated categories. See for example the proof of [1, Theorem 3.1] wherein the morphism constructed is Verdier good. \blacklozenge

Definition 3.8 A morphism of n -angles (3.5) in an n -angulated category, where we label

$$A_{1i} := A_i, \quad A_{2i} := B_i, \quad \alpha_{1i} := \alpha_i, \quad \alpha_{2i} := \beta_i, \quad \text{and} \quad \varphi_{1i} := \varphi_i,$$

is *middling good* if it can be extended to a diagram

$$\begin{array}{ccccccc}
A_{11} & \xrightarrow{\alpha_{11}} & A_{12} & \xrightarrow{\alpha_{12}} & \cdots & \xrightarrow{\alpha_{1,n-1}} & A_{1n} & \xrightarrow{\alpha_{1n}} & \Sigma A_{11} \\
\downarrow \varphi_{11} & & \downarrow \varphi_{12} & & & & \downarrow \varphi_{1n} & & \downarrow \Sigma \varphi_{11} \\
A_{21} & \xrightarrow{\alpha_{21}} & A_{22} & \xrightarrow{\alpha_{22}} & \cdots & \xrightarrow{\alpha_{2,n-1}} & A_{2n} & \xrightarrow{\alpha_{2n}} & \Sigma A_{21} \\
\downarrow \varphi_{21} & & \downarrow \varphi_{22} & & & & \downarrow \varphi_{2n} & & \downarrow \Sigma \varphi_{21} \\
\vdots & & \vdots & & & & \vdots & & \vdots \\
\downarrow \varphi_{n-1,1} & & \downarrow \varphi_{n-1,2} & & & & \downarrow \varphi_{n-1,n} & & \downarrow \Sigma \varphi_{n-1,1} \\
A_{n1} & \xrightarrow{\alpha_{n1}} & A_{n2} & \xrightarrow{\alpha_{n2}} & \cdots & \xrightarrow{\alpha_{n,n-1}} & A_{nn} & \xrightarrow{\alpha_{nn}} & \Sigma A_{n1} \\
\downarrow \varphi_{n1} & & \downarrow \varphi_{n2} & & & & \downarrow \varphi_{nn} & \Omega & \downarrow \Sigma \varphi_{n1} \\
\Sigma A_{11} & \xrightarrow{\Sigma \alpha_{11}} & \Sigma A_{12} & \xrightarrow{\Sigma \alpha_{12}} & \cdots & \xrightarrow{\Sigma \alpha_{1,n-1}} & \Sigma A_{1n} & \xrightarrow{\Sigma \alpha_{1n}} & \Sigma^2 A_{11}
\end{array} \tag{3.8}$$

where the first n columns and n rows are all n -angles and in which all squares commute when n is even, and the square Ω anticommutes when n is odd.

We should like to observe the relationships of Remark 3.5 in the n -angulated setting and, while each goodness condition has a quite clear candidate in this setting as seen above, the arguments and methods used by Verdier, Neeman, and Christensen–Frankland do not easily generalise. Even the proof that Verdier good morphisms are middling good in a 4-angulated category eludes us; a (5×5) -diagram consisting mostly of 4-angles is obtainable but, so far, we have been unable to prove whether or not the square of cone objects in the diagram commutes. For now, let us move away from the model free setting in order to say something concrete.

4 Concrete higher triangulations

It was originally proved in [8] that an $(n - 2)$ -cluster tilting subcategory in a triangulated category with suspension Σ has an n -angulation if it is stable with respect to Σ^{n-2} . We show in Section 4.1 that not only are Verdier good morphisms also middling good in this class of n -angulated categories, in fact, all morphisms of n -angles are middling good. One may wonder whether this is the case for all n -angulated categories, so in Section 4.2 we provide an example of a morphism in an exotic n -angulated category which is not middling good.

4.1 Cluster tilting categories

We begin by recalling the relevant definitions to define the n -angulated structure on $(n - 2)$ -cluster tilting categories. Following that, we make some specific observations on the nature of n -angles in this setting before moving on to the proof of the claim.

Definition 4.1 For a category \mathcal{T} and a full subcategory \mathcal{C} , an object $X \in \mathcal{T}$ has a *left \mathcal{C} -approximation* if there exists a morphism $f \in \mathcal{T}(X, C)$ with $C \in \mathcal{C}$ such that for any object $C' \in \mathcal{C}$ and map $g \in \mathcal{T}(X, C')$ there exists a map $h \in \mathcal{C}(C, C')$ such that $hf = g$.

$$\begin{array}{ccc}
X & \xrightarrow{f} & C \\
& \searrow g & \downarrow h \\
& & C'
\end{array}$$

If all objects in \mathcal{T} have this property, it is called *covariantly finite*. If \mathcal{T}^{op} is covariantly finite, then \mathcal{T} is *contravariantly finite*. Finally, if \mathcal{T} is both covariantly and contravariantly finite, we say that \mathcal{T} is *functorially finite*.

Definition 4.2 Let \mathcal{T} be a triangulated category with a class of distinguished triangles Δ and with an autoequivalence Σ . A full subcategory $\mathcal{C} \subset \mathcal{T}$ is *d-cluster tilting* if it is functorially finite and it satisfies the condition

$$\begin{aligned}
\mathcal{C} &= \{X \in \mathcal{T} \mid \mathcal{T}(\mathcal{C}, \Sigma^i X) = 0 \text{ for all } i \in \{1, \dots, d-1\}\} \\
&= \{X \in \mathcal{T} \mid \mathcal{T}(X, \Sigma^i \mathcal{C}) = 0 \text{ for all } i \in \{1, \dots, d-1\}\}.
\end{aligned}$$

Let us now recall the theorem of [8].

Theorem 4.3 (Geiss–Keller–Oppermann). *Let $(\mathcal{T}, \Sigma, \Delta)$ be a triangulated category with an $(n-2)$ -cluster tilting subcategory \mathcal{C} such that the objects of $\Sigma^{n-2}\mathcal{C}$ are contained in \mathcal{C} . Then $(\mathcal{C}, \Sigma^{n-2}, \mathcal{N})$ is an n -angulated category, where \mathcal{N} is the class of all sequences*

$$A_1 \xrightarrow{\alpha_1} A_2 \xrightarrow{\alpha_2} \dots \xrightarrow{\alpha_{n-1}} A_n \xrightarrow{\alpha_n} \Sigma^{n-2} A_1$$

such that there exists a diagram

$$\begin{array}{ccccccc}
& & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & A_4 & \xrightarrow{\alpha_4} & \dots & \xrightarrow{\alpha_{n-2}} & A_{n-1} \\
& \nearrow \alpha_1 & \searrow \alpha'_2 & \nearrow \alpha''_2 & \searrow \alpha'_3 & \nearrow \alpha''_3 & & & & \nearrow \alpha''_{n-2} & \searrow \alpha_{n-1} \\
A_1 & \xleftarrow{\alpha_2} & A_{2.5} & \xleftarrow{\alpha_3} & A_{3.5} & \xleftarrow{\alpha_4} & \dots & \xleftarrow{\alpha_{n-2}} & A_{n-1.5} & \xleftarrow{\alpha_{n-1}} & A_n
\end{array}$$

which we will refer to as a the trapez diagram of an n -angle, with $A_i \in \mathcal{T}$ for $i \notin \mathbb{Z}$, such that

$$(\alpha''_i, \alpha'_{i+1}, \alpha_n^{i+1}) \in \Delta \quad \text{for } i \in \{1, \dots, n-2\},$$

where $\alpha''_1 := \alpha_1$ and $\alpha'_{n-1} := \alpha_{n-1}$, and such that

$$\alpha_i = \alpha''_i \alpha'_i \quad \text{for } i \in \{2, \dots, n-2\}$$

and

$$\alpha_n = (\Sigma^{n-3} \alpha_n^2)(\Sigma^{n-4} \alpha_n^3) \dots (\Sigma \alpha_n^{n-2}) \alpha_n^{n-1}.$$

We call the objects $A_{i+1.5}$ the supporting objects of the n -angle for $i \in \{1, \dots, n-3\}$.

Remark 4.4. Assuming $n > 3$, in proving that the category \mathcal{C} in Theorem 4.3 satisfies axiom (N1)(c), one can construct a \mathcal{C} -resolution of A_1 inductively, starting with α_1 , by applying (TR1)(c) for α''_i ($\alpha''_1 = \alpha_1$) and then taking the left \mathcal{C} -approximations of the arising cone of α''_i , i.e. the supporting objects, to obtain the maps α'_i . Proving that this process terminates, i.e. that the cone of

α''_{n-2} is an object in \mathcal{C} , is, in our opinion, not trivial but we will not reproduce the argument here. Conversely, given a trapez diagram as in Theorem 4.3, it turns out that $\alpha''_{i+1}: A_{i+1.5} \rightarrow A_{i+2}$ must, in fact, be a left approximation. To see this, fix an i such that $1 \leq i \leq n-3$. First, given a morphism $\gamma: A_{i+1.5} \rightarrow C$ where $C \in \mathcal{C}$, since

$$\gamma \alpha'_{i+1} \alpha_i = \gamma(\alpha'_{i+1} \alpha''_i) \alpha'_i = 0,$$

we obtain the solid diagram

$$\begin{array}{ccccccccccc} A_i & \xrightarrow{\alpha_i} & A_{i+1} & \xrightarrow{\alpha_{i+1}} & A_{i+2} & \xrightarrow{\alpha_{i+2}} & \cdots & \xrightarrow{-\Sigma\alpha_{i-1}} & \Sigma A_{i-1} & \xrightarrow{-\Sigma\alpha_i} & \Sigma A_i \\ \downarrow & & \downarrow \gamma \alpha'_{i+1} & & \downarrow \bar{\gamma} & & & & \downarrow & & \downarrow \\ 0 & \longrightarrow & C & \xlongequal{\quad} & C & \longrightarrow & \cdots & \longrightarrow & 0 & \longrightarrow & \Sigma C \end{array}$$

and applying (N3) we gain a dashed map $\bar{\gamma} \in \mathcal{C}(A_{i+2}, C)$ such that

$$\bar{\gamma} \alpha_{i+1} = \gamma \alpha'_{i+1}.$$

From this identity, as $\alpha_{i+1} = \alpha''_{i+1} \alpha'_{i+1}$, we can conclude that

$$(\bar{\gamma} \alpha''_{i+1} - \gamma) \alpha'_{i+1} = 0,$$

so since $\alpha_n^{i+1}: A_{i+1.5} \rightarrow \Sigma A_{i+5}$ is a weak cokernel of α'_{i+1} in \mathcal{T} , there is a map

$$\Sigma\gamma_1: \Sigma A_{i+5} \rightarrow C$$

satisfying $(\Sigma\gamma_1) \alpha_n^{i+1} = \bar{\gamma} \alpha''_{i+1} - \gamma$. Recall that \mathcal{C} is closed under Σ^{n-2} , so $\Sigma^{n-2} A_i \in \mathcal{C}$, and therefore we get that

$$\gamma_1 \alpha'_i \in \mathcal{T}(A_i, \Sigma^{-1}C) \cong \mathcal{T}(\Sigma^{n-2} A_i, \Sigma^{n-3}C) = 0$$

by the cluster tilting condition. Note that the cluster tilting condition only guarantees this because $n-2 > 1$, and this is why we only consider $(n-2)$ -cluster tilting categories for $n > 3$. Since α_n^i is a weak cone of α'_i , we again obtain a map

$$\Sigma\gamma_2: \Sigma A_{i-.5} \rightarrow \Sigma^{-1}C$$

such that $(\Sigma\gamma_2) \alpha_n^i = \gamma_1$. We repeat this process until we get a map

$$\Sigma\gamma_i: \Sigma A_1 \rightarrow \Sigma^{1-i}C$$

such that $(\Sigma\gamma_i) \alpha_n^2 = \gamma_{i-1}$. Inductively we see that

$$(\Sigma^{i-1} \gamma_i) (\Sigma^{i-2} \alpha_n^2) (\Sigma^{i-3} \alpha_n^3) \cdots (\Sigma \alpha_n^{i-1}) \alpha_n^i = \gamma_1$$

but since $1 \leq i \leq n-3$ it follows that $1 \leq n-i-2 \leq n-3$, so

$$\gamma_i \in \mathcal{T}(A_1, \Sigma^{-i}C) \cong \mathcal{T}(\Sigma^{n-2} A_1, \Sigma^{n-i-2}C) = 0.$$

The important difference between γ_i and γ_k for $k < i$ here, is that the source of γ_i is an object in \mathcal{C} which is not necessarily true of γ_k . Thus, it follows that $\gamma_1 = 0$ and we conclude that

$$\bar{\gamma} \alpha''_{i+1} - \gamma = (\Sigma\gamma_1) \alpha_n^{i+1} = 0,$$

i.e. for all maps $\gamma: A_{i+1.5} \rightarrow C$ where $C \in \mathcal{C}$, there exists a map $\bar{\gamma}: A_{i+2} \rightarrow C$ in \mathcal{C} such that $\bar{\gamma} \alpha''_{i+1} = \gamma$, and therefore $\alpha''_{i+1}: A_{i+1.5} \rightarrow A_{i+2}$ is a left \mathcal{C} -approximation. \blacklozenge

Remark 4.5. Since any n -angle in a cluster tilting subcategory is made up of left approximations, we may, similarly to the previous remark, freely construct a morphism of n -angles from a square

$$\begin{array}{ccc} A_1 & \xrightarrow{\alpha_1} & A_2 \\ \downarrow \varphi_1 & & \downarrow \varphi_2 \\ B_1 & \xrightarrow{\beta_1} & B_2 \end{array}$$

by iterately applying (TR3) and the universal property of the left approximations to the trapez prism diagram as illustrated in Figure 4. The entire diagram of Figure 4 therefore commutes by construction. Conversely, given a morphism of n -angles, any fill-in between the support objects that arise from (TR3) applied to the constituent distinguished triangles, constitutes a commutative trapez diagram. We state this more precisely as a lemma. \blacklozenge

Figure 4: The dotted morphisms come from (TR3) while the dashed morphisms come from the universal property of left approximations.


Lemma 4.6 *Let $n > 3$ be an integer. Suppose we have a morphism of n -angles in an n -angulated $(n - 2)$ -cluster tilting subcategory*

$$\begin{array}{ccccccc} A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma^{n-2} A_1 \\ \downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_3 & & & & \downarrow \varphi_n & & \downarrow \Sigma \varphi_1 \\ B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta_2} & B_3 & \xrightarrow{\beta_3} & \cdots & \xrightarrow{\beta_{n-1}} & B_n & \xrightarrow{\beta_n} & \Sigma^{n-2} B_1. \end{array}$$

where the upper and lower row have support objects $A_{i+1.5}$ for $i \in \{1, \dots, n-3\}$ and $B_{i+1.5}$ for $i \in \{1, \dots, n-3\}$ respectively. If we inductively fill in the diagrams

$$\begin{array}{ccccccc} A_{i+1.5} & \xrightarrow{\alpha'_{i+1}} & A_{i+2} & \xrightarrow{\alpha'_{i+2}} & A_{i+2.5} & \xrightarrow{\alpha_n^{i+2}} & \Sigma A_{i+1.5} \\ \downarrow \varphi_{i+1.5} & & \downarrow \varphi_{i+2} & & \downarrow \varphi_{i+2.5} & & \downarrow \Sigma \varphi_{i+1.5} \\ B_{i+1.5} & \xrightarrow{\beta'_{i+1}} & B_{i+2} & \xrightarrow{\beta'_{i+2}} & B_{i+2.5} & \xrightarrow{\beta_n^{i+2}} & \Sigma B_{i+1.5} \end{array}$$

starting with $i = 0$ and applying the conventions $A_{1.5} := A_1$, $B_{1.5} := B_1$, $\alpha''_1 := \alpha_1$, and $\beta''_1 := \beta_1$, the trapez prism diagram



will commute.

Proof. We progress in a similar fashion to the preceding remark. Fill in the diagram

$$\begin{array}{ccccccc}
 A_1 & \xrightarrow{\alpha_1} & A_2 & \xrightarrow{\alpha'_2} & A_{2.5} & \xrightarrow{\alpha_n^2} & \Sigma A_1 \\
 \downarrow \varphi_1 & & \downarrow \varphi_2 & & \downarrow \varphi_{2.5} & & \downarrow \Sigma \varphi_1 \\
 B_1 & \xrightarrow{\beta_1} & B_2 & \xrightarrow{\beta'_2} & B_{2.5} & \xrightarrow{\beta_n^2} & \Sigma B_1
 \end{array} \quad (4.1)$$

however you like such that the diagram commutes. Of course, there always is at least one choice by (TR3). Drawing in $\varphi_{2.5}$ in the trapez prism, we know that the solid part commutes along with the two new squares that arise from (4.1) but, possibly, we might not have the commutativity relation $\varphi_3 \alpha''_2 = \beta''_2 \varphi_{2.5}$. To see that this identity also holds, note that

$$\varphi_3 \alpha''_2 \alpha'_2 = \varphi_3 \alpha_2 = \beta_2 \varphi_2 = \beta''_2 \beta'_2 \varphi_2 = \beta''_2 \varphi_{2.5} \alpha'_2$$

so

$$(\beta''_2 \varphi_{2.5} - \varphi_3 \alpha''_2) \alpha'_2 = 0,$$

and thus there is a morphism

$$\Sigma \gamma: \Sigma A_1 \rightarrow B_3$$

such that $(\Sigma \gamma) \alpha_n^2 = \beta''_2 \varphi_{2.5} - \varphi_3 \alpha''_2$. However, as before, $\Sigma \gamma = 0$ since

$$\mathcal{T}(\Sigma A_1, B_3) \cong \mathcal{T}(\Sigma^{n-2} A_1, \Sigma^{n-3} B_3) = 0,$$

so we conclude that the commutativity relation holds. Again, this follows from the fact that $n - 2 > 1$. Now suppose that $\varphi_{i+1} \alpha''_i = \beta''_i \varphi_{i+1.5}$ for some fixed $i \in \{2, \dots, n-3\}$. To see that $\varphi_{i+2} \alpha''_{i+1} = \beta''_{i+1} \varphi_{i+1.5}$, we note again that

$$(\beta''_{i+1} \varphi_{i+1.5} - \varphi_{i+2} \alpha''_{i+1}) \alpha'_{i+1} = 0$$

so there is a map

$$\Sigma \gamma_1: \Sigma A_{i+1.5} \rightarrow B_{i+2}$$

such that $(\Sigma\gamma_1)\alpha_n^{i+1} = \beta''_{i+1}\varphi_{i+1.5} - \varphi_{i+2}\alpha''_{i+1}$, and

$$(\Sigma\gamma_1)\alpha'_i \in \mathcal{T}(\Sigma A_2, B_{i+2}) \cong \mathcal{T}(\Sigma^{n-2}A_2, \Sigma^{n-3}B_{i+2}) = 0,$$

so from the cokernel property of α_n^i we get another map

$$\Sigma\gamma_2: \Sigma A_{i-.5} \rightarrow \Sigma^{-1}B_{i+2}$$

such that $(\Sigma\gamma_2)\alpha_n^i = \gamma_1$. Now repeating this process exactly as we did in the preceding remark but with B_{i+2} in place of C , we conclude that

$$\gamma_1 = 0$$

and so

$$\varphi_{i+2}\alpha''_{i+1} = \beta''_{i+1}\varphi_{i+.5}$$

for all $i \in \{2, \dots, n-3\}$. \square

We have now arrived at our main result of this paper. Since all n -angles in $(n-2)$ -cluster tilting subcategories are built from triangles in the ambient triangulated category, diagrams witnessing middling good morphisms can be built up from the triangulated structure.

Theorem 4.7 *For $n > 3$, all morphisms of n -angles in an $(n-2)$ -cluster tilting n -angulated category are middling good.*

Notation guide for the proof: We construct an $(n+1) \times (n+1)$ diagram of objects labeled A_{ij} with $A_{i\bullet}$ and $A_{\bullet j}$ representing n -angles and, in particular, when i or j are not integers, A_{ij} represents a support object. We will also adopt the convention that horizontal morphisms are labeled α_{ij} and vertical morphisms are labeled φ_{ij} with ij being the index of the source object in both cases. Thus $\alpha_{i\bullet}$ are the structure morphisms of $A_{i\bullet}$ while $\varphi_{\bullet j}$ are the structure morphisms for $A_{\bullet j}$. This will be put aside in the case of support objects, though, as we in this case opt to follow the convention of writing $\alpha''_{*,*}$, $\alpha'_{*,*}$, and $\alpha^*_{*,*}$ similarly to what we have been doing so far.

Proof. We begin with a morphism of n -angles, i.e. a commutative diagram

$$\begin{array}{ccccccccccc} A_{11} & \xrightarrow{\alpha_{11}} & A_{12} & \xrightarrow{\alpha_{12}} & A_{13} & \xrightarrow{\alpha_{13}} & \cdots & \xrightarrow{\alpha_{1,n-1}} & A_{1n} & \xrightarrow{\alpha_{1n}} & \Sigma A_{11} \\ \downarrow \varphi_{11} & & \downarrow \varphi_{12} & & \downarrow \varphi_{13} & & & & \downarrow \varphi_{1n} & & \downarrow \Sigma\varphi_{11} \\ A_{21} & \xrightarrow{\alpha_{21}} & A_{22} & \xrightarrow{\alpha_{22}} & A_{23} & \xrightarrow{\alpha_{23}} & \cdots & \xrightarrow{\alpha_{2,n-1}} & A_{2n} & \xrightarrow{\alpha_{2n}} & \Sigma A_{21} \end{array}$$

such that the rows are n -angles. Since we, by Lemma 4.6, can choose any fill-in morphism we like between the support objects of $A_{1\bullet}$ and $A_{2\bullet}$, we can choose them to be middling good. For short, we may say that the morphism has *middling good support*. For each morphism of triangles, because the third

morphism is a middling good fill-in, we have the diagram

$$\begin{array}{ccccccc}
A_{1,i+.5} & \xrightarrow{\alpha''_{1i}} & A_{1,i+1} & \xrightarrow{\alpha'_{1,i+1}} & A_{1,i+1.5} & \xrightarrow{\alpha_{1n}^{i+1}} & \Sigma A_{1,i+.5} \\
\downarrow \varphi_{1,i+.5} & & \downarrow \varphi_{1,i+1} & & \downarrow \varphi_{1,i+1.5} & & \downarrow \Sigma \varphi_{1,i+.5} \\
A_{2,i+.5} & \xrightarrow{\alpha''_{2i}} & A_{2,i+1} & \xrightarrow{\alpha'_{2,i+1}} & A_{2,i+1.5} & \xrightarrow{\alpha_{2n}^{i+1}} & \Sigma A_{2,i+.5} \\
\downarrow \varphi'_{2,i+.5} & & \downarrow \varphi'_{2,i+1} & & \downarrow \varphi'_{2,i+1.5} & & \downarrow \Sigma \varphi'_{2,i+.5} \\
A_{2.5,i+.5} & \dashrightarrow^{\alpha''_{2.5,i}} & A_{2.5,i+1} & \dashrightarrow^{\alpha'_{2.5,i+1}} & A_{2.5,i+1.5} & \dashrightarrow^{\alpha_{2.5,n}^{i+1}} & \Sigma A_{2.5,i+.5} \\
\downarrow \varphi_{i+.5,n}^2 & & \downarrow \varphi_{i+1,n}^2 & & \downarrow \varphi_{i+1.5,n}^2 & \Omega_{1i} & \downarrow \Sigma \varphi_{i+.5,n}^2 \\
\Sigma A_{1,i+.5} & \xrightarrow{\Sigma \alpha''_{1i}} & \Sigma A_{1,i+1} & \xrightarrow{\Sigma \alpha'_{1,i+1}} & \Sigma A_{1,i+1.5} & \xrightarrow{\Sigma \alpha_{1n}^{i+1}} & \Sigma^2 A_{1,i+.5}.
\end{array} \tag{4.2}$$

The dotted morphisms represent the new distinguished triangles we get from (TR1)(c) and the dashed morphisms represent the new distinguished triangle we get from middling goodness. We now fix i . Recall that the square Ω_{1i} anticommutes and that the morphism $\alpha''_{2.5,i}: A_{2.5,i+.5} \rightarrow A_{2.5,i+1}$ is a middling good fill-in between the two left-most vertical distinguished triangles in the diagram. By (N1)(c) we can produce an n -angle $A_{\bullet,i+.5}$ with base $\varphi_{1,i+.5}: A_{1,i+.5} \rightarrow A_{2,i+.5}$, and by uniqueness of cones in triangulated categories, we may assume that the object $A_{2.5,i+.5}$ is the initial support object for $A_{\bullet,i+.5}$. Similarly, we construct an n -angle $A_{\bullet,i+1}$ with initial support object $A_{2.5,i+1}$. We can further construct a morphism of n -angles $A_{\bullet,i+.5} \rightarrow A_{\bullet,i+1}$ such that the morphism between the initial support objects is $\alpha''_{2.5,i}$ by repeatedly applying the left approximation property and (TR3), and we may assume that the fill-ins between the remaining support objects are middling good. Therefore, we in general get middling good diagrams

$$\begin{array}{ccccccc}
A_{j+.5,i+.5} & \xrightarrow{\alpha''_{j+.5,i}} & A_{j+.5,i+1} & \dashrightarrow^{\alpha'_{j+.5,i+1}} & A_{j+.5,i+1.5} & \dashrightarrow^{\alpha_{j+.5,n}^{i+1}} & \Sigma A_{j+.5,i+.5} \\
\downarrow \varphi''_{j,i+.5} & & \downarrow \varphi''_{j,i+1} & & \downarrow \varphi''_{j,i+1.5} & & \downarrow \Sigma \varphi''_{j,i+.5} \\
A_{j+1,i+.5} & \xrightarrow{\alpha''_{j+1,i}} & A_{j+1,i+1} & \dashrightarrow^{\alpha'_{j+1,i+1}} & A_{j+1,i+1.5} & \dashrightarrow^{\alpha_{j+1,n}^{i+1}} & \Sigma A_{j+1,i+.5} \\
\downarrow \varphi'_{j+1,i+.5} & & \downarrow \varphi'_{j+1,i+1} & & \downarrow \varphi'_{j+1,i+1.5} & & \downarrow \Sigma \varphi'_{j+1,i+.5} \\
A_{j+1.5,i+.5} & \xrightarrow{\alpha''_{j+1.5,i}} & A_{j+1.5,i+1} & \dashrightarrow^{\alpha'_{j+1.5,i+1}} & A_{j+1.5,i+1.5} & \dashrightarrow^{\alpha_{j+1.5,n}^{i+1}} & \Sigma A_{j+1.5,i+.5} \\
\downarrow \varphi_{n,i+.5}^{j+1} & & \downarrow \varphi_{n,i+1}^{j+1} & & \downarrow \varphi_{n,i+1.5}^{j+1} & \Omega_{ji} & \downarrow \Sigma \varphi_{n,i+.5}^{j+1} \\
\Sigma A_{j+.5,i+.5} & \xrightarrow{\Sigma \alpha''_{j+.5,i}} & \Sigma A_{j+.5,i+1} & \xrightarrow{\Sigma \alpha'_{j+.5,i+1}} & \Sigma A_{j+.5,i+1.5} & \xrightarrow{\Sigma \alpha_{j+.5,n}^{i+1}} & \Sigma^2 A_{j+.5,i+.5}
\end{array}$$

where Ω_{ji} is anticommutative. We may call the above diagram Γ_{ji} . For the previously fixed i , we now graft together the diagrams Γ_{ji} vertically diagrams; the third row of Γ_{ji} is the first row of $\Gamma_{j+1,i}$, and so we extend this diagram by composing the vertical morphisms in

$$\begin{array}{ccccccc}
A_{j+1,i+.5} & \xrightarrow{\alpha''_{j+1,i}} & A_{j+1,i+1} & \xrightarrow{\alpha'_{j+1,i+1}} & A_{j+1,i+1.5} & \xrightarrow{\alpha_{j+1,n}^{i+1}} & \Sigma A_{j+1,i+.5} \\
\downarrow \varphi'_{j+1,i+.5} & & \downarrow \varphi'_{j+1,i+1} & & \downarrow \varphi'_{j+1,i+1.5} & & \downarrow \Sigma \varphi'_{j+1,i+.5} \\
A_{j+1.5,i+.5} & \xrightarrow{\alpha''_{j+1.5,i}} & A_{j+1.5,i+1} & \xrightarrow{\alpha'_{j+1.5,i+1}} & A_{j+1.5,i+1.5} & \xrightarrow{\alpha_{j+1.5,n}^{i+1}} & \Sigma A_{j+1.5,i+.5} \\
\downarrow \varphi''_{j+1,i+.5} & & \downarrow \varphi''_{j+1,i+1} & & \downarrow \varphi''_{j+1,i+1.5} & & \downarrow \Sigma \varphi''_{j+1,i+.5} \\
A_{j+2,i+.5} & \xrightarrow{\alpha''_{j+2,i}} & A_{j+2,i+1} & \xrightarrow{\alpha'_{j+2,i+1}} & A_{j+2,i+1.5} & \xrightarrow{\alpha_{j+2,n}^{i+1}} & \Sigma A_{j+2,i+.5}
\end{array}$$

to get

$$\varphi_{j+1,i+.5} = \varphi''_{j+1,i+.5} \varphi'_{j+1,i+.5} \quad \text{and} \quad \varphi_{j+1,i+1} = \varphi''_{j+1,i+1} \varphi'_{j+1,i+1}$$

vertically by construction, and

$$\varphi_{j+1,i+1.5} := \varphi''_{j+1,i+1.5} \varphi'_{j+1,i+1.5}$$

by definition. Similarly, the suspension of the third row of Γ_{ji} is the last row of $\Gamma_{j+1,i}$, so we complete our extension of Γ_{ji} by composing the vertical morphisms in

$$\begin{array}{ccccccc}
A_{j+2.5,i+.5} & \xrightarrow{\alpha''_{j+2.5,i}} & A_{j+2.5,i+1} & \xrightarrow{\alpha'_{j+2.5,i+1}} & A_{j+2.5,i+1.5} & \xrightarrow{\alpha_{j+2.5,n}^{i+1}} & \Sigma A_{j+2.5,i+.5} \\
\downarrow \varphi_n^{j+2} & & \downarrow \varphi_n^{j+2} & & \downarrow \varphi_n^{j+2} & \Omega_{j+1,i} & \downarrow \Sigma \varphi_n^{j+2} \\
\Sigma A_{j+1.5,i+.5} & \xrightarrow{\Sigma \alpha''_{j+1.5,i}} & \Sigma A_{j+1.5,i+1} & \xrightarrow{\Sigma \alpha'_{j+1.5,i+1}} & \Sigma A_{j+1.5,i+1.5} & \xrightarrow{\Sigma \alpha_{j+1.5,n}^{i+1}} & \Sigma^2 A_{j+1.5,i+.5} \\
\downarrow \Sigma \varphi_n^{j+1} & & \downarrow \Sigma \varphi_n^{j+1} & & \downarrow \Sigma \varphi_n^{j+1} & \Sigma \Omega_{ji} & \downarrow \Sigma^2 \varphi_n^{j+1} \\
\Sigma^2 A_{j+.5,i+.5} & \xrightarrow{\alpha''_{j+.5,i}} & \Sigma^2 A_{j+.5,i+1} & \xrightarrow{\alpha'_{j+.5,i+1}} & \Sigma^2 A_{j+.5,i+1.5} & \xrightarrow{\alpha_{j+.5,n}^{i+1}} & \Sigma^3 A_{j+.5,i+.5}
\end{array}$$

We refrain from naming these compositions at this point, but we notice that the square enveloping $\Omega_{j+1,i}$ and $\Sigma \Omega_{ji}$ is commutative, since the individual squares are anticommutative. Composing the diagram vertically, we will think of the enveloping square as the vertical composition of $\Omega_{j+1,i}$ and $\Sigma \Omega_{ji}$ and thus we denote it by $\Sigma \Omega_{ji} \circ_v \Omega_{j+1,i}$. All told, we obtain a commutative diagram

$$\begin{array}{ccccccc}
A_{j+.5,i+.5} & \xrightarrow{\alpha''_{j+.5,i}} & A_{j+.5,i+1} & \xrightarrow{\alpha'_{j+.5,i+1}} & A_{j+.5,i+1.5} & \xrightarrow{\alpha_{j+.5,n}^{i+1}} & \Sigma A_{j+.5,i+.5} \\
\downarrow \varphi''_{j,i+.5} & & \downarrow \varphi''_{j,i+1} & & \downarrow \varphi''_{j,i+1.5} & & \downarrow \Sigma \varphi''_{j,i+.5} \\
A_{j+1,i+.5} & \xrightarrow{\alpha''_{j+1,i}} & A_{j+1,i+1} & \xrightarrow{\alpha'_{j+1,i+1}} & A_{j+1,i+1.5} & \xrightarrow{\alpha_{j+1,n}^{i+1}} & \Sigma A_{j+1,i+.5} \\
\downarrow \varphi_{j+1,i+.5} & & \downarrow \varphi_{j+1,i+1} & & \downarrow \varphi_{j+1,i+1.5} & & \downarrow \Sigma \varphi_{j+1,i+.5} \\
A_{j+2,i+.5} & \xrightarrow{\alpha''_{j+2,i}} & A_{j+2,i+1} & \xrightarrow{\alpha'_{j+2,i+1}} & A_{j+2,i+1.5} & \xrightarrow{\alpha_{j+2,n}^{i+1}} & \Sigma A_{j+2,i+.5} \\
\downarrow \varphi'_{j+2,i+.5} & & \downarrow \varphi'_{j+2,i+1} & & \downarrow \varphi'_{j+2,i+1.5} & & \downarrow \Sigma \varphi'_{j+2,i+.5} \\
A_{j+2.5,i+.5} & \xrightarrow{\alpha''_{j+2.5,i}} & A_{j+2.5,i+1} & \xrightarrow{\alpha'_{j+2.5,i+1}} & A_{j+2.5,i+1.5} & \xrightarrow{\alpha_{j+2.5,n}^{i+1}} & \Sigma A_{j+2.5,i+.5} \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
\Sigma^2 A_{j+.5,i+.5} & \xrightarrow{\Sigma^2 \alpha''_{j+.5,i}} & \Sigma^2 A_{j+.5,i+1} & \xrightarrow{\Sigma^2 \alpha'_{j+.5,i+1}} & \Sigma^2 A_{j+.5,i+1.5} & \xrightarrow{\Sigma^2 \alpha_{j+.5,n}^{i+1}} & \Sigma^3 A_{j+.5,i+.5}.
\end{array}$$

where the dashed morphisms represent the composite maps we constructed above. Applying this specific construction to Γ_{1i} , i.e. the diagram (4.2), we denote the resulting 5×4 diagram by Γ_i^1 . We now repeat the procedure to Γ_i^1 , noticing that the fourth row of this diagram corresponds to the first row of Γ_{3i} and the suspension of the fourth row of Γ_i^1 is the last row of Γ_{3i} . This grafting procedure thus produces a 6×4 diagram which we denote by Γ_i^2 , and in which the lowest row is $\Sigma^3 A_{1\bullet}$ and the lower right hand square is anticommutative since it is the one we denote by

$$\Sigma^2 \Omega_{1i} \circ_v \Sigma \Omega_{2i} \circ_v \Omega_{3i}.$$

We now repeatedly graft on Γ_{ji} to Γ_i^{j-2} for $j \in \{3, \dots, n-2\}$ until we arrive at the $(n+1) \times 4$ diagram $\Gamma_i := \Gamma_i^{n-3}$

$$\begin{array}{ccccccc}
A_{1,i+.5} & \longrightarrow & A_{1,i+1} & \longrightarrow & A_{1,i+1.5} & \longrightarrow & \Sigma A_{1,i+.5} \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
A_{2,i+.5} & \longrightarrow & A_{2,i+1} & \longrightarrow & A_{2,i+1.5} & \longrightarrow & \Sigma A_{2,i+.5} \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
A_{3,i+.5} & \longrightarrow & A_{3,i+1} & \longrightarrow & A_{3,i+1.5} & \longrightarrow & \Sigma A_{3,i+.5} \\
\vdots & & \vdots & & \vdots & & \vdots \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
A_{n,i+.5} & \longrightarrow & A_{n,i+1} & \longrightarrow & A_{n,i+1.5} & \longrightarrow & \Sigma A_{n,i+.5} \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
\Sigma^{n-2} A_{1,i+.5} & \longrightarrow & \Sigma^{n-2} A_{1,i+1} & \longrightarrow & \Sigma^{n-2} A_{1,i+1.5} & \longrightarrow & \Sigma^{n-1} A_{1,i+.5}.
\end{array}$$

Ω_i

where Ω_i represents the square

$$\Sigma^{n-3} \Omega_{1i} \circ_v \Sigma^{n-4} \Omega_{2i} \circ_v \dots \circ_v \Sigma \Omega_{n-3,i} \circ_v \Omega_{n-2,i}.$$

Note that Ω_i is a composition of $n-2$ anticommutative squares, so Ω_i commutes if n is even and anticommutes if n is odd. Note also that the first three columns are n -angles in the cluster tilting subcategory, while the first n rows of Γ_i are distinguished triangles in the ambient triangulated category.

We now essentially repeat this procedure in the horizontal direction, grafting Γ_i and Γ_{i+1} for $i \in \{1, \dots, n-3\}$. This means that we notice that the third column of Γ_1 is the first column of Γ_2 and the suspension of the third column of Γ_1 is the last column of Γ_2 , and so a similar grafting procedure takes place to create the $(n+1) \times 5$ diagram

$$\begin{array}{ccccccccc}
A_{1,i+.5} & \longrightarrow & A_{1,i+1} & \longrightarrow & A_{1,i+2} & \longrightarrow & A_{1,i+2.5} & \longrightarrow & \Sigma^2 A_{1,i+.5} \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
A_{2,i+.5} & \longrightarrow & A_{2,i+1} & \longrightarrow & A_{2,i+2} & \longrightarrow & A_{2,i+2.5} & \longrightarrow & \Sigma^2 A_{2,i+.5} \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
A_{3,i+.5} & \longrightarrow & A_{3,i+1} & \longrightarrow & A_{3,i+2} & \longrightarrow & A_{3,i+2.5} & \longrightarrow & \Sigma^2 A_{3,i+.5} \\
\vdots & & \vdots & & \vdots & & \vdots & & \vdots \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & & \downarrow \\
A_{n,i+.5} & \longrightarrow & A_{n,i+1} & \longrightarrow & A_{n,i+2} & \longrightarrow & A_{n,i+2.5} & \longrightarrow & \Sigma^2 A_{n,i+.5} \\
\downarrow & & \downarrow & & \downarrow & & \downarrow & \xrightarrow{\Sigma\Omega_1 \circ_h \Omega_2} & \downarrow \\
\Sigma^{n-2} A_{1,i+.5} & \rightarrow & \Sigma^{n-2} A_{1,i+1} & \rightarrow & \Sigma^{n-2} A_{1,i+2} & \rightarrow & \Sigma^{n-2} A_{1,i+2.5} & \rightarrow & \Sigma^n A_{1,i+.5}.
\end{array}$$

which we will denote by Γ_1^1 . Note that either both Ω_1 and Ω_2 commute or both anticommute, so the enveloping square, which we denote as the horizontal composition $\Sigma\Omega_1 \circ_h \Omega_2$, commutes regardless of the parity of n . Continuing on in the same fashion to before, we end with the $(n+1) \times (n+1)$ diagram

$$\begin{array}{ccccccccccc}
A_{11} & \longrightarrow & A_{12} & \longrightarrow & A_{13} & \longrightarrow & \cdots & \longrightarrow & A_{1n} & \longrightarrow & \Sigma^{n-2} A_{11} \\
\downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow \\
A_{21} & \longrightarrow & A_{22} & \longrightarrow & A_{23} & \longrightarrow & \cdots & \longrightarrow & A_{2n} & \longrightarrow & \Sigma^{n-2} A_{21} \\
\downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow \\
A_{31} & \longrightarrow & A_{32} & \longrightarrow & A_{33} & \longrightarrow & \cdots & \longrightarrow & A_{3n} & \longrightarrow & \Sigma^{n-2} A_{31} \\
\vdots & & \vdots & & \vdots & & & & \vdots & & \vdots \\
\downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow \\
A_{n1} & \longrightarrow & A_{n2} & \longrightarrow & A_{n3} & \longrightarrow & \cdots & \longrightarrow & A_{n,n} & \longrightarrow & \Sigma^{n-2} A_{n1} \\
\downarrow & & \downarrow & & \downarrow & & & & \downarrow & \xrightarrow{\Omega} & \downarrow \\
\Sigma^{n-2} A_{11} & \rightarrow & \Sigma^{n-2} A_{12} & \rightarrow & \Sigma^{n-2} A_{13} & \rightarrow & \cdots & \rightarrow & \Sigma^{n-2} A_{1n} & \rightarrow & \Sigma^{2(n-2)} A_{11},
\end{array}$$

denoted $\Gamma := \Gamma_1^{n-3}$, such that the first n rows and columns are all n -angles and such that the upper most $2 \times (n+1)$ diagram is the morphism we started with. Note, lastly, that the square Ω represents the square

$$\Sigma^{n-3}\Omega_1 \circ_h \Sigma^{n-4}\Omega_2 \circ_h \cdots \circ_h \Sigma\Omega_{n-3} \circ_h \Omega_{n-2}.$$

which is a composition of $n-2$ squares that either all commute or all anticommute. As such, Ω commutes exactly when n is even and anticommutes when n is odd. \square

4.2 Exotic n -angulated categories

For this section, let R be a commutative local ring with principal non-zero maximal ideal $\mathfrak{m} = (p)$ satisfying $\mathfrak{m}^2 = 0$. It was originally proved in [3] that the category of finitely generated free R -modules can be equipped with an n -angulated structure for $n \geq 3$ if n is even, and also for n odd if $2p = 0$. We state this as Theorem 4.10. We then provide an example of a morphism of n -angles which is not middling good. First we need to recall a few definitions.

Definition 4.8 Let \mathcal{C} be an n -angulated category with suspension Σ and let

$$A_1 \xrightarrow{\alpha_1} A_2 \xrightarrow{\alpha_2} A_3 \xrightarrow{\alpha_3} \cdots \xrightarrow{\alpha_{n-1}} A_n \xrightarrow{\alpha_n} \Sigma A_1$$

be an n - Σ -sequence in \mathcal{C} . We say that A_\bullet is a *candidate n -angle* if $(\Sigma\alpha_1)\alpha_n = 0$ and $\alpha_{i+1}\alpha_i = 0$ for $1 \leq i \leq n-1$.

Definition 4.9 An n - Σ -sequences A_\bullet is *contractible* if there exists maps $\Theta_i: A_{i+1} \rightarrow A_i$ for $1 \leq i \leq n-1$ and $\Theta_n: \Sigma A_1 \rightarrow A_n$ such that

$$1_{A_i} = \Theta_i\alpha_i + \alpha_{i-1}\Theta_{i-1} \quad \text{for } 2 \leq i \leq n$$

and

$$1_{\Sigma A_1} = (\Sigma\Theta_1)(\Sigma\alpha_1) + \alpha_n\Theta_n.$$

We now have the vocabulary to state the theorem.

Theorem 4.10 (Bergh–Jasso–Thaule). *Let \mathcal{C} be the category of finitely generated free R -modules, let $n \geq 3$ be an integer, and let $\Sigma: \mathcal{C} \rightarrow \mathcal{C}$ be the identity functor. Furthermore, let \mathcal{N} be the collection of all n - Σ -sequences isomorphic to a direct sum $C_\bullet \oplus F(p)_\bullet$, where C_\bullet is a contractible candidate n -angle, F is a free module in \mathcal{C} , and $F(p)_\bullet$ is the n - Σ -sequence*

$$F \xrightarrow{p} F \xrightarrow{p} \cdots \xrightarrow{p} F \xrightarrow{p} \Sigma F.$$

Then $(\mathcal{C}, \Sigma, \mathcal{N})$ is an n -angulated category whenever n is even, or when n is odd and $2p = 0$.

Given such an n -angulated category, we claim that there is a morphism of n -angles which is not middling good. To this end, we examine n -angles with base $p: R \rightarrow R$ and $0: R \rightarrow R$.

Lemma 4.11 *Let A_\bullet be an n -angle in \mathcal{C} with base $p: R \rightarrow R$, i.e.,*

$$A_\bullet: R \xrightarrow{p} R \xrightarrow{\alpha_2} A_3 \xrightarrow{\alpha_3} \cdots \xrightarrow{\alpha_{n-1}} A_n \xrightarrow{\alpha_n} \Sigma R.$$

Then $R(p)_\bullet$ is a summand of A_\bullet . Similarly, if A_\bullet has base $0: R \rightarrow R$, the n -angle $(\Gamma R)_\bullet[1] \oplus (\Gamma R)_\bullet[-1]$ is a summand of A_\bullet .

Proof. Suppose A_\bullet has base $p: R \rightarrow R$. By (N3), we have weak isomorphisms

$$\begin{array}{ccccccccccccccc} R & \xrightarrow{p} & R & \xrightarrow{p} & R & \xrightarrow{p} & R & \xrightarrow{p} & \cdots & \xrightarrow{p} & R & \xrightarrow{p} & \Sigma R \\ \parallel & & \parallel & & \downarrow i_3 & & \downarrow i_4 & & & & \downarrow i_n & & \parallel \\ R & \xrightarrow{p} & R & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & A_4 & \xrightarrow{\alpha_4} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma R \\ \parallel & & \parallel & & \downarrow r_3 & & \downarrow r_4 & & & & \downarrow r_n & & \parallel \\ R & \xrightarrow{p} & R & \xrightarrow{p} & R & \xrightarrow{p} & R & \xrightarrow{p} & \cdots & \xrightarrow{p} & R & \xrightarrow{p} & \Sigma R. \end{array}$$

Every R -module map $R \rightarrow R$ can be interpreted as multiplication by an element in R , so we may consider $r_k i_k$ as such an element. Since R is a local ring, either $r_k i_k \in \mathfrak{m}$ or $r_k i_k$ is invertible. By commutativity of R , we see that

$$r_k i_k p = p r_{k-1} i_{k-1} = r_{k-1} i_{k-1} p = p r_{k-2} i_{k-2} = \cdots = p r_3 i_3 = p, \quad (4.3)$$

so if $r_k i_k \in \mathfrak{m}$, this would imply that

$$p = r_k i_k p = 0$$

which is a contradiction. Therefore $r_k i_k$ must be invertible for all k . Now replace $r_k: A_k \rightarrow R$ by $(r_k i_k)^{-1} r_k: A_k \rightarrow R$ and note that

$$(1, 1, (r_3 i_3)^{-1} r_3, \dots, (r_n i_n)^{-1} r_n)$$

is a morphism of n -angles, since

$$(r_k i_k)^{-1} p = p (r_{k-1} i_{k-1})^{-1}$$

by (4.3) and commutativity of R , and therefore

$$(r_k i_k)^{-1} r_k \alpha_k = (r_k i_k)^{-1} p r_{k-1} = p (r_{k-1} i_{k-1})^{-1} r_{k-1}.$$

Then we have a retraction of A_\bullet onto $R(p)_\bullet$ and thus $R(p)_\bullet$ is a direct summand of A_\bullet .

Now suppose A_\bullet has base $0: R \rightarrow R$. Again we have weak isomorphisms

$$\begin{array}{ccccccccccccccc} R & \xrightarrow{0} & R & \xrightarrow{1} & R & \longrightarrow & 0 & \longrightarrow & \cdots & \longrightarrow & \Sigma R & \xrightarrow{(-1)^n} & \Sigma R \\ \parallel & & \parallel & & \downarrow i_3 & & \downarrow & & & & \downarrow i_n & & \parallel \\ R & \xrightarrow{0} & R & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\alpha_3} & A_4 & \xrightarrow{\alpha_4} & \cdots & \xrightarrow{\alpha_{n-1}} & A_n & \xrightarrow{\alpha_n} & \Sigma R \\ \parallel & & \parallel & & \downarrow r_3 & & \downarrow & & & & \downarrow r_n & & \parallel \\ R & \xrightarrow{0} & R & \xrightarrow{1} & R & \longrightarrow & 0 & \longrightarrow & \cdots & \longrightarrow & \Sigma R & \xrightarrow{(-1)^n} & \Sigma R. \end{array}$$

and in this case it is straightforward to see that $r_3 i_3 = 1_R$ and $r_n i_n = 1_R$. Thus, we have a retraction of A_\bullet onto $(\Gamma R)_\bullet[1] \oplus (\Gamma R)_\bullet[-1]$. \square

In the first case, replacing A_3 by $R \oplus M$ for some R -module M , we see that α_2 is replaced by $\begin{pmatrix} p \\ 0 \end{pmatrix}$ since the diagram

$$\begin{array}{ccccc} & & R & & \\ & \nearrow p & \downarrow i_3 & \searrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} & \\ R & \xrightarrow{\alpha_2} & A_3 & \xrightarrow{\cong} & R \oplus M \end{array}$$

commutes. We may thus assume that any n -angle with base $p: R \rightarrow R$ is immediately followed by the morphism $\begin{pmatrix} p \\ 0 \end{pmatrix}: R \rightarrow R \oplus M$ for some module $M \in \mathcal{C}$. Similarly, we may assume that any n -angle with base $0: R \rightarrow R$ is immediately followed by the morphism $\begin{pmatrix} 1 \\ 0 \end{pmatrix}: R \rightarrow R \oplus M$ for some module $M \in \mathcal{C}$.

Example 4.12. Consider the morphism

$$\begin{array}{cccccccccccc} R & \xrightarrow{p} & R & \xrightarrow{p} & R & \xrightarrow{p} & \cdots & \xrightarrow{p} & R & \xrightarrow{p} & R & \xrightarrow{p} & \Sigma R \\ \downarrow 0 & & \downarrow 0 & & \downarrow 0 & & & & \downarrow 0 & & \downarrow p & & \downarrow 0 \\ R & \xrightarrow{p} & R & \xrightarrow{p} & R & \xrightarrow{p} & \cdots & \xrightarrow{p} & R & \xrightarrow{p} & R & \xrightarrow{p} & \Sigma R \end{array}$$

that is 0 componentwise except for on the last object, on which it is p . Suppose this can be extended to an $(n+1) \times (n+1)$ -diagram where the first n rows and columns are all n -angles. Then we may apply Lemma 4.11 and assume it will have the form

$$\begin{array}{cccccccccccc} R & \xrightarrow{p} & R & \xrightarrow{p} & R & \xrightarrow{p} & \cdots & \xrightarrow{p} & R & \xrightarrow{p} & R & \xrightarrow{p} & \Sigma R \\ \downarrow 0 & & \downarrow 0 & & \downarrow 0 & & & & \downarrow 0 & & \downarrow p & & \downarrow 0 \\ R & \xrightarrow{p} & R & \xrightarrow{p} & R & \xrightarrow{p} & \cdots & \xrightarrow{p} & R & \xrightarrow{p} & R & \xrightarrow{p} & \Sigma R \\ \downarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} & & \downarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} & & \downarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} & & & & \downarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} & & \downarrow \begin{pmatrix} p \\ 0 \end{pmatrix} & & \downarrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\ R \oplus M_1 & \xrightarrow{\gamma_1} & R \oplus M_2 & \xrightarrow{\gamma_2} & R \oplus M_3 & \xrightarrow{\gamma_3} & \cdots & \xrightarrow{\gamma_{n-2}} & R \oplus M_{n-1} & \xrightarrow{\gamma_{n-1}} & R \oplus M_n & \xrightarrow{\gamma_n} & \Sigma R \oplus \Sigma M_1 \\ \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow \\ A_{41} & \longrightarrow & A_{42} & \longrightarrow & A_{43} & \longrightarrow & \cdots & \longrightarrow & A_{4,n-1} & \longrightarrow & A_{4n} & \longrightarrow & \Sigma A_{41} \\ \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow \\ \vdots & & \vdots & & \vdots & & & & \vdots & & \vdots & & \vdots \\ \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow \\ A_{n-1,1} & \longrightarrow & A_{n-1,2} & \longrightarrow & A_{n-1,3} & \longrightarrow & \cdots & \longrightarrow & A_{n-1,n-1} & \longrightarrow & A_{n-1,n} & \longrightarrow & \Sigma A_{n-1,1} \\ \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \downarrow & & \downarrow \\ \Sigma R & \longrightarrow & \Sigma R & \longrightarrow & \Sigma R & \longrightarrow & \cdots & \longrightarrow & \Sigma R & \longrightarrow & \Sigma R & \xrightarrow{(-1)^n} & \Sigma^2 R \end{array}$$

In particular, the third row must be an n -angle. By commutativity of the diagram, we can deduce that for $i = 1, \dots, n-2$ we have

$$\gamma_i = \begin{pmatrix} p & * \\ 0 & * \end{pmatrix}$$

where $*$ indicates an unknown map, and we can further deduce that

$$\gamma_{n-1} = \begin{pmatrix} 0 & * \\ 0 & * \end{pmatrix}.$$

Suppose now that

$$\gamma_n = \begin{pmatrix} \mu_1 & * \\ \mu_2 & * \end{pmatrix}.$$

Then commutativity of the diagram further implies that

$$\mu_1 p = p \quad \text{and} \quad \mu_2 p = 0$$

which, as noted earlier, implies that $\mu_1: R \rightarrow R$ is an isomorphism because R is a local ring with maximal ideal \mathfrak{m} satisfying $\mathfrak{m}^2 = 0$. Therefore, if the original morphism of n -angles extends to a $(n+1) \times (n+1)$ -diagram as claimed, we get an n -angle

$$R \oplus M_1 \xrightarrow{\begin{pmatrix} p & * \\ 0 & * \end{pmatrix}} R \oplus M_2 \xrightarrow{\begin{pmatrix} p & * \\ 0 & * \end{pmatrix}} \cdots \xrightarrow{\begin{pmatrix} p & * \\ 0 & * \end{pmatrix}} R \oplus M_{n-1} \xrightarrow{\begin{pmatrix} 0 & * \\ 0 & * \end{pmatrix}} R \oplus M_n \xrightarrow{\begin{pmatrix} \mu_1 & * \\ \mu_2 & * \end{pmatrix}} \Sigma R \oplus \Sigma M_1$$

and thus

$$\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix} = (-1)^n \begin{pmatrix} \mu_1 & * \\ \mu_2 & * \end{pmatrix} \begin{pmatrix} p & * \\ 0 & * \end{pmatrix} = (-1)^n \begin{pmatrix} \mu_1 p & * \\ 0 & * \end{pmatrix}.$$

But this implies that $\mu_1 p = 0$ which is a contradiction. Therefore, the morphism is not middling good. \diamond

References

- [1] Emilie Arentz-Hansen, Petter Andreas Bergh, and Marius Thaule. ‘The morphism axiom for N -angulated categories’. In: *Theory Appl. Categ.* 31 (2016), Paper No. 18, 477–483.
- [2] A. A. Beilinson, J. Bernstein, and P. Deligne. ‘Faisceaux pervers’. In: *Analysis and topology on singular spaces, I (Luminy, 1981)*. Vol. 100. Astérisque. Soc. Math. France, Paris, 1982, pp. 5–171.
- [3] Petter Andreas Bergh, Gustavo Jasso, and Marius Thaule. ‘Higher n -angulations from local rings’. In: *J. Lond. Math. Soc. (2)* 93.1 (2016), pp. 123–142. ISSN: 0024-6107. DOI: 10.1112/jlms/jdv064. URL: <https://doi.org/10.1112/jlms/jdv064>.
- [4] Petter Andreas Bergh and Marius Thaule. ‘The axioms for n -angulated categories’. In: *Algebr. Geom. Topol.* 13.4 (2013), pp. 2405–2428. ISSN: 1472-2747. DOI: 10.2140/agt.2013.13.2405. URL: <https://doi.org/10.2140/agt.2013.13.2405>.
- [5] J. Daniel Christensen. ‘Ideals in triangulated categories: phantoms, ghosts and skeleta’. In: *Adv. Math.* 136.2 (1998), pp. 284–339. ISSN: 0001-8708. DOI: 10.1006/aima.1998.1735. URL: <https://doi.org/10.1006/aima.1998.1735>.

- [6] J. Daniel Christensen and Martin Frankland. ‘On good morphisms of exact triangles’. In: *J. Pure Appl. Algebra* 226.3 (2022), Paper No. 106846, 33. ISSN: 0022-4049. DOI: 10.1016/j.jpaa.2021.106846. URL: <https://doi.org/10.1016/j.jpaa.2021.106846>.
- [7] J. Daniel Christensen and Martin Frankland. ‘The Moss pairing for the Adams spectral sequence in triangulated categories’. In preparation.
- [8] Christof Geiss, Bernhard Keller, and Steffen Oppermann. ‘ n -angulated categories’. In: *J. Reine Angew. Math.* 675 (2013), pp. 101–120. ISSN: 0075-4102. DOI: 10.1515/crelle.2011.177. URL: <https://doi.org/10.1515/crelle.2011.177>.
- [9] Gustavo Jasso. ‘ n -abelian and n -exact categories’. In: *Math. Z.* 283.3-4 (2016), pp. 703–759. ISSN: 0025-5874. DOI: 10.1007/s00209-016-1619-8. URL: <https://doi.org/10.1007/s00209-016-1619-8>.
- [10] Amnon Neeman. ‘Some new axioms for triangulated categories’. In: *J. Algebra* 139.1 (1991), pp. 221–255. ISSN: 0021-8693. DOI: 10.1016/0021-8693(91)90292-G. URL: [https://doi.org/10.1016/0021-8693\(91\)90292-G](https://doi.org/10.1016/0021-8693(91)90292-G).
- [11] Amnon Neeman. *Triangulated categories*. Vol. 148. Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 2001, pp. viii+449. ISBN: 0-691-08685-0; 0-691-08686-9. DOI: 10.1515/9781400837212. URL: <https://doi.org/10.1515/9781400837212>.

DEPARTMENT OF MATHEMATICAL SCIENCES, NTNU
N-7491 TRONDHEIM, NORWAY

`sebastian.martensen@ntnu.no`

ORCID iD: 0000-0002-4477-8856