

Composite scalar boson mass dependence on the constituent mass anomalous dimension

A. Doff^a and A. A. Natale^b

^a*Universidade Tecnológica Federal do Paraná - UTFPR - DAFIS,*

R. Doutor Washington Subtil Chueire,

330 - Jardim Carvalho,

84017-220 Ponta Grossa, PR, Brazil

agomes@utfpr.edu.br

^b*Instituto de Física Teórica - UNESP,*

Rua Dr. Bento T. Ferraz, 271,

Bloco II, 01140-070,

São Paulo, SP, Brazil

natale@ift.unesp.br

Abstract

We perform a Bethe-Salpeter equation (BSE) evaluation of composite scalar boson masses in order to verify how these masses can be smaller than the composition scale. The calculation is developed with a constituent self-energy dependent on its mass anomalous dimension (γ), and we obtain a relation showing how the scalar mass decreases as γ is increased. We also discuss how fermionic corrections to the BSE kernel shall decrease the scalar mass, whose effect can be as important as the one of a large γ . An estimate of the top quark loop effect that must appear in the BSE calculation gives a lower bound on the composite scalar mass.

I. INTRODUCTION

The discovery of the Higgs boson at the LHC [1, 2] completed the Standard Model (SM), where a scalar boson sector is present as proposed long ago [3]. In many extensions of this model this scalar sector is even larger than the SM one, although experimental signals of new particles belonging to this sector are still missing. Furthermore, there are theoretical shortcomings about this scalar sector [4, 5].

The absence of signals of a large scalar boson sector in the experimental data, as well as a possible explanation of a light Higgs boson has been discussed recently in Ref.[6, 7]. Composite scalar bosons also appear in the context of Technicolor theories (TC) [8–10], which usually have a composition scale of order of $\Lambda_H \geq 1\text{TeV}$.

The construction of a realistic Technicolor model may indeed be a very precise engineering problem. In order to get around this problem different models with large mass anomalous dimension (γ) emerged, as walking technicolor[11–16] and gauged Nambu-Jona-Lasinio models[17–23]. A light composite scalar boson may be generated when the strong interaction theory (or TC) has large mass anomalous dimension (γ) as proposed by Holdom[24]. A discussion of how these scenarios can lead a light composite scalar boson became clear in the Refs.[25]. Based on these scenarios, calculations involving effective Higgs Lagrangians have led to different predictions regarding to composite scalar bosons masses.

In this case the self-energy of the new fermions (or technifermions) responsible for the composite states is characterized by a large mass anomalous dimension, resulting in mass diagrams whose calculation do not scale with the naive dimensions. This self-energy at large momenta is proportional to

$$\Sigma_C(p^2) \propto \frac{\mu_C^3}{p^2} (p^2/\mu_C^2)^{\gamma/2}. \quad (1)$$

where μ_C is the typical composition or dynamical mass scale and γ the mass anomalous dimension. It is now known that according to the mass anomalous dimension, this self-energy may vary asymptotically from a $1/p^2$ behavior, up to a very slowly decreasing logarithmic behavior with the momentum [23, 26]. It is usually assumed that large γ values appear in gauge theories with higher fermionic representations or with a large number of fermions. However, Eq.(1) can be quite modified just coupling two strongly interacting theories [26, 27].

A light composite scalar boson appears naturally when its mass is calculated with the help of Eq.(1) (see, for instance, Ref.[25]). It is interesting to see that the problem of

understanding a possible light composite Higgs boson could also be related to the case of the sigma meson. The sigma meson is the QCD scalar composite now known as $f_0(500)$. A standard Bethe-Salpeter equation (BSE) calculation gives $m_\sigma = 670\text{MeV}$ [28], which is larger than its experimental value. The detailed work of Ref.[28] deals with possible contributions that may lower the estimate of this mass. A composite $J = 0$ state may have many contributions to its mass. In the case of the sigma meson it is not even clear how much of its composition is due to different quarks, even more the amount of its mass that is due to gluons, although it is already a puzzle the fact that a simple BSE mass estimate gives a result larger than the experimental value. Therefore, it is natural to think what a calculation similar to the one of Ref.[28] would teach us about the composite Higgs mass.

As far as we know there are not in the literature detailed calculations of the composite Higgs boson mass using BSE, particularly looking for effects of different γ values or the effect of massive fermions contribution to the BSE kernel. In this work we calculate the scalar composite mass with the help of Bethe-Salpeter equations, obtaining a relation between the scalar mass and γ . We present a fit of such relation expecting that it could be tested by other methods. This calculation is shown in Section II. In Section III we perform a simple order of magnitude estimate of fermionic loop effects that contribute to the BSE kernel and can lower the scalar mass value. It is important to remark that most studies dealing with the possibility of a light composite scalar boson are based on the calculation of Schwinger-Dyson equations and related to a near conformal behavior of the theory [29]. The advantage of the BSE approach is that it takes into account all the possible bound states that contribute to the scalar mass calculation. A light scalar comes out not only due to a conformal behavior, but it does emerge when some bound states contribute negatively to the scalar BSE kernel. We verify that the effect of a large fermionic mass, particularly a bound state formed by the top quark, affect the Higgs boson mass calculation and cannot be neglected. A quite simple estimate of the top quark contribution to the BSE kernel leads to a lower bound on the composite Higgs mass of the order of 123.3GeV . As far as we know there are not in the literature detailed calculations of the composite Higgs boson mass using BSE. In this work we generalize the results obtained in [38], improving the work of Ref.[31], which is applied only to QCD bound states, introducing fermionic propagators characterized by Eq.(1) which is dependent on different anomalous mass dimension, and investigating the effect of TC bound states in order to obtain a light composite scalar mass. Section IV contains our conclusions.

II. THE SCALAR MASS: BSE AND MASS ANOMALOUS DIMENSION

The scalar boson mass can be calculated using BSE once we specify the propagators and vertices of the strong interaction that binds such scalar boson. For the scalar case the Bethe-Salpeter equation has the form[30–32]

$$\begin{aligned}\chi(p, q) &= -i \int \frac{d^4 k}{(2\pi)^4} S(q + \alpha p) K_{\rho\nu}(p, k, q) S(q - \beta p), \\ K_{\rho\nu}(p, k, q) &= \gamma_\rho \chi(p, k) \gamma_\nu G_{\rho\nu}(k - q),\end{aligned}\tag{2}$$

where $\alpha + \beta = 1$ (α and β characterize the fraction of momentum carried by the constituents), although the result is not dependent on these quantities[31]. As a first approximation we shall choose $\alpha = \beta = 1/2$. $G_{\rho\nu}$ is the gauge boson propagator in the Landau gauge. There are different models for this quantity which will be discussed ahead.

The fermion propagator is given by

$$S^{-1}(q) = \not{q} A(q^2) - B(q^2),\tag{3}$$

From now on we shall assume $A(q^2) = 1$ and $B(q^2) = \Sigma(q^2)$. Usually in this type of calculation the self-energy $\Sigma(q^2)$ is obtained from the numerical solution of the Schwinger-Dyson equation (SDE) for the fermionic propagator. In order to simplify the calculation this self-energy is going to be given by one ansatz that is a function of the mass anomalous dimension γ .

We recall that $\Sigma(q^2)$ at large momenta in QCD (or any asymptotically free non-Abelian gauge theory) is given by [33]

$$\Sigma(q^2) \propto \frac{\langle \bar{\psi}\psi \rangle_\mu}{q^2} \left(\frac{q^2}{\mu^2} \right)^{\gamma/2},\tag{4}$$

assuming for the fermionic condensate $\langle \bar{\psi}\psi \rangle_\mu \approx \mu^3$ where μ is the dynamically generated mass.

The ansatz that we assume for the self-energy has the form

$$\Sigma(q^2) = \frac{\mu^3}{q^2 + \mu^2} \left(\frac{q^2 + \mu^2}{\mu^2} \right)^\kappa\tag{5}$$

where

$$\kappa = \gamma/2.\tag{6}$$

Eq.(5) behaves in the infrared region as a mass μ and decays at large momenta as prescribed in Eq.(4), mapping the SDE in the full Euclidean space, and will allow us to obtain BSE solutions for the scalar mass as a function of γ . Note that κ is just a parameter that when $\kappa = 0$ the self-energy behaves asymptotically as $1/p^2$, and when $\kappa \rightarrow 1$ Eq.(5) behaves like

$$\Sigma(q^2) \approx \mu [1 + \delta_1 \ln [(q^2 + \mu^2)/\mu^2]]^{-\delta_2}, \quad (7)$$

where δ_1 and δ_2 are obtained from γ when expanded as a function of the running coupling $g^2(q^2)$.

The BSE solution appear as an eigenvalue problem for $p^2 = M^2$, where M is the bound state mass. The variables are p, q, k . k is integrated and we remain with a equation in q that will have a solution for $p^2 = M^2$. In the Eq.(2) χ can be projected into four coupled homogeneous integral equations, that implies for projection of the scalar component

$$\chi(p, q) = \chi_{S0} + \not{p}\chi_{S1} + \not{q}\chi_{S2} + [\not{p}, \not{q}]\chi_{S3}, \quad (8)$$

which are functions of p^2, q^2 and $p \cdot q = pq \cos \theta$.

It is possible to expand $\chi(p, q)$ in terms of Tschebyshev polynomials, and these equations can be truncated at a given order determined by the relative size of the next-order functions. In accordance with Ref.[30], a satisfactory solution can be obtained by keeping only some terms, like $\chi_{S(0,1)}^{(0)}, \chi_{S(1)}^{(0,1)}, \chi_{S(2)}^{(0,1)}$.

To set up the problem we follow closely the work of Ref. [31], where the BSE was solved for the scalar boson constituents with masses m_a and m_b , and for simplicity we assume

$$m_a = m_b = m = \Sigma(x + \frac{1}{4}p^2), \quad (9)$$

where it was assumed $\alpha = \beta$ (each constituent carries half of the momentum), and $x = q^2/\Lambda^2$, where $\Lambda = \Lambda_{QCD}$ (or $\Lambda = \Lambda_H$, the TC mass scale) the characteristic mass scale of the bounding force.

The different components of Eq.(8), $\chi_{S(i)}^{(0,1)}$ for $i = 0..3$, are given by

$$\chi_{S0}^{(0)} = 2[(x - \frac{1}{4}p^2 - m^2)J_1]I_{S0} + \Delta\chi_{S0}^{(0)}, \quad (10)$$

with

$$I_{S0} = \frac{2}{3\pi} \int dy y \chi_{S0}^{(0)} K_1, \quad (11)$$

where $y = k^2/\Lambda^2$ and

$$K_1(x, y) = \frac{3}{16\pi^2} \int d\theta \sin^2\theta G(x, y, \cos\theta), \quad (12)$$

$$J_1 = \frac{2}{\pi} \int_0^\pi d\theta \frac{\sin^2\theta}{D(p^2, q^2, pq\cos\theta)} \quad (13)$$

and in the Eq.(13)

$$D(p^2, q^2, pq\cos\theta) = \{(q + \frac{1}{2}p)^2 + m^2\}\{(q - \frac{1}{2}p)^2 + m^2\}. \quad (14)$$

In the above equation we can expand $(q + \frac{1}{2}p)^2 + m^2$ and $(q - \frac{1}{2}p)^2 + m^2$ in Taylor series. Just keeping the first-order derivative terms for m , we have that the function J_1 can be written in the form

$$J_1 = \frac{2}{c_1 c_4 + c_2 c_3} \left[\frac{c_2}{D_1} + \frac{c_4}{D_2} + d_1 \left(\frac{c_1}{D_1} - \frac{c_3}{D_2} \right) \right] \quad (15)$$

where, in our approximation, we have

$$c_1 = c_3 = x + \frac{1}{4}p^2 + m^2$$

$$c_2 = c_4 = 1 + 2mm'$$

and m' is the derivative of m with respect to the momentum. In addition, as a consequence of $\alpha = \beta$ we obtain

$$d_1 = 0$$

$$D_1 = D_2 = c_1 + \sqrt{c_1^2 - p^2 x c_2^2}.$$

In Eq.(10), the term $\Delta\chi_{S0}^{(0)}$ represent corrections to the leading-order results of $\chi_{S0}^{(0)}$, that correspond to $\chi_{S1}^{(0,1)}$, $\chi_{S2}^{(0,1)}$ and $\chi_{S3}^{(0,1)}$. With the approximations considered here we obtain

$$\begin{aligned} \Delta\chi_{S0}^{(0)} = & -\frac{4}{3\pi} m J_1 \int dy y \chi_{S2}^0 \sqrt{xy} (3K_6 - 2\sqrt{xy} K_3) - \\ & -\frac{2}{3\pi} p^2 (J_3 - J_1) \int dy y \chi_{S3}^0 \{2\sqrt{xy} K_6 - \frac{8}{3} xy K_3\} - \\ & -\frac{4}{\pi} \{(-x + \frac{1}{4}p^2 + m^2) J_2\} \int dy y \chi_{S0}^{(1)} \sqrt{\frac{y}{x}} K_6 - \\ & -\frac{2}{\pi} \{(-x + \frac{1}{4}p^2 + m^2) J_1\} \int dy y^2 \chi_{S0}^{(2)} \{\frac{4}{3} K_7 - K_1\}. \end{aligned} \quad (16)$$

In the Eq.(16), the lowest order terms $\chi_{S(1-3)}^{(0)}$ are given by

$$\chi_{S1}^{(0)} = 0 \quad (17)$$

$$\begin{aligned} \chi_{S2}^{(0)} = & -\frac{4}{\pi}mJ_1I_{S0} + \frac{2}{3\pi} \left(-x - \frac{1}{4}p^2 + m^2 \right) \times \\ & \int dy y \chi_{S2}^0 \left(3\sqrt{\frac{y}{x}}K_6 - 2yK_3 \right), \end{aligned} \quad (18)$$

and

$$\chi_{S3}^{(0)} = \frac{3}{2}J_1I_{S0}. \quad (19)$$

While the higher order terms are described by

$$\chi_{S0}^{(1)} = -6 \left(\left(-x + \frac{1}{4}p^2 + m^2 \right) \frac{J_2}{xp^2} \right) I_{S0}, \quad (20)$$

and

$$\chi_{S0}^{(2)} = -3 \left\{ \frac{1}{xp^2} \left(-x + \frac{1}{4}p^2 + m^2 \right) (4J_3 - J_1) \right\} I_{S0}. \quad (21)$$

Since we are dealing with a scalar boson case with equal mass constituents the equations of Ref.[31] are also simplified and we have:

$$J_2 = 0 \quad (22)$$

$$J_3 = \frac{1}{D_1^2} \quad (23)$$

where in the Eqs.(16-18)

$$K_3(x, y) = \frac{3}{16\pi^2} \int d\theta \frac{\sin^4\theta}{x + y - 2\sqrt{xy}\cos\theta} G(x, y, \cos\theta), \quad (24)$$

$$K_6(x, y) = \frac{3}{16\pi^2} \int d\theta \sin^2\theta \cos\theta G(x, y, \cos\theta), \quad (25)$$

$$K_7(x, y) = \frac{3}{16\pi^2} \int d\theta \sin^4\theta G(x, y, \cos\theta). \quad (26)$$

These equations can be solved making use of suitable expressions for the main Green functions (i.e. propagators and vertices). The gauge boson propagator is given by

$$G(k^2) = \frac{16\pi}{3} \left[\frac{\pi d}{k^2 \ln(x_0 + x)} \right] + G_{IR}(k^2), \quad (27)$$

where $G(x, y, \cos\theta) = \Lambda^2 G(k - q)^2$, and $G_{IR}(k^2)$ is an assumed form of the interaction at infrared momenta. In Ref.[31] this contribution was chosen to be of the form

$$G_{IR}(k^2) = \frac{16\pi}{3} a k^2 e^{-\frac{k^2}{\omega^2}}, \quad (28)$$

and in the gaussian ansatz for the $G_{IR}(k^2)$, $\omega \in [0.4, 0.6] GeV$ [34–36]. According to Ref.[31] the parameters used in the QCD case are given by

$$\begin{aligned} a &= (0.387 GeV)^{-4} \quad , \quad \omega = (0.510 GeV) \\ d &= 12/(33 - 2n_f) \quad , \quad n_f = 5 \\ \mu &\approx \Lambda_{QCD} = 0.228 GeV \quad , \quad x_0 = 10. \end{aligned} \quad (29)$$

More recent expressions for these Green functions were formulated in Refs.[34–36], where the two free parameters in Eq.(28), ω and $a = D$, are parameterized by $(\varsigma_g)^3 = D\omega = const$, and the fitted values of ς_g depend on the form that is assumed for the dressed-gluon quark vertex. Considering the recent results reported in Ref.[34], it is possible to verify that these choices do not modify substantially the numerical results.

The complete set of BS equations given by Eq.(2) was solved numerically by iteration starting with the inhomogeneous terms as input, where the function $\chi(p, 0) = \chi_{S0}^{(0)}$ is fixed to some arbitrarily chosen value, considering an interactive process in the equation below

$$\Delta(p^2) = \chi(p^2, 0) - \chi(p^2, q^2) \quad (30)$$

to find the eigenvalues $p^2 = M_S^2$, of eigenfunction $\chi(p^2, q^2)$. For a given value of p^2 , not an eigenvalue, $\Delta(p^2)$ is not zero, then after several interactions we can find p^2 such that we obtain $\Delta(p^2) = 0$, or BS wave function eigenvalues. Notice that a normalization procedure is also necessary because we cannot have arbitrary mass anomalous dimensions, as already pointed out in Refs.[38, 40].

Assuming Eqs.(8 - 28), in Fig.1 we present the results obtained for $\Delta(p^2)$, given by Eq.(30), considering the ansatz Eq.(5) for m . In this figure we normalize our results for M_S in terms of

$$M_S = 2\Lambda_{QCD},$$

associated to a negligible γ .

The choice of this normalization is based on the result described in Ref.[37], where Delbourgo and Scadron verified analytically with the help of the homogeneous Bethe-Salpeter




FIG. 1: In this figure we show the scalar masses $M_S(\gamma)$ obtained for $\Delta(p^2)$, given by Eq.(30), in the QCD case. The contextualization of the curves behavior are described in the text.

equation (BSE), that the sigma meson mass is given by $m_\sigma = 2\mu_{\text{dyn}}$. In this calculation it is assumed that the dynamically generated quark mass behaves (for large p^2) as $m_{\text{dyn}} \sim \frac{\mu_{\text{QCD}}^3}{q^2}$, which corresponds to the case where $\kappa = 0$ in the ansatz proposed in Eq.(5). In this way, based on this normalization, we can follow the behavior of how M_S resulting from Eq.(30) is influenced by γ .

In Fig.(1a) the black line corresponds exactly to the case where $\kappa = 0$, while the blue line to $\kappa = 0.2$. In Fig. (1b) we consider the cases where $\kappa = 0.3$ (red line) and $\kappa = 0.4$ (green line), the Fig.(1c) is a composition of the previous results, where we indicate for each curve the value assumed for κ . Note that in Fig1.(a-b), in the upper right corner, we describe M_S found in each case observing that for a given $z = p^2/\Lambda_{\text{QCD}}^2$ we have

$$M_S = \sqrt{z}\Lambda_{\text{QCD}}, \quad (31)$$

Finally, in Fig.(1d), we present the behavior of $M_S(\gamma)$ obtained for κ in the range $\in [0, 0.95]$.




FIG. 2: In this figure we show the scalar masses $M_S^{TC}(\gamma)$ obtained for $\Delta(p^2)$, given by Eq.(30), in the TC case. The contextualization of the curves behavior are described in the text.

We obtained a very simple fit to the data with $R^2 = 0.977$ which corresponds to

$$A(\gamma) = \frac{M_S(\gamma)}{\Lambda_{QCD}} = \frac{2.15}{(1 + \gamma/2)^{5.34}}. \quad (32)$$

It is clear that Fig.(1d) may be slightly dependent on the propagators and vertex that we assumed here. However, we expect that the behavior of this curve can be tested by other methods, and more importantly it shows how the scalar composite mass should behave as we vary the constituent mass anomalous dimension.

We can now focus on the Higgs boson case. In Fig.2 we extend the results obtained for QCD in the case of a $SU(3)$ TC model. As a first approach, since TC is based upon an analogy with the dynamics of QCD, we can use the equations obtained for QCD to determine, by appropriate rescaling, the behavior of $M_S^{TC}(\gamma)$. Hence, we can estimate ω_{TC} , a_{TC} from the QCD analogue using the following scaling relation

$$(\omega, a)_{TC} = \sqrt{\frac{N_{TC}}{3}} \frac{\Lambda_{TC}}{\Lambda_{QCD}} (\omega, a)_{QCD},$$

where $\Lambda_H = \Lambda_{TC} = 1TeV$. In this case the results for $M_S(\gamma) = M_S^{TC}(\gamma)$ follow from the normalization $M_S = 2\Lambda_{TC}$; and we include in Fig.(2d) the dot-dashed line in red, which corresponds to the observed Higgs boson mass for the purpose of comparison with the $M_S(\gamma)$ behavior. In the region where $\kappa \sim 0.8$, we recover the result obtained for the extreme walking behavior[38], where for a $SU(3)$ TC model in that reference, assuming for $\Sigma(p^2)_{TC}$ the behavior given by Eq.(7), we obtained $M_H \sim O(110)GeV$.

In Fig.(2d), we present the behavior for $M_S^{TC}(\gamma)$ obtained in the range $\kappa \in [0, 0.95]$. The fit obtained with $R^2 = 0.988$ corresponds to

$$A(\gamma) = \frac{M_S^{TC}(\gamma)}{\Lambda_{TC}} = \frac{2.1}{(1 + \gamma/2)^{5.12}}. \quad (33)$$

This result shows how the composite Higgs boson mass may vary with the constituent mass anomalous dimension. However, as we will discuss in the next section, it is not only the value of γ that modifies these estimates. Note that Eq.(33) do not differ appreciably from Eq.(32) due to the fact that we have chosen the TC gauge theory as a QCD rescaled version with self-energy given by ansatz, Eq.(5).

Notice that $\Delta(p^2)$ and the scalar masses in Fig. 1 and Fig. 2 decay faster with increasing gamma. The fact is that the BSE kernel is proportional to the fermionic and gauge boson propagators, and is an integration over these quantities as well over the coupling constants. The product of coupling and gauge boson propagator may be interpreted as the strength of the interaction, and the effect of larger (γ) values in the fermionic propagators act in the sense of diminish the interaction strength implying in smaller scalar masses, and similarly a change in the composite scalar wave functions.

III. SCALAR MASSES: THE EFFECT OF FERMIONS

In usual BSE calculations of QCD light hadronic states the effect of heavy quarks are not included. However, when performing a BSE calculation of a possible composite Higgs boson we cannot neglect the contribution of a top quark loop to the BSE kernel. There are two reasons for this; the strong coupling of the top quark to the scalar boson and the approximate values of the masses of these two particles. We are clearly interested in the case of a composite Higgs boson mass calculation, but we shall start with a simple discussion about the QCD scalar (the sigma meson), for which a more detailed calculation will be left

for a forthcoming work.

If we go back to the linear sigma model at constituent level we know that the sigma couples to fermions as

$$\mathcal{L}_\sigma = \lambda \bar{\Psi} \Psi \phi, \quad (34)$$

where ϕ stands for σ . This coupling imply that the ϕ mass obtains contributions from the BSE diagram shown in Fig.(3a), that comes with a negative sign due to the effect of a fermion loop. Eq.(34) also describe the Higgs Yukawa coupling to fermions. In particular, when we consider a composite Higgs boson the coupling to fermions is more sophisticated, we may even have fermionic contributions to the BSE like the one shown in Fig.(3b), involving the exchange of extended TC gauge bosons (ETC) [10]. However, as the gauge bosons of Fig.(3b) are very heavy, the vertex in that figure can be reduced to an effective vertex as shown in Fig.(3c) and the final BSE mass contribution can be reduced to the one of Fig.(3a).

The fermionic contribution to the BSE would be given by

$$\begin{aligned} \Pi(p^2) &= m_{f(t)}^2 \text{Tr} \int_0^\Lambda d^4 q \chi(q^2) \mathcal{F}(q, p, f(t)) \chi(q^2), \\ \mathcal{F}(q, p, f(t)) &= S_{f(t)}(p/2 + q) \chi(q^2) S_{f(t)}(p/2 - q) \end{aligned} \quad (35)$$

where the vertex of the BSE (due to a fermion f or to the top quark t) reads

$$m_{f(t)} \times \chi(p^2), \quad (36)$$

where we stress the effect of the large top quark mass in the calculation of the Higgs boson mass. A full calculation of the BSE including fermionic corrections with complete solutions of the Schwinger-Dyson equations for the self-energies is a lengthy work and is under study; it may affect the sigma as well as the Higgs boson mass estimate. In the case of the Higgs boson we can resort to a simple estimate of the loop of Fig.(3a), i.e. a correction of M_ϕ^2 represented by δM_ϕ^2 , is given by

$$i\delta M_\phi^2 = \frac{\lambda^2 N_f}{(2\pi)^4} \int d^4 q \frac{\Sigma^2(q^2)}{(q^2 - \Sigma^2(q^2))^2}, \quad (37)$$

where N_f is the number of fermions (f) in the loop, and the biggest effective coupling λ (when f= top quark) is given by

$$\lambda = m_t(\gamma)/v, \quad (38)$$

and v is the standard model vacuum expectation value ($v = 2M_W/g_w$).




FIG. 3: In Fig.(3b) we indicate the BSE diagram that should introduce fermionic corrections to M_ϕ^2 in the case of a composite Higgs boson. As the ETC gauge bosons depicted in this diagram are very heavy, the vertex appearing in Fig.(3b) can be reduced to the one of Fig.(3c). Therefore, Fig.(3a) is the result of Fig.(3b) when we assume the effective vertex, and it reduces the scalar boson mass.

Eq.(37) is enough to verify the order of magnitude that we shall obtain when solving the complete system of BSE for the scalar boson. Considering the anzats described by Eq.(5), in euclidean space we obtain the following expression for $\delta M_\phi^2(\gamma)$

$$\delta M_\phi^2(\gamma) = \frac{\lambda^2 N_f}{4\pi^2} f(\gamma) \Lambda_{TC}^2 \quad (39)$$

where

$$f(\gamma) = \frac{1-\gamma}{3-\gamma}. \quad (40)$$

The behavior of Eq.(40) with γ is an artifact of our approximation, since the γ running with momentum was not considered, and its value has to be bounded so that the scalar boson wave function is quadratically integrable [39, 40].

The contribution due to the fermion loop indicated in Fig.(3a) , particularly in the extreme walking behavior or massive top case for the self-energy (i.e. $\gamma \rightarrow 2$ (or $\kappa \rightarrow 1$)), tends

to decrease M_ϕ^2 according to Eq.(39), and in this case this contribution lowers the estimate of the composite scalar boson mass.

In Fig.(3c), the vertex λ can be approximately represented by

$$\lambda \approx \frac{g_w}{\pi} N_F \lambda_{ETC} \frac{\Lambda_{TC}}{M_W} \left(\frac{\Lambda_{TC}^2}{\Lambda_{ETC}^2} \right)^{1-\kappa}, \quad (41)$$

where N_F is the number of technifermions that couple to the fermions (f) in the loop, and we assumed the existence of an ETC gauge theory with coupling α_{ETC} . The effective charge, $\lambda_{ETC} = C_{ETC} \alpha_{ETC}$, involve the ETC coupling and the appropriate ETC Casimir operator eigenvalues C_{ETC} . The top quark makes the most significant contribution in the loop described in Fig.(3), as discussed in Refs.[41, 42] in the extreme walking behavior, its mass can approximately be expressed by $m_t(\gamma) \approx N_F \lambda_{ETC} \Lambda_{TC}$, so we can write the vertex λ as

$$\lambda \approx \frac{g_w}{\pi} \frac{m_t(\gamma)}{M_W} \left(\frac{\Lambda_{TC}^2}{\Lambda_{ETC}^2} \right)^{1-\kappa}. \quad (42)$$

In the limit when $\kappa \rightarrow 0$, we have

$$\delta M_\phi^2(0) \approx 0,$$

while in the limit when $\kappa \rightarrow 1$ (or $\gamma \rightarrow 2$), we recover the effective coupling of the top described in Eq.(38)

$$\lambda \propto \frac{g_w}{\pi} \frac{m_t(\gamma)}{M_W} \approx \frac{m_t(\gamma)}{v}$$

and we obtain

$$\begin{aligned} \delta M_\phi^2(2) &\approx -\frac{3\lambda^2}{4\pi^2} \Lambda_{TC}^2 \\ &\approx -\frac{3}{\pi^4} \left(\frac{m_t(2)}{v} \right)^2 \Lambda_{TC}^2. \end{aligned} \quad (43)$$

At this point we should highlight that in the parameterization of the estimate presented by Eq.(43), $m_t(2)$ is model dependent. In Fig.(3), the number of technifermions F (N_Q :techniquarks or N_L :technileptons) that generate the $m_t(2)$ mass depends on ETC interactions. However, we can consider as an illustrative example the model described in Ref.[42], where in Fig.(3) of that reference, we present the diagrams that contribute to $m_t(2)$ in that work. As a result of these contributions $m_t(2)$ was estimated to be on the order of $\sim 100 GeV$, what leads to

$$|\delta M_\phi(2)| \approx 70 GeV. \quad (44)$$

As we have seen, the determination of $m_t(2)$ is model dependent, however, assuming that it is possible to elaborate a more realistic ETC model, where in principle $m_t(2)$ can be of the same order of the observed top quark mass, the positivity condition of $M_S^2 > 0$, which is given by the smallest BSE solution ($M_S(2)$) minus the fermionic correction described by Eq.(43), leads to the following intriguing theoretical limit

$$\frac{M_S(2)}{\Lambda_{TC}} > \frac{\sqrt{3}}{\pi^2} \frac{m_t}{v}, \quad (45)$$

i.e., assuming $\Lambda_{TC} \approx 1\text{TeV}$ and use the known m_t and v values the bound of Eq.(45) is exactly of the order of the known Higgs boson mass, or $M_S(2) \geq 123.3\text{GeV}$.

Note that these are very rough estimates originated by the existence of radiative corrections due to TC and ETC as appear in Fig.(3). The effect of fermion loops inevitably decreases the scalar bound state mass. A full BSE calculation should also involve the dependence of all Green's functions on the mass anomalous dimensions and fermion masses of the scalar boson constituents. Actually, the result of this section can be seen only as a correction to the BSE result of the previous section, which is dependent on the ansatz proposed in Eq.(5), whereas a complete calculation should rely on a self-energy obtained from the Schwinger-Dyson equation, which is beyond the scope of the present work. It is also opportune to recall that even the sigma meson mass calculation may have corrections of similar type (generated by electroweak bosons exchange), that can lower the BSE evaluation of its mass.

IV. CONCLUSIONS

In the case of a possible composite scalar boson, we computed its mass using Bethe-Salpeter equations and assuming constituents of same mass. The calculation was performed with the help of a constituent self-energy dependent on the mass anomalous dimension. Our result indicates how the scalar masses, no matter we are talking about the sigma meson or the Higgs boson, can vary with the mass anomalous dimension as shown in Eqs.(32) and (33). We hope that this behavior can be tested by other methods.

In Section III we call attention to the fact that a full BSE calculation should include diagrams like the one of Fig.(3). The effect of such diagrams is to lower the scalar boson mass. As a simple estimate of this effect we have computed the fermionic contribution of

the radiative corrections induced by Fig.(3a). Of course, our calculation is very simple but it shows that this effect cannot be neglected. The bound of Eq.(45) is an example of the balance between the different contributions to scalar masses.

Our results using the Bethe-Salpeter equations show how scalar composite masses can be smaller than the composition scale, as long as we have large anomalous dimensions and the effect of fermions, like the top quark, included into the calculation. Actually, we may have a delicate balance between the mass anomalous dimension of the fermionic constituents and the contribution of fermions that contribute negatively to the Higgs boson mass.

If the Higgs boson is a composite particle, it is still possible that its constituents are bounded by a non-Abelian gauge strong interaction similar to QCD. In this case this new strong interaction dynamics can be scaled from the known QCD Green's functions, which nowadays are used to describe hadronic physics with high accuracy as reported in Refs.[43, 44]. Therefore, we believe that there is a systematic path to perform a realistic Higgs boson mass calculation using BSE and DSE, assuming a given TC gauge group with Green's functions scaled from QCD, and varying data such as the number of fermions and other parameters until obtaining the Higgs boson mass experimental value.

Acknowledgments

This research was partially supported by the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) under the grants 310015/2020-0 (A.D.) and 303588/2018-7 (A.A.N.) .

-
- [1] ATLAS Collaboration, Phys. Lett. B **716**, 1 (2012).
 - [2] CMS Collaboration, Phys. Lett. B **716**, 30 (2012).
 - [3] S. Weinberg, Phys.Rev.Lett. **19**, 1264 (1967).
 - [4] K. G. Wilson, Phys. Rev. D **3**, 1818 (1971).
 - [5] G. 't Hooft, "Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking," NATO Adv.Study Inst.Ser.B Phys. **59**, 135 (1980).
 - [6] E. J. Eichten and K. Lane, Phys. Rev. D **103**, 115022 (2021).

- [7] K. Lane, “The composite Higgs signal at the next big collider”, 2022 Snowmass Summer Study, arXiv: 2203.03710.
- [8] S. Weinberg, Phys. Rev. D **19**, 1277 (1979).
- [9] L. Susskind, Phys. Rev. D **20**, 2619 (1979).
- [10] E. Farhi and L. Susskind, Phys. Rept. **74**, 277 (1981).
- [11] K. D. Lane and M. V. Ramana, Phys. Rev. D **44**, 2678 (1991).
- [12] T. W. Appelquist, J. Terning and L. C. R. Wijewardhana, Phys. Rev. Lett. **79**, 2767 (1997).
- [13] Y. Aoki et al., Phys. Rev. D **85**, 074502 (2012).
- [14] T. Appelquist, K. Lane and U. Mahanta, Phys. Rev. Lett. **61**, 1553 (1988).
- [15] R. Shrock, Phys. Rev. D **89**, 045019 (2014).
- [16] M. Kurachi and R. Shrock, JHEP **0612**, 034 (2006).
- [17] V. A. Miransky and K. Yamawaki, Mod. Phys. Lett. A **4**, 129 (1989).
- [18] K.-I. Kondo, H. Mino and K. Yamawaki, Phys. Rev. D **39**, 2430 (1989).
- [19] V. A. Miransky, T. Nonoyama and K. Yamawaki, Mod. Phys. Lett. A **4**, 1409 (1989).
- [20] T. Nonoyama, T. B. Suzuki and K. Yamawaki, Prog. Theor. Phys. **81**, 1238 (1989).
- [21] V. A. Miransky, M. Tanabashi and K. Yamawaki, Phys. Lett. B **221**, 177 (1989).
- [22] K.-I. Kondo, M. Tanabashi and K. Yamawaki, Mod. Phys. Lett. A **8**, 2859 (1993).
- [23] T. Takeuchi, Phys. Rev. D **40**, 2697 (1989).
- [24] B. Holdom, Phys. Rev. D **24**, 1441 (1981).
- [25] K. Yamawaki, Prog. Theor. Phys. Suppl. **180**, 1 (2010); and hep-ph/9603293; C. T. Hill, E. H. Simmons, Phys.Rept. **381** (2003) 235-402, Phys.Rept. **390** (2004) 553-554 (erratum); e-Print: hep-ph/0203079 [hep-ph].
- [26] A. C. Aguilar, A. Doff and A. A. Natale, Phys. Rev. D **97**, 115035 (2018).
- [27] A. Doff and A. A. Natale, Phys. Rev. D **99**, 055026 (2019).
- [28] M. Blatnik, ”The sigma meson in the Bethe-Salpeter approach”, <https://inspirehep.net/literature/1365940>.
- [29] B. Holdom and R. Koniuk, J. High Energ. Phys. **2017**, 102 (2017).
- [30] P. Jain and H. J. Munczek, Phys. Rev. D **44**, 1873 (1991).
- [31] H. J. Munczek and P. Jain, Phys. Rev. D **46**, 438 (1992).
- [32] P. Jain and H. J. Munczek, Phys. Rev. D **48**, 5403 (1993).
- [33] H. D. Politzer, Nucl. Phys. B **117**, 397 (1976).

- [34] D. Binosi, L. Chang, J. Papavassiliou and C. D. Roberts, Phys. Lett. B **742**, 183-188 (2015).
- [35] Muyang Chen, Minghui Ding, Lei Chang, and Craig D. Roberts, Phys. Rev. D **98**, 091505 (2018).
- [36] Y.-Z. Xu, D. Binosi, Z.-F. Cui, B.-L. Li, C. D. Roberts, S.-S. Xu and H.-S. Zong, Phys. Rev. D **100**, 114038 (2019).
- [37] R. Delbourgo and M. D. Scadron, Phys. Rev. Lett. **48**, 379 (1982).
- [38] A. Doff, A. A. Natale and P. S. Rodrigues da Silva, Phys.Rev.D **80**, 055005 (2009).
- [39] S. Mandelstam, Proc. R. Soc. A **233**, 248 (1955).
- [40] K. Lane, Phys. Rev. D **10**, 2605 (1974).
- [41] A. Doff and A. A. Natale, Eur.Phys.J. C **32**, 417-426 (2003).
- [42] A. Doff and A. A. Natale, Eur.Phys.J. C **80**, 684 (2020).
- [43] C. D. Roberts, D. G. Richards, T. Horn and L. Chang, Prog. Part. Nucl. Phys. **120**, 103883 (2021).
- [44] C. D. Roberts, Few Body Syst. **62**, 30 (2021).