On the image of an affine subspace under the inverse function within a finite field

N. Kolomeec, D. Bykov

Sobolev Institute of Mathematics, Russia kolomeec@math.nsc.ru, den.bykov.2000i@gmail.com

Abstract

We consider the function x^{-1} that inverses a finite field element $x \in \mathbb{F}_{p^n}$ (p is prime, $0^{-1} = 0$) and affine \mathbb{F}_p -subspaces of \mathbb{F}_{p^n} such that their images are affine subspaces as well. It is proven that the image of an affine subspace L, |L| > 2, is an affine subspace if and only if $L = q\mathbb{F}_{p^k}$, where $q \in \mathbb{F}_{p^n}^*$ and $k \mid n$. In other words, it is either a subfield of \mathbb{F}_{p^n} or a subspace consisting of all elements of a subfield multiplied by q. This generalizes the results that were obtained for linear invariant subspaces in 2006. As a consequence, we propose a sufficient condition providing that a function $A(x^{-1}) + b$ has no invariant affine subspaces U of cardinality $2 < |U| < p^n$ for an invertible linear transformation $A : \mathbb{F}_{p^n} \to \mathbb{F}_{p^n}$ and $b \in \mathbb{F}_{p^n}^*$. As an example, it is shown that the condition works for S-box of AES. Also, we demonstrate that some functions of the form $\alpha x^{-1} + b$ have no invariant affine subspaces except for \mathbb{F}_{p^n} , where $\alpha, b \in \mathbb{F}_{p^n}^*$ and n is arbitrary.

Keywords: Finite fields, inversion, affine subspaces, invariant subspaces, APN functions.

1 Introduction

In this work we consider affine subspaces whose images under the function that inverses an element of a finite field are affine subspaces as well. We denote this function by σ , i.e. $\sigma : \mathbb{F}_{p^n} \to \mathbb{F}_{p^n}$ such that

$$\sigma(x) = \begin{cases} x^{-1}, & \text{if } x \neq 0\\ 0, & \text{if } x = 0 \end{cases}$$

where \mathbb{F}_{p^n} is a finite field consisting of p^n elements. Hereinafter p is a prime number. A linear subspace L of \mathbb{F}_{p^n} is its linear \mathbb{F}_p -subspace. In other words, L is an additive subgroup of \mathbb{F}_{p^n} . We denote an affine subspace of \mathbb{F}_{p^n} by a + L. For any $S \subseteq \mathbb{F}_{p^n}$, $a \in \mathbb{F}_{p^n}$ and $f : \mathbb{F}_{p^n} \to \mathbb{F}_{p^n}$ we define

$$a + S = \{a + s : s \in S\}, aS = \{as : s \in S\} \text{ and } f(S) = \{f(s) : s \in S\}.$$

It is known that \mathbb{F}_{p^k} is a subfield of \mathbb{F}_{p^n} if and only if $k \mid n$. In this case we assume that $\mathbb{F}_{p^k} \subseteq \mathbb{F}_{p^n}$. Also, $\mathbb{F}_{p^n}^* = \mathbb{F}_{p^n} \setminus \{0\}$. According to the denotations above, we consider affine subspaces U of \mathbb{F}_{p^n} such that $\sigma(U)$ are affine subspaces as well. We are interested in *nontrivial* subspaces that satisfy $2 < |U| < p^n$ since any set S of cardinality 1, 2 (for p = 2) or p^n is always an affine subspace.

These subspaces are connected with invariant subspaces. If we know all U, we can construct all invariant affine subspaces for a function that is affinely equivalent to σ . Recall that $S \subseteq \mathbb{F}_{p^n}$ is an invariant set of f if $f(S) \subseteq S$. Functions f and g are affinely equivalent if and only if g(x) = A(f(B(x) + b)) + a for all $x \in \mathbb{F}_{p^n}$, where A and B are invertible linear transformations, $a, b \in \mathbb{F}_{p^n}$. Similarly to subspaces, we consider linear transformations assuming that \mathbb{F}_{p^n} is a vector space over \mathbb{F}_p .

The function σ is interesting for cryptography. For instance, S-box of AES [8, 11] is based on σ . Its invariant linear subspaces were used to investigate group properties of the AES-like round functions [1, 2, 3]. An attack using an invariant subspace of several rounds of ciphering was introduced in [9]. A more general attack which can use an invariant set is also known [17]. Note that invariant sets for both of them may be constructed starting with S-boxes, see, for instance, [5, 7]. Note that D. Goldstein et al. and S. Mattarei proved that all invariant linear subspaces of the function σ are subfields of \mathbb{F}_{p^n} , see [6, 16]. Thus, we generalize this result in two directions: for affine subspaces and for functions that are affinely equivalent to σ .

At the same time, if f(U) is not an affine subspace for any 2-dimensional affine subspace U of \mathbb{F}_{2^n} , then f is an APN function [12]. Moreover, only APN functions satisfy this property. They are very important for cryptography. Also, there are many open questions in this area, see, for instance, [4]. If p = 2 and n is odd, then σ is an APN function.

Our contribution is the following. In Section 2, we prove that both U and $\sigma(U)$, |U| > 2, are affine subspaces if and only if $U = q\mathbb{F}_{p^k}$, where $q \in \mathbb{F}_{p^n}^*$ and $k \mid n$ (Theorem 2). Next, we show how to apply this to construct a function of the form $A(\sigma(x)) + b$ without invariant affine subspaces, where A is an invertible linear transformation and $b \in \mathbb{F}_{p^n}^*$. Section 3 contains a sufficient condition for such functions to have no nontrivial invariant affine subspaces (Theorem 3). Also, for the function $\alpha\sigma(x) + b$, where $\alpha, b \in \mathbb{F}_{p^n}^*$, we provide a criterion on the existence of invariant affine subspaces excluding the whole \mathbb{F}_{p^n} (Theorem 4). Section 4 demonstrates that the results obtained can theoretically prove that there are no nontrivial invariant affine subspaces for S-box of AES.

2 Subspaces that are mapped to subspaces

In this section we find all nontrivial affine subspaces U such that $\sigma(U)$ is an affine subspace as well. We start with the following well known property of linear subspaces.

Lemma 1. Let L be a linear subspace of \mathbb{F}_{p^n} and |L| > 2. Then $\sum_{x \in L} x = 0$. Proof. Let $s = \sum_{x \in L} x$, $s \in \mathbb{F}_{p^n}$. Let p > 2. Since (p-1)L = L, it holds

$$s = \sum_{x \in L} x = \sum_{x \in L} (p-1)x = (p-1)s$$

But $p - 1 \neq 1$. This implies that s = 0.

Let p = 2. Then dim $L \ge 2$ and $L = L' \cup (a + L')$, where L' is a $(\dim L - 1)$ -dimensional linear subspace of $L, a \in L$. Therefore,

$$s = \sum_{x \in L} x = \sum_{x \in L'} x + \sum_{x \in L'} (a + x) = 2^{\dim L - 1} a = 0.$$

The lemma is proven.

Actually, it is enough to consider only linear subspaces.

Theorem 1. Let U be an affine subspace of \mathbb{F}_{p^n} , |U| > 2 and $\sigma(U)$ be an affine subspace as well. Then U is a linear subspace.

Proof. Let U = a + L and $\sigma(U) = U' = a' + L'$, where L and L' are linear subspaces, $a, a' \in \mathbb{F}_{p^n}$.

First of all, we prove some property of U and U'. Taking into account that |U| > 2, Lemma 1 provides that

$$\sum_{y \in \sigma(U)} y = \sum_{y \in L'} (a' + y) = p^{\dim U} a' + \sum_{y \in L'} y = 0 + 0 = 0.$$

Since $\sigma(U) = \{\sigma(x+s) : s \in L\}$ for any $x \in U$, we obtain that

$$\sum_{s \in L} \sigma(x+s) = 0 \text{ for any } x \in U.$$
(1)

Let us define the following polynomial h over \mathbb{F}_{p^n} :

$$h(y) = \sum_{s \in L} \prod_{t \in L \setminus \{s\}} (y+t).$$

Now we prove by contradiction that U is a linear subspace. Suppose that U is not a linear subspace. Let x be any element of U. Therefore, $x \neq s$ for any $s \in L$. At the same time, $x + s \neq 0$ since $-s \in L$ as well. As a result,

$$h(x) = \sum_{s \in L} \prod_{t \in L \setminus \{s\}} (x+t) = \prod_{t \in L} (x+t) \sum_{s \in L} \sigma(x+s) \stackrel{(1)}{=} 0.$$

Thus, any $x \in U$ is a root of h. However, the degree of h is less than |U| = |L|. Also, $h \neq 0$ since

$$h(0) = \prod_{t \in L \setminus \{0\}} (0+t) = \prod_{t \in L \setminus \{0\}} t \neq 0.$$

In other words, h cannot have |U| roots. This is a contradiction. Hence, U is a linear subspace. The theorem is proven.

The rest of the proofs of this section use Hua's identity [13] (it can also be found in [14, 15]). They are very similar to ones proposed in [6, 16] for linear invariant subspaces of the function σ .

Lemma 2. Let L and $\sigma(L)$ be linear subspaces of \mathbb{F}_{p^n} . Then $a^2\sigma(b) \in L$ for any $a, b \in L$.

Proof. Hua's identity

$$a - (a^{-1} + (b'^{-1} - a)^{-1})^{-1} = ab'a$$

holds for any $a, b' \in \mathbb{F}_{p^n}$, $a, b' \neq 0, ab' \neq 1$. Note that it works not only for fields and can be easily proven.

Let $b' = \sigma(b)$. It is clear that $a^2 \sigma(b) \in L$ if a = 0, b' = 0 or ab' = 1. Next, we consider other cases. It means that $b' = b^{-1}$. Since $\sigma(\sigma(L)) = L$ and both L and $\sigma(L)$ are linear subspaces, the identity provides that

$$b - a = b'^{-1} - a \in L,$$

$$a^{-1} + (b - a)^{-1} \in \sigma(L),$$

$$a - (a^{-1} + (b - a)^{-1})^{-1} \in L.$$

Hence, $ab'a = a^2\sigma(b) \in L$ as well. The lemma is proven.

The following theorem describes all considered subspaces.

Theorem 2. Let U be an affine subspace of \mathbb{F}_{p^n} and |U| > 2. Then $\sigma(U)$ is an affine subspace if and only if $U = q\mathbb{F}_{p^k}$, where $k \mid n$ and $q \in \mathbb{F}_{p^n}^*$. *Proof.* First of all, $\sigma(\mathbb{F}_{p^k}) = \mathbb{F}_{p^k}$ for $k \mid n$ since it is a subfield of \mathbb{F}_{p^n} . Also, $\sigma(q\mathbb{F}_{p^k}) = q^{-1}\sigma(\mathbb{F}_{p^k})$ which is a linear subspace. The sufficient condition is proven.

Next, we prove the necessary condition. Let L = U and $\sigma(L)$ be an affine subspace. By Theorem 1, L is a linear subspace. In fact, $\sigma(L)$ is linear as well since $0 \in L$ and $\sigma(0) = 0$. Without loss of generality we assume that $1 \in L$ since we can consider any qL instead of L, $q \in \mathbb{F}_{p^n}^*$. Indeed, $\sigma(qL)$ is a linear subspace since $\sigma(qL) = q^{-1}\sigma(L)$. In light of that, we need only to prove that L is a subfield of \mathbb{F}_{p^n} .

According to Lemma 2, $a^2\sigma(b) \in L$ for any $a, b \in L$. But $1 \in L$. It means that $\sigma(b) \in L$ and $a^2 \in L$. As a consequence, $L = \sigma(L)$ and $a^2b, a^2 \in L$ for any $a, b \in L$.

We know that $0, 1 \in L$, L is an additive subgroup and all inversions belong to L. The final step is to prove that $bc \in L$ for any $b, c \in L$. There are two cases.

Case 1. p = 2. Let $b, c \in L$. Since $a^2 \in L$ for any $a \in L$, there always exists $a \in L$ such that $c = a^2$. Thus, $bc = a^2b \in L$.

Case 2. $p \neq 2$. For any $b, c \in L$ it holds that $2bc = (c+1)^2b - b - c^2b \in L$. Indeed, all these terms belong to L. Since $2 \in \mathbb{F}_p$ and L is a linear subspace, $bc \in L$ as well.

Thus, L is a subfield of \mathbb{F}_{p^n} , i.e. $L = \mathbb{F}_{p^k}$, where $k \mid n$. The theorem is proven.

Remark 1. Theorem 2 shows that for prime n > 2 and p = 2 the APN function σ maps any nontrivial affine subspace to a set that is not an affine subspace.

3 Functions without invariant subspaces

Here we show how to apply the results of Section 2 to construct functions without invariant affine subspaces of the form

$$\sigma_{A,b}(x) = A(\sigma(x)) + b,$$

where $A : \mathbb{F}_{p^n} \to \mathbb{F}_{p^n}$ is an invertible linear transformation and $b \in \mathbb{F}_{p^n}$. They are affinely equivalent to σ . Let us define

$$\mathcal{S}(\mathbb{F}_{p^n}) = \{ x \in \mathbb{F}_{p^n} : x \notin \mathbb{F}_{p^k}, \text{ where } k \mid n \text{ and } k < n \}.$$

In other words, $\mathcal{S}(\mathbb{F}_{p^n})$ contains all elements of \mathbb{F}_{p^n} which do not lie in its subfields. It is always nonempty.

Theorem 3. Let $A : \mathbb{F}_{p^n} \to \mathbb{F}_{p^n}$ be an invertible linear transformation, $b \in \mathbb{F}_{p^n}^*$ and

$$b^{-1}A(b^{-1}) \in \mathcal{S}(\mathbb{F}_{p^n}).$$
⁽²⁾

Then $\sigma_{A,b}$ has no invariant affine subspaces U such that $2 < |U| < p^n$.

Proof. Let U be affine subspace of \mathbb{F}_{p^n} and $\sigma_{A,b}(U) = U$, where |U| > 2. Theorem 2 provides that $U = q^{-1}K$ and b + A(qK) = U for some subfield K of \mathbb{F}_{p^n} and $q \in \mathbb{F}_{p^n}^*$. This is equivalent to

$$qb + qA(qK) = K.$$
(3)

Since $0 \in K$, (3) provides that $qb = qb + qA(q \cdot 0) \in K$. Thus, $q = b^{-1}u$, where $u \in K \setminus \{0\}$. According to (3),

$$qb + qA(q \cdot u^{-1}) = qb + ub^{-1}A(b^{-1}uu^{-1}) = qb + ub^{-1}A(b^{-1}) \in K.$$

At the same time, both qb and u belong to K. Hence, $b^{-1}A(b^{-1}) \in K$. As a result, $b^{-1}A(b^{-1}) \notin \bigcup_{k|n,k < n} \mathbb{F}_{p^k}$ implies that $U = K = \mathbb{F}_{p^n}$.

Remark 2. The condition (2) is equivalent to

$$b^{-1}\sigma_{A,b}(b) \in \mathcal{S}(\mathbb{F}_{p^n}).$$
 (4)

Proof. Indeed, $b^{-1}\sigma_{A,b}(b) = b^{-1}A(b^{-1}) + 1$. At the same time, $x \in \mathcal{S}(\mathbb{F}_{p^n})$ if and only if $x + 1 \in \mathcal{S}(\mathbb{F}_{p^n})$ since 1 lies in any subfield of \mathbb{F}_{p^n} .

Remark 3. Recall that it is not difficult to determine if $x \in \mathcal{S}(\mathbb{F}_{p^n})$. It is enough to check that $x^{p^k} \neq x$ for any $k \mid n, k < n$.

Even if A does not satisfy (2), we can modify it and obtain a function without nontrivial invariant subspaces.

Corollary 1. Let $A : \mathbb{F}_{p^n} \to \mathbb{F}_{p^n}$ be invertible linear transformation, $b \in \mathbb{F}_{p^n}^*$. Then the transformation $A' : \mathbb{F}_{p^n} \to \mathbb{F}_{p^n}$ such that

$$A'(x) = \alpha \beta A(x), \text{ where } \alpha \in \mathcal{S}(\mathbb{F}_{p^n}) \text{ and } \beta = (b^{-1}A(b^{-1}))^{-1},$$

together with b satisfy the conditions of Theorem 3.

Proof. A' is defined correctly since $b^{-1}A(b^{-1}) \neq 0$. Indeed, A(0) = 0 and A is invertible. It means that $A(b^{-1}) \neq 0$. Let us check the properties of A'. It is clear that A' is linear and invertible since it is a composition of two linear invertible transformations. Also, let us check (2):

$$b^{-1}A'(b^{-1}) = b^{-1}\alpha(b^{-1}A(b^{-1}))^{-1}A(b^{-1}) = \alpha.$$

As a result, A' and b satisfy the conditions of Theorem 3.

However, the proposed condition does not guarantee that $\sigma_{A,b}$ has no invariant affine subspaces U such that $1 \leq |U| \leq 2$, see, for instance, the next section. It is clear that

- any $U = \{u\} \subseteq \mathbb{F}_{p^n}$ is invariant for $\sigma_{A,b}$ if and only if u is a fixed point of $\sigma_{A,b}$,
- any $U = \{u, v\} \subseteq \mathbb{F}_{p^n}$, where $u \neq v$, is an invariant affine subspace if and only if p = 2 and either u, v are fixed point of $\sigma_{A,b}$ or $u = \sigma_{A,b}(v)$, $v = \sigma_{A,b}(u)$.

It is not easy to exclude these cases for arbitrary A. At the same time, we can do this for the simplest linear functions.

Theorem 4. Let $\alpha, b \in \mathbb{F}_{p^n}^*, \sigma_{\alpha,b} : \mathbb{F}_{p^n} \to \mathbb{F}_{p^n}$ such that

$$\sigma_{\alpha,b}(x) = \alpha \sigma(x) + b.$$

Then $\sigma_{\alpha,b}$ has no invariant affine subspaces except for \mathbb{F}_{p^n} if and only if

•
$$\alpha b^{-2} \in M_2$$
 for $p = 2$, where
 $M_2 = \{x \in \mathcal{S}(\mathbb{F}_{2^n}) : \operatorname{tr}(x) = 1\}, \operatorname{tr}(x) = x^{2^0} + x^{2^1} + \ldots + x^{2^{n-1}},$ (5)

•
$$\alpha b^{-2} + 4^{-1} \in M_p$$
 for $p \neq 2$, where

$$M_p = \{ x \in \mathcal{S}(\mathbb{F}_{p^n}) : x \neq y^2 \text{ for any } y \in \mathbb{F}_{p^n} \}.$$
 (6)

Proof. Let U be an invariant affine subspace of $\sigma_{\alpha,b}$. We consider a criterion of the existence of such U depending on |U|.

Case 1. $2 < |U| < p^n$, i.e. U is nontrivial. According to Theorem 3, $\alpha b^{-2} \in \mathcal{S}(\mathbb{F}_{p^n})$ implies that such U does not exist. Let us prove that such Uexists if $\alpha b^{-2} \notin \mathcal{S}(\mathbb{F}_{p^n})$, i.e. $\alpha b^{-2} \in K$ for some subfield K of \mathbb{F}_{p^n} , $|K| < |p^n|$. According to (3), if

$$qb + q\alpha qx \in K$$
 for any $x \in K$,

then U = qK is invariant. Let $q = b^{-1}$. Then, $1 + (\alpha b^{-2})x \in K$ for any $x \in K$. In other words, $U = b^{-1}K$ is invariant.

Thus, such U does not exist if and only if $\alpha b^{-2} \in \mathcal{S}(\mathbb{F}_{p^n})$. Also, if $p \neq 2$, we can see that $\alpha b^{-2} \in \mathcal{S}(\mathbb{F}_{p^n})$ if and only if $\alpha b^{-2} + 4^{-1} \in \mathcal{S}(\mathbb{F}_{p^n})$ since 4^{-1} lies in any subfield of \mathbb{F}_{p^n} .

Case 2. |U| = 1, i.e. we consider fixed points of $\sigma_{\alpha,b}$. Let $x \in \mathbb{F}_{p^n}^*$ since 0 is not a fixed point of $\sigma_{\alpha,b}$. Then x is a fixed point if and only if

$$\alpha x^{-1} + b = x.$$

Multiplying both parts by $xb^{-2} \neq 0$, we obtain the following equivalent equation:

$$y^2 - y - \alpha b^{-2} = 0$$
, where $y = xb^{-1}$. (7)

Case 2.1. p = 2. It is known that the equation (7) has a solution if and only if $tr(\alpha b^{-2}) = 0$, see, for instance, [10].

Case 2.2. $p \neq 2$. The equation (7) has a solution if and only if $(y - 2^{-1})^2 = 4^{-1} + \alpha b^{-2}$ has a solution. Thus, there is a fixed point if and only if $\alpha b^{-2} + 4^{-1} \in \{x^2 : x \in \mathbb{F}_{p^n}\}.$

Case 3. |U| = 2, i.e. $U = \{x, y\} \subseteq \mathbb{F}_{p^n}, x \neq y$. Let U be invariant. Then

$$\alpha \sigma(\alpha \sigma(x) + b) + b = x, \text{ i.e.}$$

$$\alpha \sigma(x) + b = \alpha \sigma(x - b). \tag{8}$$

Case 3.1. $x \in \{0, b\}$. If x = 0, then $\alpha \sigma(0) + b = b \neq 0$. Let x = b. According to (8), $\alpha \sigma(b) + b = 0 \neq b$. In other word, $U = \{0, b\}$ for both cases. Also, $\alpha \sigma(b) + b = 0$ is equivalent to $\alpha b^{-2} = -1$. Moreover, it is not difficult to see that $\sigma_{\alpha,b}(\{0, b\}) = \{0, b\}$ if and only if $\alpha b^{-2} = -1$. At the same time, $-1 \notin \mathcal{S}(\mathbb{F}_{p^n})$.

Case 3.2. $x \notin \{0, b\}$. We multiply both parts of the last equation by $b^{-2}x(x-b)$:

$$b^{-2}x(\alpha\sigma(x) + b)(x - b) = b^{-2}x\alpha, \text{ i.e.}$$

$$b^{-2}(bx^2 - \alpha b - b^2x) = b^{-2}(\alpha - \alpha\sigma(x)x)x, \text{ or, equivalently,}$$

$$(xb^{-1})^2 - xb^{-1} - \alpha b^{-2} = 0.$$

Since it is equivalent to (7), x is a fixed point of $\sigma_{\alpha,b}$, see case 2.

Finally, \mathbb{F}_{p^n} is the only invariant affine subspace of $\sigma_{\alpha,b}$ if and only if $\alpha b^{-2} \in M_2$ for p = 2 and $\alpha b^{-2} + 4^{-1} \in M_p$ for $p \neq 2$.

Remark 4. The sets M_p defined in (5) and (6) are nonempty.

Proof. It is straightforward that $|\{x \in \mathbb{F}_{2^n} : \operatorname{tr}(x) = 1\}| = 2^{n-1}$ and $|\{x \in \mathbb{F}_{p^n} : x \neq y^2 \text{ for any } y \in \mathbb{F}_{p^n}\}| = \frac{1}{2}(p^n - 1)$. At the same time, we can estimate the number of elements that lie in subfields of \mathbb{F}_{p^n} :

$$\left|\bigcup_{k|n,k< n} \mathbb{F}_{p^k}\right| < 1 + p + p^2 + \ldots + p^{n-2} = \frac{p^{n-1} - 1}{p-1}, \text{ where } n \ge 3.$$

This implies that M_p is nonempty for $n \ge 3$. It is not difficult to check that M_p is nonempty for n = 1, 2 as well.

Theorem 4 allows us to construct a permutation of an arbitrary \mathbb{F}_{p^n} without invariant affine subspaces (except for the whole \mathbb{F}_{p^n}). However, its algebraic structure is very simple. Also, if n is not prime, there always exists a linear subspace $L \neq \mathbb{F}_{p^n}$ such that $\sigma_{\alpha,b}(u+L) = v + L$ for some $u, v \in \mathbb{F}_{p^n}$. This makes the function $\sigma_{\alpha,b}$ not very good for cryptography.

4 S-box of AES

The Advanced Encryption Standard (AES) is the most popular block cipher whose S-box $S : \mathbb{F}_2^8 \to \mathbb{F}_2^8$ (see [8, Figure 7]) is based on the function $\sigma_{A,b}$ for certain invertible linear transformation A and $b \in \mathbb{F}_{2^8}$, see [8, 11]. The field \mathbb{F}_{2^8} is constructed as the quotient ring of polynomials modulo

$$m(x) = x^8 + x^4 + x^3 + x + 1$$
 over \mathbb{F}_2

and $b = x^6 + x^5 + x + 1 = 0x63$. Let us consider its invariant subspaces.

• Nontrivial affine subspaces. It can be shown that $t = b^{-1}S(b) \in S(\mathbb{F}_{2^8})$. Indeed, $S(b) = x^7 + x^6 + x^5 + x^4 + x^3 + x + 1 = 0$ xFB. It is not difficult to check that $t = x^7 + x^6 + x^3 = 0$ xC8. Also,

$$\begin{split} t^2 &= x^6 + x^5 + x^4 + 1 \neq t, \\ t^{2^2} &= x^7 + x^6 + x^4 + x^3 + x^2 + 1 \neq t, \\ t^{2^4} &= x^7 + x^4 + x^3 + 1 \neq t. \end{split}$$

Hence, $t \in \mathcal{S}(\mathbb{F}_{2^8})$. In other words, Theorem 3 and Remark 2 guarantee that S has no nontrivial invariant affine subspaces.

- Fixed points. It is known that $S(x) \neq x$ for any $x \in \mathbb{F}_2^8$ by the choice, see [11]. Therefore, S has no invariant affine subspaces of dimension 0.
- However,

$$S(0x73) = 0x8F \text{ and } S(0x8F) = 0x73.$$

This implies that

$$\{\texttt{0x73},\texttt{0x8F}\} = \{x^6 + x^5 + x^4 + x + 1, x^7 + x^3 + x^2 + x + 1\}$$

is its invariant affine subspace of dimension 1.

Thus, the results obtained allow us to prove theoretically that S has no nontrivial invariant affine subspaces. At the same time, it has an invariant affine subspace of dimension 1.

Acknowledgements The work is supported by Mathematical Center in Akademgorodok under agreement No. 075–15–2022–281 with the Ministry of Science and Higher Education of the Russian Federation.

References

- F. М. |1| A. Caranti, Volta, Sala, Imprimitive permutations the functions generated by round of key-alternating groups block ciphers and truncated differential cryptanalysis. https://arxiv.org/abs/math/0606022 (2006)
- [2] A. Caranti, F. Volta, M. Sala, An application of the O'Nan-Scott theorem to the group generated by the round functions of an AES-like cipher. Designs, Codes and Cryptography, 52, 293–301 (2009)
- [3] A. Caranti, F. Volta, M. Sala, On some block ciphers and imprimitive groups. Appl. Algebra Eng. Commun. Comput., 20, 339–350 (2009)
- [4] C. Carlet, Open Questions on Nonlinearity and on APN Functions. In: Arithmetic of Finite Fields 5th International Workshop, WAIFI 2014, Gebze, Turkey, September 27-28, 2014. Lecture Notes in Computer Science, 9061, 83-107 (2015)
- [5] D. A. Burov, About existence of the special nonlinear invariants for round functions of XSL-ciphers. Diskr. Mat., 33(2), 31–45 (2021) (in Russian)
- [6] D. Goldstein, R. Guralnick, L. Small, E. Zelmanov, Inversion-invariant additive subgroups of division rings. Pacific Journal of Mathematics, 227(2), 287–294 (2006)
- [7] D. I. Trifonov, D. B. Fomin, Invariant subspaces in SPN block cipher. Prikl. Diskr. Mat., 54, 58–76 (2021) (in Russian)
- [8] FIPS Publication 197 (NIST), Advanced Encryption Standard. https://nvlpubs.nist.gov/nistpubs/fips/nist.fips.197.pdf (2001)
- [9] G. Leander, M. A. Abdelraheem, H. AlKhzaimi, E. Zenner, A Cryptanalysis of PRINTcipher: The Invariant Subspace Attack. In: Rogaway, P. (eds) Advances in Cryptology – CRYPTO 2011, CRYPTO 2011. Lecture Notes in Computer Science, 6841, 206–221 (2011)

- [10] J. Cherly, L. Gallardo, L. Vaserstein, E. Wheland, Solving quadratic equations over polynomial rings of characteristic two. Publicacions matematiques, 42(1), 131–142 (1998)
- [11] J. Daemen, V. Rijmen, The design of Rijndael: AES the Advanced Encryption Standard. Springer-Verlag, 2002
- [12] K. Nyberg, Differentially uniform mappings for cryptography. In: Advances in Cryptology EUROCRYPT '93, Workshop on the Theory and Application of Cryptographic Techniques, Lofthus, Norway, May 23-27, 1993. LNCS, 765, 55–64 (1994)
- [13] L.-K. Hua, Some properties of a sfield. Proceedings of the National Academy of Sciences of the United States of America, 35, 533–537 (1949)
- [14] N. Jacobson, Basic Algebra, I. W. H. Freeman and Co., San Francisco, California, 1974
- [15] R. Lidl, H. Niederreiter, Finite Fields: Encyclopedia of Mathematics and its Applications, vol. 20. Addison-Wesley Publishing Company Advanced Book Program, Reading, MA, 1983. With a foreword by P. M. Cohn
- [16] S. Mattarei, Inverse-closed additive subgroups of fields. Israel Journal of Mathematics, 159, 343–347 (2007)
- [17] Y. Todo, G. Leander, Y. Sasaki, Nonlinear invariant attack: practical attack on full SCREAM, iSCREAM, and Midori64. In: ASIACRYPT 2016. LNCS, 10032, 3–33 (2016)