ASYMPTOTIC STABILITY NEAR THE SOLITON FOR QUARTIC KLEIN-GORDON IN 1D

ADILBEK KAIRZHAN AND FABIO PUSATERI

ABSTRACT. We consider the nonlinear focusing Klein-Gordon equation in 1+1 dimensions and the global space-time dynamics of solutions near the unstable soliton. Our main result is a proof of optimal decay, and local decay, for even perturbations of the static soliton originating from well-prepared initial data belonging to a subset of the stable manifold constructed in [2, 24]. Our results complement those of Kowalczyk-Martel-Muñoz [24] and confirm numerical results of Bizon-Chmaj-Szpak [3] when considering nonlinearities u^p with $p \geq 4$. In particular, we provide new information both local and global in space about asymptotically stable perturbations of the soliton under localization assumptions on the data.

Contents

| 1. | Introduction | 1 |
|------------|-------------------------|----|
| 2. | Linear Theory | 8 |
| 3. | Set-up and bootstrap | 15 |
| 4. | Quadratic term analysis | 19 |
| 5. | Quartic term analysis | 21 |
| References | | 33 |

1. Introduction

1.1. The problem. We consider the focusing quartic Klein-Gordon (NLKG) equation

$$\partial_t^2 u - \partial_x^2 u + u = u^4, (1.1)$$

where $(t, x) \in \mathbb{R} \times \mathbb{R}$ are time and space variables, and $u(t, x) \in \mathbb{R}$. Our results apply to more general power nonlinearities as well but we restrict our discussion and proof to the case p = 4 for concreteness; see Remark 1.3. The Hamiltonian associated with (1.1) is

$$\mathcal{H}(u, u_t) = \frac{1}{2} \int_{\mathbb{R}} \left[(\partial_t u)^2 + (\partial_x u)^2 + u^2 \right] dx - \frac{1}{5} \int_{\mathbb{R}} u^5 dx, \tag{1.2}$$

and for initial data in the energy space, $(u(0), u_t(0)) \in H^1(\mathbb{R}) \times L^2(\mathbb{R})$, the equation is locally well-posed, and globally well-posed for small solutions. Global existence for small energy initial data follows from standard energy estimates, see for example [4]. It is also known that finite time blow-up occurs for solutions with negative energy initial data [28] and for large positive initial data [38, 40], which follows from a convexity argument.

(1.1) admits an explicit time-independent static soliton solution

$$Q(x) = (\alpha + 1)^{\frac{1}{2\alpha}} \operatorname{sech}^{1/\alpha}(\alpha x), \quad \alpha := \frac{3}{2}.$$
 (1.3)

In this paper we study the dynamics of certain classes of global-in-time solutions ($t \ge 0$) in the vicinity of Q and, in particular, their behavior on the whole real line and their decay towards

the soliton. Letting u = Q + v denote the perturbed soliton for a real-valued and sufficiently small v, we see that

$$v_{tt} + \mathcal{L}v = N(v), \tag{1.4}$$

where

$$\mathcal{L} := -\partial_x^2 + 1 - 4Q^3, \qquad N(v) := 6Q^2v^2 + 4Qv^3 + v^4. \tag{1.5}$$

The spectral properties of the linearized operator \mathcal{L} were described in details in [5], also in the case of general powers u^p in (1.1). In particular, it is known that \mathcal{L} has continuous spectrum $[1, \infty)$ and exactly two isolated eigenvalues:

- The first eigenvalue is $\lambda_0 = -\alpha(\alpha + 2) = -\Omega^2$ ($\Omega > 0$) with corresponding even normalized eigenfunction

$$\rho(x) := c_0(\operatorname{sech}(\alpha x))^{\frac{\alpha+1}{\alpha}}, \qquad \mathcal{L}\rho = -\Omega^2 \rho, \tag{1.6}$$

where the positive c_0 is chosen to satisfy $\|\rho\|_{L^2} = 1$;

- The other eigenvalue is $\lambda_1 = 0$ with corresponding odd eigenfunction Q'(x); in particular, \mathcal{L} has no gap eigenvalues (internal mode) in (0,1).
- Moreover, the operator \mathcal{L} has no resonance at the edge of the continuous spectrum, i.e., there does not exist a bounded solution φ to $\mathcal{L}\varphi = 0$. Equivalently, the potential $V = -4Q^3$ is 'generic'.

To avoid the translation mode and having to modulate the soliton, we restrict our analysis to even perturbations, and decompose v as

$$v(t,x) = a(t)\rho(x) + \chi(t,x), \quad \langle \chi, \rho \rangle_{L_x^2} = 0, \quad \chi = P_c v, \tag{1.7}$$

where $P_c: L_x^2 \to L_x^2$ is the projection onto the continuous spectrum of \mathcal{L} .

1.2. Stable manifold and some literature. Before stating our main result we review the definition of the stable manifold around the soliton Q for even solutions of (1.1), following Kowalczyk, Martel and Muñoz in [24] (see also Bates-Jones [2]), as well as the main result in [24] concerning¹ (1.1). Rewrite the linear part of the equation (1.4) as the system

$$\begin{cases}
\partial_t v_1 = v_2, \\
\partial_t v_2 = -\mathcal{L}v_1,
\end{cases}$$
(1.8)

where $\vec{v} := (v_1, v_2) = (v, v_t)$ and the operator \mathcal{L} is defined in (1.5). The exponentially unstable and stable solutions of (1.8) are $\vec{v}_+(t, x) = e^{\Omega t} Y_+(x)$ and $\vec{v}_-(t, x) = e^{-\Omega t} Y_-(x)$, respectively, where

$$Y_{+} = (\rho, \Omega \rho)$$
 and $Y_{-} = (\rho, -\Omega \rho)$. (1.9)

We let

$$Z_{+} = \left(\rho, \Omega^{-1}\rho\right) \tag{1.10}$$

so that $\langle Y_-, Z_+ \rangle = 0$.

For every $\delta_0 > 0$, define

$$\mathcal{A}(\delta_0) := \left\{ \boldsymbol{\gamma} \in H_x^1(\mathbb{R}) \times L_x^2(\mathbb{R}) : \quad \boldsymbol{\gamma} \text{ is even, } \|\boldsymbol{\gamma}\|_{H_x^1 \times L_x^2} < \delta_0 \text{ and } \langle \boldsymbol{\gamma}, Z_+ \rangle = 0 \right\}. \tag{1.11}$$

Then, the following stability result was obtained in [2, 24].

The results in [24] cover the more general case of nonlinearities $|u|^{p-1}u$ with p>3.

Theorem 1 ([2, 24]). There exist $C, \delta_0 > 0$ and a Lipschitz function $h : \mathcal{A}(\delta_0) \to \mathbb{R}$ with h(0) = 0 and $|h(\gamma)| \le C ||\gamma||_{H^1_x \times L^2_x}^{3/2}$ such that, denoting

$$\mathcal{M}(\delta_0) := \{ (Q, 0) + \gamma + h(\gamma) Y_+ : \ \gamma \in \mathcal{A}(\delta_0) \}$$

$$(1.12)$$

the following holds:

1. If $(u_0, u_1) \in \mathcal{M}(\delta_0)$, then the solution of (1.1) with initial data (u_0, u_1) is global and satisfies $\left\| (u(t) - Q, \partial_t u(t)) \right\|_{H^1_x \times L^2_x} \le C \left\| (u_0 - Q, u_1) \right\|_{H^1_x \times L^2_x} \quad \text{for all } t \ge 0. \tag{1.13}$

2. If a global even solution $(u, \partial_t u)(t)$ of (1.1) satisfies

$$\left\| (u(t) - Q, \partial_t u(t)) \right\|_{H^1_x \times L^2_x} < \frac{1}{2} \delta_0 \quad \text{for all } t \ge 0,$$

then $(u, \partial_t u)(t) \in \mathcal{M}(\delta_0)$ for all $t \geq 0$.

The above classical theorem provides a stable manifold $\mathcal{M} = \mathcal{M}(\delta_0)$ such that data originating in \mathcal{M} is global, and stays globally-in-time close to the soliton in the energy norm. Moreover, any solution that is globally sufficiently close to Q, has to lie in \mathcal{M} . Note that, in view of the decomposition $u = Q + v = Q + a(t)\rho + \chi$, see (1.7), if one chooses sufficiently small data $a(0), \dot{a}(0)$ and $(\chi(0, x), \chi_t(0, x)) \in H_x^1 \times L_x^2$ small enough, so that $(u(0), u_t(0)) \in \mathcal{M}(\delta_0)$, then the global bound (1.13) implies that, for every $t \geq 0$, we have $||v(t)||_{H_x^1 \times L_x^2} \lesssim \delta_0$ and

$$|a(t)| + |\dot{a}(t)| + |\chi(t)|_{L_x^2} + |\partial_t \chi(t)|_{L_x^2} \lesssim \delta_0.$$
(1.14)

A natural question to ask is whether the above global solutions are also attracted to the soliton, besides remaining close to the it for all times, and what is their asymptotic behavior as $t \to \infty$.

For nonlinear equations with large enough power nonlinearities these types of asymptotic stability result have been achieved by several authors. The literature is ample so we will only mention a few recent results, and refer the readers to the cited works for more discussions and references. For the Klein-Gordon equation with power nonlinearity u^p with p > 5 asymptotic stability (on the stable manifold) was proven by Krieger-Nakanishi-Schlag [27] for data in the energy space. See also [35, 36] and reference therein, for a similar problem and more general results in higher dimensions. See also the review by Kowalczyk-Martel-Muñoz [22]. For the nonlinear Schrödinger (NLS) equation Krieger-Schlag [26] treated the case of a $|u|^{2\sigma}u$ nonlinearity with $\sigma > 2$ in 1d and Cuccagna-Pelinovsky [12] the cubic case. See also the seminal work of Schlag [37] for the cubic NLS in 3d, the recent works and survey by Cuccagna-Maeda [9, 8] and Maeda-Yamazaki [33] and references therein.

For low power nonlinearities instead (typically the energy sub-critical ones) one cannot expect similar asymptotic stability result in the energy space $H_x^1(\mathbb{R}) \times L_x^2(\mathbb{R})$, essentially because Strichartz estimates are not very effective in controlling the nonlinearity. There are then two natural different and complementary ways to look at this question: (a) studying the asymptotics for energy solutions locally in the space variable x, or (b) studying asymptotics on the whole real line in a smaller space, e.g., a weighted Sobolev space. The first approach is usually based on delicate virial type estimates, and has been very successfully employed in several problems; see for example the already mentioned [24], the recent work of Li-Lührmann [30] on the soliton of quadratic KG, Cuccagna-Maeda-Scrobogna [11] on cubic KG, and the works on kink solutions for relativistic scalar fields [23, 25, 10]. The second approach has instead been pursued mostly in the context of kinks (non-localized solitons) which naturally lead to nonlinear PDEs with potentials and low power non-localized nonlinearities; we refer

the reader to Delort-Masmoudi [14], Germain and the second author [19] and Zhang [21], Lührmann-Schlag [34], Lindblad-Lührmann-Soffer [31], and Lindblad-Lührmann-Schlag-Soffer [32].

Concerning the problem we are interested in, [24] showed the following asymptotic stability result

Theorem 2 ([24, Theorem 1]). Consider the Klein-Gordon equation

$$\partial_t^2 u - \partial_x^2 u + u = |u|^{2\alpha} u, \qquad \alpha > 1, \tag{1.15}$$

and denote its static soliton by

$$Q(x) := (\alpha + 1)^{\frac{1}{2\alpha}} \operatorname{sech}^{1/\alpha}(\alpha x). \tag{1.16}$$

Then, there exists a constant $\delta > 0$ such that if a global even solution (u, u_t) of (1.15) satisfies

$$\left\| \left(u(t) - Q, \partial_t u(t) \right) \right\|_{H^1_x(\mathbb{R}) \times L^2_x(\mathbb{R})} < \delta \quad \text{for all } t \ge 0, \tag{1.17}$$

then, for any bounded interval I,

$$\lim_{t \to \infty} \| (u(t) - Q, \partial_t u(t)) \|_{H^1_x(I) \times L^2_x(I)} = 0.$$
 (1.18)

Moreover,

$$\int_{0}^{\infty} \left\| (u(t) - Q, \partial_{t} u(t)) \right\|_{H_{x}^{1}(I) \times L_{x}^{2}(I)}^{2} dt < \infty.$$
 (1.19)

Note that a global solution satisfying (1.17) for δ small enough is on the stable manifold and therefore stays close to the soliton in view of Theorem 1. Theorem 2 then gives that these solutions actually converge as $t \to \infty$ to Q locally-in-space. Our main theorem will give a complementary asymptotic stability result; we will assume more localization on the (well-prepared) initial data and prove convergence to Q on the whole real-line with optimal time-decay rates, and local-decay estimates as well. In particular, we will show that, see (1.27),

$$\int_0^\infty \left\| u(t) - Q \right\|_{L_x^\infty(\mathbb{R})}^{2+} dt < \infty, \tag{1.20}$$

$$\int_0^\infty \left\| \langle x \rangle^{-2} (u(t) - Q) \right\|_{L_x^\infty(\mathbb{R})}^{1+} dt < \infty. \tag{1.21}$$

1.3. Main result. Recall that we write $u(t,x) = Q(x) + v(t,x) = Q(x) + a(t)\rho(x) + \chi(t,x)$, where ρ is given in (1.6), $Y_{\pm} := (\rho, \pm \Omega \rho)$ and \mathcal{M} is defined in (1.12) with (1.10). We will consider well-prepared initial data in a subset of \mathcal{M} as we now describe. Let $\gamma(x) = (\gamma_1(x), \gamma_2(x))$ be given in the form (see (1.11))

$$\gamma(x) = bY_{-}(x) + \zeta(x), \quad \zeta = (\zeta_1, \zeta_2),$$

where b is a scalar constant, and ζ is even in x, with $\langle \rho, \zeta_j \rangle = 0$ for j = 1, 2. In other words, γ has components

$$\gamma_1(x) = b\rho(x) + \zeta_1(x), \quad \gamma_2(x) = -\Omega b\rho(x) + \zeta_2(x).$$
 (1.22)

We define (see (1.5))

$$H := P_c(-\partial_x^2 - 4Q^3), \quad H + 1 = P_c \mathcal{L},$$
 (1.23)

where P_c denotes the projection onto the continuous spectrum, and require that

$$|b| + \|(\sqrt{H+1}\ \zeta_1, \zeta_2)\|_{H_x^2} + \|\langle x\rangle(\sqrt{H+1}\ \zeta_1, \zeta_2)\|_{L_x^2} \le c\epsilon_0 \tag{1.24}$$

for sufficiently small $\epsilon_0 > 0$, with c a small absolute constant (c = 1/10, say). Then, γ belongs to a strict subset of $\mathcal{A}(\epsilon_0)$. We then define the following subset of $\mathcal{M}(\epsilon_0)$:

 $\mathcal{M}_0(\epsilon_0) := \{(Q, 0) + \gamma + h(\gamma)Y_+: \gamma \text{ is even and satisfies } (1.22) - (1.24)\} \subseteq \mathcal{M}(\epsilon_0), \quad (1.25)$ where h is the Lipschitz function from Theorem 1. We are now ready to state our main result:

Theorem 3. There exists sufficiently small $\epsilon_0 > 0$ such that, for every $(u_0, u_1) \in \mathcal{M}_0(\epsilon_0)$, the equation (1.1) with initial data $(u, \partial_t u)(0) = (u_0, u_1)$ admits a unique global solution of the form

$$u(t,x) = Q(x) + a(t)\rho(x) + \chi(t,x), \quad \langle \rho, \chi \rangle_{L_x^2} = 0,$$
 (1.26)

that satisfies, for all t > 0, (1.13) and the following global-in-time decay bounds:

$$|a(t)| \lesssim \epsilon_0 \langle t \rangle^{-2}, \qquad \|\chi\|_{L^{\infty}_x} \lesssim \epsilon_0 \langle t \rangle^{-1/2}, \qquad \|\langle x \rangle^{-2} \chi\|_{L^{\infty}_x} \lesssim \epsilon_0 \langle t \rangle^{-1}.$$
 (1.27)

Moreover, χ behaves like a free solution of the Klein-Gordon equation as $t \to \infty$.

Here are a few remarks about our result:

Remark 1.1 (About the initial data). The choice of the initial data can also be interpreted as follows. We can express (u_0, u_1) as

$$u_0(x) = Q(x) + a_0 \rho(x) + \chi_0(x),$$

 $u_1(x) = a_1 \rho(x) + \chi_1(x),$

with $\langle \chi_j, \rho \rangle_{L^2_x} = 0$, j = 1, 2, $(a_0, a_1) = (a(0), \dot{a}(0))$, $(\chi_0, \chi_1) = (\chi(0), \partial_t \chi(0))$. Then,

$$(u_0, u_1) = (Q, 0) + \frac{1}{2} \left(a_0 - \frac{1}{\Omega} a_1 \right) Y_- + \frac{1}{2} \left(a_0 + \frac{1}{\Omega} a_1 \right) Y_+ + (\chi_0, \chi_1),$$

which, compared with (1.22) would give $b = (1/2)(a_0 - \Omega^{-1}a_1)$, $(\zeta_1, \zeta_2) = (\chi_0, \chi_1)$. Then $(u_0, u_1) \in \mathcal{M}_0(\epsilon_0)$ if we impose the stability condition

$$\frac{1}{2}\left(a_0 + \frac{1}{\Omega}a_1\right) = h\left(\frac{1}{2}\left(a_0 - \frac{1}{\Omega}a_1\right)Y_- + (\chi_0, \chi_1)\right),\,$$

that χ_0, χ_1 are even, and

$$|a_0| + |a_1| + \|(\sqrt{H+1} \chi_0, \chi_1)\|_{H_x^2} + \|\langle x\rangle(\sqrt{H+1} \chi_0, \chi_1)\|_{L_x^2} \le c\epsilon_0.$$
 (1.28)

for some absolute c > 0 small enough.

Remark 1.2 (Linearized operator). The potential appearing in linearized operator (1.5) is $V = -4Q^3 = -10\cosh^{-2}(3x/2)$. The corresponding Schrödinger operator is, up to a conjugation, given by $H = -\partial_x^2 + c\cosh^{-2}(x)$ with c = -40/9. In particular, V is not in the class of Pöschel-Teller potentials $V_{\ell}(x) = -\ell(\ell+1)\cosh^{-2}(x)$, $\ell = 0, 1, 2, ...$ that can be conjugated to the flat V = 0 operator; see the discussion in [34] and references therein.

Remark 1.3 (More general powers). Consider the pure power Klein-Gordon equation

$$\partial_t^2 u - \partial_x^2 u + u = |u|^{2\alpha} u, \tag{1.29}$$

and its soliton

$$Q_{\alpha}(x) = (\alpha + 1)^{\frac{1}{2\alpha}} \operatorname{sech}^{1/\alpha}(\alpha x), \tag{1.30}$$

Recall that Theorem 2 applies to all powers $\alpha > 1$. In the borderline cubic case, $\alpha = 1$, the linearized operator has an even resonance. Then, the results of [24] do not apply, and no asymptotic stability results are available to date.

In the quadratic case, $\alpha = 1/2$, the linearized operator has an internal mode (a gap eigenvalue in (0,1)). This is a more difficult case since stability does not hold at the linearized level because of the presence of periodic-in-time localized bound states. Nevertheless, a stability result like the one in Theorem 2 was recently proven by Li and Lührmann [30]. See this latter, [23], and the work of Léger and the second author [29] for more references about internal mode dynamics.

Our main result in Theorem 3, which corresponds to the case $2\alpha = 3$, can be easily extended to any α such that 2α is an integer larger 3. Non-integer powers with $\alpha \geq 3/2$ can also be treated with minor modifications. The treatment of powers $\alpha \in (1,3/2)$ may instead require additional arguments since our approach is not well suited for (relatively low) fractional powers. However, it is natural to conjecture that the same result would hold, consistently with the predictions of [3]; see also the next remark for more about this.

Remark 1.4 (Pointwise and integrated decay on the real line). Our main Theorem 3 complements known results in two ways. First, it complements [24] by showing that asymptotic stability holds on the whole real line. Second, it complements results for larger powers, such as [27], by obtaining (also for lower power nonlinearities) exact rates of convergence to the soliton for a large class of solutions trapped by it. As already mentioned, we need to require stronger assumptions on the data, the most relevant of which is the smallness of the weighted norm in (1.28).

Also note that the first and second estimate in (1.27) immediately imply the integrated decay (1.20), while the first and third give (1.21).²

Remark 1.5 (About local decay and rates of convergence). Algebraic rates of decay towards the soliton on the stable manifold had been predicted numerically already in [3]. In particular, Fig. 2 in [3] depicts the expected decay (or growth, for larger data) in the quartic case (1.1). Note that the data used for these numerical results is a Gaussian.

The predictions of [3] actually give a local-in-space rate of decay of $t^{-3/2}$, as opposed to the rate of t^{-1} which we obtain from (1.27). The discrepancy here is just due to the fact that we only require smallness of $xu(0) \in L^2$. For such data, even at a the level of the linear operator $e^{it\sqrt{H+1}}$, one cannot obtain more than t^{-1} local decay. Assuming the stronger localization $xu(0) \in L^1$, the linear decay rate is indeed $t^{-3/2}$, and one can expect to obtain this same rate of convergence to Q also for our nonlinear solutions.

1.4. Strategy of the proof. We begin by following the standard strategy of substituting the decomposition (1.7) into (1.4)-(1.5), and separating the discrete and continuous modes as follows:

$$\begin{cases} a_{tt} - \Omega^2 a = \langle N(v), \rho \rangle_{L_x^2} := F[v], \\ \chi_{tt} + \mathcal{L}\chi = P_c N(v). \end{cases}$$
 (1.31)

Following the Duhamel's principle, (1.31) is equivalent to the system of integral equations

$$\begin{cases} a(t) = \cosh(\Omega t)a(0) + \frac{1}{\Omega}\sinh(\Omega t)\dot{a}(0) + \frac{1}{\Omega}\int_{0}^{t}\sinh(\Omega(t-s))F[v](s)ds, \\ \chi(t) = \cos(\sqrt{H+1}t)\chi(0) + \frac{\sin(\sqrt{H+1}t)}{\sqrt{H+1}}\dot{\chi}(0) + \int_{0}^{t}\frac{\sin(\sqrt{H+1}(t-s))}{\sqrt{H+1}}P_{c}N(v(s))ds. \end{cases}$$
(1.32)

²Similar estimates with $\partial_t u$ replacing Q - u can also be obtained, but one would need to change slightly the assumptions on the data in terms of derivatives; for simplicity, we do not pursue this here.

Imposing the stability condition

$$\dot{a}(0) = -\Omega a(0) - \int_0^\infty e^{-\Omega s} F[v](s) ds, \tag{1.33}$$

which ensures that the contribution of the exponentially increasing term $e^{\Omega t}$ of a(t) vanishes as $t \to \infty$, (1.32) reduces to

$$\begin{cases} a(t) = e^{-\Omega t} \left[a(0) + \frac{1}{2\Omega} \int_0^\infty e^{-\Omega s} F[v](s) ds \right] - \frac{1}{2\Omega} \int_0^\infty e^{-\Omega |t-s|} F[v](s) ds, \\ \chi(t) = \cos(\sqrt{H+1}t) \chi(0) + \frac{\sin(\sqrt{H+1}t)}{\sqrt{H+1}} \dot{\chi}(0) + \int_0^t \frac{\sin(\sqrt{H+1}(t-s))}{\sqrt{H+1}} P_c N(v(s)) ds. \end{cases}$$
(1.34)

Note that (1.33) holds for solutions on the stable manifold, in view of (1.13).

To rewrite the evolution equation for $\chi(t,x)$ in (1.31) in a convenient way for our analysis, we introduce the profile function g(t,x) as

$$g(t,x) := e^{it\sqrt{H+1}}(\partial_t - i\sqrt{H+1})\chi(t,x). \tag{1.35}$$

Then the real-valued function χ is given in terms of g and \bar{g} by

$$\chi(t,x) = \frac{1}{2i\sqrt{H+1}} (e^{it\sqrt{H+1}}\bar{g}(t,x) - e^{-it\sqrt{H+1}}g(t,x)). \tag{1.36}$$

We then want to look at Duhamel's formula in terms of g on the distorted Fourier side. We refer the reader to Section 2 for the definition of the distorted Fourier Transform (dFT) and its properties. Applying the distorted Fourier transform to (1.35) gives $\tilde{g}(t,k) = e^{it\langle k \rangle}(\partial_t - i\langle k \rangle)\tilde{\chi}(t,k)$, and the second equation in (1.34) becomes

$$\partial_t \widetilde{g}(t,k) = e^{it\langle k \rangle} \widetilde{\mathcal{F}} P_c N(Q,v), \qquad N(Q,v) := P_c (6Q^2 v^2 + 4Qv^3 + v^4). \tag{1.37}$$

Integrating (1.37) over time gives

$$\widetilde{g}(t,k) = \widetilde{g}(0,k) + \int_0^t e^{is\langle k \rangle} \widetilde{\mathcal{F}} P_c N(s,k) ds.$$
 (1.38)

Equation (1.38) is the starting point for our analysis of the projection on the continuous spectrum of the solution.

To obtain the desired nonlinear estimates that will imply our main result, we introduce the following norm

$$||v||_X := ||\langle t \rangle^2 a(t)||_{L_t^{\infty}(0,\infty)} + ||\partial_k \widetilde{g}||_{L_t^{\infty}((0,\infty):L_k^2)} + ||\langle k \rangle^2 \widetilde{g}||_{L_t^{\infty}((0,\infty):L_k^2)}, \tag{1.39}$$

The first norm above is used to control the evolution of the solution in the direction of the even eigenfunction, while the other two control the evolution of the continuous spectral component. The proof of our result will follow from a priori estimates for this norm:

Proposition 1.1. For $||v||_X$ sufficiently small we have

$$||v||_X \lesssim |a(0)| + ||\partial_k \widetilde{g}(0)||_{L_k^2} + ||\langle k \rangle^2 \widetilde{g}(0)||_{L_k^2} + ||v||_X^2.$$

Proposition 1.1 follows from the three Propositions 3.1, 3.2 and 3.3. With Proposition 1.1 we can prove the main theorem.

1.5. **Proof of Theorem 3.** First, given initial data $(u_0, u_1) \in \mathcal{M}_0(\epsilon_0)$ with (1.22)-(1.24), Theorem 1 gives a global solution trapped by the soliton for all $t \geq 0$ provided ϵ_0 is chosen small enough depending on δ_0 .

The smallness condition (1.28) on the initial data with (recall (1.35)) $(-\sqrt{H+1}\chi_0, \chi_1) = (\text{Im }(g(0)), \text{Re }(g(0)))$ and the estimates (2.18) and (2.19), imply that

$$|a(0)| + \|\partial_k \widetilde{g}(0)\|_{L_k^2} + \|\langle k \rangle^2 \widetilde{g}(0)\|_{L_k^2} \le C\epsilon_0$$
 (1.40)

for some absolute constant C > 0. Then, Proposition 1.1 gives, for a possibly different C > 0, $||v||_X \le C\epsilon_0 + C||v||_X^2$.

Therefore, we see that the map defined by the first equation on the right-hand side of (1.34) and by the right-hand side of (1.38) takes a ball of radius $2C\epsilon_0$, in X, which we denote by $B_{2C\epsilon_0}^X$, into itself for some absolute C>0. Moreover, repeating the same arguments one can verify that the map is a contraction, hence, there is a unique fixed point in $B_{2C\epsilon_0}^X$.

Finally, the bound $||v||_X \lesssim \epsilon_0$ implies directly the first bound in (1.27). The other two estimates in (1.27) are obtained as a direct consequence of the pointwise and local decay estimates in Lemmas 2.5 and 2.7.

- 1.6. **Notations.** We let $\langle x \rangle := (1+x^2)^{\frac{1}{2}}$. We write $a \lesssim b$ is there is a uniform constant C such that $a \leq Cb$. Indicator functions in variable y will be denoted as $\mathbf{1}_I(y)$ meaning the function is equal to 1 whenever $y \in I$ and vanishes otherwise. To distinguish between the function h and its complex conjugate, in a number of occasions, we use $h^{(+)}(x) := h(x)$ to denote the function itself and $h^{(-)}(x) := \overline{h}(x)$ to denote the complex conjugate of h.
- 1.7. **Acknowledgments.** A. K. was supported in part by NSERC grant RGPIN 2018-04536, and thanks the Fields Institute and McMaster University for its support.
- F.P. was supported in part by a start-up grant from the University of Toronto, and NSERC grant RGPIN-2018-06487.

2. Linear Theory

In this section, we consider the second order differential operator $H := -\partial_x^2 + V$, where V(x) is a potential satisfying the following assumption

Assumption 2.1. The potential V is a smooth function exponentially decaying at infinity,

$$\lim_{|x| \to \infty} e^{\alpha|x|} V(x) = 0 \quad \text{for some } \alpha > 0, \tag{2.1}$$

and $H = -\partial_{xx} + V$ has no point spectrum, and no zero energy resonance, i.e. there is no bounded nontrivial solution in the kernel of H or, equivalently, V is 'generic' in the sense of Definition 2.1.

In our application to the equation (1.4), the role of the operator H is played by $P_c\mathcal{L}$. For an operator H as above we present the basic facts about Jost functions, reflection and transmission coefficients, and the distorted Fourier transform. This theory is based on classical results of Deift-Trubowitz [13] and Faddeev [16, 17]. See also the book of Yafaev [39].

Remark 2.1. As it was proven in [5], the potential $-4Q^3$ given in (1.5) is generic.

2.1. **Jost functions.** We define $f_+(x,k)$ and $f_-(x,k)$ as solutions to the differential equation

$$Hf_{\pm}(x,k) = k^2 f_{\pm}(x,k)$$
 (2.2)

which satisfy the asymptotic boundary conditions

$$\lim_{x \to \infty} |e^{-ikx} f_{+}(x,k) - 1| = 0 \quad \text{and} \quad \lim_{x \to -\infty} |e^{ikx} f_{-}(x,k) - 1| = 0$$
 (2.3)

We further define $m_{\pm}(x,k) := e^{\mp ikx} f_{\pm}(x,k)$. For fixed x, the functions $m_{\pm}(x,k)$ are analytic in Im k > 0 and continuous in Im $k \geq 0$. The following lemma provides with the primary estimates on m_{\pm} , which will be used later.

Lemma 2.1. For every s = 0, 1, 2 and every positive integer N, we have

$$\begin{aligned} |\partial_k^s(m_{\pm}(x,k)-1)| &\lesssim \langle x \rangle^{-12} \langle k \rangle^{-1}, & \pm x \geq -1, \\ |\partial_k^s(m_{\pm}(x,k)-1)| &\lesssim \langle x \rangle^{s+1} \langle k \rangle^{-1}, & \pm x \leq 1. \end{aligned}$$
(2.4)

Moreover, in the presence of the x derivatives, we have

$$|\partial_x^N \partial_k^s m_{\pm}(x,k)| \lesssim \langle x \rangle^{-12} \langle k \rangle^{-1+N}, \qquad \pm x \ge -1, |\partial_x^N \partial_k^s m_{\pm}(x,k)| \lesssim \langle x \rangle^{s+1} \langle k \rangle^{-1+N}, \qquad \pm x \le 1.$$
 (2.5)

Proof. The estimates follow by iterating the Volterra integral equations for m_{\pm} given by

$$m_{\pm}(x,k) = 1 \pm \int_{x}^{\pm \infty} D_{k}(\pm(y-x))V(y)m_{\pm}(y,k)dy$$
, where $D_{k}(y) = \frac{e^{2iky} - 1}{2ik}$

as in Deift-Trubowitz [13] or Appendix A in [20] under the assumption (2.1) on V.

Remark 2.2. Results similar to those in Lemma 2.1 can be derived for any $s \geq 0$, and in such case some of the bounds should be modified, see for example [6, 19]. However, we will be working only with up to second derivatives of m_{\pm} , and so are the choices of s.

The Jost functions f_{\pm} are related through so-called transmission and reflection coefficients, T(k) and $R_{\pm}(k)$, respectively [13]. That is,

$$T(k)f_{+}(x,k) = R_{-}(k)f_{-}(x,k) + f_{-}(x,-k),$$

$$T(k)f_{-}(x,k) = R_{+}(k)f_{+}(x,k) + f_{+}(x,-k),$$
(2.6)

and using the boundary conditions (2.3), we obtain

$$T(k)f_{+}(x,k) \sim R_{-}(k)e^{-ikx} + e^{ikx} \quad \text{as } x \to -\infty$$

 $T(k)f_{-}(x,k) \sim R_{+}(k)e^{ikx} + e^{-ikx} \quad \text{as } x \to \infty.$ (2.7)

The coefficients T(k) and $R_{\pm}(k)$ can be written explicitly through

$$\frac{1}{T(k)} = 1 - \frac{1}{2ik} \int V(x) m_{\pm}(x, k) dx,
\frac{R_{\pm}(k)}{T(k)} = \frac{1}{2ik} \int e^{\mp 2ikx} V(x) m_{\mp}(x, k) dx, \tag{2.8}$$

and satisfy the following relations at every k:

$$T(-k) = \overline{T(k)}, \quad R_{\pm}(-k) = \overline{R_{\pm}(k)},$$

 $|T(k)|^2 + |R_{+}(k)|^2 = 1, \quad T(k)\overline{R_{-}(k)} + \overline{T(k)}R_{+}(k) = 0.$

One distinguishes two possible asymptotics of T(k) for $k \approx 0$, depending on the value of the integral $\int_{-\infty}^{\infty} V(x)m_{+}(x,0)dx$, which is zero whenever the potential V is generic. Here are some equivalent definitions of generic potentials.

Definition 2.1. The potential V is generic if one of the following, equivalent, conditions hold:

- $\int_{-\infty}^{\infty} V(x) m_{+}(x,0) dx = 0;$
- T(0) = 0 and $R_{\pm}(0) = -1$; $H = -\partial_x^2 + V$ has no zero resonance.

For generic potentials, T(k) and $R_{\pm}(k)$ have the following asymptotics around $k \approx 0$.

Lemma 2.2. For V satisfying Assumption 2.1, the coefficients T(k) and $R_{\pm}(k)$ have the following properties:

• $as k \rightarrow 0$,

$$T(k) = \alpha k + \mathcal{O}(k^2) \quad \text{for some } \alpha \neq 0,$$

$$1 + R_{\pm}(k) = \alpha_{\pm} k + o(k).$$
(2.9)

In particular, T(0) = 0 and $R_{\pm}(0) = -1$.

• For every $k \in \mathbb{R}$ we have

$$|T(k)| + |R(k)| \lesssim 1,$$

$$|\partial_k T(k)| + |\partial_k R_+(k)| \lesssim \langle k \rangle^{-1}.$$
(2.10)

• For every $k \in \mathbb{R}$ we have

$$\left| \frac{T(k)}{k} \right| + \langle k \rangle \left| \partial_k \left(\frac{T(k)}{k} \right) \right| \lesssim \frac{1}{\langle k \rangle},$$
 (2.11)

and

$$\left| \frac{1 + R_{\pm}(k)}{k} \right| + \left| \partial_k \left(\frac{1 + R_{\pm}(k)}{k} \right) \right| \lesssim \frac{1}{\langle k \rangle}. \tag{2.12}$$

Proof. The detailed proof of the estimates (2.9)–(2.10) can be found in Deift-Trubowitz [13], so we only prove the estimates (2.11)–(2.12). It is worth pointing out that Assumption 2.1 on V is stronger than needed for this lemma to be true, and it suffices to have $\langle x \rangle^2 V \in L_x^1$.

The asymptotic of T(k) around $k \approx 0$ imply that there exists a positive constant k^* such that

$$\left| \frac{T(k)}{k} \right| \lesssim \frac{1}{\langle k \rangle}$$
 for every $|k| \leq k^*$.

On the other hand, for $|k| > k^*$, we use uniform boundedness of |T(k)| to get

$$\left| \frac{T(k)}{k} \right| \lesssim \frac{|T(k)|}{\langle k \rangle} \lesssim \frac{1}{\langle k \rangle}.$$

Combining these estimates together we obtain the first part of (2.11). To get the bound on the derivative part, we note that

$$\partial_k \left(\frac{T(k)}{k} \right) = \left(\frac{T(k)}{k} \right)^2 \left(\frac{1}{2i} \int V(x) \partial_k m_{\pm}(x,k) dx - 1 \right),$$

which leads to a desired estimate in the view of (2.4) and assumptions on V.

2.2. **Pseudo-differential bounds.** We provide some $L^2 \to L^2$ bounds for pseudodifferential operators whose symbols include $m_{\pm}(x,k)$ and their derivatives. Under milder assumptions than ours, which in particular do not require any regularity for V, these types of bounds are given in [6].

Lemma 2.3. For every function $a \in L_k^{\infty}$ we have

$$\left\| \int \mathbf{1}_{[-1,\infty)} e^{ikx} a(k) (m_+(x,k) - 1) h(x) dx \right\|_{L^2_k} \lesssim \left\| \langle x \rangle^{-11} h \right\|_{L^2_x}, \tag{2.13}$$

and

$$\left\| \int \mathbf{1}_{[-1,\infty)} e^{ikx} a(k) \partial_k m_+(x,k) h(x) dx \right\|_{L^2_k} \lesssim \|\langle x \rangle^{-10} h \|_{L^2_x}. \tag{2.14}$$

Similar estimates hold for $\mathbf{1}_{(-\infty,1]}(m_{-}(x,k)-1)$ and $\mathbf{1}_{(-\infty,1]}\partial_k m_{-}(x,k)$, respectively.

Proof. Using the estimates (2.4), we have that the left-hand side of (2.13) is bounded by

$$\left\| \int \langle x \rangle^{-12} \langle k \rangle^{-1} |h(x)| dx \right\|_{L^2_k} \lesssim \left\| \langle x \rangle^{-11} h \right\|_{L^2_x} \left\| \langle k \rangle^{-1} \right\|_{L^2_k} \lesssim \left\| \langle x \rangle^{-11} h \right\|_{L^2_x},$$

where we used the Cauchy-Schwartz inequality for dx integral. The estimate (2.14) follows in a similar fashion using (2.5).

2.3. **Distorted Fourier transform.** In this subsection, we present the generalization of the standard Fourier transform which we apply later to the perturbed Schrödinger operator $H = -\partial_x^2 + V$. Throughout the section and further in the paper, we use the notation

$$\widetilde{D} := \sqrt{-\partial_x^2 + V}, \qquad D := \sqrt{-\partial_x^2} = |\partial_x|.$$

The standard Fourier transform on the line is defined for every $\phi \in L^2$ as

$$\widehat{\mathcal{F}}\phi(k) = \widehat{\phi}(k) := \frac{1}{\sqrt{2\pi}} \int e^{-ikx} \phi(k) \, dx,$$

with

$$\widehat{\mathcal{F}}^{-1}\phi(x) := \frac{1}{\sqrt{2\pi}} \int e^{ikx} \phi(k) dx.$$

When dealing with the operator H, the function e^{-ikx} inside the standard Fourier transform can be replaced by a generalized eigenfunction $\psi(x,k)$ given in terms of the Jost functions f_{\pm} from (2.2) in the following way:

$$\psi(x,k) := \frac{1}{\sqrt{2\pi}} \begin{cases} T(k)f_{+}(x,k) & \text{for } k \ge 0, \\ T(-k)f_{-}(x,-k) & \text{for } k < 0 \end{cases}$$
 (2.15)

and we now introduce the distorted Fourier transform for $h \in \mathcal{S}$ by

$$\widetilde{\mathcal{F}}h(k) = \widetilde{h}(k) := \int \overline{\psi(x,k)}h(x) dx.$$
 (2.16)

Lemma 2.4. In above settings, $\widetilde{\mathcal{F}}: L_x^2 \to L_k^2$ is an isometry,

$$\|\widetilde{\mathcal{F}}h\|_{L_k^2} = \|P_c h\|_{L_x^2}, \quad \text{for all } h \in L^2.$$
 (2.17)

The inverse of the distorted Fourier transform is given by

$$\widetilde{\mathcal{F}}^{-1}g(x) = \int \psi(x,k)g(k)dk.$$

Setting $\widetilde{D} := \sqrt{-\partial_{xx} + V}$, we have

$$m(\widetilde{D})P_c = \widetilde{\mathcal{F}}^{-1}m(k)\widetilde{\mathcal{F}}.$$

Moreover, the distorted Fourier transform satisfies the following properties:

- $\widetilde{\mathcal{F}}\widetilde{\mathcal{F}}^{-1} = \mathrm{Id}_{L^2_k} \text{ and } \widetilde{\mathcal{F}}^{-1}\widetilde{\mathcal{F}} = P_c;$
- if $h \in L^1_x$, then \widetilde{h} is a continuous, bounded function;
- since V is generic, $\widetilde{h}(0) = 0$;
- we have

$$\|\partial_k \widetilde{h}\|_{L^2_k} \lesssim \|\langle x \rangle h\|_{L^2_x},\tag{2.18}$$

and, for s > 0

$$\|\langle k \rangle^s \widetilde{h}\|_{L^2_L} = \|P_c h\|_{H^s_x}$$
 (2.19)

Proof. The proofs can be found in Agmon [1], Dunford-Schwartz [15], Germain-Pusateri-Rousset [20] and Yafaev [39]. \Box

2.4. **Decomposition of the generalized eigenfunctions.** By the definition of the generalized eigenfunction we have

for
$$k \ge 0$$
, $\sqrt{2\pi}\psi(x,k) = T(k)f_{+}(x,k) = T(k)m_{+}(x,k)e^{ikx}$
for $k < 0$, $\sqrt{2\pi}\psi(x,k) = T(-k)f_{-}(x,-k) = T(-k)m_{-}(x,-k)e^{ikx}$ (2.20)

We introduce smooth versions of the indicator functions, σ_+ and σ_- ,

$$\sigma_{+}(x) = \int_{-\infty}^{x} \varphi(y)dy$$
, and $\sigma_{+}(x) + \sigma_{-}(x) = 1$, (2.21)

where φ is a smooth, nonnegative function, which is equal to 1 in a neighborhood of 0, vanishes outside of [-2,2] and $\int \varphi \ dx = 1$. By construction, $\sigma_+(x)$ vanishes for large negative values of x while $\sigma_-(x)$ vanishes for large positive values of x. These functions allow to decompose expressions in (2.20) as

for
$$k \ge 0$$
, $\sqrt{2\pi}\psi(x,k) = \sigma_+(x)T(k)f_+(x,k) + \sigma_-(x)\left[f_-(x,-k) + R_-(k)f_-(x,k)\right]$,
for $k < 0$, $\sqrt{2\pi}\psi(x,k) = \sigma_-(x)T(-k)f_-(x,-k) + \sigma_+(x)\left[f_+(x,k) + R_+(-k)f_+(x,-k)\right]$,

where we used the relations (2.6). We further rewrite the generalized eigenfunction above using $m_{\pm}(x,k) = f_{\pm}(x,k)e^{\mp ikx}$,

for
$$k \ge 0$$
,

$$\sqrt{2\pi}\psi(x,k) = \sigma_{+}(x)T(k)m_{+}(x,k)e^{ikx} + \sigma_{-}(x)\left[m_{-}(x,-k)e^{ikx} + R_{-}(k)m_{-}(x,k)e^{-ikx}\right]$$
for $k < 0$,

$$\sqrt{2\pi}\psi(x,k) = \sigma_{-}(x)T(-k)m_{-}(x,-k)e^{ikx} + R_{-}(k)m_{-}(x,k)e^{-ikx}$$

$$+ \sigma_{+}(x)\left[m_{+}(x,k)e^{ikx} + R_{+}(-k)m_{+}(x,-k)e^{-ikx}\right],$$
(2.22)

Remark 2.3. Based on the boundedness of $\sigma_{\pm}m_{\pm}$ in (2.4), T(k) and $R_{\pm}(k)$ in (2.10),

$$|\psi(x,k)| \lesssim 1. \tag{2.23}$$

In the view of (2.4), we even further decompose ψ into a singular and regular parts as follows

$$\sqrt{2\pi}\psi(x,k) = \psi_S(x,k) + \psi_R(x,k). \tag{2.24}$$

The singular part ψ_S is given by

$$\psi_S(x,k) := \sigma_+(x)\psi_{S,+}(x,k) + \sigma_-(x)\psi_{S,-}(x,k), \tag{2.25}$$

where, setting the indicator functions $\mathbf{1}_{+} := \mathbf{1}_{[0,\infty)}$ and $\mathbf{1}_{-} := \mathbf{1}_{(-\infty,0)}$,

$$\begin{cases}
\psi_{S,+}(x,k) := \mathbf{1}_{+}(k)T(k)e^{ikx} + \mathbf{1}_{-}(k)\left[e^{ikx} + R_{+}(-k)e^{-ikx}\right], \\
\psi_{S,-}(x,k) := \mathbf{1}_{+}(k)\left[e^{ikx} + R_{-}(k)e^{-ikx}\right] + \mathbf{1}_{-}(k)T(-k)e^{ikx}.
\end{cases} (2.26)$$

The rest is a regular part $\psi_R(x,k)$ given by

he rest is a regular part
$$\psi_R(x,k)$$
 given by
$$\psi_R(x,k) := \begin{cases} \sigma_+(x)T(k)(m_+(x,k)-1)e^{ikx} \\ + \sigma_-(x)[(m_-(x,-k)-1)e^{ikx} + R_-(k)(m_-(x,k)-1)e^{-ikx}], & k \ge 0 \end{cases}$$

$$\sigma_-(x)T(-k)(m_-(x,-k)-1)e^{ikx} + \sigma_+(x)[(m_+(x,k)-1)e^{ikx} + R_+(-k)(m_+(x,-k)-1)e^{-ikx}], & k < 0. \tag{2.27}$$

Note that the regular part is localized around x = 0 due to the estimates (2.4). On the other hand, the singular part ψ_S is a combination of oscillatory terms given by exponential functions with bounded coefficients, see (2.26). From (2.26), we can further rewrite the elements of the singular part as

$$\begin{cases}
\psi_{S,+}(x,k) := a_{+,+}(k)e^{ikx} + a_{+,-}(k)e^{-ikx}, \\
\psi_{S,-}(x,k) := a_{-,+}(k)e^{ikx} + a_{-,-}(k)e^{-ikx},
\end{cases}$$
(2.28)

where

$$a_{+,+}(k) = \mathbf{1}_{+}(k)T(k) + \mathbf{1}_{-}(k), \qquad a_{+,-}(k) = \mathbf{1}_{-}(k)R_{+}(-k), a_{-,+}(k) = \mathbf{1}_{+}(k) + \mathbf{1}_{-}(k)T(-k), \qquad a_{-,-}(k) = \mathbf{1}_{+}(k)R_{-}(k).$$
(2.29)

As it would be useful later, we also point out that, given $h \in P_cL_x^2$, the decomposition (2.24) allows to rewrite

$$h = h_S + h_R$$
, where $h_* := \frac{1}{\sqrt{2\pi}} \int \psi_*(x,k) \tilde{h}(k) dk$ for $* \in \{S,R\}$. (2.30)

Remark 2.4. We point out that the outcomes of Lemma 2.3 also apply to ψ_R . In other words, based on the construction of ψ_R in (2.27), we can verify that

$$\left\| \int \overline{\psi_R(x,k)} h(x) dx \right\|_{L^2_h} \lesssim \left\| \langle x \rangle^{-11} h \right\|_{L^2_x}, \tag{2.31}$$

and

$$\left\| \int \partial_k \overline{\psi_R(x,k)} h(x) dx \right\|_{L^2_k} \lesssim \left\| \langle x \rangle^{-10} h \right\|_{L^2_x}. \tag{2.32}$$

Moreover, the function ψ_R decays as

$$|\psi_R(x,k)| \lesssim \langle x \rangle^{-12} \langle k \rangle^{-1}.$$
 (2.33)

2.5. Linear estimates and local decay. In this section we provide linear and local decay estimates for the term

$$\langle \widetilde{D} \rangle^{-1} e^{\pm it \langle \widetilde{D} \rangle} P_c f,$$
 (2.34)

where $\langle \widetilde{D} \rangle = \sqrt{H+1} = \widetilde{\mathcal{F}}^{-1} \langle k \rangle \widetilde{\mathcal{F}}$. These estimates will be used in later sections to bound the term $\chi(t,x)$ in (1.31).

Lemma 2.5. Under Assumption 2.1 on V, for every $t \ge 0$ and every N = 0, 1, 2 we have

$$\left\| \partial_x^N \langle \widetilde{D} \rangle^{-1} e^{\pm it \langle \widetilde{D} \rangle} P_c f \right\|_{L_x^{\infty}} \lesssim \langle t \rangle^{-\frac{1}{2} + \frac{N}{5}} \left(\| \partial_k \widetilde{f} \|_{L_k^2} + \| \langle k \rangle^2 \widetilde{f} \|_{L_k^2} \right) \tag{2.35}$$

The proof of Lemma 2.5 is standard and follows from similar, in fact easier, arguments to those in [19, Proposition 3.10] since here we are not dealing with any bad frequency. In particular, using also the bound $\|\langle k \rangle h\|_{L_k^{\infty}} \lesssim \|\partial_k h\|_{L_k^2} + \|\langle k \rangle^2 h\|_{L_k^2}$, one can deduce from [19, Proposition 3.10] that

$$\left\| \langle \widetilde{D} \rangle^{-1} e^{\pm it \langle \widetilde{D} \rangle} P_c f \right\|_{L_x^{\infty}} \lesssim \langle t \rangle^{-\frac{1}{2}} \left(\|\partial_k \widetilde{f}\|_{L_k^2} + \|\langle k \rangle^2 \widetilde{f}\|_{L_k^2} \right). \tag{2.36}$$

The estimate (2.35) with N=1,2 can be proven with the same argument, or also using Gagliardo-Nirenberg-Sobolev interpolation and (2.19).

Note that in [19] the decay rates (for N=0) associated to the L^2 norms on the right-hand side of (2.35) are faster, and needed to compensate the small growth of these norms over the time. In our case, we do not have such a growth, so (2.35) suffices for our purposes.

Remark 2.5. Lemma 2.5 can be repeated for V = 0. In such case, \widetilde{D} is replaced by D and the distorted Fourier transform becomes the standard Fourier transform. The corresponding estimate is

$$\|\langle D\rangle^{-1}e^{\pm it\langle D\rangle}f\|_{L_x^{\infty}} \lesssim \langle t\rangle^{-1/2}\left(\|\partial_k \widehat{f}\|_{L_k^2} + \|\langle k\rangle^2 \widehat{f}\|_{L_k^2}\right). \tag{2.37}$$

Next, we state the local decay estimates which will be used in later sections. We start with the estimates for the standard Fourier transform under additional assumption at k = 0.

Lemma 2.6. Suppose $\hat{h} \in H_k^1$ and $\hat{h}(0) = 0$, then for every $t \geq 0$ we have

$$\left\| \langle x \rangle^{-1} \langle D \rangle^{-1} e^{\pm it \langle D \rangle} h \right\|_{L^{\infty}} \lesssim \langle t \rangle^{-3/4} \|\widehat{h}\|_{H^{1}_{k}}. \tag{2.38}$$

The proof of (2.38) is standard, and can be found in [6, Lemma 3.6] in the completely analogous case of the Schrödinger equation.

Estimates for the distorted Fourier transform have better decays rates under genericity assumption on V.

Lemma 2.7. Suppose V satisfies Assumption 2.1. Then, for every $t \geq 0$, we have

$$\left\| \langle x \rangle^{-2} \langle \widetilde{D} \rangle^{-1} e^{\pm it \langle \widetilde{D} \rangle} P_c h \right\|_{L_x^{\infty}} \lesssim \langle t \rangle^{-1} \|\widetilde{h}\|_{H_k^1}. \tag{2.39}$$

The proof of this result is also standard and can be found in [21]. An analogous estimate for the Schrödinger flow is given in [6, Lemma 3.6]; see also the references in [21, 6] for other related and more general estimates.

We can extend the result of the above lemma to cases when we deal with the singular and regular parts of the function $\langle \widetilde{D} \rangle^{-1} e^{\pm it \langle \widetilde{D} \rangle} P_c h$ according to the decomposition in (2.30). First, we have the following analogue of Lemma 2.7:

Lemma 2.8. Suppose V satisfies Assumption 2.1. Then, for every $t \geq 0$, we have

$$\left\| \langle x \rangle^{-2} \left(\langle \widetilde{D} \rangle^{-1} e^{\pm it \langle \widetilde{D} \rangle} P_c h \right)_S \right\|_{L_x^{\infty}} \lesssim \langle t \rangle^{-1} \|\widetilde{h}\|_{H_k^1}, \tag{2.40}$$

For the regular part, the function ψ_R is already decaying according to (2.33), which allows to get estimates with growing weight.

Lemma 2.9. Suppose V satisfies Assumption 2.1. Then, for every $t \geq 0$, we have

$$\left\| \langle x \rangle^{10} \left(\langle \widetilde{D} \rangle^{-1} e^{\pm it \langle \widetilde{D} \rangle} P_c h \right)_R \right\|_{L_{\infty}^{\infty}} \lesssim \langle t \rangle^{-1} \|\widetilde{h}\|_{H_k^1}. \tag{2.41}$$

3. Set-up and bootstrap

In this section we set-up the upcoming analysis of the evolution equations for a(t) and $\chi(t, x)$ in (1.31), and state our main bootstrap propositions that will imply Proposition 1.1.

We let $H := -\partial_{xx} + V(x)$ with $V(x) := -4Q^3$, where Q(x) is the soliton (1.3). Then the operator (1.5) is $\mathcal{L} = H + 1$. Recall the definition of $g(t,x) := e^{it\sqrt{H+1}}(\partial_t - i\sqrt{H+1})\chi(t,x)$ in (1.35), so that χ is given as in (1.36) Recall also the notation for our main norm (1.39),

$$||v||_X := ||\langle t \rangle^2 a(t)||_{L_t^{\infty}(0,\infty)} + ||\partial_k \widetilde{g}||_{L_t^{\infty}((0,\infty):L_k^2)} + ||\langle k \rangle^2 \widetilde{g}||_{L_t^{\infty}((0,\infty):L_k^2)}.$$
(3.1)

Remark 3.1. From Lemma 2.5 and Lemma 2.7 the following estimates for χ follow

$$\|\partial_x^N \chi(t)\|_{L_x^{\infty}} \lesssim \langle t \rangle^{-\frac{1}{2} + \frac{N}{5}} \left(\|\partial_k \widetilde{g}\|_{L_k^2} + \|\langle k \rangle^2 \widetilde{g}\|_{L_k^2} \right) \lesssim \langle t \rangle^{-\frac{1}{2} + \frac{N}{5}} \|v\|_X, \qquad N = 0, 1, 2, \tag{3.2}$$

and

$$\left\| \langle x \rangle^{-2} \chi(t) \right\|_{L_x^{\infty}} \lesssim \langle t \rangle^{-1} \|\widetilde{g}\|_{H_k^1} \lesssim \langle t \rangle^{-1} \|v\|_X. \tag{3.3}$$

Applying the distorted Fourier transform to the relation (1.35), we obtain $\widetilde{g}(t,k) = e^{it\langle k \rangle}(\partial_t - i\langle k \rangle)\widetilde{\chi}(t,k)$ and, after substitution into the second equation in (1.31),

$$\partial_t \widetilde{g}(t,k) = e^{it\langle k \rangle} \widetilde{\mathcal{F}} P_c N(Q,v), \tag{3.4}$$

where $N_c(Q, v) = P_c(6Q^2v^2 + 4Qv^3 + v^4)$ and we recall that the term v is decomposed as $v(t, x) = a(t)\rho(x) + \chi(t, x)$. Integrating the equation (3.4) over time give

$$\widetilde{g}(t,k) = \widetilde{g}(0,k) + \int_0^t e^{is\langle k \rangle} \widetilde{\mathcal{F}} P_c N(s,k) ds,$$
(3.5)

and applying the partial derivative with respect to k, we obtain

$$\partial_k \widetilde{g}(t,k) = \partial_k \widetilde{g}(0,k) + \int_0^t \partial_k \left[e^{is\langle k \rangle} \widetilde{\mathcal{F}} P_c N(s,k) \right] ds. \tag{3.6}$$

In the course of our estimates we will need to deal with $\|\partial_t \widetilde{g}(t)\|_{L^2_k}$, which can be bounded using the following

Lemma 3.1. Given $||v||_X \ll 1$, we have

$$\|\partial_t \widetilde{g}\|_{L^2_k} \lesssim \langle t \rangle^{-3/2} \|v\|_X^2 \tag{3.7}$$

Proof. From the evolution equation (3.4), since $\widetilde{\mathcal{F}}$ is an $L_x^2 \to L_k^2$ isometry, we have

$$\|\partial_t \widetilde{g}\|_{L^2_{\iota}} \lesssim \|Q^2 v^2\|_{L^2_x} + \|Q v^3\|_{L^2_x} + \|v^4\|_{L^2_x}.$$

We substitute the decomposition $v(t,x) = a(t)\rho(x) + \chi(t,x)$, and bound the first two terms using the local decay (3.3) as

$$||Q^2v^2||_{L^2_x} \lesssim |a(t)|^2 + ||\langle x\rangle^{-2}\chi||_{L^\infty}^2 \lesssim \langle t\rangle^{-2}||v||_X^2,$$

with a similar estimate for $||Qv^3||_{L^2_x}$. For the remaining term we have

$$||v^4||_{L^2_x} \lesssim |a(t)|^4 + ||\chi||_{L^2_x}^4 \lesssim \langle t \rangle^{-8} ||v||_X^4 + ||\chi||_{L^2_x}^4 ||\chi||_{L^\infty_x}^3.$$

The term $\|\chi\|_{L^{\infty}_x}$ is bounded by (3.2), while $\|\chi\|_{L^2_x} \lesssim \|\widetilde{g}\|_{L^2_k} \leq \|v\|_X$, using (2.17). Then, we have $\|v^4\|_{L^2_x} \lesssim \langle t \rangle^{-3/2} \|v\|_X^4$, and combining all the estimates we obtain the desired bound (3.7).

In the following subsections, we give estimates on the three norms in (3.1).

3.1. Estimates for the first norm in (3.1).

Proposition 3.1. Given $||v||_X \ll 1$, for every $t \geq 0$ we have

$$\langle t \rangle^2 |a(t)| \lesssim |a(0)| + ||v||_X^2.$$
 (3.8)

Proof. Using the integral equation for a(t) in (1.34), with F[v] as in (1.31), we have

$$\langle t \rangle^2 a(t) = \langle t \rangle^2 e^{-\Omega t} \left[a(0) + \frac{1}{2\Omega} \int_0^\infty e^{-\Omega s} F[v](s) ds \right]$$
$$- \frac{\langle t \rangle^2 e^{-\Omega t}}{2\Omega} \int_0^t e^{\Omega s} F[v](s) ds - \frac{\langle t \rangle^2}{2\Omega} \int_t^\infty e^{-\Omega(s-t)} F[v](s) ds$$
$$:= \mathcal{I}_1(t) + \mathcal{I}_2(t) + \mathcal{I}_3(t).$$

To bound F[v](s) we use the definition in (1.5) and the decomposition of v in (1.7) to get

$$|F[v](s)| \lesssim |\langle v^2(s), \rho \rangle| + |\langle v^3(s), \rho \rangle| + |\langle v^4(s), \rho \rangle| \lesssim ||\langle x \rangle^{-2} v(s)||_{L^{\infty}}^2 \lesssim \langle s \rangle^{-2} ||v||_X^2, \tag{3.9}$$

where for the last inequality we used (3.3). Then, using (3.9), direct computations show that

$$|\mathcal{I}_1(t)| \lesssim |a(0)| + ||v||_X^2$$
, and $|\mathcal{I}_2(t)|, |\mathcal{I}_3(t)| \lesssim ||v||_X^2$.

Combining these estimates, we obtain the desired result.

3.2. Estimates for the second norm in (3.1). Here, we give the L_k^2 bound for $\partial_k \widetilde{g}$.

Proposition 3.2. Given $||v||_X \ll 1$, for every $t \ge 0$ we have

$$\|\partial_k \widetilde{g}(t)\|_{L^2_k} \lesssim \|\partial_k \widetilde{g}(0)\|_{L^2_k} + \|v\|_X^2. \tag{3.10}$$

The proof is based on the estimates for the terms in the Duhamel expression for $\partial_k \widetilde{g}$ in (3.6). Recall that, according to the decomposition $v(s,x) = a(s)\rho(x) + \chi(s,x)$, we have

$$\widetilde{\mathcal{F}}N_c(s,k) = \widetilde{\mathcal{F}}P_c\left(6Q^2(a(s)\rho + \chi(s))^2 + 4Q(a(s)\rho + \chi(s))^3 + (a(s)\rho + \chi(s))^4\right). \tag{3.11}$$

Then the L_k^2 bound for $\partial_k \widetilde{g}$ is given by

$$\|\partial_k \widetilde{g}(t)\|_{L^2_k} \lesssim \|\partial_k \widetilde{g}(0)\|_{L^2_k} + \|A_1(t)\|_{L^2_k} + \|A_2(t)\|_{L^2_k} + \|A_3(t)\|_{L^2_k}, \tag{3.12}$$

where

$$A_{1}(t,k) := \int_{0}^{t} \partial_{k} \left[e^{is\langle k \rangle} \widetilde{\mathcal{F}} P_{c} Q^{2} (a(s)\rho + \chi(s))^{2}(k) \right] ds,$$

$$A_{2}(t,k) := \int_{0}^{t} \partial_{k} \left[e^{is\langle k \rangle} \widetilde{\mathcal{F}} P_{c} Q(a(s)\rho + \chi(s))^{3}(k) \right] ds,$$

$$A_{3}(t,k) := \int_{0}^{t} \partial_{k} \left[e^{is\langle k \rangle} \widetilde{\mathcal{F}} P_{c} (a(s)\rho + \chi(s))^{4}(k) \right] ds.$$

$$(3.13)$$

To prove Proposition 3.2, it suffices to show that $||A_j(t)||_{L^2_k} \lesssim ||v||_X$ for every j = 1, 2, 3. We will verify these estimates in the following lemmas as well as in Section 4 and Section 5.

Lemma 3.2. Given $||v||_X \ll 1$, for every $t \geq 0$ we have

$$||A_1(t)||_{L^2_k} \lesssim \left\| \int_0^t s \frac{k}{\langle k \rangle} e^{is\langle k \rangle} \widetilde{\mathcal{F}} P_c Q^2 \chi^2(s) ds \right\|_{L^2_k} + ||v||_X^2. \tag{3.14}$$

Proof. Distributing ∂_k and using Minkowski inequality, we obtain

$$||A_1(t)||_{L_k^2} \lesssim \int_0^t ||\partial_k \widetilde{\mathcal{F}} P_c Q^2(a(s)\rho + \chi(s))^2||_{L_k^2} ds$$

$$+ ||\int_0^t s \frac{k}{\langle k \rangle} e^{is\langle k \rangle} \widetilde{\mathcal{F}} P_c Q^2(a(s)\rho + \chi(s))^2 ds||_{L_k^2}$$

$$=: A_{11} + A_{12}.$$

The first term A_{11} is easily bounded using (2.18), followed by the local decay estimates (3.3):

$$A_{11} \lesssim \int_0^t \|\langle x \rangle Q^2(a(s)\rho + \chi(s))^2\|_{L_x^2} ds$$

$$\lesssim \int_0^t a^2(s) ds + \int_0^t \|\langle x \rangle^{-2} \chi(s)\|_{L_x^\infty}^2 ds \lesssim \|v\|_X^2.$$

For the second term A_{12} we bound it similarly, term by term, as follows

$$A_{12} \lesssim \int_0^t s \ a^2(s) \left\| \widetilde{\mathcal{F}} P_c Q^2 \rho^2 \right\|_{L_k^2} ds + \int_0^t s \ a(s) \left\| \widetilde{\mathcal{F}} P_c Q^2 \rho \chi(s) \right\|_{L_k^2} ds$$
$$+ \left\| \int_0^t s \frac{k}{\langle k \rangle} e^{is\langle k \rangle} \widetilde{\mathcal{F}} P_c Q^2 \chi^2(s) ds \right\|_{L_k^2}$$
$$:= B_1 + B_2 + B_3.$$

Using (2.17) and (3.3) we obtain

$$B_{1} \lesssim \|v\|_{X}^{2} \int_{0}^{t} \frac{s}{\langle s \rangle^{4}} ds \lesssim \|v\|_{X}^{2},$$

$$B_{2} \lesssim \int_{0}^{t} s \ a(s) \|\langle x \rangle^{-2} \chi(s)\|_{L_{x}^{2}} ds \lesssim \|v\|_{X}^{2} \int_{0}^{t} \frac{s}{\langle s \rangle^{3}} ds \lesssim \|v\|_{X}^{2}.$$

This completes the proof.

Remark 3.2. Note that in the proof of Lemma 3.2, the term B_3 can not be bounded simply using Minkowski's inequality, as the corresponding integral in ds would diverge when $t \to \infty$. Indeed, we would have

$$B_3 \lesssim \int_0^t s \|\langle x \rangle^{-2} \chi(s)\|_{L_k^2}^2 ds \lesssim \|v\|_X^2 \int_0^t \frac{s}{\langle s \rangle^2} ds \approx \|v\|_X^2 \log t. \tag{3.15}$$

To get a $O(\|v\|_X^2)$ bound for the term $\|A_1(t)\|_{L_k^2}$, we will need some more refined analysis, which will be provided in Section 4.

Lemma 3.3. Given $||v||_X \ll 1$, for every $t \geq 0$ we have

$$||A_2(t)||_{L^2_k} \lesssim ||v||_X^3.$$
 (3.16)

Proof. The proof is similar to the proof of Lemma 3.2. However, unlike in (3.15), each term, which appears after applying ∂_k inside the $A_2(t)$ integral in (3.13) and expanding the cubic term $(a(s)\rho + \chi(s))^3$, is directly bounded by $||v||_X^3$.

Lemma 3.4. Given $||v||_X \ll 1$, for every $t \ge 0$ we have

$$||A_3(t)||_{L^2_k} \lesssim \left\| \int_0^t \partial_k \left[e^{is\langle k \rangle} \widetilde{\mathcal{F}} P_c \chi^4(s) \right] ds \right\|_{L^2_k} + ||v||_X^4.$$
 (3.17)

Proof. The proof is similar to the proof of previous lemmas, using the fact that $\rho(x)$ is an exponentially decaying function and (3.3).

Remark 3.3. Since there are no (exponentially) decaying function in front of χ^4 we cannot just rely on the local decay estimates (3.3) to bound the quartic terms on the left-hand side of (3.17). We will bound these terms in Section 5, using some multilinear analysis in the spirit of [6, 19].

The proof of Proposition 3.2 is now straightforward assuming that both the quadratic term in (3.14) and the quartic term in (3.17) are bounded by $||v||_X^2$. These two bounds will be proven, respectively, in the upcoming Sections 4 and 5.

Proof of Proposition 3.2. Under the above assumption, Lemmas 3.2–3.4 imply

$$||A_1(t)||_{L_k^2} + ||A_2(t)||_{L_k^2} + ||A_3(t)||_{L_k^2} \lesssim ||v||_X^2.$$

This bound applied to (3.12) completes the proof.

3.3. Estimates for the third norm in (3.1). Here we give the L_k^2 bound for $\|\langle k \rangle^2 \widetilde{g}\|_{L_t^2}$.

Proposition 3.3. For every $t \ge 0$ we have

$$\|\langle k \rangle^2 \widetilde{g}(t)\|_{L^2_h} \lesssim \|\langle k \rangle^2 \widetilde{g}(0)\|_{L^2_h} + \|v\|_X^2. \tag{3.18}$$

To prove the above statement, we use the Duhamel's expression (3.5) apply Minkowski, and recall (1.5), to get

$$\|\langle k \rangle^{2} \widetilde{g}(t)\|_{L_{k}^{2}} \lesssim \|\langle k \rangle^{2} \widetilde{g}(0)\|_{L_{k}^{2}} + \int_{0}^{t} \|\langle k \rangle^{2} \widetilde{\mathcal{F}} P_{c} Q^{2} v^{2}(s)\|_{L_{k}^{2}} ds + \int_{0}^{t} \|\langle k \rangle^{2} \widetilde{\mathcal{F}} P_{c} Q v^{3}(s)\|_{L_{k}^{2}} ds + \int_{0}^{t} \|\langle k \rangle^{2} \widetilde{\mathcal{F}} P_{c} v^{4}(s)\|_{L_{k}^{2}} ds$$

$$:= \|\langle k \rangle^{2} \widetilde{g}(0)\|_{L_{k}^{2}} + \mathcal{I}_{1}(t) + \mathcal{I}_{2}(t) + \mathcal{I}_{3}(t).$$

$$(3.19)$$

Therefore, it suffices to show the following:

Lemma 3.5. For every t > 0 we have

$$\mathcal{I}_{j}(t) \lesssim ||v||_{X}^{2} \quad \text{for every } j = 1, 2, 3.$$
 (3.20)

Proof. Starting with the quadratic terms, we recall the definition of the norm (3.1), and estimate using (2.19) followed by the pointwise and local decay estimates (3.2) and (3.3):

$$\begin{split} \left\| \langle k \rangle^2 \widetilde{\mathcal{F}} P_c Q^2 v^2(s) \right\|_{L^2_k} &\lesssim \| Q^2 v^2(s) \|_{H^2_x} \lesssim \| Q v(s) \|_{H^2_x} \| Q v(s) \|_{L^\infty_x} \\ &\lesssim \sup_{j=0,1,2} \| \langle x \rangle^{-5} \partial_x^j v(s) \|_{L^2_x} \| Q v(s) \|_{L^\infty_x} \lesssim \langle s \rangle^{-1/10} \| v \|_X \langle s \rangle^{-1} \| v \|_X, \end{split}$$

which proves the desired bound for j = 1. The estimates for the cubic term \mathcal{I}_2 is simpler so we can skip it. For \mathcal{I}_3 and use the decay (3.2) to see that

$$\left\| \langle k \rangle^2 \widetilde{\mathcal{F}} P_c v^4(s) \right\|_{L^2_k} \lesssim \left\| v^4(s) \right\|_{H^2_2} \lesssim \left\| v(s) \right\|_{H^2_x} \left\| v(s) \right\|_{L^\infty_x}^3 \lesssim \langle s \rangle^{-3/2} \| v \|_X^4.$$

4. Quadratic term analysis

In this section we prove the following bound for the quadratic terms on the right-hand side of (3.14) that were not treated in the previous section.

Lemma 4.1. Given $||v||_X \ll 1$, for every $t \geq 0$ we have

$$\left\| \int_0^t s \frac{k}{\langle k \rangle} e^{is\langle k \rangle} \widetilde{\mathcal{F}} P_c Q^2 \chi^2(s) ds \right\|_{L^2_h} \lesssim \|v\|_X^2.$$

We split the integral into two integrals over separate regions $0 \le s \le 1$ and s > 1, and obtain

$$\left\| \int_{0}^{t} s \frac{k}{\langle k \rangle} e^{is\langle k \rangle} \widetilde{\mathcal{F}} P_{c} Q^{2} \chi^{2}(s) ds \right\|_{L_{k}^{2}} \lesssim \left\| \int_{0}^{1} s \frac{k}{\langle k \rangle} e^{is\langle k \rangle} \widetilde{\mathcal{F}} Q^{2} \chi^{2}(s) ds \right\|_{L_{k}^{2}} + \left\| \int_{1}^{t} s \frac{k}{\langle k \rangle} e^{is\langle k \rangle} \widetilde{\mathcal{F}} Q^{2} \chi^{2}(s) ds \right\|_{L_{k}^{2}} =: C_{1} + C_{2}.$$

$$(4.1)$$

Of course, if $t \leq 1$ then the term C_2 is not present and the term C_1 corresponds to the integration over the region 0 < s < t and can be treated in the same way below. Using the local decay (3.3) we can bound

$$C_1 \lesssim \int_0^1 s \|\widetilde{\mathcal{F}}Q^2\chi^2(s)\|_{L_k^2} ds \lesssim \int_0^1 s \|\langle x \rangle^{-2}\chi(s)\|_{L_x^\infty}^2 \lesssim \|v\|_X^2,$$
 (4.2)

To get a bound for the term C_2 we use the following smoothing-type estimate:

Lemma 4.2. Given a function $Q : \mathbb{R}_x \times \mathbb{R}_k \to \mathbb{C}$ such that, for some $\beta \in \mathbb{R}$,

$$\sup_{x \in \mathbb{R}, k \in \mathbb{R}} \left| \langle x \rangle^{\beta} \mathcal{Q}(x, k) \right| < \infty,$$

and $\phi : \mathbb{R} \to \mathbb{C}$ such that $|\phi(k)| \lesssim |k|/\langle k \rangle$, then, for all $t \geq 0$,

$$\left\| \langle x \rangle^{\beta} \int_{\mathbb{R}} \mathcal{Q}(x,k) \phi(k) e^{i\langle k \rangle s} h(k) dk \right\|_{L_{x}^{\infty} L_{s}^{2}([0,t])} \lesssim \|h\|_{L^{2}}. \tag{4.3}$$

Under the same assumptions, the following inhomogeneous estimate holds:

$$\left\| \int_0^t \left[\int_{\mathbb{R}} e^{-i\langle k \rangle s} \overline{\phi}(k) \overline{\mathcal{Q}}(x,k) F(s,x) \, dx \right] \, ds \right\|_{L_k^2} \lesssim \left\| \langle x \rangle^{-\beta} F \right\|_{L_x^1 L_s^2([0,\infty])}. \tag{4.4}$$

We now complete the proof of Lemma 4.1 using Lemma 4.2 above.

Proof of Lemma 4.1. Applying (4.4) to the term C_2 in (4.1), that is, letting $\beta = 0$, $\mathcal{Q}(y, x) = \psi(x, k)$, $\phi(k) = k/\langle k \rangle$, and $F(s, x) = s Q^2(x)\chi^2(s, x)$, we get

$$C_{2} = \left\| \int_{0}^{t} \left[\int_{\mathbb{R}} e^{i\langle k \rangle s} \frac{k}{\langle k \rangle} \overline{\psi(x,k)} F(s,x) dx \right] ds \right\|_{L_{k}^{2}}$$

$$\lesssim \left\| s Q^{2}(\cdot) \chi^{2}(s,\cdot) \right\|_{L_{x}^{1}(\mathbb{R}) L_{s}^{2}([0,\infty])}$$

$$\lesssim \left\| s Q(\cdot) \chi^{2}(s,\cdot) \right\|_{L_{s}^{2}([0,\infty]) L_{x}^{\infty}(\mathbb{R})}$$

$$\lesssim \left\| v \right\|_{X}^{2} \left\| s \langle s \rangle^{-2} \right\|_{L_{s}^{2}([0,\infty])} \lesssim \left\| v \right\|_{X}^{2},$$

$$(4.5)$$

having used the local decay estimate (3.3).

Observe that Lemma 4.2 does not use the genericity of the potential. The proof is similar to the one for the analogous case of the Schrödinger flow e^{itH} given in [7, Lemma 3.5]. We provide the details here for completeness.

Proof of Lemma 4.2. Without loss of generality we may assume $\beta = 0$ and restrict the integral in (4.3) to $k \in [0, \infty)$. Making a change of variable $\langle k \rangle = \lambda$, so that $k \langle k \rangle^{-1} dk = d\lambda$, $dk = \frac{\lambda}{\sqrt{\lambda^2 - 1}} d\lambda$ one has

$$I(s,x) := \int_0^\infty \mathcal{Q}(x,k)\phi(k)e^{i\langle k\rangle s}h(k)\,dk$$
$$= \int_1^\infty \mathcal{Q}(x,\sqrt{\lambda^2-1})\phi(\sqrt{\lambda^2-1})e^{is\lambda}h(\sqrt{\lambda^2-1})\frac{\lambda}{\sqrt{\lambda^2-1}}\,d\lambda.$$

Taking the L_s^2 norm and applying Plancherel's theorem we obtain

$$||I(\cdot,x)||_{L_{s}^{2}([0,t])}^{2} \lesssim \left\| \int_{\mathbb{R}} \mathcal{Q}(x,\sqrt{\lambda^{2}-1}) \mathbf{1}_{[1,\infty)}(\lambda) \phi(\sqrt{\lambda^{2}-1}) e^{is\lambda} h(\sqrt{\lambda^{2}-1}) \frac{\lambda}{\sqrt{\lambda^{2}-1}} d\lambda \right\|_{L_{s}^{2}(\mathbb{R})}^{2}$$

$$\lesssim \int_{1}^{\infty} \left| \mathcal{Q}(x,\sqrt{\lambda^{2}-1}) \frac{\lambda \phi(\sqrt{\lambda^{2}-1})}{\sqrt{\lambda^{2}-1}} h(\sqrt{\lambda^{2}-1}) \right|^{2} d\lambda$$

$$\lesssim \int_{1}^{\infty} \left| h(\sqrt{\lambda^{2}-1}) \right|^{2} d\lambda \lesssim \int_{0}^{\infty} |h(k)|^{2} \frac{k}{\langle k \rangle} dk \lesssim ||h||_{L^{2}}^{2}.$$

To obtain the inhomogeneous estimate (4.4) we test the expression on the left-hand side against an arbitrary function $h \in L^2$:

$$\left\langle h, \int_0^t \left[\int_{\mathbb{R}} e^{-i\langle k \rangle s} \overline{\phi}(k) \overline{\mathcal{Q}}(y, k) F(s, y) \, dy \right] ds \right\rangle_{L_k^2}$$

$$= \int_0^t \int_{\mathbb{R}} \left(\int_{\mathbb{R}} e^{i\langle k \rangle s} \phi(k) \mathcal{Q}(y, k) h(k) \, dk \right) \overline{F}(s, y) \, dy ds$$

Applying Hölder's inequality followed by the homogeneous estimate (4.3):

$$\begin{split} & \left| \int_0^t \int_{\mathbb{R}} \left(\int e^{i\langle k \rangle s} \phi(k) \mathcal{Q}(y,k) h(k) \, dk \right) \overline{F}(s,y) \, dy ds \right| \\ & \lesssim \left\| \int \mathcal{Q}(x,k) \phi(k) e^{i\langle k \rangle s} h(k) \, dk \right\|_{L_x^\infty L_s^2([0,t])} \left\| F \right\|_{L_x^1 L_s^2([0,t])} \\ & \lesssim \|h\|_{L^2} \|F\|_{L_x^1 L_s^2([0,t])} \end{split}$$

which implies (4.4) by duality.

5. Quartic term analysis

In this section we provide a proof of the following lemma:

Lemma 5.1. Given $||v||_X \ll 1$, for every $t \ge 0$ we have

$$\left\| \int_0^t \partial_k \left(e^{is\langle k \rangle} \widetilde{\mathcal{F}} \chi^4(s) \right) ds \right\|_{L^2_k} \lesssim \|v\|_X^4. \tag{5.1}$$

This lemma gives a bound for the quartic term in (3.17) and, together with Lemma 3.2 and 4.1, and Lemma 3.4, will thus complete the proof of Proposition 3.2. This is turn give the main theorem.

The term inside the integral in (5.1) can be written as

$$16e^{is\langle k\rangle}\widetilde{\mathcal{F}}\chi^4(s,k)$$

$$= \sum_{\iota_1,\iota_2,\iota_3,\iota_4 \in \{+,-\}} \iota_1 \iota_2 \iota_3 \iota_4 \int \frac{e^{is\Phi_{\iota_1 \iota_2 \iota_3 \iota_4}}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} \widetilde{g}^{(\iota_1)}(l) \widetilde{g}^{(\iota_2)}(m) \widetilde{g}^{(\iota_3)}(n) \widetilde{g}^{(\iota_4)}(p) \mu_{\iota_1 \iota_2 \iota_3 \iota_4} dlmnp$$

$$(5.2)$$

where we used the short-hand notation dlmnp := dldmdndp, the definition of the distorted Fourier transform and the relation (1.36), with the functions $\Phi_{\iota_1\iota_2\iota_3\iota_4}$ and $\mu_{\iota_1\iota_2\iota_3\iota_4}$, depending on the variables k, l, m, n, p, defined as follows

$$\Phi_{\iota_1\iota_2\iota_3\iota_4}(k, l, m, n, p) = \langle k \rangle - \iota_1 \langle l \rangle - \iota_2 \langle m \rangle - \iota_3 \langle n \rangle - \iota_4 \langle p \rangle, \tag{5.3}$$

and

$$\mu_{\iota_1\iota_2\iota_3\iota_4}(k,l,m,n,p) = \int \overline{\psi(x,k)} \psi^{(\iota_1)}(x,l) \psi^{(\iota_2)}(x,m) \psi^{(\iota_3)}(x,n) \psi^{(\iota_4)}(x,p) dx. \tag{5.4}$$

We call $\mu_{\iota_1\iota_2\iota_3\iota_4}(k,l,m,n,p)$ as the nonlinear spectral distribution. We further decompose this spectral distribution into singular and regular parts according to (2.24), and consider each part separately.

5.1. Decomposition of the nonlinear spectral distribution. Using the decomposition (2.24), we rewrite the nonlinear spectral distribution (5.4) as

$$(2\pi)^{5/2}\mu_{\iota_1\iota_2\iota_3\iota_4}(k,l,m,n,p) = \mu^S_{\iota_1\iota_2\iota_3\iota_4}(k,l,m,n,p) + \mu^R_{\iota_1\iota_2\iota_3\iota_4}(k,l,m,n,p),$$

where μ^S and μ^R denote the 'Singular' and 'Regular' parts, respectively. Below, we give the exact construction of each part. The singular part is the sum of integrals of functions which oscillate at either $-\infty$ or $+\infty$. The regular part consists instead of integrals localized or compactly supported functions.

Recalling (2.25), we define the singular part by

$$\mu_{\iota_{1}\iota_{2}\iota_{3}\iota_{4}}^{S}(k,l,m,n,p) = \mu_{\iota_{1}\iota_{2}\iota_{3}\iota_{4}}^{+}(k,l,m,n,p) + \mu_{\iota_{1}\iota_{2}\iota_{3}\iota_{4}}^{-}(k,l,m,n,p),$$

$$\mu_{\iota_{1}\iota_{2}\iota_{3}\iota_{4}}^{*}(k,l,m,n,p) = \int \nu_{*}(x)\overline{\psi_{S,*}(x,k)}\psi_{S,*}^{(\iota_{1})}(x,l)\psi_{S,*}^{(\iota_{2})}(x,m)\psi_{S,*}^{(\iota_{3})}(x,n)\psi_{S,*}^{(\iota_{4})}(x,p)dx,$$

$$(5.5)$$

where $\nu_*(x) = \sigma_*^5(x)$ for $* \in \{+, -\}$.

For the regular part, we define the set

$$Z_R := \{(A_1, A_2, A_3, A_4, A_5) : \exists j = 1, 2, 3, 4, 5 \text{ s.t. } A_j = R\} \subseteq \{S, R\}^5,$$

and write μ^R as the sum of two parts as follows:

$$\mu_{\iota_1\iota_2\iota_3\iota_4}^R(k,l,m,n,p) = \mu_{\iota_1\iota_2\iota_3\iota_4}^{R,1}(k,l,m,n,p) + \mu_{\iota_1\iota_2\iota_3\iota_4}^{R,2}(k,l,m,n,p)$$

where

$$\mu_{\iota_1\iota_2\iota_3\iota_4}^{R,1} = \sum_{(A,B,C,D,E)\in Z_R} \int \overline{\psi_A(x,k)} \psi_B^{(\iota_1)}(x,l) \psi_C^{(\iota_2)}(x,m) \psi_D^{(\iota_3)}(x,n) \psi_E^{(\iota_4)}(x,p) dx, \tag{5.6}$$

and

$$\mu_{\iota_1\iota_2\iota_3\iota_4}^{R,2} = \int \overline{\psi_S(x,k)} \psi_S^{(\iota_1)}(x,l) \psi_S^{(\iota_2)}(x,m) \psi_S^{(\iota_3)}(x,n) \psi_S^{(\iota_4)}(x,p) dx - \mu_{\iota_1\iota_2\iota_3\iota_4}^S.$$
 (5.7)

Notice that $\mu^{R,1}$ is the sum of integrals of functions localized in x around 0, and $\mu^{R,2}$ consists of integrals of functions compactly supported around 0 due to the presence of product of the form $\sigma_+(x)\sigma_-(x)$.

We decompose accordingly (5.2) and write

$$16e^{is\langle k\rangle}\widetilde{\mathcal{F}}\chi^4(s,k) = \mathcal{N}_+(s,k) + \mathcal{N}_-(s,k) + \mathcal{N}_{R,1}(s,k) + \mathcal{N}_{R,2}(s,k),$$

where

$$(2\pi)^{5/2} \mathcal{N}_{*}(s,k) = \sum_{\iota_{1},\iota_{2},\iota_{3},\iota_{4} \in \{+,-\}} \iota_{1}\iota_{2}\iota_{3}\iota_{4} \int \frac{e^{is\Phi_{\iota_{1}\iota_{2}\iota_{3}\iota_{4}}}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} \widetilde{g}^{(\iota_{1})}(l) \widetilde{g}^{(\iota_{2})}(m) \widetilde{g}^{(\iota_{3})}(n) \widetilde{g}^{(\iota_{4})}(p) \mu_{\iota_{1}\iota_{2}\iota_{3}\iota_{4}}^{*} dlmnp.$$

$$(5.8)$$

Hence, to prove Lemma 5.1 it suffices to show that bound in (5.1) holds for every term in the right-hand side of the following inequality:

$$\left\| \int_{0}^{t} \partial_{k} \left(e^{is\langle k \rangle} \widetilde{\mathcal{F}} \chi^{4}(s) \right) ds \right\|_{L_{k}^{2}} \lesssim \left\| \int_{0}^{t} \partial_{k} \mathcal{N}_{+}(s,k) ds \right\|_{L_{k}^{2}} + \left\| \int_{0}^{t} \partial_{k} \mathcal{N}_{-}(s,k) ds \right\|_{L_{k}^{2}} + \left\| \int_{0}^{t} \partial_{k} \mathcal{N}_{R,2}(s,k) ds \right\|_{L_{k}^{2}}$$

$$+ \left\| \int_{0}^{t} \partial_{k} \mathcal{N}_{R,1}(s,k) ds \right\|_{L_{k}^{2}} + \left\| \int_{0}^{t} \partial_{k} \mathcal{N}_{R,2}(s,k) ds \right\|_{L_{k}^{2}}.$$

$$(5.9)$$

In Subsection 5.2, we prove the bound for the first two terms in (5.9) corresponding to the singular part. In Subsection 5.3, we prove the bound for the rest of the terms corresponding to the regular part.

5.2. Estimates for the singular part. Our goal here is to show that for every $t \geq 0$,

$$\left\| \int_0^t \partial_k \mathcal{N}_{\pm}(s,k) ds \right\|_{L_k^2} \lesssim \|v\|_X^4. \tag{5.10}$$

We will provide the proof for \mathcal{N}_+ only, as \mathcal{N}_- case can be treated similarly. We first expand $\psi_{S,+}$ functions in (5.5) as the linear combination in (2.28), and rewrite $\mu^{S,+}$ as follows

$$\mu_{\iota_{1}\iota_{2}\iota_{3}\iota_{4}}^{S,+}(k,l,m,n,p) = \sqrt{2\pi} \sum_{\epsilon_{0},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4} \in \{+,-\}} \overline{a_{+,\epsilon_{0}}(k)} \ a_{+,\epsilon_{1}}^{(\iota_{1})}(l) \ a_{+,\epsilon_{2}}^{(\iota_{2})}(m) \ a_{+,\epsilon_{3}}^{(\iota_{3})}(n) \ a_{+,\epsilon_{4}}^{(\iota_{4})}(p) \\ \times \widehat{\nu_{+}}(\epsilon_{0}k - \iota_{1}\epsilon_{1}l - \iota_{2}\epsilon_{2}m - \iota_{3}\epsilon_{3}n - \iota_{4}\epsilon_{4}p),$$

$$(5.11)$$

where $\widehat{\nu}_{+}$ represents the (standard) Fourier transform of ν_{+} . Then, substituting the expression (5.11) into (5.8), and introducing the multilinear form

$$\mathcal{T}_{b,\epsilon_{0},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}^{\iota_{1},\iota_{2},\iota_{3},\iota_{4}}(f_{1},f_{2},f_{3},f_{4})(k) := (\iota_{1}\iota_{2}\iota_{3}\iota_{4}) \int \frac{e^{is\Phi_{\iota_{1}\iota_{2}\iota_{3}\iota_{4}}}}{\langle l\rangle\langle m\rangle\langle n\rangle\langle p\rangle} f_{1}^{(\iota_{1})}(l) f_{2}^{(\iota_{2})}(m) f_{3}^{(\iota_{3})}(n) f_{4}^{(\iota_{4})}(p)$$

$$\times b(\epsilon_{0}k - \iota_{1}\epsilon_{1}l - \iota_{2}\epsilon_{2}m - \iota_{3}\epsilon_{3}n - \iota_{4}\epsilon_{4}p) dlmnp$$

$$(5.12)$$

with $\iota_j, \epsilon_j \in \{+, -\}$ and a distribution b, the formula for \mathcal{N}_+ is equivalent to

$$4\pi^{2}\mathcal{N}_{+}(s,k) = \sum_{\substack{\iota_{1},\iota_{2},\iota_{3},\iota_{4} \in \{+,-\}\\ \epsilon_{0},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4} \in \{+,-\}}} \overline{a_{+,\epsilon_{0}}(k)} \, \mathcal{T}_{\widehat{\nu_{+}},\epsilon_{0},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}^{\iota_{1},\iota_{2},\iota_{3},\iota_{4}}(a_{+,\epsilon_{1}}\widetilde{g},a_{+,\epsilon_{2}}\widetilde{g},a_{+,\epsilon_{3}}\widetilde{g},a_{+,\epsilon_{4}}\widetilde{g})(k)$$

$$(5.13)$$

To carry on the analysis of \mathcal{N}_+ we first analyze the properties of the multilinear form \mathcal{T} in (5.12). For simplicity, we consider the particular form given by

$$\mathcal{T}_{b,\epsilon_0,\epsilon_1,\epsilon_2,\epsilon_3,\epsilon_4}(f_1, f_2, f_3, f_4)(k) := \mathcal{T}_{b,\epsilon_0,\epsilon_1,\epsilon_2,\epsilon_3,\epsilon_4}^{+,+,+,+}(f_1, f_2, f_3, f_4)(k)$$
(5.14)

as for the remaining cases, when at least one of $\iota_1, \iota_2, \iota_3, \iota_4$ is '-', the analysis is the same. We will also drop the dependence on ϵ_0 when this is not relevant (for example in Lemmas 5.2 and 5.3) with the understanding that we will be considering $\epsilon_0 = +$.

The following algebraic lemma establishes an alternative representation of $\partial_k \mathcal{T}$, which will be used later to derive the desired estimate (5.10).

Lemma 5.2. For $f_i \in \mathcal{S}$, we have

$$\langle k \rangle \partial_{k} \mathcal{T}_{b,\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1}, f_{2}, f_{3}, f_{4})(k) = is \mathcal{T}_{qb,\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1}, f_{2}, f_{3}, f_{4})(k)$$

$$+ \epsilon_{1} \mathcal{T}_{b,\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(\langle \cdot \rangle \partial_{l} f_{1}, f_{2}, f_{3}, f_{4})(k)$$

$$+ \epsilon_{2} \mathcal{T}_{b,\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1}, \langle \cdot \rangle \partial_{m} f_{2}, f_{3}, f_{4})(k)$$

$$+ \epsilon_{3} \mathcal{T}_{b,\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1}, f_{2}, \langle \cdot \rangle \partial_{n} f_{3}, f_{4})(k)$$

$$+ \epsilon_{4} \mathcal{T}_{b,\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1}, f_{2}, f_{3}, \langle \cdot \rangle \partial_{p} f_{4})(k)$$

$$+ \int \frac{\Phi e^{is\Phi}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} f_{1}(l) f_{2}(m) f_{3}(n) f_{4}(p) \partial_{q} b(q) dlmnp,$$

$$(5.15)$$

where $\Phi := \Phi_{+,+,+,+}$, see (5.3), and $q = k - \epsilon_1 l - \epsilon_2 m - \epsilon_3 n - \epsilon_4 p$.

Proof. Throughout the proof we drop the $\epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4$ indices for simplicity. Then, given $\epsilon_0 = \iota_1 = \iota_2 = \iota_3 = \iota_4 = +$ in (5.12) from the choice of \mathcal{T}_b , we differentiate (5.12) with respect to k and obtain

$$\langle k \rangle \partial_k \mathcal{T}_{b}(f_1, f_2, f_3, f_4)(k) = isk \mathcal{T}_{b}(f_1, f_2, f_3, f_4)(k) + \langle k \rangle \mathcal{T}_{\partial_q b}(f_1, f_2, f_3, f_4)(k). \tag{5.16}$$

Then, the expression (5.16) can be further rewritten as

$$\langle k \rangle \partial_{k} \mathcal{T}_{b}(f_{1}, f_{2}, f_{3}, f_{4})(k) = is \mathcal{T}_{qb}(f_{1}, f_{2}, f_{3}, f_{4})(k) + \langle k \rangle \mathcal{T}_{\partial_{q}b}(f_{1}, f_{2}, f_{3}, f_{4})(k) + \epsilon_{1} is \mathcal{T}_{b}(\cdot f_{1}, f_{2}, f_{3}, f_{4})(k) + \epsilon_{2} is \mathcal{T}_{b}(f_{1}, \cdot f_{2}, f_{3}, f_{4})(k) + \epsilon_{3} is \mathcal{T}_{b}(f_{1}, f_{2}, \cdot f_{3}, f_{4})(k) + \epsilon_{4} is \mathcal{T}_{b}(f_{1}, f_{2}, f_{3}, \cdot f_{4})(k),$$

$$(5.17)$$

where f denotes the multiplication of f by its variable, (f)(x) = xf(x). Using $isl\langle l \rangle^{-1}e^{is\Phi} = -\partial_l(e^{is\Phi})$ and $q = k - \epsilon_1 l - \epsilon_2 m - \epsilon_3 n - \epsilon_4 p$, we integrate the third term in (5.17) by parts in

l to obtain

$$is\mathcal{T}_{b}(\cdot f_{1}, f_{2}, f_{3}, f_{4})(k) = \int \frac{isl e^{is\Phi}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} f_{1}(l) f_{2}(m) f_{3}(n) f_{4}(p) b(q) dlmnp$$

$$= \int \frac{e^{is\Phi}}{\langle m \rangle \langle n \rangle \langle p \rangle} [\partial_{l} f_{1}(l)] f_{2}(m) f_{3}(n) f_{4}(p) b(q) dlmnp$$

$$+ \int \frac{e^{is\Phi}}{\langle m \rangle \langle n \rangle \langle p \rangle} f_{1}(l) f_{2}(m) f_{3}(n) f_{4}(p) \partial_{l} b(q) dlmnp$$

$$= \mathcal{T}_{b}(\langle \cdot \rangle \partial_{l} f_{1}, f_{2}, f_{3}, f_{4})(k) - \epsilon_{1} \mathcal{T}_{\partial_{q} b}(\langle \cdot \rangle f_{1}, f_{2}, f_{3}, f_{4})(k),$$

$$(5.18)$$

where we used the relation $\partial_l b(q) = -\epsilon_1 \partial_q b(q)$. The last three terms in (5.17) are manipulated similarly. Then, we substitute to the last four terms in (5.17) their alternative expressions as in (5.18), and obtain (5.15). This way the terms corresponding to $\mathcal{T}_{\partial_q b}$ are combined into the last line in (5.15).

The following lemma will be used to transfer estimates for the multilinear forms into a physical space.

Lemma 5.3. Given $f_j \in \mathcal{S}$, we have

$$\widehat{\mathcal{F}}^{-1}\left[e^{-is\langle k\rangle}\mathcal{T}_{\mathbf{b},\epsilon_1,\epsilon_2,\epsilon_3,\epsilon_4}(f_1,f_2,f_3,f_4)(k)\right](x) = \widehat{\mathcal{F}}^{-1}[\mathbf{b}](x)\prod_{j=1,2,3,4}u_j(s,\epsilon_jx),\tag{5.19}$$

where $u_j(s,x) = \langle D \rangle^{-1} e^{-is\langle D \rangle} \widecheck{f_j}(x)$ with $\langle D \rangle = \sqrt{-\partial_x^2 + 1}$ and \widecheck{f} the inverse Fourier transform.

The proof relies on explicit computations, and is left to the reader.

A direct application of Lemma 5.3 leads to

$$\|\mathcal{T}_{b,\epsilon_1,\epsilon_2,\epsilon_3,\epsilon_4}(f_1, f_2, f_3, f_4)\|_{L^2_k} \lesssim \|\widehat{\mathcal{F}}^{-1}[b] \prod_{j=1,2,3,4} u_j(s, \epsilon_j \cdot)\|_{L^2_x}$$
 (5.20)

Remark 5.1. Throughout the computations for the singular part, the distribution b is replaced by $\widehat{\nu_+}$, which is given by the formula

$$\widehat{\nu_{+}}(k) = \sqrt{\frac{\pi}{2}} \delta_0 + \frac{\widehat{\zeta}(k)}{ik} + \widehat{\overline{\omega}}(k), \qquad (5.21)$$

where ζ is an even C_c^{∞} function with integral 1 and $\overline{\omega}$ denotes a (generic) C_c^{∞} function. See for example [20]

From the formula (5.13) we see that \mathcal{N}_+ is a sum of several terms. The analysis for all these terms is similar and, therefore, in what follows we consider only the terms with fixed $\iota_1, \iota_2, \iota_3, \iota_4 = +$ and drop these from our notation. Then, to prove (5.10), it suffices to show that, for every $\epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4 \in \{+, -\}$ we have

$$\left\| \int_{0}^{t} \partial_{k} \sum_{\epsilon_{0} \in \{+,-\}} \overline{a_{+,\epsilon_{0}}}(k) \, \mathcal{T}_{\widehat{\nu_{+}},\epsilon_{0},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1}, f_{2}, f_{3}, f_{4})(k) \, ds \right\|_{L_{k}^{2}} \lesssim \|v\|_{X}^{4}, \tag{5.22}$$

where we set

$$f_j(s,\cdot) = a_{+,\epsilon_j}(\cdot)\widetilde{g}(\cdot)$$
 for $j = 1, 2, 3, 4$.

Notice that the coefficients $a_{+,\epsilon_0}(k)$ have jump discontinuities at k=0, see (2.29), which may, in principle, result in a singular δ -type contribution when differentiated. However, using

the formulas (2.29) and the vanishing at k = 0 of T(k) and $R_{\pm}(k) + 1$, see (2.9), one can see that this is not the case for the summation in (5.22). More precisely, we first write

$$\begin{split} &\sum_{\epsilon_0 \in \{+,-\}} \overline{a_{+,\epsilon_0}}(k) \, \mathcal{T}_{\widehat{\nu_+},\epsilon_0,\epsilon_1,\epsilon_2,\epsilon_3,\epsilon_4}(f_1,f_2,f_3,f_4) \\ &= \left[\overline{a_{+,+}}(k) + \overline{a_{+,-}}(k) \right] \mathcal{T}_{\widehat{\nu_+},+,\epsilon_1,\epsilon_2,\epsilon_3,\epsilon_4}(f_1,f_2,f_3,f_4) \\ &+ \overline{a_{+,-}}(k) \left[\mathcal{T}_{\widehat{\nu_+},-,\epsilon_1,\epsilon_2,\epsilon_3,\epsilon_4}(f_1,f_2,f_3,f_4) - \mathcal{T}_{\widehat{\nu_+},+,\epsilon_1,\epsilon_2,\epsilon_3,\epsilon_4}(f_1,f_2,f_3,f_4) \right], \end{split}$$

and observe that the difference in the last line vanishes at k = 0, see (5.14) and (5.12). Then, we use that, for $\iota_0 \in \{+, -\}$, $\partial_k(a_{\iota_0, +}(k) + a_{\iota_0, -}(k)) = \mathbf{1}_{\iota_0}(k)\partial_k T(\iota_0 k) + \mathbf{1}_{-\iota_0}(k)\partial_k R_{\iota_0}(-\iota_0 k)$. Therefore, using also (2.10) and Minkowski's inequality we can bound

$$\left\| \int_{0}^{t} \partial_{k} \sum_{\epsilon_{0} \in \{+,-\}} \overline{a_{+,\epsilon_{0}}} \, \mathcal{T}_{\widehat{\nu_{+}},\epsilon_{0},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1},f_{2},f_{3},f_{4}) ds \right\|_{L_{k}^{2}} \lesssim \int_{0}^{t} \left\| \mathcal{T}_{\widehat{\nu_{+}},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1},f_{2},f_{3},f_{4}) \right\|_{L_{k}^{2}} ds + \left\| \int_{0}^{t} \left| \partial_{k} \mathcal{T}_{\widehat{\nu_{+}},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1},f_{2},f_{3},f_{4}) ds \right\|_{L_{k}^{2}} := E_{1}(t) + E_{2}(t).$$

Below we give estimates for $E_1(t)$ and $E_2(t)$, and show that each of them is bounded by $||v||_X^4$, which would imply the bound (5.10).

5.2.1. Estimates for E_1 . We show that $E_1(t) \lesssim ||v||_X^4$ for every $t \geq 0$. Using (5.20),

$$E_{1}(t) \lesssim \int_{0}^{t} \|\nu_{+}u_{1}(s,\epsilon_{1}\cdot)u_{2}(s,\epsilon_{2}\cdot)u_{3}(s,\epsilon_{3}\cdot)u_{4}(s,\epsilon_{4}\cdot)\|_{L_{x}^{2}} ds$$

$$\lesssim \int_{0}^{t} \|u_{1}(s,\cdot)\|_{L_{x}^{2}} \|u_{2}(s,\cdot)\|_{L_{x}^{\infty}} \|u_{3}(s,\cdot)\|_{L_{x}^{\infty}} \|u_{4}(s,\cdot)\|_{L_{x}^{\infty}} ds,$$
(5.23)

where $u_j(s,x) = \langle D \rangle^{-1} e^{-is\langle D \rangle} \widehat{\mathcal{F}}^{-1}(a_{+,\epsilon_j}\widetilde{g})$. The formulas (2.29) and the bounds in (2.10) imply that for every $s \geq 0$ we have

$$||u_j(s)||_{L^2_x} \lesssim ||\widetilde{g}(s)||_{L^2_k} \lesssim ||v||_X.$$
 (5.24)

Using (2.37) we can bound

$$||u_j(s)||_{L_x^{\infty}} \lesssim \langle s \rangle^{-1/2} \left(||\partial_k \left(a_{+,\epsilon_j} \widetilde{g}(s) \right)||_{L_h^2} + ||\langle k \rangle^2 \widetilde{g}(s)||_{L_k^2} \right); \tag{5.25}$$

using that $\tilde{g}(s,0) = 0$, we see that no singularity arises when differentiating a_{+,ϵ_j} and, therefore,

$$\left\| \partial_k \left(a_{+,\epsilon_j} \widetilde{g}(s) \right) \right\|_{L^2_k} \lesssim \left\| \partial_k \widetilde{g}(s) \right\|_{L^2_k}; \tag{5.26}$$

then, from the definition (3.1) and (5.25)-(5.26) we have

$$||u_j(s)||_{L^{\infty}_{\infty}} \lesssim \langle s \rangle^{-1/2} ||v||_X. \tag{5.27}$$

Using (5.24) and (5.27) in (5.23) we obtain

$$E_1(t) \lesssim ||v||_X^4 \int_0^t \langle s \rangle^{-3/2} ds \lesssim ||v||_X^4.$$

5.2.2. Estimates for E_2 . We want to show that for every $t \geq 0$,

$$\left\| \int_{0}^{t} \partial_{k} \mathcal{T}_{\widehat{\nu_{+}}, \epsilon_{1}, \epsilon_{2}, \epsilon_{3}, \epsilon_{4}}(f_{1}, f_{2}, f_{3}, f_{4}) ds \right\|_{L_{k}^{2}} \lesssim \|v\|_{X}^{4}. \tag{5.28}$$

From Lemma 5.2, the left-hand side of the above expression can be written as

$$\begin{split} \left\| \int_{0}^{t} \partial_{k} \mathcal{T}_{\widehat{\nu_{+}},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1},f_{2},f_{3},f_{4})ds \right\|_{L_{k}^{2}} \\ &= \left\| \langle k \rangle^{-1} \int_{0}^{t} s \, \mathcal{T}_{q\widehat{\nu_{+}},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1},f_{2},f_{3},f_{4})ds \right\|_{L_{k}^{2}} \\ &+ \left\| \langle k \rangle^{-1} \int_{0}^{t} \mathcal{T}_{\widehat{\nu_{+}},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(\langle \cdot \rangle \partial_{l}f_{1},f_{2},f_{3},f_{4})ds \right\|_{L_{k}^{2}} \\ &+ \left\| \langle k \rangle^{-1} \int_{0}^{t} \mathcal{T}_{\widehat{\nu_{+}},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1},\langle \cdot \rangle \partial_{m}f_{2},f_{3},f_{4})ds \right\|_{L_{k}^{2}} \\ &+ \left\| \langle k \rangle^{-1} \int_{0}^{t} \mathcal{T}_{\widehat{\nu_{+}},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1},f_{2},\langle \cdot \rangle \partial_{n}f_{3},f_{4})ds \right\|_{L_{k}^{2}} \\ &+ \left\| \langle k \rangle^{-1} \int_{0}^{t} \mathcal{T}_{\widehat{\nu_{+}},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1},f_{2},f_{3},\langle \cdot \rangle \partial_{p}f_{4})ds \right\|_{L_{k}^{2}} \\ &+ \left\| \langle k \rangle^{-1} \int_{0}^{t} \int \frac{\Phi \, e^{is\Phi}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} f_{1}(l)f_{2}(m)f_{3}(n)f_{4}(p)\partial_{q}\widehat{\nu_{+}}(q)dlmnpds \right\|_{L_{k}^{2}}, \\ &=: E_{2}^{(1)}(t) + E_{2}^{(2)}(t) + E_{2}^{(3)}(t) + E_{2}^{(4)}(t) + E_{2}^{(5)}(t) + E_{2}^{(6)}(t), \end{split}$$

where we recall that $q = k - \epsilon_1 l - \epsilon_2 m - \epsilon_3 n - \epsilon_4 p$. Therefore, to prove (5.28) it suffices to show that every $E_2^{(j)}$ in (5.29) is bounded by $||v||_X^4$. This will be established in the lemmas below.

Lemma 5.4. For every $t \ge 0$, we have

$$E_2^{(1)}(t) \lesssim ||v||_X^4.$$
 (5.30)

Proof. First, according to the Fourier representation of $\widehat{\nu}_{+}$ in (5.21), we have

$$q\widehat{\nu_+} = -i\widehat{\zeta}(q) + q\widehat{\overline{\omega}}(q) = \widehat{\Psi}(q)$$

with Ψ being a Schwartz function. In the estimates below we extract several $\langle x \rangle^{-2}$ factors from Ψ , and use them to apply the local L_x^{∞} decay estimates (2.38). Then, using Minkowski inequality and (5.20), we have

$$E_{2}^{(1)} \lesssim \int_{0}^{t} s \| \mathcal{T}_{q\widehat{\nu_{+}},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}(f_{1},f_{2},f_{3},f_{4}) \|_{L_{k}^{2}} ds$$

$$\lesssim \int_{0}^{t} s \| \Psi u_{1}(s,\epsilon_{1}\cdot)u_{2}(s,\epsilon_{2}\cdot)u_{3}(s,\epsilon_{3}\cdot)u_{4}(s,\epsilon_{4}\cdot) \|_{L_{x}^{2}} ds$$

$$\lesssim \int_{0}^{t} s \| \langle x \rangle^{-2} u_{1}(s) \|_{L_{x}^{\infty}} \| \langle x \rangle^{-2} u_{2}(s) \|_{L_{x}^{\infty}} \| \langle x \rangle^{-2} u_{3}(s) \|_{L_{x}^{\infty}} \| \langle x \rangle^{-2} u_{4}(s) \|_{L_{x}^{\infty}} ds,$$

where $u_j(s,x) = \langle D \rangle^{-1} e^{-is\langle D \rangle} \widehat{\mathcal{F}}^{-1}(a_{+,\epsilon_j} \widetilde{g})$ for every j = 1, 2, 3, 4. Applying the local decay estimates (2.38), using (2.29) and the bounds in (2.10) together with $\widetilde{g}(0) = 0$, we get

$$E_2^{(1)} \lesssim \|v\|_X^4 \int_0^t s \langle s \rangle^{-3} ds \lesssim \|v\|_X^4.$$

Lemma 5.5. For every $t \ge 0$ and j = 2, 3, 4, 5, we have

$$E_2^{(j)}(t) \lesssim ||v||_X^4.$$
 (5.31)

Proof. The proofs for j = 2, 3, 4, 5 are identical so we only prove (5.31) for j = 2. Using (5.20) we can estimate as follows:

$$E_2^{(2)}(t) \lesssim \int_0^t \|u_1(s,\cdot)\|_{L_x^2} \|u_2(s,\cdot)\|_{L_x^\infty} \|u_3(s,\cdot)\|_{L_x^\infty} \|u_4(s,\cdot)\|_{L_x^\infty} ds,$$

where

$$\begin{cases} u_1(s,x) = \langle D \rangle^{-1} e^{-is\langle D \rangle} \widehat{\mathcal{F}}^{-1}(\langle \cdot \rangle \partial_l[a_{+,\epsilon_1} \widetilde{g}]), \\ u_r(s,x) = \langle D \rangle^{-1} e^{-is\langle D \rangle} \widehat{\mathcal{F}}^{-1}(a_{+,\epsilon_r} \widetilde{g}) & \text{for } r = 2, 3, 4. \end{cases}$$

For u_1 we have $||u_1(s)||_{L^2_x} \lesssim ||\widetilde{g}||_{H^1_l} \lesssim ||v||_X$, while the L^{∞}_x norms of u_r for r=2,3,4 are bounded similarly to (5.27). Combining these estimates, we get the desired result

$$E_2^{(2)} \lesssim ||v||_X^4 \int_0^t \langle s \rangle^{-3/2} ds \lesssim ||v||_X^4.$$

The next lemma will complete the proof of (5.28).

Lemma 5.6. For every $t \geq 0$, we have

$$E_2^{(6)}(t) \lesssim ||v||_X^4.$$
 (5.32)

Proof. We first consider the integral

$$\mathcal{I} := \int_0^t \int \frac{\Phi \ e^{is\Phi}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} f_1(l) f_2(m) f_3(n) f_4(p) \partial_q \widehat{\nu_+}(q) dlmnp \ ds,$$

where $q = k - \epsilon_1 l - \epsilon_2 m - \epsilon_3 n - \epsilon_4 p$. Using that $\Phi e^{is\Phi} = -i\partial_s e^{is\Phi}$, we integrate \mathcal{I} by parts in s to get

$$\mathcal{I} = -i \int \frac{e^{is\Phi}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} f_1(l) f_2(m) f_3(n) f_4(p) \partial_q \widehat{\nu_+}(q) dl m n p \bigg|_{s=0}^{s=t}
+ i \int_0^t \int \frac{e^{is\Phi}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} \partial_s [f_1(l) f_2(m) f_3(n) f_4(p)] \partial_q \widehat{\nu_+}(q) dl m n p ds
=: -i \mathcal{I}_1(t, k) + i \mathcal{I}_2(t, k).$$
(5.33)

Both integrals contain the term $\partial_q \widehat{\nu}_+(q)$, which we treat using the relations

$$\partial_q \widehat{\nu_+}(q) = -\epsilon_1 \partial_l \widehat{\nu_+}(q) = -\epsilon_2 \partial_m \widehat{\nu_+}(q) = -\epsilon_3 \partial_n \widehat{\nu_+}(q) = -\epsilon_4 \partial_p \widehat{\nu_+}(q)$$
 (5.34)

to integrate by parts and deal with $\widehat{\nu}_{+}(q)$ only.

We look at \mathcal{I}_1 first. From (5.34), we integrate by parts in l, and obtain

$$\mathcal{I}_{1} = \epsilon_{1} \int \frac{e^{is\Phi}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} \partial_{l} f_{1}(l) f_{2}(m) f_{3}(n) f_{4}(p) \widehat{\nu_{+}}(q) dlmnp \Big|_{s=0}^{s=t} \\
- \epsilon_{1} \int \frac{le^{is\Phi}}{\langle l \rangle^{3} \langle m \rangle \langle n \rangle \langle p \rangle} f_{1}(l) f_{2}(m) f_{3}(n) f_{4}(p) \widehat{\nu_{+}}(q) dlmnp \Big|_{s=0}^{s=t} \\
- i\epsilon_{1} \int \frac{sle^{is\Phi}}{\langle l \rangle^{2} \langle m \rangle \langle n \rangle \langle p \rangle} f_{1}(l) f_{2}(m) f_{3}(n) f_{4}(p) \widehat{\nu_{+}}(q) dlmnp \Big|_{s=0}^{s=t} \\
=: \mathcal{I}_{1}^{(1)}(t,k) + \mathcal{I}_{1}^{(2)}(t,k) + \mathcal{I}_{1}^{(3)}(t,k). \tag{5.35}$$

The integrals above are similar in structure, and, in fact, can be treated in the same way. We only show how to deal with $\mathcal{I}_1^{(3)}$ as it contain the potentially problematic factor s. Also, we drop the endpoint values of s and, abusing notations, only work with

$$\mathcal{I}_{1}^{(3)}(s,k) := -i\epsilon_{1} \int \frac{sle^{is\Phi}}{\langle l \rangle^{2} \langle m \rangle \langle n \rangle \langle p \rangle} f_{1}(l) f_{2}(m) f_{3}(n) f_{4}(p) \widehat{\nu_{+}}(q) dlmnp.$$

From (5.14), we have

$$\mathcal{I}_{1}^{(3)}(s,k) = -i\epsilon_{1}s\mathcal{T}_{\widehat{\nu_{+}},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}}\left(\frac{\cdot}{\langle \cdot \rangle}f_{1},f_{2},f_{3},f_{4}\right).$$

Then, using (5.20),

$$\begin{aligned} \left\| \mathcal{I}_{1}^{(3)}(s) \right\|_{L_{x}^{2}} &\lesssim s \left\| \nu_{+} u_{1}(s, \epsilon_{1} \cdot) u_{2}(s, \epsilon_{2} \cdot) u_{3}(s, \epsilon_{3} \cdot) u_{4}(s, \epsilon_{4} \cdot) \right\|_{L_{x}^{2}} \\ &\lesssim s \left\| u_{1}(s, \cdot) \right\|_{L_{x}^{2}} \left\| u_{2}(s, \cdot) \right\|_{L_{x}^{\infty}} \left\| u_{3}(s, \cdot) \right\|_{L_{x}^{\infty}} \left\| u_{4}(s, \cdot) \right\|_{L_{x}^{\infty}} \end{aligned}$$

where

$$\begin{cases} u_1(s,x) = \langle D \rangle^{-1} e^{-is\langle D \rangle} \widehat{\mathcal{F}}^{-1} \left(\frac{\cdot}{\langle \cdot \rangle} a_{+,\epsilon_1} \widetilde{g} \right), \\ u_r(s,x) = \langle D \rangle^{-1} e^{-is\langle D \rangle} \widehat{\mathcal{F}}^{-1} (a_{+,\epsilon_r} \widetilde{g}) & \text{for } r = 2, 3, 4. \end{cases}$$

Then, the dispersive decay estimates (2.37) imply

$$\left\| \mathcal{I}_{1}^{(3)}(s) \right\|_{L_{k}^{2}} \lesssim s \langle s \rangle^{-3/2} \|v\|_{X}^{4} \lesssim \|v\|_{X}^{4}.$$

Similar steps repeated for $\mathcal{I}_1^{(1)}(t,k)$ and $\mathcal{I}_1^{(2)}(t,k)$ from (5.35) give the same bound $\|v\|_X^4$ for the integrals. Combining such estimates we have $\|\mathcal{I}_1(t)\|_{L^2_k} \lesssim \|v\|_X^4$ for every $t \geq 0$, as desired.

It remains to verify $\|\mathcal{I}_2(t)\|_{L^2_k} \lesssim \|v\|_X^4$ for \mathcal{I}_2 given in (5.33). As in previous computations, based on (5.34) we use $\partial_q \widehat{\nu_+}$ to integrate by parts in one of the integration variables l, m, n or p. The choice of the variable depends on what function ∂_s hits; more precisely, when integrating by parts, we want avoid hitting the function of the type $\partial_s f_j$. Applying ∂_s to the product of functions in the \mathcal{I}_2 expression in (5.33), we obtain four terms which can all be treated in the same way, so we only look at one of them:

$$\mathcal{I}_{2}^{(1)}(t,k) := \int_{0}^{t} \int \frac{e^{is\Phi}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} [\partial_{s} f_{1}(l)] f_{2}(m) f_{3}(n) f_{4}(p) \partial_{q} \widehat{\nu}_{+}(q) dlmnp \ ds. \tag{5.36}$$

We use (5.34) to integrate by parts in m, and using (5.14) we write

$$\mathcal{I}_{2}^{(1)}(t,k) = \epsilon_{2} \int_{0}^{t} \mathcal{T}_{\widehat{\nu_{+}},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}} \left(\partial_{s} f_{1}, \partial_{m} f_{2}, f_{3}, f_{4}\right) ds
- \epsilon_{2} \int_{0}^{t} \mathcal{T}_{\widehat{\nu_{+}},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}} \left(\partial_{s} f_{1}, \frac{\cdot}{\langle \cdot \rangle^{2}} f_{2}, f_{3}, f_{4}\right) ds
- i\epsilon_{2} \int_{0}^{t} s \mathcal{T}_{\widehat{\nu_{+}},\epsilon_{1},\epsilon_{2},\epsilon_{3},\epsilon_{4}} \left(\partial_{s} f_{1}, \frac{\cdot}{\langle \cdot \rangle} f_{2}, f_{3}, f_{4}\right) ds =: J_{1}(t,k) + J_{2}(t,k) + J_{3}(t,k).$$

Note that all three integrals contain the term $\partial_s f_1$, and we eventually use it to get $\langle s \rangle^{-3/2}$ decay via (3.7) which allows to overcome the factor s in the integral J_3 without any difficulties. For the first integral we use (5.20),

$$||J_{1}(t)||_{L_{k}^{2}} \lesssim \int_{0}^{t} ||\nu_{+}u_{1}(s,\epsilon_{1}\cdot)u_{2}(s,\epsilon_{2}\cdot)u_{3}(s,\epsilon_{3}\cdot)u_{4}(s,\epsilon_{4}\cdot)||_{L_{x}^{2}} ds$$

$$\lesssim \int_{0}^{t} ||u_{1}(s,\cdot)||_{L_{x}^{\infty}} ||u_{2}(s,\cdot)||_{L_{x}^{2}} ||u_{3}(s,\cdot)||_{L_{x}^{\infty}} ||u_{4}(s,\cdot)||_{L_{x}^{\infty}} ds$$
(5.37)

where

$$\begin{cases} u_1(s,x) = \langle D \rangle^{-1} e^{-is\langle D \rangle} \widehat{\mathcal{F}}^{-1}(a_{+,\epsilon_1} \partial_s \widetilde{g}), \\ u_2(s,x) = \langle D \rangle^{-1} e^{-is\langle D \rangle} \widehat{\mathcal{F}}^{-1}(\partial_m [a_{+,\epsilon_2} \widetilde{g}]), \\ u_r(s,x) = \langle D \rangle^{-1} e^{-is\langle D \rangle} \widehat{\mathcal{F}}^{-1}(a_{+,\epsilon_r} \widetilde{g}) \quad \text{for } r = 3, 4. \end{cases}$$

To bound the L_x^{∞} norm of u_1 we use the simple relation $||u_1(s)||_{L_x^{\infty}} \lesssim ||\langle l \rangle \widehat{u}_1||_{L_t^2}$, which implies

$$||u_1(s)||_{L^{\infty}_{x}} \lesssim ||\partial_s \widetilde{g}||_{L^2_t} \lesssim \langle s \rangle^{-3/2} ||v||_X^2$$
 (5.38)

according to the bound in (3.7). From the dispersive decay estimates for the case of the standard Fourier transform (2.37), (2.10), and (5.26), we have

$$||u_2(s)||_{L_x^2} \lesssim ||v||_X \quad \text{and} \quad ||u_r(s)||_{L_x^\infty} \lesssim \langle s \rangle^{-1/2} ||v||_X.$$
 (5.39)

Using (5.38) and (5.39) into (5.37), we obtain

$$||J_1(t)||_{L^2_k} \lesssim ||v||_X^5 \int_0^t \langle s \rangle^{-5/2} ds \lesssim ||v||_X^4.$$
 (5.40)

The remaining integrals J_2 and J_3 can be treated in the same way with appropriate modifications for $u_2(s,x)$. In both cases, the L_x^2 norm of $u_2(s)$ is bounded by $||v||_X$, and this lead to a $||v||_X^4$ bound for both $||J_2(t)||_{L_k^2}$ and $||J_3(t)||_{L_k^2}$, as desired. Combining these estimates, we obtain $||\mathcal{I}_2(t)||_{L_k^2} \lesssim ||v||_X^4$. Together with the previously obtained result for \mathcal{I}_1 , we get the desired estimate (5.32).

5.3. Estimates for the regular part. Our goal here is to show that for every $t \geq 0$,

$$\left\| \int_0^t \partial_k \mathcal{N}_{R,j}(s,k) ds \right\|_{L^2_k} \lesssim \|v\|_X^4 \quad \text{for } j = 1, 2.$$
 (5.41)

We separate the analysis for j = 1 and j = 2. Recall the regular parts of the nonlinear spectral distribution given by (5.6)–(5.7) and the expression for $\mathcal{N}_{R,j}$ from (5.8).

5.3.1. Estimates for $\mathcal{N}_{R,1}$. Based on (5.8), the expression for $\mathcal{N}_{R,1}$ reads

$$(2\pi)^{5/2}\mathcal{N}_{R,1}(s,k)$$

$$= \sum_{\iota_1,\iota_2,\iota_3,\iota_4 \in \{+,-\}} \iota_1 \iota_2 \iota_3 \iota_4 \int \frac{e^{is\Phi_{\iota_1 \iota_2 \iota_3 \iota_4}}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} \widetilde{g}^{(\iota_1)}(l) \widetilde{g}^{(\iota_2)}(m) \widetilde{g}^{(\iota_3)}(n) \widetilde{g}^{(\iota_4)}(p) \mu_{\iota_1 \iota_2 \iota_3 \iota_4}^{R,1} dlmnp. \tag{5.42}$$

The analysis for each term in the above sum is similar, therefore we may only consider the term with $\iota_1 = \iota_2 = \iota_3 = \iota_4 = +$. Next, we recall that $\mu_{++++}^{R,1}$ defined in (5.6) is the sum of multiple integrals. Every integral in the sum (5.6) has at least one of the indices in (A, B, C, D, E) equal to R; they are similar in nature, and are treated essentially in the same way. In particular, it is sufficient to consider only two terms:

- the term with A = R;
- the term with A = S, B = R.

For these terms, the analysis is independent on the rest of the indices. so we do not specify them explicitly as R and S. In other words, we consider

$$\mu^{R,1,1} := \int \overline{\psi_R(x,k)} \psi_{M_1}(x,l) \psi_{M_2}(x,m) \psi_{M_3}(x,n) \psi_{M_4}(x,p) dx, \quad M_j \in \{S,R\},$$
 (5.43)

and

$$\mu^{R,1,2} := \int \overline{\psi_S(x,k)} \psi_{M_1}(x,l) \psi_{M_2}(x,m) \psi_{M_3}(x,n) \psi_R(x,p) dx, \quad M_j \in \{S,R\}.$$
 (5.44)

The corresponding integrals from (5.42) are

$$\mathcal{N}_{R,1,1}(s,k) = \int \frac{e^{is\Phi}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} \widetilde{g}(l) \widetilde{g}(m) \widetilde{g}(n) \widetilde{g}(p) \mu^{R,1,1} dlmnp,$$

and

$$\mathcal{N}_{R,1,2}(s,k) = \int \frac{e^{is\Phi}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} \widetilde{g}(l) \widetilde{g}(m) \widetilde{g}(n) \widetilde{g}(p) \mu^{R,1,2} dlmnp,$$

where $\Phi = \Phi_{++++}$ given by (5.3). After applying the ∂_k differentiation to the above integral, we get for j = 1, 2

$$\partial_{k} \mathcal{N}_{R,1,j}(s,k) = is \frac{k}{\langle k \rangle} \int \frac{e^{is\Phi}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} \widetilde{g}(l) \widetilde{g}(m) \widetilde{g}(n) \widetilde{g}(p) \mu^{R,1,j} dlmnp$$

$$+ \int \frac{e^{is\Phi}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} \widetilde{g}(l) \widetilde{g}(m) \widetilde{g}(n) \widetilde{g}(p) \partial_{k} \mu^{R,1,j} dlmnp$$

$$=: \mathcal{A}_{j}(s,k) + \mathcal{B}_{j}(s,k).$$
(5.45)

To show the bound in (5.41) it suffices to verify the bound for the L_k^2 norms of ds integral associated with A_i and B_i from (5.45). We show these estimates in the following lemmas.

Lemma 5.7. For every $t \ge 0$ we have

$$\left\| \int_0^t \mathcal{A}_1(s,k) ds \right\|_{L^2_h} \lesssim \|v\|_X^4.$$

Proof. We substitute $\mu^{R,1,1}$ from (5.43) into the expression for \mathcal{A}_1 , and expand the function $\Phi = \Phi_{++++}$ according to its definition in (5.3). As a result, using (2.30), we can rewrite \mathcal{A}_1 as

$$\mathcal{A}_1(s,k) = 4is\pi^2 \frac{k}{\langle k \rangle} e^{is\langle k \rangle} \int \overline{\psi_R(x,k)} u_{M_1}(x) u_{M_2}(x) u_{M_3}(x) u_{M_4}(x) dx,$$

where $u(x) = \langle \widetilde{D} \rangle^{-1} e^{-is\langle \widetilde{D} \rangle} g(s,x)$. Then, using the estimate (2.31), we have

$$\left\| \int_{0}^{t} \mathcal{A}_{1}(s,k) ds \right\|_{L_{k}^{2}} \lesssim \int_{0}^{t} \left\| \mathcal{A}_{1}(s,k) \right\|_{L_{k}^{2}} ds \lesssim \int_{0}^{t} s \left\| \langle x \rangle^{-11} u_{M_{1}} u_{M_{2}} u_{M_{3}} u_{M_{4}} \right\|_{L_{x}^{2}} ds$$
$$\lesssim \int_{0}^{t} s \left\| \langle x \rangle^{-2} u_{M_{1}} \right\|_{L_{x}^{\infty}} \left\| \langle x \rangle^{-2} u_{M_{2}} \right\|_{L_{x}^{\infty}} \left\| \langle x \rangle^{-2} u_{M_{3}} \right\|_{L_{x}^{\infty}} \left\| \langle x \rangle^{-2} u_{M_{4}} \right\|_{L_{x}^{\infty}} ds.$$

From (2.40) and (2.41), each of the above L_x^{∞} norms is bounded by $\langle s \rangle^{-1} \|\widetilde{g}\|_{H_k^1}$, and this completes the proof.

Lemma 5.8. For every $t \ge 0$ we have

$$\left\| \int_0^t \mathcal{B}_1(s,k) ds \right\|_{L^2_k} \lesssim \|v\|_X^4.$$

Proof. The proof is similar to the proof of Lemma 5.7. We rewrite \mathcal{B}_1 as

$$\mathcal{B}_1(s,k) = 4\pi^2 e^{is\langle k \rangle} \int \partial_k \overline{\psi_R(x,k)} u_{M_1}(x) u_{M_2}(x) u_{M_3}(x) u_{M_4}(x) dx,$$

where $u(x) = \langle \widetilde{D} \rangle^{-1} e^{-is\langle \widetilde{D} \rangle} g(s,x)$. Then the desired estimate follows from (2.32), (2.40) and (2.41).

Lemma 5.9. For every $t \ge 0$ we have

$$\left\| \int_0^t \mathcal{A}_2(s,k) ds \right\|_{L^2_k} \lesssim \|v\|_X^4.$$

Proof. Using (2.30), we rewrite A_2 as

$$\mathcal{A}_2(s,k) = 4is\pi^2 \frac{k}{\langle k \rangle} e^{is\langle k \rangle} \int \overline{\psi_S(x,k)} u_{M_1}(x) u_{M_2}(x) u_{M_3}(x) u_R(x) dx,$$

where $u(x) = \langle \widetilde{D} \rangle^{-1} e^{-is\langle \widetilde{D} \rangle} g(s,x)$. From the definition of ψ_S in (2.25), one can verify that

$$\left\| \int \overline{\psi_S(x,k)} h(x) \ dx \right\|_{L^2_k} \lesssim \|h\|_{L^2_x}. \tag{5.46}$$

Then, applying (5.46) and (2.41) we get

$$\begin{split} \left\| \int_0^t \mathcal{A}_2(s,k) ds \right\|_{L^2_k} &\lesssim \int_0^t s \left\| u_{M_1} u_{M_2} u_{M_3} u_R \right\|_{L^2_x} ds \\ &\lesssim \int_0^t s \left\| \langle x \rangle^{-2} u_{M_1} \right\|_{L^\infty_x} \left\| \langle x \rangle^{-2} u_{M_2} \right\|_{L^\infty_x} \left\| \langle x \rangle^{-2} u_{M_3} \right\|_{L^\infty_x} \left\| \langle x \rangle^{10} u_R \right\|_{L^\infty_x} ds \lesssim \|v\|_X^4 \\ \text{as desired.} \end{split}$$

Lemma 5.10. For every $t \ge 0$ we have

$$\left\| \int_0^t \mathcal{B}_2(s,k) ds \right\|_{L^2_L} \lesssim \|v\|_X^4.$$

Proof. The proof is similar to the proof of Lemma 5.9. We rewrite \mathcal{B}_2 as

$$\mathcal{B}_2(s,k) = 4\pi^2 e^{is\langle k \rangle} \int \partial_k \overline{\psi_S(x,k)} u_{M_1}(x) u_{M_2}(x) u_{M_3}(x) u_R(x) dx,$$

where $u(x) = \langle \widetilde{D} \rangle^{-1} e^{-is\langle \widetilde{D} \rangle} g(s, x)$. Similarly to (5.46), one can verify

$$\left\| \int \partial_k \overline{\psi_S(x,k)} h(x) \ dx \right\|_{L^2_k} \lesssim \|\langle x \rangle h\|_{L^2_x}. \tag{5.47}$$

Then, we obtain

$$\left\| \int_{0}^{t} \mathcal{B}_{2}(s,k) ds \right\|_{L_{k}^{2}} \lesssim \int_{0}^{t} s \left\| \langle x \rangle u_{M_{1}} u_{M_{2}} u_{M_{3}} u_{R} \right\|_{L_{x}^{2}} ds$$

$$\lesssim \int_{0}^{t} s \left\| \langle x \rangle^{-2} u_{M_{1}} \right\|_{L_{x}^{\infty}} \left\| \langle x \rangle^{-2} u_{M_{2}} \right\|_{L_{x}^{\infty}} \left\| \langle x \rangle^{-2} u_{M_{3}} \right\|_{L_{x}^{\infty}} \left\| \langle x \rangle^{10} u_{R} \right\|_{L_{x}^{\infty}} ds \lesssim \|v\|_{X}^{4}.$$

This completes the proof of (5.41) for j = 1.

5.3.2. Estimates for $\mathcal{N}_{R,2}$. Based on (5.8), the expression for $\mathcal{N}_{R,2}$ reads

$$(2\pi)^{5/2}\mathcal{N}_{R,2}(s,k)$$

$$= \sum_{\iota_1,\iota_2,\iota_3,\iota_4 \in \{+,-\}} \iota_1 \iota_2 \iota_3 \iota_4 \int \frac{e^{is\Phi_{\iota_1 \iota_2 \iota_3 \iota_4}}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} \widetilde{g}^{(\iota_1)}(l) \widetilde{g}^{(\iota_2)}(m) \widetilde{g}^{(\iota_3)}(n) \widetilde{g}^{(\iota_4)}(p) \mu_{\iota_1 \iota_2 \iota_3 \iota_4}^{R,2} dlmnp.$$
(5.48)

As previously, we only consider the contribution from $\mu_{++++}^{R,2}$. Then, using (2.25), we can rewrite (5.7) as

$$\mu_{++++}^{R,2}(k,l,m,n,p)$$

$$= \sum \int \sigma_{\epsilon_0}(x) \overline{\psi_{S,\epsilon_0}}(x,k) \sigma_{\epsilon_1}(x) \psi_{S,\epsilon_1}(x,l) \sigma_{\epsilon_2}(x) \psi_{S,\epsilon_2}(x,m) \sigma_{\epsilon_3}(x) \psi_{S,\epsilon_3}(x,n) \sigma_{\epsilon_4}(x) \psi_{S,\epsilon_4}(x,p) dx,$$

where the sum is over

$$\epsilon_0, \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4 \in \{+, -\}$$
 with $(\epsilon_0, \epsilon_1, \epsilon_2, \epsilon_3, \epsilon_4) \neq (+, +, +, +, +), (-, -, -, -, -)$.

Note that the integrand in each integral is compactly supported due to the presence of a $\sigma_{+}(x)\sigma_{-}(x)$ factor. The structure of the integrals is similar, and, therefore, it suffices to consider only the measure

$$\mu^{R,2,1}(k,l,m,n,p) := \int \sigma_{+}(x)\overline{\psi_{S,+}(x,k)}\sigma_{-}(x)\psi_{S,-}(x,l)\psi_{S}(x,m)\psi_{S}(x,n)\psi_{S}(x,p)dx, \quad (5.49)$$

and the corresponding integral

$$\mathcal{N}_{R,2,1}(s,k) = \int \frac{e^{is\Phi}}{\langle l \rangle \langle m \rangle \langle n \rangle \langle p \rangle} \widetilde{g}(l) \widetilde{g}(m) \widetilde{g}(n) \widetilde{g}(p) \mu^{R,2,1}(k,l,m,n,p) dlmnp \quad \text{with } \Phi = \Phi_{++++}.$$

Using (2.30) and (5.49), we can rewrite

$$\mathcal{N}_{R,2,1}(s,k) = 4\pi^2 e^{is\langle k \rangle} \int \sigma_+(x) \sigma_-(x) \overline{\psi_{S,+}(x,k)} u_-(x) u_S^3(x) dx, \qquad (5.50)$$

where $u(x) = \langle \widetilde{D} \rangle^{-1} e^{-is\langle \widetilde{D} \rangle} g(s, x)$ and u_{-} stands for

$$u_{-}(x):=\frac{1}{\sqrt{2\pi}}\int_{\mathbb{R}}\psi_{S,-}(x,k)\widetilde{u}(k)dk.$$

We apply the ∂_k differentiation to (5.50) and obtain

$$\partial_{k} \mathcal{N}_{R,2,1}(s,k) = 4is\pi^{2} \frac{k}{\langle k \rangle} e^{is\langle k \rangle} \int \sigma_{+}(x) \sigma_{-}(x) \overline{\psi_{S,+}(x,k)} u_{-}(x) u_{S}^{3}(x) dx$$

$$+ 4\pi^{2} e^{is\langle k \rangle} \int \sigma_{+}(x) \sigma_{-}(x) \partial_{k} \overline{\psi_{S,+}(x,k)} u_{-}(x) u_{S}^{3}(x) dx$$

$$=: \mathcal{C}_{1} + \mathcal{C}_{2}.$$

Both terms C_1 and C_2 are treated using the presence of the compactly supported function $\sigma_+(x)\sigma_-(x)$ and the local decay estimates. We also observe that, similarly to (5.46) and (5.47), we have

$$\left\| \int \partial_k^r \overline{\psi_{S,+}(x,k)} h(x) \ dx \right\|_{L^2_k} \lesssim \|\langle x \rangle^r h\|_{L^2_x}, \qquad r = 0, 1.$$

Then, using (2.40) and an analogous (2.40)-type estimate for u_{-} , we have

$$\left\| \int_{0}^{t} C_{1}(s,k) ds \right\|_{L_{k}^{2}} \lesssim \int_{0}^{t} s \left\| \sigma_{-} \sigma_{+} u_{-} u_{S}^{3} \right\|_{L_{x}^{2}} ds$$

$$\lesssim \int_{0}^{t} s \left\| \langle x \rangle^{-2} u_{-} \right\|_{L_{x}^{\infty}} \left\| \langle x \rangle^{-2} u_{S} \right\|_{L_{x}^{\infty}}^{3} ds \lesssim \|v\|_{X}^{4}.$$

Similar steps ensure that $\left\| \int_0^t C_2(s,k) ds \right\|_{L^2_k} \lesssim \|v\|_X^4$. Combining the above estimates, we get

$$\left\| \int_0^t \partial_k \mathcal{N}_{R,2,1}(s,k) ds \right\|_{L^2_k} \lesssim \|v\|_X^4$$

which completes the proof (5.41) for j = 2.

5.3.3. Proof of Lemma 5.1. The proof follows from (5.9) and the estimates (5.10) and (5.41).

References

- [1] Agmon S. Spectral properties of Schrödinger operators and scattering theory. Ann. Scuola Norm. Sup. Pisa Cl. Sci. 4 (1975) no. 2, pp. 151-218.
- [2] Bates P.W. and Jones C.K.R.T. Invariant manifolds for semilinear partial differential equations. *Dynamics reported*, Vol. 2, Dynam. Report Ser. Dynam. Systems Appl. 2, Wiley, Chichester (1989) pp. 1-38.
- [3] Bizon P., Chmaj T. and Szpak N. Dynamics near the threshold for blowup in the onedimensional focusing nonlinear Klein-Gordon equation. *J. Math. Phys.* 52, 103703 (2011) 11.
- [4] Cazenave T. Uniform estimates for solutions of nonlinear Klein-Gordon equations, *Journal* of Functional Analysis 60 (1985) pp. 36-55.
- [5] Chang S.-M., Gustafson S., Nakanishi K. and Tsai Tai-P. Spectra of Linearized Operators for NLS Solitary Waves. SIAM J. Math. Anal. 39 (2007) no. 4, pp. 1070–1111.
- [6] Chen G. and Pusateri F. The 1d nonlinear Schrödinger equation with a weighted L^1 potential. Preprint arXiv:1912.10949. To appear in *Analysis & PDE*.
- [7] Chen G. and Pusateri F. On the 1d cubic NLS with a non-generic potential. Preprint arXiv:2205.01487.

- [8] Cuccagna S. and Maeda M. A survey on asymptotic stability of ground states of nonlinear Schrödinger equations II. *Discrete and Continuous Dynamical Systems S* 14 (2021) no. 5, pp. 1693-1716.
- [9] Cuccagna S. and Maeda M. Revisiting asymptotic stability of solitons of nonlinear Schrödinger equations via refined profile method. Preprint arXiv:2111.02681.
- [10] Cuccagna S. and Maeda M. Asymptotic stability of kink with internal modes under odd perturbation. Preprint arXiv:2203.13468.
- [11] Cuccagna S., Maeda M. and Scrobogna S. Small energy stabilization for 1D Nonlinear Klein Gordon Equations. Preprint arXiv:2206.08012.
- [12] Cuccagna S. and Pelinovsky D.E. The asymptotic stability of solitons in the cubic NLS equation on the line. *Appl. Anal.* 93 (2014) pp. 791–822.
- [13] Deift P. and Trubowitz E. Inverse scattering on the line. Comm. Pure. Appl. Math 32 (1979) no. 2, pp. 121-251.
- [14] Delort J.M. and Masmoudi N. Long time dispersive estimates for perturbations of a kink solution of one dimensional cubic wave equations. < hal-02862414> (2020).
- [15] Dunford N. and Schwartz J. Linear operators. Part II. Spectral theory. Selfadjoint operators in Hilbert space. Reprint of the 1963 original. *Wiley Classics Library*. A Wiley-Interscience Publication. John Wiley & Sons, Inc., New York (1988).
- [16] Faddeev L. D. The inverse problem in the quantum theory of scattering. *Uspekhi Mat. Nauk.* 14 (1959) pp. 57-119; *J. Math. Phys.* 4 (1963) pp. 72-104.
- [17] Faddeev L. D. Properties of the S-matrix of the one-dimensional Schrödinger equation. Trudy Mat. Inst. Steklov 73 (1964) pp. 314-333; AMST 2 65, pp. 139-166.
- [18] Germain P., Hani Z. and Walsh S. Nonlinear resonances with a potential: multilinear estimates and an application to NLS. *Int. Math. Res. Not.* IMRN (2015) no. 18, pp. 8484-8544.
- [19] Germain P. and Pusateri F. Quadratic Klein-Gordon equations with a potential in one dimension. Preprint arXiv:2006.15688. To appear in *Forum Math. Pi*.
- [20] Germain P., Pusateri F. and Rousset F. The nonlinear Schrödinger equation with a potential in dimension 1. Ann. Inst. H. Poincaré C (2018) no. 6, pp. 1477-1530.
- [21] Germain P., Pusateri F. and Zhang Z. On 1d quadratic Klein-Gordon equations with a potential and symmetries Preprint arXiv:2202.13273.
- [22] Kowalczyk M., Martel Y. and Muñoz C. On asymptotic stability of nonlinear waves. Séminaire Laurent Schwartz - Équations aux dérivées partielles et applications. Année 2016-2017. Ed. Éc. Polytechnique Palaiseau (2017) Exp. No. XVIII.
- [23] Kowalczyk M., Martel Y. and Muñoz C. Kink dynamics in the ϕ^4 model: Asymptotic stability for odd perturbations in the energy space. J. Amer. Math. Soc. 30 (2017) no. 3, pp. 769-798.
- [24] Kowalczyk M., Martel Y. and Muñoz C. Soliton dynamics for the 1D NLKG equation with symmetry and in the absence of internal modes. *J. Eur. Math. Soc.* (2021), published online first.
- [25] Kowalczyk M., Martel Y., Muñoz C. and Van Den Bosch H. A sufficient condition for asymptotic stability of kinks in general (1 + 1) scalar fields. *Ann. PDE* 7 (2021) no. 1, Paper No. 10, 98 pp.
- [26] Krieger J. and Schlag W. Stable manifolds for all monic supercritical focusing nonlinear Schrödinger equations in one dimension. *J. Amer. Math. Soc.* 19 (2006) no. 4, pp. 815-920.
- [27] Krieger J., Nakanishi K. and Schlag W. Global dynamics above the ground state energy for the one-dimensional NLKG equation. *Math. Z.* 272 (2012) no. 1-2, pp. 297-316.

- [28] Levine H.A. Instability and nonexistence of global solutions of nonlinear wave equations of the form $Pu_{tt} = -Au + F(u)$, Trans. Amer. Math. Soc. 192 (1974) pp. 1-21.
- [29] Léger T. and Pusateri F. Internal modes and radiation damping for quadratic Klein-Gordon in 3D. Preprint arXiv:2112.13163.
- [30] Li Y. and Lührmann J. Soliton dynamics for the 1D quadratic Klein-Gordon equation with symmetry. Preprint *arXiv:2203.11371*.
- [31] Lindblad H., Lührmann J. and Soffer A. Asymptotics for 1D Klein-Gordon equations with variable coefficient quadratic nonlinearities. *Arch. Rational Mech. Anal.* 241 (2021), no. 3, 1459-1527.
- [32] Lindblad H., Lührmann J., Schlag W. and Soffer A. On modified scattering for 1D quadratic Klein-Gordon equation with non-generic potentials. To appear in *IMRN*.
- [33] Maeda M. and Yamazaki Y. Center stable manifold for ground states of nonlinear Schrödinger equations with internal modes. Preprint arXiv:2206.08156.
- [34] Lührmann J. and Schlag W. Asymptotic stability of the sine-Gordon kink under odd perturbations. Preprint arXiv:2106.09605.
- [35] Nakanishi K. and Schlag W. Invariant manifolds and dispersive Hamiltonian evolution equations. *Zurich Lectures in Advanced Mathematics*. European Mathematical Society (EMS), Zürich, 2011. vi+253 pp.
- [36] Nakanishi K. and Schlag W. Global dynamics above the ground state for the nonlinear Klein-Gordon equation without a radial assumption. *Arch. Ration. Mech. Anal.* 203 (2012) no. 3, pp. 809-851.
- [37] Schlag W. Stable manifolds for an orbitally unstable nonlinear Schrödinger equation. *Ann. of Math.* (2) 169 (2009) no. 1, pp. 139-227.
- [38] Wang Y.J. A sufficient condition for finite time blow up of the nonlinear Klein-Gordon equations with arbitrarily positive initial energy. *Proc. Amer. Math. Soc.* 136 (2008) pp. 3477–3482.
- [39] Yafaev D. Mathematical scattering theory. Analytic theory. *Mathematical Surveys and Monographs*, 158. American Mathematical Society, Providence, RI, 2010. xiv+444 pp.
- [40] Yang Y.B. and Xu R.Z. Finite time blow up for nonlinear Klein-Gordon equations with arbitrarily positive initial energy. *Appl. Math. Lett.* 77 (2018) pp. 21-26.

University of Toronto, Department of Mathematics, 40 St George Street, Toronto, ON, M5S 2E4, Canada.

Email address: kairzhan@math.toronto.edu

University of Toronto, Department of Mathematics, $40~\mathrm{St}$ George Street, Toronto, ON, M5S $2\mathrm{E}4$, Canada.

Email address: fabiop@math.toronto.edu