Uniform, rigid branchwise-real trees

Sam Adam-Day*

*Mathematical Institute, University of Oxford, Andrew Wiles Building, Radcliffe Observatory Quarter, Woodstock Road, Oxford, OX2 6GG, United Kingdom; adamday@maths.ox.ac.uk

10th March 2023

Abstract

A branchwise-real tree is a partial order which is a tree and in which every branch is isomorphic to a real interval. I give constructions of such trees which are both rigid (i.e. without non-trivial order-automorphisms) and uniform (in two different senses). Specifically, I show that there is a rigid branchwise-real tree in which every branching point has the same degree, one in which every point is branching and of the same degree, and finally one in which every point is branching of the same degree and which admits no order-preserving function into the reals. Trees are grown iteratively in stages, and a key technique is the construction (in ZFC) of a family of colourings of $(0, \infty)$ which is 'sufficiently generic', using these colourings to determine how to proceed with the construction.

1 Introduction

This article concerns the construction of branchwise-real trees which are rigid — i.e. without non-trivial order-automorphisms. Branchwise-real trees were first defined in [FJ04] and further elaborated on in [Ada22].¹ They are tree partial orders in which every branch is order-isomorphic to a real interval, and in which every two elements have a meet (see Section 2 for definitions).

Branchwise-real trees were originally motivated from the study of \mathbb{R} -trees. Informally, an \mathbb{R} -tree is a metric space tree in which every point is permitted to be branching. They play a role in geometric group theory, in which one considers some infinite group acting by isometries on an \mathbb{R} -tree. For more information, see [Bes97; MNO92] and the references contained in [Fab15]. Now, fixing any point p in an \mathbb{R} -tree, one can consider the *cut-point order*: the set of paths through the tree from p, ordered by path extension. The resulting partial order is a branchwise-real tree [FJ04, p. 50]. Conversely, any branchwise-real tree which admits an order-preserving function into the reals is the cut-point order of some \mathbb{R} -tree [FJ04; Ada22]. The present study of the order-automorphisms of branchwise-real trees is thus motivated by the study of the isometries of \mathbb{R} trees.

Keywords: branchwise-real tree, tree automorphism, rigid, uniform, forcing, R-tree

²⁰²⁰ Mathematics Subject Classification: 03E04, 06A07, 20E08, 54F05

¹In [FJ04] branchwise-real trees are called 'non-metric trees'.

A different motivation comes the investigation of the automorphisms of wellstratified trees (tree partial orders in which every branch is isomorphic to an ordinal) [GS64; Jen69; Jec72; Avr79; AS85; FH09]. If *X* is a well-stratified tree without uncountable branches, we can turn it into a branchwise-real tree by taking its 'road space', essentially replacing every node with a copy of the interval [0, 1), so that every branch becomes isomorphic to a real interval [Jon65; Ada22]. Every automorphism of *X* can then be extended to an automorphism of its road space (but not vice versa).

However, the class of branchwise-real trees is more general than that obtained by taking the road spaces of well-stratified trees without uncountable branches. This is because any point can be branching, so that the structure of the branching points can be far from well-stratified. Consider for instance the 'comb' structure given by taking a 'shaft' [0, 1] and adding a 'tooth' — a copy of (0, 1] — as a new branch from each point on the shaft. See Figure 1 for a picture. Every point along the shaft except the maximal one is branching of degree 2. Moreover, the task of constructing a rigid branchwise-real tree is more challenging than in the well-stratified case. Indeed, any part of the tree which looks like a real interval and which contains no branching points allows for an easy automorphism which fixes the rest of the tree and permutes the interval in some non-trivial way.




Figure 1: A representation of the comb branchwise-real tree, consisting of the shaft and a few example teeth

In this article, I construct examples of rigid branchwise-real trees subject to two uniformity conditions, the latter stronger than the former. First, I construct one in which every branching node has the same degree κ , for $2 \le \kappa \le c$. I call this condition 'weakly uniformly κ -branching'. Second, using a different technique, I construct one in which every node is branching and of the same degree κ , for $2 \le \kappa \le c^+$. This condition I call 'uniformly κ -branching'. This second tree admits an order-preserving function into the reals, and thus corresponds to the cut-point order of some \mathbb{R} -tree. The technique can be further adapted to produce such a uniformly κ -branching, rigid branchwise real tree which does not admit an order-preserving function into the reals, and thus is not related to an \mathbb{R} -tree. These results manifest the phenomenon of the marriage of two seemingly opposing notions: rigidity and uniformity. As such they continue in the vein of similar results found for well-stratified trees [Avr79; FH09].

All trees here are grown recursively in stages. We start with a single point: the root. At successor stages, we choose some number of points added at the previous stage, and add $(\kappa - 1)$ many new 'spines' — copies of the interval $(0, \infty)$ — above. (Throughout $\kappa - 1$ denotes n - 1 if $\kappa = n$ finite and κ otherwise). We never extend directly above a spine once it has been added. At limit stages there may be new branches appearing through the tree, consisting of portions added at each of the previous stages. We decide which of these to extend by adding a new point on top.

A common technique used throughout when growing rigid trees is to first produce a collection of colourings of the interval $(0, \infty)$. As the tree grows, we take for each new spine added a different, unused colouring and lay it along the spine. In this way, we also build up a colouring of the whole tree as we go. These colourings are then used to determine which points will form the bases of new spines at successor stages, and which new branches to extend at limit stages. The collection of colourings of $(0, \infty)$ is carefully constructed so as to enable the desired properties of the final tree.

The constructions of the trees in this article, though they occur in ZFC, have strong forcing flavour to them. Indeed, I first built these rigid trees using forcing notions, before realising that related constructions could be carried out in ZFC. I will briefly elaborate on two of these forcing notions. For more details on forcing, see [Hal17] and [Jec03; Kun13]. However, knowledge of forcing is not required to understand any of the ZFC constructions in this article.

2 Background

Let me begin by fixing some terminology and notation relating to partial orders. Let *P* be a partial order. A *chain* in *P* is a linearly-ordered subset. A *branch* through *P* is a maximal chain. A *ray* is a final segment of a branch. A subset $Q \subseteq P$ is *coinitial* if for every $x \in P$ there is $y \in Q$ such that $y \leq x$. Elements $x, y \in P$ are *comparable* if $x \leq y$ or $y \leq x$. Two subsets $Q, R \subseteq P$ are *comparable* if $x \leq y$ or $y \leq x$. Two subsets $Q, R \subseteq P$ are *comparable* if a set of pairwise incomparable elements. For any $x \in P$ let:

$$\downarrow(x) := \{ y \in X \mid y \leq x \}, \qquad \uparrow(x) := \{ y \in P \mid y \geq x \}$$

For any x < y in *P*, define:

$$[x, y] := \{z \in P \mid x \le z \le y\}$$

Define the other intervals [x, y), (x, y] and (x, y) analogously. A function $f: P \rightarrow Q$ between partial orders is *order-preserving* if whenever x < y we have f(x) < f(y). I will also call such a function a *Q*-grading of *P*. An *iso-morphism* between *P* and *Q* is a bijection $f: P \rightarrow Q$ which is order-preserving with order-preserving inverse.

I can now introduce the main object of study: branchwise-real trees. I follow the presentation given in [Ada22].

Definition 1. A *tree order* is a partial order *X* such that the following conditions hold.

(TO1) For every $x \in X$ the set $\downarrow(x)$ is a linear order.

(TO2) X has a minimum element, its root.

Definition 2. A *branchwise-real tree* is a tree order X subject to the following extra conditions.²

- (BR1) Every branch is order-isomorphic to a real interval.
- (BR2) For any $x, y \in X$, the set $\{z \in X \mid z \leq x, y\}$ has a maximum element $x \wedge y$, the *meet* of x and y (in other words, X is a meet-semilattice).

We will also meet trees in which every branch is well-ordered. Such trees will be called 'well-stratified trees'; in a purely set theoretic context, we would simply say 'trees'.

Definition 3. A tree order *T* is *well-stratified* if every branch is order-isomorphic to an ordinal. The *rank* of an element $x \in T$ is the order type of $\downarrow(x) \setminus \{x\}$.

In order to define the *degree* of a point in a tree order we need the notion of a connected component above that point.

Definition 4. Let *X* be a tree order and $x \in X$. A *connected component above x* is an equivalence class of $\uparrow(x) \setminus \{x\}$ under the relation:

 $y \sim_x z \iff$ there is w > x such that $w \le y, z$

Note that, when *X* is a meet-semilattice, we have $y \sim_x z$ if and only if $y \wedge z > x$. See Figure 2 for an example illustration of this relation. I will usually drop the 'connected' and refer to these as 'components above *x*'. The *degree*, deg(*x*), of *x* is the number of components above *x*. Say that *x* is *terminal* if deg(*x*) = 0. Say that *x* is *branching* if deg(*x*) > 1. The *degree* of *X* is the supremum of the degrees of its elements.




Figure 2: An example illustrating the relation \sim_x which defines components above x. We have $y_0 \sim_x z$ because $x < w \le y_0, z$, but $z \not\sim_x y_1$ because there is no element above x which lies below both y_1 and z.

The notion of *continuous gradability* is important for branchwise-real trees and is the main property studied in [Ada22].

Definition 5. Let *X* be a branchwise-real tree. An \mathbb{R} -grading $\ell: X \to \mathbb{R}$ is *continuous* if for any x < y in *X* the restriction:

$$\ell [[x, y]: [x, y] \to [\ell(x), \ell(y)]$$

is an order-isomorphism. I will usually drop the ' \mathbb{R} ' and call such functions 'continuous gradings'. Say that *X* is *continuously gradable* if it admits a continuous grading.

Theorem 6. A branchwise-real tree is continuously gradable if and only if it is \mathbb{R} -gradable.

²I make a slight change of notation compared with [Ada22], where these objects are referred to as 'branchwise-real tree orders'.

Proof sketch. See [Ada22, Theorem 30]. An arbitrary \mathbb{R} -grading f of a branchwise-real tree X may contain a number of discontinuities. Using a basic result from real analysis, it can be shown that such discontinuities must take the form of 'jumps' in value, and that any branch may contain only countably many. The proof proceeds to eliminate every jump in f, using a Zorn's Lemma style argument.

Two final pieces of notation. Throughout, tuples will be denoted using angle brackets: $\langle a, b, \ldots \rangle$. A partial function between sets *X* and *Y* will be denoted using $f : X \rightarrow Y$.

3 How to grow branchwise-real trees

The trees in this article are grown using an iterative process. We always start with a singleton as the 'root'. At a successor step, above each point introduced in the previous stage we can add any number of new 'spines'. A spine is a copy of the positive real numbers $(0, \infty)$, or the interval $(0, \infty)$, which is 'terminal', in the sense that we will never extend the tree above the end of a spine. In other words, in the final tree each spine will be a ray. Call the former type of spine an *open spine* and the latter a *closed spine*. At a limit step, we first take the union of the previous stages. There is more to do however, since although we do not extend above spines, new branches through the tree appear at the limit, which we may choose to extend or not. A newly appearing branch contains the root, follows a stage-1 spine partway, then branches onto a stage-2 spine, and so on, cofinally in the limit. Such a branch thus consists of a little piece from each previous stage of the construction, with no final piece. If we decide to extend it, we add a new point directly above. This process is illustrated in Figure 3.




Figure 3: An example of growing a branchwise-real tree for $\omega + 2$ steps. Spines beginning at the same vertical height are added at the same stage. Each stage is also represented using a different colour. At stage ω , there are two new branches through the tree, the one on the right and the one on the left. We only extend the right, and we do so by adding a point at the limit, which is then further extended in subsequent stages.

This construction may continue for ω_1 -many steps. Note however that since every branch must be order-isomorphic to a real interval, and there is no increasing ω_1 -sequence of reals, no branch through the final tree can contain a piece from every stage below ω_1 .

Remark 7. A similar construction was used by Urysohn in [Ury27] (in German) to construct a metric space which is nowhere separable. Urysohn essentially followed the process outlined above, adding c-many open spines above each point added at the previous step, for ω -many steps. Of course, Urysohn was constructing a topological space, not a partial order, but there is a clear correspondence between the two types of structure. This correspondence is elaborated in more detail in [Ada22]. In [Ber19], Urysohn's construction is defined (in English) and extended to all countable steps, by taking the Hausdorff completion at limit steps.

Now, given any branchwise-real tree *X* constructed in the fashion described above, we can naturally define a rank function $\rho: X \to \omega_1$, where $\rho(x)$ is the stage α at which *x* was introduced. As Lemma 9 below demonstrates, any branchwise-real tree can be realised using the construction above, and thus admits a rank function. Note that rank functions are not unique in general. In fact, the same tree can have rank functions with a variety of ranges, as Figure 4 demonstrates.




Figure 4: An alternative construction of the tree in Figure 3, which yields a different rank function. The root gets rank 0. Both the branch which passes through every spine on the left and the branch which passes through every spine on the right have rank 1. The rest of the tree has rank 2. While the original construction produced $\omega + 2$ ranks, this only produces 3.

Definition 8. Let *X* be a tree order. A *rank function* on *X* is an order-preserving function $\rho: X \rightarrow \gamma$, where γ is some ordinal, such that the following hold.

- (RF1) For every branch *B* in *X* the set $\{\rho(x) \mid x \in B\}$ is downwards-closed.
- (RF2) If α is 0 or a limit, then $\rho^{-1}[\{\alpha\}]$ is an antichain.
- (RF3) If α is a successor then $\rho^{-1}[\{\alpha\}]$ is the disjoint union of a family of incomparable rays.

Lemma 9. Every branchwise-real tree admits a rank function.

Proof. Let *X* be a tree order. By Zorn's Lemma, there is a maximal partial rank function ρ whose domain is a downwards closed subset $Y \subseteq X$. Suppose for a

contradiction that there is $x \in X \setminus Y$. Consider $\downarrow(x) \cap Y$. Note that $\downarrow(x) \cap Y$ is non-empty, since we can always give the root rank 0. Furthermore, we can assign any limit of ranked points the rank which is the limit of the ranks of those points. Hence $\downarrow(x) \cap Y$ has a maximum element *y*. Let *C* be the component above *y* which contains *x*, and pick a maximal ray *R* through *C* which contains *x*. Then we can extend ρ so that *x* becomes ranked by giving every element of *R* rank $\rho(y) + 1$.

Remark 10. Lemma 9 shows that the construction elaborated above of iteratively adding spines is a completely general way of building branchwise-real trees. It was necessary to permit closed spines in addition to open spines in order to allow for trees with terminal nodes. All trees produced in this article, however, are without terminal nodes. Hence all constructions from now on will use open spines exclusively.

Let us now establish some facts concerning rank functions on a branchwisereal tree *X*. Since every branch is order-isomorphic to a real interval, we can assume that the codomain of any rank function on *X* is ω_1 .

Definition 11. A rank function $\rho: X \to \omega_1$ is *bounded* if the supremum of the set $\{\rho(x) \mid x \in X\}$ is less than ω_1 .

Lemma 12. If X has a bounded rank function, then it is continuously gradable.

To prove this lemma, we make use of the following set-theoretic result due to Cantor [Can95], which in particular shows that every countable ordinal embeds into \mathbb{Q} .

Lemma 13. The set \mathbb{Q} of rationals is universal for countable linear orders: every countable linear order *X* embeds into \mathbb{Q} .

Proof. A slick proof makes use of the 'forth' part of the classical 'back-and-forth method'. Enumerate $X = \{x_n \mid n \in \omega\}$. By induction on this enumeration we then build up an embedding $r: X \to \mathbb{Q}$. Once we have $r \upharpoonright \{x_0, \dots, x_{n-1}\}$, since \mathbb{Q} is a dense linear order without endpoints, we can find $r(x_n)$ whose relative position with respect to $r(x_0), \dots, r(x_{n-1})$ is the same that of x_n with respect to x_0, \dots, x_{n-1} .

Proof of Lemma 12. Let ρ : $X \rightarrow \omega_1$ be a rank function with:

$$\gamma := \{ \rho(x) \mid x \in X \} < \omega_1$$

Let $r: \gamma \to \mathbb{Q}$ be an embedding furnished by Lemma 13. We construct an embedding $\ell: X \to \mathbb{R}$ by induction on the rank. If α is 0 or a limit, map each element in $\rho^{-1}[\{\alpha\}]$ to $r(\alpha)$. Take a successor $\beta + 1$. Then $\rho^{-1}[\{\beta + 1\}]$ is the disjoint union of a family of incomparable rays, each of which is order-isomorphic to a real interval. Take for each an isomorphism onto a sub-interval of $(r(\beta), r(\beta + 1))$, and use this to extend ℓ there. After γ -many steps, we obtain an \mathbb{R} -grading $X \to \mathbb{R}$. Then by Theorem 6, we find that X is continuously gradable.

Remark 14. The converse of Lemma 12 does not hold. A good counterexample is U_{κ} , the universal continuously gradable branchwise real tree of degree $\kappa \ge 2$, defined as follows. This is essentially the construction of the universal \mathbb{R} -tree given in [DP01]; see also [Nik89; MNO92]. Let U_{κ} be the set of functions $r: [0, a) \to \kappa$, for each nonnegative real a, which are 'piecewise constant to the right': for any $t \in [0, a)$ there is $\epsilon > 0$ such that $r \upharpoonright [t, t + \epsilon)$ is constant. Note that this condition ensures that the set of 'value-change points' — those $t \in [0, a)$ for which for every $\epsilon > 0$ there is $s \in (t - \epsilon, t)$ such that $r(s) \neq r(t)$ — is well-ordered.

Then U_{κ} becomes a branchwise-real tree under function extension. We can define a continuous grading ℓ on U_{κ} by setting $\ell(r) = a$ when $r: [0, a) \to \kappa$. With this definition, any continuously graded branchwise-real tree of degree at most κ embeds into U_{κ} in such a way that the continuous gradings agree.

Let us see briefly why U_{κ} has no bounded rank function. Suppose $\rho: X \to \gamma$ is a (surjective) rank function with $\gamma < \omega_1$. Assume that γ is a limit ordinal, the successor case being similar. By Lemma 13 we can find an order embedding $s: \gamma+1 \to [0, \infty)$ such that s(0) = 0. We can recursively define $r: [0, s(\gamma)) \to \kappa$ such that $\rho(r \upharpoonright [0, s(\alpha))) = \alpha$ for each $\alpha < \gamma$ as follows. To define r on $[s(\alpha), s(\alpha+1))$, choose a ray above $r \upharpoonright [0, s(\alpha))$ of rank $\alpha+1$. This ray is essentially a function $h_{\alpha}: [s(\alpha), \infty) \to \kappa$. Define:

$$r \upharpoonright [s(\alpha), s(\alpha+1)) := h_{\alpha} \upharpoonright [s(\alpha), s(\alpha+1))$$

At limits we take unions. But now *r* cannot have a rank (it would get rank γ), which is a contradiction. 4

In Section 5 below, we shall be considering the interaction between rank functions, connected components and order-automorphisms. The following basic facts will be useful.

Definition 15. Let $\langle X, \rho \rangle$ be a ranked tree order. Take $x \in X$ and *C* a component above *x*. The *rank* of *C* is $\rho(C) := \min\{\rho(y) \mid y \in C\}$.

Note that the rank of a connected component is always a successor.

Lemma 16. Let $\langle X, \rho \rangle$ be a ranked tree order and take $x \in X$ non-terminal.

- (1) When $\rho(x)$ is 0 or a limit, every component above x has rank $\rho(x) + 1$.
- (2) When $\rho(x)$ is a successor, there is one component above x of rank $\rho(x)$ and the rest are of rank $\rho(x) + 1$.
- Let $f: X \to X$ be an order-automorphism.
 - (3) Then f factors through a bijection of the components above x onto the components above f (x).
 - (4) If f maps the component C above x onto the component f(C) above f(x), then there is a coinitial interval $I \subseteq C$ of constant rank $\rho(C)$ which maps onto a coinitial interval $f(I) \subseteq f(C)$ of constant rank $\rho(f(C))$.
- *Proof.* (1) Let *C* be any component above *x*, and pick $y \in C$ of minimal rank. Then the restriction of ρ to $\downarrow(y)$ is surjective onto $\rho(y) + 1$, from which we conclude that $\rho(C) = \rho(y) = \rho(x) + 1$.
 - (2) Since ρ(x) is a successor, x lies on a ray all of whose elements have rank ρ(x). Then as ρ is order-preserving, the component above x through which this ray passes must have rank ρ(x). Let C be any other component above x. Then C cannot have rank ρ(x), since ρ⁻¹[{ρ(x)}] consists of the disjoint union of a family of incomparable rays. By the above argument, we have that ρ(C) = ρ(x) + 1.
 - (3) Let C be any component above x. If y', z' ∈ f(C), then there is w > x such that w ≤ f⁻¹(y'), f⁻¹(z'). Hence we have f(x) < f(w) ≤ y', z', so that y' and z' lie in the same component above f(x). By the same argument applied to f⁻¹, we see that f maps components above x onto components above f(x), and vice versa.
 - (4) There is a ray *R* through *C* all of whose elements are of rank ρ(*C*). Similarly, there is a ray *R'* through *f*(*C*) all of whose elements are of rank ρ(*f*(*C*)). Take *y'* ∈ *R'* and *z'* ∈ *f*(*R*). Then there is *w'* > *f*(*x*) such that *w'* ≤ *y'*, *z'*. This *w'* then lies on both rays. Since *f* is an automorphism, this means that the coinitial interval (*x*, *f*⁻¹(*w'*)] is mapped onto the coinitial interval (*f*(*x*), *w'*].

4 A weakly uniformly branching, rigid branchwise-real tree

In this section, I construct our first rigid branchwise-real tree. It satisfies the following uniformity condition, the stronger version of which will be considered in the next section.

Definition 17. A tree order is *weakly uniformly* κ *-branching* if every branching node has degree κ .

Theorem 18. Let $2 \le \kappa \le c$. There is a weakly uniformly κ -branching, rigid branchwise-real tree which is continuously gradable.

Proof. We first construct a collection of dense, mutually non-isomorphic sets of positive reals $\{S_A \subseteq (0, \infty) \mid A \subseteq \omega\}$, such that S_A 'encodes' A.³ The set S_A consists of the rationals in $(0, \infty)$ plus a descending sequence of real intervals with limit 0 whose endpoints are irrational, such that the *n*th interval is open when $n \notin A$ and half-open when $n \in A$. Formally, fix some descending sequence (x_n) of irrationals with limit 0. Then for any $A \subseteq \omega$ we define:

$$S_A := (\mathbb{Q} \cap (0, \infty)) \cup \bigcup_{n \in \omega} (x_{2n+1}, x_{2n}) \cup \{x_{2n} \mid n \in A\}$$

See Figure 5 for a representation of an example S_A .




Figure 5: A representation of an example S_A set. Solid regions contain every real, while dotted regions only contain the rationals. A curved boundary on a solid region signifies that it does not include that endpoint. The set $A \subseteq \omega$ is written in descending order, so as to coincide with the ordering of the real intervals. Arrows relate the elements of A with the corresponding endpoints of real intervals which code for them in S_A .

Suppose that $p: S_A \to S_B$ is an order-isomorphism for $A \neq B$. Then p must map the real intervals onto the real intervals. Moreover, the rightmost real interval in S_A must map to the rightmost real interval in S_B , and so on. But A and B disagree on some natural number, say $n \in A \setminus B$ (without loss of generality). Then the *n*th real interval from the right in S_A is half-open, and p sends it to the *n*th real interval in S_B , which is open. Since we ensured that the endpoints of the real interval in S_A has a supremum. 4

Our rigid branchwise-real tree *X* is now constructed in ω -many stages using open spines, laying along each new spine a different set *S*_{*A*}, and using this to determine which points to use as the bases of new spines. Note that we need to

³In fact, we only need that the S_A 's are mutually non-isomorphic as subsets of $(0, \infty)$, but the construction yields sets which are mutually non-isomorphic as stand-alone linear orders.

make sure that we have fresh S_A sets available at each stage of the construction: we don't want to run out. For this, we can partition $\mathcal{P}(\omega)$ into ω -many batches of size \mathfrak{c} , so that the *n*th batch is reserved for the *n*th stage.

We construct *X* in stages X_n for $n \in \omega$. Start with X_0 the singleton tree, and form X_1 by adding κ -many new open spines above the root. Now assume that X_n is constructed for $n \ge 1$. To each spine *U* added at stage *n*, associate a different, unused subset $A \subseteq \omega$. Fix an isomorphism $p_U: U \to (0, \infty)$ and above each element in $p_U^{-1}(S_A)$ add $(\kappa - 1)$ -many new open spines. Finally, let $X := \bigcup_{n \in \omega} X_n$. Define a rank function $\rho: X \to \omega$, so that for each $n \in \omega$ we have $\rho^{-1}[\{n + 1\}] = X_{n+1} \setminus X_n$.

Note that at each stage of the construction, we made sure that any branching points got degree κ . Furthermore, ρ is a bounded rank function on *X*, hence by Lemma 12 we have that *X* is continuously gradable.

Let us see that X is rigid. Suppose for a contradiction that $f: X \to X$ is a non-trivial order-automorphism. Then there is $x \in X$ such that $f(x) \neq x$. There must then be y > x such that f([x, y]) and [x, y] are disjoint. Since each S_A is dense in $(0, \infty)$, there must be $z \in [x, y]$ which is branching. Choose any spine U above z added during the construction. Then f(U) is a ray through $\uparrow(f(z))$ and $f \upharpoonright U : U \to f(U)$ is an isomorphism such that $w \in U$ is branching if and only if $f(w) \in f(U)$ is branching.

Let us now consider the pattern of branching nodes along U and f(U). Following U, this looks like some S_A : it consists in an alternating sequence of real intervals and rational intervals, which converge at the base. Hence we must have the same pattern along f(U). Now, there three possibilities for the way in which f(U) fits into our construction. (i) f(U) is a single spine added above f(z). (ii) f(U) consists of an initial segment of a spine added above f(z), followed by an initial segment of a spine added at the next stage (followed potentially by further initial segments of spines added at later stages). (iii) f(U) starts part-way up a spine added at an earlier stage. See Figure 6 for a representation of the three types of patterns of branching points which can occur along f(U). Considered as subsets of $(0, \infty)$, no two of three possible patterns of branching nodes can be order-isomorphic. Since the pattern of branching nodes along U looks like (i), the pattern along f(U) must also have this form.




Figure 6: A comparison three possible types of pattern which can occur along f(U). Solid regions contain every real, while dotted regions only contain the rationals.

Therefore, there are $A, B \subseteq \omega$ distinct, such that the pattern of branching nodes along U looks like S_A , and the pattern of branching nodes along f(U)looks like S_B . In other words, we have order-isomorphisms $p_U: U \to (0, \infty)$ and $p_{f(U)}: f(U) \to (0, \infty)$ such that $p_{f(U)} \circ f \circ p_U^{-1}: (0, \infty) \to (0, \infty)$ is an order-automorphism mapping S_A onto S_B . This is our contradiction. \pounds

Let us now consider the relationship between the construction just given and a forcing notion. For ease of exposition, we only deal with the case $\kappa =$ \aleph_0 . We force a weakly uniformly \aleph_0 -branching, continuously gradable, rigid branchwise-real tree by considering countable approximations to it. Such objects might be called sub-branchwise-real trees: we replace (BR1) with the requirement that each branch be embeddable into \mathbb{R} . In addition, we keep track of a rank function from the approximations into ω , together with a set of 'non-extension promises'. The latter is a subset of the sub-branchwise-real tree consisting of degree-1 nodes which guarantees that these nodes never become branching. The forcing conditions are thus triples (X, ρ, S) consisting of: (a) a countable sub-branchwise-real tree X (which for concreteness we can take to be a subset of ω_1), (b) a rank function $\rho: X \to \omega$, and (c) a non-extension promise set $S \subseteq X$ of degree-1 nodes. For one condition to be stronger than another, we require the sub-branchwise-real tree, rank function and promise set of the former extend those of the latter, and further that we do not add new connected components above any already-branching point, so that in particular the degrees of branching points do not decrease. To ensure this latter condition, we require that for every element x of the weaker approximation, every component above x with respect to the stronger approximation contain at most one component above x with respect to the weaker. Note that the fact that stronger conditions must extend the promise sets and that every element of the promise set must have degree 1 ensures that we keep our promises. Let \mathbb{P} be the forcing notion just described.

Theorem 19. The forcing notion \mathbb{P} is countably closed, and any \mathbb{P} -generic set determines an ω -ranked, weakly uniformly \aleph_0 -branching, rigid branchwise-real tree.

Proof sketch. That \mathbb{P} is countably closed follows immediately from the definition. Denote by \overline{X} the union of the approximations found in *G*. It is not hard to see by genericity that \overline{X} is a weakly uniformly \aleph_0 -branching branchwise-real tree, and that the union of the partial rank functions yields a total rank function $\overline{X} \to \omega$.

To see that \overline{X} is rigid, take $\dot{f}: \overline{X} \to \overline{X}$ a non-trivial automorphism. By a countable-closure argument, we can find a condition $\langle X, \rho, S \rangle$ forcing (i) that \dot{f} is a order-preserving function, (ii) that $X \subseteq \text{dom}(\dot{f})$ and that \dot{f} has a decided value on every element of X and (iii) that there is $x \in X$ such that $\dot{f}(x)$ is incomparable with x. Moreover we can assume, using another countable-closure argument, that there is $z \ge x$ in X such that both z and $\dot{f}(z)$ are branching, and furthermore that there is w > z such that $[z, w]^X$ is order-dense as subset of $[z, w]^{\overline{X}}$. Since $[z, w]^{\overline{X}}$ is countable, it has continuum-many 'holes' to fill. Any element which fills a hole has its image under \dot{f} fixed, given then density of $[z, w]^X$ is filled with a branching node while the corresponding hole in $[\dot{f}(z), \dot{f}(w)]^X$ is filled with a non-branching node which is promised to remain non-branching. Since G is generic, this means that \dot{f} cannot be an automorphism of \overline{X} , which is a contradiction. \dot{f}

5 A uniformly branching, rigid branchwisereal tree

In this section, I strengthen the result of the previous, by showing that for every κ with $2 \le \kappa \le c^+$ there is a rigid branchwise-real tree in which every point is branching of the same degree κ .

Definition 20. A tree order is *uniformly* κ *-branching* if every node has degree κ .

Moreover, we can ask that such a tree be either continuously gradable (Theorem 21) or not (Theorem 29).

Theorem 21. Let $2 \le \kappa \le c^+$. There is a uniformly κ -branching, rigid branchwisereal tree which is continuously gradable.

As in the proof of Theorem 18, our uniformly branching trees will be grown iteratively upwards. But notice that, since we require every point to have the same degree κ , at successor stages there are no choices to make: we must add (κ – 1)-many new spines above every point added at the previous stage. Were we to stop this process after ω -many steps, we would end up with a 'minimal' uniformly branching tree, which is not only non-rigid, but moreover homogeneous, as the following result shows.

Definition 22. Let κ be a cardinal. The *minimal uniformly* κ -branching branchwise-real tree, M_{κ} , is the branchwise-real tree built in ω -many steps, starting with the root, and at successors stages adding (κ – 1)-many new open spines above every point added at the previous stage.

Remark 23. The tree M_c is the (underlying partial order of) the nowhere separable metric space constructed by Urysohn in [Ury27].

Proposition 24.

- (1) M_{κ} is minimal: for every uniformly κ -branching branchwise-real tree X there is an order-preserving embedding $f: M_{\kappa} \to X$. Moreover, we can assume that f is continuous: for any x < y in M_{κ} the restriction $f \upharpoonright [x, y] : [x, y] \to [f(x), f(y)]$ is surjective.
- (2) M_{κ} is homogeneous: if $x, y \in M_{\kappa}$ are not the root, there is an orderautomorphism $f: M_{\kappa} \to M_{\kappa}$ such that f(x) = y.

Proof. First note the construction of M_{κ} yields a rank function $\rho: M_{\kappa} \to \omega$.

- We can build an order-preserving embedding M_κ → X using a Zorn's Lemma style argument, essentially following the proof of Lemma 9.
- (2) First take any $z \in M_{\kappa}$ such that $n = \rho(z) > 1$. We define an automorphism $r_z: M_{\kappa} \to M_{\kappa}$ which steps down *z*'s rank: $\rho(r_z(z)) = n - 1$. Let *w* be the greatest element of $\downarrow(z)$ of rank n-1. Then z lies on a spine added above w at stage n. Let C be the component above w which contains z(which component is of rank n). Let D be the component above w which contains the rest of the rank-(n-1) spine on which w lies (which is of rank n-1). See Figure 7 for a picture of the situation. It is easy to define an isomorphism $s: C \to D$ such that the rank-*n* spine on which *z* lies is mapped to the rest of the rank-(n-1) spine above w. Letting r_z be the automorphism which swaps *C* and *D* via *s*, we see that $\rho(r_z(z)) = n - 1$. Now take $x, y \in M_{\kappa}$ non-root. By repeating the above procedure, we can assume that both have rank 1. Moreover, by performing another 'twist', sending the component above the root containing y to that containing x, we may assume that x and y are comparable, lying on the same spine; say $x \leq y$. Finally, it is not hard to define an automorphism of M_{κ} whose restriction to the rank-1 spine containing x and y looks like an automorphism of $(0, \infty)$, and which maps *x* to *y*.

Turning back to the construction of rigid trees, in the proof of Theorem 21, as well as Theorem 29 below, the notion of a 'colouring' of the positive real numbers $(0, \infty)$ plays an auxiliary role. A colouring of the positive real numbers is simply a function with domain $(0, \infty)$; we think of elements of the range as 'colours'.




Figure 7: The situation when reducing *z*'s rank. The point *z* lies on a rank-*n* spine added above *w*, in the component *C* above *w*. The rank-(n - 1) spine on which *w* lies continues into the component *D* above *w*. The automorphism r_z swaps the components *C* and *D*.

Definition 25. Let *X* and *S* be sets. An *S*-colouring of *X* is a function $X \rightarrow S$.

In the proof of Theorem 18, the first step was essentially to construct a family of black-white colourings of $(0, \infty)$ with suitable properties. To construct a uniformly branching, rigid branchwise-real tree, we extend this idea. This time we look for a family of 2-colourings (and later ω -colourings) which is 'sufficiently generic'. This means that for any pair of distinct colourings, any order-automorphism of $(0, \infty)$ and any pair of colours, we can densely often find a point coloured with the first colour under the first colouring whose image under the automorphism is coloured with the second colour under the second colouring. This is the following result.

Lemma 26 (Generic Colouring Lemma). Let $\lambda \leq c$ be a cardinal. There is a family \mathscr{A} with size c^+ of λ -colourings $(0, \infty) \to \lambda$ of the positive real numbers such that the following holds. For any $c, d \in \mathscr{A}$ distinct, for any order-automorphism $p: (0, \infty) \to (0, \infty)$ and for any $\alpha, \beta \in \lambda$ there is a dense set of $x \in (0, \infty)$ such that $c(x) = \alpha$ and $d(p(x)) = \beta$.

To prove this we need the following basic result concerning automorphisms of the positive real numbers.

Lemma 27. There are exactly c-many order-automorphisms $(0, \infty) \rightarrow (0, \infty)$.

Proof. Every real number is the limit of rationals below it, so any orderautomorphism $(0, \infty) \rightarrow (0, \infty)$ is determined by its values on $(0, \infty) \cap \mathbb{Q}$, of which there are c-many possibilities.

Proof of Lemma 26. By Zorn's Lemma, we can take a maximal family \mathscr{A} of λ -colourings, satisfying both the property in the statement, as well as that for every $c \in \mathscr{A}$:

for every $\alpha \in \lambda$ and for every b > a > 0 there are continuum-many points $x \in (a, b)$ such that $c(x) = \alpha$ (P)

Let us first see that \mathscr{A} is non-empty. For convenience, I will construct a (0, 1)colouring satisfying (P); this can then easily be adapted to a $(0, \infty)$ -colouring. It suffices to take $\lambda = \mathfrak{c}$. Consider the elements of (0, 1) as binary ω -sequences representing binary expansions (where, to avoid ambiguity, we make some canonical choice for the binary representation of $x \in (0, 1)$, in the case that two such representations are possible). Define an equivalence relation on these, where two sequences are related if one is a tail of the other. Each equivalence class is countable, hence there are continuum-many. Moreover, each equivalence class is dense in (0, 1). Group the classes into continuum-many batches of size \mathfrak{c} , and colour all of each batch with a different element of \mathfrak{c} . The resulting colouring satisfies (P).

Now suppose for a contradiction that $|\mathcal{A}| < c^+$. We will extend \mathcal{A} by diagonalising against the previous colourings, and against every automorphism. Let Aut $(0, \infty)$ be the set of order-automorphisms $(0, \infty) \rightarrow (0, \infty)$. By Lemma 27, the set Aut $(0, \infty) \times \mathcal{A} \times \lambda \times \lambda$ has size c. Enumerate:

Aut
$$(0, \infty) \times \mathscr{A} \times \lambda \times \lambda = \{(p_{\theta}, c_{\theta}, \alpha_{\theta}, \beta_{\theta}) \mid \theta < \mathfrak{c}\}$$

We build a new colouring $d: (0, \infty) \to \lambda$ recursively in stages d_{θ} for $\theta < \mathfrak{c}$, each of which is a partial colouring with domain of size less than \mathfrak{c} . Assume we have constructed d_{μ} for $\mu < \theta$. First let $d'_{\theta} := \bigcup_{\mu < \theta} d_{\mu}$. Now, since c_{θ} satisfies (P), and since d'_{θ} has domain of size less than \mathfrak{c} , we can find a countable dense set of $X \subseteq (0, \infty)$ such that for every $x \in X$ we have $c_{\theta}(x) = \alpha_{\theta}$ while $d'_{\theta}(p_{\theta}(x))$ is undefined. Extend d'_{θ} to d_{θ} by letting $d_{\theta}(p_{\theta}(x)) := \beta_{\theta}$, for every $x \in X$. Finally, let d be the union $\bigcup_{\theta < \mathfrak{c}} d_{\theta}$, filling in any points which remain uncoloured arbitrarily. But now, $\mathcal{A} \cup \{d\}$ is a larger family, contradicting the maximality of \mathcal{A} .

With this lemma established, we can now construct our uniformly κ branching, rigid branchwise-real tree which is continuously gradable.

Proof of Theorem 21. Let \mathscr{A} be a \mathfrak{c}^+ -sized family of 2-colourings of $(0, \infty)$ as per the Generic Colouring Lemma 26. This time, we construct X in stages X_a for $\alpha < \omega^2$. When we add a new spine we lay a new colouring from \mathscr{A} along it, and thus we also build colourings $c_\alpha: X_\alpha \to 2$. At limit stages, we use these colourings to decide which branches to extend. We also keep track of the rank ρ of elements of X, so that the points added at stage α get rank α . As before, we need to make sure we never run out of colourings, so we can partition \mathscr{A} into ω^2 -many batches of size \mathfrak{c}^+ .

Start with X_0 the singleton, coloured 0. Assume that X_α and c_α are constructed, where α is 0 or a limit. To make $X_{\alpha+1}$, above all of the points of rank α , we add κ -many new open spines. For each spine added, we pick a different, unused colouring $c \in \mathcal{A}$, and use it to define $c_{\alpha+1}$ on that spine. Now assume that X_α and c_α are constructed, where α is a successor. To make $X_{\alpha+1}$, above each point of rank α , we add ($\kappa - 1$)-many new open spines, again colouring them with new colourings from \mathcal{A} .

So take $\alpha < \omega^2$ a limit, and assume we have constructed X_β and c_β for $\beta < \alpha$. First let $X'_\alpha := \bigcup_{\beta < \alpha} X_\alpha$ and $c'_\alpha := \bigcup_{\beta < \alpha} c_\alpha$. Then X'_α admits a number of new branches which appear in the limit: in other words, branches through X'_α containing elements of rank unbounded in α . It is these branches which we will decide to extend or not. Let *B* be any such branch. Then for every $\beta < \alpha$, there is a maximum element $x_\beta \in B$ of rank β . Consider the sequence $(c'_\alpha(x_0), c'_\alpha(x_1), \ldots)$ of the colours of such points. We will extend *B* if and only if this sequence has a tail consisting of 1's. To extend *B*, add a new point to X'_α lying directly above *B*, and colour it 0. Once we have carried out this procedure for every new branch through X'_α , we arrive at our new pair $\langle X_\alpha, c_\alpha \rangle$.

Finally, let $X := \bigcup_{\alpha < \omega^2} X_\alpha$ and $c := \bigcup_{\alpha < \omega^2} c_\alpha$. Note that at each stage we made sure that every point is branching of degree κ , so X is uniformly κ -branching. Moreover, $\rho : X \to \omega^2$ is a bounded rank function, so X is continuously gradable by Lemma 12.




Figure 8: Picking x_1 when $\kappa \ge 3$. The grey regions represent connected components, while thin lines represent spines. The ordinal above a spine indicates its rank.

Let us see that *X* is rigid. Suppose for a contradiction that $f: X \to X$ is a nontrivial automorphism. Then there is $x_0 \in X$ such that $f(x_0) \neq x_0$. When $\kappa \ge 3$, the proof is simpler, so let's deal with that case first. See Figure 8 for a representation of the situation. Consider the components above above x_0 and above $f(x_0)$. By Lemma 16, *f* factors through a bijection of the components above x_0 to those above $f(x_0)$. Moreover, all but at most one component above x_0 is of rank $\rho(x_0) + 1$, and similarly for $f(x_0)$. Since deg $(x_0) = \deg(f(x_0)) = \kappa \ge 3$ there is a component *C* above x_0 of rank $\rho(x_0)+1$ which maps to a component f(C) above $f(x_0)$ of rank $\rho(f(x_0)) + 1$. Furthermore, using Lemma 16 again, there is an coinitial interval $I \subseteq C$ of constant rank $\rho(x_0) + 1$ which maps to a coinitial interval $f(I) \subseteq f(C)$ of constant rank $\rho(f(x_0)) + 1$. Then *I* is an initial segment of a spine *U* added at stage $\rho(x_0) + 1$, while f(I) is an initial segment of a spine *V* added at stage $\rho(f(x_0)) + 1$.

Now, consider the colouring *c* along *I* and along *f*(*I*). By construction, these come from different colourings from \mathscr{A} . The order-isomorphism $f \upharpoonright I : I \to f(I)$ induces an isomorphism $U \to V$, which in turn induces an automorphism of $(0, \infty)$. Hence by the key property of the family \mathscr{A} guaranteed by the Generic Colouring Lemma 26, there is $x_1 \in I$ coloured $c(x_1) = 1$ such that its image under *f* is coloured $c(f(x_1)) = 0$.

Iterate this process to produce a sequence $x_0 < x_1 < \cdots$ such that $c(x_n) = 1$ and $c(f(x_n)) = 0$ for every n > 0. Then $\{x_0, x_1, \ldots\}$ determines a branch Bthrough $X'_{\rho(x_0)+\omega}$, while $\{f(x_0), f(x_1), \ldots\}$ determines a branch f(B) through $X'_{\rho(f(x_0))+\omega}$. Moreover, when deciding whether to extend B at stage $\rho(x_0) + \omega$, we use a sequence whose tail is (x_0, x_1, \ldots) , and respectively for f(B). But then the former branch gets extended, while the latter does not, contradicting that f is an automorphism. 4

Let us now turn to the case $\kappa = 2$. The issue here is that we can no longer guarantee that an initial segment of a new spine added above x_0 gets mapped to an initial segment of a new spine added above $f(x_0)$. Here's how we proceed. If the component *C* above x_0 of rank $\rho(x_0) + 1$ is mapped to the component above $f(x_0)$ of rank $\rho(f(x_0)) + 1$, proceed as before. Otherwise, the rank of f(C) must be $\rho(f(x_0))$. See Figure 9 for a representation of the situation. By Lemma 16 there is a coinitial interval $I \subseteq C$ of constant rank whose image $f(I) \subseteq f(C)$ has constant rank. So *I* is an initial segment of a spine added above x_0 at stage $\rho(x_0)+1$ which maps onto a segment of a spine which already exists




Figure 9: Picking x_1 when $\kappa = 2$ and $\rho(f(C)) = \rho(f(x_0))$. The dark and light grey regions represent connected components, while thin lines represent spines. The ordinal above a spine indicates its rank, where $\alpha_0 = \rho(x_0)$ and $\beta_0 = \rho(f(x_0))$.

at stage $\rho(f(x_0))$. Now, using the key property of the colouring \mathscr{A} , pick any element x'_0 in the interior of I with colour $c(x'_0) = 1$ whose image has colour $c(f(x'_0)) = 0$. Then we must have that the component C' above x'_0 of rank $\rho(x'_0) + 1$ is mapped to the component above $f(x'_0)$ of rank $\rho(f(x'_0)) + 1$, and so we can proceed as before to obtain $x_1 > x'_0$ coloured $c(x_1) = 1$ whose image is coloured $c(f(x_1)) = 0$. Iterating, we again build a sequence $x_0 < x'_1 < x_1 < \cdots$ through X, all of whose elements (except possibly x_0) are coloured 1 but with image coloured 0. When deciding whether to extend the corresponding branch through $X'_{\rho(x_0)+\omega}$, we use the colours of a subsequence of (x_0, x'_1, x_1, \ldots) . In the end, we find that this sequence has a limit in X, while its image does not. \pounds

With this ZFC result established, let us examine once more the connection with a forcing notion. We again work with the $\kappa = \aleph_0$ case. We will define forcing conditions similarly to the discussion at the end of Section 4. This time however, instead of making promises that some points won't become branching, we promise that certain branches won't get ever get extended. There are a number of different ways of achieving this. (1) The most direct way is to include in each condition a set of branches through the sub-branchwise-real tree, which are promised never to be extended. A technical issue which arises is that a branch through a smaller sub-branchwise-real tree need not be a branch through a larger one, since it need not be maximal as a chain any more. To make sure that the set of promises remains valid, we require that a stronger condition include a promise set which contains the unique extension of each promised branch in the weaker tree to the larger one. (2) A more elegant method is to keep track instead of an order-preserving function from the approximations into \mathbb{R} . This way the branches which are not to be extended are exactly those on which the order-preserving function is unbounded. (3) Yet a third way is make use of the universal continuously-gradable branchwise-real tree U_{κ} defined in Remark 14. Any maximal antichain A through U_{κ} defines a uniformly κ -branching branchwise-real tree: take { $x \in U_{\kappa} \mid \exists a \in A: x < a$ }. We can then consider simply the set of all antichains in U_{κ} under extension as a forcing poset. For the following result, I remain agnostic as to the exact method used.

Theorem 28. A generic tree \overline{X} for the forcing just described is a uniformly κ -

branching, rigid branchwise-real tree.

Proof sketch. The hard part is again showing that \overline{X} is rigid. Take $f: \overline{X} \to \overline{X}$ a non-trivial automorphism. By a countable-closure argument, we can find a condition p_0 forcing that \dot{f} is a non-trivial order-preserving function whose domain contains the approximation X_0 at p_0 with \dot{f} decided on every element of X_0 , and with an element $x_0 \in X_0$ such that $f(x_0)$ is incomparable with x_0 . By another countable-closure argument, it is dense to find a condition $p_1 < p_0$ such that the domain of f contains the approximation X_1 at p_1 and such that there is $x_1 \in X_1$ not less than any element in X_0 with $x_0 < x_1$. Continuing in this way, we build a chain of forcing conditions $p_0 > p_1 > \cdots$ together with $x_0 < x_1 < \cdots$ in $\bigcup_{n \in \omega} X_n$. Let p_{ω} be the limit of the p_n 's. Then we can strengthen p_{ω} so that the branch determined by $\{x_0, x_1, \ldots\}$ gets extended, while we promise never to extend the branch determined by $\{\dot{f}(x_0), \dot{f}(x_1), \ldots\}$. (In the case where we keep track of an order-preserving function into \mathbb{R} , we would need to make a tweak to ensure that one sequence remains bounded while the other becomes unbounded.) Therefore, we can densely find a contradiction to the fact that fis order-preserving. \$

Turning back to our ZFC constructions, the tree constructed in Theorem 21 is continuously gradable. By suitably modifying the method, we can construct one which is not continuously gradable.

Theorem 29. Let $2 \le \kappa \le c^+$. There is a uniformly κ -branching, rigid branchwisereal tree which is not continuously gradable.

This time, the construction must proceed for ω_1 -many steps, because Lemma 12 shows that any construction terminating after countably many steps would yield a continuously gradable tree. This then introduces a complication, as we need to ensure that there are no branches through the final tree which contain points of every rank below ω_1 . Indeed, such a branch would contain an ω_1 -sequence, violating condition (BR1) that every branch is isomorphic to a real interval. Now, as mentioned in Remark 14, not every tree constructed in ω_1 -many steps is non-continuously-gradable, so we also need to guarantee this. This is done by realising a known non-R-gradable well-stratified tree inside the final branchwise-real tree, following techniques used in [Ada22]. Lastly, these two competing requirements need to be balanced with the need for the final tree to be rigid. These desiderata are achieved following the method used in the proof of Theorem 21 above, this time utilising ω -gradings instead of 2-gradings. At limit steps, the idea is essentially to extend a branch if and only if in the sequence of colours $(c'_a(x_0), c'_a(x_1), \ldots)$ of the maximal points of each rank, every element of ω appears only finitely often.

First, let us meet the non- \mathbb{R} -gradable well-stratified tree without uncountable branches which will be found inside the final branchwise-real tree.

Lemma 30. Let *T* be the tree of functions *r* into ω , whose domain is a countable ordinal, such that when restricted to the set of successor ordinals *r* is injective into $\omega \setminus \{0\}$, and elsewhere *r* is identically 0. We consider *T* as a partial order under function extension. Then *T* is a well-stratified tree without an \mathbb{R} -grading.

Proof. The tree *T* is isomorphic to the tree Shift(In_{ω}) defined in Section 3 of [Ada22]. There, using a proof due to Baumgartner, Gavin and Laver [Bau70], it is shown that this tree is well-stratified, has no \mathbb{R} -grading, and moreover that every branch is countable.

With this to hand, we can now prove the theorem.

Proof of Theorem 29. Let \mathscr{A} be a \mathfrak{c}^+ -sized family of $(\omega \setminus \{0\})$ -colourings of $(0, \infty)$ as per the Generic Colouring Lemma 26. We construct *X* in stages X_a

for $\alpha < \omega_1$, colouring each spine appearing at a stage which is the successor of a successor using a different colouring from \mathscr{A} . As in Theorem 21, these colourings are used to determine which branches to extend at limit stages. This will ensure that the resulting tree *X* is rigid, that every branch is isomorphic to a real interval and that there is no continuous grading.

Proceed as in Theorem 21. First partition \mathscr{A} into ω_1 -many batches of size \mathfrak{c}^+ , so that we never run out. We build each X_a and colouring c_a , together with the rank function ρ , recursively. Start with X_0 the singleton coloured 0. As before, at successor stages add $(\kappa - 1)$ -many new open spines above the points added at the previous stage. If we are at the successor of 0 or a limit, colour the new spines all with 0, otherwise colour each using new colourings from \mathscr{A} . Now assume that $\alpha < \omega_1$ is a limit. First let $X'_{\alpha} := \bigcup_{\beta < \alpha} X_{\beta}$ and $c'_{\alpha} := \bigcup_{\beta < \alpha} c_{\beta}$. We need to decide which new branches through X'_{α} to extend. As before, a branch *B* is determined by its sequence (x_0, x_1, \ldots) of maximum points of each rank $\beta < \alpha$. Extend the branch *B* if and only if no non-zero colour appears infinitely often in the sequence $(c'_{\alpha}(x_0), c'_{\alpha}(x_1), \ldots)$. To extend a branch, add a new point above it coloured 0. Let X_a and c_a be result of performing this operation on every branch in X'_{α} .

Finally, let $X := \bigcup_{a < \omega_1} X_a$ and $c := \bigcup_{a < \omega_1} c_a$. Note that at each stage we made sure that every point is branching of degree κ , so X is uniformly κ -branching. To see that X is branchwise-real, take any branch B through X. Then B is determined by its sequence $(x_0, x_1, ...)$ of maximum points of each rank $\beta < \omega_1$. Consider the sequence $(c_a(x_0), c_a(x_1), ...)$. Every colour indexed by the successor of a successor is non-zero, and every non-zero colour appears at most countably many times. Hence the sequence must have countable length. Therefore, the branch B cannot contain a subset isomorphic to ω_1 , so it must be isomorphic to a real interval, as required.

To show that *X* is rigid, we follow the method in Theorem 21. If $f: X \to X$ is a non-trivial automorphism, we can pick x_0 such that $f(x_0) \neq x_0$. Note that if we consider the sequence of points below x_0 , respectively $f(x_0)$, maximal in each rank, then no non-zero colour appears infinitely often. So to determine whether a branch containing x_0 gets extended at a limit, it suffices to consider the sequence of colours of points maximal in each rank lying *above* x_0 , and likewise for $f(x_0)$. As in Theorem 21 we can construct a sequence $x_0 < x_1 < \cdots$ recursively, this time ensuring that $c(x_n) = n$ while $c(f(x_n)) = 1$, for each n > 2. Note that while any of $x_0, x_1, f(x_0), f(x_1)$ may be coloured 0, none of the rest of the points or their images can be, since they all have ranks which are the successor of a successor. Then the sequence (x_0, x_1, \ldots) has a limit in *X*, while its image does not.

Finally, to see that *X* has no \mathbb{R} -grading, we realise the tree *T* given by Lemma 30 as a subtree. For this, we recursively construct an embedding $\iota: T \to X$ such that for every $r \in T$:

• $\rho(\iota(r)) = \operatorname{rank}(r)$, and

• $c(\iota(r)) = r(\alpha)$ when rank $(r) = \alpha + 1$.

Start by sending the root \emptyset to the root.

Take $r \in T$ and assume that we have defined $\iota(r)$. There are two cases. If rank(r) is 0 or a limit, then r has exactly one immediate successor s. Moreover $\rho(\iota(r))$ is 0 or a limit, so by the way we constructed X there are κ -many spines with base $\iota(r)$ added at stage rank(r) + 1, all of which are coloured 0. Let $\iota(s)$ be any point on any one of these spines. The other case is when rank(r) is a successor. Then r has κ -many successors. Moreover, there are κ -many spines with base $\iota(r)$ added at stage rank(r) + 1, each of which are coloured according to a colouring from \mathscr{A} . In particular, every non-zero colour appears

on every spine. Place each successor *s* of *r* on a different spine, in such a way that $c(\iota(s)) = s(\operatorname{rank}(r))$.

Now take $r \in T$ with rank(r) a limit, and assume that we have defined $\iota(s)$ for all s < r. Consider the sequence $(\iota(r \upharpoonright 0), \iota(r \upharpoonright 1), ...)$. This determines a branch through $X'_{\text{rank}(r)}$. Moreover, when determining whether to extend this branch, the sequence of colours used is:

 $(c(\iota(r \mid 0)), c(\iota(r \mid 1)), \ldots) = (0, r(0), r(1), \ldots, 0, r(\omega), r(\omega + 1), \ldots)$

By definition of *T*, no non-zero colour appears more than once in this sequence. Therefore, the branch is extended, meaning that the sequence $(\iota(r \mid 0), \iota(r \mid 1), ...)$ has a supremum, which we can set as the image of *r*.

Putting it all together, we obtain an embedding $\iota: T \to X$. Any continuous grading of the latter would restrict to an \mathbb{R} -grading of the former, which by Lemma 30 is impossible.

6 Open questions

In this final section I present a number of questions left open by the preceding investigation.

The strongest result in this article is that there exists both a continuously gradable and a non-continuously-gradable uniformly κ -branching, rigid branchwise-real tree for $2 \le \kappa \le c^+$. This limit of c^+ comes from the Generic Colouring Lemma 26. It is natural to wonder if we could get past this limit using some alternative strategy.

Question 31. Does there exist a uniformly κ -branching, rigid branchwise-real tree for $\kappa > c^+$?

Now, it may be that such a question is beyond ZFC, and that the best we can hope for is a consistency result. We might try expanding the Generic Colouring Lemma by forcing both a 2^c-sized family of colourings and that $2^c > c^+$. Or alternatively we could try to force our rigid tree directly, for example by using a forcing notion similar to that presented in Section 5.

Taking a different tack, throughout the present article we have been aiming to eliminate all order-automorphisms. What if wanted more fine-grained control over the automorphism group? It is not hard to modify the methods presented here to yield a branchwise-real tree with automorphism group (i) the symmetries of κ as a set, (ii) \mathbb{Z} , (iii) the order-automorphism group of the positive reals or (iv) various combinations of these.

Question 32. What is the class of automorphism groups of (uniformly κ -branching) branchwise-real trees?

Finally, we have seen four examples of uniformly κ -branching branchwisereal trees: the minimal M_{κ} , the universal U_{κ} , the rigid continuously gradable one and the rigid non-continuously-gradable one. None of these trees are isomorphic. Moreover, each comes with a rank function, the supremum of whose values is, respectively, ω , ω_1 , ω^2 and ω_1 . Clearly, any branchwise-real tree may admit many different rank functions, but is there some canonical way in which we might stratify uniformly κ -branching branchwise-real trees in terms of their 'complexity'? One direction of investigation might be to first investigate how these tree embed into one another. We already know that M_{κ} is a minimum and U_{κ} a maximum, as least for the continuously gradable ones.

Question 33. What is the nature of the class of uniformly κ -branching (continuously gradable) branchwise-real trees under embeddability?

Alternatively, we might consider the following isomorphism-invariant of branchwise-wise real trees X: what is the minimum supremum value of a rank function on X? For M_{κ} , this is ω , and for any non-continuously gradable branchwise-real tree, this must be ω_1 (by Lemma 12). It is not immediate that the uniformly κ -branching, rigid, continuously gradable branchwise-real tree constructed in Section 5 has minimum supremum rank ω^2 ; however it is not hard to see that its minimum cannot be ω . This leads to the following questions.

Question 34. Which ordinals arise as the minimum supremum value of a rank function on a uniformly κ -branching branchwise-real tree?

7 Acknowledgements

I would like to thank my supervisor, Joel David Hamkins, for much helpful guidance and proof-reading throughout the process of doing this research and writing the present article. I also wish to thank the anonymous reviewer, whose suggestions led to many improvements of the paper. This work was supported by the EPSRC [studentship with project reference *2271793*].

References

- [Ada22] Sam Adam-Day. 'On the continuous gradability of the cut-point orders of R-trees'. In: *Topology and its Applications* 306 (2022), p. 107937. ISSN: 0166-8641. DOI: 10.1016/j.topol.2021.107937. arXiv: 2107.14718 [math.L0].
- [AS85] Uri Abraham and Saharon Shelah. 'Isomorphism types of Aronszajn trees'. In: Israel Journal of Mathematics 50 (1985), pp. 75–113.
- [Avr79] Uri Avraham. 'Construction of a rigid Aronszajn tree'. In: Proceedings of the American Mathematical Society 77.1 (1979).
- [Bau70] James E. Baumgartner. 'Results and Independence Proofs in Combinatorial Set Theory'. PhD thesis. University of California, 1970. ISBN: 9798659828038.
- [Ber19] Valerii Nikolaevich Berestovskii. 'On Urysohn's R-Tree'. In: Siberian Mathematical Journal 60.1 (2019), pp. 10–19.
- [Bes97] Mladen Bestvina. *Real trees in topology, geometry, and group* theory. 1997. arXiv: math/9712210 [math.GT].
- [Can95] Georg Cantor. 'Beiträge zur Begründung der transfiniten Mengenlehre'. In: *Mathematische Annalen* 46 (1895), pp. 481–512.
- [DP01] Anna Dyubina and Iosif Polterovich. 'Explicit Constructions of Universal R-Trees and Asymptotic Geometry of Hyperbolic Spaces'. In: Bulletin of the London Mathematical Society 33.6 (Nov. 2001), pp. 727–734. arXiv: math/9904133 [math.DG].

- [Fab15] Paul Fabel. 'A Topological Characterization of the Underlying Spaces of Complete R-Trees'. In: *Michigan Mathematical Journal* 64 (2015), pp. 881–887. DOI: 10.1307/mmj/1447878035.
- [FH09] Gunter Fuchs and Joel David Hamkins. 'Degrees of Rigidity for Souslin Trees'. In: *The Journal of Symbolic Logic* 74.2 (2009), pp. 423–454. ISSN: 00224812, 19435886. DOI: 10.2178/jsl/1243948321.
- [FJ04] Charles Favre and Mattias Jonsson. The valuative tree. Lecture notes in mathematics 1853. Berlin; London: Springer, 2004. ISBN: 9783540229841. DOI: 10.1007/b100262. arXiv: math/0210265 [math.AC].
- [GS64] Haim Gaifman and E. Specker. 'Isomorphism Types of Trees'. In: Proceedings of The American Mathematical Society 15 (Feb. 1964), pp. 1–1. DOI: 10.1090/S0002-9939-1964-0168484-2.
- [Hal17] Lorenz J. Halbeisen. Combinatorial Set Theory. With a Gentle Introduction to Forcing. 2nd ed. Springer Monographs in Mathematics. Springer Cham, 2017. ISBN: 978-3-319-60230-1. DOI: 10.1007/978-3-319-60231-8.
- [Jec03] Thomas J. Jech. *Set Theory*. The Third Millennium Edition. Springer Monographs in Mathematics. Springer-Verlag, 2003. ISBN: 978-3-540-44085-7. DOI: 10.1007/3-540-44761-X.
- [Jec72] Thomas J. Jech. 'Automorphisms of ω1-Trees'. In: *Trans*actions of the American Mathematical Society 173 (1972), pp. 57–70. ISSN: 00029947.
- [Jen69] Ronald Björn Jensen. 'Automorphism properties of Souslin continua'. In: *Notices of the American Mathematical Society* 16 (1969), p. 576.
- [Jon65] P. Burton Jones. 'Remarks on the normal Moore space metrization problem'. In: *Topology Seminar Wisconsin*. Vol. 60.
 Princeton: Princeton University Press, 1965, pp. 115–120.
 ISBN: 9781400882076. DOI: 10.1515/9781400882076-015.
- [Kun13] Kenneth Kunen. *Set theory*. Revised edition. Studies in logic 34. London: College Publications, 2013. ISBN: 9781848900509.
- [MNO92] John C. Mayer, Jacek Nikiel and Lex G. Oversteegen. 'Universal Spaces for R-Trees'. In: *Transactions of the American Mathematical Society* 334.1 (1992), pp. 411–432. ISSN: 00029947. DOI: 10.2307/2153989.
- [Nik89] Jacek Nikiel. *Topologies on pseudo-trees and applications*. Vol. 416. American Mathematical Society, 1989.
- [Ury27] Paul Urysohn. 'Beispiel eines nirgends separablen metrischen Raumes'. German. In: *Fundamenta Mathematicae* 9 (1927), pp. 119–121.