# WHY DO THE SYMMETRIES OF THE MONSTER VERTEX ALGEBRA FORM A FINITE SIMPLE GROUP?

#### SCOTT CARNAHAN

ABSTRACT. Together with the 1988 construction of the monster vertex algebra, Frenkel, Lepowsky, and Meurman showed that the largest sporadic simple group, known as the Fischer-Griess monster, forms the symmetry group of an infinite dimensional algebraic object whose construction was motivated by theoretical physics. However, the fact that the symmetry group is in fact finite and simple ultimately relied on highly non-trivial group-theoretic results used in Griess's work on the monster. We prove some properties of the automorphism group of the monster vertex algebra  $V^{\natural}$ , most notably that it is is finite and simple, using recent developments in the theory of vertex operator algebras, but mostly 19th century group theory.

#### CONTENTS

1. Introduction	1
1.1. Acknowledgements	2
2. Two Niemeier lattices	2
2.1. The Golay code construction	3
2.2. The Leech Lattice	4
3. Vertex algebraic ingredients	5
3.1. Basic definitions	5
3.2. Lattice vertex algebras	7
3.3. The diagonal octahedral group	9
3.4. Modules	10
3.5. Holomorphic orbifolds	12
4. The monster vertex algebra	14
4.1. Extensions of Conway groups	15
4.2. Finiteness	16
4.3. Simplicity	16
5. What else can we derive about the monster from vertex-algebraic methods?	17
6. Open problems	19
References	20

#### 1. INTRODUCTION

Let us imagine a world much like our own, but where finite group theory developed much more slowly than 2-dimensional conformal field theory and the theory of vertex algebras <sup>1</sup>. We might find that many finite simple groups would be discovered not as part of a classification program, but instead appear naturally from the symmetries of vertex algebras that are built with standard techniques. For example, most of (and possibly all of) the groups of Lie type could be found as "large subquotients" in the automorphism groups of easily constructed vertex algebras over finite

<sup>&</sup>lt;sup>1</sup>Alternatively, we may imagine a mathematician who, like the author, is reasonably familiar with recent developments in vertex algebras but is somewhat unclear with details about the classification of finite simple groups.

fields [Griess-Lam 2014], and we would find the monster simple group and its sporadic subquotients naturally through the study of orbifolds of rank 24 lattice vertex algebras.

In this paper, we describe the fundamentals of the construction of the monster vertex algebra  $V^{\ddagger}$  from [Frenkel-Lepowsky-Meurman 1988], and prove many key properties of its automorphism group, in particular that it is finite and simple, using recent results in vertex algebra theory but very little finite group theory or explicit calculation. That is, we will play a game, where we wrestle with an unwieldy opponent (Aut  $V^{\ddagger}$ ) with our strongest hand (20th century finite group theory) tied behind our back. Thus, our main group-theoretic tools for proving finiteness and simplicity are Sylow's theorem and the Frattini argument, both from the late 19th century, and we also use a basic fact from algebraic geometry, namely that a complex algebraic group with trivial Lie algebra is finite. In particular, our use of post-19th-century finite group theory is limited to properties of double covers of lattices, the Golay code, and the Leech lattice.

Our main result, that the automorphism group of the monster vertex algebra  $V^{\natural}$  is finite and simple, is clearly not new: it was first proved in [Frenkel-Lepowsky-Meurman 1988], some pages after the construction of  $V^{\natural}$  itself. However, the original proof relies on an identification with the monster simple group  $\mathbb{M}$ , constructed in [Griess 1982], where the fact that  $\mathbb{M}$  is finite and simple was proved using a lot of the machinery of the classification of finite simple groups. This path to proof has had some simplifications made: a simpler proof that  $\mathbb{M}$  is finite and simple is sketched in [Tits 1984], and a simplified construction of  $\mathbb{M}$  is given in [Conway 1985]. Furthermore, there are now many alternative constructions of  $V^{\natural}$ , e.g., in [Miyamoto 1997], [Chen-Lam-Shimakura 2016], [Abe-Lam-Yamada 2017], [Carnahan 2017]. However, until now, these simplifications have not been substantially combined.

We take ideas from several treatments of both the monster and  $V^{\natural}$ . The construction of  $V^{\natural}$  is fundamentally the same as in the original work [Frenkel-Lepowsky-Meurman 1988], but the analysis of the vertex algebra structure and symmetry is simplified by recent advances in the theory of holomorphic orbifolds. We replace a brutally explicit treatment of triality with a more intrinsically symmetric method inspired by Conway's construction of the monster in [Conway 1985], but we avoid use of finite nonassociative structures like the Parker loop. For the proofs of finiteness and simplicity, we originally used the outline from [Tits 1984], but we found that the vertex-algebraic description makes our proofs a small fraction of the original length, and in particular allowed us to avoid the explicit computations. We conclude by describing some additional facts about Aut  $V^{\natural}$ that are accessible with vertex algebraic techniques. It is interesting to compare our results with those in [Norton 2001], where structural features of the monster were worked out assuming the generalized moonshine conjecture, but not using vertex algebras.

One notable omission from this paper is the rich theory of Virasoro minimal models as a way to do explicit calculations. There is significant work in the analysis of  $V^{\natural}$  and the monster using these tools, in particular the existence of a vertex operator subalgebra of the form  $L(1/2, 0)^{\otimes 48}$ ([Dong-Mason-Zhu 1994]) and consequent results concerning symmetries (e.g., [Miyamoto 1997], [Shimakura 2006], [Shimakura 2010], [Griess-Lam 2011]) and constructions employing codes (e.g., [Dong-Griess-Höhn 1997], [Miyamoto 1996]).

1.1. Acknowledgements. I would like to thank Jacob Lurie and Akshay Venkatesh for inviting me to talk at the IAS moonshine seminar in March 2021. My preparations for the talk directly led to the writing of this paper. I would also like to thank John Duncan and Theo Johnson-Freyd for helpful comments during the talk that led to improvements in the exposition.

## 2. Two Niemeier lattices

Before we begin our treatment of vertex algebras, we need to briefly consider lattices.

**Definition 2.1.** A lattice is a free abelian group of finite rank equipped with an integer-valued bilinear form. A lattice L is even if  $(a, a) \in 2\mathbb{Z}$  for all  $a \in \mathbb{Z}$ , nondegenerate if (a, b) = 0 for

all  $a \in L$  implies b = 0, and **positive definite** if  $(a, a) \leq 0$  implies a = 0. The **dual lattice** of a nondegenerate lattice L is the set of all  $b \in L \otimes \mathbb{Q}$  such that  $(a, b) \in \mathbb{Z}$  for all  $a \in L$ . A lattice L is **unimodular** if it is equal to its dual lattice. A **root** in a positive definite even lattice is a vector a of norm 2, i.e., (a, a) = 2.

The first example of an even positive definite lattice is  $\sqrt{2}\mathbb{Z}$ , also known as the  $A_1$  root lattice, generated by a vector a satisfying (a, a) = 2. This is not unimodular, since the dual lattice is  $\frac{1}{\sqrt{2}}\mathbb{Z}$ , which contains  $\sqrt{2}\mathbb{Z}$  as an index 2 subgroup.

It is well-known that the rank of a positive definite even unimodular lattice is an integer multiple of 8. The first example is the  $E_8$  lattice, in rank 8. There are 2 isomorphism types in rank 16, 24 in rank 24 (classified by Niemeier in [Niemeier 1973]), and more than 10<sup>9</sup> in rank 32 [King 2000]. One can see that there is a phase transition around rank 24, beyond which lattices become difficult to classify.

A standard invariant of lattices is given by the generating function that enumerates vectors of a given length:

**Definition 2.2.** Let L be a positive definite even lattice. Then, the theta function of L is:

$$\theta_L(\tau) = \sum_{a \in L} q^{(a,a)/2}$$

where  $\tau$  describes a point in the complex upper half-plane, and  $q = e^{2\pi i \tau}$ .

We note that this sum converges normally to a holomorphic function on the upper half-plane, because there is a degree d polynomial in n that bounds the coefficient of  $q^n$  for all n. A short exercise in Poisson summation yields:

**Lemma 2.3.** If L is a positive definite even unimodular lattice of rank d, then  $\theta_L$  is a modular form of weight  $\frac{d}{2}$ . In particular,  $\theta_L(\frac{-1}{\tau}) = \tau^{d/2}\theta_L(\tau)$ .

In order to study the monster, we consider two rank 24 even unimodular lattices, called Niemeier lattices: one called  $N(A_1^{24})$ , because its root lattice is a product of 24 copies of the  $A_1$  lattice  $\sqrt{2\mathbb{Z}}$ , and the Leech lattice  $\Lambda$ , which is distinguished by the property that it has no roots.

2.1. The Golay code construction. Let us suppose there is a positive definite even unimodular lattice L of rank 24 with root system  $A_1^{24}$ . Then, the sublattice  $(\sqrt{2}\mathbb{Z})^{24}$  generated by roots has index  $2^{24}$  in its dual lattice  $(\frac{1}{\sqrt{2}}\mathbb{Z})^{24}$ . Thus, the unimodular condition implies L contains the root lattice as an index  $2^{12}$  subgroup, i.e.,  $L/(\sqrt{2}\mathbb{Z})^{24}$  is a subgroup C of order  $2^{12}$  in  $(\frac{1}{\sqrt{2}}\mathbb{Z})^{24}/(\sqrt{2}\mathbb{Z})^{24} \cong (\mathbb{Z}/2\mathbb{Z})^{24}$ . We may identify elements of this quotient group with subsets of the distinguished basis  $\{e_i = (0, \ldots, 0, \frac{1}{\sqrt{2}}, 0, \ldots, 0)\}$  of  $(\frac{1}{\sqrt{2}}\mathbb{Z})^{24}$ , and we find that for L to be even, it is necessary and sufficient that all elements of C correspond to subsets whose size is a multiple of 4. For L to contain no additional roots beyond the  $A_1^{24}$  root system, it is necessary and sufficient that C have no elements corresponding to subsets of size 4. In the language of codes, C is a doubly even binary code of length 24 with minimal weight 8. The following theorem then implies the lattice L exists, and is uniquely determined up to isomorphism by the conditions we specified.

**Theorem 2.4.** There exists a doubly even binary code C of length 24 with minimal weight 8, and it is unique up to permutation of the basis. The group of symmetries of this code is the sporadic simple group  $M_{24}$ .

This code is called the extended Golay code. Perhaps the most concise construction of this code takes as a basis the set  $\mathbb{Z}/23\mathbb{Z} \cup \{\infty\}$ , and takes  $\mathcal{C}$  as the subgroup generated by the size 12 set QR of quadratic residues modulo 23 together with all linear fractional transforms of QR. This construction makes a  $PSL_2(\mathbb{F}_{23})$  action clear, but many important properties are not so obvious.

There is an alternative construction in [Aschbacher 1994] that also yields a proof that  $M_{24}$  is simple, using iterated extension of Steiner systems, starting from the projective plane over  $\mathbb{F}_4$ . A third construction, using the hexacode, is given in [Conway-Sloane 1982] and [Griess 1998].

**Lemma 2.5.** The theta function of  $N(A_1^{24})$  is the unique weight 12 modular form with constant term 1 and q-coefficient 48:  $\theta_{\Lambda}(\tau) = 1 + 48q + 195408q^2 + \cdots$ . In particular, there are 48 vectors of norm 2 and 195408 vectors of norm 4.

The automorphisms of  $N(A_1^{24})$  form a group of type  $2^{24}.M_{24}$ , where the notation means there is a normal subgroup isomorphic to  $(\mathbb{Z}/2\mathbb{Z})^{24}$ , with quotient isomorphic to  $M_{24}$ . The normal 2-group is the Weyl group, generated by root reflections, and the quotient  $M_{24}$  comes from basis permutations that preserve the Golay code.

2.2. The Leech Lattice. There are many constructions of the Leech lattice  $\Lambda$ , but a construction starting from  $N(A_1^{24})$  is most convenient for us. The vector  $\frac{1}{4}\alpha_{\mathcal{C}} = \frac{1}{2\sqrt{2}}(1,\ldots,1) \in \mathbb{R}^{24}$  has inner product  $\pm \frac{1}{2}$  with all roots in our presentation of  $N(A_1^{24})$ , and in fact, the inner product with any element of  $N(A_1^{24})$  lies in  $\frac{1}{2}\mathbb{Z}$ , so the sublattice  $\Lambda_0$  whose vectors have integer inner product with  $\frac{1}{4}\alpha_{\mathcal{C}}$  has index 2, and has no roots.  $\Lambda_0$  has 3 non-identity cosets in its dual lattice, and only two of them yield even lattices, so we may construct  $\Lambda$  as the unique even lattice strictly containing  $\Lambda_0$  that is not  $N(A_1^{24})$ . This construction realizes  $\Lambda$  and  $N(A_1^{24})$  as "neighbors" in the sense of Kneser.

that is not  $N(A_1^{24})$ . This construction realizes  $\Lambda$  and  $N(A_1^{24})$  as "neighbors" in the sense of Kneser. The nontrivial coset contains the vector  $\left(-\frac{3}{\sqrt{2}}, \frac{1}{\sqrt{2}}, \dots, \frac{1}{\sqrt{2}}\right)$ . More intrinsically, we may present  $\Lambda$  as the lattice with vectors of the form  $(a_1, \dots, a_{24}) \in \left(\frac{1}{\sqrt{2}}\mathbb{Z}\right)^{24}$  satisfying:

- (1)  $\sqrt{2}\sum_{i} a_{i} = 4m$  for some integer m.
- (2) The set  $S = \{i | \sqrt{2}a_i \not\equiv m \mod 4\} \in \mathcal{C}$ , and furthermore,  $\sqrt{2}a_i \equiv m+2 \mod 4$  for all  $i \in S$ .

**Lemma 2.6.** The theta function of  $\Lambda$  is the unique weight 12 modular form with constant term 1 and vanishing q-coefficient:  $\theta_{\Lambda}(\tau) = 1 + 196560q^2 + \cdots$ . In particular, there are no vectors of norm 2 and 196560 vectors of norm 4.

Since it is not difficult to compute the coefficients of a spanning set of weight 12 forms, like  $E_4(\tau)^3 = (1 + 240 \sum_{n \ge 1} q^n \sum_{d \mid n} d^3)^3$  and  $\Delta(\tau) = q \prod_{n \ge 1} (1 - q^n)^{24}$ , it is also not difficult to enumerate the vectors in  $\Lambda$ .

**Definition 2.7.** We write  $Co_0$  to denote the automorphism group of  $\Lambda$ , and  $Co_1$  to denote the quotient of  $Co_0$  by its center  $\pm 1$ .

**Theorem 2.8.** ([Conway 1969]) The group  $Co_1$  is simple of order  $2^{21} \cdot 3^9 \cdot 5^4 \cdot 7^2 \cdot 11 \cdot 13 \cdot 23$ .

We briefly outline the elements of Conway's proof - perhaps the most sophisticated group theory lies in the use of Sylow's theorem and the Frattini argument, both from the late 19th century. The order can be computed by enumerating "frames" or "crosses", namely systems of 24 mutually orthogonal lines spanned by norm 8 vectors. The stabilizer of a cross in  $Co_1$  is a maximal nonnormal subgroup N of type  $2^{11}.M_{24}$ , where the normal 2-subgroup is given by products of reflections parametrized by Golay codewords, and  $M_{24}$  permutes the lines. To get the order, one then shows that  $Co_1$  acts transitively on crosses, and that any element of  $Co_1$  either stabilizes a given cross or takes it to a disjoint cross.

Conway's proof of simplicity uses the fact that  $Co_1$  acts primitively on lines spanned by norm 4 vectors, so any nontrivial normal subgroup H acts transitively on these lines, and hence has order dividing 13. By Frattini,  $Co_1$  is a product of H and the normalizer of a 13-Sylow subgroup, so at least one of them has order dividing 23. In the latter case, we get an element of order  $13 \cdot 23$ , contradicting a rank restriction coming from rational canonical form. In the former case, where  $|H| \in 23\mathbb{Z}$ , one notes that  $H \cap N$  is normal in N with an element of order 23, hence all of N, so by maximality of N we have  $H = Co_1$ .

#### 3. Vertex algebraic ingredients

## 3.1. Basic definitions.

**Definition 3.1.** Let V be a complex vector space. A field on V is a linear map  $V \to V((z))$ , where V((z)) denotes the space of formal Laurent series with coefficients in V, i.e., formal series of the form  $\sum_{n \in \mathbb{Z}} v_n z^{-n-1}$ , where  $v_n = 0$  for n sufficiently large. Two fields A(z) and B(z) on V are **mutually local** if there there is a positive integer N such that  $(z - w)^N[A(z), B(w)] = 0$  as an element of End  $V[[z, z^{-1}, w, w^{-1}]]$ .

Locality is a weak version of commutativity. For example, if we are given a commutative ring R, then for any  $a \in R$ , the "left multiplication by a" map can be viewed as a constant field  $b \mapsto ab = a_{-1}bz^0$  on R, and any two left-multiplication maps strictly commute. The identity element yields the identity field  $I(z) = \operatorname{id}_R z^0$ . However, fields in general cannot be composed in the way that left-multiplication composes in a commutative ring. We instead have a collection of composition laws parametrized by integers.

**Definition 3.2.** Given a pair A(z), B(z) of fields, the *m*th residue product is defined as:

$$A(z)_m B(z) = \operatorname{Res}_{y=0} A(y) B(z) (y-z)^m - \operatorname{Res}_{y=0} B(z) A(y) (y-z)^m$$
$$= \sum_{n \in \mathbb{Z}} \sum_{i \ge 0} (-1)^i \binom{m}{i} (A_{m-i} B_{n+i} - (-1)^m B_{m+n-i} A_i) z^{-n-1}$$

Here, we expand  $(y-z)^m$  as a formal power series by the "first field variable is big" convention: when attached to A(y)B(z), we set |y| > |z|, yielding the expansion  $y^m(1-z/y)^m = y^m \sum_{i\geq 0} {m \choose i} z^i y^{-i}$ , and when attached to B(z)A(y), we set |z| > |y|, yielding the expansion  $(-1)^m z^m (1-y/z)^m = (-1)^m z^m \sum_{i\geq 0} {m \choose i} z^{-i} y^i$ .

Residue products look somewhat cumbersome at first, but they satisfy the following convenient properties (see sections 1 and 2 of [Matsuo-Nagatomo 1997]):

- **Theorem 3.3.** (1) Residue products with the identity field satisfy  $A(z)_n I(z) = 0$  for  $n \ge 0$ , and  $A(z)_{-1}I(z) = A(z)$ .
  - (2) If A and B are mutually local fields on a complex vector space, then  $A(z)_n B(z) = 0$  for sufficiently large positive n.
  - (3) If A, B, C are mutually local fields on a complex vector space, then A is mutually local with  $B(z)_n C(z)$  for all  $n \in \mathbb{Z}$ .
  - (4) If A, B, C are mutually local fields on a complex vector space, then for any  $p, q, r \in \mathbb{Z}$ ,

$$\sum_{i\geq 0} \binom{p}{i} (A(z)_{r+i}B(z))_{p+q-i} C(z)$$
  
= 
$$\sum_{i\geq 0} (-1)^{i} \binom{r}{i} (A(z)_{p+r-i}(B(z)_{q+i}C(z)))_{-(-1)^{r}B(z)_{q+r-i}(A(z)_{p+i}C(z))).$$

The last is called "Borcherds's identity for residue products".

We now introduce the notion of vertex algebra.

**Definition 3.4.** A vertex algebra (over  $\mathbb{C}$ ) is a complex vector space V equipped with a multiplication map  $Y: V \otimes V \to V((z))$ , written  $u \otimes v \mapsto Y(u, z)v = \sum_{n \in \mathbb{Z}} u_n v z^{-n-1}$ , and a distinguished unit vector **1**, satisfying the following properties:

(1) (unit axiom)  $Y(u, z)\mathbf{1} \in u + zV[[z]]$  for all  $u \in V$ . That is,  $u_n\mathbf{1}$  vanishes for  $n \ge 0$  and  $u_{-1}\mathbf{1} = u$ .

(2) (locality) For any  $u, v \in V$ , the fields Y(u, z) and Y(v, z) are mutually local.

The power series  $Y(u, z) \in (\text{End } V)[[z, z^{-1}]]$  is called the "field attached to u".

The simplest examples of vertex algebras are commutative associative algebras, where we set  $Y(u, z)v = u_{-1}vz^0 = uv$ . Slightly more generally, if we are given a unital commutative associative algebra with derivation D, we may set  $Y(u, z)v = (e^{Dz}u)v$  to get a vertex algebra. These vertex algebras satisfy the property that Y(u, z)Y(v, w) = Y(v, w)Y(u, z) for all  $u, v \in V$ . Vertex algebras satisfying this property are called "commutative", and any commutative vertex algebra arises from a commutative associative algebra with derivation this way. Commutativity is equivalent to the property that  $Y(u, z)v \in V[[z]]$ , i.e., the power series given by products have no poles at z = 0. We may therefore think of a vertex algebra as a generalization of a commutative ring with derivation, where the products are allowed to have singularities at z = 0.

One may note the similarity between the unit axiom of vertex algebras and the behavior of residue products with the identity field. In fact, we have the following theorem [Li 1996]:

**Theorem 3.5.** Let M be a complex vector space, and let V be a collection of fields on M satisfying the following conditions:

- (1) V is closed under addition and scalar multiplication.
- (2) The identity field I(z) lies in V
- (3) V is closed under residue products.
- (4) All fields in V are mutually local.

Then, the assignments  $\mathbf{1} = I(z)$ ,  $Y(A(z), x) = \sum_{n \in \mathbb{Z}} A(z)_n x^{-n-1}$  endow V with the structure of a vertex algebra. Conversely, if V is a vertex algebra, then  $Y(u_n v, z) = Y(u, z)_n Y(v, z)$  for all  $u, v \in V$  and  $n \in \mathbb{Z}$ , so the map  $u \mapsto Y(u, z)$  is a vertex algebra isomorphism from V to a space of mutually local fields on V that satisfies the conditions listed above.

Combining this with the Borcherds identity for residue products, we obtain the "Jacobi identity":

**Theorem 3.6.** Let V be a vertex algebra. Then, for any  $u, v, w \in V$  and  $p, q, r \in \mathbb{Z}$ ,

$$\sum_{i\geq 0} \binom{p}{i} (u_{r+i}v)_{p+q-i} w$$
  
=  $\sum_{i\geq 0} (-1)^i \binom{r}{i} (u_{p+r-i}(v_{q+i}w))$   
 $-(-1)^r v_{q+r-i}(u_{p+i}w)).$ 

By making various substitutions to the Jacobi identity, one obtains many results, such as:

- (1) (commutator formula)  $[u_p, v_q] = \sum_{i>0} {p \choose i} (u_i v)_{p+q-i}$ : set r = 0.
- (2) (skew symmetry)  $Y(v, z)u = e^{Tz}Y(u, -z)v$ , where  $Tu = u_{-2}\mathbf{1}$ : set  $w = \mathbf{1}, p = -1, q = 0$ .
- (3) (weak associativity) For any  $u, v, w \in V$ , there is some N > 0 such that  $(y+z)^N Y(u, y+z)Y(v, z)w = (y+z)^N Y(Y(u, y)v, z)w$ : let N satisfy  $u_p w = 0$  for all p > N.

Before considering the non-commutative examples of interest, we introduce a convenient theorem for showing a collection of fields generate a vertex algebra.

**Theorem 3.7.** (Reconstruction theorem) Let V be a vector space, **1** an element of V, T an endomorphism of V, and let  $\{A^i(z)\}_{i\in J}$  be a set of fields in V including the identity field  $I(z) = id_V z^0$ . Suppose the following properties hold:

- (1)  $A^i(z)\mathbf{1} \in A^i + zV[[z]]$  for some vector  $A^i \in V$ , for each  $i \in J$ .
- (2)  $T\mathbf{1} = 0$  and  $[T, A^{i}(z)] = \frac{d}{dz}A^{i}(z)$
- (3) All  $A^i(z)$  are mutually local.
- (4) V is spanned by  $A_{n_1}^{i_1} \cdots A_{n_k}^{i_k} \mathbf{1}$ , where  $k \ge 0$  and each  $n_i < 0$ .

Then, the assignment  $Y(A_{n_1}^{i_1}\cdots A_{n_k}^{i_k}\mathbf{1},z) = A^{i_1}(z)_{n_1}\cdots A^{i_k}(z)_{n_k}I(z)$  defines a vertex algebra structure on V, and it is the unique vertex algebra structure such that  $Y(A_{-1}^i\mathbf{1},z) = A^i(z)$  for all  $i \in J$ .

We also note that [Frenkel-Lepowsky-Meurman 1988] introduced a variant of the notion of vertex algebra that has some additional structure and satisfies convenient finiteness conditions:

**Definition 3.8.** A vertex operator algebra of central charge  $c \in \mathbb{C}$  is a quadruple  $(V, Y, \mathbf{1}, \omega)$ , where  $(V, Y, \mathbf{1})$  is a vertex algebra, and  $\omega \in V$  is a **conformal vector** satisfying the following properties:

- (1) The coefficients of the field  $T(z) = Y(\omega, z) = \sum L_n z^{-n-2}$  satisfy  $[L_m, L_n] = (m-n)L_{m+n} + {\binom{m+1}{3}}\delta_{m,-n}\frac{c}{2} \operatorname{id}_V$ , i.e., V is endowed with the structure of a representation of the Virasoro algebra with central charge c.
- (2)  $L_0$  acts diagonalizably with integer eigenvalues that are bounded below and finite dimensional eigenspaces, and furthermore,  $L_0 \mathbf{1} = 0$  and  $L_0 \omega = 2\omega$ .
- (3)  $L_{-1}u = u_{-2}\mathbf{1}$  for all  $u \in V$ .

All of the vertex algebras we consider are vertex operator algebras for a natural choice of  $\omega$ . Sometimes, there are alternative choices of conformal vector, but this is not important in our treatment.

3.2. Lattice vertex algebras. One of the easiest noncommutative examples of a vertex algebra is the Heisenberg vertex algebra attached to a vector space  $\mathfrak{h}$  with inner product. We form the infinite dimensional Heisenberg Lie algebra  $\hat{\mathfrak{h}}$ , which decomposes into  $\mathfrak{h}[t, t^{-1}] \oplus \mathbb{C}K$  as a vector space, with Lie bracket determined by  $[ut^m, vt^n] = m(u, v)\delta_{m,-n}K$ ,  $[K, ut^n] = 0$  for all  $u, v \in \mathfrak{h}, m, n \in \mathbb{Z}$ . This is a central extension of the abelian Lie algebra  $\mathfrak{h}[t, t^{-1}]$ . We let  $\mathfrak{h}^+$  denote the Lie subalgebra  $\mathfrak{h}[t] \oplus \mathbb{C}K$ , and for each  $u \in \mathfrak{h}$ , we let  $\mathbb{C}_u$  denote the one-dimensional representation on which  $vt^n$ acts by  $(v, u)\delta_{n,0}$  and K acts by identity. Then, we define  $\pi_u$  (or  $\pi_u^{\mathfrak{h}}$  if  $\mathfrak{h}$  is ambiguous) to be the induced representation  $\operatorname{Ind}_{\mathfrak{h}^+}^{\mathfrak{h}}\mathbb{C}_u$ . Because  $\hat{\mathfrak{h}}$  decomposes as a direct sum of abelian subalgebras  $\mathfrak{h}^+$  and  $t^{-1}\mathfrak{h}[t^{-1}]$ , we see by the Poincaré-Birkhoff-Witt theorem that  $\pi_u$  is a free module of rank 1 for the universal enveloping algebra  $U(t^{-1}\mathfrak{h}[t^{-1}])$ , which is isomorphic to the symmetric algebra  $\operatorname{Sym}(t^{-1}\mathfrak{h}[t^{-1}])$ . In particular, given an ordered basis  $e^1, \ldots, e^d$  of  $\mathfrak{h}$ , we obtain a basis of  $\pi_u$  given by monomials of the form  $e^{i_1}t^{n_1}\cdots e^{i_k}t^{n_k}v_u$ , where  $n_1 \leq \cdots \leq n_k < 0$  and  $i_j < i_{j+1}$  if  $n_j = n_{j+1}$ .

**Theorem 3.9.** There is a unique vertex algebra structure on  $\pi_0$  with

$$1 = v_0$$
 and  $Y(e^i t^{-1}, z) = \sum_{n \in \mathbb{Z}} e^i t^n z^{-n-1}$ 

Furthermore, given an orthonormal basis  $e^1, \ldots, e^d$  of  $\mathfrak{h}$ , the vector  $\omega = \sum_{i=1}^d (e^i t^{-1})^2 \mathbf{1}$  endows  $\pi_0$  with a vertex operator algebra structure with central charge d.

The first claim is immediate from the reconstruction theorem, once we establish pairwise locality of the fields  $Y(e^{i}t^{-1}, z)$ . The essential relation  $(z - w)^{2}[Y(e^{i}t^{-1}, z)Y(e^{j}t^{-1}, w)] = 0$  follows from a straightforward calculation using the Lie bracket on  $\hat{\mathfrak{h}}$ . The grading on  $\pi_{0}$  by  $L_{0}$ -eigenvalues is non-negative, and the degree *n* part has dimension p(n), the number of partitions of *n* into positive integers. We can assemble the graded dimensions into a power series, called the character:

**Lemma 3.10.** The character of  $\pi_0$ , given by  $\sum_{n\geq 0} \dim(\pi_0)_n q^{n-d/24}$ , is equal to the modular form

$$\eta(\tau)^{-d} = q^{-d/24} \prod_{k=1}^{\infty} (1-q^k)^{-d}$$

The extra -d/24 in the definition, called the "vacuum energy shift" by physicists, is what we need to get a modular form.

We briefly note that the group of vertex algebra automorphisms of  $\pi_0$  is the orthogonal group of  $\mathfrak{h}$ . The action on the generating vectors  $\{ut^{-1}|u \in \mathfrak{h}\}$  extends naturally to  $\operatorname{Sym}(t^{-1}\mathfrak{h}[t^{-1}])$ .

We now move on to the lattice construction. Let L be an even integral lattice spanning a vector space  $\mathfrak{h}$ . Then, we wish to produce a vertex algebra structure on the direct sum  $\bigoplus_{u \in L} \pi_u$  extending that of  $\pi_0$ . There is a degenerate way to do this, where the multiplication map vanishes when restricted to  $\pi_u \otimes \pi_v$  for  $u, v \neq 0$ . However, the situation where this vanishing never happens is substantially more interesting.

**Theorem 3.11.** ([Borcherds 1986], [Frenkel-Lepowsky-Meurman 1988]) Given an even integral lattice L spanning a vector space  $\mathfrak{h}$ , there exists a unique (up to isomorphism) vertex algebra structure on the direct sum  $\bigoplus_{u \in L} \pi_u$ , subject to the condition that the multiplication map is nonzero when restricted to  $\pi_u \otimes \pi_v$  for any  $u, v \in L$ . Furthermore, if L is positive definite, then the vertex operator algebra structure on  $\pi_0$  extends to a vertex operator algebra structure on the direct sum.

A proof of this theorem using reconstruction can be found in Chapter 5 of [Frenkel-Ben-Zvi 2004]. It suffices to define a suitable field  $Y(v_u, z)$  for the generating vector  $v_u$  of each irreducible Heisenberg module  $\pi_u$ . However, such a field is uniquely determined up to a scalar by the requirements that  $Y(v_u, z)$  take  $\pi_v$  to  $\pi_{v+u}((z))$  for all  $v \in L$ , and that  $Y(v_u, z)$  be mutually local with the fields from  $\pi_0$ . We will not describe the field in detail here, but we remark that locality imposes a nontrivial condition on the scalars: We may choose to take them from  $\pm 1$ , but we have to choose signs in a way that corresponds to constructing a central extension  $\hat{L}$  of the lattice by the group  $\{\pm 1\}$ . We will write  $V_L$  for the lattice vertex algebra.

**Proposition 3.12.** Let *L* be an even integral lattice of rank *d*. Then,  $V_L$  is  $\mathbb{Z}$ -graded with weight given by eigenvalues of  $L_0$ . If *L* is positive definite, then the weights are non-negative, and the character of  $V_L$ , defined as  $\sum_{n\geq 0} \dim(V_L)_n q^{n-d/24}$ , is equal to the modular function  $\frac{\theta_L(\tau)}{\eta(\tau)^d}$ , for  $\tau$  in the complex upper half plane, where  $q = e^{2\pi i \tau}$ . Here,  $\theta_L(\tau) = \sum_{\alpha \in L} q^{(\alpha, \alpha)/2}$  is a modular form of weight d/2, and  $\eta(\tau) = q^{1/24} \prod_{n\geq 1} (1-q^n)$  is a modular form of weight 1/2.

We briefly consider automorphism groups of lattice vertex algebras, and here, we must restrict our view to positive definite lattices to ensure the groups are finite dimensional. The reason is that the weight 1 vectors generate derivations, and the weight 1 space is finite dimensional if and only if the lattice is positive definite. Specifically,  $\pi_u$  is nonzero in degrees  $\frac{(u,u)}{2} + n$ ,  $n \ge 0$ .

**Theorem 3.13.** ([Dong-Nagatomo 1999]) Let L be a positive definite even lattice, and let  $V_L$  denote the vertex operator algebra attached to L. Then, the automorphism group of  $V_L$  is a complex Lie group of the form GO, where:

- (1) G is the connected component of identity, which is in particular a normal subgroup. G has a natural structure of a complex Lie group whose Lie algebra is isomorphic to the weight 1 subspace of  $V_L$ , with Lie bracket  $[u, v] = u_0 v$ . That is, G is generated by  $e^{h_0}$  for h in the weight 1 subspace.
- (2) O is the finite group of orthogonal transformations of a double cover  $\pi : \hat{L} \to L$ . The precise structure of the double cover is uniquely determined by the commutator relation  $[a,b] = (-1)^{(\pi(a),\pi(b))}$  for all  $a, b \in \hat{L}$ . The group O lies in an exact sequence  $1 \to \text{Hom}(L,\pm 1) \to O \to \text{Aut } L \to 1$ .
- (3) The intersection of G and O contains a copy of  $\text{Hom}(L,\pm 1)$  lying in the distinguished maximal torus (generated by  $\{e^{h_0}, h \in \pi_0\}$ ) of G.

As we mentioned earlier, our main examples for the lattice L are the following positive definite even unimodular lattices of rank 24:

- (1) The lattice  $N(A_1^{24})$  with root system  $A_1^{24}$ .
- (2) The Leech lattice  $\Lambda$ , with no roots.

**Theorem 3.14.** The automorphism group of  $V_{\Lambda}$  has the form  $(\mathbb{C}^{\times})^{24}.Co_0$ .

Proof. This follows from Theorem 3.13 as follows: the connected component of identity G is infinitesimally generated by the weight 1 space, which is the rank 24 abelian Lie algebra. G therefore acts as scalars on the  $\Lambda$ -graded Heisenberg modules by the identification  $\operatorname{Hom}(\Lambda, \mathbb{C}^{\times}) \cong (\mathbb{C}^{\times})^{24}$ . The subgroup O is a  $2^{24}$ -cover of  $\operatorname{Aut} \Lambda = Co_0$ , and  $G \cap O$  is precisely  $\operatorname{Hom}(\Lambda, \pm 1)$ , which is identified with the  $2^{24}$  subgroup. Thus,  $(\operatorname{Aut} V_{\Lambda})/G \cong Co_0$ .

**Theorem 3.15.** The automorphism group of the vertex algebra of  $N(A_1^{24})$  has the form GO, where G is the quotient of  $(SL_2(\mathbb{C}))^{24}$  by a copy of the Golay code in the center  $(\pm I_2)^{24}$ , and O has the form  $2^{24} \cdot 2^{24} \cdot M_{24}$ . The copy of the Golay code is precisely the one whose codewords describe the cosets of  $(\sqrt{2\mathbb{Z}})^{24}$  in  $N(A_1^{24})$ .

*Proof.* Because the root system of  $N(A_1^{24})$  is  $A_1^{24}$ , the Lie algebra of G is  $\mathfrak{sl}_2^{24}$ . Thus, the connected component of identity is a quotient of  $SL_2(\mathbb{C})^{24}$  by a subgroup of its center. Given  $u = (u^1, \ldots, u^{24}) \in (\frac{1}{\sqrt{2}}\mathbb{Z})^{24}$ , a torus element  $t = (t_1, \ldots, t_{24}) \in (\mathbb{C}^{\times})^{24}$  acts on  $\pi_u$  by the

Given  $u = (u^1, \ldots, u^{24}) \in (\frac{1}{\sqrt{2}}\mathbb{Z})^{24}$ , a torus element  $t = (t_1, \ldots, t_{24}) \in (\mathbb{C}^{\times})^{24}$  acts on  $\pi_u$  by the scalar  $\prod_{a=1}^{24} t_a^{\sqrt{2}u^a}$ . If we specify  $t_i \in \pm 1$ , we find that the scalar is uniformly identity if and only if we have an even number of -1s for each Golay codeword. Equivalently, the subset of -1 indices in  $\{1, \ldots, 24\}$  is orthogonal to  $\mathcal{C}$ . Because  $\mathcal{C}$  is self-orthogonal, it is necessary and sufficient that this subset is an element of  $\mathcal{C}$  itself. This proves the claim about G.

O is the orthogonal group of the double cover of  $N(A_1^{24})$ , hence it lies in an exact sequence

$$1 \to (\pm 1)^{24} \to O \to \operatorname{Aut} N(A_1^{24}) \to 1$$

This proves the claim about O.

**Definition 3.16.** Given an automorphism  $\sigma$  of a lattice L, we say an automorphism  $\tilde{\sigma}$  of  $V_L$  is a lift of  $\sigma$  if  $\tilde{\sigma}$  sends vectors in  $\pi_u$  to  $\pi_{\sigma(u)}$  for all  $u \in L$ .

In general the obstruction to an automorphism g of order n of a lattice admitting an order n lift is n being even and  $(v, g^{n/2}v)$  being odd for some lattice vector v [Borcherds 1992]. In particular, this obstruction vanishes for the -1 automorphism of an even lattice, so -1 always admits an order 2 lift.

3.3. The diagonal octahedral group. We introduce a particular subgroup of automorphisms of  $V_{N(A_{2}^{24})}$  isomorphic to  $S_4$ , and most importantly, a pair of conjugate involutions.

**Definition 3.17.** The **binary octahedral group** 2*O* is the following group of 48 unit quaternions:

$$2O = \{\pm x, \frac{\pm 1 \pm i \pm j \pm k}{2}, \frac{\pm x \pm y}{\sqrt{2}} | x, y \in \{1, i, j, k\}, x \neq y\}.$$

We also view this group as a subgroup of  $SL_2(\mathbb{C})$  by fixing once and for all the linear representation  $i \mapsto \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, j \mapsto \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, k \mapsto \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$ 

The binary octahedral group is the normalizer of the subgroup  $Q_8 = \{\pm 1, \pm i, \pm j, \pm k\}$  in the unit quaternions. A few short calculations yield the following:

**Lemma 3.18.** Under the adjoint representation of  $SL_2(\mathbb{C})$  on its Lie algebra, the quaternion *i* acts as identity on diagonal matrices and -1 on the span of  $E = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$  and  $F = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}$ . The quaternion *j* acts as -1 on diagonal matrices, and exchanges *E* with *F*. The images of the quaternions *i*, *j*, *k* in Aut  $\mathfrak{sl}_2$  have order 2, and are the non-identity elements in a group  $V_4$  of order 4. The image of the binary octahedral group is isomorphic to  $S_4$ , which naturally permutes the elements of  $V_4$ , and yields an identification with the holomorph  $V_4 \rtimes \operatorname{Aut} V_4$ .

We may alternatively introduce 2*O* by first considering the 4-group in  $SO_3(\mathbb{R})$  generated by half-rotations about the *x*-, *y*-, and *z*-axes, defining the octahedral group as its normalizer (or just explicitly as symmetries of an octahedron or cube), and lifting to the double cover SU(2).

**Definition 3.19.** We write  $\rho$  to denote the representation of the binary octahedral group on the vertex operator algebra  $V_{N(A_1^{24})}$  by the composite  $2O \to SL_2(\mathbb{C}) \to SL_2(\mathbb{C})^{24} \to \operatorname{Aut} V_{N(A_1^{24})}$ , where the second map is the diagonal embedding.

**Lemma 3.20.** The image of the quaternion -1 under  $\rho$  is identity on  $V_{N(A_1^{24})}$ . In particular, the action of the binary octahedral group factors through the quotient  $S_4$ , and the elements  $\rho(i), \rho(j), \rho(k)$  are the mutually conjugate non-identity elements in the normal 4-subgroup of this  $S_4$ .

Proof. The quaternion -1 acts by  $(-I_2, \ldots, -I_2) \in SL_2(\mathbb{C})^{24}$ , which lies in the distinguished torus, so it acts by scalars on each  $\pi_u$  for  $u \in N(A_1^{24})$ . For any  $u = (u^1, \ldots, u^{24}) \in (\frac{1}{\sqrt{2}}\mathbb{Z})^{24}$  the scalar is  $\prod_{a=1}^{24} (-1)^{\sqrt{2}u^a}$ . For  $u \in N(A_1^{24})$ , this scalar is 1 by the evenness of the Golay code. The last claim follows immediately from the previous lemma.

**Lemma 3.21.** The quaternion j acts on  $V_{N(A_1^{24})}$  as a lift of the -1 automorphism on  $N(A_1^{24})$ . That is, j stabilizes  $\pi_0$ , and restricts to isomorphisms  $\pi_u \to \pi_{-u}$  for each  $u \in N(A_1^{24})$ . The quaternion i acts as identity on the vertex subalgebra  $V_{\Lambda_0}$  and acts as -1 on  $\pi_u$  for  $u \in N(A_1^{24}) \setminus \Lambda_0$ . In particular, the fixed point subalgebra  $V_{N(A_1^{24})}^j$  is equal to  $V_{\Lambda_0}$ .

Proof. By Lemma 3.18, *i* acts as identity on the Heisenberg subalgebra, and *j* acts as -1. The claim about *j* then follows from this property of the action. Now, recall the action of *i* is given by  $\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \ldots, \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$ , which lies in the torus so for any lattice vector  $u = (u^1, \ldots, u^{24}) \in (\frac{1}{\sqrt{2}}\mathbb{Z})^{24}$ ,  $\rho(i)$  acts as the scalar  $\prod_{a=1}^{24} i^{\sqrt{2}u^a}$  on the corresponding module  $\pi_u$ . The map  $N(A_1^{24}) \to \pm 1$  given by  $u \mapsto \prod_{a=1}^{24} i^{\sqrt{2}u^a}$  has kernel equal to  $\Lambda_0$ .

**Lemma 3.22.** The fixed point vertex subalgebra  $V_{N(A_1^{24})}^{\rho(Q_8)}$  for the action of the 4-group  $\rho(Q_8) = \langle \rho(i), \rho(j) \rangle$  inherits an action of  $S_3$  from  $V_{N(A_1^{24})}$ .

*Proof.* The action of  $\rho(2O)$  on the fixed points for  $\rho(Q_8)$  is clearly trivial when restricted to the normal subgroup  $\rho(Q_8)$ , so the action factors through the quotient. Since  $\rho(2O)$  admits a section of the quotient map, the action of this section is transported to the fixed point subalgebra.

3.4. **Modules.** Vertex algebras have a notion of module that is analogous to the situation with commutative rings.

**Definition 3.23.** Let V be a vertex algebra. A V-module is a complex vector space M together with a map  $V \otimes M \to M((z))$ , written  $u \otimes w \mapsto Y_M(u, z)w = \sum_{n \in \mathbb{Z}} u_n w z^{-n-1}$  satisfying the following conditions:

- (1) (unit)  $Y_M(\mathbf{1}, z)w = w = wz^0$  for all  $w \in W$ .
- (2) (weak associativity) For any  $u, v \in V$  and  $m \in M$ , there is some N such that  $(z + w)^N Y_M(u, z + w) Y_M(v, w) m = Y_M(Y(u, z)v, w) m$ .

We note that we get an equivalent definition if we replace the second condition with the following: For any  $u, v \in V$  and  $m \in M$ , the operator-valued power series  $Y_M(u, z)Y_M(v, w)m \in M((z))((w))$ ,  $Y_M(v, w)Y_M(u, z)m \in M((w))((z))$  and  $Y_M(Y(u, z-w)v, w)m \in M((w))((z-w))$  are expansions of a common element of  $M[[z, w]][z^{-1}, w^{-1}, (z-w)^{-1}]$ , i.e., there is some N > 0 such that multiplying these three power series by  $z^N w^N (z-w)^N$  yields equal series in M[[z, w]].

**Theorem 3.24.** Let  $\mathfrak{h}$  be a complex vector space with nondegenerate inner product. Then, any irreducible  $\pi_0$ -module has the form  $\pi_u$  for some  $u \in \mathfrak{h}$ , where  $Y_{\pi_u}(e^i t^{-1}, z) = \sum_{n \in \mathbb{Z}} e^i t^n z^{-n-1}$ .

WHY DO THE SYMMETRIES OF THE MONSTER VERTEX ALGEBRA FORM A FINITE SIMPLE GROUP? 11

While it is easy to understand the irreducible  $\pi_0$ -modules, the category of all  $\pi_0$ -modules has a complicated extension structure. In particular, the finitely generated  $\pi_0$ -modules that admit a  $\mathbb{Z}_{\geq 0}$ -grading are in natural bijection with finite length Sym  $\mathfrak{h}$ -modules, or equivalently, finite length coherent sheaves on  $\mathfrak{h}$ . To avoid these pathologies, we introduce complete reducibility hypotheses.

**Definition 3.25.** Let V be a vertex operator algebra. We say V is strongly rational if the following conditions are satisfied:

- (1) The operator  $L_0$  on V has non-negative eigenvalues, and the kernel of  $L_0$  is spanned by **1** this is sometimes called the "CFT type" condition.
- (2) Any V-module is a direct sum of irreducible V-modules.

We say V is **strongly holomorphic** if V is not only strongly rational, but also any irreducible V-module is isomorphic to V. That is, the representation category of a strongly holomorphic vertex operator algebra is equivalent to the category of complex vector spaces.

The main examples we consider are lattices:

**Theorem 3.26.** ([Dong 1993]) Let L be a positive definite even lattice, and let  $L^{\vee}$  be the dual lattice. Then,  $V_L$  is strongly rational, and moreover, the category of  $V_L$ -modules is equivalent to the category of  $L^{\vee}/L$ -graded vector spaces. The irreducible  $V_L$ -modules have the form  $V_{L+\gamma} = \bigoplus_{u \in L+\gamma} \pi_u$  for  $\gamma \in L^{\vee}$ . In particular,  $V_L$  is strongly holomorphic if and only if L is self-dual.

An important case we consider is fixed points of holomorphic vertex operator algebras under actions of finite groups. In the following theorem, the assumption that G is solvable is conjectured to be superfluous, and by [McRae 2019], it can be removed when  $V^G$  satisfies the " $C_2$ -cofiniteness" condition.

**Theorem 3.27.** Let V be a strongly holomorphic vertex operator algebra, and let G be a finite solvable group of automorphisms. Then:

- (1) The fixed point subalgebra  $V^G$  is strongly rational [Carnahan-Miyamoto 2016].
- (2) The representation category of  $V^G$  is equivalent to the category of  $D^{\alpha}(\mathbb{C}G)$ -modules, for some  $\alpha \in H^3(G, \mathbb{C}^{\times})$ , where  $D^{\alpha}(\mathbb{C}G)$  is the  $\alpha$ -twisted Drinfeld double of the group ring  $\mathbb{C}G$ [Dong-Ng-Ren 2021].

For the cases of  $V_{\Lambda}$  and  $V_{N(A_1^{24})}$  and certain involutions, we will describe the modules more concretely later. The following theorem seems to be implicit in [Frenkel-Lepowsky-Meurman 1988], but is not explicitly stated there.

**Theorem 3.28.** The fixed point vertex operator subalgebra  $V^{\theta}_{\Lambda}$  is generated by its weight 2 subspace.

Proof. We first consider the vertex operator subalgebra  $\pi_0^{\theta}$ , given by  $\theta$ -fixed vectors in the Heisenberg vertex operator subalgebra of  $V_{\Lambda}$ . By [Dong-Nagatomo 1999], such a fixed-point subalgebra is generated by its weight 2 subspace (when  $\mathfrak{h}$  has dimension at least 2, which holds for  $\mathfrak{h} = \Lambda \otimes \mathbb{C}$ ). Then,  $V_{\Lambda}^{\theta}$  decomposes into irreducible  $\pi_0^{\theta}$ -modules as  $\pi_0^{\theta} \oplus \bigoplus_{[\lambda] \in \Lambda/\pm 1} (\pi_{\lambda} \oplus \pi_{-\lambda})^{\theta}$ . The lowest-weight vectors of  $(\pi_{\lambda} \oplus \pi_{-\lambda})^{\theta}$  have the form  $v_{[\lambda]} = v_{\lambda} + \theta(v_{\lambda})$  with weight equal to half of the norm of  $\lambda$ , so the weight 2 space is spanned by these vectors for  $\lambda$  of norm 4. Thus, it suffices to show that for any  $\mu \in \Lambda$ , some scalar multiple of  $v_{[\mu]}$  can be produced as a coefficient of products of the form

$$Y(v_{[\lambda_1]}, z_1)Y(v_{[\lambda_2]}, z_2)\cdots Y(v_{[\lambda_{n-1}]}, z_{n-1})v_{[\lambda_n]}$$

for  $\lambda_1, \ldots, \lambda_n \in \Lambda$  of norm 4, and this follows (after some calculation) from the fact that  $\Lambda$  is generated by norm 4 vectors.

Vertex algebras admit a different notion of module that doesn't appear in the usual study of associative algebras:

**Definition 3.29.** Let V be a vertex algebra, and let g be an automorphism of finite order k. A g-twisted V-module is a vector space M equipped with a linear map  $V \otimes M \to M((z^{1/k}))$ , written  $u \otimes w \mapsto Y_M(u, z)w = \sum_{n \in \frac{1}{L}\mathbb{Z}} u_n w z^{-n-1}$  satisfying the following conditions:

- (1)  $Y_M(\mathbf{1}, z)w = w = wz^0$  for all  $w \in W$ .
- (2) For any  $u, v \in V$  and  $m \in M$ , the operator-valued power series  $Y_M(u, z)Y_M(v, w)m \in M((z^{1/k}))((w^{1/k})), Y_M(v, w)Y_M(u, z)m \in M((w^{1/k}))((z^{1/k}))$  and  $Y_M(Y(u, z w)v, w)m \in M((w^{1/k}))((z w))$  are expansions of a common element of  $M[[z^{1/k}, w^{1/k}]][z^{-1}, w^{-1}, (z w)^{-1}]$ , i.e., there is some N > 0 such that multiplying these three power series by  $z^N w^N(z w)^N$  yields equal series in  $M[[z^{1/k}, w^{1/k}]]$ .
- (3) If  $a \in V$  satisfies  $ga = e^{2\pi i k/n} a$  for some  $k \in \mathbb{Z}$ , then  $Y_M(a, z) \in z^{k/n}(\operatorname{End} M)[[z, z^{-1}]].$

In conformal field theory, modules are called "sectors", and are naturally attached to points on a Riemann surface. One may then compute quantities like correlation functions by choosing vectors in the modules, and applying the global geometry of the curve. In order to consider twisted modules, called "twisted sectors", one considers a Galois cover of Riemann surfaces, and attaches twisted modules to ramified points, where the monodromy matches the twisting.

The main result about twisted modules that we need is:

**Theorem 3.30.** ([Dong-Li-Mason 1997]) Let V be a strongly holomorphic vertex operator algebra, and let g be a finite order automorphism of V. Let  $T_g(\tau) = \sum_n \operatorname{Tr}(g|V_n)q^{n-c/24}$  be the graded trace of g on V, where we define  $q = e^{2\pi i \tau}$  for  $\tau$  on the complex upper half-plane. Then, there exists a unique irreducible g-twisted V-module V(g) up to isomorphism, and its character  $\sum \dim V(g)_n q^{n-c/24}$  is equal to  $T_g(-1/\tau)$  as a holomorphic function on the complex upper half-plane.

We can say more in the particular case of the -1-automorphism of positive definite even modular lattices:

**Theorem 3.31.** ([Frenkel-Lepowsky-Meurman 1988]) Let  $\theta$  be an order 2 lift of -1 on a positive definite even unimodular lattice L, and let  $\hat{L}$  be the unique central extension of L by  $\pm 1$ satisfying  $[\hat{a}, \hat{b}] = (-1)^{(a,b)}$  for any lifts  $\hat{a}, \hat{b} \in \hat{L}$  of vectors  $a, b \in L$ . Then,  $V_L(\theta)$  has the form  $T \otimes \text{Sym}(t^{-1/2}\mathfrak{h}[t^{-1}])$ , where  $\mathfrak{h} = L \otimes \mathbb{C}$ , and T is the irreducible  $\hat{L}/2L$ -module of rank  $2^{(\dim \mathfrak{h})/2}$ .

Applying this to our examples, we have the following:

**Corollary 3.32.** The graded trace of  $\rho(i)$  on  $V_{N(A_1^{24})}$  is  $\frac{2\theta_{L_0}(\tau) - \theta_{N(A_1^{24})}(\tau)}{\eta(\tau)^{24}} = \frac{\eta(\tau)^{24}}{\eta(2\tau)^{24}} = q^{-1} - 24 + 276q - 2048q^2 + \cdots$ . The irreducible  $\rho(i)$ -twisted  $V_{N(A_1^{24})}$ -module has character  $2^{12} \frac{\eta(\tau)^{24}}{\eta(\tau/2)^{24}} = 4096q^{1/2} + 98304q + \cdots$ . The same formulas hold if we replace  $\rho(i)$  with  $\rho(j)$ . For any order 2 lift  $\theta$  of the -1 automorphism of  $\Lambda$  to  $V_{\Lambda}$ , the graded trace of  $\theta$  is  $\frac{\eta(\tau)^{24}}{\eta(2\tau)^{24}}$ , i.e., the same as the graded trace of  $\rho(i)$ . The irreducible  $\theta$ -twisted  $V_{\Lambda}$  module has character  $2^{12} \frac{\eta(\tau)^{24}}{\eta(\tau/2)^{24}}$ , i.e., the same as the same as  $V_{N(A_1^{24})}(\rho(i))$ . In each case, the twisted module decomposes into the direct sum of 2 irreducible modules for the fixed-point vertex subalgebra, and these submodules are the +1 and -1 eigenspaces for the action of  $e^{2\pi i L_0}$ .

3.5. Holomorphic orbifolds. The monster vertex algebra was originally constructed using the following method:

- (1) Start with the Leech lattice vertex operator algebra  $V_{\Lambda}$ , which is strongly holomorphic (Theorem 3.26).
- (2) Take the vertex subalgebra of fixed points under an order 2 automorphism  $\theta$ .
- (3) Adjoin an irreducible module for the fixed-point subalgebra.

More generally, we may take fixed points for any group of automorphisms, and try to attach modules for the fixed point subalgebra. For finite groups of automorphisms, this is called the "orbifold construction", since in conformal field theory, it is conjectured to produce the chiral algebra attached to a quotient orbifold.

Recent progress in the representation theory of fixed points has yielded the following:

**Theorem 3.33.** ([Evans-Gannon 2018]) Let V be a strongly holomorphic vertex operator algebra, and let G be a finite solvable group of automorphisms. Then strongly holomorphic extensions of  $V^G$  are parametrized by pairs  $(K, \psi)$ , where K is a subgroup of G on which the restriction of  $\omega$  is a coboundary, and  $\psi \in Z^2(K, \mathbb{C}^{\times})$ . In particular, V corresponds to the pair ({e}, 1).

Similar to the situation with the properties of fixed points that we mentioned earlier, the assumption that G is solvable is conjecturally superfluous. The subgroup K parametrizes the set of twisted modules whose  $V^{G}$ -submodules appear in the extension.

**Definition 3.34.** Let V be a strongly holomorphic vertex operator algebra, and let G be a finite group of automorphisms. The **orbifold dual** of V with respect to G, written V//G, is the strongly holomorphic extension of  $V^G$  attached to the pair (G, 1), if it exists. If G is a cyclic group generated by an element  $g \in G$ , we write V//g instead.

In the case where G is a cyclic group generated by an element g, we can describe the orbifold dual more concretely.

**Theorem 3.35.** ([van Ekeren-Möller-Scheithauer 2015]) Let V be a strongly holomorphic vertex operator algebra, and let g be a finite order automorphism of V. The orbifold dual V//g exists if and only if the irreducible g-twisted module V(g) has  $L_0$ -spectrum in  $\frac{1}{|g|}\mathbb{Z}$ . If this condition holds, then each irreducible twisted V-module  $V(g^i)$  admits a faithful action of  $\mathbb{Z}/|g|\mathbb{Z}$  satisfying the following properties:

- (1) The decomposition of  $V(g^i)$  into eigenspaces for this action is a decomposition into |g| irreducible modules for the fixed-point subalgebra  $V^g$ .
- (2) The action on V(g) is given by  $a \mapsto e^{2\pi i a L_0}$ , meaning in particular that the fixed-point subspace is precisely the subspace on which  $L_0$  acts with integral eigenvalues.
- (3) V//g decomposes as a  $V^g$ -module into the direct sum of fixed points  $\bigoplus_{i=0}^{|g|-1} V(g^i)^{\mathbb{Z}/|g|\mathbb{Z}}$ .

Furthermore, there is an action of a cyclic group  $\langle g^* \rangle$  of order |g| on V//g, such that  $g^*$  acts on the subspace  $V(g^j)^{\mathbb{Z}/|g|\mathbb{Z}}$  by the scalar  $e^{2\pi i j/|g|}$ , and  $(V//g)//g^* \cong V$ .

**Lemma 3.36.** The three cyclic orbifolds  $V_{N(A_1^{24})}//\rho(i)$ ,  $V_{N(A_1^{24})}//\rho(j)$ , and  $V_{N(A_1^{24})}//\rho(k)$  exist and are isomorphic to  $V_{\Lambda}$ .

Proof. By Lemma 3.20, the automorphisms  $\rho(i), \rho(j), \rho(k)$  are conjugate in  $\operatorname{Aut}(V_{N(A_1^{24})})$ , so it suffices to prove the claim for  $V_{N(A_1^{24})}//\rho(i)$ . By Lemma 3.21, the fixed point vertex subalgebra is  $V_{\Lambda_0}$ , so by Theorem 3.26 (from [Dong 1993]) irreducible modules are the sums  $M_{\gamma} = \bigoplus_{u \in \Lambda_0 + \gamma} \pi_u$ for  $\gamma \in \Lambda_0^{\vee}/\Lambda_0$ . Thus, the strongly holomorphic extensions of  $V_{\Lambda_0}$  are precisely the direct sums  $V_{\Lambda_0} \oplus M_{\gamma}$  for  $\gamma \neq 0$  such that  $M_{\gamma}$  has integer weight, yielding a lattice vertex algebra. One choice of  $\gamma$  gives non-integer weight, one yields  $V_{N(A_1^{24})}$ , and the third yields  $V_{\Lambda}$ .

**Lemma 3.37.** The identification of  $V_{\Lambda}$  with  $V_{N(A_1^{24})}//\rho(i)$  induces a natural action of a 4-group, generated by a lift of the involution  $\rho(j)$  on  $V_{N(A_1^{24})}^{\rho(i)}$ , and an involution  $\rho(i)^*$  whose fixed point subalgebra is  $V_{N(A_1^{24})}^{\rho(i)}$ .

*Proof.* The fact that  $\rho(i)^*$  is order 2 with specified fixed points is given in Theorem 3.35. The involutions  $\rho(j)$  and  $\rho(k)$  coincide on this subspace, and commute with the restriction of  $\rho(i)^*$ , since it is the identity. There are 2 possible lifts of  $\rho(j)$  (seen as an order 2 lift of -1 on  $\Lambda_0$  to  $V_{\Lambda_0}$ )

to automorphisms of  $V_{\Lambda}$ , uniquely determined by where they take a lowest weight vector  $v_{\lambda} \in \pi_{\lambda}$  for  $\lambda \in \Lambda \setminus \Lambda_0$ . The image is necessarily a scalar times  $v_{-\lambda}$ , and the two choices differ by a sign, i.e., composition with  $\rho(i)^*$ . Both lifts have order 2, so we obtain a 4-group.

The following is a special case of the main result of [Shimakura 2003].

**Lemma 3.38.** Let  $\theta$  be an order 2 lift of the -1 automorphism of  $\Lambda$  to  $V_{\Lambda}$ . Then, the automorphism group of  $V^{\theta}_{\Lambda}$  has the form  $2^{24}.Co_1$ .

Proof. Any automorphism of a vertex operator algebra induces a permutation of the irreducible modules, and in this case,  $V_{\Lambda}^{\theta}$  has 4 irreducible modules, with pairwise distinct characters. The pairwise-distinct property implies for any automorphism  $\sigma$  of  $V_{\Lambda}^{\theta}$ , there exists a lift to an automorphism of  $V_{\Lambda}$ . We conclude that the automorphism group of  $V_{\Lambda}^{\theta}$  is equal to the quotient  $C_{\text{Aut }V_{\Lambda}}(\theta)/\langle\theta\rangle$ . Because  $\theta$  acts as -1 on the tangent space of the torus, the centralizer has torus part  $(\pm 1)^{24}$ , and because  $\theta$  is a lift of the central element -1 in  $Co_0$ , the quotient of the centralizer by the torus part is  $Co_0$ . Thus,  $\text{Aut }V_{\Lambda}^{\theta}$  has the form  $2^{24}.Co_0/\langle\theta\rangle$ , giving the form  $2^{24}.Co_1$ .

4. The monster vertex algebra

**Definition 4.1.** The monster vertex algebra  $V^{\natural}$  is the cyclic order 2 orbifold:

$$V^{\natural} = V_{\Lambda} / / \theta,$$

where  $\theta$  is an order 2 lift of the -1 automorphism of  $\Lambda$ . That is,  $V^{\natural}$  decomposes as the direct sum of the simple vertex operator algebra  $V_{\Lambda}^{\theta}$  and its irreducible module  $V_{\Lambda}(\theta)^{\theta}$ .

We note some basic facts.

**Lemma 4.2.** (1)  $V^{\natural}$  is a holomorphic vertex operator algebra.

(2)  $V^{\natural}$  has character  $J(\tau) = q^{-1} + 196884q + \cdots$ . In particular, the weight 1 subspace is zero. (3)  $V^{\natural}$  has an order 2 automorphism z whose fixed points are identified with  $V^{\theta}_{\Lambda}$ .

*Proof.* For the claim about the character, it suffices to work out the characters of the subspaces  $V_{\Lambda}^{\theta}$  and  $V_{\Lambda}(\theta)^{\theta}$ . These can be computed from Corollary 3.32. The order 2 automorphism z is precisely  $\theta^*$  as defined in Theorem 3.35.

From Lemma 3.36, we see that  $V^{\natural}$  can be constructed from  $V_{N(A_1^{24})}$  as a composite of 2 orbifold duals in 3 different ways: We start by taking the orbifold dual with respect to one of the three involutions  $\rho(i)$ ,  $\rho(j)$ ,  $\rho(k)$  to get  $V_{\Lambda}$ , then taking the orbifold dual with respect to a lift of one of the other involutions. This information can be organized in the following way:

**Theorem 4.3.** (Triality) We have the following diagram of orbifold duals:



where the vertex operator algebras in the middle row are isomorphic to  $V_{\Lambda}$ , the vertex operator algebra on the bottom is isomorphic to  $V^{\natural}$ , and x, y, z are commuting order 2 automorphisms of  $V^{\natural}$  that generate a 4-group, which splits  $V^{\natural}$  into a direct sum of 4 irreducible  $V_{N(A_{1}^{24})}{}^{\rho(Q_{8})}$ -modules  $V^{00}, V^{01}, V^{10}, V^{11}$ , satisfying the following properties:

- (1)  $V^{01}$  is an irreducible  $V_{N(A_1^{24})}^{\rho(Q_8)}$ -submodule of  $V_{N(A_1^{24})}(\rho(i))$ , on which z acts by 1 and x, y act by -1. It is contained in  $V_{N(A_1^{24})}//\rho(i)$ .
- (2)  $V^{10}$  is an irreducible  $V_{N(A_1^{24})}^{\rho(Q_8)}$ -submodule of  $V_{N(A_1^{24})}(\rho(j))$ , on which x acts by 1 and y, z act by -1. It is contained in  $V_{N(A_1^{24})}//\rho(j)$ .
- (3)  $V^{11}$  is an irreducible  $V_{N(A_1^{24})}^{\rho(Q_8)}$ -submodule of  $V_{N(A_1^{24})}(\rho(k))$ , on which y acts by 1 and x, z act by -1. It is contained in  $V_{N(A_1^{24})}//\rho(k)$ .
- (4) The involutions x\*, y\*, z\* are lifts of the -1 automorphism on Λ, for some identification of the corresponding orbifolds of V<sub>N(A<sup>24</sup><sub>1</sub>)</sub> with V<sub>Λ</sub>. In particular, the restriction of z\* to V<sub>Λ0</sub> is given by ρ(j), a lift of the -1 automorphism of the lattice.

Furthermore, the action of  $S_3 \subset \operatorname{Aut} V_{N(A_1^{24})}$  permuting  $\{\rho(i), \rho(j), \rho(k)\}$  induces a faithful permutation action on  $\{V^{01}, V^{10}, V^{11}\}$ , and extends to the conjugation action of a subgroup  $\Sigma \subset \operatorname{Aut} V^{\natural}$  that induces a faithful permutation action of  $S_3$  on  $\{x, y, z\}$ .

One can show that the group  $\Sigma$ , given by lifting the  $S_3$  acting on the vertex operator subalgebra  $V_{N(A_1^{24})}{}^{\rho(Q_8)}$ , can be chosen isomorphic to  $S_3$ , and together with the 4-group generates a copy of  $S_4$ . Indeed, in [Frenkel-Lepowsky-Meurman 1988] a pair of generating elements of  $S_3 \subset \text{Aut } V^{\natural}$  was explicitly constructed.

**Corollary 4.4.**  $V^{\ddagger}$  is generated by its degree 2 subspace.

Proof. By Theorem 4.3,  $V^{\natural}$  decomposes into 4 irreducible  $V_{N(A_1^{24})}{}^{\rho(Q_8)}$ -modules  $V^{00}, V^{01}, V^{10}, V^{11}$ , where  $V^{00} \oplus V^{01}$  is identified with  $V^{\theta}_{\Lambda}$ , and there is a group  $\Sigma \subset \operatorname{Aut} V^{\natural}$  whose conjugation action induces a faithful  $S_3$ -action on  $\{V^{01}, V^{10}, V^{11}\}$ . Thus, it suffices to show that  $V^{\theta}_{\Lambda}$  is generated by its degree 2 subspace, and this is precisely Theorem 3.28.

4.1. Extensions of Conway groups. We now consider the automorphisms of  $V^{\ddagger}$  that arise directly from  $V_{\Lambda}$ .

**Theorem 4.5.** We have the following diagram of finite groups where arrows are double covers:



and the groups have the following form:

- (1)  $C_0$  is the centralizer of  $\theta$  in Aut  $V_{\Lambda}$ . It has the form  $2^{24}.Co_0$ , where  $2^{24}$  naturally identifies with  $\Lambda/2\Lambda$ . This group acts faithfully on  $V_{\Lambda}$ .
- (2)  $C_1$  is the quotient of  $C_1$  by the central subgroup  $\langle \theta \rangle$ , and has the form  $2^{24}.Co_1$ . By Lemma 3.38,  $C_1$  is the full automorphism group of  $V_{\Lambda}^{\theta}$ .
- (3)  $C_T$  is defined using the projective action of  $C_1$  on the twisted module  $V_{\Lambda}(\theta)$ . The projective action on the lowest weight space T (of dimension  $2^{12}$ ) induces an injective homomorphism  $C_1 \rightarrow PGL(T)$ , and the pullback  $C_* \subset GL(T)$  has commutator subgroup  $C_T$ , which has the form  $2.2^{24}.Co_1$ . This group acts faithfully on the vertex superalgebra  $V_{\Lambda}^{\theta} \oplus V_{\Lambda}(\theta)^{\theta=-1}$ .

- (4)  $\hat{C}$  is the pullback  $C_0 \times_{C_1} C_T$ . It has the form  $2^2 \cdot 2^{24} \cdot Co_1$ , and acts faithfully on the "generalized vertex algebra"  $V_{\Lambda} \oplus V_{\Lambda}(\theta)$  constructed in [Huang 1994].  $C_0$  and  $C_T$  are the quotients of  $\hat{C}$  by two of the involutions in the central 4-group  $2^2$ .
- (5) C is the quotient of Ĉ by the remaining involution in 2<sup>2</sup>. C has the form 2<sup>1+24</sup>.Co<sub>1</sub>, i.e., there is an extraspecial normal subgroup Q, naturally identified with Â/2Λ, and the quotient by this subgroup is Co<sub>1</sub>. C acts faithfully on V<sup>β</sup>, and it is the centralizer of the distinguished involution z ∈ Aut V<sup>β</sup> that is trivial on V<sup>β</sup><sub>Λ</sub>, and -1 on V<sub>Λ</sub>(θ)<sup>θ</sup>.

4.2. Finiteness. Our proof that  $\operatorname{Aut} V^{\natural}$  is a finite group is surprisingly easy - compare with the proof in [Tits 1984] or [Griess 1982].

**Theorem 4.6.** The group  $\operatorname{Aut} V^{\natural}$  of all vertex operator algebra automorphisms of  $V^{\natural}$  is finite.

*Proof.* By Corollary 4.4,  $V^{\natural}$  is generated by its degree 2 subspace. The condition that a linear transformation on a generating subspace induces an automorphism of the vertex operator algebra is given by polynomial identities on the coefficients of multiplication, so Aut  $V^{\natural}$  embeds as a Zariski closed subgroup of  $GL_{196884}(\mathbb{C})$ .

Because the centralizers of the mutually conjugate involutions x, y, z are finite (in particular, isomorphic to C), conjugation by any of these elements acts as -1 on the Lie algebra of Aut  $V^{\ddagger}$ , hence as inversion on the connected component of identity. However, the equation xy = z implies z also acts as identity on the Lie algebra, so the Lie algebra of Aut  $V^{\ddagger}$  is trivial. We conclude that Aut  $V^{\ddagger}$  is a zero dimensional linear algebraic group, hence finite.

We note that we don't really need the explicit work that goes into Corollary 4.4, because it suffices to show that  $V^{\natural}$  is finitely generated, and by Corollary 3.4 in [Li 1998], the finitely generated property holds for all  $C_1$ -cofinite vertex operator algebras. Here,  $C_1$ -cofiniteness is a condition that holds for a large class that includes all strongly rational vertex operator algebras like  $V^{\natural}$ .

4.3. Simplicity. We now show that  $\operatorname{Aut} V^{\natural}$  is simple. Since we have a large subgroup C which has relatively few normal subgroups, we will consider the possible intersections between C and a normal subgroup of  $\operatorname{Aut} V^{\natural}$ .

## **Lemma 4.7.** Let H be a normal subgroup of Aut $V^{\natural}$ . If $H \cap C$ is trivial, then H is trivial.

Proof. If  $H \cap C = \{1\}$ , then conjugation by z is an involution on H that has no non-identity fixed points. By a standard result (using injectiveness hence bijectiveness of the map  $a \mapsto (a^z)a^{-1}$  on H), this implies H is abelian of odd order and z acts by inversion. The same is true of x and y, since they are conjugate to z and H is normal. However, xy = z, so z also acts by identity, and H is necessarily trivial.

**Lemma 4.8.** If H is a normal subgroup of Aut  $V^{\natural}$ , then  $H \cap C \neq \langle z \rangle$ .

*Proof.* z is conjugate to x, and  $x \in C$ , so  $x \notin \langle z \rangle$  contradicts normality of H.

**Lemma 4.9.** If H is a normal subgroup of  $\operatorname{Aut} V^{\natural}$ , then  $H \cap C \neq Q$ .

*Proof.* It suffices to show that there is an element of Aut  $V^{\natural}$  that conjugates an element of Q to an element of  $C \setminus Q$ . For this, we consider the involution  $\rho(\frac{i+j}{\sqrt{2}}) \in \rho(2O)$  by which conjugation switches  $\rho(i)$  with  $\rho(j)$ . Then, there exists a lift to an automorphism  $\sigma$  of  $V^{\natural}$  for which conjugation switches z with x. We will consider a suitable element in Q that commutes with x, because conjugation by  $\sigma$  will then yield an element of C.

We start by considering automorphisms of  $V_{N(A_1^{24})}$  of the form  $u = (u^1, \ldots, u^{24}) \in SL_2(\mathbb{C})^{24}$ where entries  $u^1, \ldots, u^{24}$  lie in  $\begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$ . We may write u as  $e^{2\pi i h_0}$  for  $h = (h^1, \ldots, h^{24}) \in \mathbb{R}^{24} \subset$   $\pi_0$ , where  $h^a \in \frac{1}{2\sqrt{2}}\mathbb{Z}$  and  $u^a = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}^{2\sqrt{2}h^a}$ . Given  $v \in N(A_1^{24})$ , an automorphism u of this form acts on  $\pi_v$  by the scalar  $\prod_{a=1}^{24} i^{h^a v^a}$ , and it has order dividing 2 if and only if the set of a such that  $u^a \notin \pm I_2$  is a Golay codeword.

We may lift u to an automorphism of  $V_{N(A_1^{24})}//\rho(i)$  as  $e^{2\pi i h_0}$ , and it has order dividing 2 if and only if h lies in  $\frac{1}{2}\Lambda$ . Thus, we some choose  $h \in \frac{1}{2}\Lambda_0 \setminus (\Lambda \cup N(A_1^{24}))$ , and we choose some lift  $\tilde{u} \in Q \setminus \langle z \rangle$ . We note that  $\{a | u^a \notin \pm I_2\}$  is a Golay codeword with nonzero weight.

We claim that  $\tilde{u}$  commutes with x. Indeed, restricting x to  $(V^{\natural})^{z} = V_{\Lambda}^{\theta}$  yields the automorphism that is identity on  $(\pi_{u} + \pi_{-u})^{\theta}$  for  $u \in \Lambda_{0}$  and -1 on the remaining modules, so this restriction is also the restriction of  $e^{2\pi i (\frac{1}{4}\alpha_{C})_{0}}$  on  $V_{\Lambda}$ . That is, the image of x under the double cover  $C \to 2^{24}.Co_{1}$ lies in  $2^{24}$ , and is represented by  $e^{2\pi i s_{0}}$  where  $(2s, 2h) \in 2\mathbb{Z}$ . By the commutator rule for double covers of lattices, we see that  $e^{2\pi i s_{0}}$  and  $e^{2\pi i h_{0}}$  lift to commuting elements x and  $\tilde{u}$  of Q.

Conjugating  $e^{2\pi i h_0}$  by  $\rho(\frac{i+j}{\sqrt{2}})$  yields an element  $w = (w^1, \dots, w^{24}) \in SL_2(\mathbb{C})^{24}$  with entries given  $\begin{pmatrix} u^a & u^a = \pm I_2 \end{pmatrix}$ 

by  $w^a = \begin{cases} u^a & u^a = \pm I_2 \\ \pm \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} & u^a = \pm \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}.$  We see that w acts on  $\mathbb{R}^{24}$  by a composite of root

reflections parametrized by the *a* for which  $w^a \notin \pm I_2$ . That is, this is a composite of reflections parametrized by a Golay codeword of nonzero weight. In particular, *w* acts nontrivially on  $\pi_0^{\theta}$ , so *w* does not lift to an element of *Q*. On the other hand, the lift  $\sigma \tilde{u} \sigma^{-1}$  of *w* commutes with *z*, because  $\tilde{u}$  commutes with *x*, so  $\sigma \tilde{u} \sigma^{-1}$  lies in  $C \setminus Q$ .

**Lemma 4.10.** (from [Tits 1984]) The only normal subgroup of  $\operatorname{Aut} V^{\natural}$  containing C is  $\operatorname{Aut} V^{\natural}$ .

*Proof.* Let H be a normal subgroup of Aut  $V^{\natural}$  containing C, let S be a Sylow 2-subgroup of C, and let N be the normalizer of S in H. Since the center of S is  $\langle z \rangle$ , N centralizes z, hence lies in C.

We claim that S is a Sylow 2-subgroup of H. Indeed, otherwise there would be a 2-group in H strictly containing S but contained in N, contradicting Sylow's theorem. From this, the Frattini argument yields  $H = \operatorname{Aut} V^{\natural}$ . To elaborate, because H is normal, and all Sylow 2-subgroups of H are conjugate to each other, for any  $g \in \operatorname{Aut} V^{\natural}$ , there is some  $h \in H$  such that  $S^g = S^h$ , implying  $g \in hN \subseteq H$ .

**Theorem 4.11.** Aut  $V^{\natural}$  is simple.

*Proof.* If H is a normal subgroup of Aut  $V^{\natural}$ , then the intersection  $H \cap C$  is normal in C, so it must be one of  $\{1\}$ ,  $\langle z \rangle$ , Q, and C (here, we use the simplicity of  $Co_1$  and the nonexistence of a  $Co_1$ -stable proper subgroup of Q strictly containing  $\langle z \rangle$ ). By lemmata 4.7, 4.8, 4.9, 4.10, we find that H is either trivial or equal to Aut  $V^{\natural}$ .

## 5. What else can we derive about the monster from vertex-algebraic methods?

We've proved our main objective, that  $\operatorname{Aut} V^{\natural}$  is finite and simple, using modern methods in vertex algebras and mostly 19th century methods in finite groups. However, there is much more one might want to say. For example, one may want to compute the order of this group, and figure out some of its irreducible characters. For the monster, this information was computed even before the group was constructed, using group-theoretic methods that are well beyond the scope of this paper.

We will describe some facts that we can prove "easily modulo vertex algebras and some hard manipulations with complex functions".

**Theorem 5.1.** (Monstrous moonshine minus explicit computations) For any  $g \in \operatorname{Aut} V^{\natural}$ , the McKay-Thompson series

$$T_g(\tau) = \sum_{n \ge 0} \operatorname{Tr}(g|V_n^{\natural}) q^{n-1}$$

is a completely replicable function of order |g|, whose k-th replicate is  $T_{g^k}(\tau)$ . That is, for any  $k \geq 1$ , the function  $\sum_{ad=k,0\leq b< d} T_{g^a}(\frac{a\tau+b}{d})$  is equal to the unique polynomial of degree k in  $T_g(\tau)$  that has q-expansion of the form  $q^{-k} + O(q)$ . Furthermore, each  $T_g$  is a hauptmodul, i.e., there is a subgroup  $\Gamma_g \subset SL_2(\mathbb{R})$  containing  $\Gamma_0(N)$  for some N, such that the quotient  $\mathfrak{H}/\Gamma_g$  of the upper half-plane is a genus zero Riemann surface with finitely many punctures, and  $T_g$  generates its function field.

*Proof.* The first claim is proved in [Borcherds 1992] up to the end of section 8, before the explicit identification of McKay-Thompson series with the candidate modular functions presented in [Conway-Norton 1979]. The second claim follows from the combination of two theorems: the main theorem of [Martin 1994] asserts that completely replicable series that are "J-final" are invariant under  $\Gamma_0(N)$  for some N whose prime divisors are the same as those of |g|, and the main theorem of [Cummins and Gannon 1997] asserts that completely replicable functions are either Laurent polynomials in q or hauptmoduls.

We note that the completely replicable functions of finite order were completely classified in [Alexander et al. 1992]: there are 171 monstrous functions and 157 non-monstrous functions.

**Corollary 5.2.** The primes dividing the order of Aut  $V^{\natural}$  are a subset of the 15 "supersingular primes"  $\{2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 41, 47, 59, 71\}$ 

*Proof.* The completely replicable functions of prime order p are hauptmoduls of level a power of p, and this implies the modular curve  $X_0(p)^+$  is genus zero. The 15 primes satisfying this genus zero condition were classified in [Ogg 1974].

We can also constrain the "largeness" of Aut  $V^{\natural}$  by analyzing the McKay-Thompson series. Using cyclic orbifolds, in particular Tuite's orbifold duality correspondence between fixed-point free classes in  $Co_1$  satisfying a "no massless states" condition and non-Fricke classes in the monster (conjectured in [Tuite 1993] and proved in [Carnahan 2017]), we can also narrow down the possible McKay-Thompson series to a set of 174 functions (where the true set is 171) without explicit use of the monster or structure of  $V^{\natural}$ :

**Theorem 5.3.** ([Carnahan-Komuro-Urano 2017]) Let V be a strongly holomorphic vertex operator algebra with graded dimension J, and let g be a finite order automorphism of V. Then the graded trace of g on V is one of 174 completely replicable functions.

In the other direction, we can produce lower bounds on the size of Aut  $V^{\natural}$ . This is important, because in the paper where he classified the supersingular primes, Ogg asked why the set of primes dividing the order of the monster coincided with the set of primes p for which  $X_0(p)^+$  is genus zero, offering a bottle of Jack Daniels for a satisfactory answer. Borcherds's proof of the monstrous moonshine conjecture for  $V^{\natural}$  gives half of the answer, and the other half requires an explanation for why the monster is so large. We can produce, without appealing to much group theory, a lower bound on the order that is a factor of  $11 \cdot 17 \cdot 19$  off the true size:

**Theorem 5.4.** The order of Aut  $V^{\natural}$  is divisible by  $2^{46} \cdot 3^{20} \cdot 5^9 \cdot 7^6 \cdot 11 \cdot 13^3 \cdot 23 \cdot 29 \cdot 31 \cdot 41 \cdot 47 \cdot 59 \cdot 71$ .

*Proof.* We first consider the small primes:

(1) The  $2^{46}$  comes from the order of the subgroup C, and so do the divisors 3, 5, 7, 11, 13, 23.

- (2) The  $3^{20}$  comes from an order 3 cyclic orbifold construction of  $V^{\natural}$ , starting from a fixed-point free automorphism of  $\Lambda$  [Chen-Lam-Shimakura 2016]. Such an automorphism endows  $\Lambda$ with the structure of a lattice over the ring  $\mathbb{Z}[\frac{-1+\sqrt{-3}}{2}]$  of Eisenstein integers, and yields a construction of  $\Lambda$  starting from the tertiary Golay code and passing through the  $A_2^{12}$ Niemeier lattice. The automorphism group of  $\Lambda$  as an Eisenstein integer lattice has the form 6.Suz, where Suz is the Suzuki sporadic group. The centralizer of an order 3 lift in Aut  $V_{\Lambda}$  has the form  $3^{12}.6.Suz$ , and the cyclic orbifold construction of  $V^{\natural}$  then yields an order 3 automorphism whose centralizer has the form  $3^{1+12}.2.Suz$  with order a multiple of  $3^{20}$ .
- (3) The terms 5<sup>9</sup>, 7<sup>6</sup>, and 13<sup>3</sup> also come from cyclic orbifold constructions starting from fixedpoint free automorphisms of  $\Lambda$  of orders 5, 7, 13 [Abe-Lam-Yamada 2017], from which we get subgroups of Aut  $V^{\natural}$  of the form 5<sup>1+6</sup>.2. $J_2$ , 7<sup>1+4</sup>.2. $A_7$ , and 13<sup>1+2</sup>. $S_4$ . We don't actually need to identify the group structure - it suffices to show that a suitable centralizer in  $Co_1$ has the right power of p in its order.

Then, we consider the large primes:

- (1) The weight 2 space  $V_2^{\natural}$  splits into the direct sum of a trivial representation spanned by the conformal vector  $\omega$  and the 196883-dimensional space of Virasoro primary vectors, namely those annihilated by all  $L_n$ , n > 0. This primary space decomposes into irreducible *C*-modules as 299 from Sym<sup>2</sup>  $\mathfrak{h}$ , 98280 from norm 4 vectors in  $\Lambda$ , and  $24 \cdot 2^{12}$  from the twisted module, and it is not hard to show using triality that it forms an irreducible representation of Aut  $V^{\natural}$ . Since the dimension of an irreducible representation is a divisor of the order we find that the prime factors of 196883 =  $47 \cdot 59 \cdot 71$  are therefore divisors of the order.
- (2) The primary subspaces of weight 3 and 4 are also irreducible, yielding additional prime factors 29, 31, 41 from their dimensions.

We will consider the remaining factors of 11, 17 and 19 in the "open problems" section.

#### 6. Open problems

- (1) Are there analogous vertex-algebraic methods to prove simplicity of other members of the "happy family" of sporadic groups that are subquotients of M? For example, the Baby monster  $\mathbb{B}$  has an order 2 element with centralizer  $2^{1+22}.Co_2$ , where  $Co_2$  is the stabilizer of a norm 4 vector in the automorphism group of  $\Lambda$ , suggesting the existence of a parallel construction. Furthermore, there is a vertex operator algebra with  $\mathbb{B}$  symmetry, constructed in [Höhn 1995] as the commutant of an "Ising model" Virasoro vertex subalgebra of  $V^{\ddagger}$  with central charge 1/2. There are other explicitly constructible vertex operator subalgebras with sporadic symmetry, e.g., Thompson symmetry from the commutant of a 3C pair of Ising vectors, and there are also twisted modules with natural sporadic symmetry, so there are multiple paths to proving simplicity and other properties.
- (2) Is there a vertex-algebraic way to obtain the remaining factors 11, 17, 19 in the order of the monster that were not covered in Theorem 5.4? One can in principle extract them from dimensions of irreducible representations, but higher weight primary subspaces become "very reducible", and this doesn't seem to be much easier than computing the whole character table. As an extreme example, the 16th smallest irreducible representation has dimension 5<sup>9</sup> · 7<sup>6</sup> · 11<sup>2</sup> · 17 · 19, and would solve all of our problems, but a short computation with the character table of the monster and replicability shows that this representation first appears in V<sup>b</sup><sub>60</sub>, with multiplicity 4 [GAP]. There are many other irreducible representations with useful divisibility properties that appear at lower weight, but none of them are particularly convenient. On the other hand, the factors of 17 and 19 appear in centralizers of many

other elements, of orders 2, 3, 5, 7. In particular, a central extension 2.B is the centralizer of an order 2 element  $t \in \mathbb{M}$ , and the irreducible representation of dimension  $2^{13} \cdot 5 \cdot 17 \cdot 19 \cdot 31$  appears in the weight 4 subspace of the twisted module  $V^{\natural}(t)$ . However, isolating this subspace may be nontrivial.

- (3) Is there a vertex-algebraic way to see the centralizer  $11 \times M_{12}$  or normalizer  $(11:5 \times M_{12}): 2$  of an order 11 element in the monster? This would give us a way to see the factor of  $11^2$  in the order.
- (4) How much of the character table of the monster can be extracted from the information in section 5? We have a set of 174 potential McKay-Thompson series as a sort of upper bound, but some non-conjugate classes may yield equal McKay-Thompson series. We also have concrete inputs from transporting a large number of classes from the subgroup C, and we have pure existence of more classes attached to large primes.

## References

[Abe-Lam-Yamada 2017]	T. Abe, C. Lam, H. Yamada, A Remark on $\mathbb{Z}_p$ -orbifold constructions of the moonshine vertex operator algebra ArXiv preprint: https://arxiv.org/abs/ 1705.09022
[Alexander et al. 1992]	D. Alexander, C. Cummins, J. McKay, C. Simons, <i>Completely replicable func-</i> tions Groups, Combinatorics and Geometry. eds. M.W. Liebeck and J. Saxl, (1992) 87–95, Cambridge University Press.
[Aschbacher 1994]	M. Aschbacher, Sporadic Groups Cambridge University Press (1994).
[Borcherds 1986]	R. Borcherds, Vertex algebras, Kac-Moody algebras, and the Monster Proc. Nat. Acad. Sci. USA, 83 no. 10 (1986) 3068–3071.
[Borcherds 1992]	<ul> <li>R. Borcherds, Monstrous moonshine and monstrous Lie superalgebras Invent.</li> <li>Math. 109 (1992) 405-444.</li> </ul>
[Carnahan 2017]	S. Carnahan, 51 constructions of the moonshine module Comm. in Number Theory and Physics 12 no. 2 (2018) 305-334. Available as https://arxiv. org/abs/1707.02954
[Carnahan-Komuro-Urano 2017]	S. Carnahan, T. Komuro, S. Urano, <i>Characterizing moonshine functions by</i> vertex-operator-algebraic conditions SIGMA 14 (2018) 114, 8 pages.
[Carnahan-Miyamoto 2016]	S. Carnahan, M. Miyamoto, Regularity of fixed-point vertex operator subalge- bras Arxiv preprint: https://arxiv.org/abs/1603.05645
[Chen-Lam-Shimakura 2016]	H.Y. Chen, C.H. Lam and H. Shimakura Z3-orbifold construction of the Moon- shine vertex operator algebra and some maximal 3-local subgroups of the Mon- ster Math. Zeit. 288 (2018) 75-100. Available as https://arxiv.org/abs/ 1606_05961
[Conway 1969]	J. Conway, A group of order 8,315,553,613,086,720,000 Bull. London Math. Soc. 1 (1969) 79–88.
[Conway 1985]	J. Conway, A simple construction for the Fischer-Griess monster group Invent. Math. <b>79</b> (1985) 513–540.
[Conway-Norton 1979]	J. Conway, S. Norton, <i>Monstrous Moonshine</i> Bull. Lond. Math. Soc. <b>11</b> (1979) 308–339.
[Conway-Sloane 1982]	J. Conway, <i>Twenty-three constructions for the Leech lattice</i> Proc. Royal Soc. London Ser. A <b>381</b> no. 1781 (1982) 275–283.
[Cummins and Gannon 1997]	C. Cummins, T. Gannon, Modular equations and the genus-zero property of modular functions Invent. Math. 129 (1997) 413–443.
[Dong 1993]	C. Dong, Vertex algebras associated with even lattices J. Algebra <b>160</b> (1993) 245–265.
[Dong 1994]	C. Dong, Representations of the moonshine module vertex operator algebra Contemporary Math. <b>175</b> (1994) 27–36.
[Dong-Griess 2001]	C. Dong and R. Griess, Automorphism groups and derivation algebras of finitely generated vertex operator algebras Michigan Math. J. <b>50</b> no. 2 (2002) 227–239. Available as https://arxiv.org/abs/math/0106051
[Dong-Griess-Höhn 1997]	C. Dong, R.L. Griess, and G. Höhn Framed vertex operator algebras, codes and Moonshine module Comm. Math. Phys. <b>193</b> (1998) 407–448. Available as https://arxiv.org/abs/math/9707008

WHY DO THE SYMMETRIES OF THE MONSTER VERTEX ALGEBRA FORM A FINITE SIMPLE GROUP? 21

[Dong-Li-Mason 1997]	C. Dong, H. Li, G. Mason, Modular invariance of trace functions in orbifold theory Comm. Math. Phys. 214 (2000) no. 1, 1–56. Available as https://
[Dong-Mason-Zhu 1994]	arxiv.org/abs/q-alg/9703016 C. Dong, G. Mason and Y. Zhu, Discrete series of the Virasoro algebra and the moonshine module Proc. Symp. Pure Math. AMS 56 II (1004) 205-316
[Dong-Nagatomo 1998]	C. Dong and K. Nagatomo, Automorphism groups and twisted modules for lattice vertex operator algebras Contemp. Math. 248 (1999), 117–133. Available
[Dong-Nagatomo 1999]	as https://arxiv.org/abs/math/9808088. C. Dong and K. Nagatomo, <i>Classification of irreducible modules for the ver-</i> <i>tex operator algebra</i> $M(1)^+$ , <i>II. Higher rank</i> J. Algebra <b>240</b> (2001) 289-325.
[Dong-Ng-Ren 2021]	C. Dong, S-H. Ng, L. Ren, Orbifolds and minimal modular extensions ArXiv preprint https://arxiv.org/abs/1507_08142
[van Ekeren-Möller-Scheithauer 2015]	J. van Ekeren, S. Möller, N. Scheithauer, <i>Construction and classification of vertex operator algebras</i> J. reine angew. Math. <b>2020</b> no. 759 (2020) 61–99. Available as https://arxiv.org/abs/1507.08142
[Evans-Gannon 2018]	D. Evans, T. Gannon Reconstruction and local extensions for twisted group doubles, and permutation orbifolds Available as https://arxiv.org/abs/1804.
[Frenkel-Ben-Zvi 2004]	E. Frenkel, D. Ben-Zvi, <i>Vertex algebras and Algebraic Curves</i> Mathematical Surveys and Monographs <b>88</b> American Mathematical Society, Providence, RI, (2004).
[Frenkel-Lepowsky-Meurman 1988]	I. Frenkel, J. Lepowsky, A. Meurman, <i>Vertex operator algebras and the Monster</i> Pure and Applied Mathematics <b>134</b> Academic Press, Inc., Boston, MA (1988).
[GAP]	The GAP Group, GAP - Groups, Algorithms, and Programming, Version 4.11.1 (2021) http://www.gap-system.org
[Griess 1982]	R. Griess, The Friendly Giant Invent. Math. 69 (1982) 1–102.
[Griess 1998]	R. Griess, Twelve Sporadic Groups Springer (1998)
[Griess-Lam 2011]	R. Griess, C. H. Lam, A new existence proof of the Monster by VOA theory
[Griess-Lam 2014]	Available as https://arxiv.org/abs/1103.1414 R. Griess, C. H. Lam, Groups of Lie type, vertex algebras, and modular moon-
[Höhn 1995]	Snine Elec. Res. Announc. Math. Sci. 21 (2014) 107–176. G. Höhn, Selbstduale Vertexoperatorsuperalgebren und das Babymonster Ph. D. Thesis, Universität Bonn (1995)
[Huang 1994]	YZ. Huang, A non-meromorphic extension of the moonshine module vertex operator algebra Contemporary Math., <b>193</b> AMS (1996), 123–148. Available
[King 2000]	as http://arxiv.org/abs/hep-th/9406190 O. King, A mass formula for unimodular lattices with no roots Mathematics of Computation <b>72</b> no. 242 (2003) 839-863. Available as https://arxiv.org/ abs/math/0012231.
[Lam-Yamada-Yamauchi 2004]	C. H. Lam, H. Yamada, H. Yamauchi, Vertex operator algebras, extended $E_8$ diagram, and McKay's observation on the monster simple group Trans. Amer. Math. Soc. <b>359</b> (2007) no. 9, 4107–4123. Available as https://arxiv.org/ abs/math/0403010
[Lam-Yamada-Yamauchi 2005]	C. H. Lam, H. Yamada, H. Yamauchi, <i>McKay's observation and vertex operator algebras generated by two conformal vectors of central charge 1/2</i> Int. Math. Res. Pap. (2005), no. 3, 117–181. Available as https://arxiv.org/abs/math/
[Li 1996]	H. Li, Local systems of vertex operators, vertex superalgebras and modules J. Pure Appl. Alg. <b>109</b> (1996) 143-195. Available as https://arxiv.org/abs/ math/9504022.
[Li 1998]	H. Li, Some finiteness properties of regular vertex operator algebras J. Algebra <b>212</b> (1999) 495–514. Available as https://arxiv.org/abs/math/9807077
[Martin 1994]	Y. Martin, On modular invariance of completely replicable functions Moon- shine, the Monster, and related topics (South Hadley, MA, 1994), Contemp. Math. 193 (1996), 263–286.
[Matsuo-Nagatomo 1997]	A. Matsuo, K. Nagatomo, Axioms for a Vertex Algebra and the Locality of Quantum Fields Mathematical Society of Japan Memoirs, Volume 4.

[McRae 2019]	R. McRae, Twisted modules and G-equivariantization in logarithmic conformal field theory Comm. Math. Phys. <b>383</b> no. 3 (2021) 1930–2019. Available as
	https://arxiv.org/abs/1910.13226.
[Miyamoto 1996]	M. Miyamoto, Griess algebras and conformal vectors in vertex operator algebras J. Algebra <b>179</b> (1996) 528–548.
[Miyamoto 1997]	M. Miyamoto, A New Construction of the Moonshine Vertex Operator Algebra over the Real Number Field Ann. Math. (2) <b>159</b> No. 2 (Mar., 2004) 535–596.
[Niemeier 1973]	H-V. Niemeier, Definite quadratische Formen der Dimension 24 und Diskrim- inate 1 J. Number Theory 5 no. 2 (1973) 142–178.
[Norton 2001]	S. Norton, From Moonshine to the Monster Proceedings on Moonshine and related topics (Montréal, QC, 1999), 163–171, CRM Proc. Lecture Notes 30, Amer. Math. Soc., Providence, BI (2001).
[Ogg 1974]	A. Ogg, Automorphisms de courbes modulaires Sem. Delange-Pisot-Poitou, Théorie des nombres <b>16</b> no. 1 (1974-1975), exp. no. 7, 1–8.
[Shimakura 2003]	The automorphism group of the Vertex Operator Algebra $V_L^+$ for an even lattice $L$ without roots J. Algebra, <b>280</b> no. 1 (2004) 29–57. Available as https://arxiv.org/abs/math/0311141
[Shimakura 2006]	H. Shimakura, Lifts of automorphisms of vertex operator algebras in simple current extensions Math. Z. <b>256</b> (2007) no. 3, 491–508. Available as https://arxiv.org/abs/math/0610671
[Shimakura 2010]	H. Shimakura, An E <sub>8</sub> -approach to the moonshine vertex operator algebra J. London Math. Soc. 83 (2011) 493-516. Available as https://arxiv.org/abs/ 1009.4752
[Smith 1979]	S. Smith, Large extraspecial subgroups of width 4 and 6 J. Algebra 58 (1979) 251–280.
[Tits 1984]	J. Tits, On R. Griess' "Friendly Giant" Invent. Math. 78 (1984) 491-499.
[Tuite 1993]	M. Tuite, On the relationship between monstrous Moonshine and the unique- ness of the Moonshine module Comm. Math. Phys. <b>166</b> (1995) 495–532. Avail- able as http://arxiv.org/abs/hep-th/9305057.

22