

A FINITE ANALOGUE OF FINE'S FUNCTION $F(a, b; t)$

RITIKA GOEL

ABSTRACT. We initiate a systematic development of $F_N(a, b; t)$, a finite analogue of Fine's function $F(a, b; t)$. Our results are transformations between $F_N(a, b; t)$ and $F_N(aq^\ell, bq^m; tq^n)$, where ℓ, m and n take the values 0 or 1.

1. INTRODUCTION

In [5], Nathan J. Fine extensively studied the function $F(a, b; t)$ which he defined by

$$F(a, b; t) := \sum_{n=0}^{\infty} \frac{(aq)_n}{(bq)_n} t^n, \quad (1.1)$$

where, here as well as in the rest of the paper, the following standard q -series notation is used:

$$\begin{aligned} (A)_0 &:= (A; q)_0 = 1, \\ (A)_n &:= (A; q)_n = (1 - A)(1 - Aq) \cdots (1 - Aq^{n-1}), \quad n \geq 1, \\ (A)_\infty &:= (A; q)_\infty = \lim_{n \rightarrow \infty} (A; q)_n, \quad |q| < 1, \\ (A_1, A_2, \dots, A_m; q)_n &:= (A_1; q)_n (A_2; q)_n \cdots (A_m; q)_n. \end{aligned}$$

As shown by Fine, $F(a, b; t)$ satisfies nice properties galore and which have applications in basic hypergeometric series, theory of partitions, modular forms and mock theta functions. The first chapter of Fine's book [5] is devoted to establishing functional equations satisfied by $F(a, b; t)$, for example, [5, p. 2, Equation (4.1)]

$$F(a, b; t) = \frac{1 - atq}{1 - t} + \frac{(1 - aq)(b - atq)}{(1 - bq)(1 - t)} tq F(aq, bq; tq). \quad (1.2)$$

There have been attempts to generalize the theory of $F(a, b; t)$ by considering the more general function $\sum_{n=0}^{\infty} \frac{(aq)_n (bq)_n}{(cq)_n (dq)_n} t^n$ although they have not been quite successful, see, for example, [8].

Another direction is to study finite analogues of $F(a, b; t)$. Andrews and Bell [2] studied the finite analogue

$$F(a, b, t, N) := \sum_{n=0}^N \frac{(aq)_n}{(bq)_n} t^n. \quad (1.3)$$

2010 *Mathematics Subject Classification.* Primary 33D15; Secondary 11P81.
Keywords and phrases. Fine's function, basic hypergeometric series, finite Heine transformation.

They constructed another function using $F(a, b, t, N)$, namely, $S_{M,N}(a, b, t)$, which specializes to the series $S_{M,N}$ used by Euler in his unpublished proof of his famous pentagonal number theorem. One of the properties Andrews and Bell proved in their paper [2, Lemma 2.1] is that for $N \geq 1$,

$$F(a, b, t, N) = \frac{1 - atq}{1 - t} + \frac{(1 - aq)(b - atq)}{(1 - bq)(1 - t)} t q F(a, b, t, N) + R_{1,N}(a, b, t), \quad (1.4)$$

where

$$R_{1,N}(a, b, t) = \sum_{j=0}^N \frac{(aq)_j}{(bq)_j} t^j \left(1 - \frac{(b - atq)q^j}{1 - t} \right) + \frac{b - 1}{1 - t}. \quad (1.5)$$

Clearly, we recover (1.2) from (1.4) upon letting $N \rightarrow \infty$ in the latter since $\lim_{N \rightarrow \infty} R_{1,N}(a, b, t) = 0$.

Fine iterated (1.2) to give a proof of the famous Rogers-Fine identity [5, p. 15] given by

$$F(a, b; t) = \sum_{n=0}^{\infty} \frac{(aq)_n (atq/b)_n}{(bq)_n (t)_{n+1}} (1 - atq^{2n+1}) (bt)^n q^{n^2}.$$

In the same vein, Andrews and Bell iterated (1.4) and used the resulting identity along with Euler's method to give a stronger version of the Rogers-Fine identity from which they then deduce the Rogers-Fine identity.

A new finite analogue of $F(a, b; t)$ was studied by the authors of [4, Section 9]. It is given for $N \in \mathbb{N}$ by

$$F_N(a, b; t) := \sum_{n=0}^N \begin{bmatrix} N \\ n \end{bmatrix} \frac{(aq)_n (t)_{N-n} (q)_n t^n}{(bq)_n (t)_N}. \quad (1.6)$$

Clearly, $\lim_{N \rightarrow \infty} F_N(a, b; t) = F(a, b; t)$.

They naturally encountered this function in their work concerning the theory of the restricted partition function $p(n, N)$. The latter is defined as the number of partitions on n whose parts are less than or equal to N . In this study, they derived some properties of $F_N(a, b; t)$ they needed in order to prove their finite analogue of a recent identity of Garvan [6, Equation (1.3)]. These properties are the partial fraction decomposition of $F_N(\alpha, \beta; t)$, namely,

$$F_N(a, b; t) = \frac{(1 - tq^N)(aq)_N}{(bq)_N} \sum_{n=0}^N \begin{bmatrix} N \\ n \end{bmatrix} \frac{(b/a)_n (aq)_{N-n} (aq)^n}{(aq)_N (1 - tq^n)}.$$

This generalizes Fine's partial fraction decomposition of $F(a, b; t)$ [5, p. 18, Equation (16.3)]:

$$F(a, b; t) = \frac{(aq)_{\infty}}{(bq)_{\infty}} \sum_{n=0}^{\infty} \frac{(b/a)_n}{(q)_n} \frac{(aq)^n}{1 - tq^n}. \quad (1.7)$$

They also derived a finite analogue of the Rogers-Fine identity [4, Lemma 9.2], that is, for $\beta \neq 0$,

$$F_N(a, b; t) = (1 - tq^N) \sum_{n=0}^N \begin{bmatrix} N \\ n \end{bmatrix} \frac{(aq)_n (q)_n \left(\frac{atq}{b}\right)_n (atq^2)_{N-1} (tb)^n q^{n^2} (1 - atq^{2n+1})}{(bq)_n (t)_{n+1} (atq^2)_{N+n}}.$$

These two properties were used to obtain the required finite analogue of Garvan's identity [4, Theorem 1.3].

At the end of [4], it was remarked that it would be worthwhile to develop the theory of $F_N(a, b; t)$ considering the enormous impact and applications that the theory of $F(a, b; t)$ has. We set forth such a task with a goal of systematically developing the theory of $F_N(a, b; t)$.

It is to be noted that Andrews [1] has already embarked upon this task. Even though his results are expressed as transformations between ${}_3\phi_2$, they can be equivalently written as transformations or functional equations for $F_N(a, b; t)$. This rephrasing is done in Section 2. Then in Section 3, we present new results on $F_N(a, b; t)$ we have found so far.

It should be mentioned here that Bowman and Wesley [3] have recently obtained new transformations for Fine's function $F(a, b; t)$ iterating what they call as Fine's "seed" identities. A seed identity is an identity where $F(a, b; t)$ gets transformed to $F(aq, b; t)$ or $F(a, bq; t)$ or $F(a, b; tq)$. One of the goals in this paper is obtain the seed identities for $F_N(a, b; t)$.

2. SOME RESULTS OF ANDREWS AND DEDUCTIONS FROM THEM

In [1], Andrews obtained a finite version of Heine's transformation and then derived some corollaries from it. Andrews' transformations are for the ${}_3\phi_2$ hypergeometric series

$${}_3\phi_2 \left[\begin{matrix} q^{-N}, & q, & aq \\ bq, & q^{1-N}/t \end{matrix} ; q, q \right] = F_N(a, b; t), \quad (2.1)$$

as can be easily seen from (1.6).

Thus, we can rephrase these transformations in terms of $F_N(a, b; t)$.

2.1. An identity of Andrews generalizing Equation (6.3) of [5]. From [1, Corollary 3],

$${}_3\phi_2 \left[\begin{matrix} q^{-N}, & abt/c, & b \\ bt, & bq^{1-N}/c. \end{matrix} ; q, q \right] = \frac{(c, t; q)_N}{(c/b, bt; q)_N} {}_3\phi_2 \left[\begin{matrix} q^{-N}, & a, & b \\ c, & q^{1-N}/t \end{matrix} ; q, q \right] \quad (2.2)$$

If we let $b \rightarrow q$, $t \rightarrow b$, $c \rightarrow tq$, $a \rightarrow atq/b$ in the above identity, this gives

$$F_N(a, b; t) = \frac{(1 - tq^N)(1 - b)}{(1 - bq^N)(1 - t)} \sum_{n=0}^N \frac{(q^{-N})_n (atq/b)_n (q)_n q^n}{(tq)_n (q^{1-N}/b)_n (q)_n}. \quad (2.3)$$

Using the elementary identities

$$\begin{aligned} \left(\frac{q^{-N}}{c} \right)_n &= \frac{(-1)^n (cq^{N-n+1})_n q^{\frac{n(n-1)}{2}}}{c^n q^{Nn}}, \\ \frac{(q^{-N})_n}{\left(\frac{q^{1-N}}{b} \right)_n} &= \frac{(q)_N (b)_{N-n} b^n}{(q)_{N-n} (b)_N q^n}, \end{aligned} \quad (2.4)$$

we are led to

$$F_N(a, b; t) = \frac{(1 - b)(1 - tq^N)}{(1 - t)(1 - bq^N)} \sum_{n=0}^N \left[\begin{matrix} N \\ n \end{matrix} \right] \frac{(atq/b)_n (q)_n (b)_{N-n} b^n}{(tq)_n (b)_N}. \quad (2.5)$$

Note that this identity is a finite analogue of Fine's identity [5, Equation (6.3)]:

$$F(a, b; t) = \frac{1 - b}{1 - t} F\left(\frac{at}{b}, t; b\right). \quad (2.6)$$

2.2. **Another ${}_3\phi_2$ transformation of Andrews.** Lemma 1 of [1] gives

$${}_3\phi_2 \left[\begin{matrix} q^{-N}, & a, & b \\ c, & q^{1-N}/t \end{matrix} ; q, q \right] = \frac{(at)_n}{(t)_n} {}_3\phi_2 \left[\begin{matrix} q^{-N}, & c/b, & a \\ c, & at \end{matrix} ; q, btq^N \right] \quad (2.7)$$

Letting $a \rightarrow q$, $b \rightarrow aq$, $c \rightarrow bq$ in the above identity and using (2.1), we get

$$\begin{aligned} F_N(a, b; t) &= \frac{(tq)_N}{(t)_N} {}_3\phi_2 \left[\begin{matrix} q^{-N}, & b/a, & q \\ bq, & tq \end{matrix} ; q, atq^{N+1} \right] \\ &= \frac{(1-tq^N)}{(1-t)} \sum_{n=0}^N (-1)^n \begin{bmatrix} N \\ n \end{bmatrix} \frac{(b/a)_n (q)_n (at)^n q^{n(n+1)/2}}{(bq)_n (tq)_n}. \end{aligned} \quad (2.8)$$

Letting $b = 0$ in the above result gives a finite analogue of [5, Equation (6.1)]:

$$F_N(a, 0; t) = \frac{(1-tq^N)}{(1-t)} \sum_{n=0}^N (-1)^n \begin{bmatrix} N \\ n \end{bmatrix} \frac{(q)_n (at)^n q^{n(n+1)/2}}{(tq)_n}. \quad (2.9)$$

Also, if we let $a \rightarrow 0$ in (2.8), we obtain a finite analogue of [5, Equation (12.3)]:

$$(1-t)F_N(0, b; t) = (1-tq^N) \sum_{n=0}^N \begin{bmatrix} N \\ n \end{bmatrix} \frac{(q)_n (bt)^n q^{n^2}}{(bq)_n (tq)_n}. \quad (2.10)$$

If we let $b = t^{-1} = e^{-i\theta}$ in (2.10), we get

$$(1 - e^{i\theta})F_N(0, e^{-i\theta}; e^{i\theta}) = (1 - e^{i\theta}q^N) + \sum_{n=1}^N \begin{bmatrix} N \\ n \end{bmatrix} \frac{(q)_n q^{n^2}}{(1 - 2q \cos \theta + q^2) \cdots (1 - 2q^n \cos \theta + q^{2n})}, \quad (2.11)$$

which is a finite analogue of [5, Equation (12.32)].

Also, letting $b \rightarrow t$ in (2.10) gives a finite analogue of [5, Equation (12.33)]:

$$(1-t)F_N(0, t; t) = (1-tq^N) \sum_{n=0}^N \begin{bmatrix} N \\ n \end{bmatrix} \frac{(q)_n (t)^{2n} q^{n^2}}{(tq)_n (tq)_n}.$$

Let r and m be positive integers, and replace q by q^m , t by q^r in finite analogue of [5, eqn(12.33)]. Then using the definition of finite analogue of Fine's function for the left side, We obtain the finite analogue of [5, eqn(12.331)]:

$$\begin{aligned} &\sum_{n=0}^N \begin{bmatrix} N \\ n \end{bmatrix}_{q^m} \frac{(q^m, q^m)_n (q^r)_{N-n} q^{rn}}{(q^r)_N (1-q^r)(1-q^{m+r}) \cdots (1-q^{nm+r})} \\ &= (1-q^{N+r}) \sum_{n=0}^N \begin{bmatrix} N \\ n \end{bmatrix}_{q^m} \frac{(q^m, q^m)_n q^{mn^2+2rn}}{[(1-q^r)(1-q^{m+r}) \cdots (1-q^{nm+r})]^2} \end{aligned}$$

2.3. Andrews' finite Heine transformation in terms of $F_N(a, b; t)$. Andrews' finite Heine transformation [1, Theorem 2] in terms of $F_N(a, b; t)$ is given by

$${}_3\phi_2 \left[\begin{matrix} q^{-N}, & c/b, & t & ; q, q \\ at, & q^{1-N}/b & \end{matrix} \right] = \frac{(c, t; q)_N}{(b, at; q)_N} {}_3\phi_2 \left[\begin{matrix} q^{-N}, & a, & b & ; q, q \\ c, & q^{1-N}/t & \end{matrix} \right]. \quad (2.12)$$

Let $t \rightarrow q, b \rightarrow t, c \rightarrow atq$ and $a \rightarrow b$ in the above result. Then

$$F_N(a, b; t) = \frac{(atq, q; q)_N}{(t, bq; q)_N} \sum_{n=0}^N \frac{(b)_n (t)_n q^n}{(atq)_n (q)_n}. \quad (2.13)$$

Letting $N \rightarrow \infty$ gives

$$F(a, b; t) = \frac{(atq)_\infty (q)_\infty}{(t)_\infty (bq)_\infty} \sum_{n=0}^{\infty} \frac{(b)_n (t)_n q^n}{(atq)_n (q)_n}. \quad (2.14)$$

Letting $b = 1$ in (2.13) gives

$$F_N(a, 1; t) = \frac{(atq)_N}{(t)_N}. \quad (2.15)$$

An interesting result is obtained if we multiply both sides of (2.5) by $(1-t)$ and then let $t \rightarrow 1$:

$$\begin{aligned} \lim_{t \rightarrow 1} (1-t) F_N(a, b; t) &= \frac{(1-b)(1-tq^N)}{(1-bq^N)} F_N(a/b, 1; b) \\ &= \frac{(1-b)(1-tq^N)}{(1-bq^N)} \cdot \frac{(aq)_N}{(b)_N} \\ &= \frac{(1-q^N)(aq)_N}{(bq)_N}, \end{aligned} \quad (2.16)$$

where in the last step, we used (2.15). Letting $N \rightarrow \infty$ in the above result we get Equation (6.31) in Fine's book [5]:

$$\lim_{t \rightarrow 1} (1-t) F(a, b; t) = \frac{(aq)_\infty}{(bq)_\infty}.$$

If we let $t \rightarrow 1$ in (2.10) and compare the right-hand side of the resulting identity with $a = 0$ case of (2.16), we arrive at a finite analogue of [5, Equation (12.31)]:

$$\frac{1}{(bq)_N} = (1-q^N) \sum_{n=0}^N \left[\begin{matrix} N \\ n \end{matrix} \right] \frac{(q)_n (b)^n q^{n^2}}{(bq)_n (q)_n}. \quad (2.17)$$

The $b = 1$ case of the above identity generalizes [5, Equation (12.311)].

Next, replace a by b/t in (2.13) and then let $b \rightarrow 0$ to get

$$F_N(b/t, 0; t) = \frac{(bq)_N (q)_N}{(t)_N} \sum_{n=0}^N \frac{(t)_n q^n}{(bq)_n (q)_n}, \quad (2.18)$$

which is a finite analogue of [5, Equation (7.3)].

Further, if we replace a by b/t in (2.9), we have

$$F_N(b/t, 0; t) = \frac{(1-tq^N)}{(1-t)} \sum_{n=0}^N \left[\begin{matrix} N \\ n \end{matrix} \right] \frac{(q)_n (-b)^n q^{n(n+1)/2}}{(tq)_n}. \quad (2.19)$$

Letting $t \rightarrow 0$ in (2.18) and (2.19) and comparing the right-hand sides, we get

$$\sum_{n=0}^N \frac{q^n}{(bq)_n(q)_n} = \frac{1}{(bq)_N(q)_N} \sum_{n=0}^N \begin{bmatrix} N \\ n \end{bmatrix} (q)_n (-b)^n q^{n(n+1)/2}. \quad (2.20)$$

This is a finite analogue of [5, Equation (7.32)]. If we now put $b = 1$ in the above identity, we get a finite analogue of [5, Equation (7.322)].

Letting $b = q^{-1/2}$ in (2.20) and then replacing q by q^2 , we obtain a finite analogue of [5, Equation (7.323)]:

$$\sum_{n=0}^N \frac{q^{2n}}{(q)_{2n}} = \frac{1}{(q)_{2N}} \sum_{n=0}^N \begin{bmatrix} N \\ n \end{bmatrix}_{q^2} (-1)^n (q^2; q^2)_n q^{n^2}.$$

When $a = 0$, (2.13) reduces to

$$\frac{(t)_N (bq)_N}{(q)_N} F_N(0, b; t) = \sum_{n=0}^N \frac{(b)_n (t)_n q^n}{(q)_n}, \quad (2.21)$$

which is a finite analogue of [5, Equation (11.3)].

A finite analogue of [5, Equation (11.4)] is obtained if we let $b = t^{-1}$ in (2.21):

$$\frac{(t)_N (t^{-1}q)_N}{(q)_N} F_N(0, t^{-1}; t) = \sum_{n=0}^N \frac{(t^{-1})_n (t)_n q^n}{(q)_n}. \quad (2.22)$$

3. NEW RESULTS ON $F_N(a, b; t)$

Equation (2.2) of [5] is

$$F(a, b; t) = 1 + \frac{t(1-aq)}{1-bq} F(aq, bq; t). \quad (3.1)$$

We begin with a finite analogue of this result which takes $(a, b) \rightarrow (aq, bq)$.

Theorem 3.1.

$$F_N(a, b; t) = 1 + \frac{t(1-aq)(1-q^{N+1})}{(1-bq)(1-tq^N)} F_N(aq, bq; t). \quad (3.2)$$

Proof.

$$\begin{aligned} F_N(a, b; t) &= \sum_{n=0}^N \begin{bmatrix} N \\ n \end{bmatrix} \frac{(aq)_n (t)_{N-n} (q)_n t^n}{(bq)_n (t)_N} \\ &= 1 + \frac{(1-aq)t}{1-bq} \sum_{n=1}^N \begin{bmatrix} N \\ n \end{bmatrix} \frac{(q^2a)_{n-1} (t)_{N-n} (q)_n t^{n-1}}{(bq^2)_{n-1} (t)_N} \\ &= 1 + \frac{(1-aq)t}{1-bq} \sum_{n=0}^{N-1} \frac{(q)_N (aq^2)_n (t)_{N-n-1} t^n}{(q)_{N-n-1} (bq^2)_n (t)_N} \\ &= 1 + \frac{(1-aq)t}{(1-bq)} \frac{(1-q^N)}{(1-tq^{N-1})} \sum_{n=0}^{N-1} \frac{(q)_{N-1} (t)_{N-n-1} (aq^2)_n t^n}{(q)_{N-n-1} (t)_{N-1} (bq^2)_n}. \end{aligned} \quad (3.3)$$

Using the identity [7, p. 351, (I.11)]

$$\frac{(b, q)_N (a, q)_{N-n} a^n}{(a, q)_N (b, q)_{N-n} b^n} = \frac{(q^{1-N}/b, q)_n}{(q^{1-N}/a, q)_n} \quad (3.4)$$

with $b \rightarrow q, a \rightarrow t$ and replacing N by $N-1$, we have

$$\begin{aligned} F_N(a, b; t) &= 1 + \frac{(1-aq)(1-q^N)t}{(1-bq)(1-tq^{N-1})} \sum_{n=0}^{N-1} \frac{(q^{1-N})_n (aq^2)_n q^n}{(q^{2-N}/t)_n (bq^2)_n} \\ &= 1 + \frac{(1-aq)(1-q^N)t}{(1-bq)(1-tq^{N-1})} {}_3\phi_2 \left[\begin{matrix} q^{1-N}, & aq^2, & q \\ q^{2-N}/t, & bq^2 \end{matrix} ; q, q \right] \end{aligned}$$

Replace N by $N+1$ to get

$$\begin{aligned} F_N(a, b; t) &= 1 + \frac{(1-aq)(1-q^{N+1})t}{(1-bq)(1-tq^N)} {}_3\phi_2 \left[\begin{matrix} q^{-N}, & aq^2, & q \\ q^{1-N}/t, & bq^2 \end{matrix} ; q, q \right] \\ &= 1 + \frac{(1-aq)(1-q^{N+1})t}{(1-bq)(1-tq^N)} F_N(aq, bq, t), \end{aligned} \quad (3.5)$$

where in the last step we used (2.1). \square

Our next result allows us to advance the parameter t in $F_N(a, b; t)$ to tq .

Theorem 3.2.

$$F_N(a, b; t) = \frac{(1-b)(1-tq^{N+1})}{(1-t)(1-bq^{N+1})} + \frac{(1-q^{N+1})(b-atq)}{(1-bq^{N+1})(1-t)} F_N(a, b; tq). \quad (3.6)$$

Proof. Using (2.5), we have

$$\begin{aligned} F_N(a, b; t) &= \frac{(1-b)(1-tq^N)}{(1-t)(1-bq^N)} \sum_{n=0}^N \left[\begin{matrix} N \\ n \end{matrix} \right] \frac{(atq/b)_n (q)_n (b)_{N-n} b^n}{(tq)_n (b)_N} \\ &= \frac{(1-b)(1-tq^N)}{(1-t)(1-bq^N)} + \frac{(1-b)(1-tq^N)}{(1-t)(1-bq^N)} \sum_{n=1}^N \left[\begin{matrix} N \\ n \end{matrix} \right] \frac{(atq/b)_n (q)_n (b)_{N-n} b^n}{(tq)_n (b)_N} \\ &= \frac{(1-b)(1-tq^N)}{(1-t)(1-bq^N)} + \frac{(1-b)(1-tq^N)(1-atq/b)b}{(1-t)(1-bq^N)(1-tq)} \sum_{n=1}^N \left[\begin{matrix} N \\ n \end{matrix} \right] \frac{(atq^2/b)_{n-1} (q)_n (b)_{N-n} b^{n-1}}{(tq^2)_{n-1} (b)_N} \\ &= \frac{(1-b)(1-tq^N)}{(1-t)(1-bq^N)} + \frac{(1-b)(1-tq^N)(1-atq/b)b}{(1-t)(1-bq^N)(1-tq)} \sum_{n=0}^{N-1} \left[\begin{matrix} N \\ n+1 \end{matrix} \right] \frac{(atq^2/b)_n (q)_{n+1} (b)_{N-n-1} b^n}{(tq^2)_n (b)_N} \\ &= \frac{(1-b)(1-tq^N)}{(1-t)(1-bq^N)} + \frac{(1-b)(1-tq^N)(b-atq)(1-q^N)}{(1-t)(1-bq^N)(1-bq^{N-1})(1-tq)} \sum_{n=0}^{N-1} \frac{(q)_{N-1} (atq^2/b)_n (b)_{N-n-1} b^n}{(q)_{N-n-1} (tq^2)_n (b)_{N-1}}. \end{aligned}$$

Let $b \rightarrow q, a \rightarrow b$ in (3.4), replace N by $N-1$ and substitute the resulting expression in the sum on the right-hand side of the above equation so that

$$\begin{aligned} F_N(a, b; t) &= \frac{(1-b)(1-tq^N)}{(1-t)(1-bq^N)} + \frac{(1-b)(1-tq^N)(b-atq)(1-q^N)}{(1-t)(1-bq^N)(1-bq^{N-1})(1-tq)} \sum_{n=0}^{N-1} \frac{(atq^2/b)_n (q^{1-N})_n q^n}{(q^{2-N}/b)_n (tq^2)_n} \\ &= \frac{(1-b)(1-tq^N)}{(1-t)(1-bq^N)} + \frac{(1-b)(1-tq^N)(b-atq)(1-q^N)}{(1-t)(1-bq^N)(1-bq^{N-1})(1-tq)} {}_3\phi_2 \left[\begin{matrix} q^{1-N}, & atq^2/b, & q \\ tq^2, & q^{2-N}/b \end{matrix} ; q, q \right] \end{aligned}$$

Now replace a by atq^2/b , b by q , c by tq^2 , t by b in (2.2), then replace N by $N-1$ and use the resulting identity to transform the ${}_3\phi_2$ on the right-hand side of the above equation so that

$$\begin{aligned} F_N(a, b; t) &= \frac{(1-b)(1-tq^N)}{(1-t)(1-bq^N)} \\ &\quad + \frac{(1-b)(1-tq^N)(b-atq)(1-q^N)(1-tq)(1-bq^{N-1})}{(1-t)(1-bq^N)(1-bq^{N-1})(1-tq)(1-b)(1-tq^N)} {}_3\phi_2 \left[\begin{matrix} q^{1-N}, & aq, & q \\ bq, & q^{2-N}/tq \end{matrix} ; q, q \right] \\ &= \frac{(1-b)(1-tq^N)}{(1-t)(1-bq^N)} + \frac{(1-q^N)(b-atq)(1-bq^{N-1})}{(1-bq^N)(1-bq^{N-1})(1-t)} \sum_{n=0}^{N-1} \frac{(q^{1-N})_n (aq)_n (q)_n q^n}{(bq)_n (q)_n (q^{2-N}/tq)_n}. \end{aligned}$$

Finally replace N by $N+1$ and use (2.4) to obtain

$$\begin{aligned} F_N(a, b; t) &= \frac{(1-b)(1-tq^{N+1})}{(1-t)(1-bq^{N+1})} + \frac{(1-q^{N+1})(b-atq)(1-bq^N)}{(1-bq^{N+1})(1-bq^N)(1-t)} \sum_{n=0}^N \frac{(q)_N (aq)_n (q)_n (tq)_{N-n} (tq)^n}{(q)_{N-n} (bq)_n (q)_n (tq)_N} \\ &= \frac{(1-b)(1-tq^{N+1})}{(1-t)(1-bq^{N+1})} + \frac{(1-q^{N+1})(b-atq)}{(1-bq^{N+1})(1-t)} F_N(a, b; tq). \end{aligned}$$

□

Our next result is a generalization of [5, Equation (4.3)]:

$$F(a, b; t) = \frac{1}{1-t} + \frac{(b-a)tq}{(1-t)(1-bq)} F(a, bq; tq). \quad (3.7)$$

Theorem 3.3. *We have*

$$F_N(a, b; t) = \frac{(1-tq^{N+1})}{(1-t)} + \frac{(1-q^{N+1})(b-a)tq}{(1-bq)(1-t)} F_N(a, bq; tq). \quad (3.8)$$

Proof. From (2.8),

$$\begin{aligned} F_N(a, b; t) &= \frac{1-tq^N}{1-t} + \frac{(1-tq^N)}{(1-t)} \sum_{n=1}^N (-1)^n \begin{bmatrix} N \\ n \end{bmatrix} \frac{(b/a)_n (q)_n (at)^n q^{n(n+1)/2}}{(bq)_n (tq)_n} \\ &= \frac{1-tq^N}{1-t} + \frac{(1-tq^N)(b-a)t}{(1-t)(1-bq)(1-tq)} \sum_{n=0}^{N-1} (-1)^n \frac{(q)_N (bq/a)_n (at)^n q^{(n+2)(n+1)/2}}{(q)_{N-n-1} (bq^2)_n (tq^2)_n}. \end{aligned} \quad (3.9)$$

Now (2.4) with $c = 1/q$ gives

$$(q^{1-N})_n = \frac{(-1)^n q^{n(n-1)/2} (q)_{N-1}}{(q)_{N-n-1} q^{(N-1)n}}. \quad (3.10)$$

Hence

$$\begin{aligned} F_N(a, b; t) &= \frac{1-tq^N}{1-t} + \frac{(1-tq^N)(b-a)(1-q^N)tq}{(1-t)(1-bq)(1-tq)} \sum_{n=0}^{N-1} \frac{(q^{1-N})_n q^{Nn+n} (bq/a)_n (at)^n}{(bq^2)_n (tq^2)_n} \\ &= \frac{1-tq^{N+1}}{1-t} + \frac{(1-tq^{N+1})(b-a)(1-q^{N+1})tq}{(1-t)(1-bq)(1-tq)} \sum_{n=0}^N \frac{(q^{-N})_n q^{(N+1)n+n} (bq/a)_n (at)^n}{(bq^2)_n (tq^2)_n}. \end{aligned} \quad (3.11)$$

Now let $b \rightarrow aq, c \rightarrow bq^2, a \rightarrow q$ and replace t by tq in (2.7) to obtain

$$\sum_{n=0}^N \frac{(q^{-N})_n q^{(N+1)n+n} (bq/a)_n (at)^n}{(bq^2)_n (tq^2)_n} = \frac{(1-tq)}{(1-tq^{N+1})} F_N(a, bq; tq).$$

Finally substitute the above equation in (3.11) to complete the proof. \square

In the next result, we transform $F_N(a, b; t)$ to $F_N(a, bq; t)$.

Corollary 3.4. *We have*

$$F_N(a, b; t) = \frac{1-tq^{N+1}}{1-t} - \frac{(b-a)(1-tq^{N+1})t}{(1-t)(b-at)} + \frac{(b-a)(1-bq^{N+2})t}{(1-bq)(b-at)} F_N(a, bq; t). \quad (3.12)$$

Proof. Replace b by bq in Theorem 3.2 and then substitute the resulting expression for $F_N(a, bq; tq)$ in (3.8). \square

Letting $N \rightarrow \infty$ in Corollary 3.4 gives [5, Equation (4.4)]

$$F(a, b; t) = \frac{b}{b-at} + \frac{(b-a)t}{(1-bq)(b-at)} F(a, bq; t).$$

Our next results relates $F_N(a, b; t)$ with $F_N(aq, b; t)$.

Corollary 3.5. *We have*

$$F_N(a, b; t) = 1 - \frac{b(1-aq)(1-q^{N+1})(1-tq^{N+1})}{(1-tq^N)(1-bq^{N+2})(b-aq)} + \frac{(1-aq)(1-q^{N+1})(b-aqt)}{(1-tq^N)(b-aq)(1-bq^{N+2})} F_N(aq, b; t).$$

Proof. Replace a by aq in Corollary 3.4 and then substitute the resulting expression for $F_N(aq, bq; t)$ in Theorem 3.1. \square

Letting $N \rightarrow \infty$ in Corollary 3.5 gives [5, Equation (4.5)]

$$F(a, b; t) = -\frac{(1-b)aq}{b-aq} + \frac{(1-aq)(b-atq)}{b-aq} F(aq, b; t).$$

Our next theorem transforms $F_N(a, b; t)$ to $F_N(aq, b; tq)$.

Theorem 3.6.

$$\begin{aligned} F_N(a, b; t) = 1 + & \frac{(1-aq)(1-q^{N+1})(1-tq^{N+1})}{(1-tq^N)(1-bq^{N+2})(1-t)} \left(t - \frac{(b-aqt)}{(b-aq)} + \frac{(b-aqt)(1-b)}{(b-aq)(1-bq^{N+1})} \right) \\ & + \frac{(1-aq)(1-q^{N+1})^2(b-aqt)(b-atq^2)}{(1-tq^N)(b-aq)(1-bq^{N+2})(1-bq^{N+1})(1-t)} F_N(aq, b; tq). \end{aligned}$$

Proof. Replace a by aq in Theorem 3.2 and then substitute the resulting expression for $F_N(aq, b; tq)$ in Corollary 3.5. \square

The identity in the above theorem leads to the following identity when we let $N \rightarrow \infty$:

$$F(a, b; t) = \left(\frac{1-b}{1-t} \right) \left(1 - \frac{(b-atq)}{(b-aq)} aq \right) + \frac{(1-aq)(b-atq)(b-atq^2)}{(1-t)(b-aq)} F(aq, b; tq),$$

which is Equation (4.6) from [5].

Our final result is the transformation between $F_N(a, b; t)$ and $F_N(aq, bq; tq)$.

Theorem 3.7. *We have*

$$F_N(a, b; t) = 1 + \frac{(1 - aq)(1 - q^{N+1})(1 - tq^{N+1})t}{(1 - tq^N)(1 - bq^{N+2})(1 - t)} + \frac{(1 - aq)(1 - q^{N+1})^2(b - atq)tq}{(1 - bq)(1 - tq^N)(1 - bq^{N+2})(1 - t)} F_N(aq, bq; tq).$$

Proof. Replace a and b in Theorem 3.2 by aq and bq respectively and then substitute the resulting expression for $F_N(aq, bq; tq)$ in Theorem 3.1. \square

Letting $N \rightarrow \infty$ in Theorem 3.7 gives [5, Equation (4.1)]:

$$F(a, b; t) = \frac{(1 - atq)}{(1 - t)} + \frac{(1 - aq)(b - atq)tq}{(1 - bq)(1 - t)} F(aq, bq; tq).$$

REFERENCES

- [1] G. E. Andrews, *The finite Heine transformation*, Bruce Landman (ed.) et al., Combinatorial number theory. Proceedings of the 3rd Integers Conference 2007, Carrollton, GA, USA, October 24–27, 2007. Berlin: Walter de Gruyter, Integers 9, Suppl., Article A1, 1–6.
- [2] G.E. Andrews and J. Bell, *Euler’s pentagonal number theorem and the Rogers-Fine identity*, Ann. Comb. **16** (2012), 411–420.
- [3] D. Bowman and S. Wesley, *General Fine transformations I*, Int. J. Number Theory **17** (2021), no. 2, 267—283.
- [4] A. Dixit, P. Eyyunni, B. Maji and G. Sood, *Untrodden pathways in the theory of the restricted partition function $p(n, N)$* , J. Combin. Theory Ser. A **180** (2021), 105423 (49 pages)
- [5] N. J. Fine, *Basic Hypergeometric Series and Applications*, Mathematical Surveys and Monographs, Amer. Math. Soc., 1989.
- [6] F. G. Garvan, *Weighted partition identities and divisor sums*, Ch. 12 in Frontiers in Orthogonal Polynomials and q -Series, Contemporary Mathematics and Its Applications: Monographs, Expositions and Lecture Notes: Vol. 1, M. Z. Nashed and X. Li, eds., World Scientific, 2018, pp. 239–249.
- [7] G. Gasper, M. Rahman, *Basic Hypergeometric series*, Second Edition, (Encyclopedia of Mathematics and its applications), 2004.
- [8] A. Gupta, *A study of certain transformations for a special type of ${}_3\phi_2$ series*, Ganita, 44 no. 1-2 (1993), 87-100.

DISCIPLINE OF MATHEMATICS, INDIAN INSTITUTE OF TECHNOLOGY GANDHINAGAR, PALAJ, GANDHINAGAR 382355, GUJARAT, INDIA

Email address: ritikag@iitgn.ac.in