

Solutions to the multimode quantum Rabi model for ultrafast W -state generation

Xun Gao,¹ Liwei Duan,² and Jie Peng^{1,*}

¹*Hunan Key Laboratory for Micro-Nano Energy Materials and Devices
and School of Physics and Optoelectronics, Xiangtan University, Hunan 411105, China*

²*Department of Physics, Zhejiang Normal University, Jinhua 321004, China*

We have obtained the analytical solutions of the multimode quantum Rabi model when all modes have identical frequencies. A series of dark state solutions $|\phi_K\rangle$ with at least K ($K = 1, 2, 3, \dots$) photons are found. Extended to the two-qubit case, such solutions lie closest to a dark state $|\psi_1\rangle$ with at most one photon in the spectrum, but will not affect the adiabatic evolution along $|\psi_1\rangle$ because $\langle\phi_K|\hat{H}|\psi_1\rangle = 0$, and arbitrary single photon multimode W state can be fast generated from $|\psi_1\rangle$. Interestingly, the addition of Stark-shift terms could enlarge the effective energy gap between $|\psi_1\rangle$ and its closet eigenstates, such that the W state can be ultrafast generated in a time proportional to the inverse of the resonator frequency. Moreover, the generation time and fidelity can both be the same for arbitrary $|W_M\rangle$. For this case, another kind of dark state which is a product of just qubit ground states and photonic squeezed states exist in the whole coupling regime with constant energy. The peculiarities of these dark states pave the way of deterministic and ultrafast state-generation in the ultrastrong coupling regime.

Introduction– The quantum Rabi model (QRM) [1, 2] is introduced to describe the simplest interaction between a quantized electromagnetic field mode and a two-level system. Although taking a very simple form, its analytical solution has not been found until 2011 [3]. This is basically because the coexistence of rotating and counter-rotating terms connects all the Fock states, and its general solution consists of infinite photon numbers. This complexity blocks the way to the ultrafast quantum-state generation by using ultrastrong qubit-photon couplings.

Recently, a special solution with at most one photon $|\psi_1\rangle$ was found to exist in the multiqubit multimode quantum Rabi model (MMQRM) [4]. Although the Hamiltonian still connects all the Fock states, the coherent superposition of basis states in this solution makes the photon number states trapped in $|0\rangle$ and $|1\rangle$, so it is called a special dark state [5]. Moreover, a single-photon multimode W state can be fast generated [4]. This paves the way to the fast quantum state generation using the ultrastrong coupling. However, the most important goal—ultrafast quantum-state generation is still undone. At the same time, this one-photon solution gives rise to several intriguing questions for generalized Rabi models: Are there solutions with photon number bounded from below? Why the adiabatic evolution can be so fast even though we did not optimize the adiabatic trajectory? Can the qubit part of a solution also be trapped in only one product state, e.g., the ground state? Are there possible applications for these peculiar solutions?

One interesting generalization is the Rabi-Stark model (QRSM) [6, 7] obtained by adding a nonlinear coupling term

$$H_{NL} = U\sigma_z a^\dagger a \quad (1)$$

to the QRM, where $a^\dagger(a)$ are creation (annihilation) operators, σ_z is a Pauli matrix for the two-level system and

U describes the coupling strength of the Stark term. This model can be realized by trapped ions [8] or the Cavity QED [6] with all parameters can be adjusted freely and independently. The so called Stark shift term may bring about novel physical properties to the original QRM, so it is extensively studied recently. The regular and exceptional solutions of the QRSM have been studied through the Bargmann representation approach [9–11], the Bogoliubov operators approach [12] and the tunable coherent states approach [13]. However, all these solutions do not terminate in the Fock states, in contrast to the “dark states” occurring in the QRM models [14].

Here we have found the analytical solution to the multimode quantum Rabi model when all modes have the same frequencies. One interesting dark-state solution $|\phi_K\rangle$ has at least K ($K = 1, 2, 3, \dots$) photons, although the Hamiltonian connects all the Fock states. Extending it to the multiqubit case and including the Stark-shift terms, we also find similar solutions as $|\phi_K\rangle$ and $|\psi_1\rangle$. What is more, there is another kind of dark state whose qubit part is trapped in just ground state, and the photon part can be expressed as squeezed states. Their peculiarities enable the ultrafast and deterministic generation of arbitrary single-photon multimode W states in the ultrastrong coupling regime, through an adiabatic evolution along $|\psi_1\rangle$. The reason is as follows: When we set $U = \omega$, the effective frequency of the resonator vanishes for qubit ground state. So when coupling $g = 0$, there are infinite degenerate ground states $|\downarrow \dots \downarrow n\rangle$ at $E = -\omega$, where n is the photon number. These eigenstates will always lay below $E = -\omega$ for $g > 0$ and even parity, which should have higher energies when $U < \omega$. This phenomenon enlarges the gap between them and $|\psi_1\rangle$, which has constant eigenenergy $E = \omega$. The adiabatic evolution speed is increased since it is proportional to the square of the inverse of the energy gap. On the other hand, the closest eigenstates to $|\psi_1\rangle$ are $|\phi_{1,2}\rangle$, which satisfy $\langle\phi_{1,2}|\hat{H}|\psi_1\rangle = 0$. Therefore they will not limit the adiabatic speed according to the adiabatic theorem

* jpeng@xtu.edu.cn

[15, 16], and the effective energy gap will be proportional to ω , which results in ultrafast W -state generation (only $11.7\omega^{-1}$) and high fidelity (over 99%). What is more, this generation time and fidelity will not change with mode numbers, and can be exactly the same for arbitrary W states.

Analytical solutions to the multimode quantum Rabi model—First, we study the multimode quantum Rabi model

$$H = \Delta\sigma_z + \sum_{i=1}^M \omega_i a_i^\dagger a_i + g_i \sigma_x (a_i^\dagger + a_i), \quad (2)$$

where $a_i^\dagger(a_i)$ are the i -th photon mode creation (annihilation) operators with frequency ω_i . The qubit is described by Pauli matrices σ_x and σ_z with the level energy splitting 2Δ . g_i is the coupling strength between the i -th mode and the qubit.

This model is solvable when $\omega_i = \omega$, because we can apply transformations [17]

$$\begin{aligned} b_1 &= \frac{\sum_{i=1}^M g_i a_i}{\sqrt{\sum_{i=1}^M g_i^2}}, \\ b_j &= \frac{\sum_{i=1}^{j-1} g_i g_j a_i - \sum_{i=1}^{j-1} g_i^2 a_j}{\sqrt{\sum_{i=1}^j g_i^2 \sum_{i=1}^{j-1} g_i^2}}, \quad j = 2, 3, \dots, M \end{aligned} \quad (3)$$

to transform H into

$$H' = \Delta\sigma_z + \sum_{i=1}^M \omega b_i^\dagger b_i + \left(\sum_{i=1}^M g_i^2\right)^{\frac{1}{2}} \sigma_x (b_1^\dagger + b_1), \quad (4)$$

which is a combination of a single-qubit quantum Rabi model H_R and $M - 1$ free bosonic modes, where the former has been solved. Note that the transformations Eq. (3) is equivalent to the unitary transformation Chilingaryan and Rodríguez-Lara used for the two-mode case [18]. H' will bring very interesting and useful results. First, with the solution to H_R at hand, we just need to set $\sqrt{\sum_{i=1}^N g_i^2} = g$ and change Fock state basis $(a^\dagger)^n |0\rangle$ into $(b_1^\dagger)^n |0\rangle$. E. g., $|1\rangle$ will become $|W_M\rangle = \frac{1}{\sqrt{\sum_{i=1}^M g_i^2}} \sum_{i=1}^M g_i |0_1 0_2 \dots 1_i 0_{i+1} \dots 0_M\rangle$, an arbitrary W state. Second, the solution to H takes the form of $|\psi_R\rangle |n_{b_2} n_{b_3} \dots n_{b_M}\rangle$, where $|\psi_R\rangle$ is the solution of the Rabi model, and $|n_{b_i}\rangle$ is the eigenstate of $b_i^\dagger b_i$. $|\psi_R\rangle$ consists of all Fock states, and it is generally believed so for generalized Rabi models where all Fock states are connected. However, once a $n_{b_i} \neq 0$, the solution to H will have at least n_{b_i} photons. This is surprising but the coherent superposition of some basis states indeed can eliminate certain basis connected by the Hamiltonian, which is the mechanism of the so called dark state. Taking two-mode case for example, the zero-photon part $|\uparrow 00\rangle$ vanishes when a solution with at least one photon $|\downarrow\rangle(g_2|10\rangle - g_1|01\rangle) + \dots$ is applied by H . And $g_2|10\rangle - g_1|01\rangle$ just corresponds to $b_2^\dagger|00\rangle$, so it is easy

to find a series of dark state solutions $|\phi_K\rangle$ with at least K photons.

Extended to the multiqubit case for fast W state generation— $|\phi_K\rangle$ is quite useful in quantum information processing, as we extend H to the multiqubit multimode quantum Rabi model

$$H_{pq} = \sum_{i=1}^M \omega_i a_i^\dagger a_i + \sum_{i=1}^M \sum_{j=1}^N g_{ij} \sigma_{jx} (a_i + a_i^\dagger) + \sum_{j=1}^N \Delta_j \sigma_{jz}, \quad (5)$$

where g_{ij} is the coupling strength between the i -th mode and j -th qubit. $2\Delta_j$ is the energy splitting of the j -th qubit. When $\omega_i = \omega$ and $\frac{g_{ij}}{g_{i'j}}$ is independent of j , H_{pq} can be transformed into

$$\begin{aligned} H'_{qm} &= \omega b_1^\dagger b_1 + \sum_j \sqrt{\sum_i g_{ij}^2} \sigma_{jx} (b_1^\dagger + b_1) + \sum \Delta_j \sigma_{jz} \\ &+ \omega \sum_{i=2}^M b_i^\dagger b_i, \end{aligned} \quad (6)$$

after the same transformation Eq. (3). So the solution to H'_{qm} still has the form of $|\psi_{qR}\rangle |n_{b_2} n_{b_3} \dots n_{b_M}\rangle$, where $|\psi_{qR}\rangle$ is the solution of the multiqubit quantum Rabi model.

For the two-qubit single-mode case, there is a special dark state

$$|\psi_d\rangle = (\Delta_1 - \Delta_2) |\uparrow\uparrow 0\rangle + g_1 |\downarrow\downarrow - \uparrow\downarrow\rangle |1\rangle \quad (7)$$

which exists in the whole coupling regime with constant eigenenergy ω , when $\Delta_1 + \Delta_2 = \omega$ and $g_{11} = g_{12} = g_1$, which can be used to generate $(|\downarrow\uparrow - \uparrow\downarrow\rangle)|1\rangle$ from $|\uparrow\uparrow 0\rangle$ through an adiabatic evolution along $|\psi_d\rangle$. When extended to the multimode case, we need to change $|1\rangle$ into $\sqrt{\sum_{i=1}^M g_i^2} |W_M\rangle$ in $|\psi_d\rangle$ and times it with $|n_{b_2} n_{b_3} \dots n_{b_M}\rangle$, to get the solution, where $g_{ij} = g_i$. When $n_b = \sum_{i=2}^M n_{b_i} = 0$, we easily recover $|\psi_1\rangle$. There will be more energy levels around $|\psi_1\rangle$ when mode number increases, since the lower part of the spectrum will rise up by $n_b \omega$, which seems to reduce the adiabatic speed along $|\psi_1\rangle$, because according to the adiabatic theorem, the system will evolve adiabatically along $|E_n\rangle$ if

$$\left| \frac{\langle E_m(t) | \dot{H} | E_n(t) \rangle}{(E_m - E_n)^2} \right| \ll 1, \quad m \neq n, \quad t \in [0, T]. \quad (8)$$

However, when $n_b = K \neq 0$, these solutions $|\phi_K\rangle$ has photon number bounded from below at K , satisfying $\langle \phi_K | \dot{H} | \psi_1 \rangle = 0$. So the adiabatic speed will not be affected by the mode number, in other words, the speed to generate a high dimensional entangled W state is the same as a single photon. That is why arbitrary W states can be fast generated within exactly the same time, independent of its dimension, as long as the effective coupling $\sqrt{\sum_i g_i^2}$ is fixed.

Here we prove $\langle \phi_K | \dot{H} | \psi_1 \rangle = 0$. To generate $|\downarrow\downarrow - \uparrow\downarrow\rangle |W_M\rangle$ from $|\uparrow\uparrow 0\rangle$, we must set $g_{ij} = g_i = 0$ and $\Delta_1 -$




FIG. 1. Dynamical spectrum of the two-qubit two-mode quantum Rabi model with $\Delta_1(t=0) = \omega$, $\Delta_2(t=0) = 0$, $\dot{\Delta}_1 = -0.005\omega^2 = -\dot{\Delta}_2$, $g(t=0) = g_1(t=0) = g_2(t=0) = 0$, $\dot{g} = \dot{g}_1 = \dot{g}_2 = 0.00255\omega^2$.

$\Delta_2 \neq 0$ initially. To keep the state always in $|\psi_1\rangle$, we must ensure $\dot{\Delta}_1 = -\dot{\Delta}_2$ and $\dot{g}_i/\dot{g}_{i'} = g_i/g_{i'}$, so

$$\dot{H} = \dot{\Delta}_1 \sigma_{1z} - \dot{\Delta}_2 \sigma_{2z} + \sum_i \dot{g}_i (a_i^\dagger + a_i) (\sigma_{1x} + \sigma_{2x}). \quad (9)$$

Therefore

$$\begin{aligned} \dot{H}|\psi_d(t)\rangle &= \frac{(\dot{g}_1(\Delta_1 - \Delta_2) - 2\dot{\Delta}_1 g_1)}{g_1} (|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle) \\ &\times \sum_{i=1}^M g_i |0_1 0_2 \dots 1_i 0_{i+1} \dots 0_M\rangle. \end{aligned} \quad (10)$$

It is easy to find $\langle n_{b_2} n_{b_3} \dots n_{b_M} | \dot{H} |\psi_1(t)\rangle = 0$ if $n_b \neq 0$, so $\langle \phi_K | \dot{H} |\psi_1\rangle = 0$.

Taking the two-mode case for example, if we choose $\Delta_1(t=0) = \omega$, $\Delta_2(t=0) = 0$, $\dot{\Delta}_1 = -0.005\omega^2$, $\dot{g} = \dot{g}_1 = \dot{g}_2 = 0.00255\omega^2$, then $|W_2\rangle$ can be generated in $100\omega^{-1}$ with fidelity reaching 99.9% [4]. The dynamical spectrum in this process is shown in Fig. 1. There are three energy levels close to $|\psi_1\rangle$, where the two blue lines have $n_{b_2} = 1$ and 2 respectively, and numerical results confirm they have at least n_{b_2} photons. They are parallel

to the lower blue energy levels in the spectrum, with a gap of ω or 2ω . The other energy level $|\psi_4\rangle$ close to $|\psi_1\rangle$ possess the structure of $|\downarrow\downarrow\rangle$ and two photons at $g=0$, so $\langle \psi_4 | \dot{H} |\psi_1(t)\rangle / \omega^2 \ll 1$ and decreases with $(\Delta_1 - \Delta_2)/g$. To conclude, the peculiarities of the dark state solutions with photon number upper and lower bounds give rise to the fast generation of arbitrary W states.

Ultrafast generation of W states by adding the Stark-shift terms—The most important goal of state generation utilizing the ultrastrong coupling is to realize ultrafast speed, which is still undone till now. Because there is still one energy level $|\psi_4\rangle$ close to $|\psi_1\rangle$, which is dominated by $|\downarrow\downarrow\rangle$ when $g \sim 0$. Intuitively, if we add Stark-shift terms $U\sigma_z a^\dagger a$, then they will be splitted in the spectrum. So we study the multiqubit multimode Rabi-Stark model

$$\begin{aligned} H_S &= \sum_j \Delta_j \sigma_{jz} + \sum_i (\omega + \sum_j U_{ij} \sigma_{jz}) a_i^\dagger a_i \\ &+ \sum_{ij} g_{ij} (a_i^\dagger + a_i) \sigma_{jx}, \end{aligned} \quad (11)$$

to see if there are similar dark state solutions. When $U_{ij} = U_j$, and $\frac{g_{ij}}{g_{i'j}}$ is independent of j , we can apply the same Bogoliubov transformation Eq. (3) to H_S , and obtain $H'_S = H_{RS} + \sum_{i=2}^M (\omega + \sum_j U_j \sigma_{jz}) b_i^\dagger b_i$ with

$$\begin{aligned} H_{RS} &= \sum_j \Delta_j \sigma_{jz} + (\omega + \sum_j U_j \sigma_{jz}) b_1^\dagger b_1 \\ &+ \sum_j (\sum_i g_{ij}^2)^{\frac{1}{2}} \sigma_{jx} (b_1^\dagger + b_1). \end{aligned} \quad (12)$$

So once we obtain the solution to the multiqubit Rabi-Stark model, we can easily generalize it to the multimode case by adding some free bosonic modes.

Let us start with the two-qubit one-mode case, aiming to find its solution with at most one photon. H_S Eq. (11) possesses a \mathbb{Z}_2 symmetry with the parity operator $R = \exp[i\pi a^\dagger a] \sigma_{1z} \sigma_{2z}$. Supposing there is an eigenstate with at most one photon in the even parity subspace $|\psi_1\rangle = c_1|0\uparrow\uparrow\rangle + c_2|0\downarrow\downarrow\rangle + c_3|1\downarrow\uparrow\rangle + c_4|1\uparrow\downarrow\rangle$, then the eigenenergy equation reads

$$\begin{pmatrix} \Delta_1 + \Delta_2 - E & 0 & g_1 & g_2 \\ 0 & -\Delta_1 - \Delta_2 - E & g_2 & g_1 \\ g_1 & g_2 & \omega - \Delta_1 + \Delta_2 - U_1 + U_2 - E & 0 \\ g_2 & g_1 & 0 & \omega + \Delta_1 - \Delta_2 + U_1 - U_2 - E \end{pmatrix} \begin{pmatrix} c_1 \\ c_2 \\ c_3 \\ c_4 \end{pmatrix} = 0. \quad (13)$$

This linear equation has nontrivial solutions if there are less nonzero rows than columns in the above 6×4 coefficient matrix after elementary row transformation, which can be done and the matrix becomes

$$\begin{pmatrix} g & 0 & -\Delta_1 + \Delta_2 - U_1 + U_2 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}. \quad (14)$$

when $g_1 = g_2 = g$, $\Delta_1 + \Delta_2 = \omega = E$. So we find a




FIG. 2. (a) Spectrum of the two-qubit two-mode quantum Rabi-Stark model with $\Delta_1 = 0.8\omega$, $\Delta_2 = 0.2\omega$, $U_{11} = U_{12} = U_{21} = U_{22} = 0.5\omega$ and $g_{11} = g_{12} = g_{21} = g_{22} = g$ for even parity. (b) Spectrum of the two-qubit two-mode quantum Rabi model with $\Delta_1 = 0.8\omega$, $\Delta_2 = 0.2\omega$ and $g_{11} = g_{12} = g_{21} = g_{22} = g$ for even parity.

quasi-exact solution with at most one photon

$$|\psi_{ds}\rangle = (\Delta_1 - \Delta_2 + U_1 - U_2)|0 \uparrow \uparrow\rangle + g|1\rangle(|\downarrow \uparrow - \uparrow \downarrow\rangle), \quad (15)$$

under this condition, which means there are such solutions for arbitrary g with constant energy, corresponding to a horizontal line in the spectrum. $|\psi_{ds}\rangle$ reduces to $|\psi_d\rangle$ when $U_1 = U_2$, but the adiabatic evolution along $|\psi_{ds}\rangle$ can be much faster. Because the Stark-shift terms $(U_1\sigma_{1z} + U_2\sigma_{2z})a^\dagger a$ will lift the degeneracy between $|0 \uparrow \uparrow\rangle$ and $|2 \downarrow \downarrow\rangle$, and thus enlarge the gap between $|\psi_{ds}\rangle$ and its closest eigenstates.

Extending it to the multimode case, we just need to change $|1\rangle$ into $\sqrt{\sum_{i=1}^M g_i^2} |W_M\rangle$ in $|\psi_{ds}\rangle$ to get the solution for $n_b = \sum_{i=2}^M n_{b_i} = 0$. Taking the two-mode case for example, we compare the spectrum of the two-qubit two-mode Rabi-Stark model and the corresponding Rabi model in Fig. 2. It is very clear that the gap between $|\psi_{ds}\rangle$ and its closet eigenstates is much larger in the former case.

As can be seen in Fig. 2(a), there are infinite energy levels lying below $E = -\omega$. The reason is as follows: When $U_1 + U_2 = \omega$, $\omega a^\dagger a$ are cancelled by $U_1 a^\dagger a \sigma_{1z} + U_2 a^\dagger a \sigma_{2z}$ for $|\downarrow \downarrow\rangle$, so H_{RS} represents a zero-frequency harmonic oscillator, and all $|n_{b_1} \downarrow \downarrow\rangle$ become degenerate when $g = 0$. They will always lie below $E = -\omega$ for even parity as g increases, so they have little effect on the adiabatic evolution of $|\psi_{ds}\rangle$. If $g \neq 0$, the effective Hamiltonian is $(g_1\sigma_{1x} + g_2\sigma_{2x})(b_1 + b_1^\dagger)$ for $|\downarrow \downarrow\rangle|f(b_1^\dagger)|0\rangle$. It seems to have no solution, however,

a squeezed state $|\xi\rangle$ is the eigenstate of $(b_1 + b_1^\dagger)$ when $\xi \rightarrow \infty$ with zero eigenvalue. The only term left in the Hamiltonian is $\Delta_1\sigma_{1z} + \Delta_2\sigma_{2z}$, so there is a dark state

$$|\psi_D\rangle = |\downarrow \downarrow\rangle \sum_{n=0,1,2,\dots} (-1)^n \sqrt{\frac{(2n)!}{2^{2n}(n!)^2}} |2n\rangle \quad (16)$$

with constant eigenenergy $E = -\omega$ in the whole coupling regime, for $\Delta_1 + \Delta_2 = \omega$ and $g_1 = g_2$, as shown in Fig. 2 (a). This state is dark not because the photon number is bounded, but the qubit states are restricted to only $|\downarrow \downarrow\rangle$, although H_{RS} connects it to other states. The most distinct feature is this state is totally a product state, leading to entanglement death, different from other dark states.

Ultrafast generation of arbitrary single-photon multimode W state— We can use $|\psi_{ds}\rangle$ to generate arbitrary single-photon multimode W state $|W_M\rangle$ from state $|0_M, \uparrow, \uparrow\rangle$. Since the energy gap between $|\psi_{ds}\rangle$ and its closest eigenstate reaches order of ω , we expect a ultrafast adiabatic speed. Our scheme is as follows. Two qubits are coupled to M resonators with linear coupling strength g_i ($i = 1, 2, 3, \dots$) and nonlinear coupling strengths U_1, U_2 . The resonator has frequency ω and the qubit energies satisfy $\Delta_1 + \Delta_2 = \omega$. At first, we set $g_i = 0$, and cool down the system to the ground state. Then we excite two qubits by pumping pulses to get $|0_M, \uparrow, \uparrow\rangle$, which corresponds to $|\psi_{ds}\rangle$ when $g_i = 0$ and $\Delta_1 - \Delta_2 + U_{i1} - U_{i2} \neq 0$. Then, we increase g_i to non-zero value and decrease $\Delta_1 - \Delta_2 + U_{i1} - U_{i2}$ to zero. After this adiabatic process, we generate the final state $|W_M \psi_B\rangle$, where $\psi_B = |\downarrow \uparrow - \uparrow \downarrow\rangle$. Aiming to identify its effectiveness, We solve the Schödinger equation numerically to simulate the time evolution of the system. It's showed in Fig. 3(b) that it only takes $T = 20\omega^{-1}$ to reach 99.94% fidelity $F_2 = |\langle W_M \psi_B | \psi(T) \rangle|^2$ with the parameters evolving as Fig. 3(a). In Fig. 3(d), the generation time to reach above 99% fidelity is just $11.7\omega^{-1}$, reaching the order of the inverse of the resonator frequency ($f^{-1} = 2\pi\omega^{-1}$), which indicates an ultrafast state-generation. Meanwhile, we find $U = 0.5$ brings the least generation time to reach above 99% fidelity according to Fig. 4. The generation time is not always decreasing with U because the added Stark-shift terms $U_1\sigma_{1z} + U_2\sigma_{2z}$ sometimes bring more energy levels close to $|\psi_{ds}\rangle$.

As we have discussed above, as long as $\sum_{i=1}^M g_i^2$ is fixed, the spectrum of H_S is the same, except for adding the energy of free bosonic modes $\sum_{i=2}^M (\omega + \sum_j U_j \sigma_{jz}) b_i^\dagger b_i$. These added energy levels will not affect the adiabatic evolution along $|\psi_{ds}\rangle$ because $\langle n_{b_2} n_{b_3} \dots n_{b_M} | \dot{H}_S | \psi_{ds} \rangle = 0$ once $n_b \neq 0$. So the adiabatic speed is the same for fixed $\sum_{i=1}^M g_i^2$. Once we find a combination of g_i which simultaneously possesses high fidelity and speed, we can easily find others by keeping $\sum_{i=1}^M g_i^2$ fixed. Now we use numerical results to testify this conclusion. We set all g_i ($i = 1, 2, \dots, M$) equals to g_M for mode M and $g_M = g_1/\sqrt{M}$, so that $\sum_{i=1}^M g_i^2$ is fixed, as Fig. 5(a) shows. It




FIG. 3. Time Evolution of transferring $|00 \uparrow\uparrow\rangle$ to $|W_2\psi_B\rangle$ by numerical simulation with adjustment of parameters $\Delta_1, \Delta_2, g = g_1 = g_2, U = U_{11} = U_{12} = U_{21} = U_{22}$ when $t\omega$ is set to 20 (a) and 11.7(c). (b)(d) Corresponding population of initial state $|00 \uparrow\uparrow\rangle$, target state $|W_2\psi_B\rangle$ and the special dark like state $|\psi_{2+}\rangle$.




FIG. 4. Generation time needed to reach above 99% fidelity for different $U = U_{11} = U_{12} = U_{21} = U_{22}$.

can be seen from Fig. 5(b) that the fidelity for $|W_M\rangle$ is the same for M ranging from 2 to 10, within the same generation time. Then we fix the fidelity to just above 99%, and find the time need for each M is also the same. At last, we change the ratio between g_1 and g_2 while keep $\sum_{i=1}^M g_i^2$ fixed for the two-mode case, and find the fidelity is also the same with the same generation time.

Conclusion—We have found the analytical solution of the multimode quantum Rabi model when all modes have the same frequency. There are solutions $|\phi_K\rangle$ with at least K photons. When extended to the multiqubit case, there are solutions $|\psi_1\rangle$ consisting of just vacuum and single-photon multimode W state, adjacent to two energy levels $|\phi_K\rangle$ with $K = 1, 2$, respectively. The adiabatic evolution along $|\psi_1\rangle$ is quite fast because $\langle\phi_K|\dot{H}|\psi_1\rangle = 0$,

and the effective energy gap between $|\psi_1\rangle$ and its closest eigenstates is large. When we consider the Stark-shift terms $Ua^\dagger a \sigma_z$, this gap is further enlarged to the order of resonator frequency. Consequently, arbitrary W state can be ultrafast generated with fidelity reach-




FIG. 5. Generation of W states through adiabatic evolution. (a) $g_M = g_1/\sqrt{M}$ for each mode. We set the same parameters and evolution time with the case of Fig. 3(a) except for all g_i equal to g_M in (a) with M ranging from 1 to 10 and get fidelities of $|W_M\rangle = \frac{1}{M} \sum_{i=1}^M |0_1 0_2 \cdots 1_i 0_{i+1} \cdots 0_M\rangle$ depicted in (b). (c) Generation time of prototype W states needed to reach $F_M > 99\%$ for mode M . (d) For two-mode case, Fidelity F_2 with g_1/g_2 ranging from 1 to 10 under condition $g_1^2 + g_2^2 = 0.98$ and $U = U_{11} = U_{12} = U_{21} = U_{22} = 0.5$ within evolution time $T = 11.7\omega^{-1}$.

ing 99% through the adiabatic evolution along $|\psi_1\rangle$ in $11.7\omega^{-1}$. Moreover, if $\sum_{i=1}^M g_i^2$ is fixed, the generation time and fidelity for arbitrary $|W_M\rangle$ will be exactly the same, independent of mode numbers. Finally, we find another interesting dark state solution for the multiqubit multimode Rabi-Stark model, which is a product of qubit ground state and photonic squeezed state.

ACKNOWLEDGMENTS

This work was supported by the Scientific Research Fund of Hunan Provincial Education Department (18A436, 21B0136), Natural Science Foundation of Hunan Province, China (2016JJ6020, 2018JJ3482), the National Basic Research Program of China (2015CB921103), the Program for Changjiang Scholars and Innovative Research Team in University (No. IRT13093), and the National Natural Science Foundation of China (11704320).

[1] I. I. Rabi, On the Process of Space Quantization, *Phys. Rev.* **49**, 324 (1936).

[2] I. I. Rabi, Space Quantization in a Gyating Magnetic

- Field, *Phys. Rev.* **51**, 652 (1937).
- [3] D. Braak, Integrability of the Rabi Model, *Phys. Rev. Lett.* **107**, 100401 (2011).
- [4] J. Peng, J. C. Zheng, J. Yu, P. H. Tang, G. A. Barrios, J. X. Zhong, E. Solano, F. Albarrán-Arriagada, and L. Lamata, One-Photon Solutions to the Multiqubit Multimode Quantum Rabi Model for Fast W -State Generation, *Phys. Rev. Lett.* **127**, 043604 (2021).
- [5] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, Cambridge, England, 1997), Chap. 7.
- [6] A. L. Grimsmo and S. Parkins, Cavity-QED simulation of qubit-oscillator dynamics in the ultrastrong-coupling regime, *Phys. Rev. A* **87**, 033814 (2013).
- [7] A. L. Grimsmo and S. Parkins, Open Rabi model with ultrastrong coupling plus large dispersive-type nonlinearity: Nonclassical light via a tailored degeneracy, *Phys. Rev. A* **89**, 033802 (2014).
- [8] L. Cong, S. Felicetti, J. Casanova, L. Lamata, E. Solano, and I. Arrazola, Selective interactions in the quantum Rabi model, *Phys. Rev. A* **101**, 032350 (2020).
- [9] A. J. Maciejewski and M. Przybylska and T. Stachowiak, Analytical method of spectra calculations in the Bargmann representation, *Phys. Lett. A* **378**, 3445 (2014).
- [10] A. J. Maciejewski and M. Przybylska and T. Stachowiak, An exactly solvable system from quantum optics, *Phys. Lett. A* **379**, 1503 (2015).
- [11] H.-P. Eckle and H. Johannesson, A generalization of the quantum Rabi model: exact solution and spectral structure, *J. Phys. A: Math. Theor.* **50**, 294004 (2017).
- [12] Y. F. Xie, L. w. Duan and Q. H. Chen, Quantum Rabi-Stark model: solutions and exotic energy spectra, *J. Phys. A: Math. Theor.* **52**, 245304 (2019).
- [13] Y. F. Xie and Q. H. Chen, Exact Solutions to the Quantum Rabi-Stark Model Within Tunable Coherent States, *Commun. Theor. Phys.* **71**, 623 (2019).
- [14] J. Peng, Z. Z. Ren, D. Braak, G. J. Guo, G. X. Ju, X. Zhang and X. Y. Guo, Solution of the two-qubit quantum Rabi model and its exceptional eigenstates, *J. Phys. A: Math. Theor.* **47**, 265303 (2014).
- [15] P. Ehrenfest, Adiabatische Invarianten und Quantentheorie, *Ann. Phys. (Leipzig)* **356**, 327 (1916).
- [16] M. Born and V. Fock, Beweis des Adiabatenatzes, *Z. Phys.* **51**, 165 (1928).
- [17] A. W. Chin, Á. Rivas, S. F. Huelga, and M. B. Plenio, Exact mapping between system-reservoir quantum models and semi-infinite discrete chains using orthogonal polynomials, *J. Math. Phys.* **51**, 092109 (2010).
- [18] S. A. Chilingaryan and B. M. Rodríguez-Lara, Exceptional solutions in two-mode quantum Rabi models, *J. Phys. B: At. Mol. Opt. Phys.* **48** 245501 (2015).