

OPERATORS ON THE KALTON-PECK SPACE Z_2

JESÚS M. F. CASTILLO, MANUEL GONZÁLEZ, AND RAÚL PINO

ABSTRACT. We study operators on the Kalton-Peck Banach space Z_2 from various points of view: matrix representations, examples, spectral properties and operator ideals. For example, we prove that there are non-compact, strictly singular operators acting on Z_2 , but the product of two of them is a compact operator. Among applications, we show that every copy of Z_2 in Z_2 is complemented, and each semi-Fredholm operator on Z_2 has complemented kernel and range, the space Z_2 is Z_2 -automorphic and we give a partial solution to a problem of Johnson, Lindenstrauss and Schetchman about strictly singular perturbations of operators on Z_2 .

1. INTRODUCTION

In spite of being a by now classical Banach space, many aspects of the celebrated Kalton-Peck space Z_2 introduced in [21] remain unknown. For instance, it is not known the answer to the *hyperplane problem*: is Z_2 isomorphic to its closed subspaces of codimension 1? The space is clearly isomorphic to its closed subspaces of codimension 2. For a short, sharp, shocked exposition of the few known facts about the structure of Z_2 and its subspaces we refer to [6, Section 10.8] (The Kalton-Peck spaces) and [6, Section 10.9] (Properties of Z_2 explained by itself) and, for a condensed exposition, to the remainder of this section.

In this paper we study operators on the space Z_2 from several points of view. In Section 3 we introduce a matrix representation of these operators, and the results obtained are refined for operators that admit an upper or lower triangular representation in Section 4. In Section 5 we consider an involution $T \longrightarrow T^+$ on the algebra $\mathcal{L}(Z_2)$ of bounded operators on Z_2 introduced in [19] and obtain several new results that we apply to show that every copy of Z_2 in Z_2 is complemented, that the space Z_2 is Z_2 -automorphic and that each semi-Fredholm operator on Z_2 has complemented kernel and range. Section 6 contains a list of natural examples of operators on Z_2 . In Section 7 we follow ideas of Kalton to show that the operator ideals of strictly singular, strictly cosingular and inessential operators coincide and contain each proper operator ideal of $\mathcal{L}(Z_2)$, we prove that the product of two of strictly singular operators on Z_2 is compact, although there are non-compact, strictly singular operators, and we show that the perturbation classes problem has a positive solution for operators on Z_2 . The final Section 8 outlines a few open directions of research.

Next we summarize some facts about the space Z_2 and operators on it. We refer the reader to the Preliminaries (Section 2) for any unexplained notation.

Date: July 5, 2022.

2010 Mathematics Subject Classification: 46B03, 46B10, 46M18, 47A53, 47L20.

Keywords: The Kalton-Peck space; Operators on sequence spaces; Operator ideals; Strictly singular operators.

The research of the first and third author was supported in part by project IB20038 funded by Junta de Extremadura. The research of the first two authors was supported in part by project PID2019-103961GB funded by MINCIN. The research of the third author was partially supported by project FEDER-UCA18-108415 funded by 2014-2020 ERDF Operational Programme and by the Department of Economy, Knowledge, Business and University of the Regional Government of Andalucía.

1.1. Facts about the space Z_2 . Let us make a brief description of the space Z_2 . Consider the homogeneous map $KP : \ell_2 \rightarrow \ell_\infty$ given by $KP(x) = 2x \log |x|$ for normalized $x \in \ell_2$. Then

$$Z_2 = \{(\omega, x) \in \ell_\infty \times \ell_2 : \omega - KP x \in \ell_2\}$$

endowed with the quasinorm $\|(\omega, x)\|_{Z_2} = \|\omega - KP x\|_2 + \|x\|_2$. The space Z_2 is a nontrivial *twisted Hilbert* space in the sense that it contains an uncomplemented subspace M isomorphic to ℓ_2 such that Z_2/M is again isomorphic to ℓ_2 . Indeed, there is a nontrivial exact sequence

$$(1) \quad 0 \longrightarrow \ell_2 \xrightarrow{i} Z_2 \xrightarrow{p} \ell_2 \longrightarrow 0,$$

with inclusion $iy = (y, 0)$ and quotient map $p(\omega, x) = x$. The quasinorm above is equivalent to a norm [21, Theorem 4.7] and thus Z_2 is a reflexive and ℓ_2 -saturated Banach space (see [12]) isomorphic (actually isometric, see below) to its dual [21, Theorem 5.1], something we will write as $Z_2 \simeq Z_2^*$. We will also consider the following quasi-Banach spaces:

- $\text{Dom} KP = \{x \in \ell_2 : KP x \in \ell_2\}$ endowed with the quasinorm $\|x\| = \|KP x\|_2 + \|x\|_2$. This quasinorm is equivalent to the norm of the Orlicz space ℓ_f generated by the Orlicz function $f(t) = t^2 \log^2 t$ [21]; so we can identify $\text{Dom} KP = \ell_f$.
- $\text{Ran} KP = \{\omega \in \ell_\infty : \exists x \in \ell_2, \omega - KP x \in \ell_2\}$ endowed with the quasinorm $\|\omega\|_{f^*} = \inf_{x \in \ell_2} \|\omega - KP x\|_2 + \|x\|_2$, and we can identify $\text{Ran} KP = \ell_f^*$ [5].

There is another natural nontrivial exact sequence

$$(2) \quad 0 \longrightarrow \ell_f \xrightarrow{j} Z_2 \xrightarrow{q} \ell_f^* \longrightarrow 0,$$

with inclusion $jx = (0, x)$, quotient map $q(\omega, x) = \omega$ and associated quasilinear map $KP^{-1} : \ell_f^* \rightarrow \ell_2$ which provides a description of Z_2 as a twisted sum of ℓ_f and ℓ_f^* :

$$Z_2 = \{(x, \omega) \in \ell_\infty \times \ell_f^* : x - KP^{-1} \omega \in \ell_f\}$$

endowed with the quasinorm $\|(x, \omega)\| = \|x - KP^{-1} \omega\|_{\ell_f} + \|\omega\|_{\ell_f^*}$, which is equivalent to the original one on Z_2 and thus there exists $m, M > 0$ so that

$$(3) \quad m\|(\omega, x)\|_{Z_2} \leq \|x - KP^{-1} \omega\|_{\ell_f} + \|\omega\|_{\ell_f^*} \leq M\|(\omega, x)\|_{Z_2}$$

In this representation, $\omega \rightarrow (\omega, KP^{-1} \omega)$ is a bounded homogeneous lifting for q (see [9]) and

- $\text{Dom} KP^{-1} = \{\omega \in \ell_f^* : KP^{-1} \omega \in \ell_f\}$ is endowed with the quasinorm $\|\omega\| = \|KP^{-1} \omega\|_{\ell_f} + \|\omega\|_{\ell_f^*}$. This quasinorm is equivalent to the norm of ℓ_2 so we can identify $\text{Dom} KP^{-1} = \ell_2$.
- $\text{Ran} KP^{-1} = \{x \in \ell_\infty : \exists \omega \in \ell_f^*, x - KP^{-1} \omega \in \ell_f\}$ endowed with the quasinorm $\|x\| = \inf_{\omega \in \ell_f^*} \|x - KP^{-1} \omega\|_{\ell_f} + \|\omega\|_{\ell_f^*}$, and one can identify $\text{Ran} KP^{-1} = \ell_2$.

There is no known explicit formula for KP^{-1} (see [8]). Both compositions $KP \circ KP^{-1} : \ell_f^* \rightarrow \ell_f^*$ and $KP^{-1} \circ KP : \ell_2 \rightarrow \ell_2$ are bounded maps, and condition (\star) in Theorem 3.2 for $T = I_{Z_2}$ implies that the map $I - KP \circ KP^{-1} : \ell_f^* \rightarrow \ell_2$ is also bounded.

The space Z_2 also appears as the derived space obtained by complex interpolation of the scale (ℓ_∞, ℓ_1) at $1/2$. This means that if \mathcal{C} is the corresponding Calderón space, $\delta : \mathcal{C} \rightarrow \ell_\infty$ denotes the evaluation

map at $1/2$ and $\delta' : \mathcal{C} \rightarrow \ell_\infty$ denotes the evaluation of the derivative map at $1/2$ then Z_2 is isomorphically the quotient space $\mathcal{C}/(\ker \delta \cap \ker \delta')$, which means that (isomorphically)

$$(4) \quad Z_2 = \{(\omega, x) \in \ell_\infty \times \ell_2 : \exists f \in \mathcal{C} : f(1/2) = x \text{ and } f'(1/2) = \omega\}$$

endowed with the natural quotient norm. We will use this approach in Section 6.4.

The space Z_2 is not only isometric to its dual, but one singular feature of KP is its “self-duality”, in the sense that $KP^* = -KP$ [21, Theorem 5.1], which is reflected in

$$|\langle KP x, y \rangle - \langle x, KP y \rangle| \leq 2\|x\|_2\|y\|_2.$$

This implies that $Z_2^* = \{(\omega^*, x^*) \in \ell_\infty \times \ell_2 : \omega^* - KP^* x^* \in \ell_2\}$ is the dual of Z_2 with duality formula

$$(5) \quad \langle (\omega^*, x^*), (\omega, x) \rangle = \langle \omega^*, x \rangle + \langle \omega, x^* \rangle,$$

and $D : Z_2 \rightarrow Z_2^*$ given by $D(\omega, x) = (-\omega, x)$ is a bijective isometry.

1.2. Facts about operators on Z_2 . The knowledge about operators on Z_2 is even scarcer than that about Z_2 itself and can be summarized in two results:

Theorem 1.1. [19, Theorems 7 and 8] *Let $S \in \mathfrak{L}(Z_2)$ and $T \in \mathfrak{L}(Z_2, Y)$.*

- (1) *If S is not strictly singular then there exists a subspace E of Z_2 isomorphic to Z_2 such that $S|_E$ is an isomorphism and $S(E)$ is complemented in Z_2 (hence E also is complemented);*
- (2) *If T is not strictly singular then there exists a complemented subspace F of Z_2 isomorphic to Z_2 such that $T|_F$ is an isomorphism.*

Proposition 1.2. [2, Lemma 16.15] *A scalar 2×2 matrix $A = \begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix}$ defines an operator in $\mathfrak{L}(Z_2)$ in the obvious way $A(e_n, e_m) = (\alpha e_n + \beta e_m, \delta e_n + \gamma e_m)$ if and only if $\alpha = \gamma$ and $\delta = 0$.*

2. PRELIMINARIES

2.1. General operator theory. An operator ideal [28] is a subclass \mathcal{A} of the class \mathfrak{L} of bounded operators between Banach spaces such that finite range operators belong to \mathcal{A} , $\mathcal{A} + \mathcal{A} \subset \mathcal{A}$ and $\mathfrak{L}\mathcal{A}\mathfrak{L} \subset \mathcal{A}$.

Let X and Y be Banach spaces. An operator $T \in \mathfrak{L}(X, Y)$ is *strictly singular* if no restriction of T to an infinite dimensional subspace is an isomorphism; T is *strictly cosingular* if $q_N T$ is never surjective when q_N is the quotient map onto an infinite dimensional quotient Y/N . The classes \mathfrak{S} of strictly singular operators and \mathfrak{C} of strictly cosingular operators are closed operator ideals [28, 1.9 and 1.10]. An operator $T \in \mathfrak{L}(X, Y)$ is *upper semi-Fredholm*, $T \in \Phi_+$, if it has closed range and finite dimensional kernel; it is *lower semi-Fredholm*, $T \in \Phi_-$, if its range is finite codimensional (hence closed), and $\Phi = \Phi_+ \cap \Phi_-$ is the class of Fredholm operators. Also T is *inessential*, denoted $T \in \mathfrak{I}_n$, if $I_X - AT$ is a Fredholm operator for all $A \in \mathfrak{L}(Y, X)$ or, equivalently, $I_Y - TA$ is Fredholm for all $A \in \mathfrak{L}(Y, X)$. Introduced by Kleinecke [22], \mathfrak{I}_n is a closed operator ideal containing both \mathfrak{S} and \mathfrak{C} .

2.2. Exact sequences and quasilinear maps. Let X and Y be quasi-Banach spaces with quasi-norms $\|\cdot\|_X$ and $\|\cdot\|_Y$. We suppose that Y is a subspace of some vector space Σ . A map $\Omega : X \rightarrow \Sigma$ is called *quasilinear from X to Y* with ambient space Σ and denoted $\Omega : X \curvearrowright Y$ if it is homogeneous and there exists a constant C so that for each $x, z \in X$,

$$\Omega(x+z) - \Omega x - \Omega z \in Y \quad \text{and} \quad \|\Omega(x+z) - \Omega x - \Omega z\|_Y \leq C(\|x\|_X + \|z\|_X).$$

A quasilinear map $\Omega : X \curvearrowright Y$ is said to be:

- *bounded* if there exists a constant D so that $\Omega x \in Y$ and $\|\Omega x\|_Y \leq D\|x\|_X$ for each $x \in X$.
- *trivial* if there exists a linear map $L : X \longrightarrow \Sigma$ so that $\Omega - L : X \longrightarrow Y$ is bounded.

A quasilinear map $\Omega : X \curvearrowright Y$ generates an exact sequence $0 \rightarrow Y \rightarrow Z \rightarrow X \rightarrow 0$ (namely, a diagram formed by Banach spaces and continuous operators so that the kernel of each of them coincides with the image of the previous one), as follows: $Z = \{(\beta, x) \in \Sigma \times X : \beta - \Omega x \in Y\}$ endowed with the quasi-norm $\|(\beta, x)\|_\Omega = \|\beta - \Omega x\|_Y + \|x\|_X$, with inclusion $y \longrightarrow (y, 0)$ and quotient map $(\beta, x) \longrightarrow x$. The space Z is sometimes called a *twisted sum of Y and X* and denoted $Y \oplus_\Omega X$. If Ω is bounded then $Y \oplus_\Omega X = Y \times X$ and $\|y - \Omega x\|_Y + \|x\|_X$ and $\|y\|_Y + \|x\|_X$ are equivalent quasi-norms on this space. If Ω is trivial then $Y \oplus_\Omega X$ is isomorphic to $Y \oplus X$.

The general theory of twisted sums developed in [21] establishes a correspondence between exact sequences $0 \longrightarrow Y \longrightarrow Z \longrightarrow X \longrightarrow 0$ of quasi Banach spaces and quasilinear maps $\Omega : X \curvearrowright Y$. The quasilinear map generating $0 \longrightarrow \ell_2 \longrightarrow Z_2 \longrightarrow \ell_2 \longrightarrow 0$ is KP.

3. MATRIX REPRESENTATION OF OPERATORS ON Z_2

The space Z_2 admits two canonical descriptions as a twisted sum space: (1) and (2). Operators on Z_2 can be represented by a matrix $\begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix}$, whose entries α, β, δ and γ are linear maps between sequence spaces and depend on whether one is using (1) or (2). Unless specified otherwise, we will always refer to (1); in which case the matrix above acts as

$$\begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix}(\omega, x) = (\alpha\omega + \beta x, \delta\omega + \gamma x).$$

This same operator using (2) has representing matrix $\begin{pmatrix} \gamma & \delta \\ \beta & \alpha \end{pmatrix}$ and acts in the form

$$\begin{pmatrix} \gamma & \delta \\ \beta & \alpha \end{pmatrix}(x, \omega) = (\gamma x + \delta\omega, \beta x + \alpha\omega).$$

We begin with some necessary conditions:

Lemma 3.1. [Necessary conditions] *Let $\begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix}$ be a bounded operator T on Z_2 . The following conditions are satisfied:*

- (d) $\delta : \ell_2 \longrightarrow \ell_2$ is bounded.
- (g) $\gamma : \ell_f \longrightarrow \ell_2$ and $(\beta - \text{KP} \circ \gamma) : \ell_f \longrightarrow \ell_2$ are bounded.
- (d+gK') $\delta + \gamma \circ \text{KP}^{-1} : \ell_f^* \longrightarrow \ell_2$ is a bounded map.
- (g+dK) $\gamma + \delta \circ \text{KP} : \ell_2 \longrightarrow \ell_2$ is a bounded map.

Proof. Recall that i, p, j and q are the maps appearing in the exact sequences (1) and (2).

(d) For $y \in \ell_2$, $pTi y = pT(y, 0) = p(\alpha y, \delta y) = \delta y$.

(g) For $x \in \ell_f$,

$$\|Tjx\|_{Z_2} = \|T(0, x)\|_{Z_2} = \|(\beta x, \gamma x)\|_{Z_2} = \|(\beta - \text{KP} \circ \gamma)x\|_2 + \|\gamma x\|_2 \leq \|Tj\| \cdot \|x\|_{\ell_f}.$$

Thus $\gamma : \ell_f \longrightarrow \ell_2$ and $\beta - \text{KP} \circ \gamma : \ell_f \longrightarrow \ell_2$ are bounded.

(d+gK') A bounded lifting L_q for q is given by $L_q \omega = (\omega, \text{KP}^{-1} \omega)$. Then for every $\omega \in \ell_f^*$, $\|pTL_q \omega\|_2 = \|(\delta + \gamma \circ \text{KP}^{-1})\omega\| \leq \|pT\| \cdot \|L_q\| \cdot \|\omega\|_{\ell_f^*}$. Hence $\delta + \gamma \circ \text{KP}^{-1} : \ell_f^* \longrightarrow \ell_2$ is bounded.

(g+dK) A bounded lifting L_p for p is given by $L_p y = (KP_y, y)$. Then for each $y \in \ell_2$, we have $\|pTL_p y\|_2 = \|(\gamma + \delta \circ KP)y\| \leq \|pT\| \cdot \|L_p\| \cdot \|y\|_2$. Hence $\gamma + \delta \circ KP : \ell_2 \rightarrow \ell_2$ is bounded. \square

Now we characterize the bounded operators on Z_2 .

Theorem 3.2. *The operator $T = \begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix}$ is bounded on Z_2 if and only if the four necessary conditions in Lemma 3.1 hold as well as*

$$(\star) \quad \alpha + \beta \circ KP^{-1} - KP(\delta + \gamma \circ KP^{-1}) : \ell_f^* \rightarrow \ell_2 \quad \text{is a bounded map.}$$

Proof. Condition (\star) is necessary: if T is bounded then $\|\alpha\omega + \beta x - KP(\delta\omega + \gamma x)\|_2 \leq \|T\| \|(\omega, x)\|_{Z_2}$ and the choice $(\omega, KP^{-1}\omega) = L_q\omega$ yields

$$\|\alpha\omega + \beta \circ KP^{-1}\omega - KP(\delta\omega + \gamma \circ KP^{-1}\omega)\|_2 \leq \|L_q\| \|T\| \|\omega\|_{\ell_f^*}.$$

Conversely, we will show that there exists a constant $C > 0$ so that for each $(\omega, x) \in Z_2$ we have $(\alpha\omega + \beta x, \delta\omega + \gamma x) \in Z_2$ and $\|(\alpha\omega + \beta x, \delta\omega + \gamma x)\|_{Z_2} \leq C \|(\omega, x)\|_{Z_2}$. We need to show that

- (1) $\delta\omega + \gamma x \in \ell_2$,
- (2) $\alpha\omega + \beta x - KP(\delta\omega + \gamma x) \in \ell_2$ (hence $\alpha\omega + \beta x \in \ell_\infty$), and
- (3) $\|\delta\omega + \gamma x\|_2 + \|\alpha\omega + \beta x - KP(\delta\omega + \gamma x)\|_2 \leq C \|(\omega, x)\|_{Z_2}$.

Recall that there exist $M > 0$ such that $\|x - KP^{-1}\omega\|_{\ell_f} + \|\omega\|_{\ell_f^*} \leq M \|(\omega, x)\|_{Z_2}$ for each $(\omega, x) \in Z_2$.

(1) Observe that $\omega - KP x \in \ell_2$ and $x - KP^{-1}\omega \in \ell_f$, and by Lemma 3.1, the maps $\gamma + \delta \circ KP : \ell_2 \rightarrow \ell_2$ and $\delta + \gamma \circ KP^{-1} : \ell_f^* \rightarrow \ell_2$ are bounded. Thus

$$\delta\omega + \gamma x = \frac{1}{2} \left(\delta(\omega - KP x) + \gamma(x - KP^{-1}\omega) + (\gamma + \delta \circ KP)x + (\delta + \gamma \circ KP^{-1})\omega \right) \in \ell_2$$

and

$$\begin{aligned} \|\delta\omega + \gamma x\|_2 &\leq \frac{1}{2} \left(\|\delta\| \|\omega - KP x\|_2 + \|\gamma\| \|x - KP^{-1}\omega\|_{\ell_f} + \|\gamma + \delta \circ KP\| \|x\|_2 \right. \\ &\quad \left. + \|\delta + \gamma \circ KP^{-1}\| \|\omega\|_{\ell_f^*} \right) \\ &\leq \frac{1}{2} \left(\|\delta\| + \|\gamma\| M + \|\gamma + \delta \circ KP\| + \|\delta + \gamma \circ KP^{-1}\| M \right) \|(\omega, x)\|_{Z_2}. \end{aligned}$$

To prove (2) we decompose $\alpha\omega + \beta x - KP(\delta\omega + \gamma x)$ in three pieces:

$$\begin{aligned} &\alpha\omega + \beta \circ KP^{-1}\omega - KP(\delta + \gamma \circ KP^{-1})\omega \\ &+ \beta(x - KP^{-1}\omega) - KP(\gamma x - \gamma \circ KP^{-1}\omega) \\ &+ KP(\gamma x - \gamma \circ KP^{-1}\omega) + KP(\delta + \gamma \circ KP^{-1})\omega - KP(\delta\omega + \gamma x). \end{aligned}$$

The first piece is bounded by (\star) ; and for the third piece, note that $KP : \ell_2 \hookrightarrow \ell_2$ is quasilinear, hence $KP(x + y) - KP x - KP y \in \ell_2$ and

$$\|KP(x + y) - KP x - KP y\|_2 \leq \|KP\| (\|x\|_2 + \|y\|_2)$$

for $x, y \in \ell_2$. Moreover γ and $\delta + \gamma \circ \text{KP}^{-1}$ are both bounded from ℓ_f into ℓ_2 . Thus

$$\begin{aligned} & \| \text{KP}(\gamma x - \gamma \circ \text{KP}^{-1} \omega) + \text{KP}(\delta \omega + \gamma \circ \text{KP}^{-1} \omega) - \text{KP}(\delta \omega + \gamma x) \|_2 \\ & \leq \| \gamma x - \gamma \circ \text{KP}^{-1} \omega \|_2 + \| \delta \omega + \gamma \circ \text{KP}^{-1} \omega \|_2 \\ & \leq \| \gamma \| \| x - \text{KP}^{-1} \omega \|_{\ell_f} + \| \delta + \gamma \circ \text{KP}^{-1} \| \| \omega \|_{\ell_f^*} \\ & \leq (\| \gamma \| + \| \delta + \gamma \circ \text{KP}^{-1} \|) M \| (\omega, x) \|_{Z_2}. \end{aligned}$$

For the second piece, since $x - \text{KP}^{-1} \omega \in \ell_f$, $(\beta - \text{KP} \circ \gamma) : \ell_f \rightarrow \ell_2$ is bounded and $\beta(x - \text{KP}^{-1} \omega) - \text{KP}(\gamma x - \gamma \circ \text{KP}^{-1} \omega)$ is equal to

$$\beta(x - \text{KP}^{-1} \omega) - \text{KP} \circ \gamma(x - \text{KP}^{-1} \omega) = (\beta - \text{KP} \circ \gamma)(x - \text{KP}^{-1} \omega),$$

one has

$$\begin{aligned} \| (\beta - \text{KP} \circ \gamma)(x - \text{KP}^{-1} \omega) \|_2 & \leq \| (\beta - \text{KP} \circ \gamma) \| \| x - \text{KP}^{-1} \omega \|_{\ell_f} \\ & \leq M \| (\beta - \text{KP} \circ \gamma) \| \| (\omega, x) \|_{Z_2}. \end{aligned}$$

(3) clearly follows from the arguments in the proof of (1) and (2). \square

Condition (\star) for $T = I_{Z_2}$ implies that $I - \text{KP} \circ \text{KP}^{-1} : \ell_f^* \rightarrow \ell_2$ is bounded, from which we can derive the Benyamini-Lindenstrauss characterization of Proposition 1.2: if $\begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix}$ is a bounded operator on Z_2 with $\alpha, \beta, \gamma, \delta$ scalars then the boundedness of $\gamma + \delta \text{KP} : \ell_2 \rightarrow \ell_2$ implies that $\delta = 0$ while the boundedness of $\alpha + \beta \text{KP}^{-1} - \gamma \text{KP} \circ \text{KP}^{-1} : \ell_f^* \rightarrow \ell_2$ yields that, since βKP^{-1} is also bounded for any scalar β and $\alpha - \gamma + \gamma(I - \text{KP} \circ \text{KP}^{-1})$ is bounded then also $\alpha - \gamma : \ell_f^* \rightarrow \ell_2$ is bounded, and thus $\alpha = \gamma$.

Apart from (\star) and the necessary conditions in Lemma 3.1, we have some additional ones that were not needed in the proof of Theorem 3.2:

Lemma 3.3. *If $T = \begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix} \in \mathcal{L}(Z_2)$ then the following conditions are satisfied:*

- (a) $\alpha : \ell_2 \rightarrow \ell_f^*$ is a bounded operator.
- (b) $\beta : \ell_f \rightarrow \ell_f^*$ is a bounded operator.
- (c₀) $(\alpha - \text{KP} \circ \delta) : \ell_2 \rightarrow \ell_2$ is bounded.
- (c₁) $\alpha \circ \text{KP} + \beta : \ell_2 \rightarrow \ell_f^*$ is bounded.
- (c₂) $\alpha + \beta \circ \text{KP}^{-1} : \ell_f^* \rightarrow \ell_f^*$ is bounded.
- (c₃) $\alpha \circ \text{KP} + \beta - \text{KP}(\delta \circ \text{KP} + \gamma) : \ell_2 \rightarrow \ell_2$ is bounded.
- (c₄) $\gamma - \text{KP}^{-1} \circ \beta : \ell_f \rightarrow \ell_f$ is bounded.

Proof. (a) and (b) follow from $\alpha = qTi$ and $\beta = qTj$.

(c₀) For $y \in \ell_2$, $\|Tiy\|_{Z_2} = \|T(y, 0)\|_{Z_2} = \|(\alpha y, \delta y)\|_{Z_2} = \|(\alpha - \text{KP} \circ \delta)y\|_2 + \|\delta y\|_2 \leq \|Ti\| \cdot \|y\|_2$. Therefore $(\alpha - \text{KP} \circ \delta) : \ell_2 \rightarrow \ell_2$ is bounded.

(c₁) For each $y \in \ell_2$, $\|qTL_p y\| = \|qT(\text{KP}y, y)\|_{Z_2} = \|(\alpha \circ \text{KP} + \beta)y\|_{\ell_f^*} \leq \|T\| \|L_p\| \|y\|_2$.

(c₂) and (c₃) are proved in a similar way, using that $qTL_q \omega = (\alpha + \beta \circ \text{KP}^{-1})\omega$ for each $\omega \in \ell_f^*$ and $\|TL_p y\|_{Z_2} = \|(\alpha \circ \text{KP}y, \delta \circ \text{KP}y)\|_{Z_2} \leq \|T\| \|L_p\| \|y\|_2$.

(c₄) follows from the continuity of Tj . \square

Let us see some applications.

Proposition 3.4. *Let $T = \begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix}$ be an operator on Z_2 .*

- (d*) *If $\gamma : \ell_2 \rightarrow \ell_2$ is bounded then $\delta : \ell_f^* \rightarrow \ell_2$ and $\delta \circ \text{KP} : \ell_2 \rightarrow \ell_2$ are also bounded.*
- (d') *If $\alpha : \ell_2 \rightarrow \ell_2$ is bounded then $\delta : \ell_2 \rightarrow \ell_f$ is bounded. Hence $\text{KP} \circ \delta : \ell_2 \rightarrow \ell_2$ is bounded.*
- (d**) *If $\gamma : \ell_f \rightarrow \ell_f$ is bounded then $\text{KP}^{-1} \circ \beta : \ell_f \rightarrow \ell_f$ is also bounded.*
- (d'') *If $\alpha : \ell_f^* \rightarrow \ell_f^*$ is bounded then $\beta \circ \text{KP}^{-1} : \ell_f^* \rightarrow \ell_f^*$ is bounded.*

Proof. (d*) Since $\text{KP}^{-1} : \ell_f^* \rightarrow \ell_2$ and $\gamma : \ell_2 \rightarrow \ell_2$ are bounded, so is $\gamma \circ \text{KP}^{-1} : \ell_f^* \rightarrow \ell_2$. By (d+gK') in Lemma 3.1, $\delta : \ell_f^* \rightarrow \ell_2$ is bounded, hence $\delta \circ \text{KP} : \ell_2 \rightarrow \ell_2$ is also bounded.

(d') By (c₀) in Lemma 3.3, $\alpha - \text{KP} \circ \delta : \ell_2 \rightarrow \ell_2$ is bounded. Then $\alpha : \ell_2 \rightarrow \ell_2$ bounded implies $\text{KP} \circ \delta : \ell_2 \rightarrow \ell_2$ bounded; hence $\text{KP}^{-1} \circ \text{KP} \circ \delta : \ell_2 \rightarrow \ell_f$ bounded; equivalently, $\delta : \ell_2 \rightarrow \ell_f$ is bounded. For the last equivalence observe that, by the definition of domain of KP and KP^{-1} , for $x \in \ell_2$, $\text{KP}^{-1} \circ \text{KP}x \in \ell_f \Rightarrow \text{KP}x \in \ell_2 \Rightarrow x \in \ell_f$. The assertions (d**) and (d'') can be obtained analogously. \square

Proposition 3.4 yields the boundedness of $\text{KP} \circ \delta$, $\delta \circ \text{KP}$, $\beta \circ \text{KP}^{-1}$ or $\text{KP}^{-1} \circ \beta$ but only under additional conditions that, in general, are not guaranteed. It is for that reasons surprising that one has:

Proposition 3.5. *Let $T = \begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix} \in \mathcal{L}(Z_2)$. Then:*

- (1) *$\text{KP} \circ \delta : \ell_2 \curvearrowright \ell_2$ and $\delta \circ \text{KP} : \ell_2 \curvearrowright \ell_2$ are trivial quasilinear maps.*
- (2) *$\text{KP}^{-1} \circ \beta : \ell_f \curvearrowright \ell_f$ and $\beta \circ \text{KP}^{-1} : \ell_f^* \curvearrowright \ell_f^*$ are trivial quasilinear maps.*

Proof. We prove assertion (1). Consider the pull-back diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \ell_2 & \xrightarrow{i} & Z_2 & \xrightarrow{p} & \ell_2 \longrightarrow 0 \\ & & \parallel & & \uparrow & & \uparrow \delta \\ 0 & \longrightarrow & \ell_2 & \longrightarrow & \text{PB} & \longrightarrow & \ell_2 \longrightarrow 0 \end{array}$$

Then $\text{KP} \circ \delta$ is a quasilinear map generating the lower exact sequence and $\text{KP} \circ \delta$ is trivial if and only if δ admits a bounded linear lifting $\ell_2 \rightarrow Z_2$ [6, Lemma 2.8.3]. Since $\alpha - \text{KP} \circ \delta : \ell_2 \rightarrow \ell_2$ and $\delta : \ell_2 \rightarrow \ell_2$ are bounded, $\hat{\delta}x = (\alpha x, \delta x)$ provides the lifting. Indeed, $\|(\alpha x, \delta x)\|_{Z_2} = \|(\alpha - \text{KP} \circ \delta)x\|_2 + \|\delta x\|_2$.

Analogously, if we consider the push-out diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \ell_2 & \xrightarrow{i} & Z_2 & \xrightarrow{p} & \ell_2 \longrightarrow 0 \\ & & \delta \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & \ell_2 & \longrightarrow & \text{PO} & \longrightarrow & \ell_2 \longrightarrow 0 \end{array}$$

then $\delta \circ \text{KP}$ is a quasilinear map generating the lower exact sequence which is trivial if and only if δ admits a bounded linear extension $\bar{\delta} : Z_2 \rightarrow \ell_2$ [6, Lemma 2.6.3]. In the proof of Theorem 3.2 it is showed that $\bar{\delta}(\omega, x) = \delta\omega + \gamma x$ provides such an extension.

The proof for assertion (2) follows in a similar way: to show that $KP^{-1} \circ \beta : \ell_f \curvearrowright \ell_f$ just consider the pull-back diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \ell_f & \xrightarrow{j} & Z_2 & \xrightarrow{q} & \ell_f^* \longrightarrow 0 \\ & & \parallel & & \uparrow & & \uparrow \beta \\ 0 & \longrightarrow & \ell_f & \longrightarrow & \text{PB} & \longrightarrow & \ell_f \longrightarrow 0 \end{array}$$

(where β is continuous by Proposition 3.3 (b)) and observe that $\omega \rightarrow (\beta\omega, \gamma\omega)$ is a bounded lifting for β by (c_4) . To show that to show that $\beta \circ KP^{-1} : \ell_f^* \curvearrowright \ell_f^*$ just consider the pushout diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \ell_f & \xrightarrow{j} & Z_2 & \xrightarrow{q} & \ell_f^* \longrightarrow 0 \\ & & \downarrow \beta & & \downarrow & & \parallel \\ 0 & \longrightarrow & \ell_f^* & \longrightarrow & \text{PO} & \longrightarrow & \ell_f^* \longrightarrow 0 \end{array}$$

and use (c_2) . □

4. TRIANGULAR OPERATORS

An operator $T \in \mathfrak{L}(Z_2)$ is said to be *compatible with the presentation (1)* if it satisfies $T[i[\ell_2]] \subset i[\ell_2]$. This occurs if and only if its corresponding matrix $\begin{pmatrix} \alpha & \beta \\ 0 & \gamma \end{pmatrix}$ is *upper triangular*; namely, $\delta = 0$. In that case, the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \ell_2 & \xrightarrow{i} & Z_2 & \xrightarrow{p} & \ell_2 \longrightarrow 0 \\ & & \downarrow \alpha & & \downarrow T & & \downarrow \gamma \\ 0 & \longrightarrow & \ell_2 & \xrightarrow{i} & Z_2 & \xrightarrow{p} & \ell_2 \longrightarrow 0 \end{array}$$

is commutative. The operator $T = \begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix}$ above is said to be *compatible with the presentation (2)* if it satisfies $T[j[\ell_f]] \subset j[\ell_f]$, and this occurs if and only if its matrix is *lower triangular*; namely, $\beta = 0$. In that case, the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \ell_f & \xrightarrow{j} & Z_2 & \xrightarrow{q} & \ell_f^* \longrightarrow 0 \\ & & \downarrow \gamma & & \downarrow T & & \downarrow \alpha \\ 0 & \longrightarrow & \ell_f & \xrightarrow{j} & Z_2 & \xrightarrow{q} & \ell_f^* \longrightarrow 0. \end{array}$$

is commutative. Consequently, $T : Z_2 \rightarrow Z_2$ is compatible with both presentations if and only if it is *diagonal*, in the sense that its matrix is $\begin{pmatrix} \alpha & 0 \\ 0 & \gamma \end{pmatrix}$. We show next that the additional conditions of Proposition 3.4 are always satisfied for T upper triangular.

Lemma 4.1. [Necessary conditions] *If $T = \begin{pmatrix} \alpha & \beta \\ 0 & \gamma \end{pmatrix} : Z_2 \longrightarrow Z_2$ is a bounded operator then*

- (a') $\alpha : \ell_2 \longrightarrow \ell_2$ is a bounded operator.
- (g') $\gamma : \ell_2 \longrightarrow \ell_2$ is a bounded operator.
- (k) $\alpha - \gamma$ is compact.

Proof. Since $\delta = 0$, $\alpha : \ell_2 \rightarrow \ell_2$ and $\gamma : \ell_2 \rightarrow \ell_2$ are bounded by (c_0) in Lemma 3.3 and (g+dK) in Lemma 3.1.

(k) was proved in [10, Corollary 5.9]. \square

The necessary conditions become sufficient if we add (see Theorem 1) that

$$\alpha + \beta \circ \text{KP}^{-1} - \text{KP} \circ \gamma \circ \text{KP}^{-1} : \ell_f^* \rightarrow \ell_2 \text{ is a bounded map.}$$

Equivalently, $\alpha\omega + \beta x - \text{KP} \circ \gamma x$ is a bounded map $Z_2 \rightarrow \ell_2$ (recall that $(\omega, x) \in Z_2$ means that $\omega - \text{KP}x \in \ell_2$ or $x - \text{KP}^{-1}\omega \in \ell_f$). Therefore

Theorem 4.2. *An operator $T = \begin{pmatrix} \alpha & \beta \\ 0 & \gamma \end{pmatrix}$ is bounded on Z_2 if and only if $\alpha, \gamma \in \mathfrak{L}(\ell_2)$ and the maps $\beta - \text{KP} \circ \gamma : \ell_f \rightarrow \ell_2$ and $\alpha + \beta \circ \text{KP}^{-1} - \text{KP} \circ \gamma \circ \text{KP}^{-1} : \ell_f^* \rightarrow \ell_2$ are bounded.*

A few variations of the previous result are possible:

Proposition 4.3.

- (a) $S = \begin{pmatrix} \alpha & 0 \\ \delta & 0 \end{pmatrix} \in \mathfrak{L}(Z_2)$ if and only if $\text{KP} \circ \delta - \alpha : \ell_f^* \rightarrow \ell_2$ is bounded. If so, $S \in \mathfrak{S}$. In particular, $\begin{pmatrix} \alpha & 0 \\ 0 & 0 \end{pmatrix} \in \mathfrak{L}(Z_2)$ if and only if $\alpha \in \mathfrak{L}(\ell_f^*, \ell_2)$; and $\begin{pmatrix} 0 & 0 \\ \delta & 0 \end{pmatrix} \in \mathfrak{L}(Z_2)$ if and only if $\delta \in \mathfrak{L}(\ell_f^*, \ell_f)$.
- (b) $R = \begin{pmatrix} 0 & \beta \\ 0 & \gamma \end{pmatrix} \in \mathfrak{L}(Z_2)$ if and only if $\text{KP} \circ \gamma$ is trivial and $\text{KP} \circ \gamma - \beta : \ell_2 \rightarrow \ell_2$ is bounded. If so, $R \in \mathfrak{S}$. In particular, $\begin{pmatrix} 0 & \beta \\ 0 & 0 \end{pmatrix} \in \mathfrak{L}(Z_2)$ if and only if $\beta \in \mathfrak{L}(\ell_2, \ell_2)$; and $\begin{pmatrix} 0 & 0 \\ 0 & \gamma \end{pmatrix} \in \mathfrak{L}(Z_2)$ if and only if $\gamma \in \mathfrak{L}(\ell_2, \ell_f)$.
- (c) $T = \begin{pmatrix} \alpha & \beta \\ 0 & 0 \end{pmatrix} \in \mathfrak{L}(Z_2)$ if and only if $\alpha \circ \text{KP} + \beta : \ell_2 \rightarrow \ell_2$ is bounded. If so, $T \in \mathfrak{S}$.
- (d) $M = \begin{pmatrix} 0 & 0 \\ \delta & \gamma \end{pmatrix} \in \mathfrak{L}(Z_2)$ if and only if $\delta \circ \text{KP} + \gamma : \ell_2 \rightarrow \ell_f$ is bounded. If so, $M \in \mathfrak{S}$.

Proof. Use Proposition 3.5. (a) Suppose $S \in \mathfrak{L}(Z_2)$. Since $S(\omega, x) = (\alpha\omega, \delta\omega)$, S factors through ℓ_f^* ; in fact $S = Bq$ with $B : \ell_f^* \rightarrow Z_2$ given by $B\omega = (\alpha\omega, \delta\omega)$, and B is bounded because $\text{KP} \circ \delta$ is trivial with $\text{KP} \circ \delta - \alpha$ bounded; hence $\omega \rightarrow (\alpha\omega, \delta\omega)$ is a continuous lifting $\ell_f^* \rightarrow Z_2$ for $\delta : \ell_f^* \rightarrow \ell_2$ as in the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \ell_2 & \longrightarrow & Z_2 & \xrightarrow{p} & \ell_2 \longrightarrow 0 \\ & & & & \nwarrow (\alpha, \delta) & & \uparrow \delta \\ & & & & & & \ell_f^* \end{array}$$

Moreover, since p is strictly singular, so is S . Regarding the “in particular” assertion, it is clear that if $\delta \in \mathfrak{L}(\ell_f^*, \ell_f)$ then $\text{KP} \circ \delta : \ell_f^* \rightarrow \ell_2$ is bounded and, by (a), the corresponding S is bounded. Conversely, if S is bounded on Z_2 then $\delta[\ell_f^*] \subset \ell_f$. Moreover, we have $\|\delta\omega\|_{\ell_f} = \|(0, \delta\omega)\|_{Z_2} \leq \|S\| \|(\omega, x)\|_{Z_2}$ and therefore $\delta : \ell_f^* \rightarrow \ell_f$ is bounded. The proofs of (b,c,d) are analogous. \square

Johnson, Lindenstrauss and Schechtman [18] asked whether every operator $T : Z_2 \rightarrow Z_2$ is a strictly singular perturbation of an operator sending $i[\ell_2]$ into itself. We could also consider operators sending $j[\ell_f]$ into itself and formulate the corresponding conjecture. One has the following partial result:

Proposition 4.4. *Let $T = \begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix} \in \mathfrak{L}(Z_2)$*

- (1) *If $\delta \in \mathfrak{L}(\ell_f^*, \ell_f)$ then T is a strictly singular perturbation of an upper triangular operator.*
- (2) *If $\beta \in \mathfrak{L}(\ell_2, \ell_2)$ then T is a strictly singular perturbation of a lower triangular operator.*

Proof. (1) follows from $\begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ 0 & \gamma \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ \delta & 0 \end{pmatrix}$ and Proposition 4.3 (a), and (2) follows from $\begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix} = \begin{pmatrix} \alpha & 0 \\ \delta & \gamma \end{pmatrix} + \begin{pmatrix} 0 & \beta \\ 0 & 0 \end{pmatrix}$ and Proposition 4.3(b). \square

5. COMPLEMENTED COPIES OF Z_2 IN Z_2

Kalton [19] considered the continuous alternating bilinear form $\Omega : Z_2 \times Z_2 \rightarrow \mathbb{R}$ given by

$$\Omega((\omega_1, x_1), (\omega_2, x_2)) = \langle \omega_1, x_2 \rangle - \langle \omega_2, x_1 \rangle,$$

and for every $T \in \mathfrak{L}(Z_2)$ he defined an operator $T^+ : Z_2 \rightarrow Z_2$ by $\Omega(T^+y, z) = \Omega(y, Tz)$ for $y, z \in Z_2$. Note that $(Dx)y = \Omega(x, y)$ for $y, z \in Z_2$. Hence $T^+ = D^{-1}T^*D \in \mathfrak{L}(Z_2)$, where $T^* \in \mathfrak{L}(Z_2^*)$ is the conjugate operator of T . Indeed, for $x, y \in Z_2$ we have

$$(DT^+x)y = \Omega(T^+x, y) = \Omega(x, Ty) = Dx(Ty) = (T^*Dx)y;$$

hence $DT^+ = T^*D$. Moreover, the map $T \rightarrow T^+$ is an *involution* on $\mathfrak{L}(Z_2)$ [30, Definition 11.14].

Since D is a bijective isometry, most of the properties of T^+ coincide with those of T^* . Namely, $\|T\| = \|T^+\|$, $R(T)$ is closed if and only if $R(T^+)$ is so, T is an isomorphism into if and only if T^+ is surjective, and $T \in \Phi_-$ if and only if $T^+ \in \Phi_+$. In particular, $T^+ = T$ and $T \in \Phi_+$ imply $T \in \Phi$.

It is easy to check that if $T = \begin{pmatrix} \alpha & \beta \\ \delta & \gamma \end{pmatrix}$ then $T^* = \begin{pmatrix} \gamma^* & \beta^* \\ \delta^* & \alpha^* \end{pmatrix}$ and $T^+ = \begin{pmatrix} \gamma^* & -\beta^* \\ -\delta^* & \alpha^* \end{pmatrix}$.

Definition 5.1. [19] *A subspace E of Z_2 is said to be isotropic when $\Omega(u, v) = 0$ for every $u, v \in E$. An operator $T \in \mathfrak{L}(Z_2)$ is isotropic when its range is isotropic; equivalently, when $T^+T = 0$.*

Clearly $i[\ell_2]$, $j[\ell_f]$ and $\{(x, x) \in Z_2 : x \in \ell_f\}$ are isotropic subspaces of Z_2 . Moreover, the operator $ip = \begin{pmatrix} 0 & I \\ 0 & 0 \end{pmatrix} \in \mathfrak{L}(Z_2)$ is isotropic, non-compact, strictly singular and strictly cosingular, with $R(ip) = N(ip)$ and $(ip)^+ = -ip$.

Lemma 5.2. *For each $T \in \mathfrak{L}(Z_2)$, $R(T) \cap N(T^+)$ is an isotropic subspace.*

Proof. Let $(\omega_1, x_1), (\omega_2, x_2) \in R(T) \cap N(T^+)$. Then $(\omega_2, x_2) = T(\omega, x)$ for some $(\omega, x) \in Z_2$ and

$$\Omega((\omega_1, x_1), (\omega_2, x_2)) = \Omega((\omega_1, x_1), T(\omega, x)) = \Omega(T^+(\omega_1, x_1), (\omega, x)) = 0$$

concluding the proof. \square

Proposition 5.3. *Let $T \in \mathfrak{L}(Z_2)$.*

- (a) *If T^+T is strictly singular then so is T .*
- (b) *If Ti is strictly singular then so is T .*

(c) If $Ti \in \Phi_+$ then $T \in \Phi_+$.

Proof. (a) and (b) are [19, Theorem 9 and Lemma 5].

(c) Suppose that $T \notin \Phi_+$. Then there exists an infinite dimensional subspace $M \subset Z_2$ such that $T|_M$ is compact. Since p is strictly singular, we can find a normalized basic sequence (x_n) in M and sequences (x_n^*) in Z_2^* with $C = \sup_n \|x_n^*\| < \infty$ and (y_n) in $i[\ell_2] = N(p)$ with $\|x_n - y_n\| < 2^{-n}/C$. Then the expression $Kx = \sum_{i=1}^{\infty} x_i^*(x)(x_i - y_i)$ defines a compact operator $K \in \mathfrak{L}(Z_2)$ with $\|K\| < 1$ such that, denoting by N the closed subspace generated by (x_n) , we have $(I - K)[N] \subset i[\ell_2]$. We claim that $Ti|_N$ is compact, hence $Ti \notin \Phi_+$. Indeed, if (z_n) is a bounded sequence in N , then $(T(I - K)z_n) = (Tz_n - TKz_n)$ has a convergent subsequence. Since TK is compact, (Tz_n) has a convergent subsequence. \square

Lemma 5.4. *Let $T \in \mathfrak{L}(Z_2)$. If $T \in \Phi_+$ then $T^+T \in \Phi$.*

Proof. Since $(T^+T)^+ = T^+T$, it is enough to show that $T^+T \in \Phi_+$. Suppose that $T^+T \notin \Phi_+$. Then $T^+Ti \notin \Phi_+$; hence there exists a normalized block basis sequence $(w^{(n)})$ in ℓ_2 such that, if we denote by $i_w : \ell_2 \rightarrow \ell_2$ the isometric embedding defined by $i_w e_n = w^{(n)}$, then $T^+T i_w$ is compact. Let W be the block operator associated to the sequence $(w^{(n)})$ in [19, Section 4]. Since $T^+T W i e_n = T^+T i w e_n$ for each $n \in \mathbb{N}$, $T^+T W i$ is compact; hence $T^+T W \in \mathfrak{S}$ by (b) in Proposition 5.3. Thus $W^+T^+T W \in \mathfrak{S}$, hence $TW \in \mathfrak{S}$ by (a) in Proposition 5.3, implying $T \notin \Phi_+$. \square

Observe that $T^+T \in \Phi_+$ implies $T \in \Phi_+$.

It is known [2, Theorem 16.16] that every infinite dimensional complemented subspace of Z_2 contains a further complemented subspace isomorphic to Z_2 . It is not known whether every infinite dimensional complemented subspace of Z_2 is isomorphic to Z_2 (which in particular would imply that Z_2 is isomorphic to its hyperplanes). We add now a new piece of knowledge:

Theorem 5.5. *Every subspace of Z_2 isomorphic to Z_2 is complemented.*

Proof. Let $T \in \mathfrak{L}(Z_2)$ an isomorphism into with range $R(T)$. Then $T^+T \in \Phi$, hence there exists a finite codimensional subspace N of $R(T)$ such that $T^+|_N$ is an isomorphism and $T^+[N]$ is finite codimensional, hence complemented; thus N is complemented and so is $R(T)$. \square

An extension of Theorem 5.5 in operator terms is available now:

Theorem 5.6. *Every semi-Fredholm operator on Z_2 has complemented kernel and range.*

Proof. Let T be an operator on Z_2 . If $T \in \Phi_+$ then the kernel is finite dimensional, and we can prove that $R(T)$ is complemented with the proof of Theorem 5.5. If $T \in \Phi_-$ then $R(T)$ is closed finite codimensional and $T^* \in \Phi_+$ (in Z_2^*). Since $Z_2^* \simeq Z_2$, $R(T^*)$ is complemented by the first part, hence $N(T) = {}^\perp R(T^*)$ is also complemented. \square

Recall from [1] that a Banach space X is said to be Y -automorphic if every isomorphism between two infinite codimensional subspaces of X isomorphic to Y can be extended to an automorphism of Z_2 . It is clear that ℓ_2 is ℓ_2 -automorphic and that Z_2 is not ℓ_2 -automorphic: indeed, since $Z_2 \simeq Z_2 \oplus Z_2$, an isomorphism between the subspaces $\ell_2 \oplus 0$ and $\ell_2 \oplus \ell_2$ cannot be extended to an automorphism of Z_2 . Surprisingly, one has:

Proposition 5.7. *Z_2 is Z_2 -automorphic.*

Proof. As Kalton remarks in [19, p. 110], Pełczyński's decomposition argument shows that if E is a complemented subspace of Z_2 and $E \oplus E \simeq E$ then E is isomorphic to Z_2 . Suppose that $Z_2 \simeq Z_2 \oplus F$. By (2) of Theorem 1.1 one has $F \simeq Z_2 \oplus N$, and thus $F \oplus F \simeq F \oplus Z_2 \oplus N \simeq Z_2 \oplus N \simeq F$. Hence $F \simeq Z_2$. \square

Now if E is an infinite dimensional complemented subspace of Z_2 then

$$E \simeq Z_2 \oplus E' \simeq Z_2 \oplus Z_2 \oplus E' \simeq Z_2 \oplus E$$

and thus E is isomorphic to its 2-codimensional subspaces since

$$E \oplus \mathbb{K}^2 \simeq Z_2 \oplus E \oplus \mathbb{K}^2 \simeq Z_2 \oplus E \simeq E.$$

We conjecture that $E \oplus E \simeq Z_2$.

Since $Z_2^* \simeq Z_2$ and i is strictly cosingular [21] (hence i^* strictly singular), one can easily derive the following results, by duality, from the previous ones:

Proposition 5.8. *Let $T \in \mathfrak{L}(Z_2)$.*

- (a) *If T^+T is strictly cosingular then so is T^+ .*
- (b) *If pT is strictly cosingular then so is T .*
- (c) *If $pT \in \Phi_-$ then $T \in \Phi_-$.*
- (d) *If $T^+ \in \Phi_-$ then $T^+T \in \Phi$.*

6. EXAMPLES OF OPERATORS ON Z_2

6.1. Rank-one operators. Given $(x^*, \omega^*) \in Z_2^*$ and $(u, v) \in Z_2$, the rank-one operator $(x^*, \omega^*) \otimes (u, v)$ acts on Z_2 as follows. For each $(\omega, x) \in Z_2$,

$$[(x^*, \omega^*) \otimes (u, v)](\omega, x) = \langle (x^*, \omega^*), (\omega, x) \rangle \cdot (u, v) = (\omega\omega^* + x^*x) \cdot (u, v).$$

Thus the matrix associated to $(x^*, \omega^*) \otimes (u, v)$ is $\begin{pmatrix} \omega^* \otimes u & x^* \otimes u \\ \omega^* \otimes v & x^* \otimes v \end{pmatrix}$.

6.2. Nuclear operators. Fix $u^* \in Z_2^*$ and $v \in Z_2$ and obtain the matrix representation for the one-dimensional map $u^* \otimes v$. If $u^* = \sum a_n u_n^*$ and $v = \sum b_n v_n$, set $\mathbf{u}(2n-1)^* = \sum a_{2n-1} u_{2n-1}^*$ the “odd” part of u^* and $\mathbf{u}(2n)^* = \sum a_{2n} u_{2n}^*$ the “even” part of u^* . Define in the same manner the odd and even parts $\mathbf{v}(2n-1)$ and $\mathbf{v}(2n)$ of v to obtain

$$u^* \otimes v = \begin{pmatrix} \mathbf{u}(2n-1)^* \otimes \mathbf{v}(2n-1) & \mathbf{u}(2n)^* \otimes \mathbf{v}(2n-1) \\ \mathbf{u}(2n-1)^* \otimes \mathbf{v}(2n) & \mathbf{u}(2n)^* \otimes \mathbf{v}(2n) \end{pmatrix}$$

Consequently, if $T = \sum u_k^* \otimes v_k$ is a nuclear operator on Z_2 one gets

$$T = \begin{pmatrix} \sum_k \mathbf{u}_k(2n-1)^* \otimes \mathbf{v}_k(2n-1) & \sum_k \mathbf{u}_k(2n)^* \otimes \mathbf{v}_k(2n-1) \\ \sum_k \mathbf{u}_k(2n-1)^* \otimes \mathbf{v}_k(2n) & \sum_k \mathbf{u}_k(2n)^* \otimes \mathbf{v}_k(2n) \end{pmatrix}$$

6.3. Operators acting on the scale of ℓ_p spaces. The fact that Z_2 is the derived space at ℓ_2 for the scale of ℓ_p spaces provides some elements of $\mathcal{L}(Z_2)$. We say that an operator $\alpha : \ell_2 \rightarrow \ell_2$ *acts on the scale* when there are $1 \leq p < 2 < q \leq \infty$ such that both $\alpha : \ell_p \rightarrow \ell_p$ and $\alpha : \ell_q \rightarrow \ell_q$ are bounded. In this case $\tau_\alpha = \begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix}$ is a continuous operator on Z_2 (see [9]).

Against the naive intuition, τ_α can be a bounded operator on Z_2 and still α is not necessarily an operator acting on the scale, as the following simple example shows: Let $z \in \ell_f$ such that $z \notin \ell_p$ for $p < 2$. Then $\alpha_0(x) = e_1^*(x)z$ defines a bounded operator on ℓ_2 but $\alpha_0(\ell_p) \not\subset \ell_p$ for $p < 2$; thus α_0 does not act in the scale. However τ_{α_0} is a bounded operator on Z_2 because $(0, e_1^*), (e_1^*, 0) \in Z_2^*$, $(z, 0), (0, z) \in Z_2$ and $\tau_{\alpha_0} = (0, e_1^*) \otimes (z, 0) + (e_1^*, 0) \otimes (0, z)$.

We next present several natural examples of operators α acting on the scale:

- (1) α a diagonal operator D_σ with $\sigma \in \ell_\infty$, or α a right (or left) shift operator.
- (2) α a surjective isometry on ℓ_p for some $p \neq 2$. It was proved in [24, Proposition 2.f.14] that these operators have the form $\alpha((x_n)_n) = (\varepsilon_n x_{\sigma(n)})_n$ for some permutation $\sigma : \mathbb{N} \rightarrow \mathbb{N}$ and some sequence of signs $(\varepsilon_n)_n$. The induced operator τ_α is then an isometry [20].
- (3) α the *Cesàro operator* C defined by $C((x_n)_n) = \left(\frac{1}{n} \sum_{k=1}^n x_k \right)_n$.

It is bounded on ℓ_p for $p > 1$ with $\|C\|_p = \frac{p}{p-1}$ [29]. It is not bounded on ℓ_1 since $C(e_1)$ is the harmonic series. See [4] for the properties of C as an operator on ℓ_2 .

- (4) α a *Hilbert matrix operator* H_λ defined by a Hilbert matrix

$$\left(\frac{1}{n+m+\lambda} \right)_{n,m=0}^\infty, \quad \lambda \in \mathbb{C} \setminus \{0, -1, -2, \dots\}.$$

The operator H_λ is bounded on ℓ_p for $1 < p < \infty$. See [31].

- (5) α a *Hausdorff operator* A associated to a sequence of complex scalars $\{\mu_n : n = 0, 1, 2, \dots\}$ (see [3, Section 3.4]). It has the form $A((x_k)_k) = \left(\sum_{k=0}^n a_{nk} x_k \right)_n$ with

$$a_{nk} = \begin{cases} \binom{n}{k} \Delta^{n-k} \mu_k & 0 \leq k \leq n \\ 0 & k > n \end{cases},$$

where $\Delta(\mu_n) = \mu_n - \mu_{n+1}$. Many Hausdorff operators are bounded on some spaces ℓ_p [3, 29]:

- (i) the *generalized Cesàro operator* C_a^α arising from $\mu_n = \frac{\Gamma(a+\alpha)\Gamma(n+a)}{\Gamma(a)\Gamma(n+a+\alpha)}$, where Γ is the Gamma function. For $\alpha > 0$, $a > 1/p$ the operator C_a^α is bounded on ℓ_p ($p > 1$) with norm equal to $\frac{\Gamma(a+\alpha)\Gamma(a-1/p)}{\Gamma(a+\alpha-1/p)}$. Note that $C_1^1 = C$, the Cesàro operator.
- (ii) the *Hölder operator* H_α arising from $\mu_n = (n+1)^{-\alpha}$. For $\alpha > 0$, H_α is bounded on ℓ_p with norm $\left(\frac{p}{p-1} \right)^\alpha$;
- (iii) the *Euler operator* (E, r) arising from $\mu_n = a^n$, where $0 < a < 1$, $r = \frac{1-a}{a}$ and its norm on ℓ_p is $(1+r)^{1/p}$;
- (iv) the *Gamma operator* Γ_a^α arising from $\mu_n = \left(\frac{a}{n+a} \right)^\alpha$, where $\alpha > 0$, and it is bounded on ℓ_p with norm $\left(\frac{a}{a-1/p} \right)^\alpha$ whenever $a > 1/p$.

We do not know whether every Hausdorff operator which is bounded on ℓ_2 acts on the scale.

The following result belongs to Sneiberg [32]: *Let (X_0, X_1) and (Y_0, Y_1) be two interpolation pairs such that $T : X_i \rightarrow Y_i$ is bounded for $i = 0, 1$. If $T^{-1} : X_\theta \rightarrow Y_\theta$ exists and it is bounded for some $0 < \theta < 1$, then there is $\varepsilon > 0$ such that $T^{-1} : X_s \rightarrow Y_s$ exists and it is bounded for $|s - \theta| < \varepsilon$. We can infer from that:*

Proposition 6.1. *For an operator $\alpha : \ell_2 \rightarrow \ell_2$ acting on the scale, the following statements are equivalent:*

- (i) τ_α is an isomorphism.
- (ii) $\alpha : \ell_2 \rightarrow \ell_2$ is an isomorphism.
- (iii) There exists $\varepsilon > 0$ such that $\alpha : \ell_p \rightarrow \ell_p$ is an isomorphism for all $|2 - p| < \varepsilon$.

Consequently $\sigma(\tau_\alpha) = \sigma(\alpha)$.

Proof. If α acts on the couple (ℓ_p, ℓ_{p^*}) then the operator $\alpha - \lambda I$ also acts on that same couple. Moreover, if τ_α is an isomorphism then $\alpha : \ell_2 \rightarrow \ell_2$ is an isomorphism. And that if both α and α^{-1} act on some scale (ℓ_p, ℓ_{p^*}) then τ_α is an isomorphism on Z_2 . \square

If α is an operator acting on the scale, its spectrum on ℓ_p maybe independent of p , as it is the case of diagonal operators or the left and right shift operators, but it also may vary with p : Leibowitz [23] proved that, for $1 < p < \infty$ the Cesàro operator C on ℓ_p has no eigenvalues and its spectrum is

$$\sigma(C) = \{\lambda \in \mathbb{C} : |\lambda - p^*/2| \leq p^*/2\}.$$

Moreover, $\lambda I - C$ is a Fredholm operator with index -1 for $|\lambda - p^*/2| < p^*/2$, and has dense proper range for $|\lambda - p^*/2| = p^*/2$ [15].

The Hilbert matrix operator H_1 on ℓ_2 has no eigenvalues and $\sigma(H_1)$ is the interval $[0, \pi]$, see [25], while the spectrum on ℓ_p varies with p , and it has eigenvalues for $p > 2$ and residual points for $p < 2$ [31]. Since the matrix representing H_1 is symmetric, the conjugate operator of $H_1 : \ell_p \rightarrow \ell_p$ is $H_1 : \ell_{p^*} \rightarrow \ell_{p^*}$. Thus the spectra of H_1 on ℓ_p and ℓ_{p^*} coincide.

6.4. Operators on the Calderón space. We obtain operators on Z_2 by picking operators on the Calderón space $T : \mathcal{C} \rightarrow \mathcal{C}$ such that $T[\ker \delta \cap \ker \delta'] \subset \ker \delta \cap \ker \delta'$. The simplest way to do that is to pick an operator τ on the scale and then set $T(f)(z) = \tau(f(z))$. If $\varphi : \mathbb{S} \rightarrow \mathbb{D}$ is a conformal map, then the operator $S(f)(z) = \tau(\varphi(z)f(z))$ induces $\begin{pmatrix} 0 & \tau \\ 0 & 0 \end{pmatrix}$. Therefore, given two operators α, ϕ on the

scale, $T(f)(z) = \alpha(f(z)) + \beta(\varphi(z)f(z))$ induces the upper triangular operator $\begin{pmatrix} \alpha & \beta \\ 0 & \alpha \end{pmatrix}$ on Z_2 .

6.5. Diagonal operators. The continuity of diagonal operators D_σ on Z_2 is related with the unconditional structure of the space. Recall that the sequence $(u_n)_{n \in \mathbb{N}}$ given by $u_{2n-1} = (e_n, 0)$ and $u_{2n} = (0, e_n)$, where $(e_n)_n$ is the canonical basis of ℓ_2 , is a basis on Z_2 which is not unconditional. Therefore not all $\sigma \in \ell_\infty$ define a diagonal operator on Z_2 . Let us denote $a = (\sigma_{2n-1})$ and $b = (b_n = \sigma_{2n})$. If $D_\sigma = \begin{pmatrix} D_a & 0 \\ 0 & D_b \end{pmatrix}$ is an operator on Z_2 then $D_a - D_b = D_{a-b}$ is compact by Lemma 4.1; thus $a - b \in c_0$. It is startling that an additional condition is required:

Proposition 6.2. *A diagonal operator $D_\sigma : Z_2 \rightarrow Z_2$ defined by a monotone decreasing sequence σ is bounded if and only if $(|\sigma_{2k-1} - \sigma_{2k}|)_{k \in \mathbb{N}} = O\left(\frac{1}{\log n}\right)$.*

Proof. Cabello and García showed in [7] that a diagonal operator $D_a : \ell_2 \rightarrow \ell_2$ can be lifted to Z_2 if and only if the decreasing rearrangement sequence $(a_n^*)_n$ is $O\left(\frac{1}{\log n}\right)$. The self duality of the Kalton-Peck space yields the result. \square

6.6. Block operators. Let $U = (u_n)$ be a bounded sequence of disjointly supported blocks in ℓ_2 . We define a bounded operator $u : \ell_2 \rightarrow \ell_2$ by $ux = \sum x_n v_n$. Kalton [19] defined the operator T_U on Z_2 by $T_U(e_n, 0) = u_n$ and $T_U(0, e_n) = (KPu_n, u_n)$, and proved it is an isometry. Let us call T_U a *block operator*. As we said in 6.3, if α is an operator acting on the scale then τ_α is an upper triangular operator on Z_2 . In general, a perturbation of τ_α is required to make it an upper triangular operator. In particular, the operator u defined by a sequence of disjointly supported normalized blocks in ℓ_1 is not an operator on the scale. In [11, Theorem 4.6] it is explained how to obtain the required perturbation and how this perturbation yields the Kalton block operator $T_U = \begin{pmatrix} u & KPu \\ 0 & u \end{pmatrix}$ mentioned above.

7. OPERATOR IDEALS ON Z_2

The classes \mathfrak{S} and \mathfrak{C} are not dual to each other. In general $T^* \in \mathfrak{S} \implies T \in \mathfrak{C}$ and $T^* \in \mathfrak{C} \implies T \in \mathfrak{S}$. But since Z_2 is reflexive, it turns out that an operator $T : Z_2 \rightarrow Z_2$ is strictly singular (resp. cosingular) if and only if $T^* : Z_2^* \rightarrow Z_2^*$ is strictly cosingular (resp. singular). One moreover has:

Theorem 7.1. *One has the identities*

$$\mathfrak{S}(Z_2) = \mathfrak{C}(Z_2) = \mathfrak{In}(Z_2).$$

Moreover, that set contains every proper ideal of $\mathfrak{L}(Z_2)$.

Proof. By the first part of Theorem 1.1, if $S \in \mathfrak{L}(Z_2) \setminus \mathfrak{S}(Z_2)$ then there exists $A, B \in \mathfrak{L}(Z_2)$ so that $ASB = I_{Z_2}$; hence S does not belong to any proper operator ideal, and $\mathfrak{In}(Z_2) \subset \mathfrak{S}(Z_2)$. Since $\mathfrak{S}(X)$ and $\mathfrak{C}(X)$ are contained in $\mathfrak{In}(X)$ for each X and $Z_2 \simeq Z_2^*$, the equalities follow. \square

Observe that $\mathfrak{S}(Z_2, \ell_\infty) \neq \mathfrak{L}(Z_2, \ell_\infty) = \mathfrak{In}(Z_2, \ell_\infty)$: let $T \in \mathfrak{L}(Z_2, \ell_\infty)$. For every $A \in \mathfrak{L}(\ell_\infty, Z_2) = \mathfrak{S}(\ell_\infty, Z_2)$, $I - AT$ is Fredholm. Similarly, $\mathfrak{C}(\ell_1, Z_2) \neq \mathfrak{L}(\ell_1, Z_2) = \mathfrak{In}(\ell_1, Z_2)$.

The next result is a dual version of the second part of Theorem 1.1.

Proposition 7.2. *Let $T \in \mathfrak{L}(X, Z_2)$. If $T \notin \mathfrak{C}$ then there exists a complemented subspace N of Z_2 with Z_2/N isomorphic to Z_2 such that $Q_N T$ is surjective, where $Q_N : Z_2 \rightarrow Z_2/N$ is the quotient map.*

Proof. If $T \in \mathfrak{L}(X, Z_2)$ is not in \mathfrak{C} then $T^* \in \mathfrak{L}(Z_2^*, X^*)$ is not in \mathfrak{S} . Since $Z_2^* \simeq Z_2$, by Theorem 1.1 there exists a complemented subspace M of Z_2^* isomorphic to Z_2^* such that $T^*|_M$ is an isomorphism. Then $N = {}^\perp M$ is a subspace of Z_2 satisfying the required conditions. \square

Let \mathfrak{K} be the class of compact operators and let $L_p \equiv L_p(0, 1)$ for $1 \leq p \leq \infty$. Then $\mathfrak{S}(L_p) \neq \mathfrak{K}(L_p)$ for $p \neq 2$ [26], but $T \in \mathfrak{S}(L_p)$ implies $T^2 \in \mathfrak{K}(L_p)$ [14].

Theorem 7.3. *We have $\mathfrak{S}(Z_2) \neq \mathfrak{K}(Z_2)$, but $S, T \in \mathfrak{S}(Z_2)$ implies $ST \in \mathfrak{K}$.*

Proof. As we mentioned before, $ip \in \mathfrak{L}(Z_2)$ is strictly singular but not compact.

For the remaining part, recall that an operator S acting on a reflexive space X is compact if and only if for every normalized weakly null sequence (x_n) in X , (Sx_n) has a norm null subsequence; and it was proved in [21, Theorem 5.4] that every normalized weakly null sequence (x_n) in Z_2 has a subsequence equivalent either to the (usual) basis of ℓ_2 or to the (usual) basis on ℓ_f .

Let $S, T \in \mathfrak{S}(Z_2)$ and let (x_n) be a normalized weakly null sequence Z_2 . If (x_{n_k}) is a subsequence equivalent to the basis of ℓ_f then (Tx_{n_k}) has no subsequence equivalent to the basis of ℓ_f because T is strictly singular; hence (Tx_{n_k}) has a subsequence equivalent to the basis of ℓ_2 or it is norm null. Also, if (x_{n_k}) is a subsequence equivalent to the basis of ℓ_2 then (Tx_{n_k}) has no subsequence equivalent to the basis of ℓ_2 , and has no subsequence equivalent to the basis of ℓ_f because $\ell_f \subsetneq \ell_2$: a bounded operator cannot take the unit basis of ℓ_2 to the unit basis of ℓ_f ; hence it is norm null. In each case, (STx_n) has a norm null subsequence. \square

The *perturbation class* of a class of operators $\mathcal{A} \subset \mathfrak{L}$ is defined by its components as follows when $\mathcal{A}(X, Y) \neq \emptyset$:

$$P\mathcal{A}(X, Y) = \{L \in \mathfrak{L}(X, Y) : T + L \in \mathcal{A}(X, Y) \text{ for all } T \in \mathcal{A}(X, Y)\}.$$

Kato and Vladimirkii (see [28, Section 26.6]) proved that $\mathfrak{S} \subset P\Phi_+$ and $\mathfrak{C} \subset P\Phi_-$, and it is known that $\mathfrak{In} = P\Phi$. The *perturbation classes problem* asks whether $\mathfrak{S} = P\Phi_+$ and $\mathfrak{C} = P\Phi_-$. This problem has a positive answer under certain conditions but not in general (see [16]), and also for Z_2 although this space does not verify those conditions.

Proposition 7.4. *We have $P\Phi(Z_2) = P\Phi_+(Z_2) = P\Phi_-(Z_2) = \mathfrak{S}(Z_2)$.*

Proof. In general, $\mathfrak{S}(X) \subset P\Phi_+(X) \subset P\Phi(X) = \mathfrak{In}(X)$ and $\mathfrak{C}(X) \subset P\Phi_-(X) \subset \mathfrak{In}(X)$, but Theorem 7.1 implies $\mathfrak{S}(Z_2) = \mathfrak{C}(Z_2) = \mathfrak{In}(Z_2)$. So all these inclusions are equalities for $X = Z_2$. \square

8. FURTHER DIRECTIONS OF RESEARCH

The overall tone of these suggestions is to determine which properties of operators on ℓ_2 are valid for operators on Z_2 and which are not.

8.1. The convolution on $\mathfrak{L}(Z_2)$. Here we consider the relation between T and T^+ as operators in $\mathfrak{L}(\ell_2)$.

Question 1. Suppose that $T \in \mathfrak{L}(Z_2)$ is an isomorphism into. Is T^+T bijective? What does it mean $T^+T = I$ or $TT^+T = T$ for $T \in \mathfrak{L}(Z_2)$?

Question 2. Is $\mathfrak{L}(Z_2)/\mathfrak{S}(Z_2)$ isomorphic to a C^* -algebra?

Clearly $\mathfrak{L}(Z_2)$ is not isomorphic to a C^* -algebra since there exists $T \neq 0$ such that $T^+T = 0$. However $T^+T \in \mathfrak{S}(Z_2)$ implies $T \in \mathfrak{S}(Z_2)$, and $T \in \mathfrak{S}(Z_2)$ if and only if $T^+ \in \mathfrak{S}(Z_2)$.

8.2. Polynomially bounded operators. A *contraction* is an operator T with $\|T\| \leq 1$, and for a polynomial p we denote $\|p\|_\infty = \sup_{|z| < 1} |p(z)|$.

Question 3. Is every contraction in $\mathfrak{L}(Z_2)$ polynomially bounded? Equivalently,

$$(6) \quad \exists C > 0 \text{ so that } \|T\| \leq 1 \Rightarrow \|p(T)\| \leq C\|p\|_\infty \text{ for every polynomial } p?$$

Note that (6) with $C = 1$ isometrically characterizes Hilbert spaces. Moreover, if X is isomorphic to a Hilbert space clearly (6) holds for some $C \geq 1$; however the converse implication fails (see [33]).

8.3. The group of invertible operators. We denote by $\mathfrak{GL}(X)$ the group of invertible operators on a Banach space X . It is known that $\mathfrak{GL}(\ell_2)$ is connected in the complex case, while $\mathfrak{GL}(\ell_p \times \ell_q)$ is not connected for $1 \leq p < q < \infty$ [13, 27].

Question 4. In the case $\mathbb{K} = \mathbb{C}$, is $\mathfrak{GL}(Z_2)$ connected?

Is the subgroup $\{T \in \mathfrak{GL}(Z_2) : T \text{ is upper triangular}\}$ connected?

The latter question could be tackled by obtaining a characterization of the invertible operators $T \in \mathfrak{GL}(Z_2)$ in terms of the components $\alpha, \beta, \delta, \gamma$ of the matrix representation of T .

8.4. Representations of Z_2 . A basic question whose meaning is not even clear is whether there are other “natural” presentations of Z_2 beyond the ℓ_2 and the ℓ_f presentations considered in this paper. In homological terms, since $Z_2 \simeq Z_2 \oplus Z_2$ one could obtain other nontrivial representations such as $0 \longrightarrow \ell_2 \longrightarrow Z_2 \longrightarrow \ell_2 \oplus Z_2 \longrightarrow 0$, $0 \longrightarrow \ell_f \longrightarrow Z_2 \longrightarrow \ell_f^* \oplus Z_2 \longrightarrow 0$, etc. Or even the trivial one

$$0 \longrightarrow Z_2 \longrightarrow Z_2 \longrightarrow Z_2 \longrightarrow 0.$$

None of these representations are even “isomorphic” to either (1) or (2).

REFERENCES

- [1] A. Aviles, F. Cabello Sánchez, J.M.F. Castillo, M. González, Y. Moreno, *Separable injective Banach spaces*, Lecture Notes in Mathematics 2132 (2016) Springer-Verlag.
- [2] Y. Benyamini, J. Lindenstrauss, *Geometric nonlinear functional analysis, Vol. 1*, Amer. Math. Soc., Providence, 1999.
- [3] J. Boos, *Classical and modern methods in sumbability*, Oxford Univ. Press, 2000.
- [4] A. Brown, P.H. Halmos, A.L. Shields, *Cèsaro operators*, Acta Sci. Math. 26 (1965) 125–137
- [5] F. Cabello Sánchez, *Nonlinear centralizers with values in L_0* , Nonlinear Anal. 88 (2014) 42–50.
- [6] F. Cabello, J.M.F. Castillo, *Homological methods in Banach space theory*, Cambridge Studies in Advanced Mathematics 203, Cambridge Univ. Press 2022. Online ISBN 9781108778312.
- [7] F. Cabello Sánchez, R. García, *The twisted Hilbert space ideals*, Integral Equations Operator Theory (2022) 94:21. arXiv:2112.03852v1.
- [8] J.M.F. Castillo, W.H.G. Correa, V. Ferenczi, M. González, *Differential processes generated by two interpolators*, RACSAM 114, paper 183 (2020). <https://doi.org/10.1007/s13398-020-00920-5>
- [9] J.M.F. Castillo, W.H.G. Correa, V. Ferenczi, M. González, *Interpolator symmetries and new Kalton-Peck spaces*, arXiv:2111.10640
- [10] J.M.F. Castillo, W. Cuellar, V. Ferenczi, Y. Moreno, *Complex structures on twisted Hilbert spaces*, Israel J. Math. 222 (2017) 787–814.
- [11] J.M.F. Castillo, V. Ferenczi, *Group actions on twisted sums of Banach spaces*, arXiv:2003.09767v2.
- [12] J.M.F. Castillo, M. González, *Three-space problems in Banach space theory*, Lecture Notes in Math. 1667, Springer 1997.
- [13] I.S. Edelstein, B.S. Mityagin, *Homotopy type of linear groups of two classes of Banach spaces*, Funct. Anal. Appl. 4 (1970) 221–231.
- [14] I.C. Gohberg, A.S. Markus, I.A. Fel’dman, *Normally solvable operators and ideals associated with them*. (Russian) Bul. Akad. Științe RSS Moldoven 76 (1960) 51–70. Translation in Amer. Math. Soc. Transl. (2) Vol. 61. Amer. Math. Soc. Providence, R.I. 1967.
- [15] M. González, *The fine spectrum of the Cesàro operator in ℓ_p ($1 < p < \infty$)*, Arch. Math. (Basel) 44 (1985) 355–358.
- [16] M. González, *The perturbation classes problem in Fredholm theory*, Journal of Functional Analysis 200 (2003) 65–73.
- [17] M. González, A. Martínez-Abejón, *Tauberian operators*, Birkhäuser, 2010.
- [18] W.B. Johnson, J. Lindenstrauss, G. Schechtman, *On the relation between several notions of unconditional structure*, Israel J. Math. 37 (1980) 120–128.
- [19] N. J. Kalton, *The space Z_2 viewed as a symplectic Banach space*, in Proc. Research Workshop on Banach space theory (Iowa City 1981), pp. 97–111, Bor-Luh Lin, ed. Univ. of Iowa, 1982.

- [20] N.J. Kalton, *Differentials of complex interpolation processes for Köthe function spaces*, Trans. Amer. Math. Soc. 333 (1992) 479–529.
- [21] N.J. Kalton, N.T. Peck, *Twisted sums of sequence spaces and three space problem*, Trans. Amer. Math. Soc. 225 (1979) 1–30.
- [22] D. Kleinecke, *Almost-finite, compact, and inessential operators*, Proc. Amer. Math. Soc. 14 (1963) 863–838.
- [23] G. Leibowitz, *Spectra of discrete Cesàro operators*, Tamkang J. Math. 3 (1972) 123–132.
- [24] J. Lindenstrauss, L. Tzafriri, *Classical Banach spaces I, sequence spaces*, Springer, 1977.
- [25] W. Magnus, *On the spectrum of the Hilbert matrix*, Amer. J. Math. 72 (1950) 699–704.
- [26] V.D. Milman, *Operators of class C_0 and C_0^** (Russian). Teor. Funkcii Funkcional. Anal. i Priložen 10 (1970) 15–26.
- [27] G. Neubauer, *The homotopy type of the group of automorphisms in the spaces ℓ_p and c_0* , Math. Ann. 174 (1967) 33–40.
- [28] A. Pietsch, *Operator ideals*, North Holland, 1980.
- [29] B.E. Rhoades, *Spectra of some Hausdorff operators*, Acta Sci. Math. 32 (1971) 91–100.
- [30] W. Rudin, *Functional analysis*, 2nd. ed., McGraw–Hill, 1991.
- [31] B. Silberman, *On the spectrum of the Hilbert matrix operator*, Integr. Eq. Oper. Theory (2021) 93:21.
- [32] I.Y. Sneiberg, *Spectral properties of linear operators in interpolation families of Banach spaces*, Math. Issled. 9 (1974) 214–229.
- [33] M. Zarrabi, *On polynomially bounded operators acting on a Banach space*, Journal of Functional Analysis 225 (2005) 147–166.

UNIVERSIDAD DE EXTREMADURA, INSTITUTO DE MATEMÁTICAS IMUEX, E-06011 BADAJOZ, SPAIN.
Email address: castillo@unex.es

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE CANTABRIA, E-39071 SANTANDER, SPAIN.
Email address: manuel.gonzalez@unican.es

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD DE EXTREMADURA, E-06011 BADAJOZ, SPAIN.
Email address: rpino@unex.es