

## Cherenkov radiation in chiral media

E. Barredo and L.F. Urrutia

*High Energy Physics Department, Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Ciudad de México, 04510, México*

In the framework of Carroll-Field-Jackiw electrodynamics we calculate the spectral distribution of the Cherenkov radiation (CHR) produced by a charge moving at constant velocity in a chiral medium. We find zero, one or two Cherenkov angles according to the relation between the velocity of the particle and the refraction index of the medium.

### 1. Introduction

In 1989 Carroll, Field and Jackiw introduced a modifications of  $3 + 1$  Maxwell electrodynamics (CFJ-ED) by adding a Chern-Simons Lagrangian density, which required the presence of a constant four-vector  $\tilde{b}_\mu = (b_0/c, \mathbf{b})$ <sup>1</sup>,

$$\mathcal{L} = -\frac{1}{16\pi}F_{\mu\nu}F^{\mu\nu} - \frac{1}{c}J^\mu A_\mu + \frac{1}{8\pi}\tilde{b}_\mu\tilde{F}^{\mu\nu}A_\nu. \quad (1)$$

Since the CFJ model breaks Lorentz invariance explicitly we recognize it as a CPT-odd contribution of the photon sector of the SME<sup>2</sup>. Remarkably, CFJ-ED describes the electromagnetic response of Weyl semimetals (chiral media) in condensed matter physics<sup>3</sup>. In the Brillouin zone, the LIV coefficients  $b_0$  and  $\mathbf{b}$  account for the separation of the corresponding Weyl nodes in energy and momentum, respectively.<sup>4</sup> CFJ-ED is also a particular case of axion ED which was previously shown to produce reversed CHR in topological insulators.<sup>5</sup>

### 2. CHR in chiral media

Let us consider a charge  $q$  moving at constant velocity  $v$  in the direction parallel to  $\mathbf{b} = (0, 0, b)$ , from  $-\xi$  to  $+\xi$  with  $\xi \rightarrow \infty$  at the end of the calculation. To determine the electromagnetic fields we start from the modified Maxwell equations for CFJ electrodynamics in vacuum and at the end of

the calculation introduce the refraction index  $n$  of the media. In the Lorentz gauge we have

$$\left(\eta^{\mu\nu}\partial^2 - \epsilon^{\mu\nu\rho\sigma}\tilde{b}_\rho\partial_\sigma\right)A_\nu = \frac{4\pi}{c}J^\mu, \quad (2)$$

which describes an infinite chiral vacuum defined by  $\tilde{b}_\mu = (0, 0, 0, b)$ ,  $b > 0$ . The Green's function (GF) of the operator in Eq. (2) is

$$G^{\mu\nu}(\mathbf{x}, \mathbf{x}'; \omega) = - \left[ (k_0^2 + \nabla^2)g^{\mu\nu} + ik_0b\epsilon^{\mu\nu 30} - b\epsilon^{\mu\nu 3i}\partial_i + \tilde{b}^\mu\tilde{b}^\nu \right] f(\mathbf{x}, \mathbf{x}'; \omega),$$

$$f(\mathbf{x}, \mathbf{x}'; \omega) = \frac{i}{8\pi} \int_0^\infty \frac{k_\perp dk_\perp}{k_\parallel} J_0(R_\perp k_\perp) \sum_{\eta=\pm 1} \frac{\eta}{b} \frac{e^{i\sqrt{k_\parallel^2 + \eta b k_\parallel} Z}}{\sqrt{k_\parallel^2 + \eta b k_\parallel}}, \quad (3)$$

Here we have  $k_0 = \omega/c$ ,  $k_\parallel = +\sqrt{k_0^2 - k_\perp^2}$ ,  $Z = |x_3 - x'_3|$ ,  $\mathbf{R} = \mathbf{x} - \mathbf{x}'$ ,  $R = |\mathbf{R}|$ . The subindex  $\perp$  in a vector indicates its projection on the plane perpendicular to the direction  $x_3$ . The function  $f(\mathbf{x}, \mathbf{x}'; \omega)$  is evaluated by the stationary phase method in the radiation zone. A further approximation to first order in  $bc/\omega$  is introduced to determine the stationary phase point. From now on the refraction index  $n$  of the material is restored. The final result for the electromagnetic potential  $A^\mu(\mathbf{x}, \omega) = \int d^3\mathbf{x}' G^\mu{}_\nu(\mathbf{x}, \mathbf{x}'; \omega) J^\nu(\mathbf{x}', \omega)$  is

$$A^\mu(\mathbf{x}, \omega) = \sum_{\eta=\pm 1} W_\eta^{\mu\nu}(\hat{\mathbf{n}}) \mathcal{J}_\nu(\mathbf{k}_\eta, \omega) \left( \frac{1}{r} e^{ik_0 C_\eta(\theta)r} \right), \quad (4)$$

where  $W_\eta^{\mu\nu}(\hat{\mathbf{n}})$  and  $\mathcal{J}_\nu(\mathbf{k}_\eta, \omega)$  are complicated functions of the observation angles  $\hat{\mathbf{n}}(\theta, \phi)$  that we do not write. For the purpose of discussing the Cherenkov angles is it enough to consider in detail

$$C_\eta(\theta) = \sin^2 \theta + \cos^2 \theta \sqrt{1 + \eta \alpha(n) \sec \theta}, \quad \alpha(n) = bc/(\omega n^2). \quad (5)$$

After calculating  $\mathbf{E}$  and  $\mathbf{B}$  in the radiation zone we obtain the spectral distribution of the radiation

$$\frac{d^2 E}{d\Omega d\omega} = \frac{n\omega^2 q^2}{4\pi^2 c^3} \sum_{\eta=\pm 1} \frac{\sin^2[\xi \tilde{\Xi}_\eta]}{\tilde{\Xi}_\eta^2} \mathcal{U}_\eta, \quad \tilde{\Xi}_\eta(\theta) = \frac{\omega}{v} \left( 1 - \frac{nv}{c} \tilde{C}_\eta(\theta) \cos \theta \right). \quad (6)$$

Again, the function  $\mathcal{U}_\eta$  is not relevant for our discussion. The main point in Eq. (6) is the property

$$\lim_{\xi \rightarrow \infty} \frac{\sin[\xi \tilde{\Xi}_\eta]}{\tilde{\Xi}_\eta} = \pi \delta \left( \frac{\omega}{v} \left( 1 - \frac{nv}{c} \cos \theta C_\eta(\theta) \right) \right), \quad (7)$$

yielding the condition  $H_\eta(\theta) \equiv \cos \theta C_\eta(\omega, \theta) = c/nv$ , which determines the Cherenkov angles as the intersection of the curves  $H_\eta(\theta)$  with the horizontal




Fig. 1. Plot of the functions  $H_{\text{clas}}(\theta)$  (dashed line),  $H_+(\theta)$  (solid line) and  $H_-(\theta)$  (dash-dotted line), for  $\alpha(n) = 0.5$ ,  $n = 2$ .  $H_{\text{clas}}$  corresponds to  $b = 0$  and it is shown for comparison. The horizontal lines labeled by  $u = v/c$  correspond to  $H = 1/(nu)$  in the ordinate.

lines  $c/nv$  for a given material and particle velocity. Figure (1) shows the appearance of a different number of Cherenkov angles according to the particle velocity. The lower limit in the horizontal lines is  $1/n$  corresponding to  $u = 1$ . When  $\alpha(n) < 1$ , the curve  $H_-(\theta)$  ranges over  $0 < \theta < \arccos \alpha(n)$  and it is forbidden otherwise. A similar plot results in the case of vacuum CHR ( $n = 1, b \neq 0$ ).<sup>6</sup>

### Acknowledgments

The authors acknowledge support from the project CF/2019/428214.

### References

1. S.M. Carroll, G.B. Field, and R. Jackiw, Phys. Rev. D **41**, 1231 (1990).
2. D. Colladay, and V.A. Kostelecky Phys. Rev. D **58**, 116002 (1998).
3. A.A. Zyuzin and A.A. Burkov, Phys. Rev. B **86**, 115133 (2012).
4. M.M. Vazifeh and M. Franz, Phys. Rev. Lett. **111** 027201 (2013).
5. O.J. Franca, L.F. Urrutia and O. Rodríguez-Tzompantzi, Phys. Rev. D **99**, 116020 (2019).
6. R. Lehnert and R. Potting, Phys. Rev. Lett. **93** 110402 (2014).