## FIEDLER LINEARIZATIONS OF MULTIVARIABLE STATE-SPACE SYSTEM AND ITS ASSOCIATED SYSTEM MATRIX

NAMITA BEHERA \* AND AVISEK BIST †

Abstract. Linearization is a standard method in the computation of eigenvalues and eigenvectors of matrix polynomials. In the last decade a variety of linearization methods have been developed in order to deal with algebraic structures and in order to construct efficient numerical methods. An important source of linearizations for matrix polynomials are the so called *Fiedler pencils*, which are generalizations of the Frobenius companion form and these linearizations have been extended to regular rational matrix function which is the transfer function of LTI State-space system in [1, 6]. We consider a multivariable state-space system and its associated system matrix  $S(\lambda)$ . We introduce Fiedler pencils of  $S(\lambda)$  and describe an algorithm for their construction. We show that Fiedler pencils are linearizations of the system matrix  $S(\lambda)$ .

Key words. rational matrix valued function, matrix polynomial, linearization, linearization, Rosenbrock system matrix.

AMS subject classifications. 65F15, 15A21, 65L80, 65L05, 34A30.

1. Introduction. We denote by  $\mathbb{C}[\lambda]$  the polynomial ring over the complex field  $\mathbb{C}$ . Further, we denote by  $\mathbb{C}^{m \times n}$  and  $\mathbb{C}[\lambda]^{m \times n}$ , respectively, the vector spaces of  $m \times n$  matrices and matrix polynomials over  $\mathbb{C}$ .

Consider a matrix polynomial  $P(\lambda) = \sum_{j=0}^{m} \lambda^j A_j$ ,  $A_j \in \mathbb{C}^{n \times n}$ . Then a matrix polynomial  $P(\lambda)$  is said to be regular if  $\det(P(\lambda)) \neq 0$  for some  $\lambda \in \mathbb{C}$ . Linearization is a standard method for solving polynomial eigenvalue problems  $P(\lambda)x = 0$ . Let  $P(\lambda)$ be an  $n \times n$  matrix polynomial (regular or singular) of degree m. Then an  $mn \times mn$ matrix pencil  $L(\lambda) := X + \lambda Y$  is said to be a *linearization* [12] of  $P(\lambda)$  if there are  $mn \times mn$  unimodular matrix polynomials  $U(\lambda)$  (the determinant of  $U(\lambda)$ , is a nonzero constant for all  $\lambda \in \mathbb{C}$ .) and  $V(\lambda)$  such that

$$U(\lambda)L(\lambda)V(\lambda) = \operatorname{diag}(I_{(m-1)n}, P(\lambda))$$

for all  $\lambda \in \mathbb{C}$ , where  $I_k$  denotes the  $k \times k$  identity matrix. Linearizations of matrix polynomials have been studied extensively over the years, see [12, 15] and references therein. Recently, a new family of linearizations of matrix polynomials referred to as Fiedler linearizations (or *Fiedler pencils*) has been introduced and is an active area of research, see [5, 11] and references therein. One of the distinctive features of a Fiedler pencil  $L(\lambda)$  of the matrix polynomial  $P(\lambda)$  is that its construction is operation free, that is, block entries of  $L(\lambda)$  are either 0 or  $I_n$  or the coefficient matrices of  $P(\lambda)$ , and that  $L(\lambda)$  allows an easy (operation free) recovery of eigenvectors of  $P(\lambda)$  from the eigenvectors of  $L(\lambda)$  [11, 5].

In this paper we extend the concept of Fiedler linearization from LTI state-space system to general multivariable state-space system and associated system matrix. In particular, in this paper we discuss the solution (finding *eigenvalues*  $\lambda \in \mathbb{C}$  and

<sup>\*</sup>Department of Mathematics, Sikkim University, Sikkim-737102, IN-DIA,(niku.namita@gmail.com, nbehera@cus.ac.in)

<sup>&</sup>lt;sup>†</sup>Department of Mathematics, Sikkim University, Sikkim-737102, INDIA, (abist.21pdmt01@sikkimuniversity.ac.in, avisek.bista@gmail.com)

eigenvectors  $v \in \mathbb{C}^n$ ) of multivariable state-space system  $\Sigma$ 

$$A\left(\frac{d}{dt}\right)x(t) = Bu(t),$$

$$y(t) = Cx(t) + D\left(\frac{d}{dt}\right)u(t) \quad t \ge 0,$$
(1.1)

such that  $\mathcal{S}(\lambda)v = 0$ , where  $A(\lambda) = \sum_{j=0}^{d_A} \lambda^j A_j \in \mathbb{C}[\lambda]^{n \times n}$  is a regular matrix polynomial of degree  $d_A$ ,  $D(\lambda) = \sum_{j=0}^{d_D} \lambda^j D_j \in \mathbb{C}[\lambda]^{m \times m}$  is a matrix polynomial of degree  $d_D$ , and  $C \in \mathbb{C}^{m \times n}$ ,  $B \in \mathbb{C}^{n \times m}$ , and its associate Rosenbrock system matrix  $\mathcal{S}(\lambda)$ ,

$$S(\lambda) = \left[\begin{array}{c|c} A(\lambda) & -B \\ \hline C & D(\lambda) \end{array}\right] \mathbb{C}[\lambda]^{n+m,n+m}$$
(1.2)

and the associated transfer function

$$R(\lambda) = D(\lambda) + CA(\lambda)^{-1}B \in \mathbb{C}(\lambda)^{m,m}.$$
(1.3)

Next, consider a more general linear multivariable time invariant state-space system  $\Sigma_1$  on the positive half line  $\mathbb{R}_+$  in the representation

$$0 = A\left(\frac{d}{dt}\right)x(t) + B\left(\frac{d}{dt}\right)u(t),$$
  
$$y(t) = C\left(\frac{d}{dt}\right)x(t) + D\left(\frac{d}{dt}\right)u(t).$$
 (1.4)

The function  $u : \mathbb{R}^+ \to \mathbb{R}^m$  is the input vector,  $x : \mathbb{R}_+ \to \mathbb{R}^r$  is the state vector,  $y : \mathbb{R}_+ \to \mathbb{R}^p$  is the output vector, and for  $M(\lambda) = \sum_{i=0}^d M_i \lambda^i \in \mathbb{C}[\lambda]^{p,p}$  we use  $M(\frac{d}{dt})$  to denote the differential operator  $\sum_{i=0}^{\ell} M_i \frac{d^i}{dt^i}$ , where  $\frac{d}{dt}$  denotes time-differentiation. The associated matrix polynomial is

$$S(\lambda) := \begin{bmatrix} A(\lambda) & B(\lambda) \\ \hline C(\lambda) & D(\lambda) \end{bmatrix} \in \mathbb{C}[\lambda]^{(p+m),(p+m)}.$$
(1.5)

The associate transfer function is defined by

$$R(\lambda) := D(\lambda) - C(\lambda)A(\lambda)^{-1}B(\lambda) \in \mathbb{C}(\lambda)^{p,p},$$
(1.6)

where, denoting by  $\mathbb{C}[\lambda]^{p,p}$  the vector space of  $p \times p$  matrix polynomials, we assume that  $A(\lambda) \in \mathbb{C}[\lambda]^{m,m}, B(\lambda) \in \mathbb{C}[\lambda]^{m,p}, C(\lambda) \in \mathbb{C}[\lambda]^{p,m}, D(\lambda) \in \mathbb{C}[\lambda]^{p,p}.$ 

Notice that in (1.2) we consider B and C are constant matrices.

Rational eigenvalue problems arise in many applications, see e.g. [10, 18, 19, 22] and the references therein. Rational matrix value functions of this form arise e.g. in linear system theory, see e.g. [21].

If  $A(\lambda)$  is regular, i.e., det  $A(\lambda)$  does not vanish identically, then performing a Schur complement, one obtains the rational matrix function (1.6) which, in frequency domain, describes the *transfer function* from the Laplace transformed input to the Laplace transformed output of the system. In this case  $S(\lambda)$  is called a *Rosenbrock* system matrix, see [20].

Conversely, if one has a given rational matric function of the form (1.6), then one can always interpret it as originating from a Rosenbrock system matrix of the form (1.5). Such rational matrix valued functions arise from realizations of input-output data, see e.g. [16], or in model order reduction, see e.g. [4, 13].

We consider the general square polynomial eigenvalue problem

$$S(\lambda_0) \begin{bmatrix} x_0 \\ u_0 \end{bmatrix} := \begin{bmatrix} A(\lambda) & B(\lambda) \\ \hline C(\lambda) & D(\lambda) \end{bmatrix} \begin{bmatrix} x_0 \\ u_0 \end{bmatrix} = 0$$
(1.7)

If A and D are square and regular, then one can form the rational function  $R(\lambda)$ as in (1.6) and, since det  $S(\lambda) = \det A(\lambda) \det R(\lambda)$ , it is clear that the eigenvalues of  $S(\lambda)$  are the eigenvalues of A and R combined and the eigenvalues of  $A(\lambda)$  are the poles of R. We restrict ourselves to rational functions of the form (1.6) with regular  $A(\lambda)$  and we assume for simplicity that B, C are constant matrices in  $\lambda$ . All the results can be extended (with a lot of technicalities) to the case that B, C depend on  $\lambda$ .

The system  $\Sigma_1$  given in (1.4) is said to be in *state-space form* if it is given by

$$\begin{aligned} E\dot{x}(t) &= Ax(t) + Bu(t)\\ y(t) &= Cx(t) + P(\lambda)u(t), \end{aligned} \tag{1.8}$$

where  $P(\lambda) \in \mathbb{C}[\lambda]^{n \times n}$  is a matrix polynomial and  $A, E \in \mathbb{C}^{m \times m}$  with E being nonsingular,  $B \in \mathbb{C}^{n \times m}, C \in \mathbb{C}^{m \times n}$  are constant matrices, see [21]. For linear time invariant (LTI) state-space system given in (1.8), there is a state-space framework developed in [1] to study the zeros of LTI system in state space form.

For computing zeros of a linear time-invariant system  $\Sigma$  in state-space form, it has been introduced Fiedler-like pencils and Rosenbrock linearization of the Rosenbrock system polynomial  $S(\lambda)$  associated with  $\Sigma$ . Also, it has shown that the Fiedler-like pencils are Rosenbrock linearizations of the system polynomial  $S(\lambda)$ , see [1, 2, 3, 7]

Next, for the higher order linear time invariant (LTI) state-space system  $\Sigma_1$  given by

$$P\left(\frac{d}{dt}\right)x(t) = Bu(t),$$

$$y(t) = Cx(t) + Du(t),$$
(1.9)

where  $P(\lambda) = \sum_{j=0}^{m} \lambda^j A_j \in \mathbb{C}[\lambda]^{n \times n}$  is regular matrix polynomial of degree m and  $D \in \mathbb{C}^{r \times r}, C \in \mathbb{C}^{r \times n}, B \in \mathbb{C}^{n \times r}$ , there is a state-space framework developed in [8] to study the zeros of  $\Sigma_1$ . To study the eigenvalues and eigenvectors of the system matrix associated with  $\Sigma_1$  recently, it has been introduced Fiedler-like pencils and Rosenbrock linearization system polynomial associated with  $\Sigma_1$ . Also, it has shown that the Fiedler-like pencils are Rosenbrock linearizations of the system polynomial, see [8]

In this paper we study the relationship between the eigenvalues and eigenvectors of a rational eigenvalue problem given in the form of a transfer function (1.6), its polynomial representation as a Rosenbrock matrix and associated linearizations. we introduce Fiedler linearizations of the system matrix  $S(\lambda)$  given in (1.2) which is also helpful to study zeros of the system  $\Sigma$  given in (1.1). This problem has recently has been studied for higher order state-space system in [8] and we will extend these results to the Multivariable state-space case.

The paper is organized as follows. In section 2 we recall the definition and some properties of matrix which we need throughout this paper. In section 3 we extend

the results of Fiedler pencils for Rosenbrock system from [1, 6] to multivariable state space system. That is, we show that the state-space framework so developed in [1] could be gainfully used to linearize (Fiedler linearizations) a multivariable state-space system. In particular, we define Fiedler pencils for  $S(\lambda)$  given in (1.2) and describe an algorithm for their construction. In Section 4 we prove that these are linearizations for associate system matrix  $S(\lambda)$ .

**Notation.** An  $m \times n$  rational matrix function  $R(\lambda)$  is an  $m \times n$  matrix whose entries are rational functions of the form  $p(\lambda)/q(\lambda)$ , where  $p(\lambda)$  and  $q(\lambda)$  are scalar polynomials in  $\mathbb{C}[\lambda]$ . We denote the *j*-th column of the  $n \times n$  identity matrix  $I_n$  by  $e_j$ and the transpose of a matrix A by  $A^T$ .

**2. Basic Concepts.** DEFINITION 2.1. [17] Let  $A \in \mathbb{C}^{m \times n}$ ,  $B \in \mathbb{C}^{p \times q}$ . Then the Kronecker product (tensor product) of A and B is defined by

$$A \otimes B = \begin{bmatrix} a_{11}B & \cdots & a_{1n}B \\ \vdots & \ddots & \vdots \\ a_{m1}B & \cdots & a_{mm}B \end{bmatrix} \in \mathbb{C}^{mp \times nq}$$

One of the properties of Kronecker product is as follows: Let  $A \in \mathbb{C}^{m \times n}, B \in \mathbb{C}^{r \times s}, C \in \mathbb{C}^{n \times p}, D \in \mathbb{C}^{s \times t}$ . Then  $(A \otimes B)(C \otimes D) = (AC \otimes BD) \in \mathbb{C}^{mr \times pt}$ .

In order to systematically generate the Fiedler linearizations for Rosenbrock system matrices, we need a few concepts introduced in [11].

DEFINITION 2.2. Let  $\sigma : \{0, 1, \dots, p-1\} \rightarrow \{1, 2, \dots, p\}$  be a bijection.

(1) For j = 0, ..., p-2, the bijection is said to have a consecution at j if  $\sigma(j) < \sigma(j+1)$  and  $\sigma$  has an inversion at j if  $\sigma(j) > \sigma(j+1)$ .

- (2) The tuple CISS(σ) := (c<sub>1</sub>, i<sub>1</sub>, c<sub>2</sub>, i<sub>2</sub>, ..., c<sub>l</sub>, i<sub>l</sub>) is called the consecution-inversion structure sequence of σ, where σ has c<sub>1</sub> consecutive consecutions at 0, 1, ..., c<sub>1</sub>-1; i<sub>1</sub> consecutive inversions at c<sub>1</sub>, c<sub>1</sub> + 1, ..., c<sub>1</sub> + i<sub>1</sub> 1 and so on, up to i<sub>l</sub> inversions at p 1 i<sub>l</sub>, ..., p 2.
- (3) The total number of consecutions and inversions in  $\sigma$  is denoted by  $c(\sigma)$  and  $i(\sigma)$ , respectively, i.e.,  $c(\sigma) = \sum_{j=1}^{l} c_j$ ,  $i(\sigma) = \sum_{j=1}^{l} i_j$ , and  $c(\sigma) + i(\sigma) = p 1$ . DEFINITION 2.3. [11] Let  $P(\lambda) = A_0 + \lambda A_1 + \dots + \lambda^m A_m$  be a matrix polyno-

DEFINITION 2.3. [11] Let  $P(\lambda) = A_0 + \lambda A_1 + \cdots + \lambda^m A_m$  be a matrix polynomial of degree m. For  $k = 0, \ldots m$ , the degree k Horner shift of  $P(\lambda)$  is the matrix polynomial  $P_k(\lambda) := A_{m-k} + \lambda A_{m-k+1} + \cdots + \lambda^k A_m$ . These Horner shifts satisfy the following:

$$P_0(\lambda) = A_m, \ P_{k+1}(\lambda) = \lambda P_k(\lambda) + A_{m-k-1}, \ for \ 0 \le k \le m-1, \ P_m(\lambda) = P(\lambda).$$

DEFINITION 2.4. [15] Matrix polynomial  $U(\lambda)$  is said to be unimodular if det  $U(\lambda)$ is a nonzero constant, independent of  $\lambda$ . Two matrix polynomials  $P(\lambda)$  and  $Q(\lambda)$  are said to be equivalent if there exists unimodular matrices  $U(\lambda)$  and  $V(\lambda)$ , such that  $Q(\lambda) = U(\lambda)P(\lambda)V(\lambda)$ . If  $U(\lambda), V(\lambda)$  are constant matrices, then  $P(\lambda)$  and  $Q(\lambda)$  are said to be strictly equivalent.

Let  $X(\lambda) \in \mathbb{C}(\lambda)^{m \times n}$  be a rational matrix function. The normal rank [20] of  $X(\lambda)$  is denoted by nrank(X) and is given by nrank $(X) := \max_{\lambda} \operatorname{rank}(X(\lambda))$  where the maximum is taken over all  $\lambda \in \mathbb{C}$  which are not poles of the entries of  $X(\lambda)$ . If nrank(X) = n = m then  $X(\lambda)$  is said to be regular, otherwise  $X(\lambda)$  is said to be singular.

A complex number  $\lambda$  is said to be an eigenvalue of the system matrix  $S(\lambda)$  if  $\operatorname{rank}(S(\lambda)) < \operatorname{nrank}(S)$ . An eigenvalue  $\lambda$  of  $S(\lambda)$  is called an invariant zero of the system  $\Sigma$ . We denote the set of eigenvalues of  $S(\lambda)$  by  $\operatorname{Sp}(S)$ .

Let  $G(\lambda) \in \mathbb{C}(\lambda)^{m \times n}$  be a rational matrix function and let

$$\mathbf{SM}(G(\lambda)) = \operatorname{diag}\left(\frac{\phi_1(\lambda)}{\psi_1(\lambda)}, \cdots, \frac{\phi_k(\lambda)}{\psi_k(\lambda)}, 0_{m-k,n-k}\right)$$

be the Smith-McMillan form [14, 20] of  $G(\lambda)$ , where the scalar polynomials  $\phi_i(\lambda)$  and  $\psi_i(\lambda)$  are monic, are pairwise coprime and,  $\phi_i(\lambda)$  divides  $\phi_{i+1}(\lambda)$  and  $\psi_{i+1}(\lambda)$  divides  $\psi_i(\lambda)$ , for i = 1, 2, ..., k - 1. The polynomials  $\phi_1(\lambda), ..., \phi_k(\lambda)$  and  $\psi_1(\lambda), ..., \psi_k(\lambda)$  are called *invariant zero polynomials* and *invariant pole polynomials* of  $G(\lambda)$ , respectively. Define

$$\phi_G(\lambda) := \prod_{j=1}^k \phi_j(\lambda) \text{ and } \psi_G(\lambda) := \prod_{j=1}^k \psi_j(\lambda).$$

A complex number  $\lambda$  is said to be a zero of  $G(\lambda)$  if  $\phi_G(\lambda) = 0$  and a complex number  $\lambda$  is said to be a pole of  $G(\lambda)$  if  $\psi_G(\lambda) = 0$ . The **spectrum** Sp(G) of  $G(\lambda)$  is given by Sp(G) := { $\lambda \in \mathbb{C} : \phi_G(\lambda) = 0$ }. That is Sp(G) is the set of zeros of  $G(\lambda)$ , see [1].

3. Fiedler pencils for Rosenbrock system matrix. In this section we define Fiedler pencils for system polynomial  $S(\lambda)$  and describe an algorithm for their construction. Let us consider a Rosenbrock system of the form (1.2) with B, C constant in  $\lambda$ ,

$$S(\lambda) = \begin{bmatrix} A(\lambda) & -B \\ \hline C & D(\lambda) \end{bmatrix} \mathbb{C}[\lambda]^{n+m,n+m}$$
(3.1)

and the associated transfer function

$$R(\lambda) = D(\lambda) + CA(\lambda)^{-1}B \in \mathbb{F}(\lambda)^{m,m}$$

where for  $A(\lambda) = \sum_{i=0}^{d_A} \lambda^i A_i \in \mathbb{C}[\lambda]^{n,n}$  is regular and  $D(\lambda) = \sum_{j=0}^{d_D} \lambda^j D_j \in \mathbb{C}[\lambda]^{m,m}$ . Our aim is to study linearizations of  $R(\lambda)$  and its relation to linearizations of  $S(\lambda)$ .

The most simple way to perform a direct linearization is to consider a first companion form

$$\mathcal{C}_1(\lambda)w := (\lambda X + Y)w = 0, \qquad (3.2)$$

where

$$X := \begin{bmatrix} A_{d_A} & & & & \\ & I_n & & & \\ & & \ddots & & \\ & & & I_n & & \\ & & & I_n & & \\ & & & & I_m & \\ & & & & \ddots & \\ & & & & & I_m \end{bmatrix},$$

$$Y := \begin{bmatrix} A_{d_A-1} & A_{d_A-2} & \cdots & A_0 & 0 & \cdots & 0 & -B \\ -I_n & 0 & \cdots & 0 & 0 & 0 & 0 \\ & \ddots & \ddots & & & \ddots & \vdots \\ & & -I_n & 0 & & & 0 \\ 0 & \cdots & 0 & C & D_{d_D-1} & D_{d_D-2} & \cdots & D_0 \\ & 0 & 0 & -I_m & 0 & \cdots & 0 \\ & & \ddots & \vdots & & \ddots & \ddots & \vdots \\ & & & 0 & & & -I_m & 0 \end{bmatrix},$$

and

$$w = \begin{bmatrix} \lambda^{d_A - 1}(A(\lambda)^{-1})Bx \\ \lambda^{d_A - 2}(A(\lambda)^{-1})Bx \\ \vdots \\ (A(\lambda)^{-1})Bx \\ \hline \lambda^{d_D - 1}x \\ \lambda^{d_D - 2}x \\ \vdots \\ x \end{bmatrix}$$

It is easy to see that if  $\lambda$  is an eigenvalue of  $R(\lambda)$  then  $R(\lambda)x = 0$  if and only if  $C_1(\lambda)w = 0$ 

An important class of linearizations (which include the first companion form (3.2) as special case) that has recently received a lot of attention are the Fielder pencils, [5, 9, 11]. Introducing *Fiedler matrices*  $M_i$ ,  $i = 0, 1, \ldots, d_A$  associated with  $A(\lambda) = \sum_{i=0}^{d_A} \lambda^i A_i \in \mathbb{C}[\lambda]^{n,n}$  of degree  $d_A$ , defined by

$$M_{d_{A}} := \begin{bmatrix} A_{d_{A}} & & \\ & I_{(d_{A}-1)n} \end{bmatrix}, \ M_{0} := \begin{bmatrix} I_{(d_{A}-1)n} & & \\ & -A_{0} \end{bmatrix},$$
$$M_{i} := \begin{bmatrix} I_{(d_{A}-i-1)n} & & \\ & -A_{i} & I_{n} & \\ & & I_{n} & 0 & \\ & & & & I_{(i-1)n} \end{bmatrix}, \ i = 1, \dots, d_{A} - 1,$$
(3.3)

see e.g. [11].

If  $\sigma : \{0, 1, \ldots, m-1\} \rightarrow \{1, 2, \ldots, m\}$  is a bijection, then one furthermore defines the products  $M_{\sigma} := M_{\sigma^{-1}(1)}M_{\sigma^{-1}(2)}\cdots M_{\sigma^{-1}(m)}$ . Note that  $\sigma(i)$  describes the position of the factor  $M_i$  in the product  $M_{\sigma}$ ; i.e.,  $\sigma(i) = j$  means that  $M_i$  is the *j*th factor in the product.

Based on the Fiedler matrices, then for given  $A(\lambda) \in \mathbb{F}[\lambda]^{n,n}$  of degree  $d_A$  and a bijection  $\sigma$ , in [11] the associated *Fiedler pencil* is defined as the  $d_An \times d_An$  matrix pencil

$$L_{\sigma}(\lambda) := \lambda M_{d_A} - M_{\sigma^{-1}(1)} \cdots M_{\sigma^{-1}(d_A)} = \lambda M_{d_A} - M_{\sigma}.$$

$$(3.4)$$

This concept was extended in [1, 2, 3, 7, 6, 8] for square Rosenbrock systems of the state-space form (1.8) and (1.9). In [1] also a multiplication-free algorithm is presented to construct Fiedler pencils for square system polynomials and it is shown

that these Fiedler pencils are linearizations of the system polynomial and as well as of the associated transfer functions under some appropriate conditions.

Extending the definition of [1], based on the idea of the companion like form (3.2), we define  $nd_A \times nd_A$  Fiedler matrices associated with  $A(\lambda) \in \mathbb{F}[\lambda]^{n,n}$  as in (3.3), and Fiedler matrices associated with the matrix polynomial  $D(\lambda) \in \mathbb{F}[\lambda]^{m,m}$  by

$$N_{d_D} := \begin{bmatrix} D_{d_D} & & \\ & I_{(d_D-1)m} \end{bmatrix}, \ N_0 := \begin{bmatrix} I_{(d_D-1)m} & & \\ & -D_0 \end{bmatrix},$$
$$N_i := \begin{bmatrix} I_{(d_D-i-1)m} & & \\ & -D_i & I_p & \\ & I_m & 0 & \\ & & & I_{(i-1)m} \end{bmatrix}, \ i = 1, \dots, d_D - 1.$$
(3.5)

Based on the Fiedler matrices, then for given  $D(\lambda) \in \mathbb{F}[\lambda]^{m,m}$  of degree  $d_D$  and a bijection  $\sigma$ , in [11] the associated *Fiedler pencil* is defined as the  $d_Dm \times d_Dm$  matrix pencil

$$T_{\sigma}(\lambda) := \lambda N_{d_D} - N_{\sigma^{-1}(1)} \cdots N_{\sigma^{-1}(d_D)} = \lambda N_{d_D} - N_{\sigma}.$$

$$(3.6)$$

Note that  $M_iM_j = M_jM_i$ ,  $N_iN_j = N_jN_i$  for |i-j| > 1 and except for the terms with index 0,  $d_A$  and  $d_D$ , respectively, each  $M_i$  and  $N_i$  is invertible. We then have the following definition of Fiedler matrices for Rosenbrock matrices  $S(\lambda) \in \mathbb{C}[\lambda]^{n+m,n+m}$  given in (1.2).

DEFINITION 3.1. Consider a system polynomial  $S(\lambda)$  as in (1.2) and let  $d = \max\{d_A, d_D\}, r = \min\{d_A, d_D\}$ . Define  $(d_An + d_Dm) \times (d_An + d_Dm)$  matrices  $\mathbb{M}_0, \ldots, \mathbb{M}_d$  by

$$\begin{split} \mathbb{M}_{i} &:= \begin{bmatrix} I_{(d_{A}-i-1)n} & & & \\ & -A_{i} & I_{n} & \\ & I_{n} & 0 & \\ \hline & & I_{(i-1)n} & \\ \hline & & I_{d_{D}m} \end{bmatrix} \\ &= \begin{bmatrix} M_{i} & & \\ \hline & I_{d_{D}m} \end{bmatrix}, \ i = r, r+1, \dots, d_{A}-1, \ \text{if} \ d_{D} < d_{A}, \\ \mathbb{M}_{i} &:= \begin{bmatrix} I_{d_{A}n} & & \\ & I_{(d_{D}-i-1)m} & \\ & & -D_{i} & I_{p} & \\ & & I_{(i-1)m} \end{bmatrix} \\ &= \begin{bmatrix} I_{d_{A}n} & & \\ \hline & & N_{i} \end{bmatrix}, \ i = d_{A}, d_{A}+1, \dots, d_{D}-1, \ \text{if} \ d_{D} > d_{A}. \end{split}$$

Observe that as in [1] one has  $\mathbb{M}_i\mathbb{M}_j = \mathbb{M}_j\mathbb{M}_i$  for |i-j| > 1 and all  $\mathbb{M}_i$  (except possibly  $\mathbb{M}_0$ ,  $\mathbb{M}_d$ ) are invertible.

The associated *Fiedler pencils* are then defined as follows.

DEFINITION 3.2. Consider a system polynomial  $S(\lambda)$  as in (1.2) and let  $d = \max\{d_A, d_D\}$ ,  $r = \min\{d_A, d_D\}$ . Let  $\mathbb{M}_0, \ldots, \mathbb{M}_d$  be Fiedler matrices associated with  $S(\lambda)$  as in Definition 3.1. Given any bijection  $\sigma : \{0, 1, \ldots, d-1\} \rightarrow \{1, 2, \ldots, d\}$ , the matrix pencil

$$\mathbb{L}_{\sigma}(\lambda) := \lambda \mathbb{M}_d - \mathbb{M}_{\sigma^{-1}(1)} \cdots \mathbb{M}_{\sigma^{-1}(m)} =: \lambda \mathbb{M}_d - \mathbb{M}_{\sigma}.$$
(3.7)

is called the Fiedler pencil of  $S(\lambda)$  associated with  $\sigma$ . We also refer to  $\mathbb{L}_{\sigma}(\lambda)$  as a Fiedler pencil of  $R(\lambda)$ .

The companion like form given in (3.2), then is

$$\mathcal{C}_1(\lambda) = \lambda \mathbb{M}_d - \mathbb{M}_{d-1} \cdots \mathbb{M}_1 \mathbb{M}_0$$
8

and

and the associated second companion form of  $\mathcal{S}(\lambda)$  is

$$C_{2}(\lambda) = \lambda \mathbb{M}_{d} + \mathbb{M}_{0}\mathbb{M}_{1} \cdots \mathbb{M}_{d-2}\mathbb{M}_{d-1}$$

$$= \lambda \begin{bmatrix} A_{d_{A}} & & & & \\ & I_{n} & & & \\ & & \ddots & & \\ & & I_{n} & & \\ & & & I_{m} & \\ & & & \ddots & \\ & & & & I_{m} \end{bmatrix}$$

$$+ \begin{bmatrix} A_{d_{A}-1} & -I_{n} & & 0 & & \\ & A_{d_{A}-2} & 0 & \ddots & & \\ & \vdots & \ddots & -I_{n} & 0 & & \\ & \vdots & \ddots & & \\ & A_{0} & \cdots & 0 & 0 & -B & 0 & \cdots & 0 \\ \hline 0 & & & & D_{d_{D}} & -I_{m} & \cdots & 0 \\ & \vdots & \ddots & & & D_{d_{D}-1} & & \vdots \\ & 0 & \ddots & & & \vdots & & -I_{m} \\ & 0 & 0 & \cdots & 0 & D_{0} & 0 & \cdots & 0 \end{bmatrix} .$$
(3.9)

EXAMPLE 3.1. Let  $R(\lambda) = A(\lambda) + CD(\lambda)^{-1}B \in \mathbb{C}^{n \times n}$  with  $A(\lambda) = A_0 + \lambda A_1 + \lambda^2 A_2 + \lambda^3 A_3, A_i \in \mathbb{C}^{n \times n}$  be a matrix polynomial of degree 3 and  $D(\lambda) = D_0 + \lambda D_1 + \lambda^2 D_2, D_i \in \mathbb{C}^{m \times m}$  be a matrix polynomial of degree 2. Here  $d_A > d_D$ , r = 2 and d = 3. Let  $\sigma_1 = (1, 3, 2)$  and  $\sigma_2 = (2, 3, 1)$  be bijections from  $\{0, 1, 2\}$  to  $\{1, 2, 3\}$ . Then  $\mathbb{L}_{\sigma_1}(\lambda) = \lambda \mathbb{M}_3 - \mathbb{M}_0 \mathbb{M}_2 \mathbb{M}_1$  and  $\mathbb{L}_{\sigma_2}(\lambda) = \lambda \mathbb{M}_3 - \mathbb{M}_2 \mathbb{M}_0 \mathbb{M}_1$ . Then the Fiedler matrices for  $G(\lambda)$  are given by

$$\mathbb{M}_{0} = \begin{bmatrix} I_{n} & 0 & 0 & 0 & 0 \\ 0 & I_{n} & 0 & 0 & 0 \\ 0 & 0 & -A_{0} & 0 & B \\ \hline 0 & 0 & 0 & I_{m} & 0 \\ 0 & 0 & -C & 0 & -D_{0} \end{bmatrix} \qquad \mathbb{M}_{1} = \begin{bmatrix} I_{n} & 0 & 0 & 0 \\ 0 & -A_{1} & I_{n} & 0 \\ \hline 0 & I_{n} & 0 & 0 \\ \hline & & & & I_{m} & 0 \\ \hline & & & & I_{m} & 0 \end{bmatrix}$$

$$\mathbb{M}_{2} = \begin{bmatrix} -A_{2} & I_{n} & 0 & & \\ I_{n} & 0 & 0 & & \\ 0 & 0 & I_{n} & & \\ & & & I_{m} \end{bmatrix} \qquad \mathbb{M}_{3} = \begin{bmatrix} A_{3} & & & & \\ & I_{n} & & \\ & & & I_{n} & \\ & & & I_{m} \end{bmatrix}.$$

Then

$$\mathbb{M}_{\sigma_1} = \begin{bmatrix} -A_2 & -A_1 & I_n & 0 & 0 \\ I_n & 0 & 0 & 0 & 0 \\ 0 & -A_0 & 0 & B & 0 \\ 0 & 0 & 0 & -D_1 & I_m \\ 0 & -C & 0 & -D_0 & 0 \end{bmatrix}.$$

By using the commutativity relation it is easy to check that  $\mathbb{L}_{\sigma_1}(\lambda) = \mathbb{L}_{\sigma_2}(\lambda)$ .

EXAMPLE 3.2. Let  $G(\lambda) = A(\lambda) + CD(\lambda)^{-1}B \in \mathbb{C}^{n \times n}$  with  $A(\lambda) = A_0 + \lambda A_1 + \lambda^2 A_2 + \lambda^3 A_3$ ,  $A_i \in \mathbb{C}^{n \times n}$  and  $D(\lambda) = D_0 + \lambda D_1 + \lambda^2 D_2 + \lambda^3 D_3 + \lambda^4 D_4$ ,  $D_i \in \mathbb{C}^{m \times m}$ . Here,  $d_A = 3$ ,  $d_D = 4$ ,  $d_A < d_D$  and r = 3, d = 4. Consider  $\mathbb{L}_{\sigma}(\lambda) = \lambda \mathbb{M}_4 - \mathbb{M}_2 \mathbb{M}_0 \mathbb{M}_1 \mathbb{M}_3$ . Then the Fiedler matrices for  $G(\lambda)$  are given by

$$\mathbb{M}_0 = \begin{bmatrix} I_n & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & I_n & 0 & 0 & 0 & 0 & 0 & B \\ \hline 0 & 0 & -A_0 & 0 & 0 & I_m & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & I_m & 0 & 0 & 0 \\ 0 & 0 & 0 & -C & 0 & 0 & I_m & 0 & 0 \\ \hline 0 & 0 & 0 & -C & 0 & 0 & 0 & -D_0 \end{bmatrix}, \\ \mathbb{M}_1 = \begin{bmatrix} I_n & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & I_n & 0 & 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_m & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_n & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_n & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_n & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_n & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_n & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_n & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_n & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_n & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_n & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_n & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_m & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_m & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_m & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_m & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_m & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_m & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_m & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_m & 0 \\ \hline 0 & 0 & 0 & 0 & I_m & 0 \\ \hline 0 & 0 & 0 & 0 & I_m & 0 \\ \hline 0 & 0 & 0 & 0 & I_m & 0 \\ \hline 0 & 0 & 0 & 0 & I_m & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & I_m \\ \hline 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_m \\ \hline 0 & 0 & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 \\$$

Then

$$\mathbb{M}_{\sigma} = \begin{bmatrix} -A_2 & -A_1 & I_n & 0 & 0 & 0 & 0 \\ I_n & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & -A_0 & 0 & 0 & 0 & B & 0 \\ \hline 0 & 0 & 0 & -D_3 & I_m & 0 & 0 \\ 0 & 0 & 0 & -D_2 & 0 & -D_1 & I_m \\ 0 & 0 & 0 & I_m & 0 & 0 & 0 \\ 0 & -C & 0 & 0 & 0 & -D_0 & 0 \end{bmatrix}$$

EXAMPLE 3.3. Let  $G(\lambda) = A(\lambda) + CD(\lambda)^{-1}B \in \mathbb{C}^{n \times n}$  with  $A(\lambda) = A_0 + \lambda A_1 + \lambda^2 A_2 + \lambda^3 A_3$  where  $A_i \in \mathbb{C}^{n \times n}$  and  $D(\lambda) = D_0 + \lambda D_1 + \lambda^2 D_2 + \lambda^3 D_3$  where  $D_i \in \mathbb{C}^{m \times m}$ . Here,  $d_A = 3$ ,  $d_D = 3$ , r = 3 and d = 3. Consider  $\mathbb{L}_{\sigma}(\lambda) = \lambda \mathbb{M}_3 - \mathbb{M}_{\sigma} = \lambda \mathbb{M}_3 - \mathbb{M}_2 \mathbb{M}_0 \mathbb{M}_1$ . Then the Fiedler matrices for  $G(\lambda)$  are given by

$$\mathbb{M}_{0} = \begin{bmatrix} I_{n} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & I_{n} & 0 & 0 & 0 & 0 & B \\ \hline 0 & 0 & -A_{0} & 0 & 0 & B & 0 \\ \hline 0 & 0 & 0 & I_{m} & 0 & 0 \\ 0 & 0 & 0 & 0 & I_{m} & 0 & 0 \\ \hline 0 & 0 & 0 & C & 0 & 0 & -D_{0} \end{bmatrix}, \ \mathbb{M}_{1} = \begin{bmatrix} I_{n} & 0 & 0 & 0 & 0 & 0 \\ 0 & -A_{1} & I_{n} & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_{m} & 0 & 0 \\ \hline 0 & 0 & 0 & I_{m} & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_{m} & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_{m} & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_{m} & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_{m} & 0 & 0 \\ \hline 0 & 0 & 0 & I_{n} & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_{n} & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_{n} & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_{n} & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & I_{n} & 0 & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_{m} & 0 & 0 \\ \hline 0 & 0 & 0 & 0 & I_{m} & 0 \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0 & 0 & 0 & 0 & I_{m} \\ \hline 0 & 0$$

Then,

$$\mathbb{M}_{\sigma} = \begin{bmatrix} -A_2 & -A_1 & I_n & 0 & 0 & 0 \\ I_n & 0 & 0 & 0 & 0 & 0 \\ 0 & -A_0 & 0 & 0 & B & 0 \\ \hline 0 & 0 & 0 & -D_2 & -D_1 & I_m \\ 0 & 0 & 0 & I_m & 0 & 0 \\ 0 & -C & 0 & 0 & -D_0 & 0 \\ \end{bmatrix}$$

Having introduced the basic idea of generating Fiedler pencils for Rosenbrock system polynomials given in (1.2), now we will analyze these constructed pencils.

THEOREM 3.3. Let  $S(\lambda)$  be given in (1.2). Let  $d = \max(d_A, d_D)$  and  $\sigma : \{0, 1, \ldots, d-1\} \rightarrow \{1, 2, \ldots, d\}$  be a bijection. Let  $L_{\sigma}(\lambda)$ ,  $T_{\sigma}(\lambda)$  and  $\mathbb{L}_{\sigma}(\lambda)$  be the Fiedler pencils of  $A(\lambda)$  of degree  $d_A$ ,  $D(\lambda)$  of degree  $d_D$  and  $S(\lambda)$ , respectively, associated with  $\sigma$ , that is,  $L_{\sigma}(\lambda) := \lambda M_{d_A} - M_{\sigma}$ ,  $T_{\sigma}(\lambda) := \lambda N_{d_D} - N_{\sigma}$  and  $\mathbb{L}_{\sigma}(\lambda) := \lambda \mathbb{M}_d - \mathbb{M}_{\sigma}$ . If  $\sigma^{-1} = (\sigma_1^{-1}, 0, \sigma_2^{-1})$  for some bijections  $\sigma_1$  and  $\sigma_2$ , then

$$\mathbb{L}_{\sigma}(\lambda) = \begin{bmatrix} L_{\sigma}(\lambda) & -M_{\sigma_1}(e_{d_A}e_{d_D}^T \otimes B)N_{\sigma_2} \\ N_{\sigma_1}(e_{d_D}e_{d_A}^T \otimes C)M_{\sigma_2} & T_{\sigma}(\lambda) \end{bmatrix}.$$

Further, if  $CISS(\sigma) = (c_1, i_1, \ldots, c_l, i_l)$  then

$$\mathbb{L}_{\sigma}(\lambda) = \begin{bmatrix} L_{\sigma}(\lambda) & | -e_{d_A} e_{d_D - c_1}^T \otimes B \\ \hline e_{d_D} e_{d_A - c_1}^T \otimes C & | T_{\sigma}(\lambda) \end{bmatrix}, \quad \text{if } c_1 > 0$$

and

$$\mathbb{L}_{\sigma}(\lambda) = \begin{bmatrix} L_{\sigma}(\lambda) & | -e_{(d_A - i_1)}e_{d_D}^T \otimes B \\ \hline e_{(d_D - i_1)}e_{d_A}^T \otimes C & T_{\sigma}(\lambda) \end{bmatrix}, \quad \text{if } c_1 = 0.$$

Thus the map (Fiedler(A), Fiedler(D))  $\longrightarrow$  Fiedler(S),  $(\lambda M_{d_A} - M_{\sigma}, \lambda N_{d_D} - N_{\sigma}) \mapsto \lambda \mathbb{M}_m - \mathbb{M}_{\sigma}$  is a bijection, where Fiedler(A), Fiedler(D) and Fiedler(S), respectively, denote the set of Fiedler pencils of  $A(\lambda)$ ,  $D(\lambda)$  and  $S(\lambda)$ .

*Proof.* We have  $\mathbb{L}_{\sigma}(\lambda) = \lambda \mathbb{M}_m - \mathbb{M}_{\sigma} = \lambda \mathbb{M}_m - \mathbb{M}_{\sigma_1} \mathbb{M}_0 \mathbb{M}_{\sigma_2}$ 

$$= \lambda \begin{bmatrix} M_{d_A} & 0 \\ 0 & N_{d_D} \end{bmatrix} - \begin{bmatrix} M_{\sigma_1} & 0 \\ 0 & N_{\sigma_1} \end{bmatrix} \begin{bmatrix} M_0 & (e_{d_A} e_{d_D}^T) \otimes B \\ -(e_{d_D} e_{d_A}^T) \otimes C & N_0 \end{bmatrix} \begin{bmatrix} M_{\sigma_2} & 0 \\ 0 & N_{\sigma_2} \end{bmatrix}$$

$$= \lambda \begin{bmatrix} M_m & 0 \\ 0 & N_{d_D} \end{bmatrix} - \begin{bmatrix} M_{\sigma_1} M_0 M_{\sigma_2} & M_{\sigma_1} (e_{d_A} e_{d_D}^T \otimes B) N_{\sigma_2} \\ -N_{\sigma_1} (e_{d_D} e_{d_A}^T \otimes C) M_{\sigma_2} & N_{\sigma_1} N_0 N_{\sigma_2} \end{bmatrix}$$

$$= \begin{bmatrix} L_{\sigma}(\lambda) & -M_{\sigma_1} (e_{d_A} e_{d_D}^T \otimes B) N_{\sigma_2} \\ N_{\sigma_1} (e_{d_D} e_{d_A}^T \otimes C) M_{\sigma_2} & T_{\sigma}(\lambda) \end{bmatrix}.$$

Now suppose that  $CISS(\sigma) = (c_1, i_1, \ldots, c_l, i_l).$ 

Case I: Suppose that  $c_1 > 0$ . Then by commutativity relation we have  $\mathbb{M}_{\sigma} = \mathbb{M}_{\sigma_1}\mathbb{M}_0\mathbb{M}_1\cdots\mathbb{M}_{c_1}$  with  $c_1+1 \in \sigma_1$ . Thus  $\mathbb{M}_{\sigma} = \mathbb{M}_{\sigma_1}\mathbb{M}_0\mathbb{M}_{\sigma_2}$ , where  $\mathbb{M}_{\sigma_2} = \mathbb{M}_1\cdots\mathbb{M}_{c_1}$ . Hence

$$\mathbb{M}_{\sigma} = \begin{bmatrix} M_{\sigma_1} \\ N_{\sigma_1} \end{bmatrix} \begin{bmatrix} M_0 & (e_{d_A} e_{d_D}^T) \otimes B \\ \hline -(e_{d_D} e_{d_A}^T) \otimes C & N_0 \end{bmatrix} \begin{bmatrix} M_{\sigma_2} & 0 \\ \hline 0 & N_{\sigma_2} \end{bmatrix} \\
= \begin{bmatrix} M_{\sigma_1} M_0 M_{\sigma_2} & M_{\sigma_1} (e_{d_A} e_{d_D}^T \otimes B) N_{\sigma_2} \\ \hline -N_{\sigma_1} (e_{d_D} e_{d_A}^T \otimes C) M_{\sigma_2} & N_{\sigma_1} N_0 N_{\sigma_2} \end{bmatrix}.$$

Since  $j \in \sigma_1$  implies that  $j \ge c_1 + 1$ , we have  $M_{\sigma_1} = \begin{bmatrix} * & & \\ & & I_{c_1n} \end{bmatrix}$  and  $N_{\sigma_1} = \begin{bmatrix} * & & \\ & & I_{c_1m} \end{bmatrix}$ . This shows that  $M_{\sigma_1}(e_{d_A} \otimes I_n) = e_{d_A} \otimes I_n$  and  $N_{\sigma_1}(e_{d_D} \otimes I_m) = e_{d_D} \otimes I_m$ . So, we have  $M_{\sigma_1}(e_{d_A} \otimes B) = e_{d_A} \otimes B$ . Next, we have  $N_{\sigma_1}(e_{d_D}e_{d_A}^T \otimes C)M_{\sigma_2} = I_{\sigma_1}$ .  $N_{\sigma_1}(e_{d_D} \otimes I_m)(e_{d_A}^T \otimes C)M_{\sigma_2}$  and  $M_{\sigma_1}(e_{d_A}e_{d_D}^T \otimes B)N_{\sigma_2} = M_{\sigma_1}(e_{d_A} \otimes B)(e_{d_D}^T \otimes I_m)N_{\sigma_2}$ . Now, we have

$$(e_{d_A}^T \otimes I_n) M_1 = (e_{d_A}^T \otimes I_n) \begin{bmatrix} I_{(d_A-2)n} & & \\ & -A_1 & I_n \\ & I_n & 0 \end{bmatrix} = (e_{d_A-1}^T \otimes I_n),$$

$$(e_{d_A}^T \otimes I_n) M_1 M_2 = (e_{d_A-1}^T \otimes I_n) \begin{bmatrix} I_{(d_A-3)n} & & \\ & -A_2 & I_n \\ & I_n & 0 \\ & & & I_n \end{bmatrix} = (e_{d_A-2}^T \otimes I_n),$$

and so on. Thus  $(e_{d_A}^T \otimes I_n)M_1M_2 \cdots M_{c_1} = (e_{d_A-c_1}^T \otimes I_n)$ . Hence  $(e_{d_A}^T \otimes I_n)M_{\sigma_2} = (e_{d_A-c_1}^T \otimes I_n)$  and  $(-(e_{d_A-c_1}^T \otimes C)M_{\sigma_2} = -(e_{d_A-c_1}^T \otimes C)$ . Similarly, we have

$$(e_{d_D}^T \otimes I_m) N_1 = (e_{d_D}^T \otimes I_m) \begin{bmatrix} I_{(d_D - 2)m} & & \\ & -D_1 & I_m \\ & I_m & 0 \end{bmatrix} = (e_{d_D - 1}^T \otimes I_m),$$

$$(e_{d_D}^T \otimes I_m) N_1 N_2 = (e_{d_D - 1}^T \otimes I_m) \begin{bmatrix} I_{(d_D - 3)m} & & \\ & -D_2 & I_m \\ & I_m & 0 \\ & & & I_m \end{bmatrix} = (e_{d_D - 2}^T \otimes I_m)$$

and so on. Thus  $(e_{d_D}^T \otimes I_m)N_1N_2 \cdots N_{c_1} = (e_{d_D-c_1}^T \otimes I_m)$ . Hence  $(e_{d_D}^T \otimes I_m)N_{\sigma_2} = (e_{d_D-c_1}^T \otimes I_m)$ . Now, we have  $N_{\sigma_1}(e_{d_D}e_{d_A}^T \otimes I_n)M_{\sigma_2} = N_{\sigma_1}(e_{d_D} \otimes I_m)(e_{d_A}^T \otimes C)M_{\sigma_2} = (e_{d_D}e_{d_A-c_1}^T \otimes I_n)$  and  $-(N_{\sigma_1}e_{d_D}e_{d_A}^T \otimes C)M_{\sigma_2} = -(e_{d_D}e_{d_A-c_1}^T \otimes C)$ . Similarly,  $M_{\sigma_1}(e_{d_A}e_{d_D}^T \otimes B)N_{\sigma_2} = (e_{d_A}e_{d_D-c_1}^T \otimes B)$ . Consequently, we have

$$\mathbb{L}_{\sigma}(\lambda) = \lambda \mathbb{M}_m - \mathbb{M}_{\sigma} = \begin{bmatrix} L_{\sigma}(\lambda) & -e_{d_A} e_{d_D - c_1}^T \otimes B \\ \hline e_{d_D} e_{d_A - c_1}^T \otimes C & T_{\sigma}(\lambda) \end{bmatrix}.$$

Case II: Suppose that  $c_1 = 0$ . Then  $\sigma$  has  $i_1$  inversions at 0. Hence by commutativity relations we have  $\mathbb{M}_{\sigma} = \mathbb{M}_{i_1} \cdots \mathbb{M}_1 \mathbb{M}_0 \mathbb{M}_{\sigma_2} =: \mathbb{M}_{\sigma_1} \mathbb{M}_0 M_{\sigma_2}$  with  $i_1 + 1 \in \sigma_2$ . Hence

$$\mathbb{M}_{\sigma} = \left[ \begin{array}{c|c} M_{\sigma_1} \\ \hline \\ N_{\sigma_1} \end{array} \right] \left[ \begin{array}{c|c} M_0 \\ \hline \\ -(e_{d_D}e_{d_A}^T) \otimes C \\ \hline \\ N_0 \end{array} \right] \left[ \begin{array}{c|c} M_{\sigma_2} \\ \hline \\ 0 \\ \hline \\ N_{\sigma_2} \end{array} \right] \\
= \left[ \begin{array}{c|c} M_{\sigma_1} M_0 M_{\sigma_2} \\ \hline \\ -N_{\sigma_1} (e_{d_D}e_{d_A}^T \otimes C) M_{\sigma_2} \\ \hline \\ N_{\sigma_1} N_0 N_{\sigma_2} \end{array} \right].$$

Since  $j \in \sigma_2$  implies that  $j \ge i_1 + 1$ , we have  $M_{\sigma_2} = \begin{bmatrix} * & \\ & I_{i_1n} \end{bmatrix}$  and  $N_{\sigma_2} = \begin{bmatrix} * & \\ & I_{i_1m} \end{bmatrix}$ . This shows that  $(e_{d_A}^T \otimes I_n)M_{\sigma_2} = e_{d_A}^T \otimes I_n$  and  $(e_{d_D}^T \otimes I_m)N_{\sigma_2} = e_{d_D}^T \otimes I_m$ .

Hence  $(-e_{d_A}^T \otimes C)M_{\sigma_2} = -e_{d_A}^T \otimes C$ . Next, we have

$$M_{1}(e_{d_{A}} \otimes I_{n}) = \begin{bmatrix} I_{(m-2)n} & & \\ & -A_{1} & I_{n} \\ & I_{n} & 0 \end{bmatrix} (e_{d_{A}} \otimes I_{n}) = (e_{d_{A}-1} \otimes I_{n}),$$
$$M_{2}M_{1}(e_{d_{A}} \otimes I_{n}) = \begin{bmatrix} I_{(d_{A}-3)n} & & \\ & -A_{2} & I_{n} \\ & & I_{n} & 0 \\ & & & & I_{n} \end{bmatrix} (e_{d_{A}-1} \otimes I_{n}) = (e_{d_{A}-2} \otimes I_{n}).$$

Thus  $M_{i_1} \cdots M_2 M_1(e_{d_A} \otimes I_n) = (e_{d_A - i_1} \otimes I_n)$ . Hence  $M_{\sigma_1}(e_{d_A} \otimes I_n) = (e_{(d_A - i_1)} \otimes I_n)$ and  $M_{\sigma_1}(e_{d_A} \otimes B) = (e_{(d_A - i_1)} \otimes B)$ . Similarly, we have

$$N_{1}(e_{d_{D}} \otimes I_{m}) = \begin{bmatrix} I_{(d_{D}-2)m} & & \\ & -D_{1} & I_{m} \\ & I_{m} & 0 \end{bmatrix} (e_{d_{D}} \otimes I_{m}) = (e_{d_{D}-1} \otimes I_{m}),$$

$$N_{2}N_{1}(e_{d_{D}} \otimes I_{m}) = \begin{bmatrix} I_{(d_{D}-3)m} & & \\ & -D_{2} & I_{m} \\ & I_{m} & 0 \\ & & & I_{m} \end{bmatrix} (e_{d_{D}-1} \otimes I_{m}) = (e_{d_{D}-2} \otimes I_{m}).$$

Thus  $N_{i_1} \cdots N_2 N_1(e_{d_D} \otimes I_m) = (e_{d_D - i_1} \otimes I_m)$ . Hence  $N_{\sigma_1}(e_{d_D} \otimes I_m) = (e_{(d_D - i_1)} \otimes I_m)$ . Now, we have  $N_{\sigma_1}(e_{d_D}e_{d_A}^T \otimes I_n)M_{\sigma_2} = N_{\sigma_1}(e_{d_D} \otimes I_m)(e_{d_A}^T \otimes C)M_{\sigma_2} = (e_{d_D - i_1}e_{d_A}^T \otimes I_n)$ and  $-(N_{\sigma_1}e_{d_D}e_{d_A}^T \otimes C)M_{\sigma_2} = -(e_{d_D - i_1}e_{d_A}^T \otimes C)$ . Similarly,  $M_{\sigma_1}(e_{d_A}e_{d_D}^T \otimes B)N_{\sigma_2} = (e_{d_A - i_1}e_{d_D}^T \otimes B)$ . Consequently, we have

$$\mathbb{L}_{\sigma}(\lambda) = \lambda \mathbb{M}_{m} - \mathbb{M}_{\sigma} = \begin{bmatrix} L_{\sigma}(\lambda) & -e_{(d_{A}-i_{1})}e_{d_{D}}^{T} \otimes B \\ \hline e_{(d_{D}-i_{1})}e_{d_{A}}^{T} \otimes C & T_{\sigma}(\lambda) \end{bmatrix}$$

Note that for each  $i, j \in \sigma$ , we have  $M_i M_j = M_j M_i$ ,  $N_i N_j = N_j N_i \Leftrightarrow \mathbb{M}_i \mathbb{M}_j = \mathbb{M}_j \mathbb{M}_i$ . Hence it follows that  $\#(\operatorname{Fiedler}(A), \operatorname{Fiedler}(D)) = \#(\operatorname{Fiedler}(S))$ . This completes the proof.  $\square$ 

THEOREM 3.4. Let  $S(\lambda)$  be in (1.2) with  $A(\lambda) = \sum_{i=0}^{d_A} \lambda^i A_i, A_i \in \mathbb{C}^{n \times n}, \sum_{i=0}^{d_D} \lambda^i D_i, D_i \in \mathbb{C}^{m \times m}$ . Suppose that  $d_A > d_D$ . Let  $\sigma : \{0, 1, \ldots, d_A - 1\} \rightarrow \{1, 2, \ldots, d_A\}$  be a bijection. The following algorithm constructs a sequence of matrices  $\{\mathbb{W}_0, \mathbb{W}_1, \ldots, \mathbb{W}_{d_A-2}\}$ , where each matrix  $\mathbb{W}_i$  for  $i = 1, 2, \ldots, d_A - 2$  is partitioned into blocks in such a way that the blocks of  $\mathbb{W}_{i-1}$  are blocks of  $W_i$ .

**Algorithm 1** Construction of  $\mathbb{M}_{\sigma}$  for  $\mathbb{L}_{\sigma}(\lambda) := \lambda \mathbb{M}_{d_A} - \mathbb{M}_{\sigma}$ .

**Input:**  $S(\lambda) = \begin{bmatrix} \frac{d_A}{\sum_{i=0}^{d} \lambda^i A_i} & -B \\ \hline C & \sum_{i=0}^{d_D} \lambda^i D_i \end{bmatrix}$  and a bijection  $\sigma$  :  $\{0, 1, \dots, d_A - 1\} \rightarrow$ 

 $\{1, 2, \ldots, d_A\}.$ Output:  $\mathbb{M}_{\sigma}$ 

 $\begin{aligned} \mathbf{if} \ \sigma \ \text{has a consecution at } 0 \ \mathbf{then} \\ \mathbb{W}_0 &:= \begin{bmatrix} -A_1 & I_n & 0 & 0 \\ -A_0 & 0 & B & 0 \\ 0 & 0 & -D_1 & I_m \\ -C & 0 & -D_0 & 0 \end{bmatrix} \\ \mathbf{else} \\ \mathbb{W}_0 &:= \begin{bmatrix} -A_1 & -A_0 & 0 & B \\ I_n & 0 & 0 & 0 \\ 0 & 0 & -C & -D_1 & -D_0 \\ 0 & 0 & I_m & 0 \end{bmatrix} \\ \mathbf{end} \ \mathbf{if} \end{aligned}$ 

If  $d_A > d_D$ 

for  $i = 1 : d_D - 2$  do

$$\begin{split} & \text{if } \sigma \text{ has a consecution at } i \text{ then}} \\ & \mathbb{W}_i := \begin{bmatrix} \frac{-A_{i+1}}{W_{i-1}(1:i+1,1)} & 0 & \mathbb{W}_{i-1}(1:i+1,2:i+1) & \mathbb{W}_{i2}}{0} \\ \hline \mathbb{W}_{i-1}(3:i+3,1) & 0 & \mathbb{W}_{i-1}(3:i+3,2:i+1) & \mathbb{W}_{i4} \end{bmatrix}, \text{ where} \\ & \mathbb{W}_{i2} = \begin{bmatrix} 0 & 0 & 0 \\ \mathbb{W}_{i-1}(1:i+1,i+2) & 0 & \mathbb{W}_{i-1}(1:i+1,i+3:2i+2) \end{bmatrix}, \\ & \mathbb{W}_{i4} = \begin{bmatrix} -D_{i+1} & I_m & 0 \\ \mathbb{W}_{i-1}(3:i+3,i+2) & 0 & \mathbb{W}_{i-1}(3:i+3,i+3:2i+2) \end{bmatrix}. \\ & \text{else} \\ & \mathbb{W}_i := \begin{bmatrix} -A_{i+1} & \mathbb{W}_{i-1}(1,1:i+1) & 0 & \mathbb{W}_{i-1}(1,3:i+3) \\ 0 & \mathbb{W}_{i-1}(2:i+1,1:i+1) & 0 & \mathbb{W}_{i-1}(2:i+1,3:i+3) \\ \hline 0 & \mathbb{W}_{i-1}(i+2,1:i+1) & -D_{i+1} & \mathbb{W}_{i-1}(i+2,3:i+3) \\ & 0 & \mathbb{W}_{i-1}(i+3:2i+2,1:i+1) & 0 & \mathbb{W}_{i-1}(i+3:2i+2,3:i+3) \\ \end{bmatrix} \\ & \text{end if} \end{split}$$

end for for  $i = d_D - 1 : d_A - 2$  do if  $\sigma$  has a consecution at i then  $W_i := \begin{bmatrix} -A_{i+1} & I_n & 0 & 0 \\ W_{i-1}(:, 1) & 0 & W_{i-1}(:, 2:i+1) & W_{i-1}(:, i+2:2i+2) \end{bmatrix}$ else  $W_i := \begin{bmatrix} -A_{i+1} & W_{i-1}(1, :) \\ I_n & 0 & 0 \\ 0 & W_{i-1}(2:i+1, :) \\ 0 & W_{i-1}(i+2:2i+2, :) \end{bmatrix}$ end if end for  $M_{\sigma} := W_{d_A-2}$ 

*Proof.* We prove the result by induction on the degree  $d_A = \max(d_A, d_D)$ . Then the rest of the proof follows directly from proof of Theorem 3.11 in [1].  $\Box$ 

THEOREM 3.5. Let  $S(\lambda)$  be in (1.2) with  $A(\lambda) = \sum_{i=0}^{d_A} \lambda^i A_i, A_i \in \mathbb{C}^{n \times n}, \sum_{i=0}^{d_D} \lambda^i D_i, D_i \in \mathbb{C}^{m \times m}$ . Suppose that  $d_A < d_D$ . Let  $\sigma : \{0, 1, \dots, d_D - 1\} \rightarrow \{1, 2, \dots, d_D\}$  be a bijection. The following algorithm constructs a sequence of matrices  $\{\mathbb{W}_0, \mathbb{W}_1, \dots, \mathbb{W}_{d_D-2}\}$ , where each matrix  $\mathbb{W}_i$  for  $i = 1, 2, \dots, d_D - 2$  is partitioned into blocks in such a way

that the blocks of  $\mathbb{W}_{i-1}$  are blocks of  $W_i$ .

Algorithm 2 Construction of  $\mathbb{M}_{\sigma}$  for  $\mathbb{L}_{\sigma}(\lambda) := \lambda \mathbb{M}_{d_{D}} - \mathbb{M}_{\sigma}$ .  $\frac{-B}{\sum_{i=0}^{d_D} \lambda^i D_i}$  $\underline{\lambda_{i=0}^{a_A}} \lambda^i A_i$ Input:  $S(\lambda) =$ and a bijection  $\sigma: \{0, 1, \ldots, d_D - 1\} \rightarrow$  $\{1, 2, \ldots, d_D\}$ Output:  $\mathbb{M}_{\sigma}$ if  $\sigma$  has a consecution at 0 then  $-A_1$  $I_n \\ 0$ B0  $-D_1$ else 0 0 0  $W_0 =$ end if If  $d_D > d_A$ for  $i = 1: d_A - 2$  do if  $\sigma$  has a consecution at i then  $-A_{i+1}$  $\mathbb{W}_{i-1}(\underline{1}:i+1,1)$ 0  $W_{i-1}(1:i+1,2:i)$ , where  $\mathbb{W}_i :=$ 0 0  $\mathbb{W}_{i-1}(3:i+3,1)$  $W_{i4}$ 0 W. -1(3:i+3,2:i0 0 0  $W_{i2} =$  $\mathbb{N}_{i-1}(1:i+1,i+2)$ (1:i+1,i+3:2i+2)WV. 0  $-D_{i+1}I_m$ 0 0  $W_{i4} =$ (3:i+3,i)+2)0  $\mathbb{W}_i$  $_{1}(3:i+3,i+3:2i+2)$ else 0 0  $\mathbb{W}_{i-1}(2:i+1,1:i+1)$ 0  $\mathbb{W}_i =$  $\mathbb{W}_{i-1}(i+2,1:i+1)$  $D_{i+1}$  $\mathbb{W}_{i-1}(i+$  $I_m$ +3:2i+2.1:i+1 $\mathbb{W}_{i-1}(i+3:2i+2,3:i+3)$ end if end for for  $i = d_A - 1 : d_D - 2$  do if  $\sigma$  has a consecution at *i* then  $W_{i-1}(1:i+1)$  $W_i =$ 1 m 0  $\mathbb{W}_{i-1}(i+2:2i+2,i+2)$  $\mathbb{W}_{i-1}(i+2:2i+2,2i)$ 1(i+2:2i+2,i:else  $\frac{\mathbb{W}_{i-1}(1:i+1,i+2:2i+2)}{\mathbb{W}_{i-1}(i+2,i+2:2i+2)}_0$ 0  $\begin{array}{c} -D_{i+1} \\ I_m \\ 0 \end{array}$ 0 0  $\mathbb{W}_i =$  $\mathbb{W}_{i-1}(2i+1:2i+2,i+2:2i+2)$  $\mathbb{W}_{i-1}(2i+1:2i+2,1:i+1)$ end if end for  $\mathbb{M}_{\sigma} := \mathbb{W}_{d_D-2}$ 

*Proof.* We prove the result by induction on the degree  $d_D = \max(d_A, d_D)$ . Then the rest of the proof follows directly from proof of Theorem 3.11 in [1].  $\Box$ 

4. Fiedler linearizations of Rosenbrock system matrix. In this section we show that the constructed Fiedler pencils associated with Rosenbrock systems are indeed linearizations. To do this we have to recall a few basic facts.

DEFINITION 4.1 (System equivalence). Let  $S_1(\lambda)$  and  $S_2(\lambda)$  be  $(n+m) \times (n+m)$  Rosenbrock system polynomials of the form (1.5), partitioned conformably. Then  $S_1(\lambda)$  is said to be system equivalent to  $S_2(\lambda)$  (denoted as  $S_1(\lambda) \sim_{se} S_2(\lambda)$ ), if there exist unimodular matrix polynomials  $U(\lambda), V(\lambda) \in \mathbb{F}^{n,n}$ ,  $\widetilde{U}(\lambda) \in \mathbb{F}^{m,m}$ , and  $\widetilde{V}(\lambda) \in$ 

 $\mathbb{F}^{m,m}$  such that for all  $\lambda \in \mathbb{C}$  we have

$$\begin{bmatrix} U(\lambda) & 0\\ 0 & \widetilde{U}(\lambda) \end{bmatrix} S_1(\lambda) \begin{bmatrix} V(\lambda) & 0\\ 0 & \widetilde{V}(\lambda) \end{bmatrix} = S_2(\lambda).$$
(4.1)

DEFINITION 4.2 (Rosenbrock linearization). Let  $S(\lambda)$  be an  $(n+m) \times (n+m)$ system polynomial of the form (1.5) with degree  $d = \max\{d_A, d_D\}$ . A linear matrix polynomial  $L(\lambda)$  is called a Rosenbrock linearization of  $S(\lambda)$ , if it has the form

$$L(\lambda) := \begin{bmatrix} \mathcal{A}(\lambda) & \mathcal{B} \\ \hline \mathcal{C} & \mathcal{D}(\lambda) \end{bmatrix}$$

with matrix polynomials  $\mathcal{A}(\lambda), \mathcal{D}(\lambda)$  of degree less than or equal to 1, (constant in  $\lambda$ ) matrices  $\mathcal{B}, \mathcal{C}$ , and  $\mathcal{S}(\lambda)$  is system equivalent to

$$\tilde{\mathcal{S}}(\lambda) := \begin{bmatrix} U(\lambda) & 0\\ 0 & \tilde{U}(\lambda) \end{bmatrix} L(\lambda) \begin{bmatrix} V(\lambda) & 0\\ 0 & \tilde{V}(\lambda) \end{bmatrix} = \begin{bmatrix} I_{(d-1)n} & 0\\ 0 & S(\lambda) \end{bmatrix}.$$
(4.2)

If, in addition,  $U(\lambda)$  and  $V(\lambda)$  in (4.2) are constant matrices, then  $S(\lambda)$  is said to be a strict Rosenbrock linearization of  $S(\lambda)$ .

Let  $E := (E_{ij})$  be a block  $m \times n$  matrix with  $p \times q$  blocks  $E_{ij}$ . The block transpose of E, see [11], denoted by  $E^{\mathcal{B}}$ , is the block  $n \times m$  matrix with  $p \times q$  blocks defined by  $(E^{\mathcal{B}})_{ij} := E_{ji}$ . We slightly modify this definition for the special structure of Rosenbrock linearizations.

DEFINITION 4.3 (Rosenbrock block transpose). Consider a Rosenbrock system matrix of the from (1.5) and let S be an  $(d_A n+m) \times (d_A n+m)$  Rosenbrock linearization of the form (4.2), where  $\mathcal{B} := -(e_i e_j^T) \otimes B$  and  $\mathcal{C} := (e_k e_\ell^T) \otimes C$  with  $B \in \mathbb{F}^{n,m}$ ,  $C \in \mathbb{F}^{p,n}$ , where  $\mathcal{A} := [\mathcal{A}_{ij}]$  is an  $d_A \times d_A$  block matrix with  $\mathcal{A}_{ij} \in \mathbb{F}^{n \times n}$ , and where  $\mathcal{D}$  is a  $d_D \times d_D$  block matrix with  $D_{ij} \in \mathbb{F}^{p \times m}$ , or  $\mathbb{F}^{m \times m}$  or  $\mathbb{F}^{p,p}$ . The Rosenbrock block transpose of S, denoted by  $S^{\mathbb{B}}$  is defined by

$$\mathcal{S}^{\mathbb{B}} := \left[ \begin{array}{c|c} A(\lambda)^{\mathcal{B}} & -(e_{\ell}e_{k}^{T}) \otimes B \\ \hline (e_{j}e_{i}^{T}) \otimes C & D(\lambda)^{\mathcal{B}} \end{array} \right].$$

For  $C_1(\lambda)$  and  $C_2(\lambda)$  given in (3.2) and (3.9), respectively, we have  $C_2(\lambda) = C_1(\lambda)^{\mathbb{B}}$ .

Extending [[11], Definition 4.2] we define auxiliary matrix polynomials associated with Horner shifts for system polynomials.

DEFINITION 4.4. Let  $A(\lambda) = \sum_{i=0}^{d_A} \lambda^i A_i \in \mathbb{C}[\lambda]^{n,n}$  be of degree  $d_A$  and let  $P_i(\lambda)$  be the degree *i* Horner shift of  $A(\lambda)$ . For  $1 \leq i \leq d_A - 1$ , define the matrix polynomials

$$Q_i(\lambda) := \begin{bmatrix} I_{(i-1)n} & & & \\ & I_n & \lambda I_n & \\ & 0_n & I_n & \\ & & & I_{(d_A - i - 1)n} \end{bmatrix},$$

$$R_i(\lambda) := \begin{bmatrix} I_{(i-1)n} & & & \\ & 0_n & I_n & \\ & -I_n & P_i(\lambda) & \\ & & & I_{(d_A - i - 1)n} \end{bmatrix}$$

$$T_{i}(\lambda) := \begin{bmatrix} 0_{(i-1)n} & & & \\ & 0_{n} & \lambda P_{i-1}(\lambda) & & \\ & & \lambda I_{n} & \lambda^{2} P_{i-1}(\lambda) & & \\ & & & 0_{(d_{A}-i-1)n} \end{bmatrix},$$
$$D_{i}(\lambda) := \begin{bmatrix} 0_{(i-1)n} & & & \\ & P_{i-1}(\lambda) & 0_{n} & & \\ & & 0_{n} & I_{n} & \\ & & & I_{(d_{A}-i-1)n} \end{bmatrix},$$

and  $D_{d_A}(\lambda) := \text{diag}\left[0_{(d_A-1)n}, P_{d_A-1}(\lambda)\right]$ . For simplicity, we often write  $Q_i, R_i, T_i, D_i$ in place of  $Q_i(\lambda), R_i(\lambda), T_i(\lambda), D_i(\lambda)$ . Note that  $D_1(\lambda) = M_{d_A}$ , and  $Q_i(\lambda), R_i(\lambda)$  are unimodular for all  $i = 1, \ldots, d_A - 1$ . Also note that  $R_i^{\mathcal{B}}(\lambda) = R_i(\lambda)$ .

The auxiliary matrices satisfy the following relations.

LEMMA 4.5 ([11], Lemma 4.3). Let  $Q_i, R_i, T_i, D_i$  be as in Definition 4.4 and  $M_i$ 's be Fiedler matrices associated with  $A(\lambda)$ . Then the following relations hold for  $i = 1, \ldots, d_A - 1$ .

(a)  $Q_i^{\mathcal{B}}(\lambda D_i)R_i = \lambda D_{i+1} + T_i$ , and  $Q_i^{\mathcal{B}}(M_{d_A-(i+1)}M_{d_A-i})R_i = M_{d_A-(i+1)} + T_i$ . (b)  $R_i^{\mathcal{B}}(\lambda D_i)Q_i = \lambda D_{i+1} + T_i^{\mathcal{B}}$ , and  $R_i^{\mathcal{B}}(M_{d_A-i}M_{d_A-(i+1)})Q_i = M_{d_A-(i+1)} + T_i^{\mathcal{B}}$ . (c)  $T_iM_j = M_jT_i = T_i$  and  $T_i^{\mathcal{B}}M_j = M_jT_i^{\mathcal{B}} = T_i^{\mathcal{B}}$  for all  $j \leq d_A - i - 2$ .

DEFINITION 4.6. Let  $D(\lambda) = \sum_{i=0}^{d_D} \lambda^i D_i$  be an  $m \times m$  matrix polynomial, and let  $P_i(\lambda)$  be the degree *i* Horner shift of  $D(\lambda)$ . For  $1 \le i \le d_D - 1$ , define the following  $md_D \times md_D$  matrix polynomials:

$$\begin{split} Z_{i}(\lambda) &:= \begin{bmatrix} I_{(i-1)m} & & & \\ & I_{m} & \lambda I_{m} & \\ & 0_{m} & I_{m} & \\ & & I_{(d_{D}-i-1)m} \end{bmatrix}, \\ J_{i}(\lambda) &:= \begin{bmatrix} I_{(i-1)m} & & & \\ & 0_{m} & I_{m} & \\ & I_{m} & P_{i}(\lambda) & \\ & & I_{(d_{D}-i-1)m} \end{bmatrix}, \\ H_{i}(\lambda) &:= \begin{bmatrix} 0_{(i-1)m} & & & \\ & 0_{m} & \lambda P_{i-1}(\lambda) & \\ & \lambda I_{m} & \lambda^{2} P_{i-1}(\lambda) & \\ & & 0_{(d_{D}-i-1)m} \end{bmatrix}, \\ E_{i}(\lambda) &:= \begin{bmatrix} 0_{(i-1)m} & & & \\ & P_{i-1}(\lambda) & 0_{m} & \\ & & 0_{m} & I_{m} & \\ & & & I_{(d_{D}-i-1)m} \end{bmatrix}, \end{split}$$

and  $E_{d_D}(\lambda) := \text{diag}\left[0_{(d_D-1)m}, P_{d_D-1}(\lambda)\right]$ . For simplicity, we often write  $Z_i, J_i, H_i, E_i$ in place of  $Z_i(\lambda), J_i(\lambda), H_i(\lambda), E_i(\lambda)$ . Note that  $E_1(\lambda) = M_{d_D}$ , and  $Z_i(\lambda), J_i(\lambda)$  are unimodular for all  $i = 1, \ldots, d_D - 1$ . Also note that  $J_i^{\mathcal{B}}(\lambda) = J_i(\lambda)$ .

REMARK 4.7. Consider the auxiliary matrices  $Z_i(\lambda), J_i(\lambda), H_i(\lambda)$ , and  $E_i(\lambda)$ given in Definition 4.6. Then the Lemma 4.5 also holds for  $Z_i(\lambda), J_i(\lambda), H_i(\lambda)$ , and  $E_i(\lambda)$ .

DEFINITION 4.8 (Auxiliary system polynomials). Let  $Q_i(\lambda), R_i(\lambda), T_i(\lambda)$ , and  $D_i(\lambda)$  be as in Definition 4.4. Let  $Z_i(\lambda), J_i(\lambda), H_i(\lambda)$ , and  $E_i(\lambda)$  be as in Definition

4.6. Let  $d = \max\{d_A, d_D\}$ , and  $r = \min\{d_A, d_D\}$ . For  $i = 1, \ldots, d-1$ , define  $(nd_A + md_D) \times (nd_A + md_D)$  system polynomials:

$$\mathcal{Q}_{i}(\lambda) = \begin{cases} \begin{bmatrix} Q_{i}(\lambda) & 0 \\ 0 & Z_{i}(\lambda) \\ \end{bmatrix}, & \text{for } 1 \leq i \leq r-1 \\ \begin{bmatrix} Q_{i}(\lambda) & 0 \\ 0 & I_{d_{D}m} \\ \end{bmatrix}, & \text{for } r \leq i \leq d-1 \text{ and } d_{A} > d_{D} \\ \begin{bmatrix} I_{d_{A}n} & 0 \\ 0 & Z_{i}(\lambda) \end{bmatrix}, & \text{for } r \leq i \leq d-1 \text{ and } d_{A} < d_{D} \end{cases}$$

$$\mathcal{R}_{i}(\lambda) = \begin{cases} \begin{bmatrix} R_{i}(\lambda) & 0 \\ 0 & J_{i}(\lambda) \\ \hline 0 & J_{d_{D}m} \\ \hline 0 & I_{d_{D}m} \\ \hline 1 & 0 \\ \hline 0 & J_{i}(\lambda) \end{bmatrix}, & \text{for } r \leq i \leq d-1 \text{ and } d_{A} > d_{D} \\ \hline 1 & d_{A}n & 0 \\ \hline 0 & J_{i}(\lambda) \end{bmatrix}, & \text{for } r \leq i \leq d-1 \text{ and } d_{A} < d_{D} \end{cases}$$

$$\mathcal{T}_{i}(\lambda) = \begin{cases} \begin{bmatrix} T_{i}(\lambda) & 0\\ 0 & H_{i}(\lambda) \end{bmatrix}, & \text{for } 1 \leq i \leq r-1\\ \begin{bmatrix} T_{i}(\lambda) & 0\\ 0 & I_{d_{D}m} \end{bmatrix}, & \text{for } r \leq i \leq d-1 \text{ and } d_{A} > d_{D}\\ \begin{bmatrix} I_{d_{A}n} & 0\\ 0 & H_{i}(\lambda) \end{bmatrix}, & \text{for } r \leq i \leq d-1 \text{ and } d_{A} < d_{D} \end{cases}$$

$$\mathcal{D}_{i}(\lambda) = \begin{cases} \begin{bmatrix} D_{i}(\lambda) & 0 \\ 0 & E_{i}(\lambda) \\ \end{bmatrix}, & \text{for } 1 \leq i \leq r-1 \\ \begin{bmatrix} D_{i}(\lambda) & 0 \\ 0 & I_{d_{D}m} \\ \end{bmatrix}, & \text{for } \leq i \leq d-1 \text{ and } d_{A} > d_{D} \\ \begin{bmatrix} I_{d_{A}n} & 0 \\ 0 & E_{i}(\lambda) \end{bmatrix}, & \text{for } r \leq i \leq d-1 \text{ and } d_{A} < d_{D} \end{cases}$$

and 
$$\mathcal{D}_d(\lambda) := \begin{bmatrix} D_{d_A}(\lambda) & 0\\ 0 & E_{d_D}(\lambda) \end{bmatrix}$$
, where  $d = \max\{d_A, d_D\}$ 

Note that  $\mathcal{D}_1(\lambda) = \begin{bmatrix} D_1(\lambda) & 0 \\ 0 & E_1(\lambda) \end{bmatrix} = \begin{bmatrix} M_{d_A} & 0 \\ 0 & N_{d_D} \end{bmatrix} = \mathbb{M}_d$  and that  $\mathcal{Q}_i(\lambda)$  and  $\mathcal{R}_i(\lambda)$  are unimodular matrix polynomials for  $i = 1, \ldots, d-1$ . Also, note that  $\mathcal{R}_i^{\mathbb{B}}(\lambda) = \mathcal{R}_i(\lambda)$  for  $i = 1, \ldots, d-1$ .

The auxiliary system polynomials satisfy the following relations.

LEMMA 4.9. Let  $Q_i, \mathcal{R}_i, \mathcal{T}_i, \mathcal{D}_i$  be the system polynomials given in Definition 4.8 and  $\mathbb{M}_i$ 's be Fiedler matrices associated with  $S(\lambda)$ . Then the following system equivalence relations hold for  $i = 1, \ldots, d-1$ .

(a) 
$$\mathcal{Q}_{i}^{\mathbb{B}}(\lambda \mathcal{D}_{i})\mathcal{R}_{i} = \lambda \mathcal{D}_{i+1} + \mathcal{T}_{i}$$
, and  $\mathcal{Q}_{i}^{\mathbb{B}}(\mathbb{M}_{d-(i+1)}\mathbb{M}_{d-i})\mathcal{R}_{i} = \mathbb{M}_{d-(i+1)} + \mathcal{T}_{i}$ .  
(b)  $\mathcal{R}_{i}^{\mathbb{B}}(\lambda \mathcal{D}_{i})\mathcal{Q}_{i} = \lambda \mathcal{D}_{i+1} + \mathcal{T}_{i}^{\mathbb{B}}$ , and  $\mathcal{R}_{i}^{\mathbb{B}}(\mathbb{M}_{d-i}\mathbb{M}_{d-(i+1)})\mathcal{Q}_{i} = \mathbb{M}_{d-(i+1)} + \mathcal{T}_{i}^{\mathbb{B}}$ .  
(c)  $\mathcal{T}_{i}\mathbb{M}_{j} = \mathbb{M}_{j}\mathcal{T}_{i} = \mathcal{T}_{i}$  and  $\mathcal{T}_{i}^{\mathbb{B}}\mathbb{M}_{j} = \mathbb{M}_{j}\mathcal{T}_{i}^{\mathbb{B}} = \mathcal{T}_{i}^{\mathbb{B}}$  for all  $j \leq d-i-2$ .  
Proof.

(a) We have

$$\begin{aligned} \mathcal{Q}_{i}^{\mathbb{B}}(\lambda \mathcal{D}_{i})\mathcal{R}_{i} &= \left[ \begin{array}{c|c} Q_{i}^{\mathcal{B}} & 0\\ \hline 0 & Z_{i}^{\mathcal{B}} \end{array} \right] \left[ \begin{array}{c|c} \lambda D_{i} \\ \hline \lambda E_{i} \end{array} \right] \left[ \begin{array}{c|c} R_{i} & 0\\ \hline 0 & J_{i} \end{array} \right] \\ &= \left[ \begin{array}{c|c} Q_{i}^{\mathcal{B}}(\lambda D_{i})R_{i} \\ \hline & Z_{i}^{\mathcal{B}}(\lambda E_{i})J_{i} \end{array} \right] \\ &= \left[ \begin{array}{c|c} \lambda D_{i+1} + T_{i} \\ \hline & \lambda E_{i+1} + H_{i} \end{array} \right] \text{ (By Lemma 4.5(a) and Remark 4.7)} \\ &= \left[ \begin{array}{c|c} \lambda D_{i+1} \\ \hline & \lambda E_{i+1} \end{array} \right] + \left[ \begin{array}{c|c} T_{i} \\ \hline & H_{i} \end{array} \right] = \lambda \mathcal{D}_{i+1} + \mathcal{T}_{i} \text{ and} \end{aligned}$$

$$\begin{aligned} \mathcal{Q}_{i}^{\mathbb{B}}\mathbb{M}_{d-(i+1)}\mathbb{M}_{d-i}\mathcal{R}_{i} &= \\ &= \left[\begin{array}{c|c} Q_{i}^{\mathcal{B}} & 0\\ \hline 0 & Z_{i}^{\mathcal{B}} \end{array}\right] \left[\begin{array}{c|c} M_{d-(i+1)} \\ \hline D_{d-(i+1)} \end{array}\right] \left[\begin{array}{c|c} M_{d-i} \\ \hline D_{d-i} \end{array}\right] \left[\begin{array}{c|c} R_{i} & 0\\ \hline 0 & J_{i} \end{array}\right] \\ &= \left[\begin{array}{c|c} Q_{i}^{\mathcal{B}}M_{d-(i+1)}M_{d-i}R_{i} \\ \hline Z_{i}^{\mathcal{B}}D_{m-(i+1)}D_{m-1}J_{i} \end{array}\right] \\ &= \left[\begin{array}{c|c} M_{d-(i+1)} + T_{i} \\ \hline D_{d-(i+1)} + H_{i} \end{array}\right] \text{ (Lemma 4.5(a) and Remark 4.7)} \\ &= \left[\begin{array}{c|c} M_{d-(i+1)} \\ \hline D_{d-(i+1)} \end{array}\right] + \left[\begin{array}{c|c} T_{i} \\ \hline H_{i} \end{array}\right] = \mathbb{M}_{d-(i+1)} + \mathcal{T}_{i}. \end{aligned}$$

(b) We have

$$\begin{aligned} \mathcal{R}_{i}^{\mathbb{B}}(\lambda \mathcal{D}_{i})\mathcal{Q}_{i} &= \left[ \begin{array}{c|c} R_{i}^{\mathcal{B}} & 0\\ \hline 0 & J_{i}^{\mathcal{B}} \end{array} \right] \left[ \begin{array}{c|c} \lambda D_{i} \\ \hline \lambda E_{i} \end{array} \right] \left[ \begin{array}{c|c} Q_{i} & 0\\ \hline 0 & Z_{i} \end{array} \right] \\ &= \left[ \begin{array}{c|c} R_{i}^{\mathcal{B}}(\lambda D_{i})Q_{i} \\ \hline J_{i}^{\mathcal{B}}(\lambda E_{i})Z_{i} \end{array} \right] \\ &= \left[ \begin{array}{c|c} \lambda D_{i+1} + T_{i}^{\mathcal{B}} \\ \hline \lambda E_{i+1} + H_{i}^{\mathcal{B}} \end{array} \right] \text{ (From Lemma 4.5(b) and Remark 4.7)} \\ &= \left[ \begin{array}{c|c} \lambda D_{i+1} \\ \hline \lambda E_{i+1} \end{array} \right] + \left[ \begin{array}{c|c} T_{i}^{\mathcal{B}} \\ \hline H_{i}^{\mathcal{B}} \end{array} \right] = \lambda \mathcal{D}_{i+1} + \mathcal{T}_{i}^{\mathbb{B}} \text{ and} \end{aligned}$$

$$\begin{aligned} \mathcal{R}_{i}^{\mathbb{B}}\mathbb{M}_{d-i}\mathbb{M}_{d-(i+1)}\mathcal{Q}_{i} &= \\ \begin{bmatrix} \underline{R_{i}^{\mathcal{B}}} & 0\\ \hline 0 & J_{i}^{\mathcal{B}} \end{bmatrix} \begin{bmatrix} \underline{M_{d-i}} & \\ \hline D_{d-i} \end{bmatrix} \begin{bmatrix} \underline{M_{d-(i+1)}} & \\ \hline D_{d-(i+1)} \end{bmatrix} \begin{bmatrix} \underline{Q_{i}} & 0\\ \hline 0 & Z_{i} \end{bmatrix} \\ &= \begin{bmatrix} \underline{R_{i}^{\mathcal{B}}M_{d-i}M_{d-(i+1)}Q_{i}} & \\ \hline J_{i}^{\mathcal{B}}D_{d-i}D_{d-(i+1)}Z_{i} \end{bmatrix} \\ &= \begin{bmatrix} \underline{M_{d-(i+1)} + T_{i}^{\mathcal{B}}} & \\ \hline D_{d-(i+1)} + H_{i}^{\mathcal{B}} \end{bmatrix} \text{ (By Lemma 4.5(b) and Remark 4.7)} \\ &= \begin{bmatrix} \underline{M_{d-(i+1)}} & \\ \hline D_{d-(i+1)} \end{bmatrix} + \begin{bmatrix} \underline{T_{i}^{\mathcal{B}}} & \\ \hline H_{i}^{\mathcal{B}} \end{bmatrix} = \mathbb{M}_{d-(i+1)} + \mathcal{T}_{i}^{\mathbb{B}}. \end{aligned}$$

(c) We have

$$\mathcal{T}_{i}\mathbb{M}_{j} = \begin{bmatrix} T_{i} \\ H_{i} \end{bmatrix} \begin{bmatrix} M_{j} \\ D_{j} \end{bmatrix} = \begin{bmatrix} T_{i}M_{j} \\ H_{i}D_{j} \end{bmatrix}$$
$$= \begin{bmatrix} M_{j}T_{i} \\ D_{j}H_{i} \end{bmatrix} \text{ (by Lemma 4.5(c) and Remark 4.7)}$$
$$= \begin{bmatrix} M_{j} \\ D_{j} \end{bmatrix} \begin{bmatrix} T_{i} \\ H_{i} \end{bmatrix} = \mathbb{M}_{j}\mathcal{T}_{i}$$

$$\mathcal{T}_{i}^{\mathbb{B}}\mathbb{M}_{j} = \begin{bmatrix} T_{i}^{\mathcal{B}} & \\ \hline & H_{i}^{\mathcal{B}} \end{bmatrix} \begin{bmatrix} M_{j} & \\ \hline & D_{j} \end{bmatrix} = \begin{bmatrix} T_{i}^{\mathcal{B}}M_{j} & \\ \hline & H_{i}^{\mathcal{B}}D_{j} \end{bmatrix}$$
$$= \begin{bmatrix} M_{j}T_{i}^{\mathcal{B}} & \\ \hline & D_{j}H_{i}^{\mathcal{B}} \end{bmatrix} \text{ (by Lemma 4.5(c) and Remark 4.7)}$$
$$= \begin{bmatrix} M_{j} & \\ \hline & D_{j} \end{bmatrix} \begin{bmatrix} T_{i}^{\mathcal{B}} & \\ \hline & H_{i}^{\mathcal{B}} \end{bmatrix} = \mathbb{M}_{j}\mathcal{T}_{i}^{\mathbb{B}}.$$

DEFINITION 4.10. Let  $\mathbb{L}_{\sigma}(\lambda) = \lambda \mathbb{M}_d - \mathbb{M}_{\sigma}$  be the Fiedler pencil of  $\mathcal{S}(\lambda)$  given in (1.2) associated with a bijection  $\sigma$ . For j = 1, 2, ..., d, define

$$\mathbb{M}_{\sigma}^{(j)} := \prod_{\sigma^{-1}(i) \le d-j} \mathbb{M}_{\sigma^{-1}(i)},$$

where the factors  $\mathbb{M}_{\sigma^{-1}(i)}$  are in the same relative order as they are in  $\mathbb{M}_{\sigma}$ . Note that  $\mathbb{M}_{\sigma}^{(1)} = \prod_{\sigma^{-1}(i) \leq d-1} \mathbb{M}_{\sigma^{-1}(i)} = \mathbb{M}_{\sigma}$  and that  $\mathbb{M}_{\sigma}^{(d)} = \mathbb{M}_0$ . Also for  $j = 1, 2, \ldots, d$ define the  $(nd_A + md_D) \times (nd_A + md_D)$  system pencils  $\mathbb{L}_{\sigma}^{(j)}(\lambda) := \lambda \mathcal{D}_j(\lambda) - \mathbb{M}_{\sigma}^{(j)}$ . Observe that  $\mathbb{L}_{\sigma}^{(1)}(\lambda) = \lambda \mathcal{D}_1 - \mathbb{M}_{\sigma}^{(1)} = \lambda \mathbb{M}_m - \mathbb{M}_{\sigma} = \mathbb{L}_{\sigma}$  and that

$$\mathbb{L}_{\sigma}^{(d)}(\lambda) = \lambda \mathcal{D}_{d} - \mathbb{M}_{\sigma}^{(d)} = \lambda \left[ \begin{array}{c|c} D_{d_{A}} & 0\\ \hline 0 & -E_{d_{D}} \end{array} \right] - \mathbb{M}_{0}$$

$$= \begin{bmatrix} -I_{(d_A-1)n} & -(e_{d_A}e_{d_D}^T) \otimes B \\ \hline A(\lambda) & & \\ \hline (e_{d_D}e_{d_A}^T) \otimes C & -I_{(d_D-1)m} \\ & & D(\lambda) \end{bmatrix}$$

The next result shows that  $\mathbb{L}_{\sigma}^{(i)}(\lambda) \sim_{se} \mathbb{L}_{\sigma}^{(i+1)}(\lambda)$  for i = 1, 2, ..., d-1. LEMMA 4.11. We have  $\mathbb{L}_{\sigma}^{(i)}(\lambda) \sim_{se} \mathbb{L}_{\sigma}^{(i+1)}(\lambda)$  for i = 1, 2, ..., d-1. More precisely, if  $Q_i$  and  $\mathcal{R}_i$  are the system polynomials given in Definition 4.8, then

$$\mathbb{L}_{\sigma}^{(i+1)}(\lambda) = \begin{cases} \mathcal{Q}_{i}^{\mathbb{B}} \mathbb{L}_{\sigma}^{(i)}(\lambda) \mathcal{R}_{i}, & \text{if } \sigma \text{ has a consecution at } d-i-1 \\ \mathcal{R}_{i}^{\mathbb{B}} \mathbb{L}_{\sigma}^{(i)} \mathcal{Q}_{i}, & \text{if } \sigma \text{ has an inversion at } d-i-1. \end{cases}$$

*Proof.* The proof is exactly the same as that of Lemma 4.5 in [11].  $\Box$ 

It is now immediate that a Fiedler pencil is a Rosenbrock linearization of  $\mathcal{S}(\lambda)$ .

THEOREM 4.12 (Rosenbrock linearization). Let  $S(\lambda)$  be an  $(n+m) \times (n+m)$ system polynomial (regular or singular) given in (1.2). Then a Fiedler pencil  $\mathbb{L}_{\sigma}(\lambda)$ of the system polynomial  $S(\lambda)$  is a Rosenbrock linearization of  $S(\lambda)$ .

*Proof.* By Lemma 4.11, we have d - 1 system equivalences

$$\mathbb{L}_{\sigma}(\lambda) = \mathbb{L}_{\sigma}^{(1)}(\lambda) \sim_{se} \mathbb{L}_{\sigma}^{(2)}(\lambda) \sim_{se} \cdots \sim_{se} \mathbb{L}_{\sigma}^{(d)}(\lambda)$$
$$= \left[ \begin{array}{c|c} -I_{(d_{A}-1)n} & -(e_{d_{A}}e_{d_{D}}^{T}) \otimes B \\ \hline (e_{d_{D}}e_{d_{A}}^{T}) \otimes C & -I_{(d_{D}-1)m} \\ \hline & D(\lambda) \end{array} \right],$$

where  $\mathbb{L}_{\sigma}^{(i)}(\lambda)$  is as in Lemma 4.11. This shows that  $\mathbb{L}_{\sigma}(\lambda) \sim_{se} I_{(d_A-1)n} \oplus S(\lambda) \oplus I_{(d_D-1)m}$ .

COROLLARY 4.13. Let  $\mathbb{L}_{\sigma}(\lambda)$  be the Fiedler pencil of  $\mathcal{S}(\lambda)$  given in (1.2) associated with a bijection  $\sigma$ , and  $\mathcal{Q}_i, \mathcal{R}_i$  for  $i = 1, 2, \ldots d - 1$ , be as in Definition 4.8. Then

$$\mathcal{U}(\lambda)\mathbb{L}_{\sigma}(\lambda)\mathcal{V}(\lambda) = \begin{bmatrix} -I_{(d_A-1)n} & -(e_{d_A}e_{d_D}^T) \otimes B \\ \hline & A(\lambda) \\ \hline & & \\ \hline & (e_{d_D}e_{d_A}^T) \otimes C & & -I_{(d_D-1)m} \\ & & & D(\lambda) \end{bmatrix}$$

$$\sim_{se} I_{(d_A-1)n} \oplus \mathcal{S}(\lambda) \oplus I_{(d_D-1)m},$$

where  $\mathcal{U}(\lambda)$  and  $\mathcal{V}(\lambda)$  are  $(nd_A + md_D) \times (nd_A + md_D)$  unimodular system polynomials given by

$$\mathcal{U}(\lambda) := \mathcal{U}_0 \mathcal{U}_1 \cdots \mathcal{U}_{d-3} \mathcal{U}_{d-2}, \text{ with } \mathcal{U}_i = \begin{cases} \mathcal{Q}_{d-(i+1)}^{\mathbb{B}}, & \text{if } \sigma \text{ has a consecution at } i, \\ \mathcal{R}_{d-(i+1)}^{\mathbb{B}}, & \text{if } \sigma \text{ has an inversion at } i, \end{cases}$$

 $\mathcal{V}(\lambda) := \mathcal{V}_{d-2}\mathcal{V}_{d-3}\cdots\mathcal{V}_{1}\mathcal{V}_{0}, \text{ with } \mathcal{V}_{i} = \begin{cases} \mathcal{R}_{d-(i+1)}, & \text{if } \sigma \text{ has a consecution at } i, \\ \mathcal{Q}_{d-(i+1)}, & \text{if } \sigma \text{ has an inversion at } i. \end{cases}$ 

The indexing of  $\mathcal{U}_i$  and  $\mathcal{V}_i$  factors in  $\mathcal{U}(\lambda)$  and  $\mathcal{V}(\lambda)$ , respectively, in Corollary 4.13 has been chosen for simplification of notation and has no other special significance.

REMARK 4.14. If we consider  $D(\lambda)$  is a matrix polynomial of degree 1 then the Fiedler pencils  $\mathbb{L}_{\sigma}(\lambda)$  are linearizations of the system matrix of LTI state-space system, see [1].

REMARK 4.15. Consider the system matrix  $S(\lambda)$  and associated transfer function  $R(\lambda)$  given in (1.2) and (1.3), respectively. Given an eigenvector x of  $\mathbb{L}_{\sigma}(\lambda)$  one can determine an eigenvector of  $S(\lambda)$  from x. That is, one can recover eigenvectors of  $R(\lambda)$  and  $S(\lambda)$  from those of the Fiedler pencils of  $R(\lambda)$ . It is directly follows from the Theorem 4.10 and Theorem 4.11 in [8].

5. Conclusions. We have considered a multivariable state-space system and its associated system matrix  $S(\lambda)$ . We have introduced Fiedler pencils of  $S(\lambda)$  and described an algorithm for their construction. Finally, we have shown that Fiedler pencils are linearizations of  $S(\lambda)$ .

## REFERENCES

- R. ALAM AND N. BEHERA, Linearizations for Rational Matrix Functions and Rosenbrock System Polynomials, SIAM J. Matrix Analysis Appl., 37(2016), 354–380.
- [2] R. ALAM AND N. BEHERA, Recovery of eigenvectors of rational matrix functions from Fiedlerlike linearizations, Linear Algebra Appl., 510(2016), 373–394.
- R. ALAM AND N. BEHERA, Generalized Fielder pencils for Rational Matrix functions, SIAM J. MATRIX ANAL. APPL., 39(2018), 587–610.
- [4] ANTOULAS, ATHANASIOS C., Approximation of large-scale dynamical systems, Advances in Design and Control, Society for Industrial and Applied Mathematics (SIAM), 2005.
- [5] E.N. ANTOULAS AND S. VOLOGIANNIDIS, A new family of companion forms of polynomial matrices, Electron. J. Linear Algebra, 11(2004), 78–87.
- [6] N. BEHERA, Fiedler linearizations for LTI state-space systems and for rational eigenvalue problems, PhD Thesis, IIT Guwahati, 2014.
- [7] N. BEHERA, Generalized Fielder pencils with repetition for rational matrix functions, Filomat, 34, (11) (2020), 3529–3552.
- [8] N. BEHERA, Fiedler linearizations for higher order state-space systems, JIMS, Accepted (2022).

 M. I. BUENO AND F. DE TERÁN, Eigenvectors and minimal bases for some families of Fiedlerlike linearizations, Linear and Multilinear Algebra, (2013), 1–24.

- [10] C. CONCA, J. PLANCHARD, AND M. VANNINATHAN, Existence and location of eigenvalues for fluid-solid structures, Comput. Methods Appl. Mech. Engrg., 77(1989), pp.253-291.
- [11] F. DE TERÁN, F. M. DOPICO, AND D. S. MACKEY, Fiedler companion linearizations and the recovery of minimal indices, SIAM J. Matrix Anal. Appl., 31(2009/10), 2181–2204.
- [12] I. GOHBERG, P. LANCASTER AND L. RODMAN, *Matrix polynomials*, Academic Press Inc., New York London, 1982.
- [13] GUGERCIN, S. AND ANTOULAS, A.C., A survey of model reduction by balanced truncation and some new results, Int. J. of Control, 77 (2004), 748–766.
- [14] T. KAILATH, Linear systems, Prentice-Hall Inc., Englewood Cliffs, N.J., 1980.
- [15] D. S. MACKEY, N. MACKEY, C. MEHL, AND V. MEHRMANN, Vector spaces of linearizations for matrix polynomials, SIAM J. Matrix Anal. Appl., 28 (2006), 971–1004
- [16] A. J. Mayo and A. C. Antoulas, A FRAMEWORK FOR THE SOLUTION OF THE GENERALIZED REALIZATION PROBLEM, Linear Algebra and its Applications, 425 (2007), 634–662.
- [17] Steeb, Willi-Hans, MATRIX CALCULUS AND KRONECKER PRODUCT WITH APPLICATIONS AND C++ PROGRAMS, World Scientific Publishing Co., Inc., River Edge, NJ, 1997.
- [18] L. MAZURENKO AND H. VOSS, Low rank rational perturbations of linear symmetric eigenproblems, ZAMM Z. Angew. Math. Mech., 86(2006), 606–616.
- [19] J. PLANCHARD, Eigenfrequencies of a tube bundle placed in a confined fluid, Comput. Methods Appl. Mech. Engrg., 30(1982), pp.75-93.
- [20] ROSENBROCK, H. H., State-space and multivariable theory, John Wiley & Sons, Inc., New York, 1970.
- [21] A. I. G. VARDULAKIS, Linear multivariable control, John Wiley & Sons Ltd., 1991.
- [22] H. Voss, A rational spectral problem in fluid-solid vibration, Electron. Trans. Numer. Anal., 16(2003), 93–105.