

CHARACTERIZING CANDIDATES FOR CANNON'S CONJECTURE FROM GEOMETRIC MEASURE THEORY

TAMUNONYE CHEETHAM-WEST, ALEXANDER NOLTE¹
RICE UNIVERSITY

ABSTRACT. We show that recent work of Song implies that torsion-free hyperbolic groups with Gromov boundary S^2 are realized as fundamental groups of closed 3-manifolds of constant negative curvature if and only if the solution to an associated spherical Plateau problem for group homology is isometric to such a 3-manifold, and suggest some related questions.

1. INTRODUCTION

Cannon's conjecture [CS98] states that a Gromov-hyperbolic group Γ with $\partial\Gamma \cong S^2$ that acts effectively on its boundary is a Kleinian group. For torsion-free groups Γ , this is equivalent to the existence of a closed orientable hyperbolic 3-manifold Y so that $\pi_1(Y) \cong \Gamma$.

Cannon's conjecture was first suggested as part of a program to prove Thurston's hyperbolization conjecture [Thu82], but remains open despite Perelman's proof of the Geometrization Theorem [Per02], [Per03a], [Per03b]. A large body of work surrounds Cannon's conjecture including, for instance, an equivalent condition in terms of ε -Fuchsian subgroups due to Marković [Mar13], a formulation in terms of the Ahlfors regular conformal dimension of the Gromov boundary by Bonk and Kleiner [BK05], and a resolution of analogous questions in some larger dimensions due to Bartels, Lück, and Weinberger [BLW10].

In this brief note, we present an equivalent formulation of Cannon's conjecture for torsion-free groups Γ (Theorem 1.1) in terms of the structure and regularity of solutions to a minimization problem studied in recent work of Song [Son22b]. In this view, the challenge of Cannon's conjecture is changed from producing a distinguished action on \mathbb{H}^3 from a vast family to an analysis of specific candidate geometric spaces that are characterizing in the following sense. If the candidate space for a particular torsion-free group Γ is a solution to Cannon's conjecture for Γ , then by Mostow Rigidity it is the unique such solution up to isometry; if it is not, then no solution to Cannon's conjecture exists for Γ .

We work in the framework of the spherical Plateau problem, as described in [Son22b] (see also Section 2). Specifically, given a torsion-free countable group Γ , a Hilbert manifold classifying space $\mathcal{S}(\Gamma)/\Gamma$ for Γ is constructed with Hilbert-Riemannian metric g_{Hil} . The group homology $H_*(\Gamma, \mathbb{Z})$ is then identified with the singular homology of $\mathcal{S}(\Gamma)/\Gamma$. For

¹This material is based upon work supported by the National Science Foundation under Grant No. 1842494 and Grant No. 2005551.

$h \in H_k(\Gamma, \mathbb{Z})$, a solution to the spherical Plateau problem for h is an integral current space $(X_\infty, d_\infty, S_\infty)$ that is the intrinsic-flat limit (see [SW11]) of a sequence C_i of k -dimensional integral currents in the sense of Ambrosio and Kirchheim [AK00] in $\mathcal{S}(\Gamma)/\Gamma$ representing h whose mass approaches the infimum of the masses of such currents. The immediately relevant feature of integral current spaces here is that (X_∞, d_∞) is a complete metric space. Solutions to the spherical Plateau problem always exist as a consequence of Wenger compactness [Wen11].

The observation of this note is that the results in [Son22b] and work of Bestvina and Mess [BM91] imply that solutions to appropriate spherical Plateau problems provide distinguished candidates for Cannon’s conjecture, constructed using only the group Γ . This is in line with the theme (e.g. [MY82], [Per03b], [Has83]) that geometric-analytic methods can often furnish fruitful candidates in existence questions in geometry and topology.

Theorem 1.1. *Let Γ be a torsion-free hyperbolic group with $\partial\Gamma \cong S^2$ that acts effectively on its boundary, let $(X_\infty, d_\infty, S_\infty)$ be any solution of the spherical Plateau problem for a generator h_Γ of $H_3(\Gamma, \mathbb{Z}) \cong \mathbb{Z}$, and let d_∞^L be the path metric of d_∞ . There exists a closed orientable hyperbolic 3-manifold with fundamental group isomorphic to Γ if and only if $(X_\infty, \sqrt{3}d_\infty^L)$ is isometric to a closed orientable hyperbolic 3-manifold (Y, g_{hyp}) with $\pi_1(Y) \cong \Gamma$.*

By Theorem 1.1, to disprove Cannon’s conjecture it would suffice to exhibit the presence of a singularity, failure of compactness, or incompatibility of fundamental group in an appropriate spherical Plateau solution. On the other hand, the Hyperbolization Theorem implies these are the only ways Cannon’s conjecture can fail: to prove Cannon’s conjecture via the approach suggested by Theorem 1.1 it suffices to show that the Plateau solutions X_∞ are closed topological 3-manifolds with $\pi_1(X_\infty) \cong \Gamma$.

One expects that such an approach would need to navigate substantial challenges: in analogous finite-dimensional Plateau problems, solutions can have singularities of codimension at least two [Alm83]. So an affirmative resolution to Cannon’s conjecture through Theorem 1.1 would require stronger regularity results in this setting than are expected in the general case.

We remark that Theorem 0.4 in [Son22b] ensures that the Plateau solutions appearing in Theorem 1.1 are at least nonempty, which rules out the most dramatic possible failure of Cannon’s conjecture from this perspective.

1.1. Some Questions. We conclude the introduction with some open questions on the spherical Plateau problem, selected with an eye towards developing tools to approach Cannon’s conjecture and related topics. A range of further questions appears in [Son22b].

From the perspective of Theorem 1.1, the regularity of spherical Plateau solutions plays a central role in Cannon’s conjecture. The development of an appropriate theory analogous to that in finite dimensions (e.g. [Alm83]) is expected to be a substantial undertaking: in the finite-dimensional setting, known proofs of regularity results are technical and involved.

In the present case, the existing theory is contained in [Son22b], and addresses questions of (non)-triviality for Plateau solutions under conditions on the group Γ . It is not yet clear what sorts of singularities to expect in spherical Plateau solutions for homology classes of hyperbolic groups. So a natural first question is:

Question 1.2. *Study examples of singularity formation in spherical Plateau solutions. Develop models for singularities that occur for spherical Plateau problems for homology classes of hyperbolic groups.*

For instance, it would be interesting to understand the possibilities for singularity formation within the class of spherical Dehn fillings studied by Song in [Son22b]. A long-term goal would be to classify possible singularities in restricted families of examples.

A basic structural feature of spherical Plateau solutions is that they can be broken up into “collapsed” and “non-collapsed” parts, which depend on a number of choices (see [Son22b], Section 5). The analysis of collapsed parts of spherical Plateau solutions is expected to be more complicated than that of non-collapsed parts. One step towards an analysis of regularity of Plateau solutions in our setting would be to determine if these Plateau solutions can always be taken to have trivial collapsed parts. One can see using Poisson embeddings (e.g. the proof of [Son22b] Theorem 3.3) that this holds for fundamental classes of closed hyperbolic 3-manifolds.

Question 1.3. *If Γ is a torsion-free hyperbolic group with $\partial\Gamma \cong S^2$, can it always be arranged for a spherical Plateau solution for a generator of $H_3(\Gamma, \mathbb{Z})$ to have trivial collapsed part?*

In a different direction, by Theorem 1.1 and work of Marković [Mar13], one class of groups for which spherical Plateau solutions solve Cannon’s conjecture are groups that satisfy the hypotheses of Cannon’s conjecture and contain sufficiently many quasi-convex surface subgroups. Here, “sufficiently many” means that for all distinct $p, q \in \partial\Gamma$, there is a quasi-convex surface subgroup H so that p and q lie in different connected components of $\partial\Gamma - \partial H$. With the hope of developing methods in setting of the spherical Plateau problem, it would be interesting to

Question 1.4. *Prove Marković’s criterion for Cannon’s conjecture [Mar13] in the framework of the spherical Plateau problem.*

We now turn to questions less directly concerned with regularity. Song shows in [Son22b] (Theorem 3.3) that results in [BCG95] and [BCG96] imply that after a homothety by a fixed factor, the spherical volume of the fundamental class of a closed connected oriented hyperbolic n -manifold agrees with the volume of the manifold. In order to assess potential counterexamples, a direction of investigation that seems productive is:

Question 1.5. *Develop methods of placing bounds on spherical volumes of Plateau solutions. Produce algorithms that allow for computational tools to find explicit bounds on spherical volumes of examples.*

One expects for upper bounds to be constructed through building examples, and for demonstration of lower bounds to be more challenging. We remark that one potential, though unlikely-appearing, application of spherical volume bounds is that were an appropriate group homology class's Plateau solution to have spherical volume beneath that of the Weeks manifold, results in [GMM09] would disprove Cannon's conjecture.

Next, two aspects of the approach to Cannon's conjecture suggested by Theorem 1.1 seem likely to be able to be addressed independently of the smoothness of Plateau solutions.

Question 1.6. *Let Γ be a torsion-free hyperbolic group with Gromov boundary S^2 and let $h \in H_3(\Gamma, \mathbb{Z})$ be a generator with associated spherical Plateau solution $(X_\infty, d_\infty, S_\infty)$. Denote the associated path-metric by d_∞^L .*

- (a) *Suppose that the metric space $(X_\infty, \sqrt{3}d_\infty^L)$ is isometric to an orientable hyperbolic 3-manifold (Y, g_{hyp}) . Must Y be closed?*
- (b) *Suppose further that Y is closed. Is $\pi_1(Y) \cong \Gamma$?*

A cause for interest in Question 1.6 is that affirmative answers would represent partial progress towards a proof of Cannon's conjecture through the perspective of Theorem 1.1, and a negative answer to either point would give a disproof of Cannon's conjecture.

Finally, one direction of inquiry communicated to the authors by Song [Son22a] is to determine to what extent the Plateau solutions for a group Γ inform the algebraic structure of Γ . For instance, one question related to Question 1.6, statement (b) in this direction is

Question 1.7. *Suppose that Γ is a countable group and $h \in H_n(\Gamma, \mathbb{Z})$ has a path-connected spherical Plateau solution $(X_\infty, d_\infty, S_\infty)$ with path-metric d_∞^L isometric to the path-metric of a closed Riemannian n -manifold (M, g) of constant negative sectional curvature. Must Γ contain a subgroup isomorphic to $\pi_1(M)$?*

Acknowledgements. This note exists thanks to an inspiring talk of Antoine Song and subsequent conversations at the conference Recent Developments on Geometric Measure Theory and its Applications (on the occasion of the retirement of Bob Hardt). A.N. thanks Mike Wolf and T.C-W thanks Alan Reid for their support, and the authors thank Christos Mantoulidis, Chris Leininger, and Antoine Song for thoughtful comments on previous drafts of this note.

2. THE SPHERICAL PLATEAU PROBLEM

We briefly review the spherical Plateau problem in the torsion-free setting, as defined in [Son22b], where proofs and detailed definitions pertaining to the following can be found.

Let Γ be a countably infinite torsion-free discrete group, \mathbf{H} a separable Hilbert space, and denote the unit sphere in $\ell^2(\Gamma, \mathbf{H})$ by $\mathcal{S}(\Gamma)$. The left-regular action of Γ , specified by $(\gamma f)(v) := f(\gamma^{-1}v)$, is a proper and free action by isometries of Γ on the contractible space $\mathcal{S}(\Gamma)$, so that $\mathcal{S}(\Gamma)/\Gamma$ is a classifying space for Γ .

Fix a homology class $h \in H_k(\Gamma, \mathbb{Z})$, let $\mathcal{C}(h)$ denote the collection of k -dimensional integral currents in the sense of [AK00] representing h , and denote the mass of a current $C \in \mathcal{C}(h)$ by $\mathbf{M}(C)$. Define the spherical volume $\text{SphereVol}(h)$ of h to be $\inf_{C \in \mathcal{C}(h)} \mathbf{M}(C)$.

We remark that in the case where h corresponds to the fundamental class $[\mathbb{1}_M]$ of a closed oriented Riemannian manifold (M, g) , this definition is equivalent to the spherical volume of Besson, Courtois, and Gallot [BCG91] ([Son22b], Remark 2.6). The essential observation here is that the setup of [BCG91] can be phrased independently of reference geometric data: if D_M is a fundamental domain for the universal cover $(\widetilde{M}, \widetilde{g})$ of (M, g) , then $L^2(\widetilde{M}, d\text{Vol}_{\widetilde{g}}) \cong \ell^2(\pi_1(M), L^2(D_M, d\text{Vol}_{\widetilde{g}}))$ and $\ell^2(\pi_1(M), \mathbf{H})$ are $\pi_1(M)$ -equivariantly isometrically identified.

A solution to the spherical Plateau problem is defined to be an integral current space C_∞ that is the intrinsic-flat limit (see [SW11]) of a sequence $C_i \in \mathcal{C}(h)$ so that $\lim_{i \rightarrow \infty} \mathbf{M}(C_i) = \text{SphereVol}(h)$.

Part of the data of an integral current space C is a complete metric space (X, d) , which for appropriate path-connected spaces X induces a path metric d^L on X . The main theorem we shall use from [Son22b] is that if (Y, g_{hyp}) is a closed hyperbolic 3-manifold, then for any Plateau solution C_∞ corresponding to the fundamental class $[\mathbb{1}_Y]$, the path-metric d_∞^L associated to the metric space (X_∞, d_∞) is a metric on X_∞ and $(X_\infty, \sqrt{3}d_\infty^L)$ is isometric to (Y, g_{hyp}) (Theorem 3.4). Song's proof of this is based on an analysis of barycenter maps associated to the currents C_i converging to C_∞ , as bounds on the Jacobians of these barycenter maps grow sharp (c.f. [BCG95]).

3. PROOFS

The contribution of this note is the observation that because group homology classes analogous to fundamental classes of closed oriented hyperbolic 3-manifolds exist for all torsion-free hyperbolic groups Γ satisfying the hypotheses of Cannon's conjecture, Song's work implies that the Plateau solutions for these homology classes coincide up to homothety with the closed oriented hyperbolic 3-manifold Y with $\pi_1(Y) \cong \Gamma$ whenever such a manifold exists.

To begin, we compute the group homology $H_3(\Gamma, \mathbb{Z})$ for a group Γ satisfying the hypotheses of Theorem 1.1.

Lemma 3.1. *Let Γ be a torsion-free hyperbolic group with Gromov boundary S^2 . Then $H_3(\Gamma, \mathbb{Z}) \cong \mathbb{Z}$.*

Proof. By Corollary 1.3(c) of [BM91], since $\partial\Gamma$ is S^2 , Γ is an integral Poincaré duality group of dimension 3. The rest is a standard computation: $H_0(\Gamma, \mathbb{Z}) \cong \mathbb{Z}$ so by duality $H^3(\Gamma, \mathbb{Z}) \cong \mathbb{Z}$, and by the Universal Coefficient Theorem the top homology group is \mathbb{Z} as well. \square

We now establish our result.

Proof of Theorem 1.1. Let Γ be a group satisfying the hypotheses of Theorem 1.1. Lemma 3.1 ensures that $H_3(\Gamma, \mathbb{Z}) \cong \mathbb{Z}$; let h be a generator of $H_3(\Gamma, \mathbb{Z})$ and let C_∞ be a spherical Plateau solution for h .

Suppose that (Y, g_{hyp}) is a closed orientable hyperbolic 3-manifold with $\pi_1(Y) \cong \Gamma$. As Y is aspherical, its homology is isomorphic to that of $\mathcal{S}(\Gamma)/\Gamma$. Choose an orientation on Y so that the isomorphism $\Gamma \cong \pi_1(Y)$ induces an equivariant isometry $\mathcal{S}(\Gamma) \rightarrow \mathcal{S}(\pi_1(Y))$ whose induced map on $H_3(\mathcal{S}(\Gamma)/\Gamma, \mathbb{Z})$ maps h to the class corresponding to $[\mathbb{1}_Y]$. Now apply Theorem 3.4 in [Son22b]. The other implication is trivial. \square

REFERENCES

- [AK00] Luigi Ambrosio and Bernd Kirchheim, *Currents in metric spaces*, Acta Math. **185** (2000), 1–80.
- [Alm83] Frederick J. Almgren Jr., *Q valued functions minimizing Dirichlet’s integral and the regularity of area minimizing rectifiable currents up to codimension two*, Bull. Amer. Math. Soc. (N.S.) **8** (1983), 327–328.
- [BCG91] Gérard Besson, Gilles Courtois, and Sylvestre Gallot, *Volume et entropie minimale des espaces localement symétriques*, Invent. Math. **103** (1991), 417–445.
- [BCG95] ———, *Entropies et rigidités des espaces localement symétriques de courbure strictement négative*, Geom. Funct. Anal. **5** (1995), 731–799.
- [BCG96] ———, *Minimal entropy and Mostow’s rigidity theorems*, Ergodic Theory Dynam. Systems **16** (1996), 623–649.
- [BK05] Mario Bonk and Bruce Kleiner, *Conformal dimension and Gromov hyperbolic groups with 2–sphere boundary*, Geom. Topol. (2005), 219–246.
- [BLW10] Arthur Bartels, Wolfgang Lück, and Shmuel Weinberger, *On hyperbolic groups with spheres as boundary*, J. Differential Geom. **86** (2010), 1–16.
- [BM91] Mladen Bestvina and Geoffrey Mess, *The boundary of negatively curved groups*, J. Amer. Math. Soc. (1991), 469–481.
- [CS98] James W. Cannon and Eric L. Swenson, *Recognizing constant curvature discrete groups in dimension 3*, Trans. Amer. Math. Soc. (1998), 809–849.
- [GMM09] David Gabai, Robert Meyerhoff, and Peter Milley, *Minimum volume cusped hyperbolic three-manifolds*, J. Amer. Math. Soc. **22** (2009), 1157–1215.
- [Has83] Joel Hass, *The geometry of the slice-ribbon problem*, Math. Proc. Cambridge Philos. Soc. **94** (1983), 101–108.
- [Mar13] Vladimir Marković, *Criterion for Cannon’s conjecture*, Geom. Funct. Anal. **23** (2013), 1035–1061.
- [MY82] William H. Meeks III and Shing Tung Yau, *The classical Plateau problem and the topology of three-dimensional manifolds. The embedding of the solution given by Douglas-Morrey and an analytic proof of Dehn’s lemma*, Topology (1982), 409–442.
- [Per02] Grisha Perelman, *The entropy formula for the Ricci flow and its geometric applications*, arXiv:math/0211159, 2002.
- [Per03a] ———, *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, arXiv:math/0307245, 2003.
- [Per03b] ———, *Ricci flow with surgery on three-manifolds*, arXiv:math/0303109, 2003.
- [Son22a] Antoine Song, *Candidates for Cannon’s conjecture*, private communication (2022).
- [Son22b] ———, *The spherical Plateau problem for group homology*, arXiv:math/2202.10636, 2022.

- [SW11] Christina Sormani and Stefan Wenger, *The intrinsic flat distance between Riemannian manifolds and other integral current spaces*, J. Differential Geom. **87** (2011), 117–199.
- [Thu82] William Thurston, *Three-dimensional manifolds, Kleinian groups and hyperbolic geometry*, Bull. Amer. Math. Soc. (N.S.) (1982), 357–381.
- [Wen11] Stefan Wenger, *Compactness for manifolds and integral currents with bounded diameter and volume*, Calc. Var. Partial Differential Equations **40** (2011), 423–448.