

A CONSTRUCTIVE PROOF OF NC FEJÉR-RIESZ THEOREM

PALAK ARORA

ABSTRACT. In this paper we present a constructive proof of Popescu's non-commutative Fejér-Riesz theorem for non-commuting polynomials. We are considering non-commuting polynomial in left-creation and left-annihilation multi-Toeplitz operators.

1. INTRODUCTION

The classical Fejér-Riesz states the following: if a trigonometric polynomial

$$w(e^{it}) = \sum_{j=-m}^m c_j e^{ijt}$$

is nonnegative for all real t , then it is expressible in the form

$$w(e^{it}) = |p(e^{it})|^2$$

for some analytic polynomial $p(z) = \sum_{j=0}^m a_j z^j$. For proof refer to Lemma 2.1 in [1]. There is also an operator version where the coefficients of w are matrices or operators ([2]). The Fejér-Riesz theorem can be reformulated as a statement about Toeplitz operators: the function w may be interpreted as the symbol of a *Toeplitz operator* T_w ; in particular if S denotes the unilateral shift on $\ell^2(\mathbb{N})$ then T_w is the operator defined by

$$T_w = c_0 I + \sum_{k=1}^m c_k S^k + \sum_{k=1}^m c_{-k} S^{*k},$$

and then the factorization $w = |p|^2$ is equivalent to the factorization of operators

$$T_w = T_p^* T_p$$

where $T_p = \sum_{k=0}^m a_k S^k = p(S)$. (The equivalence of the two formulations follows easily from the fact that S is an isometry ($S^* S = I$)). It turns out this operator formulation admits a generalization, in the noncommutative setting, to so-called *multi-Toeplitz* operators, where the single isometry S is replaced by a *row isometry*, for example the d tuple of left shifts, (L_1, L_2, \dots, L_d) or d tuple of right shifts, (R_1, R_2, \dots, R_d) . Precise definitions are given in the next section. Following an idea of Dritschel and Woerdeman [3], this paper develops a constructive proof of Riesz-Fejér theorem in the non commutative setting. We have a non negative multi-Toeplitz polynomial operator

$$T_Q := Q_0 \otimes I_{\mathcal{F}_d^2} + \sum_{0 < |v| \leq n} Q_v \otimes L_v + \sum_{0 < |v| \leq n} Q_v^* \otimes L_v^*.$$

We then find the multi-Toeplitz operator factorization of this polynomial,

$$T_Q := T_F^* T_F$$

where $T_F := \sum_{0 \leq |v| \leq n} F_v \otimes L_v$ [This is a slight rewording of the theorem 1.6 [4]].

2. PRELIMINARIES

Let us recall a few definitions required for the following section:

Definition 2.1. Let \mathcal{F}_d^+ denote the word set which is a monoid formed from the letters, $1, 2, \dots, d$. We say that **Fock space** is $\ell^2(\mathcal{F}_d^+)$.

The Fock space, $\ell^2(\mathcal{F}_d^+)$ is the Hilbert space with orthonormal basis $\{\xi_w\}_{w \in \mathcal{F}_d^+}$.

Definition 2.2. The **left-creation operator** L_j is defined as

$$L_j \xi_w = \xi_{jw}$$

for $j = 1, \dots, d$ and can be extended linearly. Similarly, **left-annihilation operator** L_j^* is defined as

$$L_j^* \xi_w = \begin{cases} \xi_v, & w = jv \\ 0, & \text{otherwise} \end{cases}$$

for $j = 1, \dots, d$. Thus $\{L_j | j = 1, \dots, d\}$ form a system of isometries with orthogonal ranges:

$$L_i^* L_j = \delta_{ij} I.$$

Note that same holds for the right shift operators as well.

From the above definition we get that (L_1, L_2, \dots, L_d) and (R_1, R_2, \dots, R_d) are row-isometries. For any $w \in \mathcal{F}_d^+$, $w = i_1 i_2 \dots i_n$ we denote $L_w = L_{i_1} L_{i_2} \dots L_{i_n}$. So for any $w = i_1 i_2 \dots i_n$ and $v = j_1 j_2 \dots j_m$ in \mathcal{F}_d^+ , $L_w^* L_v = L_{i_n}^* \dots L_{i_2}^* L_{i_1}^* L_{j_1} \dots L_{j_m}$. Thus

$$L_w^* L_v = \begin{cases} L_x, & \text{if } v = wx \\ L_y^*, & \text{if } w = xy \\ 0, & \text{otherwise} \end{cases}$$

Definition 2.3. In the classical setting, T is said to be a **Toeplitz operator** if $S^* T S = T$ where S is unilateral shift. An operator T is **L -multi-Toeplitz** if $L_i^* T L_j = \delta_{ij} T$ where L_j is left-creation operator. Similarly, T is called **R -multi-Toeplitz** if $R_i^* T R_j = \delta_{ij} T$ where R_j is right-creation operator.

Example 1. Any left-creation operator L_w is R -multi-Toeplitz. Since L_i and R_j commute with each other, R_j commutes with L_w for all w and thus we have

$$\begin{aligned} R_i^* L_w R_j &= R_i^* R_j L_w \\ &= \delta_{ij} L_w \end{aligned}$$

Similarly, L_v^* is R -multi-Toeplitz for any word v .

Therefore for any non commutative polynomials f, g , we have that $f(L)^* + g(L)$ is R -multi-Toeplitz.

Next let us consider a R -multi-Toeplitz operator say,

$$(1) \quad T := \sum_{0 \leq |v| \leq n} q_v L_v + \sum_{0 < |v| \leq n} q_v^* L_v^*.$$

Then corresponding to the Fock space basis $\{\xi_v\}_{v \in \mathcal{F}_d^+}$ we get its matrix representation which is a multi-Toeplitz matrix:

q_0	q_1^*	\cdots	q_d^*	\cdots	$q_{11\dots 1}^*$	\cdots	$q_{1d\dots d}^*$	\cdots	$q_{d1\dots 1}^*$	\cdots	$q_{dd\dots d}^*$	0	\cdots	0	\cdots
q_1	q_0	\cdots	0	\cdots	$q_{11\dots 1}^*$	\cdots	0	\cdots	$q_{d1\dots 1}^*$	\cdots	0	$q_{11\dots 1}^*$	\cdots	0	\cdots
\vdots	\vdots	\ddots	\vdots		\vdots	\ddots	\vdots	\ddots	\vdots	\ddots	\vdots	\vdots	\ddots	\vdots	\ddots
q_d	0	\cdots	q_0	\cdots	0	\cdots	$q_{11\dots 1}^*$	\cdots	0	\cdots	$q_{dd\dots d}^*$	0	\cdots	$q_{dd\dots d}^*$	\ddots
\vdots	\vdots	\ddots	\vdots	\ddots	\vdots	\ddots	\vdots	\ddots	\vdots	\ddots	\vdots	\vdots	\ddots	\vdots	\ddots
$q_{11\dots 1}$	$q_{1\dots 1}$	\cdots	0	\cdots	q_0	\cdots	0	\cdots	0	\cdots	0	q_1^*	\cdots	0	\ddots
\vdots	\vdots	\ddots	\vdots	\vdots	\vdots	\ddots	\vdots		\vdots		\vdots	\vdots	\ddots	\vdots	\ddots
$q_{1d\dots d}$	0	\cdots	$q_{1\dots 1}$	\cdots	0	\cdots	q_0	\cdots	0	\cdots	0	0		0	
\vdots		\ddots		\ddots	\vdots			\ddots			\vdots	\vdots	\ddots	\vdots	\ddots
$q_{d1\dots 1}$	$q_{d\dots d}$	\cdots	0	\cdots	0	\cdots	0	\cdots	q_0	\cdots	0	0		0	
\vdots	\vdots	\ddots	\vdots	\ddots	\vdots				\ddots		\vdots	\vdots	\ddots	\vdots	\ddots
$q_{dd\dots d}$	0	\cdots	$q_{d\dots d}$	\cdots	0	\cdots	0	\cdots	0	\cdots	q_0	0	\cdots	q_d^*	
0	$q_{11\dots 1}$	\cdots	0	\cdots	q_1	\cdots	0	\cdots	0	\cdots	0	q_0	\cdots	0	
\vdots		\ddots		\ddots	\vdots	\ddots		\ddots		\ddots	\vdots	\vdots	\ddots	\vdots	\ddots
0	0	\cdots	$q_{dd\dots d}$	\cdots	0	\cdots	0	\cdots	0	\cdots	q_d	0	\cdots	q_0	
\vdots		\ddots		\ddots		\ddots		\ddots		\ddots			\ddots		\ddots

Here we have used the lexicographic ordering for ordering the elements of the word set, \mathcal{F}_d^+ .

Now we do some relabeling of the indexes here and define for $d = 1, \dots, n$:

$$q_k := \text{col}(q_w)_{w \in \mathcal{F}_d^+; |w|=k} \text{ and } q_{-k} := \text{row}(q_w^*)_{w \in \mathcal{F}_d^+; |w|=k}$$

and also identifying, $q_1 \otimes I_d :=$

$$\begin{pmatrix} q_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & q_1 \\ \vdots & \ddots & \vdots \\ \vdots & \ddots & \vdots \\ q_d & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & q_d \end{pmatrix} \text{ and so on.}$$

Thus we get the following compact form of T , which makes it easier to see multi-Toeplitz form of the matrix:

$$T = \begin{bmatrix} q_0 & q_{-1} & q_{-2} & \cdots & q_{-n} & 0 & 0 & \cdots \\ q_1 & q_0 \otimes I_d & q_{-1} \otimes I_d & \cdots & q_{-(n-1)} \otimes I_d & q_{-n} \otimes I_d & 0 & \cdots \\ q_2 & q_1 \otimes I_d & q_0 \otimes I_d \otimes I_d & \cdots & q_{-(n-2)} \otimes I_d \otimes I_d & q_{-(n-1)} \otimes I_d \otimes I_d & q_{-n} \otimes I_d^{\otimes 2} & \cdots \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots \\ q_n & q_{n-1} \otimes I_d & q_{n-2} \otimes I_d \otimes I_d & \cdots & q_0 \otimes I_d \otimes \cdots \otimes I_d & q_{-1} \otimes I_d^{\otimes n} & q_{-2} \otimes I_d^{\otimes n} & \ddots \\ 0 & q_n \otimes I_d & q_{n-1} \otimes I_d \otimes I_d & \cdots & q_1 \otimes I_d^{\otimes n} & q_0 \otimes I_d^{\otimes (n+1)} & q_{-1} \otimes I_d^{\otimes (n+1)} & \ddots \\ 0 & 0 & q_n \otimes I_d \otimes I_d & \cdots & q_2 \otimes I_d^{\otimes n} & q_1 \otimes I_d^{\otimes (n+1)} & q_0 \otimes I_d^{\otimes (n+2)} & \ddots \\ \vdots & \vdots & \vdots & \ddots & \ddots & \ddots & \ddots & \ddots \end{bmatrix}.$$

We are going to use a schur complement technique from Dritschel and Woerdeman [3] in the proof of the main theorem. So let us define the following:

Definition 2.4. If \mathcal{H}_1 and \mathcal{H}_2 are Hilbert spaces and

$$M = \begin{pmatrix} A & B^* \\ B & C \end{pmatrix} : \mathcal{H}_1 \oplus \mathcal{H}_2 \rightarrow \mathcal{H}_1 \oplus \mathcal{H}_2$$

is a positive semidefinite operator then there exists a unique contraction $G : \overline{\text{ran}}(C) \rightarrow \overline{\text{ran}}(A)$ such that $B = A^{1/2}GC^{1/2}$. The **Schur complement of M supported on \mathcal{H}_1** is defined to be positive semidefinite operator $A^{1/2}(1 - GG^*)A^{1/2}$.

An alternative way to define the schur complement of M supported on \mathcal{H}_1 is via

$$\langle Sf, f \rangle = \inf \left\{ \left\langle \begin{pmatrix} A & B \\ B^* & C \end{pmatrix} \begin{pmatrix} f \\ g \end{pmatrix}, \begin{pmatrix} f \\ g \end{pmatrix} \right\rangle : g \in \mathcal{H}_2 \right\}$$

that is, $S : \mathcal{H}_1 \rightarrow \mathcal{H}_1$ is the largest positive semidefinite operator which may be subtracted from A in M such that the resulting operator matrix remains positive semidefinite.

Remark. Consider any positive semidefinite operator matrix, M , say

$$M = \begin{pmatrix} A & B^* \\ B & C \end{pmatrix} : \mathcal{H}_1 \oplus \mathcal{H}_2 \rightarrow \mathcal{H}_1 \oplus \mathcal{H}_2$$

and let S_M be the Schur complement of M supported on \mathcal{H}_1 . Then for a positive semidefinite matrix, $M \otimes I_d$, the Schur complement supported on $\mathcal{H}_1 \otimes \mathbb{C}^d$ is $S_{M \otimes I_d} = S_M \otimes I_d$.

If Q is an operator from \mathcal{H} to \mathcal{H} for some hilbert space \mathcal{H} , then $Q \otimes I_d$ takes values from $\mathcal{H} \otimes \mathbb{C}^d$ and outputs in $\mathcal{H} \otimes \mathbb{C}^d$.

Let us denote $\mathcal{H}_i = \mathcal{H} \otimes (\mathbb{C}^d)^{\otimes i}$ for $i \geq 0$.

We make use of the following notation for the next section from [3]: Typically we will index rows and columns of an $n \times n$ matrix with $0, \dots, n-1$. For $\Lambda \subseteq \{0, \dots, n-1\}$ and an $n \times n$ matrix M , we write $S(M; \Lambda)$, or $S(\Lambda)$ when there is no chance of confusion, for the Schur complement supported on the rows and columns labeled by elements of Λ . It is usual to view $S(\Lambda)$ as an $m \times m$ matrix, where $m = \text{card} \Lambda$, however it is often useful to take $S(\Lambda)$ as an $n \times n$ matrix by padding rest of the entries in this $n \times n$ matrix with zeros. For notational convenience we have used $S(m)$ for Schur complement supported on rows and columns labeled by $\{0, \dots, m\}$.

3. MAIN THEOREM

The following Theorem 3.1 and Corollary 3.2 are multi-Toeplitz versions of Proposition 3.1 and corollary 3.2 from [3].

Theorem 3.1. Consider the positive semidefinite multi-Toeplitz operator matrix

$$T_Q = \begin{bmatrix} Q_0 & Q_{-1} & Q_{-2} & \cdots & \cdots \\ Q_1 & Q_0 \otimes I_d & Q_{-1} \otimes I_d & \cdots & \cdots \\ Q_2 & Q_1 \otimes I_d & Q_0 \otimes I_d \otimes I_d & \cdots & \cdots \\ \vdots & \vdots & \vdots & \ddots & \ddots \end{bmatrix}$$

acting on $\mathcal{H}_0 \oplus \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \cdots$. Then Schur complement of T_Q satisfies the recurrence relation:

$$S(m) = \begin{bmatrix} A & B^* \\ B & S(m-1) \otimes I_d \end{bmatrix}$$

for appropriate choice of $A : \mathcal{H}_0 \rightarrow \mathcal{H}_0$ and $B^* : \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \cdots \rightarrow \mathcal{H}$. When $Q_j = 0$ for $j \geq m+1$, then $A = Q_0$ and $B = \text{col}(Q_i)_{i=1}^m$.

Remark. Given

$$T_Q = \begin{bmatrix} Q_0 & Q_{-1} & Q_{-2} & \cdots & \cdots \\ Q_1 & Q_0 \otimes I_d & Q_{-1} \otimes I_d & \cdots & \cdots \\ Q_2 & Q_1 \otimes I_d & Q_0 \otimes I_d \otimes I_d & \cdots & \cdots \\ \vdots & \vdots & \vdots & \ddots & \end{bmatrix},$$

we observe that T_Q can be identified with

$$T_Q = \begin{bmatrix} Q_0 & \text{row}(Q_{-j})_{j \geq 1} \\ \text{col}(Q_j)_{j \geq 1} & T_Q \otimes I_d \end{bmatrix}.$$

Proof. Let us write

$$S(m) = \begin{bmatrix} A & B^* \\ B & C \end{bmatrix} : \mathcal{H}_0 \oplus \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \cdots \oplus \mathcal{H}_m \rightarrow \mathcal{H}_0 \oplus \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \cdots \oplus \mathcal{H}_m.$$

By definition of Schur complement we have that

$$T_Q - \begin{bmatrix} S(m) & 0 \\ 0 & 0 \end{bmatrix} \geq 0.$$

That is, we have that

$$(2) \quad \left[\begin{array}{c|c} Q_0 & \text{row}(Q_{-j})_{j \geq 1} \\ \hline \text{col}(Q_j)_{j \geq 1} & T_Q \otimes I_d \end{array} \right] - \left[\begin{array}{c|c} A & B^* & 0 \\ \hline B & C & 0 \\ 0 & 0 & 0 \end{array} \right] \geq 0.$$

Then leaving out 0th row and column in (2) we get,

$$T_Q \otimes I_d - \begin{bmatrix} C & 0 \\ 0 & 0 \end{bmatrix} \geq 0.$$

So we have from theorem 3.1, $C \leq S(m-1) \otimes I_d$.

Now leaving out the rows and columns $1, \dots, m$ in (2), we get

$$\left[\begin{array}{c|c} Q_0 - A & \text{row}(Q_j^*)_{j \geq m+1} \\ \hline \text{col}(Q_j)_{j \geq m+1} & T_Q \otimes I_d^{m+1} \end{array} \right] \geq 0.$$

That is,

$$A \leq S \left(\left[\begin{array}{c|c} Q_0 & \text{row}(Q_j^*)_{j \geq m+1} \\ \hline \text{col}(Q_j)_{j \geq m+1} & T_Q \otimes I_d^{m+1} \end{array} \right]; 0 \right) := \tilde{A}.$$

Note that when $Q_j = 0$, $j \geq m+1$ then $\tilde{A} = Q_0$.

Again considering the following operator matrix

$$(3) \quad \left[\begin{array}{c|c|c} Q_0 - \tilde{A} & X & \text{row}(Q_j^*)_{j \geq m+1} \\ \hline X^* & (Q_{i-j} \otimes I_d)_{i=1, j=1}^m - S(m-1) \otimes I_d & (Q_{i-j} \otimes I_d)_{i=1, j=m+1}^{m+1, \infty} \\ \hline \text{col}(Q_j)_{j \geq m+1} & (Q_{i-j} \otimes I_d)_{i=m+1, j=1}^{\infty, m} & T_Q \otimes I_d^{m+1} \end{array} \right]$$

The existence of an operator X making this into a positive semidefinite matrix is a variant of a standard operator matrix completion problem, (see Theorem XVI.3.1 in [5]) so there always exists such an X . Note that when $\tilde{A} = Q_0$ we have necessarily that $X = 0$. We fix such an X . Now (3) is positive semidefinite, we obtain that

$$\begin{bmatrix} \tilde{A} & \text{row}(Q_j^*)_{j=1}^m - X \\ \text{col}(Q_j)_{j=1}^m - X & S(m-1) \otimes I_d \end{bmatrix} \leq S(m) = \begin{bmatrix} A & B^* \\ B & C \end{bmatrix}.$$

This implies that $\tilde{A} \leq A$ and $S(m-1) \otimes I_d \leq C$. From above we also have $A \leq \tilde{A}$ and $C \leq S(m-1) \otimes I_d$, thus the equalities $A = \tilde{A}$ and $C = S(m-1) \otimes I_d$ follow.

Moreover, if $Q_j = 0$ for $j \geq m+1$, we have that $\tilde{A} = Q_0$ and $X = 0$, and thus $B = \text{col}(Q_i)_{i=1}^m$. \square

Corollary 3.2. Consider the positive semidefinite multi-Toeplitz operator matrix

$$T_Q = \begin{bmatrix} Q_0 & Q_{-1} & Q_{-2} & \cdots & \cdots \\ Q_1 & Q_0 \otimes I_d & Q_{-1} \otimes I_d & \cdots & \cdots \\ Q_2 & Q_1 \otimes I_d & Q_0 \otimes I_d \otimes I_d & \cdots & \cdots \\ \vdots & \vdots & \vdots & \ddots & \ddots \end{bmatrix}$$

acting on $\mathcal{H}_0 \oplus \mathcal{H}_1 \oplus \mathcal{H}_2 \oplus \cdots$. Then for each $m \geq 0$, there exist operators F_0, F_1, \dots with

$$F_{m-k} \otimes I_d^{\otimes k} : \mathcal{H}_k \longrightarrow \overline{\text{ran}} F_0 \otimes \mathbb{C}^m \subseteq \mathcal{H} \otimes \mathbb{C}^m$$

for $0 \leq k \leq (m-1)$ so that the Schur complements $S(m)$ of T_Q satisfy

$$S(m) = \begin{bmatrix} F_0^* & F_1^* & F_2^* & \cdots & F_m^* \\ & F_0^* \otimes I_d & F_1^* \otimes I_d & \cdots & F_{m-1}^* \otimes I_d \\ & & F_0^* \otimes I_d^{\otimes 2} & \cdots & F_{m-2}^* \otimes I_d^{\otimes 2} \\ & & & \ddots & \vdots \\ & & & & F_0^* \otimes I_d^{\otimes m} \end{bmatrix} \begin{bmatrix} F_0 & F_1 & F_2 & \vdots & F_m \\ F_1 & F_0 \otimes I_d & F_1 \otimes I_d & \vdots & F_{m-1} \otimes I_d \\ F_2 & F_1 \otimes I_d & F_0 \otimes I_d^{\otimes 2} & \vdots & F_{m-2} \otimes I_d^{\otimes 2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ F_m & F_{m-1} \otimes I_d & F_{m-2} \otimes I_d^{\otimes 2} & \cdots & F_0 \otimes I_d^{\otimes m} \end{bmatrix}.$$

Proof. We will prove this by induction on m .

Base step: $S(0)$ being a positive semidefinite operator, we can write $S(0) = F_0^* F_0$ where $F_0 = (S(0))^{1/2}$.

Induction hypothesis: Let us assume that the result holds for $S(m-1)$.

By [3], Proposition 3.1, we have that $(S(m))_{m,m} = (S(m-1))_{m-1,m-1} \otimes I_d = F_0^* F_0 \otimes I_d^{\otimes(m-1)} \otimes I_d = F_0^* F_0 \otimes I_d^{\otimes m}$.

From [3] Corollary 2.3, we have that $S(m-1) = S(S(m); m-1)$. Thus applying Lemma 2.1 from [3] to

$$(4) \quad P = \begin{bmatrix} F_0 & F_1 & F_2 & \vdots & F_{m-1} \\ F_1 & F_0 \otimes I_d & F_1 \otimes I_d & \vdots & F_{m-2} \otimes I_d \\ F_2 & F_1 \otimes I_d & F_0 \otimes I_d^{\otimes 2} & \vdots & F_{m-3} \otimes I_d^{\otimes 2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ F_{m-1} & F_{m-2} \otimes I_d & F_{m-3} \otimes I_d^{\otimes 2} & \cdots & F_0 \otimes I_d^{\otimes(m-1)} \end{bmatrix}, R = F_0 \otimes I_d^{\otimes m}$$

there exist $(G_m \cdots G_1)$ so that

$$S(m) = \begin{bmatrix} F_0^* & F_1^* & F_2^* & \cdots & G_m^* \\ & F_0^* \otimes I_d & F_1^* \otimes I_d & \cdots & G_{m-1}^* \\ & & F_0^* \otimes I_d^{\otimes 2} & \cdots & G_{m-2}^* \\ & & & \ddots & \vdots \\ & & & & F_0^* \otimes I_d^{\otimes m} \end{bmatrix} \begin{bmatrix} F_0 & F_1 & F_2 & \vdots & G_m \\ F_1 & F_0 \otimes I_d & F_1 \otimes I_d & \vdots & G_{m-1} \\ F_2 & F_1 \otimes I_d & F_0 \otimes I_d^{\otimes 2} & \vdots & G_{m-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ G_m & G_{m-1} & G_{m-2} & \cdots & F_0 \otimes I_d^{\otimes m} \end{bmatrix},$$

and $\text{ran}(G_m \cdots G_1) \subseteq \overline{\text{ran}}F_0 \otimes (\mathbb{C}^d)^{\otimes m}$. Comparing with $S(m) = \begin{bmatrix} A & B^* \\ B & S(m-1) \otimes I_d \end{bmatrix}$ along with the induction hypothesis yields, $S(m-1) \otimes I_d$ factors into

$$\begin{aligned} & \begin{bmatrix} F_0^* \otimes I_d & F_1^* \otimes I_d & \cdots & F_{m-2}^* \otimes I_d & G_{m-1}^* \\ F_0^* \otimes I_d^{\otimes 2} & \cdots & F_{m-3}^* \otimes I_d^{\otimes 2} & G_{m-2}^* & \\ & \ddots & \vdots & \vdots & \\ & & F_0^* \otimes I_d^{\otimes(m-1)} & G_1^* & \\ & & & F_0^* \otimes I_d^m & \end{bmatrix} \begin{bmatrix} F_0 \otimes I_d & & & & \\ F_1 \otimes I_d & F_0 \otimes I_d^{\otimes 2} & & & \\ F_2 \otimes I_d & F_1 \otimes I_d^{\otimes 2} & F_0 \otimes I_d^{\otimes 3} & & \\ \vdots & \vdots & \vdots & \ddots & \\ G_{m-1} & G_{m-2} & G_{m-3} & \cdots & F_0 \otimes I_d^{\otimes m} \end{bmatrix} \\ &= \begin{bmatrix} F_0^* \otimes I_d & F_1^* \otimes I_d & \cdots & F_{m-2}^* \otimes I_d & F_{m-1}^* \otimes I_d \\ & F_0^* \otimes I_d^{\otimes 2} & \cdots & F_{m-3}^* \otimes I_d^{\otimes 2} & F_{m-2}^* \otimes I_d^{\otimes 2} \\ & & \ddots & \vdots & \vdots \\ & & & F_0^* \otimes I_d^{\otimes(m-1)} & F_1^* \otimes I_d^{\otimes(m-1)} \\ & & & & F_0^* \otimes I_d^{\otimes m} \end{bmatrix} \begin{bmatrix} F_0 \otimes I_d & & & & \\ F_1 \otimes I_d & F_0 \otimes I_d^{\otimes 2} & & & \\ F_2 \otimes I_d & F_1 \otimes I_d^{\otimes 2} & F_0 \otimes I_d^{\otimes 3} & & \\ \vdots & \vdots & \vdots & \ddots & \\ F_{m-1} \otimes I_d & F_{m-2} \otimes I_d^{\otimes 2} & F_{m-3} \otimes I_d^{\otimes 3} & \cdots & F_0 \otimes I_d^{\otimes m} \end{bmatrix} \end{aligned}$$

and thus we have

$$F_0^* \otimes I_d^{\otimes m} \begin{pmatrix} G_{m-1} & G_{m-2} & \cdots & G_1 \end{pmatrix} = F_0^* \otimes I_d^{\otimes m} \begin{pmatrix} F_{m-1} \otimes I_d & F_{m-2} \otimes I_d^{\otimes 2} & \cdots & F_1 \otimes I_d^{\otimes(m-1)} \end{pmatrix}.$$

As

$$\text{ran} \begin{pmatrix} G_{m-1} & G_{m-2} & \cdots & G_1 \end{pmatrix} \subseteq \overline{\text{ran}}F_0 \otimes (\mathbb{C}^d)^{\otimes m}$$

and

$$\text{ran} \begin{pmatrix} F_{m-1} \otimes I_d & F_{m-2} \otimes I_d^{\otimes 2} & F_{m-3} \otimes I_d^{\otimes 3} & \cdots & F_1 \otimes I_d^{\otimes(m-1)} \end{pmatrix} \subseteq \overline{\text{ran}}F_0 \otimes (\mathbb{C}^d)^{\otimes m}$$

it follows that $G_j = F_j \otimes I_d^{\otimes(m-j)}$ for $j = 1, \dots, m-1$. By setting $F_m := G_m$, we have our result. \square

Theorem 3.3. If (L_1, L_2, \dots, L_d) is the left d -shift, \mathcal{H} a Hilbert space and $\{Q_w\}_{|w| \leq n}$ are operators $Q_w : \mathcal{H} \rightarrow \mathcal{H}$ such that the operator $T_Q : \mathcal{H} \otimes \ell^2(\mathcal{F}_d^+) \rightarrow \mathcal{H} \otimes \ell^2(\mathcal{F}_d^+)$ given by

$$T_Q := Q_0 \otimes I_{\ell^2(\mathcal{F}_d^+)} + \sum_{0 < |v| \leq n} Q_v \otimes L_v + \sum_{0 < |v| \leq n} Q_v^* \otimes L_v^*$$

is positive, then there exist operators $F_0, \dots, F_w : \mathcal{H} \rightarrow \mathcal{H}$, $|w| \leq n$, such that for $T_F := \sum_{0 \leq |v| \leq n} F_v \otimes L_v$ we have $T_Q = T_F^* T_F$.

Proof. Note that this operator, T_Q is R multi-Toeplitz operator. Let us consider the matrix representation of this operator, T_Q corresponding to basis, $\beta = \{e_i \otimes \xi^v\}_{i,v}$ where $\{e_i\}_i$ is some orthonormal basis for \mathcal{H} and $v \in \mathcal{F}_d^+$. It is a multi-toeplitz operator matrix (corresponding to polynomial (1)) as was defined on page 3 with operator entries Q_j for $j \in \mathcal{F}_d^+$ instead. Then as done before, after relabeling the indexes we get the multi-Toeplitz matrix:

$$T_Q = \begin{bmatrix} Q_0 & Q_{-1} & Q_{-2} & \cdots & Q_{-n} & 0 & 0 & \cdots \\ Q_1 & Q_0 \otimes I_d & Q_{-1} \otimes I_d & \cdots & Q_{-(n-1)} \otimes I_d & Q_{-n} \otimes I_d & 0 & \cdots \\ Q_2 & Q_1 \otimes I_d & Q_0 \otimes I_d \otimes I_d & \cdots & Q_{-(n-2)} \otimes I_d \otimes I_d & Q_{-(n-1)} \otimes I_d \otimes I_d & Q_{-n} \otimes I_d^{\otimes 2} & \cdots \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots \\ Q_n & Q_{n-1} \otimes I_d & Q_{n-2} \otimes I_d \otimes I_d & \cdots & Q_0 \otimes I_d \otimes \cdots \otimes I_d & Q_{-1} \otimes I_d^{\otimes n} & Q_{-2} \otimes I_d^{\otimes n} & \ddots \\ 0 & Q_n \otimes I_d & Q_{n-1} \otimes I_d \otimes I_d & \cdots & Q_1 \otimes I_d^{\otimes n} & Q_0 \otimes I_d^{\otimes (n+1)} & Q_{-1} \otimes I_d^{\otimes (n+1)} & \ddots \\ 0 & 0 & Q_n \otimes I_d \otimes I_d & \cdots & Q_2 \otimes I_d^{\otimes n} & Q_1 \otimes I_d^{\otimes (n+1)} & Q_0 \otimes I_d^{\otimes (n+2)} & \ddots \\ \vdots & \vdots & \vdots & \ddots & \ddots & \ddots & \ddots & \ddots \end{bmatrix}.$$

In this case we have that $Q_j = 0$ for $|j| \geq n+1$.

Now consider the schur complement, $S(n) = \begin{bmatrix} A & B^* \\ B & C \end{bmatrix}$ (that is, supported on first $n+1$ rows and columns of T_Q).

Thus from theorem 3.1 we get that

$$A = Q_0 \text{ and } B = \text{col}(Q_i)_{i=1}^n.$$

Comparing the first row of $S(n)$ above with the first row of the product in corollary 3.2 factorization we get that:

$$Q_0 = \sum_{j=0}^n F_j^* F_j;$$

$$Q_{-j} = \sum_{k=j}^n F_k^* (F_{k-j} \otimes I_d^{\otimes j}) \text{ for } 1 \leq j \leq n.$$

These F_i for $i = 0, \dots, n$ are known from corollary 3.2.

Due to the Toeplitz structure and self adjointness of the matrix, T_Q , the above equations give us the following factorization:

$$T_Q = T_F^* T_F$$

where

$$T_F = \begin{pmatrix} F_0 & & & & & & & \\ F_1 & F_0 \otimes I_d & & & & & & \\ F_2 & F_1 \otimes I_d & F_0 \otimes I_d^{\otimes 2} & & & & & \\ \vdots & \vdots & \vdots & \ddots & & & & \\ F_n & F_{n-1} \otimes I_d & F_{n-2} \otimes I_d^{\otimes 2} & \cdots & F_0 \otimes I_d^{\otimes n} & & & \\ 0 & F_n \otimes I_d & F_{n-1} \otimes I_d^{\otimes 2} & \cdots & F_1 \otimes I_d^{\otimes n} & F_0 \otimes I_d^{\otimes (n+1)} & & \\ 0 & 0 & F_n \otimes I_d^{\otimes 2} & \cdots & F_2 \otimes I_d^{\otimes n} & F_1 \otimes I_d^{\otimes (n+1)} & F_0 \otimes I_d^{\otimes (n+2)} & \\ \vdots & \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots \end{pmatrix}.$$

In compact representation of T_F , recall $F_i = \text{col}(F_v)_{|v|=i}$. Thus the operator corresponding to the above matrix is,

$$T_F = \sum_{0 \leq |v| \leq n} F_v \otimes L_v.$$

Therefore, we have a multi-toeplitz factorization for a multi-toeplitz positive semidefinite matrix, T_Q which is of the form $T_F^* T_F$. \square

Remark: The above polynomial operator, T_Q (since its a polynomial in the left shifts, L) is R -multi-Toeplitz operator. Analogously, we have similar result for the polynomial operators in right shifts.

REFERENCES

- [1] V. Paulsen, *Completely Bounded Maps and Operator Algebras*, ser. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2003.
- [2] M. Rosenblum, “Vectorial Toeplitz operators and the Fejér-Riesz theorem,” *J. Math. Anal. Appl.*, vol. 23, pp. 139–147, 1968. [Online]. Available: [https://doi.org/10.1016/0022-247X\(68\)90122-4](https://doi.org/10.1016/0022-247X(68)90122-4)
- [3] M. Dritschel and H. Woerdeman, “Outer factorizations in one and several variables,” *Transactions of the American Mathematical Society*, vol. 357, no. 11, pp. 4661–4679, 2005.
- [4] G. Popescu, “Multi-analytic operators on Fock spaces,” *Math. Ann.*, vol. 303, no. 1, pp. 31–46, 1995. [Online]. Available: <https://doi.org/10.1007/BF01460977>
- [5] C. Foias and A. E. Frazho, *The commutant lifting approach to interpolation problems*, ser. Operator Theory: Advances and Applications. Birkhäuser Verlag, Basel, 1990, vol. 44. [Online]. Available: <https://doi.org/10.1007/978-3-0348-7712-1>