

Uncertainty relations from graph theory

Carlos de Gois,^{*} Kiara Hansenne,^{*} and Otfried Gühne

Naturwissenschaftlich-Technische Fakultät, Universität Siegen, Walter-Flex-Straße 3, 57068 Siegen, Germany

Quantum measurements are inherently probabilistic and quantum theory often forbids to precisely predict the outcomes of simultaneous measurements. This phenomenon is captured and quantified through uncertainty relations. Although studied since the inception of quantum theory, the problem of determining the possible expectation values of a collection of quantum measurements remains, in general, unsolved. By constructing a close connection between observables and graph theory, we derive uncertainty relations valid for any set of dichotomic observables. These relations are in many cases tight and related to the size of the maximum clique of the associated graph. As applications, our results can be straightforwardly used to formulate entropic uncertainty relations, separability criteria and entanglement witnesses.

Introduction.— Physical systems are characterised through the results of measurements, and in quantum theory measurements are described by observables. Mathematically, these are linear operators with real eigenvalues, each of which represents a possible result of a measurement. Born's rule allows to compute for each of these outcomes the probability of happening, depending solely on the state and the observable in question. The mathematical structure of quantum theory, however, enforces constraints on the probabilities, for instance if a pair of non-commuting observables is considered. These constraints frequently manifest themselves as uncertainty relations [1–8]. More generally, significant research effort has been devoted to the problem of determining the set of possible expectation values for sets of observables [9–13], which is of high relevance for various applications, such as quantum cryptography, entanglement characterisation and quantum metrology.

The main problem can be explained with a simple, yet fundamental example. Consider a single spin-1/2 particle (or qubit) and let X, Y, Z denote the Pauli matrices [14]. Then, it is well known that for any quantum state the relation

$$\langle X \rangle^2 + \langle Y \rangle^2 + \langle Z \rangle^2 \leq 1 \quad (1)$$

holds [15], but it also gives rise to the uncertainty relation $\Delta^2(X) + \Delta^2(Y) + \Delta^2(Z) \geq 2$, where $\Delta^2(A) = \langle A^2 \rangle - \langle A \rangle^2$ denotes the variance. This uncertainty relation is not only of fundamental interest, but it also found application in entanglement characterisation [16] and quantum cryptography [17]. But how can this relation be generalised?

To do that, it is natural to consider the expression $\mathbb{E} = \sum_i \langle A_i \rangle_\rho^2$, where each A_i is an observable with spectrum ± 1 , and ρ a quantum state. As a simple generalisation of Eq. (1) it is known that, for any set of pairwise anticommuting observables, $\mathbb{E} \leq 1$ [5, 9, 13]. This inequality led to the derivation of entropic uncertainty relations for multiple observables [13, 18], Bell monogamy inequalities [19] and quantum network state compatibility criteria [20], among others. Moreover, it leads to an uncertainty relation for multiple observables, $\sum_{i=1}^n \Delta^2(A_i) \geq n - 1$. Similar uncertainty relations have been previously studied




FIG. 1. For any set $\{A_i\}_{i=1}^n$ of observables, we draw a graph G with one vertex corresponding to each observable and an edge $\{i, j\} \in G$ if and only if the anticommutator $\{A_i, A_j\} \neq 0$. Its complement, \bar{G} , is called anticommutativity graph, and the Lovász number $\vartheta(\bar{G})$ provides an upper bound to \mathbb{E} .

[21–26], but the results were mainly applicable to two or three observables and not to larger sets.

Generalisations of \mathbb{E} to other sets of observables are thus scarce. One trivial fact is that $\mathbb{E} \leq n$, where n is the number observables employed. A more insightful fact is that, when $\{A_i\}$ contains d^2 operators that span the space of hermitian operators acting on a d -dimensional Hilbert space, with $\text{tr}(A_i A_j) = d\delta_{ij}$, then $\mathbb{E} \leq d$. This follows from the positivity of the density operator and can be interpreted as an upper bound on the length of its Bloch vector [27, 28]. Further than combinations of these facts (for instance, one may divide a set of observables into subsets, such that in each subset the observables anticommute), not much is known.

In this paper we derive nontrivial bounds on \mathbb{E} for any set of dichotomic observables by connecting this problem to graph theory. More specifically, to any set of observables we associate a graph encoding the anticommutativity relations (see also Fig. 1) and show that graph-theoretic quantities, such as the Lovász number or the order of a maximum clique have a physical interpretation. For

plenty of cases — for instance, for all perfect graphs — the obtained bounds and uncertainty relations are tight. Then, we discuss how our method leads to generalisations of entropic uncertainty relations [13, 18, 29], to applications in entanglement detection via local uncertainty relations [16, 30], to a simple procedure for generating entanglement witnesses [31] and their entropic counterparts [32] and to links with the sector lengths problem [33, 34].

The main connection. — Without loss of generality, we will consider observables that square to the identity, as any dichotomic observable can be rescaled to have a ± 1 -spectrum. To find an upper bound for \mathbb{E} , we start by defining E as a linear combination of the given observables depending on a state ϱ , that is, $E_\varrho = \sum_{i=1}^n a_i A_i$, with $a_i = \langle A_i \rangle_\varrho \in \mathbb{R}$. The variance of this observable on the same state ϱ is non-negative, hence $\langle E_\varrho^2 \rangle \geq \langle E_\varrho \rangle^2$ (here and later on, we will omit the dependency of expectation values on ϱ). While the right-hand-side is equal to $[\sum_i \langle A_i \rangle^2]^2 = \|\vec{a}\|^2 \sum_i \langle A_i \rangle^2$, the l.h.s. can be rewritten as

$$\sum_{ij} a_i a_j \langle A_i A_j \rangle = \sum_{ij: \{A_i, A_j\} \neq 0} a_i a_j \Re(\langle A_i A_j \rangle), \quad (2)$$

where we used that $\langle A_j A_i \rangle = \langle A_i A_j \rangle^*$. This sum is upper bounded by $\|\vec{a}\|^2 \Lambda(\mathcal{A})$, where $\Lambda(\mathcal{A})$ is the largest eigenvalue of the matrix \mathcal{A} whose elements \mathcal{A}_{ij} are $\Re(\langle A_i A_j \rangle)$.

Let us further specify \mathcal{A} . When $\{A_i, A_j\} = 0$ (i.e., a pair of observables anticommute), then $\mathcal{A}_{ij} = 0$. As the observables are required to have ± 1 -spectra, they square to the identity, thus the diagonal elements of \mathcal{A} are equal to one. Furthermore, the matrix \mathcal{A} is real and positive semidefinite. So, for any state ϱ ,

$$\sum_{i=1}^n \langle A_i \rangle^2 \leq \max_{\mathcal{A}} \Lambda(\mathcal{A}) \quad (3a)$$

$$\text{subject to: } \mathcal{A} \in \mathbb{R}^{n \times n} \text{ and } \mathcal{A} \succeq 0, \quad (3b)$$

$$\mathcal{A}_{ii} = 1, \forall i \in \{1, \dots, n\}, \quad (3c)$$

$$\mathcal{A}_{ij} = 0, \forall \{i, j\} : \{A_i, A_j\} = 0. \quad (3d)$$

To establish the connection to graph theory, we define the *anticommutativity graph* \bar{G} as the one with n vertices — each corresponding to an observable — connected by an edge if and only if $\{A_i, A_j\} = 0$, see also Fig. 1. For general graphs, there are several quantities of interest. The *clique number* $\omega(G)$ is the size of the largest subgraph of G that is fully connected and the *chromatic number* $\chi(G)$ is the smallest number of colours needed to colour the vertices of G such that no connected vertices share the same colour. The *Lovász number*, also called ϑ -function, was originally introduced as an upper bound to the so-called Shannon capacity of a graph [35], and can be defined in different equivalent ways (see also below). Remarkably, it can also be used to estimate the previously mentioned quantities: The sandwich theorem [35] states that $\omega(G) \leq \vartheta(\bar{G}) \leq \chi(G)$. Here, G and \bar{G} are complementary graphs, i.e. vertices which are connected in G are disconnected in

\bar{G} and vice versa. The sandwich theorem is of particular interest since, whereas determining $\omega(G)$ and $\chi(G)$ is an NP-complete problem [36], $\vartheta(\bar{G})$ can be computed through a semidefinite program [35]. A graph G for which these three quantities are equal for all its induced subgraphs is called a perfect graph.

Among the many, equivalent definitions of the Lovász number $\vartheta(G)$, one is particularly interesting for our problem. As shown in Section 33 of [37] (up to a typo in Eq. 33.3¹),

$$\vartheta(G) = \max_B \Lambda(B) \quad (4a)$$

$$\text{subject to: } B \in \mathbb{R}^{n \times n} \text{ and } B \succeq 0 \quad (4b)$$

$$B_{ii} = 1, \forall i \in \{1, \dots, n\} \quad (4c)$$

$$B_{ij} = 0, \forall \{i, j\} \in G. \quad (4d)$$

This exactly matches the quantity in Eq. (3), leading to a generalisation of Eq. (1) as our main result:

$$\sum_{i=1}^n \langle A_i \rangle^2 \leq \vartheta(\bar{G}). \quad (5)$$

Note that these bounds are state independent; when the purity of the states is known, the Lovász number can also be employed to bound similar expressions in various ways [38].

Immediate consequences. — Complete anticommutativity graphs originate from observables that all anticommute pairwise. In this case, in Eq. (4), B is the identity matrix. Then, $\vartheta(\bar{G}) = 1$, recovering the result derived in Refs. [9, 13]. Oppositely, the Lovász number of an empty graph is n , leading to the trivial bound $\sum_i \langle A_i \rangle^2 \leq n$.

As stated in the introduction, the smallest number of pairwise anticommuting subsets of $\{A_i\}$ is an upper bound for \mathbb{E} . In fact, this corresponds to the chromatic number of the associated G , which is always an upper bound on the Lovász number. Since there exist graphs for which $\vartheta(\bar{G}) < \chi(G)$ (such as the pentagon in Fig. 1) pairwise anticommutation relations alone do not always lead to the best upper bounds.

If one considers tensor products of Pauli matrices, the observables A_i have the additional property that they can only commute or anticommute. In this case, one can clearly achieve $\sum_i \langle A_i \rangle^2 = \omega(G)$ by choosing a common eigenstate for the largest commuting set. So, if one has a perfect graph for this scenario, then $\omega(G) = \vartheta(\bar{G})$ and the bound in Eq. (5) is tight. Interesting perfect graphs include all bipartite graphs, even cycles, and chordal graphs. In fact, from the *strong perfect graph theorem*, a graph is imperfect if and only if it contains some odd cycle larger or equal than a pentagon (or the complement

¹ Therein, the “dual feasible for G ” is defined under three conditions, one of which states that $B_{uv} = 0$ whenever the edge $\{i, j\}$ is *not* in G . The correct condition, however, is for when the edge *is* in G .

of one) as an induced subgraph [39]. Therefore, these are the only cases where the bound in Eq. (5) may not be tight — the smallest example of such a possibility being the pentagon itself.

Also notably, there is a similar connection between linear expressions on squares of the expectation values, $\mathbb{E}_{\vec{w}} = \sum_{i=1}^n w_i \langle A_i \rangle^2$, where $\vec{w} = (w_1, \dots, w_n)$, $w_i > 0$, and the *weighted Lovász number* $\vartheta(\bar{G}, \vec{w})$. This is evidenced from the fact that, when considering $\mathbb{E}_{\vec{w}}$, Eq. (3c) (and only that) changes to $\mathcal{A}_{ii} = w_i$. That is precisely how the representation of the Lovász number given in Eq. (4) changes in its weighted variation. Consequently,

$$\sum_{i=1}^n w_i \langle A_i \rangle^2 \leq \vartheta(\bar{G}, \vec{w}). \quad (6)$$

Additionally, notice that observables with arbitrary spectrum (i.e., any set of hermitian operators) can also be addressed by the weighted Lovász number. Indeed, in this case, the elements \mathcal{A}_{ii} are no longer equal to one but are upper bounded by $\Lambda(A_i^2)$. This leads to $\sum_{i=1}^n \langle A_i \rangle^2 \leq \vartheta(\bar{G}, \vec{w})$, where $\vec{w} = (\Lambda(A_1^2), \dots, \Lambda(A_n^2))$. Clearly, the case for dichotomic observables is recovered by unit weights. Computing $\vartheta(\bar{G}, \vec{w})$ is also a small instance of a semidefinite program, thus the bound is easily computable [37].

A stronger conjecture. — Backed by extensive numerical evidence, we conjecture that, for the case where all observables are dichotomic and each pair of them either commute or anticommute, the tight bound for \mathbb{E} is actually $\omega(G)$, i.e., the size of the largest set of pairwise commuting observables in the set.

As an example, consider the three-qubit observables $X\mathbb{1}\mathbb{1}$, $\mathbb{1}X\mathbb{1}$, $Z\mathbb{1}X$, $ZZ\mathbb{1}$ and $\mathbb{1}ZZ$ (we omitted the tensor product for readability). Their anticommutativity graph \bar{G} is a pentagon (Fig. 1), up to reordering of the vertices, for which $\vartheta(\bar{G}) = \sqrt{5}$. Hence we get $\mathbb{E} \leq \sqrt{5}$, which is better than the bound $\mathbb{E} \leq 3$ obtained by exploiting the chromatic number. One can also compute \mathbb{E} by maximising $S = \sum_i A_i \otimes A_i$ over all product states $\varrho_A \otimes \varrho_B$. This maximisation can be relaxed by optimising over states with a positive partial transpose [40], which is a semidefinite program with a certified solution. In this case this proves that, up to numerical precision, the quantum maximum is $\mathbb{E} = 2$, which is equal to $\omega(G)$. This can be achieved, for instance, with the state $|000\rangle$, which is a common eigenstate of $ZZ\mathbb{1}$ and $\mathbb{1}ZZ$.

The conjecture is in line with the upper bound on the length of the Bloch vector mentioned in the introduction, since for $d \times d$ -dimensional operators, the largest number of linearly independent pairwise commuting observables (i.e., $\omega(G)$) is d .

Another noteworthy example is taking the twenty-seven observables $\{X, Y, Z\}^{\otimes 3}$. The largest subset of pairwise commuting observables is of size 4, whereas $\vartheta(\bar{G}) = 5$. It was previously shown that the actual tight bound for

these observables is $\mathbb{E} \leq 4$ [41], in agreement with our conjecture.

More generally, our conjecture is also in line with tight bounds on full-body sector lengths for any number of qubits [33]. These quantities are defined for N -qubit states as $L_N = \sum_{\sigma \in \{X, Y, Z\}^{\otimes N}} \langle \sigma \rangle^2$ and quantify the amount of full body correlations in quantum states. Trivially, we get $\omega(G) \leq \max L_N$. Furthermore, it is known that $\max L_N$ is achieved by the N -qubit GHZ state, as it is stabilised by exactly $\max L_N$ full body elements [41, 42]. Since stabiliser elements pairwise commute, it follows that $\omega(G) \geq \max L_N$. Hence we have $\max L_N = \omega(G)$. The bound can be expressed as $\max L_N = 2^{N-1} + 1/2(1 + (-1)^N)$.

Application: Entanglement characterisation. — Recall that a two-party state ϱ is separable ($\varrho \in \text{SEP}$) if it can be written as a convex combination of product states, $\varrho = \sum_k p_k \varrho_k^A \otimes \varrho_k^B$. Rewriting Eq. (5) and making use of the Cauchy–Schwarz inequality, we get

$$\max_{\varrho \in \text{SEP}} \sum_{i=1}^n \langle A_i \otimes B_i \rangle \leq \sqrt{\vartheta(\bar{G}_A) \vartheta(\bar{G}_B)} =: \vartheta_{AB}, \quad (7)$$

where $\vartheta(\bar{G}_A)$ is the Lovász number of the anticommutativity graph of $\{A_i\}$ and similarly for $\vartheta(\bar{G}_B)$. This separability criterion naturally leads to entanglement witnesses of the form [31]

$$W_{\mathbb{E}} = \vartheta_{AB} \mathbb{1} - \sum_i A_i \otimes B_i. \quad (8)$$

These operators are required to have $\text{tr}(W_{\mathbb{E}} \varrho) \geq 0$ for separable ϱ , which directly follows, and to have $\text{tr}(W_{\mathbb{E}} \varrho) < 0$ for at least one entangled state, which is fulfilled when the observables are chosen insightfully. For cases where the $\vartheta(\bar{G}_A)$ and $\vartheta(\bar{G}_B)$ bounds are tight, $W_{\mathbb{E}}$ is a weakly optimal entanglement witness, i.e., there exists at least one separable state ϱ_s such that $\text{tr}(W_{\mathbb{E}} \varrho_s) = 0$.

As an example, let us take the two-quart PPT entangled state

$$\varrho_I = \frac{1}{6} (\varrho_{\mathbb{1}Y} + \varrho_{XX} + \varrho_{YZ} + \varrho_{ZX} + \varrho_{ZY} + \varrho_{ZZ}), \quad (9)$$

where ϱ_{AB} is the projector onto the pure state $|\psi_{AB}\rangle = 1/2 \left(\mathbb{1} \mathbb{1} AB \sum_{i=0}^3 |i\rangle |i\rangle \right)$, $A, B \in \{\mathbb{1}, X, Y, Z\}$ [43]. Physically, this state can be seen as a mixture of six pairs of Bell states ($|\beta_i\rangle$) of the form $|\beta_i\rangle_{A'B'} \otimes |\beta_j\rangle_{A''B''}$, where one party holds the qubits A' and A'' , and the other B' and B'' [43, 44]. As sets of observables, we take

$$\{A_i/B_i\} = \{\pm \mathbb{1}X, \mathbb{1}Y, \pm \mathbb{1}Z, \pm X\mathbb{1}, XX, \pm XY, \pm XZ, Y\mathbb{1}, YX, YY, \pm YZ, Z\mathbb{1}, \pm ZX, \pm ZY, \pm ZZ\}, \quad (10)$$

where the upper signs are for $\{A_i\}$. These sets have the same anticommutativity graph, for which $\omega(G) = \vartheta(\bar{G}) = 3$. The witness based on these observables detects

$\nu \varrho_I + (1 - \nu)^{1/16}$ to be entangled for $\nu > 3/5$, which is exactly the separability bound [44].

Application: Uncertainty relations.— As explained already in the introduction, our result in Eq. (5) gives rise to the uncertainty relation

$$\sum_{i=1}^n \Delta^2(A_i) \geq n - \vartheta(\bar{G}). \quad (11)$$

Using the approach of so-called local uncertainty relations [16] this can be turned into an entanglement criterion: Suppose that a local uncertainty relation $\sum_{i=1}^n \Delta^2(A_i)_{\varrho_A} \geq C_A$ is true for all d_A -dimensional quantum states ϱ_A and $\sum_{i=1}^n \Delta^2(B_i)_{\varrho_B} \geq C_B$, similarly, for any state of dimension d_B . Then, if we define on the bipartite system $M_i \equiv A_i \otimes \mathbb{1} + \mathbb{1} \otimes B_i$, for separable states, $\sum_{i=1}^n \Delta^2(M_i)_{\varrho} \geq C_A + C_B$ holds. Combining this method with Eq. (11) leads to nonlinear witnesses of the form

$$\begin{aligned} & \frac{1}{2} [\vartheta(\bar{G}_A) + \vartheta(\bar{G}_B)] - \sum_{i=1}^n \langle A_i \otimes B_i \rangle - \\ & - \frac{1}{2} \sum_{i=1}^n \langle A_i \otimes \mathbb{1} - \mathbb{1} \otimes B_i \rangle_{\text{SEP}}^2 \geq 0, \end{aligned} \quad (12)$$

where \bar{G}_A and \bar{G}_B are the anticommutativity graphs corresponding to $\{A_i\}$ and $\{B_i\}$, respectively. If $\vartheta(\bar{G}_A) = \vartheta(\bar{G}_B)$ this criterion is obviously stronger than Eq. (8), see also Ref. [45].

From the variance uncertainty relation in Eq. (11), it is also possible to derive several entropic URS. Entropic URS are sometimes considered to be a more fundamental formulation of URS, and can be used in security proofs for quantum key distribution and as entanglement witnesses, among other applications [6].

As an example, when $\vartheta(\bar{G})$ is an integer (e.g., for perfect graphs), Theorem 6 in Ref. [18] is straightforwardly generalisable to our case and leads to the tight bound

$$\sum_{i=1}^n S(A_i|\varrho) \geq [n - \vartheta(\bar{G})] S_0. \quad (13)$$

Here, S is any entropy that is concave in $\langle A_i \rangle^2$,² $S(A_i|\varrho)$ is the entropy of the probability distribution gotten from measuring A_i on ϱ and S_0 is the entropy value of the flat probability distribution. The proof of the statement proceeds just as in Ref. [18], with the difference that the allowed domain of values for $\langle A_1 \rangle^2, \dots, \langle A_n \rangle^2$, has now as extremal points all vectors with up to $\vartheta(\bar{G})$ entries “1” and zeroes elsewhere. Prominent examples of entropies that have the desired concavity property, are the Shannon

² Concavity in $\langle A_i \rangle^2$ here means that the function $\tilde{S}(x) := S(A_i|\varrho) = S(P_{\pm})$ is concave in $x = \langle A_i \rangle^2$, where $P_{\pm} = (1 \pm \sqrt{x}/2, 1 \mp \sqrt{x}/2)$ are the two possible probability distribution arising from measuring A_i .

entropy and the Tsallis entropies S_q^T for $1 < q < 2$ and $3 < q$ [18].

Interestingly, Eq. (13) can easily be turned into entropic criteria for steering and entanglement. To understand how, suppose that $\sum_{i=1}^n S(A_i|\varrho_A) \geq C$ holds for all d_A -dimensional quantum states ϱ_A . Then, for all separable bipartite $d_A d_B$ -dimensional states and any set of n observables $\{B_i\}$, $\min_{\varrho \in \text{SEP}} \sum_{i=1}^n S(A_i \otimes B_i|\varrho) \geq C$ [32]. Even more strongly, all probability distributions arising from measuring any unsteerable state fulfil the same condition, thus these same expressions can also be used as entropic steering criteria [46].

As a closing comment we recall that, in the two-dimensional case, $\Delta^2(X) + \Delta^2(Y) + \Delta^2(Z) \geq 2$ completely characterises the set of states, in the sense that any probability distribution obeying it can originate from a valid quantum state. This also holds for the extension to larger sets of pairwise anticommuting sets of observables but is, however, not true in general for the constraints in Eq. (11). To build an example, start from the Pauli basis in dimension $d = 4$, given by $\{A_i\} = \{\mathbb{1}, X, Y, Z\}^{\otimes 2}$. In this case, $\vartheta(\bar{G}) = 4$, and from Eq. (5), $\sum_{i=1}^{16} \langle A_i \rangle_{\varrho}^2 \leq 4$, implying that the length of the Bloch vector of any ququart is bounded by 4. But it is well-known that there are vectors of length 4 which do not represent quantum states [27, 28], therefore there must be higher-order constraints not encoded in Eq. (11). Finding such relations is an interesting topic for further research.

Discussion.— Through exploring a link between observables and graph theory, we have proven that the sum of squares of expectation values of any set of dichotomic observables is bounded by the Lovász number of their anticommutativity graph. This found further applications for the characterisation of entanglement and quantum steering, and also in various uncertainty relations.

Regarding extensions of our results, proving or disproving the conjecture discussed after Eq. (6) is an interesting open problem in itself. Our current evidence is based on exhaustive studies of sets of Pauli observables. Another intriguing idea is to consider observables with more than two outcomes. This would lead to more general uncertainty relations, of which not much is currently known. A related, perhaps more practically-minded concern is to investigate how the ϑ -bound changes in the presence of a perturbation such that, for some pairs of observables, $|\{A_i, A_j\}| < \epsilon$. This is in line with approaches to devise entanglement tests with not perfectly calibrated observables [44, 47, 48].

It could also be insightful to consider other definitions of the Lovász number. Our choice, presented in Eq. (4), was based on the straightforward connection with Eq. (3). However, others may provide further physical intuition. Of particular interest is the representation given in Theorem 5 of [35], which can be more closely related to quantum structures, such as state vectors and measurements. Indeed, it was already shown to be connected to quantum

bounds in contextuality inequalities [49]. Similarly, it could also be interesting to find closer correspondences with other approaches to similar problems, such as the one via joint numerical ranges (e.g., [23, 25]).

Applications-wise, the entanglement witnesses we have shown are just simple examples, and much in the way of tailoring them to particular families of states can be done. Apart from standard entanglement theory, bounds for special cases of \mathbb{E} were previously applied to the network entanglement compatibility problem [20], and we believe our more general bounds can be further employed in network scenarios and lead to stronger certificates. Moreover, our results imply bounds on the trace of covariance matrices, which can be used as network entanglement criteria as well. Finally, establishing bounds for the case of arbitrary weights in Eq. (6) would be a potentially very fruitful generalisation of the theory presented here.

Note added. — A counterexample to the discussion in the section “A stronger conjecture” was recently found by Z.-P. Xu and R. Schwonnek [50].

Acknowledgments. — Many thanks to J. Carrasco, D. García-Martín, F. Huber, B. Kraus and A. Perez-Salinas (among many other participants of the “Entanglement in Action” conference in Benasque), and also to J. L. Bönsel, M. Terra Cunha, L. Ligthart, H. C. Nguyen, T. Ohst, M. Plávala, R. Schwonnek, L. Vandr e, R. Werner, Z.-P. Xu, and B. Yadin for interesting discussions. This work was financially supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, project numbers 447948357 and 440958198), the Sino-German Center for Research Promotion (Project M-0294), and the House of Young Talents of the University of Siegen.

* These authors contributed equally to this paper.

- [1] W. Heisenberg, *Z. Physik* **43**, 172 (1927).
- [2] E. H. Kennard, *Z. Physik* **44**, 326 (1927).
- [3] H. P. Robertson, *Phys. Rev.* **34**, 163 (1929).
- [4] H. P. Robertson, *Phys. Rev.* **46**, 794 (1934).
- [5] S. Wehner and A. Winter, *New J. Phys.* **12**, 025009 (2010).
- [6] P. J. Coles, M. Berta, M. Tomamichel, and S. Wehner, *Rev. Mod. Phys.* **89**, 015002 (2017).
- [7] P. Busch, P. Lahti, and R. F. Werner, *Rev. Mod. Phys.* **86**, 1261 (2014).
- [8] R. Schwonnek, *Uncertainty relations in quantum theory*, Ph.D. thesis, Gottfried Wilhelm Leibniz Universit at Hannover (2018).
- [9] G. T oth and O. G uhne, *Phys. Rev. A* **72**, 022340 (2005).
- [10] K. Szymański, S. Weis, and K. Życzkowski, *Linear Algebra Appl.* **545**, 148 (2018).
- [11] T. Schulte-Herbr uggen, G. Durr, U. Helmke, and S. J. Glaser, *Lin. Multilin. Alg.* **56**, 3 (2008).
- [12] P. Gawron, Z. Puchała, J. A. Miszczak, L. Skowronek, and K. Życzkowski, *J. Math. Phys.* **51**, 102204 (2010).
- [13] S. Wehner and A. Winter, *J. Math. Phys.* **49**, 062105 (2008).
- [14] W. Pauli, *Z. Physik* **43**, 601 (1927).
- [15] R. P. Feynman, F. L. Vernon Jr, and R. W. Hellwarth, *J. Appl. Phys.* **28**, 49 (1957).
- [16] H. F. Hofmann and S. Takeuchi, *Phys. Rev. A* **68**, 032103 (2003).
- [17] C. Portmann and R. Renner, *Rev. Mod. Phys.* **94**, 025008 (2022).
- [18] S. Niekamp, M. Kleinmann, and O. G uhne, *J. Math. Phys.* **53**, 012202 (2012).
- [19] P. Kurzyński, T. Paterek, R. Ramanathan, W. Laskowski, and D. Kaszlikowski, *Phys. Rev. Lett.* **106**, 180402 (2011).
- [20] K. Hansenne, Z.-P. Xu, T. Kraft, and O. G uhne, *Nat. Commun.* **13**, 1 (2022).
- [21] P. Busch, P. Lahti, and R. F. Werner, *Phys. Rev. A* **89**, 012129 (2014).
- [22] L. Maccone and A. K. Pati, *Phys. Rev. Lett.* **113**, 260401 (2014).
- [23] K. Szymański and K. Życzkowski, *J. Phys. A: Math. Theor.* **53**, 015302 (2019).
- [24] A. A. Abbott, P.-L. Alzieu, M. J. W. Hall, and C. Branciard, *Mathematics* **4** (2016).
- [25] R. Schwonnek, L. Dammeier, and R. F. Werner, *Phys. Rev. Lett.* **119**, 170404 (2017).
- [26] P. Giorda, L. Maccone, and A. Riccardi, *Phys. Rev. A* **99**, 052121 (2019).
- [27] G. Kimura, *Phys. Lett. A* **314**, 339 (2003).
- [28] M. S. Byrd and N. Khaneja, *Phys. Rev. A* **68**, 062322 (2003).
- [29] J. Kaniewski, M. Tomamichel, and S. Wehner, *Phys. Rev. A* **90**, 012332 (2014).
- [30] O. G uhne, *Phys. Rev. Lett.* **92**, 117903 (2004).
- [31] O. G uhne and G. T oth, *Phys. Rep.* **474**, 1 (2009).
- [32] O. G uhne and M. Lewenstein, *Phys. Rev. A* **70**, 022316 (2004).
- [33] H. Aschauer, J. Calsamiglia, M. Hein, and H. J. Briegel, *Quantum. Inf. Comput.* **4**, 383 (2004).
- [34] N. Wyderka and O. G uhne, *J. Phys. A: Math. Theor.* **53**, 345302 (2020).
- [35] L. Lov asz, *IEEE Trans. Inf. Theory* **25**, 1 (1979).
- [36] R. M. Karp, in *Complexity of computer computations* (Springer, 1972) pp. 85–103.
- [37] D. E. Knuth, *Electron. J. Comb.* **1**, A1 (1994).
- [38] Z.-P. Xu, Private communication (2022).
- [39] M. Chudnovsky, N. Robertson, P. Seymour, and R. Thomas, *Ann. of Math.* **51** (2006).
- [40] A. Peres, *Phys. Rev. Lett.* **77**, 1413 (1996).
- [41] M. C. Tran, B. Dakić, W. Laskowski, and T. Paterek, *Phys. Rev. A* **94**, 042302 (2016).
- [42] C. Eltschka and J. Siewert, *Quantum* **4**, 229 (2020).
- [43] F. Benatti, R. Floreanini, and M. Piani, *Open Syst. Inf. Dyn.* **11**, 325 (2004).
- [44] T. Moroder and O. Gittsovich, *Phys. Rev. A* **85**, 032301 (2012).
- [45] O. G uhne, M. Mechler, G. T oth, and P. Adam, *Phys. Rev. A* **74**, 010301 (2006).
- [46] A. C. S. Costa, R. Uola, and O. G uhne, *Phys. Rev. A* **98**, 050104 (2018).
- [47] M. Seevinck and J. Uffink, *Phys. Rev. A* **76**, 042105 (2007).
- [48] S. Morelli, H. Yamasaki, M. Huber, and A. Tavakoli, *Phys. Rev. Lett.* **128**, 250501 (2022).
- [49] A. Cabello, S. Severini, and A. Winter, *Phys. Rev. Lett.* **112**, 040401 (2014).
- [50] Z.-P. Xu and R. Schwonnek, In preparation (2022).