

Noncrystalline topological superconductors

Sourav Manna,¹ Sanjib Kumar Das,² and Bitan Roy²

¹*Department of Condensed Matter Physics, Weizmann Institute of Science, Rehovot 7610001, Israel*

²*Department of Physics, Lehigh University, Bethlehem, Pennsylvania, 18015, USA*

Topological insulators, featuring bulk-boundary correspondence, have been realized on a large number of noncrystalline materials, among which amorphous network, quasicrystals and fractal lattices are the most prominent ones. By contrast, topological superconductors beyond the realm of quantum crystals are yet to be harnessed, as their nucleation take place around a well-defined Fermi surface with a Fermi momentum, the existence of which rests on the underlying translational symmetry. Here we identify a family of noncrystalline Dirac materials, devoid of translational symmetry, on which suitable local or on-site pairings yield topological superconductors. We showcase this outcome on all the above mentioned noncrystalline platforms embedded on a two-dimensional flat space. The resulting noncrystalline topological superconductors possess a quantized topological invariant (Bott index) and harbor robust one-dimensional Majorana edge modes.

Introduction. Topological insulators constitute the foundation of topological classification of quantum crystals. In particular, when the electronic wavefunction of a quantum insulator acquires nontrivial geometry in the Brillouin zone, robust gapless modes of charged quasiparticles appear at the interface of topological crystals [1–14]. Such boundary modes manifest the bulk topological invariant of electronic wavefunction, a phenomenon named the bulk-boundary correspondence, with quantum Hall insulators standing as its paradigmatic examples [15, 16]. Furthermore, topological classification extends beyond the realm of charged fermions, and it is equally applicable to neutral Bogoliubov de-Gennes (BdG) quasiparticles, opening a rich landscape of topological superconductors [17, 18]. The bulk-boundary correspondence is also germane to such paired states, resulting in gapless boundary modes of Majorana fermions.

As over the time it became evident that topological insulators can be realized on noncrystalline materials, such as amorphous networks [19–25], quasicrystals [26–29] and fractal lattices [30–40], a question of fundamental and practical importance arises naturally. *Can topological superconductors be realized on noncrystalline materials?* The subtlety of this quest stems from the fact that condensation of electrons into a macroscopic number of Cooper pairs in any real material takes place around a Fermi surface, defined via a Fermi momentum, which roots in the underlying translational symmetry, as in crystals. Here we provide an affirmative answer to this question, by identifying a family of noncrystalline Dirac materials (NCDMs), encompassing amorphous network, quasicrystals and fractal lattices, on which suitable local or on-site pairings give rise to nontrivial topology of BdG quasiparticles [Fig. 1], and the resulting noncrystalline topological superconductors (NCTSCs) support topological boundary modes of Majorana fermions [Fig. 2].

Although here we exclusively focus on two-dimensional (2D) NCDMs, the underlying mechanism behind the robust realization of NCTSCs is sufficiently general, which can be straightforwardly generalized to three dimensions. Therefore, with the recent experimental realizations of topological insulator in three-dimensional amor-




FIG. 1. Cuts of the global phase diagram of NCTSCs on (a) (M, Δ) plane for zero chemical doping ($\mu = 0$) and (b) (M, μ) plane for a fixed pairing amplitude $\Delta = 1$ [Eq. (4)]. Throughout we set $t = t_0 = 1$ and $r_0 = 1$ [Eq. (3)]. We find identical phase diagrams on a square lattice and an amorphous network both containing 400 lattice sites, Penrose and Ammann-Beenker quasicrystals with 481 and 357 lattice sites, respectively, and third generation Sierpiński carpet and fifth generation glued Sierpiński triangle fractals, containing 512 and 454 lattice sites, respectively, with open boundary condition. This is so because the hopping elements in the normal state are infinitely long ranged, washing out microscopic details of the underlying noncrystalline order. We find NCTSCs with the topological Bott indices $B = -1$ and -2 [Eq. (5)], both of which support one-dimensional Majorana edge modes [Fig. 2], besides the trivial one with $B = 0$.

phous Bi_2Se_3 [41], our predicted NCTSCs can be identified in real materials, especially when noncrystalline (topological) insulators are doped, yielding finite number of charged carriers in the normal state, which at low temperatures can form a Cooper condensate. These findings should promote NCDMs onto a promising platform of Majorana fermion based applications: *Majotronics*.

Key results. To exemplify these general outcomes, here we consider a 2D time reversal symmetry breaking Dirac insulator, constituting the normal state. Depending on the parameter values, such a system can describe either a topological, namely quantum anomalous Hall insulator or a trivial/normal insulator. However, for the discussion of superconductivity and emer-




FIG. 2. Energy spectra of $H_{\text{pair}}^{\text{real}}$ [Eq. (4)] on (a) a square lattice containing 400 sites, where r_0 is the nearest-neighbor (NN) distance, (b) an amorphous network with 600 sites with $r_0 = 0.05 \times$ linear dimension of the system in each direction, (c) Penrose (481 sites) and (d) Ammann-Beenker (617 sites) quasicrystals, where r_0 is the arm length, (e) third generation Sierpiński carpet (512 sites) and (f) fifth generation glued Sierpiński triangle (454 sites) fractal lattices, where r_0 is the NN distance, with periodic (black) and open (red) boundary conditions, for $M = 2$, $t = t_0 = 1$ and $\Delta = 0.5$. We then realize a NCTSC in all these systems with the Bott index $B = -1$ [Fig. 1]. Local density of states (LDOS) for two closest to zero energy states (normalized by its maximum value) with open boundary condition on (g) square lattice, (i) Penrose and (j) Ammann-Beenker quasicrystals, (k) Sierpiński carpet and (l) glued Sierpiński triangle fractals, showing sharp edge localization. In (h) we show the total LDOS (normalized by its maximum value) for all the near zero energy states that can only be found with open boundary condition, displaying sharp edge localization. On two fractal lattices the spectra remain qualitatively unchanged with periodic and open geometries due to the inner edges. The LDOS of two closest to zero energy states in open (periodic) fractal lattices shows sharp localization dominantly around the outer (inner) edges. See also Fig. 3. We arrive at qualitatively similar results for NCTSCs with $B = -2$. The trivial paired states with $B = 0$ is devoid of any near zero energy edge modes.

gent topology therein, the normal state topology is unimportant. We implement the corresponding model Hamiltonian on a variety of 2D noncrystalline materials besides the square lattice, such as an amorphous network, Penrose and Ammann-Beenker quasicrystals, and Sierpiński carpet and glued Sierpiński triangle fractal lattices. Subsequently, we introduce the only local or on-site pairing in this model and show that it translates into a first-order topological pairing in all the noncrystalline materials, a phase we name as NCTSC. The corresponding global phase diagrams for zero and finite chemical doping are shown in Figs. 1(a) and (b), respectively, which besides unfolding NCTSCs also feature trivial pairing.

The NCTSCs are characterized by quantized Bott index (B). The global phase diagram supports NCTSCs with $B = -1$ and -2 , besides a trivial paired state with $B = 0$. All NCTSCs showcase the bulk-boundary correspondence in terms of one-dimensional (1D) Majorana edge modes in systems with open geometry [Fig. 2]. However, exclusively on two fractal lattices with open (periodic) geometry, the edge modes dominantly localize near the outer (inner) edges, as shown in Fig. 2 (Fig. 3). This unique phenomenon on fractal lattices solely stems from its self-similarity symmetry, resulting in inner boundaries in the interior of the system [42].

The edge localization of topological boundary modes is further anchored from the inverse participation ratio (IPR) [Fig. 4]. On crystal, amorphous network, quasicrystals containing N sites, the number of sites at the edges scales as \sqrt{N} . Consequently, the IPR for the near

zero energy modes scales (almost) linearly with \sqrt{N} , ensuring their sharp edge localization. By contrast, due to the inner boundaries there is no such connection between N and the number edge sites on fractals, and the IPR of closest to zero energy modes does not show linear scaling with \sqrt{N} on Sierpiński carpet or glued Sierpiński triangle fractal lattices. This outcome reconciles with (i) energy spectra of NCTSCs on fractal lattices are qualitatively sensitive to boundary conditions [Fig. 2(e), (f)] and (ii) on periodic fractal lattices the closest to energy modes dominantly reside on their inner edges [Fig. 3].

Model. The Hamiltonian for a 2D time-reversal sym-




FIG. 3. The LDOS for two closest to zero energy states on a (a) Sierpiński carpet and (b) glued Sierpiński triangle fractal lattices with periodic boundary condition, displaying their localization dominantly around the inner edges. Compare with Fig. 2(k) and (l), respectively.




FIG. 4. Inverse participation ratio (IPR), denoted by P (in units of 10^{-3} [Eq. (6)]), on (a) square lattice, (b) amorphous network, (c) Penrose and (d) Ammann-Beenker quasicrystals, (e) Sierpiński carpet and (f) glued Sierpiński triangle fractal lattices, containing N sites (see legends). In each subfigure, panel (1) shows IPR of all the states, panel (2) shows the scaling of the IPR for the closest to zero energy state (edge mode) as a function of \sqrt{N} , and panel (3) shows the scaling of a large energy state (bulk mode) as a function of N . Except on two fractal lattices, IPR in (2) scales linearly with \sqrt{N} , confirming sharp outer edge localization of near zero energy mode [Fig. 2]. Absence of such a linear scaling on fractal lattices stems from the existence of the inner boundaries, which host near zero energy modes with periodic geometry [Fig. 3].

metry breaking Dirac insulator is [5, 6]

$$H_{\text{DI}} = t \sum_{j=1,2} S_j \tau_j + \left[M - 2t_0 \sum_{j=1,2} [1 - C_j] \right] \tau_3, \quad (1)$$

where $S_j \equiv \sin(k_j a)$, $C_j \equiv \cos(k_j a)$, $\mathbf{k} = (k_1, k_2)$ is the spatial momenta, a is the lattice spacing, and the vector Pauli matrix $\boldsymbol{\tau} = (\tau_1, \tau_2, \tau_3)$ operates on the orbital space. Using Fourier transformation this model can be implemented on a square lattice. However, when we sacrifice the underlying crystalline order, as in NCDMs, the above model needs to be generalized such that it can be implemented on an arbitrary lattice system. This is accomplished by symmetry analyses of each term appearing in H_{DI} under reflections about the x and y directions, and four-fold rotation in the xy plane. We then replace each momentum dependent factor from H_{DI} by its corresponding term in the real space that transforms identically under all the symmetry operations, leading to

$$H_{\text{DI}}^{\text{real}} = \sum_{j \neq k} \frac{F(r_{jk})}{2} c_j^\dagger \left[-it(C_{jk}\tau_1 + S_{jk}\tau_2) + 2t_0 \tau_3 \right] c_k + \sum_j c_j^\dagger \left[(M - 4t_0) \tau_3 \right] c_j, \quad (2)$$

where $C_{jk} \equiv \cos \phi_{jk}$, $S_{jk} \equiv \sin \phi_{jk}$, $c_j = [c_{\alpha j}, c_{\beta j}]$, $c_{\tau j}$ is the fermion annihilation operator on site j with orbital $\tau = \alpha, \beta$. Hopping strengths between sites j and k , respectively placed at \mathbf{r}_j and \mathbf{r}_k , are accompanied by

$$F(r_{jk}) = \Theta(r_{jk} - R) \exp \left[1 - \frac{r_{jk}}{r_0} \right], \quad (3)$$

ensuring that all the sites are well connected. Here $r_{jk} = |\mathbf{r}_j - \mathbf{r}_k|$ is the distance and ϕ_{jk} is the azimuthal angle between them, R controls the range of hopping, and r_0 is the decay length. For the sake of simplicity, here we typically consider R to be larger than the system size, such that the hopping elements are effectively infinitely long ranged. This model on any lattice system supports both quantum anomalous Hall and normal insulators, which we do not discuss in any further details.

NCTSCs. In principle, H_{DI} can describe a paired state with suitable choice of the spinor basis [43]. For example, the term proportional to t describes a topological $p_x + ip_y$ pairing, when the term proportional to τ_3 yields a Fermi surface, as it happens for $0 < M/t_0 < 8$ on a square lattice. It is therefore tempting to conclude that $H_{\text{DI}}^{\text{real}}$ captures topological pairing on noncrystalline materials. However, such an attempt suffers physical shortcomings. Firstly, H_{DI} does not reveal any microscopic origin of the $p_x + ip_y$ pairing. Even more importantly, the pairing term in $H_{\text{DI}}^{\text{real}}$ becomes infinitely long-ranged connecting all the sites with decaying amplitude of the Cooper pairs, which is unphysical. Finally, the notion of a Fermi surface in the absence of an underlying translational symmetry becomes moot. To bypass these limitations, we search for a suitable material platform where *on-site* or *local* pairings give rise to topological superconductors even in the absence of the underlying translational symmetry.

NCDMs, described by the Hamiltonian $H_{\text{DI}}^{\text{real}}$, serve this purpose. To accommodate pairings, we Nambu double the theory and introduce the only local or on-site or momentum independent pairing with amplitude Δ . The

effective single particle Hamiltonian then reads

$$H_{\text{pair}}^{\text{real}} = \sum_{j \neq k} \frac{F(r_{jk})}{2} c_j^\dagger \left[-it(C_{jk}\Gamma_{01} + S_{jk}\Gamma_{02}) + 2t_0\Gamma_{03} \right] c_k + \sum_j c_j^\dagger \left[(M - 4t_0)\Gamma_{03} + \Delta\Gamma_{13} - \mu\Gamma_{30} \right] c_j, \quad (4)$$

where $\Gamma_{\rho\nu} = \eta_\rho\tau_\nu$, with $\rho, \nu = 0, \dots, 3$. The newly introduced Pauli matrices $\{\eta_\rho\}$ operate on the Nambu or particle-hole index. The chemical potential μ is measured from the zero energy. The Nambu doubled spinor is $c_j = [c_{\alpha j}, c_{\beta j}, c_{\beta j}^\dagger, c_{\alpha j}^\dagger]$. We implement the above Hamiltonian on a square lattice, as well as on amorphous network, Penrose and Ammann-Beenker quasicrystals, and Sierpiński carpet and glued Sierpiński triangle fractal lattices, paradigmatic members of the NCDM family.

Bott index. NCTSCs on 2D NCDMs can be identified from the Bott index (B). To proceed, we place all the sites of NCDMs within a unit square, and denote their coordinates by $x_i \in [0, 1]$ and $y_i \in [0, 1]$. Then in terms of two diagonal matrices $X_{i,j} = x_i\delta_{i,j}$ and $Y_{i,j} = y_i\delta_{i,j}$, we define two diagonal unitary matrices $U_x = \exp(2\pi i X)$ and $U_y = \exp(2\pi i Y)$. Finally, in terms of the projector (\mathcal{P}) onto the filled eigenstates of $H_{\text{pair}}^{\text{real}}$ up to the Fermi energy (μ), defined as $\mathcal{P} = \sum_{E < \mu} |E\rangle\langle E|$, we compute

$$B = \frac{1}{2\pi} \text{Im} [\text{Tr} [\ln [V_x V_y V_x^\dagger V_y^\dagger]]], \quad (5)$$

where $V_j = I - \mathcal{P} + \mathcal{P}U_j\mathcal{P}$ for $j = x$ and y [44]. Two instructive cuts of the global phase diagram of $H_{\text{pair}}^{\text{real}}$ in the (M, Δ) plane for $\mu = 0$ and (M, μ) plane for a fixed Δ , as shown in Fig. 1(a) and (b), respectively, unveil NCTSCs with $B = -1$ and -2 , besides the trivial paired state with $B = 0$. Also notice that as the chemical doping is increased, the NCTSCs occupy a larger portion of the phase diagram. Hence, increasing chemical doping is conducive for the nucleation of NCTSCs, which is thus promising to be observed in doped NCDMs.

The bulk-boundary correspondence for NCTSCs can be established by diagonalizing $H_{\text{pair}}^{\text{real}}$ on various noncrystalline lattices with periodic and open boundary conditions. Except on two fractal lattices, the energy spectra with these two geometries show clear distinction, especially near the zero energy. Namely, near zero energy modes can only be found for NCTSCs, characterized by $B = -1$ and -2 , with open geometry. See Fig. 2 (top row). The LDOS of these modes are highly localized near the outer edges of these systems. See Fig. 2 (bottom row). By contrast, on fractal lattices the energy spectra are qualitatively insensitive to the boundary conditions and near zero energy modes display strong localization close to the outer (inner) edges of these systems with open (periodic) geometry. See Fig. 3. Therefore, NCTSCs like their crystalline counterparts on a square lattice manifest the bulk-boundary correspondence.

IPR. To further pin the edge localization of near zero energy modes, we compute the IPR for generic wavefunction of $H_{\text{pair}}^{\text{real}}$ with energy E_j , given by $|E_j\rangle = \sum_i \psi_{i,j} |i\rangle$

$$P(j) = \sum_i |\psi_{i,j}|^4, \quad (6)$$

where the index i involves the site, orbital and Nambu indices. The results are shown in Fig. 4. Except on two fractal lattices, the IPR for the near zero energy states scales linearly with \sqrt{N} in a system containing N sites. This observation confirms the outer edge localization of these modes as the number of edge sites in all NCDMs scales as \sqrt{N} . By contrast, no such scaling of IPR of near zero energy modes can be observed on two fractal lattices due to the inner edges and the localization of near zero energy modes near it with periodic geometry [Fig. 3].

Summary and discussions. To summarize, here we show that a family of time-reversal symmetry breaking 2D NCDMs, encompassing amorphous network, quasicrystals and fractal lattices, can accommodate 2D NCTSCs that arise from the only allowed (by Pauli exclusion principle) local or on-site pairing. Consequently, such local NCTSCs are immune to the lack of translational symmetry or structural disorder in NCDMs. They possess a genuine bulk topological invariant, namely quantized Bott index, encoded by 1D topological edge modes of neutral Majorana fermions, manifesting the bulk-boundary correspondence. We also note that chemical doping boosts the propensity of electrons toward the formation of Cooper pairs in the topological channel.

Electronic designer materials [45–48], constitute the most promising platform where our proposed NCTSCs (proximity or phonon mediated) can be observed at low temperatures. On this class of systems, Penrose quasicrystal [46] and Sierpiński triangle fractal lattice [48] have been realized, where hopping elements can be tuned at least to a certain degree. Molecular quantum materials are also promising for NCTSCs, where Sierpiński triangle fractal lattices have been engineered [49].

The proposed mechanism is sufficiently general, and it can be extended to NCDMs belonging to arbitrary symmetry class in arbitrary dimensions to demonstrate NCTSCs therein. In particular, our conclusions should be applicable to recently realized three-dimensional topological insulator in amorphous Bi_2Se_3 [41], especially when it is doped. This specific material is promising as its crystalline counterpart, when doped, possibly harbors odd-parity topological paired states [50–57] that stems from a local or on-site pairing, the exact nature of which, however, still remains a subject of debate [58–62].

Acknowledgments. S.M. thanks Weizmann Institute of Science, Israel Deans fellowship through Feinberg Graduate School for financial support. S.K.D and B.R. were supported by the Startup grant of B.R. from Lehigh University. We are thankful to András L. Szabó for technical support.

-
- [1] M. Z. Hasan and C. L. Kane, *Colloquium: Topological insulators*, Rev. Mod. Phys. **82**, 3045 (2010).
- [2] X.-L. Qi and S. C. Zhang, Topological insulators and superconductors, Rev. Mod. Phys. **83**, 1057 (2011).
- [3] C.-K. Chiu, J. C. Y. Teo, A. P. Schnyder, and S. Ryu, Classification of topological quantum matter with symmetries, Rev. Mod. Phys. **88**, 035005 (2016).
- [4] C. L. Kane and E. J. Mele, Z_2 Topological Order and the Quantum Spin Hall Effect, Phys. Rev. Lett. **95**, 146802 (2005).
- [5] X.-L. Qi, Y.-S. Wu, and S.-C. Zhang, Topological quantization of the spin Hall effect in two-dimensional paramagnetic semiconductors, Phys. Rev. B **74**, 085308 (2006).
- [6] B. A. Bernevig, T. L. Hughes, and S.-C. Zhang, Quantum Spin Hall Effect and Topological Phase Transition in HgTe Quantum Wells, Science **314**, 1757 (2006).
- [7] M. König, S. Wiedmann, C. Brüne, A. Roth, H. Buhmann, L. W. Molenkamp, X.-L. Qi, and S.-C. Zhang, Quantum spin Hall insulator state in HgTe quantum wells, Science **318**, 766 (2007).
- [8] D. Hsieh, D. Qian, L. Wray, Y. Xia, Y. S. Hor, R. J. Cava, and M. Z. Hasan, A topological Dirac insulator in a quantum spin Hall phase, Nature **452**, 970 (2008).
- [9] L. Fu, Topological Crystalline Insulators, Phys. Rev. Lett. **106**, 106802 (2011).
- [10] R.-J. Slager, A. Mesaros, V. Juričić, and J. Zaanen, The space group classification of topological band-insulators, Nat. Phys. **9**, 98 (2013).
- [11] K. Shiozaki and M. Sato, Topology of crystalline insulators and superconductors, Phys. Rev. B **90**, 165114 (2014).
- [12] J. Kruthoff, J. de Boer, J. van Wezel, C. L. Kane, R.-J. Slager, Topological Classification of Crystalline Insulators through Band Structure Combinatorics, Phys. Rev. X **7**, 041069 (2017).
- [13] B. Bradlyn, L. Elcoro, J. Cano, M. G. Vergniory, Z. Wang, C. Felser, M. I. Aroyo, and B. A. Bernevig, Topological quantum chemistry, Nature **547**, 298 (2017).
- [14] F. Tang, H. C. Po, A. Vishwanath, and X. Wang, Comprehensive search for topological materials using symmetry indicators, Nature **566**, 486 (2019).
- [15] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, Quantized Hall Conductance in a Two-Dimensional Periodic Potential, Phys. Rev. Lett. **49**, 405 (1982).
- [16] F. D. M. Haldane, Model for a Quantum Hall Effect without Landau Levels: Condensed-Matter Realization of the “Parity Anomaly”, Phys. Rev. Lett. **61**, 2015 (1988).
- [17] A. Altland and M. R. Zirnbauer, Nonstandard symmetry classes in mesoscopic normal-superconducting hybrid structures, Phys. Rev. B **55**, 1142 (1997).
- [18] G. E. Volovik, *The Universe in a Helium Droplet* (Oxford University Press, New York, 2003).
- [19] A. Agarwala and V. B. Shenoy, Topological Insulators in Amorphous Systems, Topological Insulators in Amorphous Systems, Phys. Rev. Lett. **118**, 236402 (2017).
- [20] N. P. Mitchell, L. M. Nash, D. Hexner, A. M. Turner and W. T. M. Irvine, Amorphous topological insulators constructed from random point sets, Nat. Phys. **14**, 380 (2018).
- [21] Y.-B. Yang, T. Qin, D.-L. Deng, L.-M. Duan, and Y. Xu, Topological Amorphous Metals, Phys. Rev. Lett. **123**, 076401 (2019).
- [22] M. Costa, G. R. Schleder, M. B. Nardelli, C. Lewenkopf, and A. Fazzio, Toward Realistic Amorphous Topological Insulators, Nano Lett. **19**, 8941 (2019).
- [23] P. Zhou, G.-G. Liu, X. Ren, Y. Yang, H. Xue, L. Bi, L. Deng, Y. Chong and B. Zhang, Photonic amorphous topological insulator, Light Sci Appl **9**, 133 (2020).
- [24] A. Agarwala, V. Juričić and B. Roy, Higher-order topological insulators in amorphous solids, Phys. Rev. Research **2**, 012067 (2020).
- [25] P. Mukati, A. Agarwala, and S. Bhattacharjee, Topological and conventional phases of a three-dimensional electronic glass, Phys. Rev. B **101**, 035142 (2020).
- [26] H. Huang and F. Liu, Quantum Spin Hall Effect and Spin Bott Index in a Quasicrystal Lattice, Phys. Rev. Lett. **121**, 126401 (2018).
- [27] H. Huang and F. Liu, Theory of spin Bott index for quantum spin Hall states in nonperiodic systems, Phys. Rev. B **98**, 125130 (2018).
- [28] R. Chen, C.-Z. Chen, J.-H. Gao, B. Zhou, and D.-H. Xu, Higher-Order Topological Insulators in Quasicrystals, Phys. Rev. Lett. **124**, 036803 (2020).
- [29] C.-B. Hua, R. Chen, B. Zhou, and D.-H. Xu, Higher-order topological insulator in a dodecagonal quasicrystal, Phys. Rev. B **102**, 241102(R) (2020).
- [30] M. Brzezińska, A. M. Cook, and T. Neupert, Topology in the Sierpiński-Hofstadter problem, Phys. Rev. B **98**, 205116 (2018).
- [31] S. Pai and A. Prem, Topological states on fractal lattices, Phys. Rev. B **100**, 155135 (2019).
- [32] A. A. Iliasov, M. I. Katsnelson, and S. Yuan, Hall conductivity of a Sierpiński carpet, Phys. Rev. B **101**, 045413 (2020).
- [33] S. Manna, B. Pal, W. Wang, and A. E. B. Nielsen, Anyons and fractional quantum Hall effect in fractal dimensions, Phys. Rev. Research **2**, 023401 (2020).
- [34] M. Fremling, M. van Hooft, C. M. Smith, and L. Fritz, Existence of robust edge currents in Sierpiński fractals, Phys. Rev. Research **2**, 013044 (2020).
- [35] Z. Yang, E. Lustig, Y. Lumer, and M. Segev, Photonic Floquet topological insulators in a fractal lattice, Light: Science & Applications **9**, 128 (2020).
- [36] S. Manna, C. W. Duncan, C. A. Weidner, J. F. Sherson, and A. E. B. Nielsen, Anyon braiding on a fractal lattice with a local Hamiltonian, Phys. Rev. A **105**, L021302 (2022).
- [37] S. Manna, S. Nandy, and B. Roy, Higher-order topological phases on fractal lattices, Phys. Rev. B **105**, L201301 (2022).
- [38] S. Manna, B. Jaworowski, and A. E. B. Nielsen, Many-body localization on finite generation fractal lattices, arXiv:2111.13516
- [39] M. N. Ivaki, I. Sahlberg, K. Pöyhönen, T. Ojanen, Topological Random Fractals, arXiv:2112.08824
- [40] S. Manna and B. Roy, Inner Skin Effects on Non-Hermitian Topological Fractals, arXiv:2202.07658
- [41] P. Corbae, S. Ciocys, D. Varjas, E. Kennedy, S. Zeltmann, M. Molina-Ruiz, S. Griffin, C. Jozwiak, Z. Chen, L.-W. Wang, A. M. Minor, M. Scott, A. G. Grushin, A. Lanzara, and F. Hellman, arXiv:1910.13412

- [42] B. B. Mandelbrot, *The Fractal Geometry of Nature* (Times Book, 2nd ed., 1982).
- [43] I. C. Fulga, D. I. Pikulin, and T. A. Loring, Aperiodic Weak Topological Superconductors, *Phys. Rev. Lett.* **116**, 257002 (2016).
- [44] T. A. Loring and M. B. Hastings, Disordered Topological Insulators via C^* -Algebras, *Europhys. Lett.* **92**, 67004 (2010).
- [45] K. K. Gomes, W. Mar, W. Ko, F. Guinea, and H. C. Manoharan, Designer Dirac fermions and topological phases in molecular graphene, *Nature* **483**, 306 (2012).
- [46] L. C. Collins, T. G. Witte, R. Silverman, D. B. Green, and K. K. Gomes, Imaging quasiperiodic electronic states in a synthetic Penrose tiling, *Nat. Commun.* **8**, 15961 (2017).
- [47] S. N. Kempkes, M. R. Slot, J. J. van den Broeke, P. Capiod, W. A. Benalcazar, D. Vanmaekelbergh, D. Bercioux, I. Swart, and C. Morais Smith, Robust zero-energy modes in an electronic higher-order topological insulator, *Nat. Mater.* **18**, 1292 (2019).
- [48] S. N. Kempkes, M. R. Slot, S. E. Freeney, S. J. M. Zevenhuizen, D. Vanmaekelbergh, I. Swart, and C. Morais Smith, Design and characterization of electrons in a fractal geometry, *Nat. Phys.* **15**, 127 (2019).
- [49] J. Shang, Y. Wang, M. Chen, J. Dai, X. Zhou, J. Kuttner, G. Hilt, X. Shao, J. M. Gottfried, and K. Wu, Assembling molecular Sierpiński triangle fractals, *Nat. Chem.* **7**, 389 (2015).
- [50] A. L. Wray, S.-Y. Xu, Y. Xia, Y. S. Hor, D. Qian, A. V. Fedorov, H. Lin, A. Bansil, R. J. Cava, and M. Z. Hasan, Observation of topological order in a superconducting doped topological insulator, *Nat. Phys.* **6**, 855 (2010).
- [51] M. Kriener, K. Segawa, Z. Ren, S. Sasaki, and Y. Ando, Bulk Superconducting Phase with a Full Energy Gap in the Doped Topological Insulator $\text{Cu}_x\text{Bi}_2\text{Se}_3$, *Phys. Rev. Lett.* **106**, 127004 (2011).
- [52] S. Sasaki, Z. Ren, A. A. Taskin, K. Segawa, L. Fu, and Y. Ando, Odd-Parity Pairing and Topological Superconductivity in a Strongly Spin-Orbit Coupled Semiconductor, *Phys. Rev. Lett.* **109**, 217004 (2012).
- [53] T. Kirzhner, E. Lahoud, K. B. Chaska, Z. Salman, and A. Kanigel, Point-contact spectroscopy of $\text{Cu}_{0.2}\text{Bi}_2\text{Se}_3$ single crystals, *Phys. Rev. B* **86**, 064517 (2012).
- [54] K. Matano, M. Kriener, K. Segawa, Y. Ando, and G.-Q. Zheng, Spin-rotation symmetry breaking in the superconducting state of $\text{Cu}_x\text{Bi}_2\text{Se}_3$, *Nat. Phys.* **12**, 852 (2016).
- [55] S. Yonezawa, K. Tajiri, S. Nakata, Y. Nagai, Z. Wang, K. Segawa, Y. Ando, and Y. Maeno, Thermodynamic evidence for nematic superconductivity in $\text{Cu}_x\text{Bi}_2\text{Se}_3$, *Nat. Phys.* **13**, 123 (2017).
- [56] T. Asaba, B. J. Lawson, C. Tinsman, L. Chen, P. Corbae, G. Li, Y. Qiu, Y. S. Hor, L. Fu, and Lu Li, Rotational Symmetry Breaking in a Trigonal Superconductor Nb-doped Bi_2Se_3 , *Phys. Rev. X* **7**, 011009 (2017).
- [57] Y. Pan, A. M. Nikitin, G. K. Araizi, Y. K. Huang, Y. Matsushita, T. Naka, and A. de Visser, Rotational symmetry breaking in the topological superconductor $\text{Sr}_x\text{Bi}_2\text{Se}_3$ probed by upper-critical field experiments, *Sci. Rep.* **6**, 28632 (2016).
- [58] L. Fu and E. Berg, Odd-Parity Topological Superconductors: Theory and Application to $\text{Cu}_x\text{Bi}_2\text{Se}_3$, *Phys. Rev. Lett.* **105**, 097001 (2010).
- [59] L. Fu, Odd-parity topological superconductor with nematic order: Application to $\text{Cu}_x\text{Bi}_2\text{Se}_3$, *Phys. Rev. B* **90**, 100509(R) (2014).
- [60] P. Goswami and B. Roy, Axionic superconductivity in three-dimensional doped narrow-gap semiconductors, *Phys. Rev. B* **90**, 041301(R) (2014).
- [61] J. W. F. Venderbos, V. Kozii, and L. Fu, Odd-parity superconductors with two-component order parameters: Nematic and chiral, full gap, and Majorana node, *Phys. Rev. B* **94**, 180504(R) (2016).
- [62] B. Roy, Y. Alavirad, and J. D. Sau, Global Phase Diagram of a Three-Dimensional Dirty Topological Superconductor, *Phys. Rev. Lett.* **118**, 227002 (2017).