

A LOCAL DOUGLAS FORMULA FOR HIGHER ORDER WEIGHTED DIRICHLET-TYPE INTEGRALS

SOUMITRA GHARA, RAJEEV GUPTA, AND MD. RAMIZ REZA

ABSTRACT. We prove a local Douglas formula for higher order weighted Dirichlet-type integrals. With the help of this formula, we study the multiplier algebra of the associated higher order weighted Dirichlet-type spaces \mathcal{H}_μ , induced by an m -tuple $\mu = (\mu_1, \dots, \mu_m)$ of finite non-negative Borel measures on the unit circle. In particular, it is shown that any weighted Dirichlet-type space of order m , for $m \geq 3$, forms an algebra under pointwise product. We also prove that every non-zero closed M_z -invariant subspace of \mathcal{H}_μ , has codimension 1 property if $m \geq 3$ or μ_2 is finitely supported. As another application of local Douglas formula obtained in this article, it is shown that for any $m \geq 2$, weighted Dirichlet-type space of order m does not coincide with any de Branges-Rovnyak space $\mathcal{H}(b)$ with equivalence of norms.

1. INTRODUCTION

The symbols \mathbb{Z} , \mathbb{N} and $\mathbb{Z}_{\geq 0}$ will denote the set of integers, positive integers and non-negative integers respectively. We use the symbols \mathbb{T} and \mathbb{D} to denote the unit circle and the open unit disc in the complex plane \mathbb{C} respectively. The notation $\mathcal{M}_+(\mathbb{T})$ stands for the set of all finite non-negative Borel measures on \mathbb{T} . Let $\mathcal{O}(\mathbb{D})$ denote the space of all complex valued holomorphic functions on \mathbb{D} . For each $\alpha \in \mathbb{R}$, let \mathcal{D}_α denote the subspace of $\mathcal{O}(\mathbb{D})$ given by

$$\mathcal{D}_\alpha := \left\{ f = \sum_{k=0}^{\infty} a_k z^k \mid \|f\|_\alpha^2 := \sum_{k=0}^{\infty} (k+1)^\alpha |a_k|^2 < \infty \right\},$$

see [32] for a detailed study of the space \mathcal{D}_α . Many classical function spaces arise as \mathcal{D}_α spaces. For examples \mathcal{D}_{-1} coincides with the Bergman space, the space \mathcal{D}_0 coincides with the Hardy space H^2 and \mathcal{D}_1 coincides with the Dirichlet space \mathcal{D} . Recall that the Dirichlet space \mathcal{D} consists of all functions f in $\mathcal{O}(\mathbb{D})$ for which the Dirichlet integral $D(f)$, given by

$$D(f) := \int_{\mathbb{D}} |f'(z)|^2 dA(z),$$

is finite. Here $A(z)$ denotes the *normalized Lebesgue area measure* on \mathbb{D} . It turns out that \mathcal{D} is contained in the Hardy space H^2 . In his study of minimal surfaces, Douglas [13] derived and used the following formula

$$D(f) = \int_{\mathbb{T}} \int_{\mathbb{T}} \frac{|f^*(\zeta) - f^*(\lambda)|^2}{|\zeta - \lambda|^2} d\sigma(\zeta) d\sigma(\lambda), \quad f \in \mathcal{D},$$

where $\sigma(\zeta)$ denotes the *normalized arc length measure* on \mathbb{T} and $f^*(\lambda)$ denotes the non-tangential limit of f at $\lambda \in \mathbb{T}$ (which exists a.e. on \mathbb{T}). This and similar formulae turns out to be very useful in understanding the boundary behaviour of functions in \mathcal{D} , see [6]. Later, Richter and Sundberg (see [19, p. 356]) introduced the notion of the local Dirichlet-type integral $D_\lambda(f)$ of a function f in H^2 at a point $\lambda \in \mathbb{T}$ in the following manner:

$$D_\lambda(f) := \int_{\mathbb{T}} \frac{|f^*(\zeta) - f^*(\lambda)|^2}{|\zeta - \lambda|^2} d\sigma(\zeta).$$

The first author was supported by the Fields Post-doctoral Fellowship of the Fields Institute for Research in Mathematical Sciences, Canada and by a post-doctoral fellowship of Laval University, Canada. The second author was supported through the INSPIRE faculty grant (Ref. No. DST/INSPIRE/04/2017/002367).

They established various remarkable formulae for the local Dirichlet-type integral in [19, Sec. 3]. One of the motivations for studying the local Dirichlet-type integral was to understand the weighted Dirichlet-type integral $D_\mu(f)$ for a function $f \in \mathcal{O}(\mathbb{D})$ and a measure $\mu \in \mathcal{M}_+(\mathbb{T})$, a notion introduced and studied by Richter in order to describe all cyclic analytic 2-isometries, see [27, Theorem 5.1]. The weighted Dirichlet-type integral $D_\mu(f)$ turns out to be a key ingredient in describing the structure of the invariant subspaces of the Dirichlet shift, that is, the operator M_z of multiplication by the co-ordinate function z on the Dirichlet space \mathcal{D} , see [20, Theorem 3.2]. For any $m \in \mathbb{Z}_{\geq 0}$, the operator M_z on \mathcal{D}_m turns out to be an analytic $(m+1)$ isometry, see [1], [2], [3] for a detailed study of m -isometries. Thus if one desires to describe the structure of the invariant subspaces of M_z on \mathcal{D}_m , it is natural to look for model for analytic $(m+1)$ isometries, see [28]. Following [28], for a measure μ in $\mathcal{M}_+(\mathbb{T})$, $f \in \mathcal{O}(\mathbb{D})$ and for a positive integer n , we consider the *weighted Dirichlet-type integral of f of order n* by

$$D_{\mu,n}(f) := \frac{1}{n!(n-1)!} \int_{\mathbb{D}} |f^{(n)}(z)|^2 P_\mu(z) (1-|z|^2)^{n-1} dA(z).$$

Here $f^{(n)}(z)$ denotes the n -th order derivative of f at z , and $P_\mu(z)$ is the Poisson integral of the measure μ , that is,

$$P_\mu(z) := \int_{\mathbb{T}} \frac{1-|z|^2}{|z-\zeta|^2} d\mu(\zeta), \quad z \in \mathbb{D}.$$

For a function $f \in \mathcal{O}(\mathbb{D})$, it will be also useful for us to consider the *weighted Dirichlet-type integral of order 0* defined by

$$D_{\mu,0}(f) := \lim_{R \rightarrow 1^-} \int_{\mathbb{T}} |f(R\zeta)|^2 P_\mu(R\zeta) d\sigma(\zeta),$$

provided the limit exists. Note that if $\mu = \sigma$ then the integral $D_{\sigma,0}(f)$ coincides with the square of the Hardy norm of f . In case of $n = 1$, the integral $D_{\mu,1}(f)$ coincides with the weighted Dirichlet-type integral $D_\mu(f)$ for every $\mu \in \mathcal{M}_+(\mathbb{T})$. In particular if $\mu = \sigma$ then $D_{\sigma,1}(f)$ coincides with $D(f)$. If μ is δ_λ , the Dirac measure at the point $\lambda \in \mathbb{T}$, we adopt a simpler notation $D_{\lambda,n}(\cdot)$ in place of $D_{\delta_\lambda,n}(\cdot)$ and we refer it as a *local Dirichlet-type integral of order n at λ* . We would like to mention that a similar notion, namely, local Dirichlet-type integral of order n of a function at a point $\lambda \in \mathbb{T}$ is introduced and studied recently in [17, p. 3]. Nevertheless, we will see in Remark 5.4 that this notion differs from that of local Dirichlet-type integral of order n introduced in this article.

For a measure $\mu \in \mathcal{M}_+(\mathbb{T})/\{0\}$ and for each $n \in \mathbb{Z}_{\geq 0}$, we consider the semi-inner product space $\mathcal{H}_{\mu,n}$ given by

$$\mathcal{H}_{\mu,n} := \{f \in \mathcal{O}(\mathbb{D}) : D_{\mu,n}(f) < \infty\},$$

associated to the semi-norm $\sqrt{D_{\mu,n}(\cdot)}$. In case μ is δ_λ , $\lambda \in \mathbb{T}$, we will use a simpler notation $\mathcal{H}_{\lambda,n}$ in place of $\mathcal{H}_{\delta_\lambda,n}$ and we refer it as a *local Dirichlet space of order n at λ* . If $\mu = 0$, we set $\mathcal{H}_{\mu,n} = H^2$ for every $n \in \mathbb{Z}_{\geq 0}$. If $\mu = \sigma$, by a straightforward computation it follows that for a holomorphic function $f = \sum_{k=0}^{\infty} a_k z^k$ in $\mathcal{O}(\mathbb{D})$, we have

$$D_{\sigma,n}(f) = \sum_{k=n}^{\infty} \binom{k}{n} |a_k|^2, \quad n \in \mathbb{Z}_{\geq 0},$$

where $\binom{k}{n} := \frac{k!}{n!(k-n)!}$ for any $k \geq n$. It follows that for each $n \in \mathbb{Z}_{\geq 0}$, the space $\mathcal{H}_{\sigma,n}$ coincides as a set with the space \mathcal{D}_n . The reader is referred to [16] (see also [28]) for several properties of the spaces $\mathcal{H}_{\mu,n}$ for an arbitrary non-negative measure μ and a positive integer n .

Richter and Sundberg had shown that, for any $\lambda \in \mathbb{T}$, $D_{\lambda,1}(f) = D_\lambda(f)$, for every f in H^2 , see [19, Proposition 2.2]. In these notations, we note that

$$D_{\lambda,1}(f) = D_{\sigma,0}\left(\frac{f(z) - f^*(\lambda)}{z - \lambda}\right),$$

for every $f \in \mathcal{H}_{\lambda,1}$, see also [29, Proposition 1] for an alternative proof. This formula is often known as a *local Douglas formula*, see [14, Section 7.2]. In this article, we establish a local Douglas formula for higher order local Dirichlet-type integrals, namely, we prove the following theorem.

Theorem 1.1. *Let n be a positive integer, $\lambda \in \mathbb{T}$, and $f \in \mathcal{O}(\mathbb{D})$. Then $f \in \mathcal{H}_{\lambda,n}$ if and only if $f = \alpha + (z - \lambda)g$ for some g in $\mathcal{H}_{\sigma,n-1}$ and $\alpha \in \mathbb{C}$. Moreover, in this case, the following statements hold:*

- (i) $D_{\lambda,n}(f) = D_{\sigma,n-1}(g)$,
- (ii) $f(z) \rightarrow \alpha$ as $z \rightarrow \lambda$ in each oricyclic approach region $|z - \lambda| < \kappa(1 - |z|^2)^{\frac{1}{2}}$, $\kappa > 0$. In particular, $f^*(\lambda)$ exists and is equal to α .

Thus one can rewrite the local Douglas formula for higher order Dirichlet-type integrals in the following form:

$$D_{\lambda,n}(f) = D_{\sigma,n-1}\left(\frac{f(z) - f^*(\lambda)}{z - \lambda}\right), \quad f \in \mathcal{H}_{\lambda,n}, \quad n \in \mathbb{N}. \quad (1)$$

We explore various applications of this generalized local Douglas formula (1) in this article. For $0 < r < 1$, let f_r be the r -dilation of a function $f \in \mathcal{O}(\mathbb{D})$ given by $f_r(z) := f(rz)$, $z \in \mathbb{D}$. As a first application of the formula (1), we prove in Theorem 2.6 that for each $n \geq 2$ and $\mu \in \mathcal{M}_+(\mathbb{T})$,

$$D_{\mu,n}(f_r) \leq \frac{4^{n-1}(2-r)r^{2n}}{(1+r)^{2n-2}} D_{\mu,n}(f) \quad (2)$$

for every $f \in \mathcal{H}_{\mu,n}$ and $0 < r < 1$. Note that for any $n \in \mathbb{N}$ and $0 < r < 1$, we have $\frac{4^{n-1}(2-r)r^{2n}}{(1+r)^{2n-2}} \leq 1$. Thus the inequality (2) is an improvement of [16, Theorem 1.1], where only the contractivity of the map $f \mapsto f_r$ was shown. Analogous results are known in the case of $n = 1$, see [19], [29], [14], [15], [18] and references therein.

In the study of function spaces, it is an important problem to determine the associated multiplier algebra. Let $\mu \in \mathcal{M}_+(\mathbb{T})$ and n be a non-negative integer. The multiplier algebra of $\mathcal{H}_{\mu,n}$ is defined by

$$\text{Mult}(\mathcal{H}_{\mu,n}) := \{\varphi \in \mathcal{O}(\mathbb{D}) : \varphi f \in \mathcal{H}_{\mu,n} \text{ for every } f \in \mathcal{H}_{\mu,n}\}.$$

Several studies have been done in characterizing the multipliers of the Dirichlet space \mathcal{D} , and more generally for the multipliers of the weighted Dirichlet-type space $\mathcal{H}_{\mu,1}$ for any $\mu \in \mathcal{M}_+(\mathbb{T})$, see [31],[20],[9] and references therein. In Theorem 3.5, we characterize $\text{Mult}(\mathcal{H}_{\lambda,n})$ for every $\lambda \in \mathbb{T}$ and $n \in \mathbb{N}$. This characterization also helps us to study the $\text{Mult}(\mathcal{H}_{\mu,n})$ for any $\mu \in \mathcal{M}_+(\mathbb{T})$ and for every $n \in \mathbb{N}$. It is well known that the space \mathcal{D}_n forms an algebra under pointwise product for every $n \geq 2$ and consequently $\text{Mult}(\mathcal{D}_n) = \mathcal{D}_n$ for each $n \geq 2$, see [32, Remark, p. 239]. Even though \mathcal{D}_1 , the classical Dirichlet space, is not an algebra under pointwise product (see [14, Theorem 1.3.1]), the subspace of bounded functions in it turns out to be an algebra, see [14, Theorem 1.3.2]. More generally, it is well known that $\mathcal{H}_{\mu,1}$ is never an algebra under pointwise product for any $\mu \in \mathcal{M}_+(\mathbb{T})$, see [14, Sec. 7.1, Exercise 4], but $\mathcal{H}_{\mu,1} \cap H^\infty$ forms an algebra under pointwise product, see [20, p. 170], where H^∞ is the Banach algebra of bounded holomorphic functions on \mathbb{D} . We extend this result in the following theorem for higher order weighted Dirichlet-type spaces.

Theorem 1.2. *Let $\mu \in \mathcal{M}_+(\mathbb{T})$ and $n \in \mathbb{N}$. Then $\mathcal{H}_{\mu,n} \cap \text{Mult}(\mathcal{H}_{\sigma,n-1})$ is an algebra.*

As a consequence of this theorem, we find that $\mathcal{H}_{\mu,n}$ also forms an algebra under pointwise product and $\text{Mult}(\mathcal{H}_{\mu,n}) = \mathcal{H}_{\mu,n}$ for every $\mu \in \mathcal{M}_+(\mathbb{T})/\{0\}$ and for each $n \geq 3$, see Corollary 3.6. The answer to the question of whether $\mathcal{H}_{\mu,2}$ is an algebra or not, is of mixed nature. We have found that when $\mu = \delta_\lambda$ with $\lambda \in \mathbb{T}$, then $\mathcal{H}_{\lambda,2}$ is not an algebra under pointwise product. On the other hand we have $\mathcal{H}_{\sigma,2} = \mathcal{D}_2$ forms an algebra under pointwise product, see [32, Remark, p. 239].

Let $m \in \mathbb{N}$ and $\boldsymbol{\mu} = (\mu_1, \dots, \mu_m)$ be an m -tuple of non-negative measures in $\mathcal{M}_+(\mathbb{T})$. Let $\mathcal{H}_{\boldsymbol{\mu}}$ be the linear subspace of $\mathcal{O}(\mathbb{D})$ given by $\mathcal{H}_{\boldsymbol{\mu}} := \bigcap_{j=1}^m \mathcal{H}_{\mu_j, j}$. Note that for each $j = 1, \dots, m$, the space $\mathcal{H}_{\mu_j, n}$ is contained in the Hardy space H^2 for any $n \in \mathbb{N}$, see [16, Corollary 2.5]. For $f \in \mathcal{H}_{\boldsymbol{\mu}}$, we associate the norm $\|f\|_{\boldsymbol{\mu}}$ given by

$$\|f\|_{\boldsymbol{\mu}}^2 := \|f\|_{H^2}^2 + \sum_{j=1}^m D_{\mu_j, j}(f).$$

The space $\mathcal{H}_{\boldsymbol{\mu}}$ with respect to the norm $\|\cdot\|_{\boldsymbol{\mu}}$ turns out to be a reproducing kernel Hilbert space, see [16, p. 13]. In particular, it follows that $\mathcal{H}_{\mu, n}$ is a Hilbert space with respect to the norm $\|\cdot\|_{\mu, n}$, given by

$$\|f\|_{\mu, n}^2 = \|f\|_{H^2}^2 + D_{\mu, n}(f), \quad f \in \mathcal{H}_{\mu, n}.$$

The operator M_z of multiplication by the co-ordinate function on $\mathcal{H}_{\boldsymbol{\mu}}$ turns out to be a cyclic analytic $(m+1)$ -isometry, see [16, Theorem 4.1, Corollary 5.3]. Let $\text{Lat}(M_z, \mathcal{H}_{\boldsymbol{\mu}})$ denote the lattice of all closed M_z -invariant subspaces of $\mathcal{H}_{\boldsymbol{\mu}}$. A subspace \mathcal{W} in $\text{Lat}(M_z, \mathcal{H}_{\boldsymbol{\mu}})$ is said to have codimension k property if $\dim(\mathcal{W} \ominus (z - \lambda)\mathcal{W}) = k$, for each $\lambda \in \mathbb{D}$, see [25, Definition 2.12]. In order to have a Beurling-type description of $\text{Lat}(M_z, \mathcal{H}_{\boldsymbol{\mu}})$, it is expected that \mathcal{W} has codimension 1 property for every non-zero \mathcal{W} in $\text{Lat}(M_z, \mathcal{H}_{\boldsymbol{\mu}})$, see [25, Proposition 2.13]. In the case of $m = 1$, that is, when $\boldsymbol{\mu} = \mu_1$ and $\mu_1 \in \mathcal{M}_+(\mathbb{T})$, it is well known that any non-zero \mathcal{W} in $\text{Lat}(M_z, \mathcal{H}_{\boldsymbol{\mu}})$ has codimension 1 property, see [20, Theorem 3.2]. We find that for any $m \geq 3$ and any non-zero \mathcal{W} in $\text{Lat}(M_z, \mathcal{H}_{\boldsymbol{\mu}})$ has codimension 1 property. For the remaining case $m = 2$, that is, when $\boldsymbol{\mu} = (\mu_1, \mu_2)$, we show that the result remains true provided μ_2 is finitely atomic, see Theorem 4.5.

There is an intimate connection between the local Dirichlet spaces \mathcal{H}_{λ} and the de Branges-Rovnyak spaces $\mathcal{H}(b)$ (see [8] for basic theory of $\mathcal{H}(b)$ spaces). For any b in the unit ball of H^∞ , the de Branges-Rovnyak space $\mathcal{H}(b)$ is the reproducing kernel Hilbert space with the reproducing kernel $\frac{1-b(z)\bar{b}(w)}{1-z\bar{w}}$, $z, w \in \mathbb{D}$. In [29], Sarason showed that any local Dirichlet space \mathcal{H}_{λ} coincides with a de Branges-Rovnyak space $\mathcal{H}(b)$, with equality of norms, where b can be chosen to be $b(z) = \frac{(1-r)\bar{\lambda}z}{1-r\bar{\lambda}z}$, $z \in \mathbb{D}$, with $r = \frac{3-\sqrt{5}}{2}$. In [10, Theorem 3.1], Chevrot, Guillot and Ransford showed that, in fact, if any \mathcal{H}_{μ} space is equal to some de Branges-Rovnyak space $\mathcal{H}(b)$, with equality of norms, then μ has to be a positive multiple of a point mass measure for some $\lambda \in \mathbb{T}$ and further b is identified explicitly. Thereafter, in [12], the problem of characterizing μ and b such that $\mathcal{H}_{\mu} = \mathcal{H}(b)$ as a set was addressed by obtaining certain necessary and some sufficient conditions for $\mathcal{H}_{\mu} = \mathcal{H}(b)$. In this paper, we study the same question for the spaces $\mathcal{H}_{\boldsymbol{\mu}}$ and $\mathcal{H}(b)$, where $\boldsymbol{\mu} = (\mu_1, \dots, \mu_m)$, $m \geq 2$, is an arbitrary m -tuple of positive measures in $\mathcal{M}_+(\mathbb{T})$.

This article is organized as follows. Section 2 is devoted to prove Theorem 1.1. Section 3 deals with a few applications of generalized local Douglas Formula. In this section, we prove Theorem 1.2 and as a corollary it is observed that $\mathcal{H}_{\mu, n}$ is an algebra for any $\mu \in \mathcal{M}_+(\mathbb{T})/\{0\}$ and $n \geq 3$, see Corollary 3.6. In Section 4, we discuss about the codimension 1 property for any closed M_z -invariant subspaces of $\mathcal{H}_{\mu, n}$. In Section 5, it is shown that, for $n \geq 2$, any weighted Dirichlet-type space of order n does not coincide with de Branges-Rovnyak space $\mathcal{H}(b)$, with equivalence of norms, for any b in the unit ball of H^∞ .

2. A GENERALIZED LOCAL DOUGLAS FORMULA

In this section, we shall prove a local Douglas formula for local Dirichlet spaces of order n . For any $\lambda \in \mathbb{T}$ and $n \in \mathbb{N}$, let $\nu_{\lambda, n}$ denote the positive weighted area measure on \mathbb{D} given by

$$d\nu_{\lambda, n}(z) = \frac{1}{n!(n-1)!} P_{\delta_{\lambda}}(z) (1 - |z|^2)^{n-1} dA(z),$$

where $P_{\delta_\lambda}(z) = \frac{1-|z|^2}{|z-\lambda|^2}$, $z \in \mathbb{D}$, is the *Poisson integral* of the Dirac measure δ_λ . Let $A^2(d\nu_{\lambda,n})$ be the weighted Bergman space corresponding to the measure $\nu_{\lambda,n}$, that is,

$$A^2(d\nu_{\lambda,n}) := \{f \in \mathcal{O}(\mathbb{D}) : \|f\|^2 = \int_{\mathbb{D}} |f(z)|^2 d\nu_{\lambda,n}(z) < \infty\}.$$

Lemma 2.1. *For any $\lambda \in \mathbb{T}$ and $n \in \mathbb{N}$, $A^2(d\nu_{\lambda,n})$ is a reproducing kernel Hilbert space with the reproducing kernel*

$$(n+1)!(n-1)! \frac{(z-\lambda)(\bar{w}-\bar{\lambda})}{(1-z\bar{w})^{n+2}}, \quad z, w \in \mathbb{D}.$$

Proof. Let $A^2((1-|z|^2)^n dA(z))$ be the weighted Bergman space corresponding to the weighted area measure $(1-|z|^2)^n dA(z)$ on \mathbb{D} , that is,

$$A^2((1-|z|^2)^n dA(z)) := \{f \in \mathcal{O}(\mathbb{D}) : \|f\|^2 = \int_{\mathbb{D}} |f(z)|^2 (1-|z|^2)^n dA(z) < \infty\}.$$

It is evident that the operator $\sqrt{n!(n-1)!}M_{z-\lambda}$ is a unitary from $A^2((1-|z|^2)^n dA(z))$ onto $A^2(d\nu_{\lambda,n})$, where $M_{z-\lambda}$ denotes the operator of multiplication by the function $(z-\lambda)$. By a direct computation, it is verified that

$$\left\{ \sqrt{(n+1) \binom{n+k+1}{k}} z^k : k \in \mathbb{Z}_{\geq 0} \right\}$$

forms an orthonormal basis of $A^2((1-|z|^2)^n dA(z))$. Therefore, by [24, Theorem 2.4], $A^2(d\nu_{\lambda,n})$ has the reproducing kernel

$$n!(n-1)!(n+1)(z-\lambda)(\bar{w}-\bar{\lambda}) \sum_{k=0}^{\infty} \binom{n+k+1}{k} z^k \bar{w}^k = (n+1)!(n-1)! \frac{(z-\lambda)(\bar{w}-\bar{\lambda})}{(1-z\bar{w})^{n+2}}, \quad z, w \in \mathbb{D}.$$

This completes the proof. \square

Lemma 2.2. *Let $\lambda \in \mathbb{T}$, $n \in \mathbb{N}$, and $f \in \mathcal{H}_{\sigma,n-1}$. Then $((z-\lambda)f)^{(n)}(z) = 0$ for all $z \in \mathbb{D}$ if and only if $D_{\sigma,n-1}(f) = 0$.*

Proof. Suppose that $f \in \mathcal{H}_{\sigma,n-1}$ and $((z-\lambda)f)^{(n)}(z) = 0$ for all $z \in \mathbb{D}$. Then $(z-\lambda)f$ is a polynomial of degree at most $n-1$. Note that $\mathcal{H}_{\sigma,n-1}$ is contained in the Hardy space H^2 (see [16, Corollary 2.5], [32, Corollary 1, Theorem 7]). Therefore, in view of [14, Lemma 1.5.4], we find that $\lim_{|z| \rightarrow 1^-} (1-|z|^2)|f(z)|^2 = 0$. Hence it follows that $\lim_{r \rightarrow 1^-} (r\lambda - \lambda)f(r\lambda) = 0$. This gives us that the polynomial $(z-\lambda)f$ must have a zero at λ and consequently f is a polynomial of degree at most $n-2$. Therefore, $D_{\sigma,n-1}(f) = 0$. Conversely, assume that $D_{\sigma,n-1}(f) = 0$. This implies f is a polynomial of degree at most $n-2$, and therefore $((z-\lambda)f)^{(n)}(z) = 0$ for all $z \in \mathbb{D}$. \square

Let $j \in \mathbb{Z}_{\geq 0}$. Note that, in general $\sqrt{D_{\sigma,j}(\cdot)}$ represents a seminorm on $\mathcal{H}_{\sigma,j}$ (unless $j = 0$). Consider, the subspace $\mathcal{U}_{\sigma,j}$ of $\mathcal{H}_{\sigma,j}$ given by

$$\mathcal{U}_{\sigma,j} = \{f \in \mathcal{O}(\mathbb{D}) : f(z) = \sum_{k=j}^{\infty} a_k z^k, D_{\sigma,j}(f) < \infty\}.$$

It is straightforward to verify that $\sqrt{D_{\sigma,j}(\cdot)}$ represents a norm on $\mathcal{U}_{\sigma,j}$ which is induced by an inner product. Moreover $\mathcal{U}_{\sigma,j}$ is a reproducing kernel Hilbert space in which the monomials given by $\{\binom{k}{j}^{-\frac{1}{2}} z^k : k \geq j\}$ forms an orthonormal basis. The reproducing kernel $K_{\sigma,j}$ of $\mathcal{U}_{\sigma,j}$ is given by

$$K_{\sigma,j}(z, w) = \sum_{k=j}^{\infty} \binom{k}{j}^{-1} z^k \bar{w}^k, \quad z, w \in \mathbb{D}. \quad (3)$$

For any $\alpha > 0$ and $j \in \mathbb{Z}_{\geq 0}$, the Pochhammer symbol $(\alpha)_j$ is defined by

$$(\alpha)_j = \begin{cases} \alpha(\alpha+1) \cdots (\alpha+j-1), & j \geq 1 \\ 1, & j = 0. \end{cases}$$

Proposition 2.3. *Let n be a positive integer, $\lambda \in \mathbb{T}$, and $T : \mathcal{O}(\mathbb{D}) \rightarrow \mathcal{O}(\mathbb{D})$ be the linear map given by*

$$(Tf)(z) = ((z - \lambda)f)^{(n)}(z), \quad z \in \mathbb{D}, \quad f \in \mathcal{O}(\mathbb{D}).$$

Then T maps the Hilbert space $(\mathcal{U}_{\sigma, n-1}, \sqrt{D_{\sigma, n-1}}(\cdot))$ isometrically onto $A^2(d\nu_{\lambda, n})$.

Proof. Consider the space $\mathcal{W} := T(\mathcal{U}_{\sigma, n-1})$ endowed with the norm

$$\|T(f)\|_{\mathcal{W}} := \sqrt{D_{\sigma, n-1}(f)}, \quad f \in \mathcal{H}_{\sigma, n-1}.$$

By Lemma 2.2, $\|\cdot\|_{\mathcal{W}}$ is a well-defined norm on \mathcal{W} . Clearly T maps $\mathcal{U}_{\sigma, n-1}$ isometrically onto \mathcal{W} . This gives us that the space \mathcal{W} endowed with the norm $\|\cdot\|_{\mathcal{W}}$ is a Hilbert space. Now we show that \mathcal{W} is a reproducing kernel Hilbert space. This is equivalent to showing that for each $w \in \mathbb{D}$, the evaluation map $E_w : \mathcal{W} \rightarrow \mathbb{C}$, defined by $E_w(g) = g(w)$, is continuous. Let $w \in \mathbb{D}$, and $T(f_n)$ converges to $T(f)$ in \mathcal{W} , where $f_n, f \in \mathcal{U}_{\sigma, n-1}$. Clearly f_n converges to f in $\mathcal{U}_{\sigma, n-1}$. Since $\mathcal{U}_{\sigma, n-1}$ is a reproducing kernel Hilbert space, it follows that f_n converges to f uniformly on compact subsets of \mathbb{D} . Therefore $T(f_n)$ also converges to $T(f)$ uniformly on compact subsets of \mathbb{D} . This, in particular, shows that E_w is continuous on \mathcal{W} , proving that \mathcal{W} is a reproducing kernel Hilbert space.

The proposition will now be established by showing that the reproducing kernels of \mathcal{W} and $A^2(d\nu)$ coincide (see [24, Proposition 2.3]). As the operator T is unitary, it maps the orthonormal basis $\left\{\left(\binom{k}{n-1}\right)^{-\frac{1}{2}} z^k : k \geq n-1\right\}$ of $\mathcal{U}_{\sigma, n-1}$ to an orthonormal basis of \mathcal{W} . Thus the reproducing kernel of \mathcal{W} is given by

$$K_{\mathcal{W}}(z, w) = \frac{\partial^{2n}}{\partial z^n \partial \bar{w}^n} \left((z - \lambda)(\bar{w} - \bar{\lambda}) K_{\sigma, n-1}(z, w) \right), \quad z, w \in \mathbb{D}. \quad (4)$$

In view of (3), we see that the expression in (4) is equal to

$$\begin{aligned} & \frac{\partial^{2n}}{\partial z^n \partial \bar{w}^n} \left(\sum_{k=n-1}^{\infty} \binom{k}{n-1}^{-1} \left(z^{k+1} \bar{w}^{k+1} + z^k \bar{w}^k - \lambda z^k \bar{w}^{k+1} - \bar{\lambda} z^{k+1} \bar{w}^k \right) \right) \\ &= \sum_{k=n-1}^{\infty} \frac{1}{\binom{k}{n-1}} ((k+2-n)_n)^2 z^{k+1-n} \bar{w}^{k+1-n} + \sum_{k=n}^{\infty} \frac{1}{\binom{k}{n-1}} ((k+1-n)_n)^2 z^{k-n} \bar{w}^{k-n} \\ & \quad - \sum_{k=n}^{\infty} \frac{1}{\binom{k}{n-1}} (k+2-n)_n (k+1-n)_n \left(\lambda z^{k-n} \bar{w}^{k+1-n} + \bar{\lambda} z^{k+1-n} \bar{w}^{k-n} \right) \\ &= \sum_{k=0}^{\infty} ((k+1)_n)^2 \left(\binom{k+n-1}{n-1}^{-1} + \binom{k+n}{n-1}^{-1} \right) z^k \bar{w}^k \\ & \quad - \sum_{k=0}^{\infty} \binom{k+n}{n-1}^{-1} (k+2)_n (k+1)_n \left(\lambda z^k \bar{w}^{k+1} + \bar{\lambda} z^{k+1} \bar{w}^k \right) \\ &= (n-1)! \left(\sum_{k=0}^{\infty} (k+1)_n (2k+n+1) z^k \bar{w}^k - \sum_{k=0}^{\infty} (k+1)_{n+1} \left(\lambda z^k \bar{w}^{k+1} + \bar{\lambda} z^{k+1} \bar{w}^k \right) \right). \end{aligned}$$

Also, by a straightforward calculation, we see that

$$\begin{aligned}
& \frac{(z - \lambda)(\bar{w} - \bar{\lambda})}{(1 - z\bar{w})^{n+2}} \\
&= \sum_{k=0}^{\infty} \binom{n+k+1}{k} \left(z^{k+1}\bar{w}^{k+1} + z^k\bar{w}^k - \lambda z^k\bar{w}^{k+1} - \bar{\lambda} z^{k+1}\bar{w}^k \right) \\
&= 1 + \sum_{k=1}^{\infty} \left(\binom{n+k}{k-1} + \binom{n+k+1}{k} \right) z^k\bar{w}^k - \sum_{k=0}^{\infty} \binom{n+k+1}{k} \left(\lambda z^k\bar{w}^{k+1} + \bar{\lambda} z^{k+1}\bar{w}^k \right) \\
&= 1 + \frac{1}{(n+1)!} \left(\sum_{k=1}^{\infty} (k+1)_n (2k+n+1) z^k\bar{w}^k - \sum_{k=0}^{\infty} (k+1)_{n+1} \left(\lambda z^k\bar{w}^{k+1} + \bar{\lambda} z^{k+1}\bar{w}^k \right) \right) \\
&= \frac{1}{(n+1)!} \left(\sum_{k=0}^{\infty} (k+1)_n (2k+n+1) z^k\bar{w}^k - \sum_{k=0}^{\infty} (k+1)_{n+1} \left(\lambda z^k\bar{w}^{k+1} + \bar{\lambda} z^{k+1}\bar{w}^k \right) \right) \\
&= \frac{1}{(n+1)!(n-1)!} K_w(z, w).
\end{aligned}$$

Use of Lemma 2.1 now completes the proof. \square

For any holomorphic function f on the unit disc \mathbb{D} and a complex number λ in \mathbb{T} , let $f^*(\lambda)$ denote the boundary value of f at λ , defined by

$$f^*(\lambda) := \lim_{r \rightarrow 1^-} f(r\lambda),$$

whenever it exists. The following theorem (main theorem of this section) is deduced as a consequence of Lemma 2.3. This theorem, in particular, tells us that for any f in $\mathcal{H}_{\lambda,n}$, the boundary value of f at λ exists. As another consequence of Theorem 1.1 and [16, Proposition 2.6], we see that $\mathcal{H}_{\lambda,n}$ is contained in $\mathcal{H}_{\sigma,n-1}$.

Proof of Theorem 1.1: Assume that $f \in \mathcal{H}_{\lambda,n}$. Thus we have $f^{(n)} \in A^2(d\nu_{\lambda,n})$. Now, by Proposition 2.3, there exists a \tilde{g} in $\mathcal{U}_{\sigma,n-1}$ such that $D_{\lambda,n}(f) = D_{\sigma,n-1}(\tilde{g})$ and $((z - \lambda)\tilde{g})^{(n)} = f^{(n)}$. Therefore,

$$f(z) = (z - \lambda)\tilde{g}(z) + p(z), \quad z \in \mathbb{D},$$

for some polynomial p of degree at most $n - 1$. Thus, with $q(z) = \frac{p(z) - p(\lambda)}{z - \lambda}$, we have

$$f(z) = p(\lambda) + (z - \lambda)g(z), \quad z \in \mathbb{D}$$

where $g = \tilde{g} + q$. Clearly, $D_{\sigma,n-1}(g) = D_{\sigma,n-1}(\tilde{g}) = D_{\lambda,n}(f)$, which is finite. Hence $g \in \mathcal{H}_{\sigma,n-1}$. Conversely, assume that $f = \alpha + (z - \lambda)g$ for some g with $g \in \mathcal{H}_{\sigma,n-1}$. Then it follows from Lemma 2.3 that $f^{(n)} \in A^2(d\nu_{\lambda,n})$, and hence $f \in \mathcal{H}_{\lambda,n}$. This completes the proof of the first of the theorem. Note that, for $f \in \mathcal{H}_{\lambda,n}$, the equality $D_{\lambda,n}(f) = D_{\sigma,n-1}(g)$ is already established. To complete the proof, suppose that $f = \alpha + (z - \lambda)g$ for some complex number α and g in $\mathcal{H}_{\sigma,n-1}$. Since $\mathcal{H}_{\sigma,n-1}$ is contained in the Hardy space H^2 , by [14, Lemma 1.5.4] we have $\lim_{|z| \rightarrow 1} (1 - |z|^2)^{\frac{1}{2}} |g(z)| = 0$. If $|z - \lambda| < \kappa(1 - |z|^2)^{\frac{1}{2}}$, $\kappa > 0$, then

$$|f(z) - \alpha| < \kappa(1 - |z|^2)^{\frac{1}{2}} |g(z)|.$$

Hence it follows that $f(z) \rightarrow \alpha$ as $z \rightarrow \lambda$ in each oricyclic approach region $|z - \lambda| < \kappa(1 - |z|^2)^{\frac{1}{2}}$. This completes the proof of this Theorem.

Corollary 2.4. *Let n be a positive integer and μ be a finite positive Borel measure on the unit circle \mathbb{T} . Then for any f in $\mathcal{H}_{\mu,n}$, $f^*(\lambda)$ exists for μ -almost every $\lambda \in \mathbb{T}$, and the following holds:*

$$D_{\mu,n}(f) = \int_{\mathbb{T}} \int_{\mathbb{D}} \left| \left(\frac{f(z) - f^*(\lambda)}{z - \lambda} \right)^{(n-1)}(z) \right|^2 (1 - |z|^2)^{n-1} dA(z) d\mu(\lambda)$$

Proof. By Fubini's theorem,

$$D_{\mu,n}(f) = \frac{1}{n!(n-1)!} \int_{\mathbb{D}} |f^{(n)}(z)|^2 P_{\mu}(z)(1-|z|^2)^{n-1} dA(z) = \int_{\mathbb{T}} D_{\lambda,n}(f) d\mu(\lambda).$$

An application of Theorem 1.1 completes the proof of the corollary. \square

We finish this section with a few applications of Theorem 1.1. The first application shows that for any $n > 1$ the map $f \mapsto f_r$, $0 \leq r < 1$, is contractive on $\mathcal{H}_{\mu,n}$. Here for any $f \in \mathcal{O}(\mathbb{D})$, the r th dilation f_r of f is defined by $f_r(z) := f(rz)$, $z \in \mathbb{D}$. We point out that this result is already obtained in [29, Proposition 3] for $n = 1$, and [16, Theorem 1.1] for every $n \geq 1$. One of the consequences of this result is that $\lim_{r \rightarrow 1^-} D_{\mu,n}(f_r - f) = 0$ for any $f \in \mathcal{H}_{\mu,n}$. We start with the following lemma.

Lemma 2.5. *Let \mathcal{H} be a semi-inner product space of holomorphic functions on the unit disc \mathbb{D} with the semi-norm $\|\cdot\|$. Suppose that $zf \in \mathcal{H}$ whenever $f \in \mathcal{H}$. If $\|zf\| \geq \|f\|$ for all $f \in \mathcal{H}$ then for $\lambda \in \mathbb{T}$, $0 \leq r < 1$, we have*

$$\|(z - \lambda)f\| \leq \frac{2}{1+r} \|(z - r\lambda)f\|, \quad f \in \mathcal{H}.$$

Proof. Let $f \in \mathcal{H}$. Since $\|zf\| \geq \|f\|$, it follows that

$$|\operatorname{Re} \lambda \langle f, zf \rangle| \leq |\lambda \langle f, zf \rangle| \leq \|f\| \|zf\| \leq \|zf\|^2,$$

where for any complex number w , $\operatorname{Re} w$ denotes the real part of w . Hence

$$\begin{aligned} & \frac{4}{(1+r)^2} \|(z - r\lambda)f\|^2 - \|(z - \lambda)f\|^2 \\ &= \left(\frac{4}{(1+r)^2} - 1 \right) \|zf\|^2 - \left(1 - \frac{4r^2}{(1+r)^2} \right) \|f\|^2 + 2 \left(1 - \frac{4r}{(1+r)^2} \right) \operatorname{Re} \lambda \langle f, zf \rangle \\ &\geq \left(\left(\frac{4}{(1+r)^2} - 1 \right) - \left(1 - \frac{4r^2}{(1+r)^2} \right) - 2 \left(1 - \frac{4r}{(1+r)^2} \right) \right) \|zf\|^2 \\ &= 0. \end{aligned}$$

This completes the proof. \square

Theorem 2.6. *Let $n \geq 1$ be an integer, and μ be in $\mathcal{M}_+(\mathbb{T})$. Then for any f in $\mathcal{O}(\mathbb{D})$, we have*

$$D_{\mu,n}(f_r) \leq \frac{4^{n-1}(2-r)r^{2n}}{(1+r)^{2n-2}} D_{\mu,n}(f), \quad 0 \leq r < 1. \quad (5)$$

Proof. We will establish the proof by induction on n . The base case of $n = 1$ follows from [18, Corollary 1.5]. Assume that the statement holds for every $f \in \mathcal{O}(\mathbb{D})$, $0 \leq r < 1$, and $n = 1, \dots, k-1$ for $k \geq 2$. Let $f \in \mathcal{O}(\mathbb{D})$ and $0 \leq r < 1$. Since $D_{\mu,k}(f) = \int_{\mathbb{T}} D_{\lambda,k}(f) d\mu(\lambda)$, in order to establish (5) for $n = k$, it suffices to show that the (5) holds for $n = k$ and $\mu = \delta_{\lambda}$ for each $\lambda \in \mathbb{T}$. Fix a λ in \mathbb{T} and assume that $D_{\lambda,k}(f) < \infty$. Then by Theorem 1.1 we obtain that $f(z) = f^*(\lambda) + (z - \lambda)g(z)$ for some g in $\mathcal{H}_{\sigma,k-1}$ and $D_{\lambda,k}(f) = D_{\sigma,k-1}(g)$. By a straightforward computation, we see that

$$f_r(z) = f_r^*(\lambda) + (z - \lambda)h_r(z), \quad (6)$$

where

$$h_r(z) = r \frac{(z - \lambda)g(z) - (r\lambda - \lambda)g(r\lambda)}{z - r\lambda}.$$

Note that

$$h(z) = r \frac{(z - \lambda)(g(z) - g(r\lambda)) + (z - r\lambda)g(r\lambda)}{z - r\lambda}.$$

Using this together with Lemma 2.5 we get that

$$\begin{aligned}
D_{\sigma,k-1}(h) &= r^2 D_{\sigma,k-1} \left(\frac{(z-\lambda)(g(z) - g(r\lambda))}{z - r\lambda} \right) \\
&\leq \frac{4r^2}{(1+r)^2} D_{\sigma,k-1}(g(z) - g(r\lambda)) \\
&= \frac{4r^2}{(1+r)^2} D_{\sigma,k-1}(g).
\end{aligned} \tag{7}$$

By induction hypothesis, we have that

$$D_{\sigma,k-1}(h_r) \leq \frac{4^{k-2}(2-r)r^{2k-2}}{(1+r)^{2k-4}} D_{\sigma,k-1}(h). \tag{8}$$

Using Theorem 1.1, together with (6), (7) and (8) we get

$$\begin{aligned}
D_{\lambda,k}(f_r) &= D_{\sigma,k-1}(h_r) \\
&\leq \frac{4^{k-2}(2-r)r^{2k-2}}{(1+r)^{2k-4}} D_{\sigma,k-1}(h) \\
&\leq \frac{4^{k-1}(2-r)r^{2k}}{(1+r)^{2k-2}} D_{\sigma,k-1}(g) = \frac{4^{k-1}(2-r)r^{2k}}{(1+r)^{2k-2}} D_{\lambda,k}(f).
\end{aligned}$$

This completes the proof. \square

The following theorem which is a generalization of [21, Theorem 3.1] (see also [12, Lemma 4.2]) describes the weighted Dirichlet-type space $\mathcal{H}_{\mu,n}$ when μ is finitely supported.

Theorem 2.7. *Let n, k be two positive integers, and $\mu = \sum_{j=1}^k c_j \delta_{\lambda_j}$ for some $c_j > 0$ and $\lambda_j \in \mathbb{T}$, $j = 1, \dots, k$. Then $f \in \mathcal{H}_{\mu,n}$ if and only if there exists a polynomial p of degree at most $k-1$ and a function $g \in \mathcal{H}_{\sigma,n-1}$ such that*

$$f = p + g \prod_{j=1}^k (z - \lambda_j).$$

Proof. Note that $f \in \mathcal{H}_{\mu,n}$ if and only if $f \in \mathcal{H}_{\lambda_j,n}$ for each $j = 1, \dots, k$. Suppose that $f = p + g \prod_{j=1}^k (z - \lambda_j)$ for some polynomial p of degree at most $k-1$ and a function $g \in \mathcal{H}_{\sigma,n-1}$. Since z is a multiplier of $\mathcal{H}_{\sigma,n-1}$, it follows that for any $i = 1, \dots, k$, $g \prod_{j=1, j \neq i}^k (z - \lambda_j)$ is in $\mathcal{H}_{\sigma,n-1}$. Therefore, by Theorem 1.1, the function $g \prod_{j=1}^k (z - \lambda_j) \in \mathcal{H}_{\lambda_i,n}$ for each $i = 1, \dots, k$. Hence $f \in \mathcal{H}_{\mu,n}$.

Conversely, suppose that $f \in \mathcal{H}_{\mu,n}$ and assume that, without loss of generality, $\lambda_1, \dots, \lambda_k$ are distinct. Then $f \in \mathcal{H}_{\lambda_j,n}$, and applying Theorem 1.1 we see that $f^*(\lambda_j)$ exists for all $j = 1, \dots, k$. Let p be the polynomial of degree $k-1$ such that $p(\lambda_j) = f^*(\lambda_j)$, $j = 1, \dots, k$. Again, by Theorem 1.1, it follows that $\frac{f-p}{z-\lambda_j}$ belongs to $\mathcal{H}_{\sigma,n-1}$ for $j = 1, \dots, k$. Since $\lambda_1, \dots, \lambda_k$ are distinct, using partial fraction formula, we see that

$$\frac{f-p}{\prod_{j=1}^k (z - \lambda_j)} = \sum_{j=1}^k \alpha_j \frac{f-p}{z - \lambda_j} \tag{9}$$

for some $\alpha_1, \dots, \alpha_k \in \mathbb{C}$. As the right hand side in (9) belongs to $\mathcal{H}_{\sigma,n-1}$, this completes the proof. \square

3. MULTIPLIERS OF $\mathcal{H}_{\mu,n}$

In the study of any function space, it is an important problem to characterize the multipliers of the associated function space. Let μ be a non-zero measure in $\mathcal{M}_+(\mathbb{T})$ and n be a positive integer. The set of multipliers of $\mathcal{H}_{\mu,n}$, is denoted by $\text{Mult}(\mathcal{H}_{\mu,n})$ and is defined by

$$\text{Mult}(\mathcal{H}_{\mu,n}) = \{\varphi \in \mathcal{O}(\mathbb{D}) : \varphi f \in \mathcal{H}_{\mu,n} \text{ for every } f \in \mathcal{H}_{\mu,n}\}.$$

In this section, we deduce many properties of a multiplier of $\mathcal{H}_{\mu,n}$ and give some descriptions of the multiplier space $\text{Mult}(\mathcal{H}_{\mu,n})$ in several different ways. Since the co-ordinate function z is a multiplier of $\mathcal{H}_{\mu,n}$ (see [16, Proposition 2.6]), it follows that the set of all polynomials is contained in $\text{Mult}(\mathcal{H}_{\mu,n})$. Let $\mathcal{O}(\overline{\mathbb{D}})$ denote the set of functions which are holomorphic in some neighbourhood of the closed unit disc $\overline{\mathbb{D}}$. As an application of the following proposition we show that if $\varphi \in \mathcal{O}(\overline{\mathbb{D}})$ then $\varphi \in \text{Mult}(\mathcal{H}_{\mu,n})$. Before proving the proposition, we start with the following useful lemma.

Lemma 3.1. *Let $\mu \in \mathcal{M}_+(\mathbb{T}) \setminus \{0\}$ and $n \in \mathbb{N}$. Then for any $f \in \mathcal{H}_{\mu,n}$ and $j = 0, \dots, n$, the integral*

$$I_{\mu,j,n}(f) := \int_{\mathbb{D}} |f^{(j)}(z)|^2 P_{\mu}(z)(1 - |z|^2)^{n-1} dA(z)$$

is finite.

Proof. Let $A^2(d\rho)$ be the weighted Bergman space with the weighted area measure ρ on the unit disc given by $d\rho(z) = P_{\mu}(z)(1 - |z|^2)^{n-1} dA(z)$. Note that, a function f in $\mathcal{O}(\mathbb{D})$ satisfies $I_{\mu,j,n}(f) < \infty$ if and only if $f^{(j)} \in A^2(d\rho)$. Let $f \in \mathcal{H}_{\mu,n}$. By definition it follows that $f^{(n)} \in A^2(d\rho)$. Let $0 \leq k \leq n-1$. The proof will be established by showing that if $f^{(j)} \in A^2(d\rho)$ for all $j = k+1, \dots, n$, then $f^{(k)} \in A^2(d\rho)$. Note that, by the Leibniz's product rule for differentiation

$$(z^{n-k}f)^{(n)} = z^{n-k}f^{(n)} + c_0 z^{n-k-1}f^{(n-1)} + \dots + c_{n-k} z f^{(k+1)} + \binom{n}{n-k} (n-k)! f^{(k)} \quad (10)$$

for some constants c_0, \dots, c_{n-k} . Since z is a multiplier of $\mathcal{H}_{\mu,n}$ (see [16, Proposition 2.6]), we see that $(z^{n-k}f)^{(n)} \in A^2(d\rho)$. Since $f^{(j)} \in A^2(d\rho)$ for all $j = k+1, \dots, n$, it follows from (10) that $f^{(k)} \in A^2(d\rho)$. □

Proposition 3.2. *Let $\mu \in \mathcal{M}_+(\mathbb{T})$ and $n \in \mathbb{N}$. Suppose $\varphi \in \mathcal{O}(\mathbb{D})$ such that $\varphi^{(n)} \in H^{\infty}$. Then $\varphi \in \text{Mult}(\mathcal{H}_{\mu,n})$.*

Proof. Let $\varphi \in \mathcal{O}(\mathbb{D})$ satisfying $\varphi^{(n)} \in H^{\infty}$. By repeated applications of the Mean Value Theorem, it follows that $\varphi^{(j)} \in H^{\infty}$ for each $j = 0, 1, \dots, n$. By the Leibniz's product rule for differentiation and the Cauchy-Schwarz's inequality, note that, for any $f \in \mathcal{H}_{\mu,n}$,

$$|(\varphi f)^{(n)}(z)|^2 \leq \left(\sum_{j=0}^n |\varphi^{(n-j)}(z)|^2 |f^{(j)}(z)|^2 \right) \left(\sum_{j=0}^n \binom{n}{j}^2 \right), \quad z \in \mathbb{D}.$$

Let $C_n = \sum_{j=0}^n \binom{n}{j}^2$ and $\|g\|_{\infty} := \sup\{|g(z)| : z \in \mathbb{D}\}$ for any $g \in H^{\infty}$. It follows that for any $f \in \mathcal{H}_{\mu,n}$,

$$D_{\mu,n}(\varphi f) \leq \frac{C_n}{n!(n-1)!} \sum_{j=0}^n \|\varphi^{(n-j)}\|_{\infty}^2 \int_{\mathbb{D}} |f^{(j)}(z)|^2 P_{\mu}(z)(1 - |z|^2)^{n-1} dA(z).$$

Applying Lemma 3.1, we obtain that $D_{\mu,n}(\varphi f) < \infty$. Hence it follows that $\varphi \in \text{Mult}(\mathcal{H}_{\mu,n})$. □

Corollary 3.3. *Let $\mu \in \mathcal{M}_+(\mathbb{T})$ and $n \in \mathbb{N}$. Then $\mathcal{O}(\overline{\mathbb{D}}) \subset \text{Mult}(\mathcal{H}_{\mu,n})$.*

Proof. Note that for any $\varphi \in \mathcal{O}(\overline{\mathbb{D}})$, we have $\varphi^{(j)} \in H^{\infty}$ for every $j \in \mathbb{N}$. An application of Proposition 3.2 now completes the proof. □

Let $\boldsymbol{\mu} = (\mu_1, \dots, \mu_m)$, where $\mu_j \in \mathcal{M}_+(\mathbb{T})$ for every $j = 1, \dots, m$. Recall that the linear space $\mathcal{H}_{\boldsymbol{\mu}} = \cap_{j=1}^m \mathcal{H}_{\mu_j, j}$ is a Hilbert space with respect to the norm $\|\cdot\|_{\boldsymbol{\mu}}$, given by

$$\|f\|_{\boldsymbol{\mu}}^2 = \|f\|_{H^2}^2 + \sum_{j=1}^m D_{\mu_j, j}(f), \quad f \in \mathcal{H}_{\boldsymbol{\mu}}.$$

Furthermore, the evaluation functional ev_z , defined by $ev_z(f) := f(z)$, $f \in \mathcal{H}_{\boldsymbol{\mu}}$, is bounded for every $z \in \mathbb{D}$. Thus $\mathcal{H}_{\boldsymbol{\mu}}$ is a reproducing kernel Hilbert space of holomorphic functions on the unit disc, see [5], [24] for definition and other basic properties of a reproducing kernel Hilbert space. In particular, it follows that for any finite positive Borel measure μ on \mathbb{T} and a positive integer n , the linear space $\mathcal{H}_{\mu, n}$ is a reproducing kernel Hilbert space with respect to the norm $\|\cdot\|_{\mu, n}$, given by

$$\|f\|_{\mu, n}^2 = \|f\|_{H^2}^2 + D_{\mu, n}(f), \quad f \in \mathcal{H}_{\mu, n}.$$

Many properties of $\text{Mult}(\mathcal{H}_{\mu, n})$ can be derived using the reproducing kernel Hilbert space structure of $\mathcal{H}_{\mu, n}$. From now onwards we will often use the notation $(\mathcal{H}_{\mu, n}, \|\cdot\|_{\mu, n})$ to denote the Hilbert space $\mathcal{H}_{\mu, n}$ equipped with the norm $\|\cdot\|_{\mu, n}$.

In the following part of this section, we consider a general reproducing kernel Hilbert space having some special properties and derive few properties of the associated multiplier space.

Let \mathcal{H}_{κ} be a reproducing kernel Hilbert space of \mathbb{C} -valued holomorphic functions defined on the unit disc \mathbb{D} and let $\kappa : \mathbb{D} \times \mathbb{D} \rightarrow \mathbb{C}$ be the sesqui-analytic reproducing kernel for \mathcal{H}_{κ} , that is, $\kappa(\cdot, w) \in \mathcal{H}_{\kappa}$ and

$$\langle f, \kappa(\cdot, w) \rangle_{\mathcal{H}_{\kappa}} = f(w), \quad f \in \mathcal{H}_{\kappa}, \quad w \in \mathbb{D}.$$

In the remaining part of this section, we shall assume the following properties on κ :

- (A1) κ is normalized at the origin, that is, $\kappa(w, 0) = 1$ for every $w \in \mathbb{D}$,
- (A2) z is a multiplier for \mathcal{H}_{κ} , that is, $zf \in \mathcal{H}_{\kappa}$ for every $f \in \mathcal{H}_{\kappa}$,
- (A3) $\mathbb{C}[z]$, the set of all polynomials, is dense in \mathcal{H}_{κ} .

We refer to the reproducing kernel Hilbert space \mathcal{H}_{κ} satisfying (A1)-(A3) as the *functional Hilbert space with property (A)*.

For example take $\boldsymbol{\mu} = (\mu_1, \dots, \mu_m)$, where $\mu_j \in \mathcal{M}_+(\mathbb{T})$ for every $j = 1, \dots, m$ and consider the associated reproducing kernel Hilbert space $\mathcal{H}_{\boldsymbol{\mu}}$. Since $\langle f, 1 \rangle = f(0)$ for every $f \in \mathcal{H}_{\boldsymbol{\mu}}$, the kernel function associated to $(\mathcal{H}_{\boldsymbol{\mu}}, \|\cdot\|_{\boldsymbol{\mu}})$ is normalized at the origin. This together with [16, Theorem 4.1, Proposition 4.2] gives us that $(\mathcal{H}_{\boldsymbol{\mu}}, \|\cdot\|_{\boldsymbol{\mu}})$ is a functional Hilbert space with property (A). In particular, it follows that, for any finite positive Borel measure μ on \mathbb{T} and for any $n \in \mathbb{N}$, the Hilbert space $(\mathcal{H}_{\mu, n}, \|\cdot\|_{\mu, n})$ is also a functional Hilbert space with property (A).

For a reproducing kernel Hilbert space \mathcal{H}_{κ} , which consists of holomorphic functions on the unit disc \mathbb{D} , the multipliers space of \mathcal{H}_{κ} , is defined by

$$\text{Mult}(\mathcal{H}_{\kappa}) := \{\varphi \in \mathcal{H}_{\kappa} : \varphi f \in \mathcal{H}_{\kappa} \text{ for every } f \in \mathcal{H}_{\kappa}\}.$$

Note that $\text{Mult}(\mathcal{H}_{\kappa})$ forms an algebra with respect to the pointwise product, that is, $\varphi\psi \in \text{Mult}(\mathcal{H}_{\kappa})$ for every $\varphi, \psi \in \text{Mult}(\mathcal{H}_{\kappa})$. Moreover one can identify $\text{Mult}(\mathcal{H}_{\kappa})$ as a closed subalgebra of $\mathcal{B}(\mathcal{H}_{\kappa})$ in the following manner. Given $\varphi \in \text{Mult}(\mathcal{H}_{\kappa})$, we write M_{φ} for the multiplication operator acting on \mathcal{H}_{κ} given by

$$M_{\varphi}(f) = \varphi f, \quad f \in \mathcal{H}_{\kappa}.$$

By an application of the Closed Graph Theorem, it follows that for any $\varphi \in \text{Mult}(\mathcal{H}_{\kappa})$, the operator M_{φ} is bounded from \mathcal{H}_{κ} into itself. Consider the norm $\|\cdot\|_{op}$ on $\text{Mult}(\mathcal{H}_{\kappa})$ defined by

$$\|\varphi\|_{op} := \|M_{\varphi}\| = \sup\{\|\varphi f\|_{\mathcal{H}_{\kappa}} : \|f\|_{\mathcal{H}_{\kappa}} = 1\}, \quad \varphi \in \text{Mult}(\mathcal{H}_{\kappa}).$$

It follows that $(\text{Mult}(\mathcal{H}_\kappa), \|\cdot\|_{op})$ is a commutative Banach algebra with identity. Let \mathcal{H}_κ be a functional Hilbert space with property (A). It follows that $\text{Mult}(\mathcal{H}_\kappa)$ can be identified as the commutant of the operator M_z of multiplication by the co-ordinate function on the Hilbert space \mathcal{H}_κ . Indeed if $T \in \mathcal{B}(\mathcal{H}_\kappa)$ satisfying $TM_z = M_zT$, then a routine argument using the polynomial density of the functional Hilbert space \mathcal{H}_κ , it can be shown that T is given by $T = M_\varphi$ for some $\varphi \in \text{Mult}(\mathcal{H}_\kappa)$ where $\varphi = T(1)$.

Let \mathcal{H}_κ be a functional Hilbert space with property (A). By assumption (A1), we have $\langle \kappa(\cdot, 0), \kappa(\cdot, w) \rangle = \kappa(w, 0) = 1$ for each $w \in \mathbb{D}$. Therefore $\kappa(\cdot, w) \neq 0$ for each $w \in \mathbb{D}$. Hence it follows that the multiplier algebra $\text{Mult}(\mathcal{H}_\kappa)$ is a subalgebra of H^∞ , the algebra of bounded analytic functions on the unit disc, for every functional Hilbert space \mathcal{H}_κ with property (A), see [23, Problem 68], [24, Corollary 5.22], [30, Proposition 3]. Thus, we obtain that $\text{Mult}(\mathcal{H}_{\mu,n}) \subseteq H^\infty$ for every measure μ in $\mathcal{M}_+(\mathbb{T})$ and $n \in \mathbb{N}$.

Among the measures in $\mathcal{M}_+(\mathbb{T})$, there are two special measures. First one is the Lebesgue measure σ and the second one is the Dirac delta measure δ_λ for $\lambda \in \mathbb{T}$. The space $\mathcal{H}_{\sigma,n}$ coincides with the weighted Dirichlet-type space \mathcal{D}_n for each $n \in \mathbb{Z}_{\geq 0}$. The set of multipliers of \mathcal{D}_n is well studied, see for example [32, 31] and references therein. It is known that, for every $n \geq 2$, the space \mathcal{D}_n forms an algebra, that is, $fg \in \mathcal{D}_n$, for every $f, g \in \mathcal{D}_n$, see [32, Theorem 7]. Consequently it follows that $\text{Mult}(\mathcal{H}_{\sigma,n}) = \mathcal{H}_{\sigma,n}$ for every $n \geq 2$. Now we concentrate our study to characterize the multiplier of $\mathcal{H}_{\lambda,n}$ for any positive integer n . The following lemma will be useful in describing the set $\text{Mult}(\mathcal{H}_{\lambda,n})$. This lemma can be thought as a generalization of [19, Lemma 5.3].

Lemma 3.4. *Let n be a positive integer, $\lambda \in \mathbb{T}$ and $f \in \mathcal{H}_{\lambda,n}$ such that $D_{\lambda,n}(f) \neq 0$. Suppose $\varphi \in \text{Mult}(\mathcal{H}_{\sigma,n-1}) \cap \mathcal{H}_{\lambda,n}$, then*

$$\begin{aligned} D_{\lambda,n}(\varphi f) &\leq 2\|\varphi\|_{s(\sigma,n-1)}^2 D_{\lambda,n}(f) + 2|f^*(\lambda)|^2 D_{\lambda,n}(\varphi) \quad \text{and} \\ |f^*(\lambda)|^2 D_{\lambda,n}(\varphi) &\leq 2\|\varphi\|_{s(\sigma,n-1)}^2 D_{\lambda,n}(f) + 2D_{\lambda,n}(\varphi f), \end{aligned}$$

where

$$\|\varphi\|_{s(\sigma,n-1)} := \sup \left\{ \sqrt{D_{\sigma,n-1}(\varphi h)} : D_{\sigma,n-1}(h) = 1 \right\}.$$

Proof. For a function $g \in H^2$, if the non-tangential limit $g^*(\lambda)$ of g at λ exists, then by the generalized local Douglas formula (see Theorem 1.1), we have

$$D_{\lambda,n}(g) = D_{\sigma,n-1} \left(\frac{g(z) - g(\lambda)}{z - \lambda} \right).$$

Since $f, \varphi \in \mathcal{H}_{\lambda,n}$, the non-tangential limits $f^*(\lambda)$ and $\varphi^*(\lambda)$, of f and φ at λ respectively, exist, see Theorem 1.1. It follows that

$$D_{\lambda,n}(\varphi f) = D_{\sigma,n-1} \left(\frac{\varphi f(z) - (\varphi f)^*(\lambda)}{z - \lambda} \right).$$

Using the identity

$$\frac{\varphi f(z) - \varphi f(\lambda)}{z - \lambda} = \varphi(z) \frac{f(z) - f^*(\lambda)}{z - \lambda} + f^*(\lambda) \frac{\varphi(z) - \varphi(\lambda)}{z - \lambda}, \quad z \in \mathbb{D}, \quad (11)$$

it follows that

$$\begin{aligned} D_{\lambda,n}(\varphi f) &\leq 2D_{\sigma,n-1} \left(\varphi \frac{f(z) - f^*(\lambda)}{z - \lambda} \right) + 2|f^*(\lambda)|^2 D_{\sigma,n-1} \left(\frac{\varphi(z) - \varphi(\lambda)}{z - \lambda} \right), \\ &\leq 2\|\varphi\|_{s(\sigma,n-1)}^2 D_{\sigma,n-1} \left(\frac{f(z) - f^*(\lambda)}{z - \lambda} \right) + 2|f^*(\lambda)|^2 D_{\sigma,n-1} \left(\frac{\varphi(z) - \varphi^*(\lambda)}{z - \lambda} \right), \\ &\leq 2\|\varphi\|_{s(\sigma,n-1)}^2 D_{\lambda,n}(f) + 2|f^*(\lambda)|^2 D_{\lambda,n}(\varphi). \end{aligned}$$

Using the identity (11) one more time, we obtain that

$$\begin{aligned}
|f^*(\lambda)|^2 D_{\lambda,n}(\varphi) &= D_{\sigma,n-1} \left(f^*(\lambda) \frac{\varphi(z) - \varphi(\lambda)}{z - \lambda} \right), \\
&= D_{\sigma,n-1} \left(\frac{\varphi f(z) - \varphi f(\lambda)}{z - \lambda} - \varphi(z) \frac{f(z) - f^*(\lambda)}{z - \lambda} \right), \\
&\leq 2D_{\sigma,n-1} \left(\frac{\varphi f(z) - (\varphi f)^*(\lambda)}{z - \lambda} \right) + 2D_{\sigma,n-1} \left(\varphi(z) \frac{f(z) - f^*(\lambda)}{z - \lambda} \right), \\
&\leq 2D_{\lambda,n}(\varphi f) + 2\|\varphi\|_{s(\sigma,n-1)}^2 D_{\sigma,n-1} \left(\frac{f(z) - f^*(\lambda)}{z - \lambda} \right), \\
&= 2D_{\lambda,n}(\varphi f) + 2\|\varphi\|_{s(\sigma,n-1)}^2 D_{\lambda,n}(f).
\end{aligned}$$

This completes the proof. \square

The following proposition describes the multiplier algebra of $\mathcal{H}_{\lambda,n}$ for any $n \in \mathbb{N}$ and $\lambda \in \mathbb{T}$.

Theorem 3.5. *Let $n \in \mathbb{N}$ and $\lambda \in \mathbb{T}$. Then, $\text{Mult}(\mathcal{H}_{\lambda,n}) = \mathcal{H}_{\lambda,n} \cap \text{Mult}(\mathcal{H}_{\sigma,n-1})$.*

Proof. By Lemma 3.4, it follows that $\varphi \in \text{Mult}(\mathcal{H}_{\lambda,n})$ whenever $\varphi \in \mathcal{H}_{\lambda,n} \cap \text{Mult}(\mathcal{H}_{\sigma,n-1})$. For the converse, let $\varphi \in \text{Mult}(\mathcal{H}_{\lambda,n})$. Since the constant function $1 \in \mathcal{H}_{\lambda,n}$, it follows that $\varphi \in \mathcal{H}_{\lambda,n}$. Consider the map $L_\lambda : \mathcal{H}_{\lambda,n} \rightarrow \mathcal{H}_{\sigma,n-1}$ defined by

$$L_\lambda(f)(z) := \frac{f(z) - f^*(\lambda)}{z - \lambda}, \quad z \in \mathbb{D}.$$

By the generalized local Douglas formula in Theorem 1.1, we have L_λ is an isometry w.r.t the associated semi-norms, that is, $D_{\sigma,n-1}(L_\lambda f) = D_{\lambda,n}(f)$ for every $f \in \mathcal{H}_{\lambda,n}$. Furthermore it also follows that L_λ is onto. Let $f \in \mathcal{H}_{\lambda,n}$. An application of the identity (11) gives us

$$\begin{aligned}
D_{\sigma,n-1} \left(\varphi(z) \frac{f(z) - f^*(\lambda)}{z - \lambda} \right) &= D_{\sigma,n-1} \left(\frac{\varphi f(z) - (\varphi f)^*(\lambda)}{z - \lambda} - f^*(\lambda) \frac{\varphi(z) - \varphi^*(\lambda)}{z - \lambda} \right), \\
&\leq 2D_{\sigma,n-1} \left(\frac{\varphi f(z) - (\varphi f)^*(\lambda)}{z - \lambda} \right) + 2|f^*(\lambda)|^2 D_{\sigma,n-1} \left(\frac{\varphi(z) - \varphi^*(\lambda)}{z - \lambda} \right), \\
&= 2D_{\lambda,n}(\varphi f) + 2|f^*(\lambda)|^2 D_{\lambda,n}(\varphi),
\end{aligned}$$

Thus we have $D_{\sigma,n-1}(\varphi L_\lambda f) < \infty$ for every $f \in \mathcal{H}_{\lambda,n}$. Since L_λ is onto, we obtain that $D_{\sigma,n-1}(\varphi g) < \infty$ for every $g \in \mathcal{H}_{\sigma,n-1}$. Hence $\varphi \in \text{Mult}(\mathcal{H}_{\sigma,n-1})$. This completes the proof. \square

Proof of Theorem 1.2 : Let $\varphi, \psi \in \mathcal{H}_{\mu,n} \cap \text{Mult}(\mathcal{H}_{\sigma,n-1})$. Since $\text{Mult}(\mathcal{H}_{\sigma,n-1})$ is an algebra, it is sufficient to show that $\varphi\psi \in \mathcal{H}_{\mu,n}$. Note that $D_{\mu,n}(g) = \int_{\mathbb{T}} D_{\lambda,n}(g) d\mu(\lambda)$, for any $g \in \mathcal{H}_{\mu,n}$. Hence we obtain that $\varphi, \psi \in \mathcal{H}_{\lambda,n}$ for $[\mu]$ a.e. λ in \mathbb{T} . By Lemma 3.4 we have

$$\begin{aligned}
D_{\lambda,n}(\varphi\psi) &\leq 2\|\varphi\|_{s(\sigma,n-1)}^2 D_{\lambda,n}(\psi) + 2|\psi^*(\lambda)|^2 D_{\lambda,n}(\varphi), \\
&\leq 2\|\varphi\|_{s(\sigma,n-1)}^2 D_{\lambda,n}(\psi) + 2\|\psi\|_\infty^2 D_{\lambda,n}(\varphi), \quad \lambda \in \mathbb{T}/A.
\end{aligned}$$

Here in the last line we used the fact that $\psi \in \text{Mult}(\mathcal{H}_{\sigma,n-1}) \subseteq H^\infty$. Integrating both sides w.r.t $\mu(\lambda)$ we obtain

$$D_{\mu,n}(\varphi\psi) \leq 2\|\varphi\|_{s(\sigma,n-1)}^2 D_{\mu,n}(\psi) + 2\|\psi\|_\infty^2 D_{\mu,n}(\varphi), \quad \lambda \in \mathbb{T}.$$

This completes the proof of the theorem.

The above theorem helps us to characterize the multipliers of $\mathcal{H}_{\mu,n}$ for any $\mu \in \mathcal{M}_+(\mathbb{T})$ and for any positive integer $n \geq 3$. In this case, it turns out that $\mathcal{H}_{\mu,n}$ forms an algebra and consequently the multipliers of $\mathcal{H}_{\mu,n}$ is itself.

Corollary 3.6. *Let $n \geq 3$ be a positive integer and $\mu \in \mathcal{M}_+(\mathbb{T})$. Then $\mathcal{H}_{\mu,n}$ is an algebra and consequently $\text{Mult}(\mathcal{H}_{\mu,n}) = \mathcal{H}_{\mu,n}$.*

Proof. We have $\text{Mult}(\mathcal{H}_{\sigma,n-1}) = \mathcal{H}_{\sigma,n-1}$ for any $n \geq 3$. Thus applying Theorem 3.6, it follows that $\mathcal{H}_{\mu,n} \cap \mathcal{H}_{\sigma,n-1}$ is an algebra. By [16, Lemma 2.4], we also have that $\mathcal{H}_{\mu,n} \subseteq \mathcal{H}_{\sigma,n-1}$. Hence $\mathcal{H}_{\mu,n} \cap \mathcal{H}_{\sigma,n-1} = \mathcal{H}_{\mu,n}$. Thus $\mathcal{H}_{\mu,n}$ is an algebra and consequently $\text{Mult}(\mathcal{H}_{\mu,n}) = \mathcal{H}_{\mu,n}$. \square

Corollary 3.7. *Let $\mu = (\mu_1, \dots, \mu_m)$ be an m -tuple of finite positive Borel measures on \mathbb{T} with $\mu_m \neq 0$. Then $\mathcal{H}_\mu \cap \text{Mult}(\mathcal{H}_{\sigma,m-1})$ is an algebra. In particular, \mathcal{H}_μ is an algebra for $m \geq 3$.*

Proof. It is well known that for every $n \in \mathbb{N}$, $\text{Mult}(\mathcal{H}_{\sigma,n}) \subset \text{Mult}(\mathcal{H}_{\sigma,n-1})$, see [32, p. 233, Corollary 1]. So, we have

$$\begin{aligned} \mathcal{H}_\mu \cap \text{Mult}(\mathcal{H}_{\sigma,m-1}) &= \mathcal{H}_{\mu_1,1} \cap \dots \cap \mathcal{H}_{\mu_m,m} \cap \text{Mult}(\mathcal{H}_{\sigma,m-1}), \\ &= (\mathcal{H}_{\mu_1,1} \cap \text{Mult}(\mathcal{H}_{\sigma,0})) \cap \dots \cap (\mathcal{H}_{\mu_m,m} \cap \text{Mult}(\mathcal{H}_{\sigma,m-1})). \end{aligned}$$

By an application of Theorem 3.6, $\mathcal{H}_{\mu_j,j} \cap \text{Mult}(\mathcal{H}_{\sigma,j-1})$ is an algebra for each $j = 1, \dots, m$. Hence $\mathcal{H}_\mu \cap \text{Mult}(\mathcal{H}_{\sigma,m-1})$ is an algebra. It also follows from Theorem 3.6 that the space $\mathcal{H}_{\mu,j} \cap \text{Mult}(\mathcal{H}_{\sigma,j-1}) = \mathcal{H}_{\mu,j}$ for any $j \geq 3$. Hence $\mathcal{H}_\mu \cap \text{Mult}(\mathcal{H}_{\sigma,m-1}) = \mathcal{H}_\mu$ for any $m \geq 3$. This completes the proof. \square

Although $\mathcal{H}_{\mu,n}$ is an algebra for every $n \geq 3$ and for any $\mu \in \mathcal{M}_+(\mathbb{T})$, it is well known that $\mathcal{H}_{\mu,1}$ is never an algebra, see [14, Exercise 7.1.4]. So it is natural to ask the following question.

Question 3.8. Characterize all $\mu \in \mathcal{M}_+(\mathbb{T})$ such that $\mathcal{H}_{\mu,2}$ is an algebra?

Example 3.9. We now point out few cases when $\mathcal{H}_{\mu,2}$ is an algebra and when it is not.

- (i) The space $\mathcal{H}_{\lambda,2}$ is not an algebra for any $\lambda \in \mathbb{T}$. If possible assume that $\mathcal{H}_{\lambda,2}$ is an algebra for some $\lambda \in \mathbb{T}$. Then we must have $\mathcal{H}_{\lambda,2} = \text{Mult}(\mathcal{H}_{\lambda,2}) \subset H^\infty$. We now show that there exists a function in $\mathcal{H}_{\lambda,2}$ which is not bounded on \mathbb{D} . Choose a $\tilde{\lambda} \neq \lambda \in \mathbb{T}$ and a function $g \in \mathcal{H}_{\sigma,1}$, the classical Dirichlet space, such that $\lim_{r \rightarrow 1^-} g(r\tilde{\lambda}) = \infty$. Then by Theorem 1.1, $(z - \lambda)g \in \mathcal{H}_{\lambda,2}$. But $\lim_{r \rightarrow 1^-} (r\tilde{\lambda} - \lambda)g(r\tilde{\lambda}) = \infty$. Hence $(z - \lambda)g \notin H^\infty$.
- (ii) Note that $\mathcal{H}_{\sigma,2} = \mathcal{D}_2$ and \mathcal{D}_2 forms an algebra, see [32, Theorem 7]. In fact, if $\mu = (\mu_1, \mu_2 + \sigma)$, where $\mu_1, \mu_2 \in \mathcal{M}_+(\mathbb{T})$, then \mathcal{H}_μ is an algebra. To see this, note that $\mathcal{H}_\mu = \mathcal{H}_{\mu_1} \cap \mathcal{H}_{\sigma,2} \cap \mathcal{H}_{\mu,2}$. Since $\mathcal{H}_{\sigma,2} \subset \text{Mult}(\mathcal{H}_{\sigma,1})$, we obtain that $\mathcal{H}_\mu \subset \text{Mult}(\mathcal{H}_{\sigma,1})$. The conclusion now follows from Theorem 1.2.

The following theorem is a generalization of [14, Theorem 8.3.2]. As a corollary to this theorem we obtain a stronger version of Theorem 1.2 of this section.

Theorem 3.10. *Let $\mu \in \mathcal{M}_+(\mathbb{T}) \setminus \{0\}$, $n \in \mathbb{N}$. Let \mathcal{W} be a closed M_z -invariant subspace of $\mathcal{H}_{\mu,n}$ and $\phi \in \text{Mult}(\mathcal{H}_{\sigma,n-1})$. Suppose $f \in \mathcal{W}$ be such that $\phi f \in \mathcal{H}_{\mu,n}$, then $\phi f \in \mathcal{W}$.*

Proof. Let ϕ be a multiplier of $\mathcal{H}_{\sigma,n-1}$. For every $0 < r < 1$, define $\phi_r(z) := \phi(rz)$. Note that for every $0 < r < 1$, $\phi_r \in \mathcal{O}(\overline{\mathbb{D}})$ and hence $\phi_r \in \text{Mult}(\mathcal{H}_{\mu,n})$. Fix a r satisfying $0 < r < 1$. Choose a sequence of polynomials $\{p_j\}_{j \in \mathbb{N}}$ such that $D_{\mu,n}(p_j - \phi_r) \rightarrow 0$ as $j \rightarrow \infty$. Let $f \in \mathcal{W}$. Since \mathcal{W} is a M_z -invariant subspace of $\mathcal{H}_{\mu,n}$, we have that $p_j f \in \mathcal{W}$ for all $j \in \mathbb{N}$. As $p_n f \rightarrow \phi_r f$ in $\mathcal{H}_{\mu,n}$, we have that $\phi_r f \in M$. Since r is an arbitrary number between 0 and 1, we conclude that $\phi_r f \in M$ for every $0 < r < 1$. To show $\phi f \in M$, it is enough to show that

$$\sup\{D_{\mu,n}(\phi_r f) : 0 < r < 1\} < \infty.$$

Since if this is the case then $\{\phi_r f\}$ turns out to be a norm bounded family of function in the Hilbert space $\mathcal{H}_{\mu,n}$. And therefore there exists a sequence r_n converging to 1 such that $\phi_{r_n} f$ converges to ϕf weakly and hence $\phi f \in M$. Using Lemma 3.4, we find that

$$\begin{aligned} D_{\lambda,n}(\phi_r f) &\leq 2\|\phi_r\|_{s(\sigma,n-1)}^2 D_{\lambda,n}(f) + 2|f(\lambda)|^2 D_{\lambda,n}(\phi_r) \\ &\leq 6\|\phi\|_{s(\sigma,n-1)}^2 D_{\lambda,n}(f) + 4D_{\lambda,n}(\phi f) \end{aligned}$$

Integrating with respect to μ on both the sides of the above inequality, we obtain that

$$\sup\{D_{\mu,n}(\phi_r f) : 0 < r < 1\} < \infty.$$

This completes the proof. \square

Corollary 3.11. *Let $\mu \in \mathcal{M}_+(\mathbb{T}) \setminus \{0\}$ and $n \in \mathbb{N}$. If \mathcal{W} is a closed M_z -invariant subspace of $\mathcal{H}_{\mu,n}$ then $\mathcal{W} \cap \text{Mult}(\mathcal{H}_{\sigma,n-1})$ is an algebra.*

Proof. Let $\varphi, f \in \mathcal{W} \cap \text{Mult}(\mathcal{H}_{\sigma,n-1})$. From Theorem 1.2, we know that $\varphi f \in \mathcal{H}_{\mu,n}$. Now by Theorem 3.10, we deduce that $\varphi f \in \mathcal{W}$. This completes the proof. \square

We have already seen that $\mathcal{H}_{\mu,n}$ is an algebra for every $n \geq 3$ and for any $\mu \in \mathcal{M}_+(\mathbb{T})$. There are many interesting consequences of a reproducing kernel Hilbert space \mathcal{H}_κ being an algebra under pointwise product, see for example [30, Proposition 31, pp. 94–95]. In this part of the section, we note down some of these consequences of \mathcal{H}_κ being an algebra. Let $\mathcal{C}(\overline{\mathbb{D}})$ denote the algebra of continuous functions on the closed unit disc $\overline{\mathbb{D}}$. In the following proposition, we first derive that under certain conditions, an abelian Banach algebra contained in $\mathcal{O}(\mathbb{D})$ must lie in the disc algebra $\mathcal{O}(\mathbb{D}) \cap \mathcal{C}(\overline{\mathbb{D}})$.

Proposition 3.12. *Let $\mathcal{A} \subseteq \mathcal{O}(\mathbb{D})$ be an abelian Banach algebra under pointwise product and the set of all polynomials is dense in \mathcal{A} . Suppose $\sigma_{\mathcal{A}}(z)$, the spectrum of the co-ordinate function z in \mathcal{A} , is equal to the closed unit disc $\overline{\mathbb{D}}$. Then it follows that $\mathcal{A} \subseteq \mathcal{C}(\overline{\mathbb{D}})$ and for every $\lambda \in \overline{\mathbb{D}}$, the evaluation map $ev_\lambda : \mathcal{A} \rightarrow \mathbb{C}$ defined by $ev_\lambda(f) = f(\lambda)$, $f \in \mathcal{A}$, is continuous. Furthermore $\sigma_{\mathcal{A}}(f) = f(\overline{\mathbb{D}})$ holds for every $f \in \mathcal{A}$.*

Proof. Let Σ be the maximal ideal space of \mathcal{A} , that is, the collection of all non-zero homomorphisms of \mathcal{A} into \mathbb{C} . Consider Σ with the weak* topology that it inherits as a subset of \mathcal{A}^* . It is well known that Σ is a compact Hausdorff space, see [11, Ch. VII, Theorem 8.6]. Since $\mathcal{A} \subseteq \mathcal{O}(\mathbb{D})$, it follows that ev_λ belong to Σ for every $\lambda \in \mathbb{D}$. By our assumption the co-ordinate function z is a generator for the algebra \mathcal{A} . By [11, Ch. VII, Proposition 8.10], we have that there is a homeomorphism $\tau : \Sigma \rightarrow \overline{\mathbb{D}}$ such that $\tau(ev_\lambda) = \lambda$. Furthermore, it follows that the Gelfand transform $\gamma : \mathcal{A} \rightarrow \mathcal{C}(\overline{\mathbb{D}})$, satisfies the relation $\gamma(p) = p$ for every polynomial p in $\mathbb{C}[z]$. Now let $f \in \mathcal{A}$. For each $\lambda \in \mathbb{D}$, we have that

$$f(\lambda) = ev_\lambda(f) = (\tau^{-1}(\lambda))(f).$$

So for each $\lambda \in \mathbb{T}$, it is natural to define $f(\lambda) := (\tau^{-1}(\lambda))(f)$. Since τ is a homeomorphism, it follows that f extends as a function in $\mathcal{C}(\overline{\mathbb{D}})$. Note that $ev_\lambda = \tau^{-1}(\lambda) \in \Sigma$ for every $\lambda \in \overline{\mathbb{D}}$. Hence the continuity of ev_λ , for each $\lambda \in \overline{\mathbb{D}}$, follows from [11, Ch. VII, Proposition 8.4]. Finally, in view of [11, Ch. VII, Theorem 8.6], it follows that

$$\sigma_{\mathcal{A}}(f) = \{h(f) : h \in \Sigma\} = \{\tau^{-1}(\lambda)(f) : \lambda \in \overline{\mathbb{D}}\} = \{f(\lambda) : \lambda \in \overline{\mathbb{D}}\}.$$

This completes the proof. \square

As an application of the aforementioned proposition, we obtain that every functional Hilbert space \mathcal{H}_κ with property (A), which is also an algebra under pointwise product, must lie in the disc algebra. Furthermore, for such instances, one can describe all the cyclic vectors in \mathcal{H}_κ . Recall that a vector f in \mathcal{H}_κ is said to be cyclic if \mathcal{H}_κ is the smallest closed M_z -invariant subspace containing f . The following results can be thought as a straightforward generalization of the results in [30, Corollary 2, Corollary 3, p. 95].

Theorem 3.13. *Let \mathcal{H}_κ be a functional Hilbert space with property (A). Suppose \mathcal{H}_κ is an algebra under pointwise product and the spectrum $\sigma(M_z) = \overline{\mathbb{D}}$. Then $\mathcal{H}_\kappa \subseteq \mathcal{C}(\overline{\mathbb{D}})$ and every closed M_z -invariant subspace of \mathcal{H}_κ is an ideal. Furthermore, for a vector $f \in \mathcal{H}_\kappa$, the following are equivalent:*

- (a) f is cyclic.
- (b) f is invertible in $\text{Mult}(\mathcal{H}_\kappa)$.
- (c) f has no zero in $\overline{\mathbb{D}}$.

Proof. Note that \mathcal{H}_κ is an algebra under pointwise product if and only if $\mathcal{H}_\kappa = \text{Mult}(\mathcal{H}_\kappa)$. Since $\|1\|_{\mathcal{H}_\kappa} = \kappa(0,0) = 1$, it follows that $\|f\|_{\mathcal{H}_\kappa} \leq \|M_f\| = \|f\|_{op}$ for every $f \in \mathcal{H}_\kappa$. This shows that the inclusion map from $(\text{Mult}(\mathcal{H}_\kappa), \|\cdot\|_{op})$ onto $(\mathcal{H}_\kappa, \|\cdot\|_{\mathcal{H}_\kappa})$ is continuous. By an application of open mapping theorem it follows that the two norms $\|\cdot\|_{op}$ and $\|\cdot\|_{\mathcal{H}_\kappa}$ on \mathcal{H}_κ are equivalent. By assumption (A3), it now follows that the set of all polynomials is dense in $(\text{Mult}(\mathcal{H}_\kappa), \|\cdot\|_{op})$ as well. Since $\text{Mult}(\mathcal{H}_\kappa)$ coincides with the commutant of M_z in $\mathcal{B}(\mathcal{H}_\kappa)$, it follows that the spectrum of the co-ordinate function z in the multiplier algebra $\text{Mult}(\mathcal{H}_\kappa)$ is equal to the spectrum of M_z in $\mathcal{B}(\mathcal{H}_\kappa)$. Now applying Proposition 3.12 to the abelian Banach algebra $\mathcal{A} = (\text{Mult}(\mathcal{H}_\kappa), \|\cdot\|_{op})$, we obtain that $\mathcal{H}_\kappa = \text{Mult}(\mathcal{H}_\kappa) \subseteq \mathcal{C}(\overline{\mathbb{D}})$. For the proof of the second part, consider an element $f \in \mathcal{H}_\kappa$. Let $\mathcal{I}(f)$ denote the smallest closed M_z -invariant subspace of \mathcal{H}_κ , containing f , that is,

$$\mathcal{I}(f) = \bigvee \{z^n f : n \in \mathbb{Z}_{\geq 0}\}.$$

To show every closed M_z -invariant subspace is an ideal, it is sufficient to show that $\mathcal{I}(f)$ is an ideal. As the two norms $\|\cdot\|_{op}$ and $\|\cdot\|_{\mathcal{H}_\kappa}$ on \mathcal{H}_κ are equivalent and the set of all polynomials is dense in \mathcal{H}_κ , it follows that $\mathcal{I}(f)$ is an ideal. Since $\mathcal{I}(f)$ is a closed ideal in the Banach algebra $\mathcal{A} = (\text{Mult}(\mathcal{H}_\kappa), \|\cdot\|_{op})$, then either one of the following two mutually exclusive event must holds.

- (i) $\mathcal{I}(f) = \mathcal{H}_\kappa$,
- (ii) $\mathcal{I}(f)$ is contained in a maximal ideal, that is, $\mathcal{I}(f) \subseteq \ker(ev_\lambda)$ for some $\lambda \in \overline{\mathbb{D}}$. In particular $f(\lambda) = 0$ for some $\lambda \in \overline{\mathbb{D}}$.

Hence we obtain that an element $f \in \mathcal{H}_\kappa$, is cyclic, that is, $\mathcal{I}(f) = \mathcal{H}_\kappa$ if and only if f has no zeros in $\overline{\mathbb{D}}$. Applying Proposition 3.12, we also have that $\sigma_{\mathcal{A}}(f) = \{f(\lambda) : \lambda \in \overline{\mathbb{D}}\}$. Thus f has no zeros in $\overline{\mathbb{D}}$ if and only if f is invertible in $\text{Mult}(\mathcal{H}_\kappa)$. This completes the proof. \square

4. CODIMENSION k PROPERTY OF INVARIANT SUBSPACES

This section is an attempt to study the codimension k property of non-zero closed M_z -invariant subspaces of \mathcal{H}_μ for any $\mu = (\mu_1, \dots, \mu_m)$ with $\mu_j \in \mathcal{M}_+(\mathbb{T})$ for $j = 1, \dots, m$. Since $M_z|_{\mathcal{M}}$ is an analytic, $(m+1)$ -isometry for every non-zero \mathcal{M} in $\text{Lat}(M_z, \mathcal{H}_\mu)$, it turns out that the dimension of $\ker((M_z|_{\mathcal{M}})^* - \bar{\lambda})$, that is, the dimension of $\mathcal{M} \ominus (z - \lambda)\mathcal{M}$, is a positive integer independent of $\lambda \in \mathbb{D}$, see [16, Proposition 6.4]. Thus any \mathcal{M} in $\text{Lat}(M_z, \mathcal{H}_\mu)$ has codimension k property if and only if $\dim(\mathcal{M} \ominus z\mathcal{M}) = k$. In case of $m = 1$, that is, $\mu = \mu_1$ and $\mu_1 \in \mathcal{M}_+(\mathbb{T})$, it is well known that for any non-zero \mathcal{M} in $\text{Lat}(M_z, \mathcal{H}_\mu)$, \mathcal{M} has codimension 1 property, see [20, Theorem 3.2]. Here, firstly we show that for any $m \geq 3$ and any non-zero \mathcal{M} in $\text{Lat}(M_z, \mathcal{H}_\mu)$ has codimension 1 property.

Theorem 4.1. *Let $\mu = (\mu_1, \dots, \mu_m)$ be an m -tuple of finite non-negative Borel measure on \mathbb{T} . Suppose \mathcal{H}_μ is an algebra under pointwise product, that is, $\mathcal{H}_\mu = \text{Mult}(\mathcal{H}_\mu)$. Then for every non-zero \mathcal{M} in $\text{Lat}(M_z, \mathcal{H}_\mu)$, we have $\dim(\mathcal{M} \ominus z\mathcal{M}) = 1$. In particular if $m \geq 3$ and $\mu_m \neq 0$, then $\dim(\mathcal{M} \ominus z\mathcal{M}) = 1$ for any \mathcal{M} in $\text{Lat}(M_z, \mathcal{H}_\mu)$.*

Proof. Suppose \mathcal{H}_μ is an algebra under pointwise product, that is, $\mathcal{H}_\mu = \text{Mult}(\mathcal{H}_\mu)$. By an application of Closed Graph Theorem the multipliers norm $\|\cdot\|_{op}$ and the Hilbertian norm $\|\cdot\|_\mu$ on \mathcal{H}_μ are equivalent. In view of [16, Corollary 5.3], we obtain that the set of all polynomials is dense in \mathcal{H}_μ w.r.t the multiplier norm $\|\cdot\|_{op}$. For any $f \in \mathcal{H}_\mu$, let $[f]$ denotes the smallest closed M_z -invariant subspace containing f in \mathcal{H}_μ . Hence it follows that $\varphi[f] \subseteq [f]$ for every $\varphi, f \in \mathcal{H}_\mu$. Applying [7, Proposition 1] to \mathcal{H}_μ , we get that \mathcal{H}_μ is a cellular-indecomposable space. Now the first part of the theorem follows immediately from [7, Theorem 1]. The last part follows using the first part together with Corollary 3.7. \square

For the remaining case $m = 2$, that is, $\mu = (\mu_1, \mu_2)$ with $\mu_1, \mu_2 \in \mathcal{M}_+(\mathbb{T})$, we answer this question partially in Theorem 4.5. We start with a basic lemma which will be useful in the proof of Theorem 4.5.

Lemma 4.2. *Let H be a Hilbert space and V be a closed subspace of H with $\dim V^\perp = n$. Let W be another subspace of H , then $\dim(W \ominus (V \cap W)) \leq n$.*

Proof. Let P denote the orthogonal projection of H onto V^\perp . Suppose $X = W \ominus (V \cap W)$. We claim that $P|_X$ is an injective operator. To verify the claim, let $x \in X$ be such that $P|_X(x) = 0$. Since $\ker P = V$, $x \in V$. This implies $x = 0$. Thus claim stands verified and therefore $\dim(W \ominus (V \cap W)) \leq \dim V^\perp = n$. \square

Lemma 4.3. *Let μ be a finite positive Borel measure on the unit circle \mathbb{T} . Let $n \in \mathbb{N}$, $\lambda_1, \dots, \lambda_n \in \mathbb{T}$ be distinct points and c_1, \dots, c_n be positive real numbers. Let $\boldsymbol{\mu}$ be the tuple $(\mu, \sum_{j=1}^n c_j \delta_{\lambda_j})$. Then the subspace $\ker(M_z^{*2} M_z^2 - 2M_z^* M_z + I) = \mathcal{V}$ (say) is a closed M_z -invariant subspace of $\mathcal{H}_{\boldsymbol{\mu}}$ and the operator $M_z|_{\mathcal{V}}$ is a cyclic analytic 2-isometry. Moreover, $\dim \mathcal{V}^\perp = n$.*

Proof. From [16, Theorem 4.1], we know that M_z on $\mathcal{H}_{\boldsymbol{\mu}}$ is a cyclic analytic 3-isometry. Thus, by [1, Proposition 1.6], it follows that \mathcal{V} is a closed M_z -invariant subspace of $\mathcal{H}_{\boldsymbol{\mu}}$ and the operator $M_z|_{\mathcal{V}}$ is a 2-isometry. Since M_z on $\mathcal{H}_{\boldsymbol{\mu}}$ is analytic, $M_z|_{\mathcal{V}}$ is also analytic. In view of [26, Theorem 1], to show $M_z|_{\mathcal{V}}$ is cyclic, it is sufficient to show that $\dim(\mathcal{V} \ominus z\mathcal{V}) = 1$.

Before proving that $\dim(\mathcal{V} \ominus z\mathcal{V}) = 1$, first we show that $\dim \mathcal{V}^\perp = n$. In this regard, note that $\mathcal{H}_{\boldsymbol{\mu}} \subseteq \mathcal{H}_{\lambda_j, 2}$ for each $j = 1, \dots, n$. Thus, by Theorem 1.1, $f^*(\lambda_j)$ exists for all $f \in \mathcal{H}_{\boldsymbol{\mu}}$ and for every $j = 1, \dots, n$. Since $\mathcal{H}_{\boldsymbol{\mu}}$ is a reproducing kernel Hilbert space on \mathbb{D} , the linear functional $f \mapsto f(r\lambda_j)$ is bounded on $\mathcal{H}_{\boldsymbol{\mu}}$ for any $0 < r < 1$ and for each $j = 1, \dots, n$. An application of uniform boundedness principle gives that $f \mapsto f^*(\lambda_j)$ is a bounded linear functional on $\mathcal{H}_{\boldsymbol{\mu}}$ for each $j = 1, \dots, n$. Therefore, by Riesz representation theorem, there exist $h_1, \dots, h_n \in \mathcal{H}_{\boldsymbol{\mu}}$ such that

$$f^*(\lambda_j) = \langle f, h_j \rangle, \quad f \in \mathcal{H}_{\boldsymbol{\mu}}, \quad j = 1, \dots, n. \quad (12)$$

Since M_z on $\mathcal{H}_{\boldsymbol{\mu}}$ is a 3-isometry, by [1, Proposition 1.5], the operator $M_z^{*2} M_z^2 - 2M_z^* M_z + I$ is positive. Thus a function f of $\mathcal{H}_{\boldsymbol{\mu}}$ is in \mathcal{V} if and only if $\|z^2 f\|_{\boldsymbol{\mu}}^2 - 2\|zf\|_{\boldsymbol{\mu}}^2 + \|f\|_{\boldsymbol{\mu}}^2 = 0$. By [16, Proposition 2.7], this is equivalent to $D_{\tau, 0}(f) = 0$, where $\tau = \sum_{j=1}^n c_j \delta_{\lambda_j}$. Since $D_{\tau, 0}(f) = \sum_{j=1}^n c_j |f^*(\lambda_j)|^2$, we see that

$$\mathcal{V} = \{f \in \mathcal{H}_{\boldsymbol{\mu}} : f^*(\lambda_j) = 0, 1 \leq j \leq n\}.$$

Hence, by (12), it follows that $\mathcal{V}^\perp = \text{span}\{h_1, \dots, h_n\}$. Since λ_j 's are distinct points on \mathbb{T} , it follows that h_1, \dots, h_n are linearly independent. This shows that $\dim \mathcal{V}^\perp = n$. To finish the proof, note that M_z on $\mathcal{H}_{\boldsymbol{\mu}}$ is in the Cowen-Douglas class $B_1(\mathbb{D})$, see [16, Corollary 6.1]. Since \mathcal{V} is a closed M_z -invariant subspace of finite co-dimension, it follows from [22, p. 71] that $\dim(\mathcal{V} \ominus z\mathcal{V}) = 1$. This completes the proof. \square

Remark 4.4. By Remark 3.9(ii), the space $\mathcal{H}_{\boldsymbol{\mu}}$ is an algebra, where $\boldsymbol{\mu} = (\mu_1, \mu_2 + \sigma)$, $\mu_1, \mu_2 \in \mathcal{M}_+(\mathbb{T})$. Hence in view of Theorem 4.1 we obtain that for any $\mathcal{M} \in \text{Lat}(M_z, \mathcal{H}_{\boldsymbol{\mu}})$, $\dim(\mathcal{M} \ominus z\mathcal{M}) = 1$.

Theorem 4.5. *Let μ be a finite non-negative Borel measure on the unit circle \mathbb{T} . Let $n \in \mathbb{N}$, $\lambda_1, \dots, \lambda_n \in \mathbb{T}$ be distinct points and c_1, \dots, c_n be positive real numbers. Suppose \mathcal{M} is a non-zero subspace in $\text{Lat}(M_z, \mathcal{H}_{\boldsymbol{\mu}})$, where $\boldsymbol{\mu} = (\mu, \sum_{j=1}^n c_j \delta_{\lambda_j})$. Then $\dim(\mathcal{M} \ominus z\mathcal{M}) = 1$.*

Proof. Let \mathcal{V} be the subspace $\ker(M_z^{*2} M_z^2 - 2M_z^* M_z + I)$. By Lemma 4.3, the operator $T := M_z|_{\mathcal{V}}$ is a cyclic analytic 2-isometry. As $\mathcal{V} \cap \mathcal{M} = \mathcal{N}$ (say) is an invariant subspace for T , from [20, Theorem 3.2], it follows that

$$\dim(\mathcal{N} \ominus z\mathcal{N}) \leq 1. \quad (13)$$

Let \mathcal{L} be a subspace of \mathcal{M} such that $\mathcal{M} = \mathcal{N} \oplus \mathcal{L}$. Then from Lemma 4.2, it follows that $\dim \mathcal{L} \leq n$. Since M_z is an injective map, $z\mathcal{N} \cap z\mathcal{L} = \{0\}$ and $\dim \mathcal{L} = \dim z\mathcal{L}$ as \mathcal{L} is finite

dimensional. From (13), we now have

$$\begin{aligned} \dim(\mathcal{M} \ominus z\mathcal{M}) &= \dim((\mathcal{N} \oplus \mathcal{L}) \ominus z(\mathcal{N} \oplus \mathcal{L})) \\ &= \dim((\mathcal{N} \oplus \mathcal{L}) \ominus (z\mathcal{N} + z\mathcal{L})) \\ &\leq 1. \end{aligned}$$

This completes the proof of the theorem. \square

5. RELATIONSHIP BETWEEN \mathcal{H}_μ AND $\mathcal{H}(b)$

In what follows, for two reproducing kernel Hilbert spaces \mathcal{H}_1 and \mathcal{H}_2 consisting of holomorphic functions on the unit disc \mathbb{D} , we write $\mathcal{H}_1 = \mathcal{H}_2$ to mean that \mathcal{H}_1 and \mathcal{H}_2 are equal as a set but the norms are not necessarily same. In case $\mathcal{H}_1 = \mathcal{H}_2$, using closed graph theorem, it can be shown that the norms on \mathcal{H}_1 and \mathcal{H}_2 must be equivalent.

For a positive measure μ in $\mathcal{M}_+(\mathbb{T})$, we set the notation (as in [12])

$$V_\mu(z) := \int_{\mathbb{T}} \frac{d\mu(\lambda)}{|1 - \lambda\bar{z}|^2}, \quad z \in \mathbb{D}.$$

With the help of generalized local Douglas formula, for a fixed $w \in \mathbb{D}$, the following lemma computes the semi-norm of Szegő kernel $S(z, w) := \frac{1}{1 - z\bar{w}}$ in $\mathcal{H}_{\mu,n}$, in terms of $V_\mu(w)$.

Lemma 5.1. *Let μ be a finite positive Borel measure on \mathbb{T} and n be a positive integer. Then*

$$D_{\mu,n}\left(\frac{1}{1 - z\bar{w}}\right) = \frac{|w|^{2n}}{(1 - |w|^2)^n} V_\mu(w), \quad w \in \mathbb{D}.$$

Proof. Let $\lambda \in \mathbb{T}$. Then by Theorem 1.1, we have

$$\begin{aligned} D_{\lambda,n}\left(\frac{1}{1 - z\bar{w}}\right) &= D_{\sigma,n-1}\left(\frac{\frac{1}{1 - z\bar{w}} - \frac{1}{1 - \lambda\bar{w}}}{z - \lambda}\right) \\ &= D_{\sigma,n-1}\left(\frac{\bar{w}}{(1 - z\bar{w})(1 - \lambda\bar{w})}\right) \\ &= \frac{|w|^2}{|1 - \lambda\bar{w}|^2} D_{\sigma,n-1}\left(\frac{1}{1 - z\bar{w}}\right). \end{aligned}$$

To complete the proof for the case μ is point mass measure δ_λ at the point λ , we claim that

$$D_{\sigma,n-1}\left(\frac{1}{1 - z\bar{w}}\right) = \frac{|w|^{2n-2}}{(1 - |w|^2)^n}. \quad (14)$$

Since $\frac{1}{1 - z\bar{w}}$ is the reproducing kernel for the Hardy space H^2 , it is easy to verify (14) for the case $n = 1$. Suppose $n > 1$, then by definition of $D_{\sigma,n-1}(\cdot)$ and the fact that $\frac{1}{(1 - z\bar{w})^n}$ is the reproducing kernel of the weighted Bergman space $A^2((n-1)(1 - |z|^2)^{n-2}dA(z))$, we get

$$\begin{aligned} D_{\sigma,n-1}\left(\frac{1}{1 - z\bar{w}}\right) &= (n-1)|w|^{2n-2} \int_{\mathbb{D}} \frac{(1 - |z|^2)^{n-2}}{|1 - z\bar{w}|^{2n}} dA(z) \\ &= \frac{|w|^{2n-2}}{(1 - |w|^2)^n}. \end{aligned}$$

Hence equation(14) stands verified and consequently we have

$$D_{\lambda,n}\left(\frac{1}{1 - z\bar{w}}\right) = \frac{|w|^{2n}}{|1 - \lambda\bar{w}|^2(1 - |w|^2)^n}.$$

Integrating both the sides with respect to μ , we get

$$D_{\mu,n}\left(\frac{1}{1 - z\bar{w}}\right) = \int_{\mathbb{T}} D_{\lambda,n}\left(\frac{1}{1 - z\bar{w}}\right) d\mu(\lambda) = \frac{|w|^{2n}}{(1 - |w|^2)^n} V_\mu(w).$$

This completes the proof of the lemma. \square

In the following main result of this section, we prove that the spaces \mathcal{H}_μ ($m \geq 2$) and $\mathcal{H}(b)$ can not be same even as sets.

Theorem 5.2. *Let $m \geq 2$ and $\mu = (\mu_1, \dots, \mu_m)$ be an m -tuple of positive measures in $\mathcal{M}_+(\mathbb{T})$. If $\mathcal{H}_\mu = \mathcal{H}(b)$ for some $b \in H^\infty$ with $\|b\|_\infty \leq 1$ then*

$$\mu_2 = \dots = \mu_m = 0.$$

Proof. Suppose that there exists a $b \in H^\infty$ with $\|b\|_\infty \leq 1$ such that $\mathcal{H}(b) = \mathcal{H}_\mu$. Then, by an application of closed graph theorem, there exist constants $\alpha > 0$ and $\beta > 0$ such that

$$\beta \|f\|_{\mathcal{H}(b)} \leq \|f\|_\mu \leq \alpha \|f\|_{\mathcal{H}(b)}, \quad f \in \mathcal{H}(b). \quad (15)$$

Since \mathcal{H}_μ contains all functions holomorphic on a neighbourhood of $\overline{\mathbb{D}}$, it follows from [12, Theorem 2.1] that b is not an extreme point of the unit ball of H^∞ . Hence there exists a outer function a such that (b, a) is a pair. Note that the function $\frac{1}{1-z\bar{w}}$ is holomorphic on a neighbourhood of $\overline{\mathbb{D}}$ for all $w \in \mathbb{D}$. By [12, Lemma 3.4], we have

$$\left\| \frac{1}{1-z\bar{w}} \right\|_{\mathcal{H}(b)}^2 = \frac{1 + \left| \frac{b(w)}{a(w)} \right|^2}{1 - |w|^2}, \quad w \in \mathbb{D}.$$

Also by Lemma 5.1 we have

$$\left\| \frac{1}{1-z\bar{w}} \right\|_\mu^2 = \frac{1}{1 - |w|^2} + \sum_{j=1}^{m-1} \frac{|w|^{2j}}{(1 - |w|^2)^j} V_{\mu_j}(w), \quad w \in \mathbb{D}.$$

Hence by (15) we obtain

$$\frac{1}{1 - |w|^2} + \sum_{j=1}^{m-1} \frac{|w|^{2j}}{(1 - |w|^2)^j} V_{\mu_j}(w) \leq \alpha \frac{1 + \left| \frac{b(w)}{a(w)} \right|^2}{1 - |w|^2}, \quad w \in \mathbb{D}.$$

Thus for $2 \leq j \leq m-1$, we have

$$\frac{|w|^{2j}}{(1 - |w|^2)^j} V_{\mu_j}(w) \leq \alpha \frac{1 + \left| \frac{b(w)}{a(w)} \right|^2}{1 - |w|^2}.$$

Since $V_{\mu_j}(w) \geq \frac{\mu_j(\mathbb{T})}{4}$ for all $w \in \mathbb{D}$, we get that

$$\mu_j(\mathbb{T}) \leq 4\alpha(1 - |w|^2)^{j-1} \frac{1 + \left| \frac{b(w)}{a(w)} \right|^2}{|w|^{2j}}.$$

Let ζ be any point in \mathbb{T} such that $\frac{b^*(\zeta)}{a^*(\zeta)}$ exists. Since $j \geq 2$, putting $w = r\zeta$ in the above inequality and taking limit $r \rightarrow 1^-$, we get that $\mu_j(\mathbb{T}) = 0$, completing the proof of the theorem. \square

Remark 5.3. Let $\mathbf{B} : \mathbb{D} \rightarrow (\ell^2)_1$ be an analytic map with $\mathbf{B}(z) = (b_i(z))_{i=0}^\infty$, where $(\ell^2)_1$ is the closed unit ball in ℓ^2 . The space $\mathcal{H}(\mathbf{B})$ corresponding to \mathbf{B} is defined to be the reproducing kernel Hilbert space with the kernel function $\frac{1 - \sum_{i=0}^\infty b_i(z)\overline{b_i(w)}}{1 - z\bar{w}}$, $z, w \in \mathbb{D}$. The space $\mathcal{H}(\mathbf{B})$ is said to be of finite rank if there exists a $N \in \mathbb{N}$ and an analytic map $\mathbf{C} : \mathbb{D} \rightarrow (\ell^2)_1$, $\mathbf{C}(z) = (c_i(z))_{i=0}^\infty$ such that $c_i = 0$ for all $i > N$ and $\mathcal{H}(\mathbf{B}) = \mathcal{H}(\mathbf{C})$, with equality of norms. Since the backward shift operator L , defined by $(Lf)(z) = \frac{f(z) - f(0)}{z}$, $z \in \mathbb{D}$, is contractive on \mathcal{H}_μ (see [16, Lemma 2.9]), it follows from [4, Proposition 2.1] that \mathcal{H}_μ coincides with the space $\mathcal{H}(\mathbf{B})$ for some \mathbf{B} with $\mathbf{B}(0) = 0$. In the case of finite rank $\mathcal{H}(\mathbf{B})$ spaces with $\mathbf{B}(0) = 0$, if the operator M_z is bounded, then it turns out that $\overline{\text{ran}}(M_z^* M_z - I)$ is finite dimensional, (see [17, Lemma 5.1]). On the contrary, for the space \mathcal{H}_μ , where $\mu = (\mu_1, \dots, \mu_m)$, $m \geq 2$, it is worth noting that the space $\ker(M_z^* M_z - I)$ is finite dimensional. This follows easily from [16, Proposition 2.7]. Therefore, in this case, $\overline{\text{ran}}(M_z^* M_z - I)$ is infinite dimensional. Consequently,

we conclude that if $m \geq 2$, then for any $\mu = (\mu_1, \dots, \mu_m)$, $\mu_m \neq 0$, \mathcal{H}_μ can not be equal to any $\mathcal{H}(\mathbf{B})$ space of finite rank with equality of norms. It will be interesting to know whether for a fixed $\mu = (\mu_1, \dots, \mu_m)$, does there exists a $\mathcal{H}(\mathbf{B})$ space of finite rank such that $\mathcal{H}_\mu = \mathcal{H}(\mathbf{B})$ with just equivalence of norms.

Remark 5.4. The last part of [17, Theorem 1.1] says that if M_z on $\mathcal{H}(b)$ is a strict $2m$ -isometry for some $m \in \mathbb{N}$ then there exists a $\zeta \in \mathbb{T}$ such that $\mathcal{H}(b)$ is equal to \mathcal{D}_ζ^m -space defined in [17, page 3] with equivalence of norms. Theorem 5.2 indicates that the local Dirichlet space of order m in the present article differs from that in [17].

REFERENCES

- [1] J. Agler and M. Stankus, m -isometric transformations of Hilbert space, I, Integral Equations Operator Theory, **21** (4) (1995), 383–429.
- [2] J. Agler and M. Stankus, m -isometric transformations of Hilbert space, II, Integral Equations Operator Theory, **23** (1) (1995), 1–48.
- [3] J. Agler and M. Stankus, m -isometric transformations of Hilbert space, III, Integral Equations Operator Theory, **24** (4) (1996), 379–421.
- [4] A. Aleman, and B. Malman, Hilbert spaces of analytic functions with a contractive backward shift, Journal of Functional Analysis, **277** (1) (2019), 157–199.
- [5] N. Aronszajn, Theory of reproducing kernels, Transactions of the American mathematical society **68** (3) (1950), 337–404.
- [6] A. Beurling, Exceptional sets, Acta mathematica, **72** (1940), 1–13.
- [7] P. Bourdon, Cellular-indecomposable operators and Beurling’s theorem, Michigan Math. J. **33**(2) (1986), 187–193.
- [8] L. De Branges and J. Rovnyak, Square summable power series, Courier Corporation, 2015.
- [9] R. Chartrand, Multipliers and Carleson measures for $D(\mu)$, Integral Equations and Operator Theory 45.3 (2003): 309–318.
- [10] N. Chevrot, D. Guillot and T. Ransford, De Branges-Rovnyak spaces are Dirichlet spaces, J. Funct. Anal. **9** (259) (2010), 2366–2383.
- [11] J. B. Conway, A course in functional analysis, **96**. Springer, 2019.
- [12] C. Costara and T. Ransford, Which de Branges-Rovnyak spaces are Dirichlet spaces (and vice versa)?, J. Funct. Anal. **265**(12), (2013), 3204–3218.
- [13] J. Douglas, Solution of the problem of Plateau, Trans. Amer. Math. Soc. (1931), 263–321.
- [14] O. El-Fallah, K. Kellay, J. Mashreghi and Thomas Ransford, A primer on the Dirichlet space, **203**, (2014), Cambridge University Press.
- [15] O. El-Fallah, K. Kellay, H. Klaja, J. Mashreghi, and T. Ransford, Dirichlet spaces with superharmonic weights and de Branges-Rovnyak spaces, Complex Analysis and Operator Theory, **10**(1), (2016), 97–107.
- [16] S. Ghara, R. Gupta, Md. R. Reza, Analytic m -isometries and weighted Dirichlet-type spaces, to appear in Journal of Operator Theory.
- [17] S. Luo, and C. Gu, and S. Richter, Higher order local Dirichlet integrals and de Branges-Rovnyak spaces, Adv. Math. **385**, (2021), Paper No. 107748, 47.
- [18] J. Mashreghi and T. Ransford, Hadamard multipliers on weighted Dirichlet spaces, Integral Equations and Operator Theory, **91**(6), (2019), 1–13.
- [19] S. Richter and C. Sundberg, A formula for the local Dirichlet integral, Michigan Math. J **38** (3), (1991), 355–379.
- [20] S. Richter and C. Sundberg, Multipliers and invariant subspaces in the Dirichlet space, Journal of Operator Theory **28** (1), (1992), 167–186.
- [21] G. Chacón, E. Fricain, and M. Shabankhah, Carleson measures and reproducing kernel thesis in Dirichlet-type spaces, Algebra i Analiz **24** (6) (2012), 0234-0852.
- [22] M. J. Cowen and R. G. Douglas, On moduli for invariant subspaces, Invariant subspaces and other topics (Timisoara and Herculane, 1981), Operator Theory: Advances and Applications **6** (1982), pp. 65–73, Birkhäuser, Basel-Boston, Mass.
- [23] P. R. Halmos, A Hilbert space problem book, Vol. 19. Springer Science & Business Media, 2012.
- [24] V. Paulsen and M. Raghupathi, An introduction to the theory of reproducing kernel Hilbert spaces, Vol. 152. Cambridge University Press, 2016.
- [25] S. Richter, Invariant subspaces in Banach spaces of analytic functions, Transactions of the American Mathematical Society, **304.2** (1987), 585–616.
- [26] S. Richter, Invariant subspaces of the Dirichlet shift, J. reine angew. Math **386** (1988), 205–220.
- [27] S. Richter, A representation theorem for cyclic analytic two-isometries, Transactions of the American Mathematical Society **328** (1991), 325–349.

- [28] E. Rydhe, Cyclic m -isometries and Dirichlet type spaces, Journal of the London Mathematical Society, **99** (3), (2019), 733–756.
- [29] D. Sarason, Local Dirichlet spaces as de Branges-Rovnyak spaces, Proceedings of the American Mathematical Society, **125** (7), (1997), 2133–2139.
- [30] A. L. Shields, Weighted shift operators and analytic function theory, Topics in operator theory, Amer. Math. Soc., Providence, R.I., Math. Surveys, **13** (1974), 49–128.
- [31] D. A. Stegenga, Multipliers of the Dirichlet space, Illinois J. Math., **24**(1), (1980), 113–139.
- [32] G. D. Taylor, Multipliers on D_α , Transactions of the American Mathematical Society (1966), 229–240.

(S. Ghara) DEPARTMENT OF MATHEMATICS AND STATISTICS, LAVAL UNIVERSITY, QUEBEC CITY - G1V 0A6

(R. Gupta) SCHOOL OF MATHEMATICS AND COMPUTER SCIENCE, INDIAN INSTITUTE OF TECHNOLOGY GOA, GOA - 403401

(Md. R. Reza) SCHOOL OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH THIRUVANANTHAPURAM, KERALA - 695551

Email address, S. Ghara: `ghara90@gmail.com`

Email address, R. Gupta: `rajeev.iisc@live.com`

Email address, M. R. Reza: `ramiz.md@gmail.com`