

ON APPROXIMATION SPACES AND GREEDY-TYPE BASES

PABLO M. BERNÁ, HÙNG VIỆT CHU, AND EUGENIO HERNÁNDEZ

ABSTRACT. The purpose of this paper is to introduce ω -Chebyshev-greedy and ω -partially greedy approximation classes and to study their relation with ω -approximation spaces, where the latter are a generalization of the classical approximation spaces. The relation gives us sufficient conditions of when certain continuous embeddings imply different greedy-type properties. Along the way, we generalize a result by P. Wojtaszczyk as well as characterize semi-greedy Schauder bases in quasi-Banach spaces, generalizing a previous result by the first author.

CONTENTS

1. Background and main results	1
2. Preliminary results	5
3. Semi-greedy Schauder bases in quasi-Banach spaces	10
4. Approximation classes and (semi-) greedy bases	12
5. Approximation classes and partially greedy bases	17
6. Examples	22
7. Annex: Lower and upper regularity properties	27
Acknowledgments	29
References	29

1. BACKGROUND AND MAIN RESULTS

Let $(\mathbb{X}, \|\cdot\|)$ be a quasi-Banach space; that is, \mathbb{X} is a complete vector space over $\mathbb{F} = \mathbb{R}$ or \mathbb{C} , with a quasi-norm $\|\cdot\| : \mathbb{X} \rightarrow [0, \infty)$ such that

- (C1) $\|x\| = 0$ if and only if $x = 0$,
- (C2) $\|tx\| = |t|\|x\|$ for all $t \in \mathbb{F}$ and all $x \in \mathbb{X}$, and
- (C3) there is a constant $\kappa \geq 1$ so that

$$\|x + y\| \leq \kappa(\|x\| + \|y\|), \forall x, y \in \mathbb{X}.$$

Given $0 < p \leq 1$, a p -norm is a map $\|\cdot\| : \mathbb{X} \rightarrow [0, \infty)$ satisfying (C1), (C2), and

- (C4) $\|x + y\|^p \leq \|x\|^p + \|y\|^p$ for all $x, y \in \mathbb{X}$.

The first and third authors were supported by Grants PID2019-105599GB-I00 (Agencia Estatal de Investigación, Spain). This work has been supported by the Madrid Government (Spain) under the multi-annual Agreement with UAM in the line for the Excellence of the University Research Staff in the context of the V PRICIT (Regional Programme of Research and Technological Innovation).

It follows that the condition (C4) implies (C3) with $\kappa = 2^{1/p-1}$. A quasi-Banach space whose quasi-norm is a p -norm shall be called a p -Banach space. Thanks to the Aoki-Rolewicz's Theorem (see [2, 17]), we know that any quasi-Banach space \mathbb{X} is p -convex for some $0 < p \leq 1$, i.e., there is a constant C such that

$$\left\| \sum_{j=1}^n x_j \right\|^p \leq C \sum_{j=1}^n \|x_j\|^p, \forall n \in \mathbb{N}, \forall x_j \in \mathbb{X}.$$

This way, a quasi-Banach space becomes p -Banach under a suitable renorming.

We say that $\mathcal{B} = (e_n)_{n=1}^\infty$ is a semi-normalized Markushevich basis (or an M-basis or simply a basis) of \mathbb{X} if the following holds

- a1) There exists a (unique) collection $(e_n^*)_{n=1}^\infty \subset (\mathbb{X}^*, \|\cdot\|_*)$, called the biorthogonal functionals, such that $e_i^*(e_j) = \delta_{i,j}$.
- a2) There exist $c_1, c_2 > 0$ such that $0 < c_1 \leq \{\|e_n\|, \|e_n^*\|_*\} \leq c_2 < \infty$ for all $n \in \mathbb{N}$.
- a3) $\mathbb{X} = \overline{\text{span}\{e_n : n \in \mathbb{N}\}}$.
- a4) If $e_n^*(x) = 0$ for all $n \in \mathbb{N}$, then $x = 0$.

Under these conditions, every $x \in \mathbb{X}$ is represented by $x \sim \sum_{n=1}^\infty e_n^*(x)e_n$ and $\lim_{n \rightarrow \infty} e_n^*(x) = 0$. Also, if we consider the algorithm of partial sums $(S_m)_{m=1}^\infty$ defined as $S_m(x) = \sum_{n=1}^m e_n^*(x)e_n$, we say that \mathcal{B} is a Schauder basis if there is a positive constant C such that

$$\|S_m(x)\| \leq C\|x\|, \forall m \in \mathbb{N}, \forall x \in \mathbb{X}. \quad (1.1)$$

We denote by K_b the smallest constant in (1.1), which is called the basis constant.

One of the main objects studied in Approximation Theory is the nonlinear approximation spaces $\mathcal{A}_q^\alpha(\mathbb{X}, \mathcal{B})$: for $\alpha > 0$ and $0 < q < \infty$,

$$\mathcal{A}_q^\alpha(\mathcal{B}, \mathbb{X}) = \mathcal{A}_q^\alpha := \left\{ x \in \mathbb{X} : \|x\|_{\mathcal{A}_q^\alpha} := \|x\| + \left[\sum_{n=1}^\infty (n^\alpha \sigma_n(x))^q \frac{1}{n} \right]^{1/q} < \infty \right\}$$

and

$$\mathcal{A}_\infty^\alpha(\mathcal{B}, \mathbb{X}) = \mathcal{A}_\infty^\alpha := \left\{ x \in \mathbb{X} : \|x\|_{\mathcal{A}_\infty^\alpha} := \|x\| + \sup_{n \geq 1} n^\alpha \sigma_n(x) < \infty \right\},$$

where $\sigma_m(x)$ is the m -term error of approximation:

$$\sigma_m(\mathcal{B}, x)_\mathbb{X} = \sigma_m(x) := \inf \left\{ \left\| x - \sum_{n \in A} b_n e_n \right\| : |A| = m, b_n \in \mathbb{F} \right\}. \quad (1.2)$$

There have been studies on embeddings among these approximation spaces and Lorentz spaces (see [12, 14, 15, 18]). One of the recent results is given in [13]:

Theorem 1.1. [13, Theorem 1.4] *Let \mathbb{X} be a quasi-Banach space with an unconditional M-basis \mathcal{B} . Assume that $h_l(m)$ is a doubling function. Then for $\alpha > 0$ and $q \in (0, \infty]$,*

we have the following continuous embeddings

$$\ell_{k^\alpha h_r(k)}^q(\mathcal{B}, \mathbb{X}) \hookrightarrow \mathcal{A}_q^\alpha(\mathcal{B}, \mathbb{X}) \hookrightarrow \ell_{k^\alpha h_l(k)}^q(\mathcal{B}, \mathbb{X})^1,$$

where $h_l(m)$ and $h_r(m)$ are the so-called democracy functions.

Also in [13], the authors introduced a new class, called the greedy class, using the Thresholding Greedy Algorithm (TGA) [16]: a greedy sum of x of order m is given by

$$G_m^\pi(x) = \sum_{n=1}^m e_{\pi(n)}^*(x) e_{\pi(n)},$$

where π is a greedy ordering, i.e., $\pi : \mathbb{N} \rightarrow \mathbb{N}$ is a permutation such that $\text{supp}(x) \subset \pi(\mathbb{N})$ and $|e_{\pi(i)}^*(x)| \geq |e_{\pi(j)}^*(x)|$ whenever $i \leq j$. Also, $A = \{\pi(n) : 1 \leq n \leq m\}$ is called a greedy set of x of order m . The m th greedy error for $x \in \mathbb{X}$ is the quantity

$$\gamma_m(x) := \sup_{\pi} \|x - G_m^\pi(x)\|. \quad (1.3)$$

The greedy class \mathcal{G}_q^α is defined as: for $\alpha > 0$ and $q \in (0, \infty)$,

$$\mathcal{G}_q^\alpha(\mathcal{B}, \mathbb{X}) = \mathcal{G}_q^\alpha := \left\{ x \in \mathbb{X} : \|x\|_{\mathcal{G}_q^\alpha} := \|x\| + \left[\sum_{n=1}^{\infty} (n^\alpha \gamma_n(x))^q \frac{1}{n} \right]^{1/q} < \infty \right\}.$$

The class $\mathcal{G}_\infty^\alpha$ is defined in the same way as $\mathcal{A}_\infty^\alpha$ with σ_n replaced by γ_n . One of the results in [13] is that if an M-basis \mathcal{B} in a quasi-Banach space is greedy, then $\mathcal{A}_q^\alpha \approx \mathcal{G}_q^\alpha$. The converse (in the context of Banach spaces) was given by Wojtaszczyk [19]:

Theorem 1.2. [19, Theorem 3.1 (restated)] *Let \mathcal{B} be an unconditional M-basis in a Banach space \mathbb{X} .*

- (1) *If \mathcal{B} is greedy, then $\mathcal{A}_q^\alpha \approx \mathcal{G}_q^\alpha$ for all $q \in (0, \infty]$ and $\alpha > 0$.*
- (2) *If $\mathcal{A}_q^\alpha \approx \mathcal{G}_q^\alpha$ for some $q \in (0, \infty]$ and $\alpha > 0$, then \mathcal{B} is greedy.*

Statement (1) in Theorem 1.2 follows from definitions, while proving statement (2) is considerably more involved. The purpose of this paper is to study the above result in the more general context of weights and for different greedy-type bases. In particular, given $q \in (0, \infty]$ and a weight $\omega = (\omega(n))_{n=1}^\infty$ satisfying certain conditions (see definitions in Subsection 2.3), we define the following ω -approximation spaces: for $0 < q < \infty$,

$$\mathcal{A}_q^\omega := \left\{ x \in \mathbb{X} : \|x\|_{\mathcal{A}_q^\omega} := \|x\| + \left[\sum_{n=1}^{\infty} (\omega(n) \sigma_n(x))^q \frac{1}{n} \right]^{1/q} < \infty \right\};$$

for $q = \infty$,

$$\mathcal{A}_\infty^\omega := \left\{ x \in \mathbb{X} : \|x\|_{\mathcal{A}_\infty^\omega} := \|x\| + \sup_{n \geq 1} \omega(n) \sigma_n(x) < \infty \right\}.$$

These spaces were recently considered in [9]. When $\omega(n) = n^\alpha$, we recover the classical approximation space \mathcal{A}_q^α . We shall introduce three different greedy approximation classes. The first one is a generalization of \mathcal{G}_q^α : if $0 < q < \infty$ and ω is a weight,

¹Throughout this paper, $X \hookrightarrow Y$ means $X \subset Y$ and there exists $C > 0$ such that $\|x\|_Y \leq C\|x\|_X$ for all $x \in X$. Moreover, $X \approx Y$ means $X \hookrightarrow Y$ and $Y \hookrightarrow X$.

$$\mathcal{G}_q^\omega := \left\{ x \in \mathbb{X} : \|x\|_{\mathcal{G}_q^\omega} := \|x\| + \left[\sum_{n=1}^{\infty} (\omega(n)\gamma_n(x))^q \frac{1}{n} \right]^{1/q} < \infty \right\},$$

and for $q = \infty$,

$$\mathcal{G}_\infty^\omega := \left\{ x \in \mathbb{X} : \|x\|_{\mathcal{G}_\infty^\omega} := \|x\| + \sup_{n \geq 1} \omega(n)\gamma_n(x) < \infty \right\}.$$

For $\omega(n) = n^\alpha$, we recover the original greedy class introduced in [13].

In [11], Dilworth, Kalton, Kutzarova, and Temlyakov introduced a new algorithm that is an enhancement of the rate of convergence of the greedy algorithm. For $x \in \mathbb{X}$, the Thresholding Chebyshev Greedy Algorithm (TCGA) $(CG_m^\pi)_{m=1}^\infty$ is defined as:

$$\|x - CG_m^\pi(x)\| = \inf_{(a_n) \subset \mathbb{F}} \left\| x - \sum_{n=1}^m a_n e_{\pi(n)} \right\|.$$

The m th Chebyshev-greedy error for $x \in \mathbb{X}$ is

$$\vartheta_m(x) := \sup_{\pi} \|x - CG_m^\pi(x)\|.$$

Using this error, we define the Chebyshev-greedy approximation class: for $0 < q < \infty$ and a weight ω ,

$$\mathcal{CG}_q^\omega := \left\{ x \in \mathbb{X} : \|x\|_{\mathcal{CG}_q^\omega} := \|x\| + \left[\sum_{n=1}^{\infty} (\omega(n)\vartheta_n(x))^q \frac{1}{n} \right]^{1/q} < \infty \right\},$$

and for $q = \infty$, we consider the usual modification as in $\mathcal{G}_\infty^\omega$. Obviously, we have the following continuous embeddings: for any $q > 0$ and a weight ω ,

$$\mathcal{G}_q^\omega \hookrightarrow \mathcal{CG}_q^\omega \hookrightarrow \mathcal{A}_q^\omega.$$

Finally, we introduce the partially greedy class \mathcal{PG}_q^ω : for $q \in (0, \infty)$ and a weight ω ,

$$\mathcal{PG}_q^\omega = \left\{ x \in \mathbb{X} : \|x\|_{\mathcal{PG}_q^\omega} := \|x\| + \left(\sum_{n=1}^{\infty} (\omega(n)\beta_n(x))^q \frac{1}{n} \right)^{1/q} < \infty \right\},$$

where $\beta_m(x) := \|x - S_m(x)\|$.

For two functions $f(a_1, a_2, \dots)$ and $g(a_1, a_2, \dots)$, we write $f \lesssim g$ to indicate that there exists an absolute constant $C > 0$ (independent of a_1, a_2, \dots) such that $f \leq Cg$. Similarly, $f \gtrsim g$ means that $Cf \geq g$ for some constant C . Furthermore, $f \asymp g$ means that $f \lesssim g$ and $f \gtrsim g$. For two sets $A, B \subset \mathbb{N}$, we write $A < B$ to mean that $\max A < \min B$. We are ready to state the three main results of this paper.

Theorem 1.3. *Let ω be a doubling weight with $i_\omega > 0$. Let \mathcal{B} be a quasi-greedy M -basis in a quasi-Banach space \mathbb{X} .*

- (1) *If \mathcal{B} is semi-greedy, then $\mathcal{A}_q^\omega \approx \mathcal{CG}_q^\omega$ for all $q \in (0, \infty]$.*
- (2) *If \mathcal{B} is Schauder with Property (W) and $\mathcal{A}_q^\omega \approx \mathcal{CG}_q^\omega$ for some $q \in (0, \infty]$, then \mathcal{B} is semi-greedy.*

The techniques used to prove the Theorem 1.3 allow us to generalize Theorem 1.2.

Theorem 1.4. *Let ω be a doubling weight with $i_\omega > 0$. Let \mathcal{B} be an unconditional M -basis in a quasi-Banach space \mathbb{X} .*

- (1) *If \mathcal{B} is greedy, then $\mathcal{A}_q^\omega \approx \mathcal{G}_q^\omega$ for all $q \in (0, \infty]$.*
- (2) *If $\mathcal{A}_q^\omega \approx \mathcal{G}_q^\omega$ for some $q \in (0, \infty]$, then \mathcal{B} is greedy.*

Our final result is related to partially greedy bases, first introduced by Dilworth et al. [11]

Theorem 1.5. *Let ω be a doubling weight with $i_\omega > 0$. Let \mathcal{B} be a quasi-greedy M -basis in a quasi-Banach space \mathbb{X} .*

- (1) *If \mathcal{B} is partially greedy, then $\mathcal{PG}_q^\omega \hookrightarrow \mathcal{G}_q^\omega$ for all $q \in (0, \infty]$.*
- (2) *If \mathcal{B} is Schauder with Property (I) and Property (W^*) and $\mathcal{PG}_q^\omega \hookrightarrow \mathcal{G}_q^\omega$ for some $q \in (0, \infty]$, then \mathcal{B} is partially greedy.*

The above-mentioned properties, the dilation index i_ω , and other terminologies will be defined later.

Remark 1.6. Note that for no basis we have $\mathcal{G}_q^\omega \hookrightarrow \mathcal{PG}_q^\omega$. Indeed, for $0 < q < \infty$, given $m \in \mathbb{N}$, we have

$$\|e_{m+1}\|_{\mathcal{G}_q^\omega} = \|e_{m+1}\|,$$

but

$$\|e_{m+1}\|_{\mathcal{PG}^\omega} = \|e_{m+1}\| + \left(\sum_{n=1}^m \left(\omega(n) \|e_{m+1}\| \frac{1}{n} \right)^q \right)^{\frac{1}{q}} \geq \|e_{m+1}\| \omega(1) \left(\sum_{n=1}^m \frac{1}{n} \right)^{\frac{1}{q}}.$$

Hence,

$$\frac{\|e_{m+1}\|_{\mathcal{G}_q^\omega}}{\|e_{m+1}\|_{\mathcal{PG}_q^\omega}} \leq \frac{1}{\omega(1) \left(\sum_{n=1}^m \frac{1}{n} \right)^{\frac{1}{q}}} \xrightarrow{m \rightarrow \infty} 0.$$

A similar computation gives the result for $q = \infty$.

2. PRELIMINARY RESULTS

2.1. Convexity. One of the arguably most important properties of Banach spaces is convexity. In the case of p -Banach spaces, we will use the following result that is an extension of classical results of convexity. Given $0 < p \leq 1$, we put

$$\mathbf{A}_p = \frac{1}{(2^p - 1)^{1/p}}. \quad (2.1)$$

Proposition 2.1 ([1, Corollary 2.3]). *Let \mathbb{X} be a p -Banach space for some $0 < p \leq 1$. Let $(x_j)_{j \in J} \subset \mathbb{X}$ with J finite, and $g \in \mathbb{X}$. Then*

- (1) *For any scalars $(a_j)_{j \in J}$ with $0 \leq a_j \leq 1$, we have*

$$\left\| g + \sum_{j \in J} a_j x_j \right\| \leq \mathbf{A}_p \sup \left\{ \left\| g + \sum_{j \in A} x_j \right\| : A \subset J \right\}.$$

(2) For any scalars $(a_j)_{j \in J}$ with $|a_j| \leq 1$, we have

$$\left\| g + \sum_{j \in J} a_j x_j \right\| \leq \mathbf{A}_{\mathbf{p}} \sup \left\{ \left\| g + \sum_{j \in J} \varepsilon_j x_j \right\| : |\varepsilon_j| = 1 \right\}.$$

2.2. Greedy-type bases. In [16], Konyagin and Temlyakov introduced the TGA $(G_m^\pi)_{m=1}^\infty$ as we described in the previous section and defined greedy bases in Banach spaces.

Definition 2.2. An M-basis \mathcal{B} in a quasi-Banach space is greedy if $\gamma_m(x) \asymp \sigma_m(x)$.

Moreover, they characterized greedy bases in terms of unconditionality and democracy in the context of Banach spaces. In [1], the authors proved the same characterization of greediness for quasi-Banach spaces. Recall that an M-basis is K -unconditional if

$$K := \sup_{A \subset \mathbb{N}, |A| < \infty} \|P_A\| < \infty,$$

where $P_A(x) = \sum_{n \in A} e_n^*(x) e_n$ is the projection operator. To discuss democracy, we need the indicator sum

$$1_{\varepsilon A} = 1_{\varepsilon A}[\mathcal{B}, \mathbb{X}] := \sum_{n \in A} \varepsilon_n e_n,$$

where $\varepsilon = (\varepsilon_n)_{n \in A}$ with $|\varepsilon_n| = 1$ for all $n \in A$. We use the notation $|\varepsilon| = 1$.

Definition 2.3. We say that \mathcal{B} is a super-democratic basis in a quasi-Banach space if there is a positive constant C such that

$$\|1_{\varepsilon A}\| \leq C \|1_{\delta B}\|, \quad (2.2)$$

for any $|A| \leq |B| < \infty$ and $|\varepsilon| = |\delta| = 1$. The smallest constant verifying (2.2) is denoted by C_{sd} and we say that \mathcal{B} is C_{sd} -super-democratic. If $\varepsilon \equiv \delta \equiv 1$, we say that \mathcal{B} is C_d -democratic.

Equivalently, to define super-democracy, we can use the democracy functions: for each $m = 1, 2, \dots$,

$$h_r(m) := \sup_{|A| \leq m, |\varepsilon|=1} \|1_{\varepsilon A}\| \text{ and } h_l(m) := \inf_{|A| \geq m, |\varepsilon|=1} \|1_{\varepsilon A}\|.$$

Then \mathcal{B} is super-democratic if and only if

$$\sup_{m \geq 1} \frac{h_r(m)}{h_l(m)} < \infty.$$

Remark 2.4. For each $m \in \mathbb{N}$ in a p -Banach space,

$$2^{1-1/p} h_r(m) \leq \mathbf{h}_r(m) := \sup_{|A|=m, |\varepsilon|=1} \|1_{\varepsilon A}\| \leq h_r(m).$$

Indeed, if $|A| \leq N$, take any $B \subset \mathbb{N}$ such that $A \subset B$ and $|B| = N$. Indeed, we have

$$\begin{aligned} \|1_{\varepsilon A}\|^p &= \left\| \frac{1}{2}(1_{\varepsilon A} + 1_{B \setminus A}) + \frac{1}{2}(1_{\varepsilon A} - 1_{B \setminus A}) \right\|^p \\ &\leq \frac{1}{2^p} \|1_{\varepsilon A} + 1_{B \setminus A}\|^p + \frac{1}{2^p} \|1_{\varepsilon A} - 1_{B \setminus A}\|^p \leq 2^{1-p} (\mathbf{h}_r(N))^p. \end{aligned}$$

If \mathcal{B} is Schauder with basis constant K_b ,

$$h_l(m) \leq \mathbf{h}_l(m) := \inf_{|A|=m, |\varepsilon|=1} \|1_{\varepsilon A}\| \leq K_b h_l(m).$$

Moreover, Dilworth, Kalton, and Kutzarova [10] introduced the concept of semi-greedy bases.

Definition 2.5. We say that \mathcal{B} is a semi-greedy basis in a quasi-Banach space \mathbb{X} if there is a positive constant C such that

$$\vartheta_m(x) \leq C \sigma_m(x), \forall x \in \mathbb{X}, \forall m \in \mathbb{N}. \quad (2.3)$$

The smallest constant verifying (2.3) is denoted by C_{sg} , and we say that \mathcal{B} is C_{sg} -semi-greedy.

Semi-greedy Schauder bases were first characterized in Banach spaces with finite cotype in terms of quasi-greediness and democracy [10], and later on, the first author of this paper [4] (see also [5]) removed the condition of finite cotype. The notion of quasi-greediness was introduced in [16]:

Definition 2.6. We say that \mathcal{B} is a quasi-greedy basis in a quasi-Banach space \mathbb{X} if there is a positive constant C such that

$$\gamma_m(x) \leq C \|x\|, \forall x \in \mathbb{X}, \forall m \in \mathbb{N}. \quad (2.4)$$

The smallest constant verifying (2.4) is denoted by C_q , and we say that \mathcal{B} is C_q -quasi-greedy.

The characterization of semi-greediness in the context of quasi-Banach spaces is unknown and we shall prove that the same characterization of semi-greediness in Banach spaces holds for quasi-Banach spaces (see Section 3.)

In the setting of Banach spaces, Dilworth, Kalton, Kutzarova, and Temlyakov [11] introduced partially greedy bases and characterized them as being quasi-greedy and conservative. Berná [3] showed that the same result holds for quasi-Banach spaces (under a stronger notion of partially greediness.)

Definition 2.7. We say that a basis \mathcal{B} in a quasi-Banach space is partially greedy if $\gamma_m(x) \lesssim \beta_m(x)$.

Definition 2.8. We say that \mathcal{B} is a super-conservative basis in a quasi-Banach space if there is a positive constant C such that

$$\|1_{\varepsilon A}\| \leq C \|1_{\delta B}\|, \quad (2.5)$$

for any $|A| \leq |B| < \infty$, $\max A < \min B$, and $|\varepsilon| = 1, |\delta| = 1$. The smallest constant verifying (2.5) is denoted by C_{sc} and we say that \mathcal{B} is C_{sc} -super-conservative. If $\varepsilon \equiv \delta \equiv 1$, we say that \mathcal{B} is C_c -conservative.

In both Definitions 2.3 and 2.8, we require $|A| \leq |B|$. By p -convexity, this requirement can be replaced by $|A| = |B|$ (but the super-democratic (conservative) constants may differ.)

Theorem 2.9 (Berná [3]). *A Schauder basis \mathcal{B} of a quasi-Banach space is partially greedy if and only if it is quasi-greedy and is super-conservative.*

Remark 2.10. In [3], it was actually proved that an M -basis of a quasi-Banach space is strongly partially greedy if and only if it is quasi-greedy and is super-conservative. In the context of Schauder bases, being strongly partially greedy is equivalent to being partially greedy (see [3, Remark 3.5].)

2.3. Weight classes. A weight is any sequence $\omega = (\omega(n))_{n=1}^{\infty}$ of nonnegative numbers with $\omega(1) > 0$. We use the following notation:

- (1) \mathbb{W} for the set of nondecreasing and positive weights: $0 < \omega(1) \leq \omega(2) \leq \dots$, and $\lim_{n \rightarrow \infty} \omega(n) = \infty$.
- (2) \mathbb{W}_d is the set of doubling weights, i.e., $\omega \in \mathbb{W}$ and there is a positive constant $\theta \geq 1$ such that $\omega(2n) \leq \theta \cdot \omega(n)$ for all $n \in \mathbb{N}$. We say that ω is θ -doubling.

Associated with a weight ω , we consider the summing weight (see [6]): for each $m = 1, 2, \dots$,

$$\tilde{\omega}(m) := \sum_{n=1}^m \frac{\omega(n)}{n}.$$

Remark 2.11. Thanks to [6, Proposition 2.4], we know that if $\omega \in \mathbb{W}_d$, then $\tilde{\omega} \in \mathbb{W}_d$. Specifically, if ω is θ -doubling, then $\tilde{\omega}$ is $\frac{3\theta}{2}$ -doubling.

A desired relation between a weight ω and its summing weight $\tilde{\omega}$ is that $\omega(n) \asymp \tilde{\omega}(n)$. To obtain the sufficient condition for this relation to hold, we need to define the so-called dilation indices.

Definition 2.12. The lower and upper dilation sequences associated with a positive sequence $\omega = (\omega(n))_{n=1}^{\infty}$ are given by

$$\varphi_{\omega}(M) := \inf_{k \geq 1} \frac{\omega(Mk)}{\omega(k)} \text{ and } \Phi_{\omega}(M) := \sup_{k \geq 1} \frac{\omega(Mk)}{\omega(k)}, \quad M = 1, 2, 3, \dots \quad (2.6)$$

Observe that $\varphi_{\omega}(M) \leq \frac{\omega(M)}{\omega(1)} \leq \Phi_{\omega}(M)$ and

$$\varphi_{\omega}(M_1)\varphi_{\omega}(M_2) \leq \varphi_{\omega}(M_1M_2) \leq \Phi_{\omega}(M_1M_2) \leq \Phi_{\omega}(M_1)\Phi_{\omega}(M_2), \quad (2.7)$$

that is, φ_{ω} is super-multiplicative and Φ_{ω} is sub-multiplicative. If $\omega \in \mathbb{W}$, then φ_{ω} and Φ_{ω} are nondecreasing and $\varphi_{\omega}(M) \geq 1$ for all $M \in \mathbb{N}$.

Proposition 2.13. *Let $\omega \in \mathbb{W}$. The following statements are equivalent:*

- (1) $\omega \in \mathbb{W}_d$.
- (2) $\Phi_{\omega}(M) < \infty$ for all $M \in \mathbb{N}$.
- (3) $\Phi_{\omega}(M_0) < \infty$ for some $M_0 \in \mathbb{N}_{\geq 2}$.

Proof. The implications (1) \Rightarrow (2) \Rightarrow (3) are obvious. To prove (3) \Rightarrow (2), fix $M \in \mathbb{N}$. Choose $k \in \mathbb{N}$ such that $M \leq M_0^k$. Since Φ_{ω} is nondecreasing and sub-multiplicative, we have

$$\Phi_{\omega}(M) \leq \Phi_{\omega}(M_0^k) \leq \Phi_{\omega}(M_0)^k < \infty.$$

Now, we show that (2) implies (1). Taking into account that

$$\frac{\omega(2k)}{\omega(k)} \leq \Phi_{\omega}(2) < \infty,$$

we have the inequality $\omega(2k) \leq \Phi_{\omega}(2)\omega(k)$, so $\omega \in \mathbb{W}_d$ and the proof is done. \square

Definition 2.14. If $\omega \in \mathbb{W}$, we define the associated lower and upper dilation indices, respectively, by

$$i_\omega = \sup_{M>1} \frac{\ln(\varphi_\omega(M))}{\ln M} \quad \text{and} \quad I_\omega = \inf_{M>1} \frac{\ln(\Phi_\omega(M))}{\ln M}. \quad (2.8)$$

Observe from (2.7) that

$$(\varphi_\omega(M))^n \leq \varphi_\omega(M^n) \leq \Phi_\omega(M^n) \leq (\Phi_\omega(M))^n,$$

and therefore,

$$\frac{\ln(\varphi_\omega(M))}{\ln M} \leq \frac{\ln(\varphi_\omega(M^n))}{\ln M^n} \leq \frac{\ln(\Phi_\omega(M^n))}{\ln M^n} \leq \frac{\ln(\Phi_\omega(M))}{\ln M}. \quad (2.9)$$

Hence, one can replace “sup” and “inf” in (2.8) by “lim sup” and “lim inf.”

Proposition 2.15. If $\omega \in \mathbb{W}$, then $0 \leq i_\omega \leq I_\omega \leq \infty$, and

$$i_\omega = \lim_{M \rightarrow \infty} \frac{\ln(\varphi_\omega(M))}{\ln M} \text{ and } I_\omega = \lim_{M \rightarrow \infty} \frac{\ln(\Phi_\omega(M))}{\ln M}. \quad (2.10)$$

Moreover, $\omega \in \mathbb{W}_d$ if and only if $I_\omega < \infty$.

Proof. Note that $\omega \in \mathbb{W}$ implies that $1 \leq \varphi_\omega \leq \Phi_\omega$. Then (2.10) follows easily from (2.9). Now, we prove the last assertion. Assuming that $I_\omega < \infty$, we have

$$\ln(\Phi_\omega(M)) \leq I_\omega \ln M, \quad M = 1, 2, \dots$$

Thus, for every M , $\Phi_\omega(M) < \infty$ and invoking Proposition 2.13, $\omega \in \mathbb{W}_d$. For the converse, if $\omega \in \mathbb{W}_d$, by Proposition 2.13, $\Phi_\omega(M) < \infty$, hence $\frac{\ln(\Phi_\omega(M))}{\ln(M)} < \infty$ for every natural $M > 1$. Hence, $I_\omega < \infty$.^o

The last assertion is a direct consequence of Proposition 2.13: \square

Lemma 2.16. If $\omega \in \mathbb{W}_d$ with doubling constant θ and $i_\omega > 0$, then for any $q \in (0, \infty)$, $\omega^q \in \mathbb{W}_d$ with doubling constant θ^q and $i_{\omega^q} > 0$.

Proof. Trivially, $i_{\omega^q} > 0$. Also, since $\omega(2n) \leq \theta\omega(n)$, it follows that $\omega^q(2n) \leq \theta^q\omega^q(n)$ for every $n = 1, 2, \dots$ \square

Proposition 2.17. [6, Proposition 2.4] Let $\omega \in \mathbb{W}_d$ with constant θ . Then

$$\omega(N) \leq \frac{\theta}{\ln(2)} \tilde{\omega}(N), \quad N = 1, 2, \dots$$

Proposition 2.18. [6, Proposition 2.5] Let $\omega \in \mathbb{W}$. Then, $\sup_{N \geq 1} \frac{\tilde{\omega}(N)}{\omega(N)} < \infty$ if and only if $i_\omega > 0$.

Proposition 2.19. Let $\omega \in \mathbb{W}_d$ with $i_\omega > 0$. Then $\omega(N) \asymp \tilde{\omega}(N)$.

Proof. This follows immediately from Propositions 2.17 and 2.18. \square

Next, we prove an auxiliary result that shall be used in due course.

Lemma 2.20. Let $\omega \in \mathbb{W}_d$ and $\alpha > I_\omega$. There exists $C_\alpha > 0$ such that

$$\omega(Mk) \leq C_\alpha M^\alpha \omega(k), \quad \forall M, k \geq 1.$$

Proof. Let $\omega \in \mathbb{W}_d$ and $\alpha > I_\omega$. By (2.10), there exists $M_\alpha \geq 2$ such that

$$\frac{\ln(\Phi_\omega(M))}{\ln(M)} \leq \alpha, \forall M \geq M_\alpha.$$

Therefore, $\Phi_\omega(M) \leq M^\alpha$ for all $M \geq M_\alpha$. Now, by definition of $\Phi_\omega(M)$, one has

$$\omega(Mk) \leq M^\alpha \omega(k), \forall M \geq M_\alpha, k \geq 1.$$

For $M < M_\alpha$ and $k \geq 1$,

$$\omega(Mk) \leq \omega(M_\alpha k) \leq M_\alpha^\alpha \omega(k) \leq M_\alpha^\alpha M^\alpha \omega(k).$$

□

Remark 2.21. In Section 7, we will give a relation between the dilation indices and the so-called Upper and Lower Regularity Properties. The relation is useful in proving our main theorems.

2.4. The truncation operator. For each $x \in \mathbb{X}$ and each finite set $A \subset \mathbb{N}$, we define the restricted truncation operator and the truncation operator as follows:

$$\begin{aligned} \mathcal{U}(x, A) &:= \min_{n \in A} |e_n^*(x)| \sum_{n \in A} \text{sgn}(e_n^*(x)) e_n, \\ \mathcal{T}(x, A) &:= \mathcal{U}(x, A) + P_{A^c}(x). \end{aligned}$$

If A is empty or is infinite, then we use the convention that $\mathcal{U}(x, A) = 0$.

Similar operators were introduced in [10]. Write the quantity:

$$\begin{aligned} \Gamma &= \sup\{\|\mathcal{U}(x, A)\| : A \text{ is a greedy set of } x, \|x\| \leq 1\}, \\ \Upsilon &= \sup\{\|\mathcal{T}(x, A)\| : A \text{ is a greedy set of } x, \|x\| \leq 1\}. \end{aligned}$$

For quasi-Banach spaces, the authors in [1] proved the following result regarding the boundedness of these operators.

Theorem 2.22. [1, Theorem 4.12 and Theorem 4.13] *Let \mathcal{B} be a C_q -quasi-greedy basis of a quasi-Banach space \mathbb{X} . Then the restricted truncation operator is uniformly bounded, i.e., $\Gamma < \infty$. Also, if \mathbb{X} is p -Banach,*

$$\Gamma \leq C_q^2 \eta_p(C_q), \text{ and } \Upsilon \leq (C_q^p + \Gamma^p)^{1/p},$$

where, if $u > 0$,

$$\eta_p(u) := \min_{0 < t < 1} (1 - t^p)^{-1/p} (1 - (1 + \mathbf{A}_p^{-1} u^{-1} t)^{-p})^{-1/p}.$$

3. SEMI-GREEDY SCHAUDER BASES IN QUASI-BANACH SPACES

The first author [4] showed that a Schauder basis in a Banach space is semi-greedy if and only if it is quasi-greedy and super-democratic. We generalized the result to the setting of quasi-Banach spaces.

Theorem 3.1. *Assume that \mathcal{B} is a Schauder basis in a quasi-Banach space. Then \mathcal{B} is quasi-greedy and super-democratic if and only if \mathcal{B} is semi-greedy. Moreover, if \mathbb{X} is a p -Banach space,*

$$\begin{aligned} C_{sd} &\leq K_b(1 + K_b)C_{sg}^2, \\ C_q &\leq K_b C_{sg}(1 + (1 + K_b)^p C_{sg}^p)^{1/p}, \end{aligned}$$

$$C_{sg} \leq (2(C_q^2 \eta_p(C_q))^p + (2\mathbf{A}_p C_{sd} C_q^2 \eta(C_q))^p)^{1/p}.$$

Proof. Assume that \mathcal{B} is C_q -quasi-greedy and C_{sd} -super-democratic. To show the semi-greediness, we proceed as in [10]. Take $x \in \mathbb{X}$ and $m \in \mathbb{N}$. Choose $z = \sum_{n \in B} b_n e_n$ with $|B| = m$, A a greedy set of x with cardinality m and $\varepsilon \equiv \text{sgn}(e_n^*(x - z))$. Set $\alpha := \max_{n \notin A} |e_n^*(x)|$ and $\Delta_\alpha := \{n : |e_n^*(x - z)| > \alpha\}$. Thus, the set Δ_α is a greedy set for $x - z$ and $\Delta_\alpha \subset A \cup B$. Define

$$h := P_A(x) - P_A(\mathcal{T}(x - z, \Delta_\alpha)).$$

It is easy to verify that

$$x - h = \mathcal{T}(x - z, \Delta_\alpha) + P_{B \setminus A}(x - \mathcal{T}(x - z, \Delta_\alpha)).$$

Let Λ be a greedy set of $x - z$ with $|\Lambda| = |B \setminus A| = |A \setminus B|$ and $\min_{n \in \Lambda} |e_n^*(x - z)| \geq \min_{n \in A \setminus B} |e_n^*(x - z)| \geq \alpha$. On the one hand, Theorem 2.22 gives

$$\|\mathcal{T}(x - z, \Delta_\alpha)\| \leq 2^{1/p} C_q^2 \eta_p(C_q) \|x - z\|. \quad (3.1)$$

On the other hand, by Proposition 2.1, super-democracy, and Theorem 2.22,

$$\begin{aligned} \|P_{B \setminus A}(x - \mathcal{T}(x - z, \Delta_\alpha))\| &\leq 2\mathbf{A}_p \alpha \sup_{|\eta|=1} \|1_{\eta(B \setminus A)}\| \\ &\leq 2\mathbf{A}_p C_{sd} \min_{n \in \Lambda} |e_n^*(x - z)| \|1_{\varepsilon \Lambda}\| \\ &\leq 2\mathbf{A}_p C_{sd} C_q^2 \eta_p(C_q) \|x - z\|. \end{aligned} \quad (3.2)$$

Hence,

$$\|x - h\|^p \leq [2(C_q^2 \eta_p(C_q))^p + (2\mathbf{A}_p C_{sd} C_q^2 \eta(C_q))^p] \|x - z\|^p.$$

Thus, since this works for any finite greedy set A of x , the basis is C_{sg} -semi-greedy with

$$C_{sg} \leq (2(C_q^2 \eta_p(C_q))^p + (2\mathbf{A}_p C_{sd} C_q^2 \eta(C_q))^p)^{1/p}.$$

Assume now that \mathcal{B} is C_{sg} -semi-greedy. First, we prove that \mathcal{B} is super-democratic. Take A, B such that $|A| \leq |B|$, $|\varepsilon| = |\delta| = 1$, $C > (A \cup B)$ with $|C| = |A|$. Define the element $y := 1_{\varepsilon A} + 1_C$. Hence, if $m = |C|$, then C is a greedy set of y of order m . Applying the Chebyshev Greedy Algorithm for a corresponding greedy ordering π ,

$$y - CG_m^\pi(y) = 1_{\varepsilon A} + \sum_{n \in C} c_n e_n,$$

for some scalars $(c_n) \subset \mathbb{F}$. Hence, using the basis constant and semi-greediness, we have

$$\|1_{\varepsilon A}\| \leq K_b \|y - CG_m^\pi(y)\| \leq K_b C_{sg} \sigma_m(y) \leq K_b C_{sg} \|1_C\|.$$

To estimate $\|1_{\varepsilon B}\|$, we define the element $z := 1_C + 1_{\delta B}$. Arguing as before, if $m = |B|$, then B is a greedy set corresponding to a greedy ordering π' and

$$\|1_C\| \leq (1 + K_b) \|z - CG_m^{\pi'}(z)\| \leq (1 + K_b) C_{sg} \sigma_m(z) \leq (1 + K_b) C_{sg} \|1_{\delta B}\|.$$

Thus, \mathcal{B} is C_{sd} -super-democratic with

$$C_{sd} \leq K_b(1 + K_b) C_{sg}^2.$$

Next, we prove that \mathcal{B} is quasi-greedy. Let $x \in \mathbb{X}$ with finite support, A be finite greedy set of cardinality m of x and $C > \text{supp}(x)$ such that $|C| = m$. Define the

element $z := (x - P_A(x)) + \alpha 1_C$, with $\alpha = \max_{n \notin A} |e_n^*(x)|$. Hence, C is a greedy set of z , and, there is a greedy ordering π such that

$$z - CG_m^\pi(z) = (x - P_A(x)) + \sum_{n \in C} d_n e_n,$$

for some $(d_n) \subset \mathbb{F}$. Thus,

$$\begin{aligned} \|x - P_A(x)\|^p &\leq K_b^p \|z - CG_m^\pi(z)\|^p \\ &\leq K_b^p C_{sg}^p \|x + \alpha 1_C\|^p \\ &\leq K_b^p C_{sg}^p (\|x\|^p + \|\alpha 1_C\|^p). \end{aligned}$$

If we consider the element $y := x + \alpha 1_C$, a greedy set of y is A . Hence, for some greedy ordering π' , $y - CG_m^{\pi'}(y) = (x - P_A(x)) + \sum_{n \in A} a_n e_n + \alpha 1_C$. Then

$$\|\alpha 1_C\| \leq (1 + K_b) \|y - CG_m^{\pi'}(y)\| \leq (1 + K_b) C_{sg} \sigma_m(y) \leq (1 + K_b) C_{sg} \|x\|.$$

Therefore, the basis is C_q -quasi-greedy with $C_q \leq K_b C_{sg} (1 + (1 + K_b)^p C_{sg}^p)^{1/p}$. \square

4. APPROXIMATION CLASSES AND (SEMI-) GREEDY BASES

Before proving Theorem 1.3, we need the following technical propositions that generalize [13, Proposition 7.1] and [19, Lemma 3.3].

Proposition 4.1. *Let $\omega \in \mathbb{W}_d$. Let \mathcal{B} be a basis of a quasi-Banach space and f, g be two nondecreasing functions with $f, g : \mathbb{N} \rightarrow (0, \infty)$, g doubling with constant d , and*

$$\limsup_{n \rightarrow \infty} \frac{f(n)}{g(n)} = \infty. \quad (4.1)$$

Then there exist $\eta_j \geq k_j \geq 1$, $j = 1, 2, \dots$, such that

$$\lim_{j \rightarrow \infty} \frac{\eta_j}{k_j} = \infty \text{ and } \frac{f(k_j)}{g(\eta_j)} \geq \frac{\omega(\eta_j)}{\omega(k_j)}.$$

Proof. By (4.1), there exists an increasing sequence $(z_n)_{n=1}^\infty$ with $z_n \rightarrow \infty$ such that

$$\lim_{n \rightarrow \infty} \frac{f(z_n)}{g(z_n)} = \infty. \quad (4.2)$$

Given z_n , define $r : \mathbb{N} \rightarrow \mathbb{N}$ such that $2^{r(n)-1} \leq z_n < 2^{r(n)}$. Since g is doubling, for any $n, M \in \mathbb{N}$,

$$g(z_n M) \leq g(2^{r(n)} M) \leq d^{r(n)} g(M). \quad (4.3)$$

Since $\omega \in \mathbb{W}_d$, by Proposition 2.15, $I_\omega < \infty$, and we can fix $\alpha > I_\omega$. By (4.2), we take an increasing sequence $(k_j)_{j=1}^\infty$ where each k_j is some z_n such that

$$\frac{f(k_j)}{g(k_j)} \geq d^{r(j)} C_\alpha z_j^\alpha, \quad (4.4)$$

where C_α is as in Lemma 2.20. Using Lemma 2.20 and (4.4), we obtain

$$\frac{f(k_j)}{g(k_j)} \geq d^{r(j)} C_\alpha z_j^\alpha \geq d^{r(j)} \frac{\omega(z_j k_j)}{\omega(k_j)}. \quad (4.5)$$

Define $\eta_j = z_j k_j$. Since $z_j \rightarrow \infty$,

$$\lim_{j \rightarrow \infty} \frac{\eta_j}{k_j} = \infty,$$

so the first part of this proposition is proved. For the second,

$$\frac{f(k_j)}{g(\eta_j)} = \frac{f(k_j)}{g(z_j k_j)} \stackrel{(4.3)}{\geq} \frac{f(k_j)}{d^{r(j)} g(k_j)} \stackrel{(4.5)}{\geq} \frac{\omega(z_j k_j)}{\omega(k_j)} = \frac{\omega(\eta_j)}{\omega(k_j)}.$$

The proof is done. \square

Definition 4.2. Let \mathcal{B} be a basis in a quasi-Banach space. We say that \mathcal{B} has Property (W) if there exists a positive constant C such that for all $n \in \mathbb{N}$ and for all $m \geq n$, there exist $A \subset \mathbb{N}_{>m}$ with $|A| = n$ and $|\varepsilon| = 1$ such that

$$h_r(n) \leq C \|1_{\varepsilon A}\|. \quad (4.6)$$

The smallest constant in (4.6) is denoted by \mathcal{K} , and we say that \mathcal{B} has the \mathcal{K} -Property (W).

Proposition 4.3. If a basis \mathcal{B} in a p -Banach space is C_{sc} -super-conservative, then \mathcal{B} has $2^{1/p} C_{sc}$ -Property (W).

Proof. Fix $n \in \mathbb{N}$ and $m \geq n$. Choose B and $|\varepsilon| = 1$ such that $|B| = n$ and $h_r(n) \leq 2^{1/p} \|1_{\varepsilon B}\|$. Let $A = \{m + \max B + 1, \dots, m + \max B + n\}$. Since \mathcal{B} is C_{sc} -super-conservative, we have

$$h_r(n) \leq 2^{1/p} \|1_{\varepsilon B}\| \leq 2^{1/p} C_{sc} \|1_A\|.$$

Then, \mathcal{B} has $2^{1/p} C_{sc}$ -Property (W). \square

Example 4.4. All bases in Section 6 have Property (W). Subsection 6.4 gives an example of a non-conservative basis that has Property (W).

Proposition 4.5. Let \mathcal{B} be a Schauder basis with \mathcal{K} -Property (W) in a p -Banach space \mathbb{X} . Let $\omega \in \mathbb{W}_d$ with $i_\omega > 0$. Assume that there exist two sequences of integers $\eta_j \geq k_j \geq 1$, $j = 1, 2, \dots$, such that

$$\lim_{j \rightarrow \infty} \frac{\eta_j}{k_j} = \infty \text{ and } \frac{h_r(k_j)}{h_l(\eta_j)} \geq \frac{\omega(\eta_j)}{\omega(k_j)}. \quad (4.7)$$

Then $\mathcal{A}_q^\omega \hookrightarrow \mathcal{CG}_q^\omega$ does not hold for any $q \in (0, \infty]$.

Proof. Assume first that $0 < q < \infty$. For each $j \in \mathbb{N}$, choose $\Gamma_{l,j} \subset \mathbb{N}$ with $|\Gamma_{l,j}| = \eta_j$ and $|\varepsilon| = 1$ such that

$$\mathbf{h}_l(\eta_j) := \inf_{|A|=\eta_j, |\delta|=1} \|1_{\delta A}\| \geq \frac{\|1_{\varepsilon \Gamma_{l,j}}\|}{2}.$$

Since $h_l(\eta_j) \leq \mathbf{h}_l(\eta_j) \leq K_b h_l(\eta_j)$,

$$h_l(\eta_j) \gtrsim \|1_{\varepsilon \Gamma_{l,j}}\|. \quad (4.8)$$

By Property (W), there exist $|\delta| = 1$ and $\Gamma_{r,j}$ with $|\Gamma_{r,j}| = k_j \leq \eta_j$ such that $\Gamma_{r,j} \supset \Gamma_{l,j}$ and

$$h_r(k_j) \leq \mathcal{K} \|1_{\delta \Gamma_{r,j}}\|. \quad (4.9)$$

Define the element $x_j := 2 \cdot 1_{\varepsilon\Gamma_{l,j}} + 1_{\delta\Gamma_{r,j}}$. We have

$$\|x_j\|^p \leq 2^p \|1_{\varepsilon\Gamma_{l,j}}\|^p + \|1_{\delta\Gamma_{r,j}}\|^p \stackrel{(4.8)}{\leq} 2^{2p} K_b^p (h_l(\eta_j))^p + (h_r(k_j))^p. \quad (4.10)$$

Since ω is nondecreasing and $\eta_j \geq k_j$, $\omega(\eta_j) \geq \omega(k_j)$. Hence, using our hypothesis,

$$h_l(\eta_j) \leq h_r(k_j) \frac{\omega(k_j)}{\omega(\eta_j)} \leq h_r(k_j). \quad (4.11)$$

By (4.10) and (4.11), we obtain

$$\|x_j\| \lesssim h_r(k_j). \quad (4.12)$$

Now, if $n \in \{1, \dots, \eta_j\}$, a greedy set of x_j of order n is a subset of $\Gamma_{l,j}$; hence,

$$\vartheta_n(x_j) \geq \frac{\|1_{\delta\Gamma_{r,j}}\|}{1 + K_b} \geq \frac{h_r(k_j)}{\mathcal{K}(1 + K_b)}. \quad (4.13)$$

Hence,

$$\begin{aligned} \|x_j\|_{\mathcal{CG}_q^\omega} &\stackrel{(4.13)}{\gtrsim} \left(\sum_{n=1}^{\eta_j} \frac{1}{n} (\omega(n) h_r(k_j))^q \right)^{1/q} \\ &= h_r(k_j) (\tilde{\zeta}(\eta_j))^{1/q}, \end{aligned} \quad (4.14)$$

where $\zeta(j) = (\omega(j))^q$ and $\tilde{\zeta}$ is the summing weight corresponding to ζ . Applying Lemma 2.16 and Corollary 2.19, we know that

$$\zeta(n) \lesssim \tilde{\zeta}(n) \lesssim \zeta(n). \quad (4.15)$$

Thus,

$$\|x_j\|_{\mathcal{CG}_q^\omega} \gtrsim h_r(k_j) \omega(\eta_j) \quad (4.16)$$

Regarding the error $\sigma_n(x_j)$, for any $1 \leq n \leq \eta_j + k_j$,

$$\sigma_n(x_j) \leq \|x_j\| \stackrel{(4.12)}{\lesssim} h_r(k_j), \quad (4.17)$$

but if $n > k_j$, we get

$$\sigma_n(x_j) \leq 2 \|1_{\varepsilon\Gamma_{l,j}}\| \stackrel{(4.8)}{\lesssim} h_l(\eta_j). \quad (4.18)$$

Due to (4.12), (4.17), and (4.18), we obtain

$$\begin{aligned} \|x_j\|_{\mathcal{A}_q^\omega} &\lesssim h_r(k_j) + \left(\sum_{n=1}^{k_j} \frac{1}{n} (\omega(n) h_r(k_j))^q + \sum_{n=k_j+1}^{k_j+\eta_j} \frac{1}{n} (\omega(n) h_l(\eta_j))^q \right)^{1/q} \\ &\leq h_r(k_j) + \left(\tilde{\zeta}(k_j) (h_r(k_j))^q + \tilde{\zeta}(k_j + \eta_j) (h_l(\eta_j))^q \right)^{1/q} \\ &\stackrel{(4.15), \omega \in \mathbb{W}_d}{\lesssim} h_r(k_j) + ((\omega(k_j) h_r(k_j))^q + (\omega(\eta_j) h_l(\eta_j))^q)^{1/q} \\ &\stackrel{\text{hypothesis}}{\leq} h_r(k_j) + ((\omega(k_j) h_r(k_j))^q + (\omega(k_j) h_r(k_j))^q)^{1/q} \\ &\lesssim h_r(k_j) \omega(k_j). \end{aligned} \quad (4.19)$$

Writing $\eta_j = s_j k_j$ and using our hypothesis, we have that $s_j \rightarrow \infty$, and applying Theorem 7.5 gives

$$\frac{\omega(\eta_j)}{\omega(k_j)} = \frac{\omega(s_j k_j)}{\omega(k_j)} \geq C_\alpha (s_j)^\alpha \rightarrow \infty, \quad (4.20)$$

where α is positive and strictly smaller than i_ω . By (4.16), (4.19), and (4.20),

$$\frac{\|x_j\|_{\mathcal{CG}_q^\omega}}{\|x_j\|_{\mathcal{A}_q^\omega}} \gtrsim \frac{h_r(k_j)\omega(\eta_j)}{h_r(k_j)\omega(k_j)} = \frac{\omega(\eta_j)}{\omega(k_j)} \rightarrow \infty \text{ as } j \rightarrow \infty.$$

Consider now $q = \infty$. By (4.13),

$$\|x_j\|_{\mathcal{CG}_\infty^\omega} \geq \sup_{1 \leq n \leq \eta_j} \omega(n) \vartheta_n(x_j) \gtrsim \omega(\eta_j) h_r(k_j). \quad (4.21)$$

Arguing as in (4.19), we obtain

$$\begin{aligned} \|x_j\|_{\mathcal{A}_\infty^\omega} &\stackrel{(4.12)}{\lesssim} h_r(k_j) + \sup_{1 \leq n \leq k_j} \omega(n) \sigma_n(x_j) + \sup_{k_j < n \leq k_j + \eta_j} \omega(n) \sigma_n(x_j) \\ &\stackrel{(4.17), (4.18)}{\lesssim} h_r(k_j) + \omega(k_j) h_r(k_j) + \omega(\eta_j) h_l(\eta_j) \\ &\stackrel{\text{hypothesis}}{\lesssim} h_r(k_j) \omega(k_j). \end{aligned} \quad (4.22)$$

We conclude that

$$\frac{\|x_j\|_{\mathcal{CG}_\infty^\omega}}{\|x_j\|_{\mathcal{A}_\infty^\omega}} \gtrsim \frac{h_r(k_j)\omega(\eta_j)}{h_r(k_j)\omega(k_j)} = \frac{\omega(\eta_j)}{\omega(k_j)} \rightarrow \infty \text{ as } j \rightarrow \infty.$$

This completes our proof. \square

Proposition 4.6. *Let \mathcal{B} be a Schauder basis in a p -Banach space and $\omega \in \mathbb{W}_d$ with $i_\omega > 0$. If $\mathcal{A}_q^\omega \hookrightarrow \mathcal{CG}_q^\omega$ for some $q \in (0, \infty]$, then h_l is a doubling function.*

Proof. Assume that $q \in (0, \infty)$ and that h_l is not doubling.

Step 1: set up. Take sufficiently large $s > 1$. Then there exists $n_s \in \mathbb{N}$ such that

$$s h_l(n_s) \leq h_l(2n_s).$$

Choose a set M_s with $|M_s| = n_s$ and $|\varepsilon| = 1$ such that

$$\|1_{\varepsilon M_s}\|^p - \frac{1}{s^p} \leq \mathbf{h}_l^p(n_s) \leq K_b^p h_l^p(n_s) \leq K_b^p \|1_{\varepsilon M_s}\|^p.$$

Let $D \subset \mathbb{N}$ such that $M_s < D$ and $|D| = n_s$. Then

$$s^p \left(\|1_{\varepsilon M_s}\|^p - \frac{1}{s^p} \right) \leq (K_b s)^p h_l^p(n_s) \leq K_b^p h_l^p(2n_s) \leq K_b^p (\|1_{\varepsilon M_s}\|^p + \|1_D\|^p).$$

Hence,

$$(s^p - K_b^p) \|1_{\varepsilon M_s}\|^p - 1 \leq K_b^p \|1_D\|^p. \quad (4.23)$$

Partition $D = \cup_{i=1}^r V_i$ with $r = \lfloor s^{p/2} \rfloor$, and each set V_i has cardinality $\lceil n_s/r \rceil$ or $\lfloor n_s/r \rfloor$. Let V_s be a set in the partition such that $\|1_{V_s}\| = \max_{1 \leq i \leq r} \|1_{V_i}\|$. Dividing each term of (4.23) by r , we obtain

$$\frac{(s^p - K_b^p)}{r} \|1_{\varepsilon M_s}\|^p \leq K_b^p \|1_{V_s}\|^p + \frac{1}{r}.$$

Thus,

$$\|1_{\varepsilon M_s}\|^p \leq \frac{1 + rK_b^p}{s^p - K_b^p} \|1_{V_s}\|^p. \quad (4.24)$$

For sufficiently large s , (4.24) implies that $\|1_{\varepsilon M_s}\| \leq \|1_{V_s}\|$. Define $x_s := 2 \cdot 1_{\varepsilon M_s} + 1_{V_s}$. Then

$$\|x_s\|^p \leq \|1_{V_s}\|^p + 2^p \|1_{\varepsilon M_s}\|^p \leq (1 + 2^p) \|1_{V_s}\|^p. \quad (4.25)$$

Step 2: bound $\|x_s\|_{\mathcal{CG}_q^\omega}$. If $k \leq |M_s| = n_s$, a greedy set of x_s is a subset $M_{1,s}$ of M_s and so,

$$\|1_{V_s}\| \leq (1 + K_b) \left\| 1_{V_s} + 2 \cdot 1_{\varepsilon(M_s \setminus M_{1,s})} + \sum_{n \in M_{1,s}} a_n e_n \right\| \leq (1 + K_b) \vartheta_k(x_s), \quad (4.26)$$

for some $(a_n) \subset \mathbb{F}$. Let $\zeta(j) = (\omega(j))^q$ and $\tilde{\zeta}$ is the summing weight corresponding to ζ . Applying lemma 2.16 and Proposition 2.19, we know that

$$\zeta(n) \lesssim \tilde{\zeta}(n) \lesssim \zeta(n). \quad (4.27)$$

Hence, for $0 < q < \infty$,

$$\|x\|_{\mathcal{CG}_q^\omega} \geq \left(\sum_{k=1}^{n_s} (\omega(k) \vartheta_k(x_s))^q \frac{1}{k} \right)^{1/q} \stackrel{(4.26)}{\gtrsim} (\tilde{\zeta}(n_s))^{1/q} \|1_{V_s}\| \stackrel{(4.27)}{\gtrsim} \omega(n_s) \|1_{V_s}\|. \quad (4.28)$$

Step 3: bound $\|x_s\|_{\mathcal{A}_q^\omega}$. If $k \leq |V_s|$,

$$\sigma_k(x_s) \leq \|x_s\| \stackrel{(4.25)}{\lesssim} \|1_{V_s}\|. \quad (4.29)$$

If $k > |V_s|$,

$$\sigma_k(x_s) \leq 2 \|1_{\varepsilon M_s}\| \stackrel{(4.24)}{\lesssim} \left(\frac{1 + rK_b^p}{s^p - K_b^p} \right)^{1/p} \|1_{V_s}\|. \quad (4.30)$$

Define $C(s, p) := \left(\frac{1 + rK_b^p}{s^p - K_b^p} \right)^{1/p}$. Since $i_\zeta > 0$, letting $\alpha = i_\zeta/2$ and using Theorem 7.5 give

$$\frac{\zeta(n)}{\zeta(m)} \geq C_\alpha \left(\frac{n}{m} \right)^\alpha, \forall m \leq n. \quad (4.31)$$

We have

$$\begin{aligned}
\|x_s\|_{\mathcal{A}_q^\omega} &= \|x_s\| + \left(\sum_{k=1}^{|V_s|} (\omega(k)\sigma_k(x_s))^q \frac{1}{k} + \sum_{k=|V_s|+1}^{2n_s} (\omega(k)\sigma_k(x_s))^q \frac{1}{k} \right)^{1/q} \\
&\stackrel{(4.25),(4.29),(4.30)}{\lesssim} \|1_{V_s}\| \left(1 + \left(\tilde{\zeta}(|V_s|) + C(s,p)\tilde{\zeta}(2n_s) \right)^{1/q} \right) \\
&\stackrel{(4.27)}{\lesssim} \|1_{V_s}\| \left(1 + (\omega^q(|V_s|) + C(s,p)\omega^q(2n_s))^{1/q} \right) \\
&\stackrel{\omega \in \mathbb{W}_d}{\lesssim} \|1_{V_s}\| \left(1 + (\omega^q(|V_s|) + C(s,p)\omega^q(n_s))^{1/q} \right) \\
&= \|1_{V_s}\| \omega(n_s) \left(\frac{1}{\omega(n_s)} + \left(\frac{\omega^q(|V_s|)}{\omega^q(n_s)} + C(s,p) \right)^{1/q} \right) \\
&\stackrel{(4.31)}{\leq} \|1_{V_s}\| \omega(n_s) \left(\frac{1}{\omega(n_s)} + \left(\frac{1}{C_\alpha} \left(\frac{|V_s|}{n_s} \right)^\alpha + C(s,p) \right)^{1/q} \right) \\
&\stackrel{(4.28)}{\lesssim} \|x_s\|_{\mathcal{CG}_q^\omega} \left(\frac{1}{\omega(n_s)} + \left(\frac{1}{C_\alpha} \left(\frac{|V_s|}{n_s} \right)^\alpha + C(s,p) \right)^{1/q} \right). \quad (4.32)
\end{aligned}$$

Therefore,

$$\frac{\|x_s\|_{\mathcal{CG}_q^\omega}}{\|x_s\|_{\mathcal{A}_q^\omega}} \gtrsim \left(\frac{1}{\omega(n_s)} + \left(C_\alpha \left(\frac{|V_s|}{n_s} \right)^\alpha + C(s,p) \right)^{1/q} \right)^{-1} \rightarrow \infty \text{ as } s \rightarrow \infty.$$

The case when $q = \infty$ is similar. We have that $\mathcal{A}_q^\omega \hookrightarrow \mathcal{CG}_q^\omega$ does not hold for any $q \in (0, \infty]$. \square

Proof of Theorem 1.3. Item (1) follows directly from the definitions of \mathcal{A}_q^ω , \mathcal{CG}_q^ω , and semi-greediness. We prove item (2). By Theorem 3.1, it suffices to prove that \mathcal{B} is super-democratic. Assume otherwise. By Proposition 4.6, h_l is doubling. Apply Proposition 4.1 with $f = h_r$ and $g = h_l$ to obtain sequences (k_j) and (n_j) satisfying (4.7). Now Proposition 4.5 implies that $\mathcal{A}_q^\omega \hookrightarrow \mathcal{CG}_q^\omega$ does not hold, which contradicts our hypothesis. \square

Proof of Theorem 1.4. Item (1) follows directly from the definitions of \mathcal{A}_q^ω , \mathcal{G}_q^ω , and greediness. We prove item (2) using the exact argument as in the proof of Theorem 1.3. However, unlike Theorem 1.3, we do not need Property (W). In fact, we only have to consider the element $x_j = 1_{\varepsilon\Gamma_{r,j}} + 2 \cdot 1_{\varepsilon\Gamma_{l,j} \setminus (\Gamma_{r,j} \cap \Gamma_{l,j})}$ defined in [13, Proposition 7.1] and apply unconditionality instead of the Property (W) in (4.13). \square

5. APPROXIMATION CLASSES AND PARTIALLY GREEDY BASES

Our goal of this section is to prove Theorem 1.5. Recall that bounding $\sigma_m(x)$ effectively is crucial in the proof of Theorem 1.3 for greedy bases. However, for partially greedy bases, establishing an effective bound for $\beta_m(x)$ is considerably more difficult. For example, if $|\text{supp}(x)| = k$, then $\sigma_m(x) = 0$ for all $m > k$, but the same conclusion does not necessarily hold for $\beta_m(x)$. Furthermore, we have more freedom in choosing

the vector y in (1.2) to estimate $\sigma_m(x)$, while $S_m(x)$ in the definition of $\beta_m(x)$ is fixed. Hence, to have the equivalences as in Theorem 1.5, we require our bases to satisfy certain properties that allow us to estimate $\beta_m(x)$ more effectively. First, we need some definitions.

Set $\mathbb{D} = \{(m, u) \in \mathbb{N} \times \mathbb{N} : m \leq u\}$. Define the left and right restricted democracy functions as follows:

$$h_{R,l}(m, u) := \sup_{\substack{|A|=m, \max A \leq u \\ |\varepsilon|=1}} \|1_{\varepsilon A}\| \text{ and } h_{R,r}(m, u) := \inf_{\substack{|A|=m, \min A > u \\ |\varepsilon|=1}} \|1_{\varepsilon A}\|.$$

where $h_{R,r}(m, u)$ is defined on $\mathbb{N} \times \mathbb{N}$ and $h_{R,l}(m, u)$ is defined on \mathbb{D} .

Proposition 5.1. (1) *For a Schauder basis \mathcal{B} , it holds that*

$$K_b h_{R,r}(m_1, u) \geq h_{R,r}(m_2, u), \forall m_1 \geq m_2. \quad (5.1)$$

(2) *For a C_q -quasi-greedy basis \mathcal{B} , it holds that*

$$C_q h_{R,l}(m_1, u) \geq h_{R,l}(m_2, u), \forall u \geq m_1 \geq m_2, \forall u \in \mathbb{N}.$$

(3) *There exists a non-quasi-greedy Schauder basis such that*

$$\sup_{m_2 \leq m_1 \leq u} \frac{h_{R,l}(m_2, u)}{h_{R,l}(m_1, u)} = \infty.$$

Proof. (1) Let $A \subset \mathbb{N}$, $|A| = m_1$, $\min A > u$ and $|\varepsilon| = 1$. Choose B to be the set of m_2 smallest numbers in A . We have

$$K_b \|1_{\varepsilon A}\| \geq \|1_{\varepsilon B}\| \geq h_{R,r}(m_2, u).$$

Taking the inf over all sets A and $|\varepsilon| = 1$ gives the desired conclusion.

(2) Let $A \subset \mathbb{N}$, $|A| = m_2$, $\max A \leq u$ and $|\varepsilon| = 1$. Choose B to be a subset of $\{1, \dots, u\}$ such that $B \cap A = \emptyset$ and $|B| = m_1 - m_2$. Consider $x = 1_{\varepsilon A} + 1_B$. Since B is a greedy set of x , we obtain

$$\|1_{\varepsilon A}\| = \|x - 1_B\| \leq C_q \|x\| \leq C_q h_{R,l}(m_1, u).$$

Taking the sup over all sets A and $|\varepsilon| = 1$ finishes the proof.

(3) See Subsection 6.2. □

Proposition 5.2. *A basis \mathcal{B} is super-conservative if and only if*

$$\sup_{(m,u) \in \mathbb{D}} \frac{h_{R,l}(m, u)}{h_{R,r}(m, u)} < \infty.$$

Proof. Assume that \mathcal{B} is C_{sc} -super-conservative. Fix $(m, u) \in \mathbb{D}$. Choose $A, B \subset \mathbb{N}$ with $|A| = |B| = m$ and $\max A \leq u < \min B$. For any $|\varepsilon| = |\delta| = 1$, we have

$$\frac{\|1_{\varepsilon A}\|}{\|1_{\delta B}\|} \leq C_{sc},$$

so taking the sup over all A , $|\varepsilon| = 1$ and the inf over all B , $|\delta| = 1$ gives

$$\frac{h_{R,l}(m, u)}{h_{R,r}(m, u)} \leq C_{sc}.$$

Taking the sup over $(m, u) \in \mathbb{D}$ completes the proof.

Now assume that

$$\sup_{(m,u) \in \mathbb{D}} \frac{h_{R,l}(m, u)}{h_{R,r}(m, u)} < C,$$

for some constant C . Choose $|\varepsilon| = |\delta| = 1$ and $A, B \subset \mathbb{N}$ with $|A| = |B|$, $\max A < \min B$. We have

$$\|1_{\varepsilon A}\| \leq h_{R,l}(|A|, \max A) \text{ and } \|1_{\delta B}\| \geq h_{R,r}(|B|, \max A).$$

Hence,

$$\frac{\|1_{\varepsilon A}\|}{\|1_{\delta B}\|} \leq \frac{h_{R,l}(|A|, \max A)}{h_{R,r}(|B|, \max A)} < C.$$

Hence, \mathcal{B} is super-conservative. \square

The next definition expedites our introduction of Property (I).

Definition 5.3. A function $\psi : \mathbb{N} \rightarrow \mathbb{N}$ is called a characteristic function of $h_{R,l}$ if $(n, \psi(n)) \in \mathbb{D}$ for all $n \in \mathbb{N}$ and

$$\sup_u h_{R,l}(m, u) \lesssim h_{R,l}(m, \psi(m)).$$

Clearly, such a function ψ exists but is not unique.

Example 5.4. A characteristic function of $h_{R,l}$ is found as follows: let $\psi(m)$ be the smallest integer at least m such that $h_r(m) \leq 2h_{R,\ell}(m, \psi(m))$.

Definition 5.5. A basis \mathcal{B} is said to have Property (I) if

(1) we have

$$\sup_{\substack{\ell \in \mathbb{N} \cup \{0\} \\ u \in \mathbb{N}}} \frac{h_{R,r}(2^{\ell+1}u, u)}{h_{R,r}(2^\ell u, u)} < \infty, \text{ and}$$

(2) there is a characteristic function ψ of $h_{R,l}$ such that $h_{R,r}(u, u) \lesssim h_{R,r}(m, u)$ whenever $u \leq \psi(m)$.

Definition 5.6. A basis \mathcal{B} is said to have Property (W*) if there is constants $C_1, C_2 > 0$ such that for every $m \in \mathbb{N}$, there exist $A \subset \mathbb{N}_{\leq C_1 m}$ and $|\varepsilon| = 1$ such that $|A| = m$ and $\|1_{\varepsilon A}\| \leq C_2 h_l(m)$. We say \mathcal{B} has (C_1, C_2) -Property (W*).

Example 5.7. All bases in Section 6 have Property (I) and Property (W*). In particular, Subsection 6.3 gives an unconditional and conservative basis with Property (I) and Property (W*), but the basis is not democratic. Subsection 6.3 gives an unconditional basis with Property (I) and Property (W*) but is not conservative.

Proposition 5.8. Let $w \in \mathbb{W}_d$. If a Schauder basis \mathcal{B} has property (I) and is not conservative, then there exist $\eta_j \geq u_j \geq k_j \geq 1$, $j = 1, 2, \dots$, such that

$$\lim_{j \rightarrow \infty} \frac{\eta_j}{u_j} \rightarrow \infty \text{ and } \frac{h_{R,l}(k_j, u_j)}{h_{R,r}(\eta_j, u_j)} \gtrsim \frac{\omega(\eta_j)}{\omega(u_j)}.$$

Proof. Since \mathcal{B} is not conservative, Proposition 5.2 gives

$$\sup_{(m,u) \in \mathbb{D}} \frac{h_{R,l}(m, u)}{h_{R,r}(m, u)} = \infty.$$

Let $(z_n, v'_n) \in \mathbb{D}$ be chosen such that

$$\lim_{n \rightarrow \infty} \frac{h_{R,l}(z_n, v'_n)}{h_{R,r}(z_n, v'_n)} = \infty. \quad (5.2)$$

Let ψ be a characteristic function of $h_{R,l}$ that is also in the definition of Property (I). By the definition of characteristic functions, we have

$$h_{R,l}(z_n, v'_n) \lesssim h_{R,l}(z_n, \psi(z_n)). \quad (5.3)$$

We now build a sequence (v_n) :

$$v_n = \begin{cases} \psi(z_n) & \text{if } v'_n > \psi(z_n), \\ v'_n & \text{if } v'_n \leq \psi(z_n). \end{cases}$$

Note that if $v'_n > \psi(z_n)$, we have

$$\frac{h_{R,l}(z_n, v_n)}{h_{R,r}(z_n, v_n)} = \frac{h_{R,l}(z_n, \psi(z_n))}{h_{R,r}(z_n, \psi(z_n))} \stackrel{(5.3)}{\gtrsim} \frac{h_{R,l}(z_n, v'_n)}{h_{R,r}(z_n, v'_n)},$$

which, along with (5.2), implies that

$$\lim_{n \rightarrow \infty} \frac{h_{R,l}(z_n, v_n)}{h_{R,r}(z_n, v_n)} = \infty \text{ and } v_n \leq \psi(z_n). \quad (5.4)$$

Observe that $h_{R,r}(z_n, v_n) \geq K_b^{-1} \inf_n \|e_n\|$ and $h_{R,l}(z_n, v_n) \leq z_n^{1/p} \sup_n \|e_n\|$, so we can assume that $z_n, v_n \rightarrow \infty$. Let $r : \mathbb{N} \rightarrow \mathbb{N}$ be such that $2^{r(n)-1} \leq z_n \leq 2^{r(n)}$. Since $\omega \in \mathbb{W}_d$, Proposition 2.15 gives $I_\omega < \infty$. Choose $\alpha > I_\omega$ and C_α as in Lemma 2.20. Choose (k_j, u_j) to be some (z_n, v_n) such that

$$\frac{h_{R,l}(k_j, u_j)}{h_{R,r}(k_j, u_j)} \geq K_b d^{r(j)} C_\alpha z_j^\alpha \geq K_b d^{r(j)} \frac{\omega(u_j z_j)}{\omega(u_j)}, \quad (5.5)$$

where d is equal to the sup in item (1) of Property (I). Set $\eta_j = u_j z_j$. We obtain

$$\begin{aligned} h_{R,r}(\eta_j, u_j) &= h_{R,r}(u_j z_j, u_j) \stackrel{(5.1)}{\leq} K_b h_{R,r}(2^{r(j)} u_j, u_j) \\ &\leq K_b d^{r(j)} h_{R,r}(u_j, u_j) \lesssim K_b d^{r(j)} h_{R,r}(k_j, u_j), \end{aligned} \quad (5.6)$$

where the last two inequalities are due to Property (I) and the fact that $u_j \leq \psi(k_j)$. Clearly, $\eta_j / u_j \rightarrow \infty$. Furthermore,

$$\frac{h_{R,l}(k_j, u_j)}{h_{R,r}(\eta_j, u_j)} \stackrel{(5.6)}{\gtrsim} K_b^{-1} d^{-r(j)} \frac{h_{R,l}(k_j, u_j)}{h_{R,r}(k_j, u_j)} \stackrel{(5.5)}{\geq} \frac{\omega(\eta_j)}{\omega(u_j)}.$$

□

If our basis is quasi-greedy, we have an immediate corollary.

Corollary 5.9. *Let $w \in \mathbb{W}_d$. If a quasi-greedy Schauder basis \mathcal{B} has property (I) and is not conservative, then there exist $\eta_j \geq u_j \geq 1$, $j = 1, 2, \dots$, such that*

$$\lim_{j \rightarrow \infty} \frac{\eta_j}{u_j} \rightarrow \infty \text{ and } \frac{h_{R,l}(u_j, u_j)}{h_{R,r}(\eta_j, u_j)} \gtrsim \frac{\omega(\eta_j)}{\omega(u_j)}.$$

Proof. Use Propositions 5.1 item (2) and 5.8. □

Proposition 5.10. *Let \mathcal{B} be a Schauder basis with Property (I) and Property (W^*) of a p -Banach space. Let $\omega \in \mathbb{W}_d$ with $i_\omega > 0$. If \mathcal{B} is not conservative, then $\mathcal{PG}_q^\omega \hookrightarrow \mathcal{G}_q^\omega$ does not hold for any $q \in (0, \infty]$.*

Proof. We assume that $q \in (0, \infty)$. (The case $q = \infty$ is similar.)

Step 1: set up. By Proposition 5.8, there exist $\eta_j \geq u_j \geq k_j \geq 1$, $j = 1, 2, \dots$, such that

$$\lim_{j \rightarrow \infty} \frac{\eta_j}{u_j} \rightarrow \infty \text{ and } \frac{h_{R,l}(k_j, u_j)}{h_{R,r}(\eta_j, u_j)} \gtrsim \frac{\omega(\eta_j)}{\omega(u_j)}. \quad (5.7)$$

Assume that \mathcal{B} has (C_1, C_2) -Property (W^*). Choose $|\varepsilon| = 1$, $\Gamma'_{r,j} \subset \mathbb{N}_{\leq C_1 \eta_j}$ with $|\Gamma'_{r,j}| = \eta_j$ and

$$\|1_{\varepsilon \Gamma'_{r,j}}\| \leq C_2 h_l(\eta_j). \quad (5.8)$$

Define $\Gamma_{r,j} = \Gamma'_{r,j} \cap [u_j + 1, \infty)$. Then

$$\|1_{\varepsilon \Gamma_{r,j}}\| \leq (K_b + 1) \|1_{\varepsilon \Gamma'_{r,j}}\| \stackrel{(5.8)}{\leq} C_2 (K_b + 1) h_{R,r}(\eta_j, u_j). \quad (5.9)$$

Choose $|\delta| = 1$, $\Gamma_{l,j} \subset \mathbb{N}_{\leq u_j}$ with $|\Gamma_{l,j}| = k_j$ and

$$\|1_{\delta \Gamma_{l,j}}\| = h_{R,l}(k_j, u_j). \quad (5.10)$$

Set $x_j := 1_{\delta \Gamma_{l,j}} + 2 \cdot 1_{\varepsilon \Gamma_{r,j}}$. We have

$$\begin{aligned} \|x_j\|^p &\leq \|1_{\delta \Gamma_{l,j}}\|^p + 2^p \|1_{\varepsilon \Gamma_{r,j}}\|^p \stackrel{(5.9), (5.10)}{\lesssim} (h_{R,l}(k_j, u_j))^p + (h_{R,r}(\eta_j, u_j))^p \\ &\stackrel{(5.7)}{\lesssim} (h_{R,l}(k_j, u_j))^p, \end{aligned} \quad (5.11)$$

which gives

$$\|x_j\| \lesssim h_{R,l}(k_j, u_j). \quad (5.12)$$

For $n \in \{1, \dots, \eta_j - u_j\}$, a greedy set of order n of x_j is a subset of $\Gamma_{r,j}$. Therefore,

$$\|\gamma_n(x_j)\| \geq \frac{1}{K_b} \|1_{\delta \Gamma_{l,j}}\| = \frac{1}{K_b} h_{R,l}(k_j, u_j). \quad (5.13)$$

Step 2: bound $\|x_j\|_{\mathcal{G}_q^\omega}$. Let $\zeta = \omega^q$ and $\tilde{\zeta}(m) = \sum_{n=1}^m \zeta(n)/n$, $m = 1, 2, \dots$. Then

$$\begin{aligned} \|x_j\|_{\mathcal{G}_q^\omega} &\geq \left(\sum_{n=1}^{\eta_j - u_j} \frac{1}{n} (\omega(n) \|\gamma_n(x_j)\|)^q \right)^{1/q} \\ &\stackrel{(5.13)}{\gtrsim} \left(\sum_{n=1}^{\eta_j - u_j} \frac{1}{n} (\omega(n) h_{R,l}(k_j, u_j))^q \right)^{1/q} \\ &= (\tilde{\zeta}(\eta_j - u_j))^{1/q} h_{R,l}(k_j, u_j) \\ &\stackrel{(4.27)}{\gtrsim} \omega(\eta_j - u_j) h_{R,l}(k_j, u_j). \end{aligned} \quad (5.14)$$

Step 3: bound $\|x_j\|_{\mathcal{PG}_q^\omega}$. For $n \leq u_j$,

$$\|\beta_n(x_j)\| \leq (1 + K_b) \|x_j\| \stackrel{(5.12)}{\lesssim} h_{R,l}(k_j, u_j), \quad (5.15)$$

and for $u_j < n \leq C_1\eta_j$,

$$\|\beta_n(x_j)\| \leq (1 + K_b)\|1_{\Gamma_{r,j}}\| \stackrel{(5.9)}{\lesssim} h_{R,r}(\eta_j, u_j). \quad (5.16)$$

We have

$$\begin{aligned} & \|x_j\|_{\mathcal{PG}_q^\omega} \\ & \leq \|x_j\| + \left(\sum_{n=1}^{u_j} \frac{(\omega(n)\beta_n(x_j))^q}{n} + \sum_{n=u_j+1}^{\lceil C_1\eta_j \rceil} \frac{(\omega(n)\beta_n(x_j))^q}{n} \right)^{1/q} \\ & \stackrel{(5.15), (5.16)}{\lesssim} \|x_j\| + \left(\sum_{n=1}^{u_j} \frac{(\omega(n)h_{R,l}(k_j, u_j))^q}{n} + \sum_{n=u_j+1}^{\lceil C_1\eta_j \rceil} \frac{(\omega(n)h_{R,r}(\eta_j, u_j))^q}{n} \right)^{1/q} \\ & \stackrel{(5.12)}{\lesssim} h_{R,l}(k_j, u_j) + \left((h_{R,l}(k_j, u_j))^q \tilde{\zeta}(u_j) + (h_{R,r}(\eta_j, u_j))^q \tilde{\zeta}(\lceil C_1\eta_j \rceil) \right)^{1/q} \\ & \stackrel{(4.27), \omega \in \mathbb{W}_d}{\lesssim} h_{R,l}(k_j, u_j) + ((\omega(u_j)h_{R,l}(k_j, u_j))^q + (\omega(\eta_j)h_{R,r}(\eta_j, u_j))^q)^{1/q} \\ & \stackrel{(5.7)}{\lesssim} \omega(u_j)h_{R,l}(k_j, u_j). \end{aligned} \quad (5.17)$$

By (5.14) and (5.17), we obtain

$$\frac{\|x_j\|_{\mathcal{G}_q^\omega}}{\|x_j\|_{\mathcal{PG}_q^\omega}} \gtrsim \frac{\omega(\eta_j - u_j)}{\omega(u_j)} \rightarrow \infty \text{ due to Theorem 7.5 and } \frac{\eta_j - u_j}{u_j} \rightarrow \infty.$$

Hence, $\mathcal{PG}_q^\omega \hookrightarrow \mathcal{G}_q^\omega$ does not hold. \square

Proof of Theorem 1.5. Item (1) follows from the definitions of \mathcal{PG}_q^ω , \mathcal{G}_q^ω , and partial greediness. We prove item (2). Since \mathcal{B} is quasi-greedy and Schauder, by Theorem 2.9, it suffices to show that \mathcal{B} is conservative, which is clearly true due to Proposition 5.10. \square

6. EXAMPLES

All examples we consider are Banach spaces over real scalars having a Schauder basis.

6.1. The summing basis of c_0 . Let $\mathcal{B} = (e_n)$ be the canonical basis in c_0 and consider the collection $(x_n) = (\sum_{i=1}^n e_i)$, $n = 1, 2, \dots$, which is a conditional Schauder basis of c_0 . We have

$$\left\| \sum_{n=1}^{\infty} a_n x_n \right\| = \sup_{N \geq 1} \left| \sum_{n=1}^N a_n \right|.$$

6.1.1. Calculating democracy functions. It is easy to see that $h_l(N) = 1$ and $h_r(N) = N$, $\forall N \in \mathbb{N}$. Similarly, $h_{R,l}(m, u) = m$, $\forall (m, u) \in \mathbb{D}$ and $h_{R,r}(m, u) = 1$, $\forall (m, u) \in \mathbb{N} \times \mathbb{N}$. Therefore, \mathcal{B} is not conservative.

6.1.2. *Properties.* We verify each desired property below.

- (1) Property (W): for every $m, n \in \mathbb{N}$ with $m \geq n$, we let $A = \{m+1, m+2, \dots, m+n\}$. Clearly, $\|1_A\| = n = h_r(n)$.
- (2) Property (W*): for every $m \in \mathbb{N}$, let $A = \{1, 2, \dots, m\}$ and $\varepsilon = ((-1)^n)_{n=1}^m$ to have $\|1_{\varepsilon A}\| = 1 = h_l(m)$.
- (3) Property (I) is due to the fact that $h_{R,r} \equiv 1$.

6.2. **The difference basis in ℓ_1 .** Consider $\mathcal{B} = (e_n)$, the canonical basis in $\ell_1(\mathbb{N})$ and consider the following vectors:

$$x_1 = e_1, x_n = e_n - e_{n-1}, n = 2, 3, \dots$$

The collection $(x_n)_{n=1}^\infty$ is a monotone conditional Schauder basis in ℓ_1 . For $(a_n)_{n=1}^N \subset \mathbb{R}$,

$$\left\| \sum_{n=1}^N a_n x_n \right\| = \sum_{n=1}^{N-1} |a_n - a_{n+1}| + |a_N|.$$

6.2.1. *Calculating democracy functions.* We have

$$\begin{aligned} h_{R,l}(2N, 2N) &= \left\| \underbrace{(1, 1, \dots, 1, \dots)}_{\text{length } 2N} \right\| = 1 \text{ and} \\ h_{R,l}(N, 2N) &\geq \left\| \underbrace{(0, 1, 0, \dots, 0, 1, 0, \dots)}_{\text{length } 2N} \right\| = 2N. \end{aligned}$$

Hence, $h_{R,l}(2N, 2N) = o(h_{R,l}(N, 2N))$, which illustrates item (3) in Proposition 5.1.

Due to [6, Lemma 8.1], we know that

$$h_r(N) = 2N \text{ and } h_l(N) = 1.$$

Proposition 6.1. *For (x_n) as above, we have*

$$h_{R,l}(m, u) = \begin{cases} 2m & \text{if } u \geq 2m, \\ 2u - 2m + 1 & \text{if } m \leq u < 2m, \end{cases}$$

and

$$h_{R,r}(m, u) = 2.$$

Proof. If $u \geq 2m$, we have

$$h_{R,l}(m, u) \geq \left\| \underbrace{(0, 1, 0, \dots, 0, 1, 0, \dots)}_{\text{length } 2m} \right\| = 2m.$$

On the other hand, $h_{R,l}(m, u) \leq h_r(m) = 2m$. Hence, if $u \geq 2m$, $h_{R,l}(m, u) = 2m$.

If $m \leq u < 2m$, assume that u is even. The case for odd u is similar. Consider the problem of distributing m 1's among the first u spots to maximize the norm. We distribute $u/2$ 1's as follows:

$$\left\| \underbrace{(0, 1, 0, \dots, 0, 1, 0, \dots)}_{\text{length } u} \right\| = u.$$

There are $(m - u/2)$ 1's remaining to be distributed. Note that putting an 1 into the first spot reduces the norm by 1, while putting an 1 into other spots reduces the norm by 2.

Case 1: if $m - u/2 = 1$, we put the only remaining 1 into the first spot to obtain norm $u - 1 = 2u - 2m + 1$.

Case 2: if $m - u/2 \geq 2$, we put an 1 into the first spot and put other 1's into other unfilled spots to obtain norm $u - 1 - 2(m - u/2 - 1) = 2u - 2m + 1$.

In conclusion, $h_{R,l}(m, u) = 2u - 2m + 1$ if $m \leq u < 2m$.

It is easy to see that $h_{R,r}(m, u) = 2$. \square

Corollary 6.2. *The difference basis \mathcal{B} is not conservative.*

6.2.2. *Properties.* We verify each desired property :

- (1) Property (W): for every $m, n \in \mathbb{N}$ with $m \geq n$, we let $A = \{m + 1, m + 3, \dots, m + 2n - 1\}$. Clearly, $\|1_A\| = 2n = h_r(n)$.
- (2) Property (W*): for every $m \in \mathbb{N}$, let $A = \{1, 2, \dots, m\}$ to have $\|1_A\| = 1 = h_l(m)$.
- (3) Property (I) is due to the fact that $h_{R,r} \equiv 2$.

6.3. **A modification of the Schreier space.** We shall use the same example as in [8, Proposition 6.10], which is a modification of the Schreier space. Let \mathbb{S} be the completion of c_{00} under the following norm:

$$\|(x_1, x_2, x_3, \dots)\|_{\mathbb{S}} = \sup_{F \in \mathcal{F}} \sum_{i \in F} |x_i|,$$

where $\mathcal{F} = \{F \subset \mathbb{N} : \sqrt{\min F} \geq |F|\}$. It is easy to check that the canonical basis \mathcal{B} is an 1-unconditional and 1-conservative monotone Schauder basis of \mathbb{S} .

6.3.1. *Calculating democracy functions.* Let $N \in \mathbb{N}$. By the definition of $\|\cdot\|_{\mathbb{S}}$, we have

$$h_r(N) = N \text{ and } h_l(N) = \|(\underbrace{1, \dots, 1}_N, 0, \dots)\|_{\mathbb{S}}.$$

For $M \in \mathbb{N}$ and $N \in \mathbb{N} \cup \{0\}$, set $x_{N,M} := \sum_{n=N+1}^{N+M} e_n$ and write $x_{N,M} = (x_1, x_2, x_3, \dots)$.

Proposition 6.3. *We have*

$$\|x_{N,M}\|_{\mathbb{S}} \lesssim \sqrt{N + M}.$$

Proof. Let $F \in \mathcal{F}$ with $\min F = N + j_0$ for some $1 \leq j_0 \leq M$.

Case 1: $\min F + \lfloor \sqrt{\min F} \rfloor - 1 \leq N + M$; equivalently, $j_0 + \lfloor \sqrt{N + j_0} \rfloor \leq M + 1$, which implies that

$$j_0 \leq M + \frac{5}{2} - \sqrt{M + N + 9/4} =: f(M, N). \quad (6.1)$$

If $f(M, N) < 1$, then Case 1 cannot happen. If $f(M, N) \geq 1$, then

$$\sum_{i \in F} |x_i| \leq \lfloor \sqrt{\min F} \rfloor \leq \sqrt{N + j_0} \stackrel{(6.1)}{\lesssim} \sqrt{N + M}.$$

Case 2: $\min F + \lfloor \sqrt{\min F} \rfloor - 1 \geq N + M$; equivalently, $j_0 + \lfloor \sqrt{N + j_0} \rfloor \geq M + 1$, which implies that

$$j_0 \geq M + \frac{3}{2} - \sqrt{M + N + 5/4} =: g(M, N). \quad (6.2)$$

Hence,

$$\begin{aligned} \sum_{i \in F} |x_i| &\leq (N + M) - \min F + 1 = M - j_0 + 1 \\ &\stackrel{(6.2)}{\leq} M - g(M, N) + 1 \lesssim \sqrt{M + N}. \end{aligned}$$

□

Proposition 6.4. *For $M \in \mathbb{N}$ and $N \in \mathbb{N} \cup \{0\}$,*

$$\|x_{N,M}\|_{\mathbb{S}} \gtrsim \begin{cases} M & \text{if } N \geq M^2 - 1, \\ \sqrt{N + M} & \text{if } N \leq M^2 - 1. \end{cases}$$

Proof. Let $F = \{N + j_0, \dots, N + M\}$, where $1 \leq j_0 \leq M$ is the smallest such that $\sqrt{N + j_0} \geq M - j_0 + 1$, i.e., $F \in \mathcal{F}$. Equivalently,

$$j_0 \geq M + 3/2 - \sqrt{M + N + 5/4} =: g(M, N).$$

Case 1: If $g(M, N) \leq 1$; equivalently, $N \geq M^2 - 1$, we choose $j_0 = 1$ and obtain $\|x_{N,M}\|_{\mathbb{S}} \geq M$.

Case 2: If $g(M, N) \geq 1$; equivalently, $N \leq M^2 - 1$, we choose $j_0 = \lceil g(M, N) \rceil$. Then

$$\|x_{N,M}\|_{\mathbb{S}} \geq M - j_0 + 1 \geq M - g(M, N) \gtrsim \sqrt{M + N}.$$

□

The following corollaries are immediate from Propositions 6.3 and 6.4.

Corollary 6.5. *We have*

$$h_l(N) = \|\underbrace{(1, \dots, 1)}_N, 0, \dots\|_{\mathbb{S}} \asymp \sqrt{N},$$

and so \mathcal{B} is not democratic.

Corollary 6.6. *For $(m, u) \in \mathbb{D}$, we have*

$$h_{R,l}(m, u) \lesssim \sqrt{u} \text{ and } h_{R,l}(m, u) \gtrsim \begin{cases} m & \text{if } u \geq m^2 + m - 1 \\ \sqrt{u} & \text{if } u \leq m^2 + m - 1. \end{cases} \quad (6.3)$$

Proof. Observe that $h_{R,l}(m, u) = \|x_{u-m,m}\|$ and apply Propositions 6.3 and 6.4. □

Corollary 6.7. *For $(m, u) \in \mathbb{N} \times \mathbb{N}$, we have*

$$h_{R,r}(m, u) \lesssim \sqrt{u + m} \text{ and } h_{R,r}(m, u) \gtrsim \begin{cases} m & \text{if } u \geq m^2 - 1 \\ \sqrt{u + m} & \text{if } u \leq m^2 - 1. \end{cases} \quad (6.4)$$

Proof. Observe that $h_{R,r}(m, u) = \|x_{u,m}\|$ and apply Propositions 6.3 and 6.4. □

6.3.2. Properties.

- (1) Property (W): for every $m, n \in \mathbb{N}$ with $m \geq n$, we let $A = \{m^2 + 1, m^2 + 2, \dots, m^2 + n\}$. Clearly, $\|1_A\| = n = h_r(n)$.
- (2) Property (W*): for every $m \in \mathbb{N}$, let $A = \{1, 2, \dots, m\}$ to have $\|1_A\| \asymp \sqrt{m} \asymp h_l(m)$.
- (3) Property (I): fix $u \in \mathbb{N}$ and $\ell \in \mathbb{N} \cup \{0\}$. We can assume that $u \leq 2^{2\ell}u^2 - 1$ since the only case that the inequality fails is when $u = 1$ and $\ell = 0$. We have

$$h_{R,r}(2^{\ell+1}u, u) \lesssim \sqrt{2^{\ell+1}u + u} \leq \sqrt{2}\sqrt{2^\ell u + u} \lesssim h_{R,r}(2^\ell u, u),$$

where the last inequality is due to (6.4) and the fact that $u \leq 2^{2\ell}u^2 - 1$. Next, we choose a suitable characteristic function of $h_{R,l}$. The function $\psi(m) = (m^2 - 1) \vee 1$ is a candidate because

$$\sup_u h_{R,l}(m, u) \leq m \stackrel{(6.3)}{\lesssim} h_{R,l}(m, (m^2 - 1) \vee 1).$$

We check that $h_{R,r}(u, u) \lesssim h_{R,r}(m, u)$ whenever $u \leq \psi(m)$: for $m \geq 2$, $\psi(m) = m^2 - 1$,

$$h_{R,r}(u, u) \stackrel{(6.4)}{\lesssim} \sqrt{u} < \sqrt{u + m} \stackrel{(6.4)}{\lesssim} h_{R,r}(m, u).$$

Therefore, the canonical basis has Property (I).

6.4. An unconditional basis with Property (I) and Property (W*) but is not conservative. We shall consider the example of a basis given in [7, Subsection 5.4]. Let $s_n = \sum_{i=1}^n \frac{1}{i}$, and Π be the set of all permutations of \mathbb{N} . Let \mathbb{X} be the completion of c_{00} with respect to the following norm:

$$\|(x_1, x_2, x_3, \dots)\| = \max \left\{ \sup_n \frac{1}{\sqrt{s_n}} \sup_{\pi \in \Pi} \sum_{i=1}^n \frac{|x_{\pi(i)}|}{i^{1/2}}, \left(\sum_i |x_{2i}|^2 \right)^{1/2} \right\}.$$

The canonical basis \mathcal{B} is an 1-unconditional and non-conservative Schauder basis. Indeed, letting $A_N = \{2, 4, 6, \dots, 2N\}$ and $B_N = \{2N + 1, 2N + 3, \dots, 4N - 1\}$, we have

$$\frac{\|1_{A_N}\|}{\|1_{B_N}\|} \gtrsim \frac{\sqrt{N}}{\sqrt{N}/\sqrt{\ln(N+1)}} = \sqrt{\ln(N+1)} \rightarrow \infty.$$

Hence, \mathcal{B} is not conservative.

6.4.1. Calculating democracy functions. For $N \in \mathbb{N}$, it is obvious that

$$h_r(N) = \sqrt{N} \text{ and } h_\ell(N) \asymp \frac{\sqrt{N}}{\sqrt{\ln(N+1)}}.$$

Similarly, for $(m, u) \in \mathbb{D}$,

$$h_{R,l}(m, u) \asymp \sqrt{m};$$

for $(m, u) \in \mathbb{N} \times \mathbb{N}$,

$$h_{R,r}(m, u) \asymp \frac{\sqrt{m}}{\sqrt{\ln(m+1)}}.$$

6.4.2. *Properties.* We verify each desired property below.

- (1) Property (W): for every $m, n \in \mathbb{N}$ with $m \geq n$, we let $A = \{j, j+2, \dots, j+2(n-1)\}$, where j is the smallest even integer greater than m . Clearly, $\|1_A\| = \sqrt{n} = h_r(n)$.
- (2) Property (W*): for every $n \in \mathbb{N}$, let $A = \{1, 3, \dots, 2n-1\}$ to have $\|1_A\| \asymp h_l(n)$.
- (3) Property (I): Pick $u \in \mathbb{N}$ and $\ell \in \mathbb{N} \cup \{0\}$. Then

$$\frac{h_{R,r}(2^{\ell+1}u, u)}{h_{R,r}(2^\ell u, u)} \asymp \frac{\sqrt{2^{\ell+1}u}/\sqrt{\ln(2^{\ell+1}u)}}{\sqrt{2^\ell u}/\sqrt{\ln(2^\ell u)}} = O(1).$$

Furthermore, choose the characteristic function $\psi(m) = m$. For all $u \leq m = \psi(m)$, we have

$$h_{R,r}(u, u) \asymp \frac{\sqrt{u}}{\sqrt{\ln(u+1)}} \leq \frac{\sqrt{m}}{\sqrt{\ln(m+1)}} \asymp h_{R,r}(m, u).$$

Corollary 6.8. *For the basis \mathcal{B} above, we have*

$$\mathcal{PG}_q^\omega \not\rightarrow \mathcal{G}_q^\omega, \forall \omega \in \mathbb{W}_d \text{ with } i_\omega > 0, \forall q \in (0, \infty].$$

7. ANNEX: LOWER AND UPPER REGULARITY PROPERTIES

Duality for greedy-type bases has been studied in [11]. Some of their results depend on two properties: the upper regularity property (URP) and lower regularity property (LRP). The purpose of this section is to show that URP and LRP are equivalent to the upper dilation index of the parameter been smaller than 1 and the lower one being greater than zero, respectively.

Definition 7.1. A positive sequence $\omega = (\omega(n))_{n=1}^\infty$ has the lower regularity property, denoted by $\omega \in LRP$, if there exist $\alpha > 0$ and $C_\alpha > 0$ such that

$$\omega(N) \geq C_\alpha \left(\frac{N}{k}\right)^\alpha \omega(k), \forall N \geq k. \quad (7.1)$$

A positive sequence ω has the upper regularity property, denoted by $\omega \in URP$, if there exist $\beta < 1$ and $C_\beta \geq 1$ such that

$$\omega(N) \leq C_\beta \left(\frac{N}{k}\right)^\beta \omega(k), \forall N \geq k. \quad (7.2)$$

We write $\omega \in LRP(\alpha)$ and $\omega \in URP(\beta)$ when (7.1) and (7.2) hold, respectively.

Proposition 7.2. *Let $\omega \in \mathbb{W}$. Then*

- (1) $\omega \in LRP(\alpha)$ if and only if there exists $c > 0$ such that

$$\varphi_\omega(M) \geq cM^\alpha, \forall M \geq 1.$$

- (2) $\omega \in URP(\beta)$ if and only if there exists $c > 0$ such that

$$\Phi_\omega(M) \leq cM^\beta, \forall M \geq 1.$$

Proof. We prove (1). (The proof of (2) is similar.) If $\omega \in LRP(\alpha)$, then

$$\omega(Mk) \geq C_\alpha M^\alpha \omega(k), \forall M, k \in \mathbb{N}.$$

Therefore, $\varphi_\omega(M) \geq C_\alpha M^\alpha$ for all M .

Conversely, suppose that $c' := \inf_{M \geq 1} \varphi_\omega(M)/M^\alpha > 0$. Let $N > k$ and choose $M \geq 1$ such that $Mk < N \leq (M+1)k$. Then

$$\omega(N) \geq \omega(Mk) \geq c' M^\alpha \omega(k) \geq c' \left(\frac{M+1}{2} \right)^\alpha \omega(k) \geq c' 2^{-\alpha} (N/k)^\alpha \omega(k).$$

Therefore, $\omega \in LRP(\alpha)$ with constant $c'/2^\alpha$. \square

Proposition 7.3. *For $\omega \in \mathbb{W}$, the following are equivalent:*

- (1) $\omega \in LRP$.
- (2) $\lim_{M \rightarrow \infty} \varphi_\omega(M) = \infty$.
- (3) *There exists $M_0 \geq 2$ such that $\varphi_\omega(M_0) > 1$.*

Proof. (1) \Rightarrow (2) follows from Proposition 7.2, and (2) \Rightarrow (3) is trivial. We show that (3) \Rightarrow (1). Let $\lambda = \varphi_\omega(M_0) > 1$ and $\alpha = \ln \lambda / \ln M_0$. Then

$$\omega(M_0 k) \geq \lambda \omega(k) = M_0^\alpha \omega(k), \forall k \in \mathbb{N}.$$

Hence, if $k < N$, we can find $j \in \mathbb{N}_0$ such that $M_0^j \leq N/k < M_0^{j+1}$ to have

$$\omega(N) \geq \omega(M_0^j k) \geq M_0^{j\alpha} \omega(k) \geq M_0^{-\alpha} (N/k)^\alpha \omega(k).$$

Thus, we have shown that $\omega \in LRP(\alpha)$. \square

Similarly, one can prove the following.

Proposition 7.4. *For $\omega \in \mathbb{W}$, the following are equivalent:*

- (1) $\omega \in URP$
- (2) $\omega \in \mathbb{W}_d$ and $\lim_{M \rightarrow \infty} \Phi_\omega(M)/M = 0$
- (3) *There exists $M_0 \geq 2$ such that $\Phi_\omega(M_0)/M_0 < 1$.*

We now deduce the main result in this section.

Theorem 7.5. *Let $\omega \in \mathbb{W}$. Then*

- (1) $\omega \in LRP$ if and only if $i_\omega > 0$. Moreover,

$$i_\omega = \sup\{\alpha > 0 : \omega \in LRP(\alpha)\}. \quad (7.3)$$

- (2) $\omega \in URP$ if and only if $I_\omega < 1$. Moreover,

$$I_\omega = \inf\{\beta < 1 : \omega \in URP(\beta)\}. \quad (7.4)$$

Proof. We prove (1). (The proof of (2) is similar.) If $\omega \in LRP(\alpha)$, then Proposition 7.2.i gives $i_\omega \geq \alpha$. Hence, $\omega \in LRP$ implies $i_\omega > 0$. Conversely, if $i_\omega > 0$, by (2.8), there exists $M_0 \geq 2$ such that $\varphi_\omega(M_0) > 1$. So $\omega \in LRP$ by Proposition 7.3. Finally, it remains to prove “ \leq ” in (7.3). That is, we must show that if $0 < \alpha < i_\omega$ then $\omega \in LRP(\alpha)$. Indeed, by definition of i_ω , there exists $M_\alpha \geq 2$ such that $\ln(\varphi_\omega(M))/\ln M \geq \alpha$ for all $M \geq M_\alpha$. Then we conclude using Proposition 7.2 item (1). \square

Remark 7.6. The sequence $\omega(n) = \sqrt{n}(\ln(n+1))^\gamma$, $\gamma \in \mathbb{R}$, shows that, in general, it may not hold that $\omega \in LRP(i_\omega)$ or $\omega \in URP(I_\omega)$. Indeed, in this case $i_\omega = I_\omega = 1/2$, but

$$\varphi_\omega(M) \asymp \sqrt{M}/(\ln(M+1))^{\gamma^-} \text{ and } \Phi_\omega(M) \asymp \sqrt{M}(\ln(M+1))^{\gamma^+}.$$

Hence, by Proposition 7.2, $\omega \notin LRP(1/2)$ if $\gamma < 0$, and $\omega \notin URP(1/2)$ if $\gamma > 0$.

ACKNOWLEDGMENTS

The authors would like to thank O. Blasco, G. Garrigós, and T. Oikhberg for the permission to write the results of Subsection 2.3 since these notes were in a previous version of [6] that was never published. Also, thanks to Miguel Berasategui for some discussions during the elaboration of the paper.

REFERENCES

- [1] F. Albiac, J. L. Ansorena, P. M. Berná, and P. Wojtaszczyk, Greedy approximation for biorthogonal systems in quasi-Banach spaces, *Dissertationes Math.* **560** (2021), 1–88.
- [2] T. Aoki, On a certain class of linear metric spaces, *Proc. Imp. Acad. Tokyo* **18** (1942), 588–594.
- [3] P. M. Berná, A note on partially-greedy bases in quasi-Banach spaces, *Studia Math.* **259** (2021), 225–239.
- [4] P. M. Berná, Equivalence between almost-greedy and semi-greedy bases, *J. Math. Anal. Appl.* **470** (2019), 218–225.
- [5] P. M. Berná, Characterization of weight-semi-greedy bases, *J. Fourier Anal. Appl.* **26** (2020), 1–21.
- [6] P. M. Berná, Ó. Blasco, G. Garrigós, E. Hernández, and T. Oikhberg, Embeddings and Lebesgue inequalities for greedy algorithms, *Constr. Approx.* **48** (2018), 415–451.
- [7] P. M. Berná and H. V. Chu, On some characterizations of greedy-type bases, *Expositiones Mathematicae* **40**, no. 4 (2022), 1135–1158.
- [8] P. M. Berná, S. J. Dilworth, D. Kutzarova, T. Oikhberg, and B. Wallis, The weighted property (A) and the greedy algorithm, *J. Approx. Theory* **248** (2019), 1–18.
- [9] F. Cobos, O. Domínguez, and T. Kühn, Approximation and entropy numbers of embeddings between approximation spaces, *Constr. Approx.* **47** (2018), 453–486.
- [10] S. J. Dilworth, N. J. Kalton, and D. Kutzarova, On the existence of almost greedy bases in Banach spaces, *Studia Math.* **159** (2003), 67–101.
- [11] S. J. Dilworth, N. J. Kalton, D. Kutzarova, and V. N. Temlyakov, The thresholding greedy algorithm, greedy bases, and duality, *Constr. Approx.* **19** (2003), 575–597.
- [12] R. A. DeVore and V. N. Temlyakov, Some remarks on greedy algorithms, *Adv. Comp. Math.* **5** (1996), 113–187.
- [13] G. Garrigós, E. Hernández, and M. de Natividade, Democracy functions and optimal embeddings for approximation spaces, *Adv. Comput. Math.* **37** (2012), 255–283.
- [14] R. Gribonval and M. Nielsen, Some remarks on nonlinear approximation approximation with Schauder bases, *East. J. of Approximation* **7** (2001), 1–19.
- [15] G. Kerkycharian and D. Picard, Nonlinear approximation and Muckenhoupt weights, *Constr. Approx.* **24** (2006), 123–156.
- [16] S. V. Konyagin and V. N. Temlyakov, A remark on greedy approximation in Banach spaces, *East. J. Approx.* **5** (1999), 365–379.
- [17] S. Rolewicz, Locally bounded linear topolnical spaces, *Bull. Acad. Polon. Sci.* **5** (1957), 471–473.
- [18] J. Schreier, Ein gegenbeispiel zur theorie der schwachen konvergentz, *Studia Math.* **2** (1962), 58–62.
- [19] P. Wojtaszczyk, On left democracy function, *Funct. Approx. Comment. Math.* **50** (2014), 207–214.

Email address: pablo.berna@cunef.edu

DEPARTAMENTO DE MÉTODOS CUANTITATIVOS, CUNEF UNIVERSIDAD, 28040 MADRID, SPAIN

Email address: hungchu2@illinois.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA,
IL 61820, USA

Email address: eugenio.hernandez@uam.es

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA DE MADRID, 28049 MADRID,
SPAIN