Integral Transforms and \mathcal{PT} -symmetric Hamiltonians

M. W. AlMasri¹ CyberSecurity and Systems Research Unit, ISI-USIM, Bandar Baru Nilai, 71800 Nilai, Malaysia

M. R. B. Wahiddin CyberSecurity and Systems Research Unit, ISI-USIM Pusat Tamhidi, USIM , Bandar Baru Nilai, 71800 Nilai, Negeri Sembilan. Malaysia

Abstract

Motivated by the fact that twice the Fourier transform plays the role of parity operator. We systematically study integral transforms in the case of \mathcal{PT} -symmetric Hamiltonian. First, we obtain a closed analytical formula for the exponential Fourier transform of a general \mathcal{PT} -symmetric Hamiltonian. Using the Segal-Bargmann transform, we investigate the effect of the Fourier transform on the eigenfunctions of the original Hamiltonian. As an immediate application, we comment on the holomorphic representation of non-Hermitian spin chains, in which the Hamiltonian operator is written in terms of analytical phase-space coordinates and their partial derivatives in the Bargmann space rather than matrices in the vector Hilbert space. Finally, we discuss the effect of integral transforms in the study of the Swanson Hamiltonian.

¹Email: mwalmasri2003@gmail.com

1 Introduction

In [1], Bender and Boettcher proved the reality of energy eigenvalues for a large class of non-Hermitian Hamiltonians numerically. The common property between these Hamiltonians is their invariance under the action of both time-reversal \mathcal{T} and parity \mathcal{P} operations, i.e., $[\mathcal{PT}, H] = 0$. Since then, many studies have been done in this area with potential applications in optics and other areas of physics such as, open quantum systems, supersymmetric quantum mechanics, topological matter, cold atoms, and magnonic waveguides [2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25].

In quantum mechanics, wave functions can be written both in the position (x-space) and momentum (p-space) spaces using the exponential Fourier transform [26, 27, 28],

$$\phi(p) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{-ipx/\hbar} \psi(x) \, dx, \qquad (1.1)$$

and its inverse transform

$$\psi(x) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} e^{ipx/\hbar} \phi(p) \, dp. \tag{1.2}$$

The Schrödinger equation in the momentum space reads

$$(E - \frac{p^2}{2m})\phi(p) = \frac{1}{\sqrt{2\pi\hbar}} \int_{-\infty}^{\infty} \tilde{V}(p - p') \ \phi(p') \ dp'$$
(1.3)

which is an integral equation. In some situations, it is more convenient to work in the momentum-space representation such as delta-potential, i.e., $V = A \, \delta(x)[27, 28, 29, 30]$. Furthermore, in one version of non-commutative quantum mechanics, we have the canonical commutation relations defined as $[p_i, p_j] = 0, [x_i, x_j] = i\theta_{ij}, [x_i, p_j] = i\hbar\delta_{ij}$ where θ_{ij} is an antisymmetric tensor with dimension $(\text{length})^2[31]$. In this case, the ordinary product of position-dependent functions is promoted to the Moyal star-product $f \star g(\mathbf{x}) = e^{i\theta_{ij}\partial_i^{(1)}\partial_j^{(2)}} f(\mathbf{x}_1)g(\mathbf{x}_2)|_{\mathbf{x}_1=\mathbf{x}_2=\mathbf{x}}[32, 33, 34]$. Consequently, working in the momentum-space might be more efficient since the momentumdependent functions are multiplied in the usual way. Noncommutative quantum mechanics appears naturally in the study of charged particle motion under the effect of magnetic field and thus it has deep connection with Landau levels, anomalous Hall effect in ferromagnetic metals and in the study of magnetic skyrmions [35, 36, 37, 38, 39, 40, 41, 42, 43]. In this work, we present a systematic study of \mathcal{PT} -symmetric Hamiltonians under the action of integral transforms such as the Fourier transform. Moreover, we use the Segal-Bargmann transform to explore the effect of the Fourier transform on eigenfunctions. Furthermore, the Bargmann representation of spin operators can be used in the study of non-Hermitian spin chains. In this case, spin operators are written in terms of orthonormal phase-space coordinates obtained from the Segal-Bargmann transform of the coordinate-dependent wavefunctions. Finally, we apply the developed formalism to the case of Swanson Hamiltonians.

2 Integral Transforms of \mathcal{PT} -symmetric Hamiltonians

The integral equation problem is to find the solution to:

$$h(x)f(x) = g(x) + \lambda \int_{a}^{b} K(x,y)f(y)dx,$$
(2.1)

We are given h(x), g(x), K(x, y) and want to determine f(x). The quantity λ is a parameter that may be complex in general, and K(x, y) is called the kernel of the integral equation. If h(x) = 0, we may take $\lambda = -1$ without loss of generality, and we end up with a Fredholm equation of the first kind or an integral transform.

Definition 2.1. The integral transform of a function f(x) in the interval $a \le x \le b$ denoted by $\mathcal{I}{f(x)} = F(k)$ is [44]

$$\mathcal{I}\{f(x)\} = F(k) = \int_a^b K(x,k)f(x)dx, \qquad (2.2)$$

where K(x,k) is called the kernel of the transform and \mathcal{I} is the transform operator.

The integral transform for a function $f(\mathbf{x})$ with several variables is

$$\mathcal{I}\{f(\mathbf{x})\} = F(\mathbf{k}) = \int_D K(\mathbf{x}, \mathbf{k}) f(\mathbf{x}) d\mathbf{x}, \qquad (2.3)$$

where $\mathbf{x} = (x_1, x_2, \dots, x_n)$, $\mathbf{k} = (k_1, k_2, \dots, k_n)$ and $D \in \mathbb{R}^n$. Apparently \mathcal{I} is a linear operator since

$$\mathcal{I}\{\alpha f(x) + \beta g(x)\} = \int_{a}^{b} \{\alpha f(x) + \beta g(x)\} K(x,k) dx$$
(2.4)

$$= \alpha \mathcal{I}\{f(x)\} + \beta \mathcal{I}\{g(x)\}.$$
(2.5)

Let $f \in L^1(\mathbb{R})$, the Fourier transform of f(x) in $L^1(\mathbb{R})$ is a linear transform which has eigenfunctions obeying

$$\mathcal{F}[\psi] = \lambda \psi, \tag{2.6}$$

with $\lambda \in \mathbb{C}$. Specifying the Fourier transform between position and momentum spaces, we define the transform as an integral of the form

$$\mathcal{F}_x[f(x)](k) = F(k) = \int_{-\infty}^{\infty} f(x) \ e^{-ikx} dx.$$
(2.7)

where the kernel of the Fourier transform (in our convention) is e^{-ikx} . Clearly, $|\int_{-\infty}^{\infty} e^{-ikx} f(x) dx| \leq \int_{-\infty}^{\infty} |f(x)| dx$. This implies that the integral 2.7 exist for all $k \in \mathbb{R}$. One interesting feature of the Fourier transform is the generalized Parseval relation. Let $f, g \in L^2(\mathbb{R})$, then

$$\langle f|g\rangle = \int_{-\infty}^{\infty} f\overline{g} \, dx = \frac{1}{2\pi} \int_{-\infty}^{\infty} F\overline{G} \, dk = \frac{1}{2\pi} \langle F|G\rangle \tag{2.8}$$

Furthermore, if we define the convolution f * g of two functions to be the integral

$$f * g(x) = \int_{-\infty}^{\infty} f(x - y)g(y)dy.$$
 (2.9)

Then, the Fourier transform is

$$\mathcal{F}[f * g(x)] = \mathcal{F}[f]\mathcal{F}[g]. \tag{2.10}$$

In other words, the Fourier transform of a convolution of two functions f, g is the product of their Fourier transforms. Other useful identities are

$$\mathcal{F}[fg] = \mathcal{F}[f] * \mathcal{F}[g], \qquad (2.11)$$

$$\mathcal{F}^{-1}[\mathcal{F}(f)\mathcal{F}(g)] = f * g, \qquad (2.12)$$

$$\mathcal{F}^{-1}[\mathcal{F}(f) * \mathcal{F}(g)] = fg. \tag{2.13}$$

It is not difficult to realize that Fourier transform can be viewed as a special case of the bilateral Laplace transform with imaginary argument,

$$F(k) = \mathcal{F}_x[f(x)] = \mathcal{L}\{f(x)\}|_{s=ik}$$

$$(2.14)$$

where $\mathcal{L}{f(x)} = \int_{-\infty}^{\infty} e^{-sx} f(x) dx$. The definition 2.2 of integral transforms holds for quantum mechanical systems where the arguments of these transforms are in principle functions of canonical variables $F(\hat{q}_i, \hat{p}_i)$ satisfying in general the canonical commutation relation $[\hat{q}_i, \hat{p}_j] = i\hbar \delta_{ij}$ (in our case, we simply chose these canonical variables to be the position and momentum operators x and p defined in the continuous Hilbert space).

The Fourier transform is a four-periodic operator and has an intimate connection with the parity operator $\mathcal{P}f(x) = f(-x)$ since

$$\mathcal{F}^0 = \mathrm{id},\tag{2.15}$$

$$\mathcal{F}^1 = \mathcal{F}, \tag{2.16}$$

$$\mathcal{F}^2 = \mathcal{P},\tag{2.17}$$

$$\mathcal{F}^3 = \mathcal{F}^{-1} = \mathcal{F} \circ \mathcal{P} = \mathcal{P} \circ \mathcal{F}, \qquad (2.18)$$

$$\mathcal{F}^4 = \mathrm{id},\tag{2.19}$$

where id is the identity operator. From the above relations, we find that applying twice the Fourier transform is identical to the action of the parity operator.

A Hamiltonian is called pseudo-Hermitian if there exists a Hermitian operator η such that [9]

$$H^{\dagger} = \eta H \eta \tag{2.20}$$

The operator η is sometimes referred to as the metric operator. In this case, the Hermitian indefinite inner product $\langle \langle | \rangle \rangle_{\eta}$ is defined by [9]

$$\langle \langle \psi_1 | \psi_2 \rangle \rangle_{\eta} := \langle \psi_1 | \eta | \psi_2 \rangle \tag{2.21}$$

which is invariant under the time translation generated by the Hamiltonian H if and only if H is η -pseudo Hermitian. Obviously, when η is the identity operator, i.e., $\eta = \text{id}$, the condition 2.20 reduces to the Hermitian case. Since the parity operator is Hermitian, it can be used as a metric operator. In this case, the condition 2.20 becomes

$$H^{\dagger} = \mathcal{P}H\mathcal{P} \tag{2.22}$$

Hence, in terms of the Fourier operator, it becomes

$$H^{\dagger} = \mathcal{F}^2 H \mathcal{F}^2 \tag{2.23}$$

The next step is to derive the expression for the Fourier transform of a general \mathcal{PT} -symmetric Hamiltonian. The exponential Fourier transform \mathcal{F}

of the operator $\hat{f}(x)$ is [45, 46]

$$\mathcal{F}_x[\hat{f}(x)] = \hat{F}(k) = \int_{-\infty}^{\infty} \hat{f}(x)e^{-ik\cdot x}dx \qquad (2.24)$$

and the inverse Fourier transform is defined by

$$\mathcal{F}_{k}^{-1}[\hat{F}(k)] = \hat{f(x)} = 2\pi \int_{-\infty}^{\infty} \hat{F}(k)e^{ikx}dk$$
(2.25)

where x, k are operators in the continuous Hilbert space. Under parity (spatial reflection) $p \to -p, x \to -x$ and under time-reversal $p \to -p, x \to x$ and $i \to -i$, the general \mathcal{PT} -symmetric Hamiltonian (in units $\hbar = 2m = 1$ for simplicity) is

$$H = p^{2} + V_{\rm Re}(x) + iV_{\rm Im}(x)$$
(2.26)

where the real part of the potential $V_{\text{R}e}$ is even and the imaginary part $V_{\text{I}m}$ is odd under parity transformation. Note that both $V_{\text{R}e}$, $V_{\text{I}m} \in \mathbb{R}$. We define their exponential Fourier transforms as

$$\tilde{V}_{\mathrm{R}e}(p) = 2 \int_0^\infty V_{\mathrm{R}e}(x) \cos(p \cdot x) \, dx, \qquad (2.27)$$

$$\tilde{V}_{\mathrm{I}m}(p) = -2i \int_0^\infty V_{\mathrm{I}m}(x) \sin(p \cdot x) \, dx. \tag{2.28}$$

We plug 3.1,3.2 in the Schrödinger equation

$$\tilde{H}\phi(p) = p^{2}\phi(p) + 2\pi \int_{-\infty}^{\infty} \tilde{V}(p-p')\phi(p')dp',$$
(2.29)

where

$$\tilde{V}(p) = 2 \int_0^\infty \left(V_{\rm Re}(x) \cos(x \cdot p) + V_{\rm Im}(x) \sin(x \cdot p) \right) dx.$$
(2.30)

Interestingly, performing the exponential Fourier transform of the Hamiltonian 2.26 removes the imaginary unit *i* in front of $V_{\rm Im}$ and thus the Hamiltonian becomes apparently Hermitian in the momentum-space representation. Therefore, the non-Hermitian \mathcal{PT} -symmetric Hamiltonian becomes Hermitian after performing the exponential Fourier transform, as seen in 2.29. Due to the unitarity of the Fourier transform, one could easily prove that in the case of broken \mathcal{PT} -symmetry, complex pairs appear in the Hamiltonian written in the momentum space, and thus the discrete symmetries of a given Hamiltonian are unaffected by unitary transformations as expected. In the above we have considered Hamiltonian with one-dimensional position spaces. However, the generalization to higher-dimensional spaces is straightforward.

Next, we need to investigate the effect of the exponential Fourier transform on the orthonormal basis attached to a quantum system . Consider, for example, the Hermite polynomials ,

$$H_n(x) = (-)^n e^{x^2} \left(\frac{d}{dx}\right)^n (e^{-x^2})$$
(2.31)

which appears in many quantum mechanical problems, such as quantum harmonic oscillator. The Fourier transform of Hermite polynomials is [44, 47]

$$\mathcal{F}\{e^{-x^2/2}H_n(x)\} = (-i)^n e^{-p^2/2}H_n(p)$$
(2.32)

It is worthy to mention that the imaginary coefficient that appears in the front of the second-hand side $(-i)^n$ has no effect on the measured energy eigenvalues. To prove this we compute the Segal-Bargmann transform of $H_n(p)$. Let f(p) be a function in the momentum Hilbert space $L^2(\mathbb{R}^d)$, we define the Segal-Bargmann transform with respect to p as [48, 49, 50, 51, 52]

$$\mathcal{B}(f(z)) = \int_{\mathbb{R}^d} e^{-(z \cdot z - 2\sqrt{2}z \cdot p - p \cdot p)/2} f(p) \, dp. \tag{2.33}$$

The Segal-Bargmann transform of Hermite polynomials H_n are simply complex monomials of the form $\{\frac{z^n}{\sqrt{n!}}\}$. These monomials form an orthonormal basis since

$$\frac{1}{\pi} \int_{\mathbb{C}} dz \; e^{-|z|^2} \; \overline{z}^n z^m = n! \; \delta_{mn}, \tag{2.34}$$

where z is arbitrary complex coordinate (physically, it is the phase-space coordinate z = x + ip). As we emphasized in [53], both $\frac{z^n}{\sqrt{n!}}$ and $\frac{(-iz)^n}{\sqrt{n!}}$ correspond to the same energy eigenvalue for each level characterized by the same quantum number n and the effect of $(-i)^n$ in front of Hermite polynomials has no effect on the measured energy eigenvalues.

Definition 2.2. The Bargmann space, also known as the Fock-Bargmann or Segal-Bargmann space, denoted by $\mathcal{H}L^2(\mathbb{C}^n,\mu)$, is the space of holomorphic functions with Gaussian integral measure $\mu(z) = (\pi)^{-n} e^{-|z|^2}$ and $|z|^2 = |z_1|^2 + \cdots + |z_n|^2$. Any entire analytic function f(z) in this space satisfy a square-integrability condition of the type [51]

$$||f||^{2} := \langle f|f\rangle_{\mu} = (\pi)^{-n} \int_{\mathbb{C}^{n}} |f(z)|^{2} e^{-|z|^{2}} dz < \infty, \qquad (2.35)$$

where dz is the 2*n*-dimensional Lebesgue measure on $\mathbb{C}^n = \mathbb{R}^{2n}$.

Remark: The inner-product between analytic functions f(z) and g(z) in the Bargmann space $\mathcal{H}L^{2}(\mathbb{C}^{n},\mu)$ is

$$\langle f|g\rangle_{\mu} = (\pi)^{-n} \int_{\mathbb{C}^n} \overline{f}(z) g(z) e^{-|z|^2} dz.$$
(2.36)

Definition 2.3. The Segal-Bargmann transform is an integral transformation from $L^2(\mathbb{R}^n)$ to $\mathcal{H}L^2(\mathbb{C}^n,\mu)$ defined as

$$\mathcal{B}(f(z)) = \int_{\mathbb{R}^n} K_n(z, x) f(x) d^n x \quad \text{for } z \in \mathbb{C}^n \text{ and } f(x) \in \mathbb{R}^n \qquad (2.37)$$

with $K_n(z,x) = e^{(z \cdot z - 2\sqrt{2}z \cdot x - x \cdot x)/2}$ the kernel integral.

Theorem 2.1 (Unitarity of the Segal-Bargmann transform). The Segal-Bargmann transform defined by 2.37 is a unitary mapping between the Hilbert space $L^2(\mathbb{R}^n)$ and $\mathcal{H}L^2(\mathbb{C}^n,\mu)$.

In Bargmann space $\mathcal{H}L^2(\mathbb{C},\mu)$, the creation and annihilation operators of the harmonic oscillator are z and $\frac{\partial}{\partial z}$ since

$$\left[\frac{\partial}{\partial z}, z\right] f(z) = \frac{\partial}{\partial z} (zf(z)) - z\frac{\partial f(z)}{\partial z} = 1$$
(2.38)

similar to the canonical commutation relation $[\hat{a}, \hat{a}^{\dagger}] = 1$. In this representation, the harmonic oscillator Hamiltonian operator is $H = \hbar \omega \left(z \frac{d}{dz} + \frac{1}{2} \right)$. Applying to energy eigenstates (Fock states), we find

$$H|n\rangle = \hbar\omega \left(z\frac{d}{dz} + \frac{1}{2}\right)\frac{z^n}{\sqrt{n!}} = \hbar\omega \left(n + \frac{1}{2}\right)|n\rangle.$$
(2.39)

3 Non-Hermitian Spin Chains

For a given entire analytic function $f_{\alpha,\beta}(z,w) = \frac{z^{\alpha}}{\sqrt{\alpha!}} \frac{w^{\beta}}{\sqrt{\beta!}}$, we define the spin operators in the two-dimensional Bargmann space $\mathcal{H}L^2(\mathbb{C}^2,\mu)$ using

Bargmann representation of the Jordan-Schwinger map as [53]

$$S_x = \frac{\hbar}{2} \left(z \frac{\partial}{\partial w} + w \frac{\partial}{\partial z} \right), \tag{3.1}$$

$$S_y = \frac{\hbar}{2i} \left(z \frac{\partial}{\partial w} - w \frac{\partial}{\partial z} \right), \tag{3.2}$$

$$S_z = \frac{\hbar}{2} \left(z \frac{\partial}{\partial z} - w \frac{\partial}{\partial w} \right). \tag{3.3}$$

These operators belong to the SU(2) Lie algebra and obey the commutation relations $[S_i, S_j] = i\hbar \varepsilon_{ijk}S_k$ since the only non-trivial commutators between z, w and their partial derivatives are $\left[\frac{\partial}{\partial z}, z\right] = \left[\frac{\partial}{\partial w}, w\right] = 1$. The total number operator is

$$N = n_z + n_w, ag{3.4}$$

where $n_z = z \frac{\partial}{\partial z}$ and $n_w = w \frac{\partial}{\partial w}$.

The functions $\{f_{\alpha,\beta}\}$ form an orthonormal basis in the two-dimensional Bargmann space since

$$\langle f_{\alpha',\beta'}(z,w)|f_{\alpha,\beta}(z,w)\rangle_{\mu}$$
(3.5)
= $\int dz dw \exp[-|z|^2 - |w|^2] f_{\alpha',\beta'}(z,w) f_{\alpha,\beta}(z,w)$
= $\pi^2 \alpha! \beta! \, \delta_{\alpha',\alpha} \delta_{\beta',\beta}$

where $\alpha, \beta \in \mathbb{N}$ is the set of all natural numbers including zero i.e. $\{0, 1, 2, ...\}$.

One advantage of Bargmann representation of spin operators appears clearly in the study of spin chains. Consider the non-Hermitian XX spin chain Hamiltonian [54],

$$\hat{H}_s = \frac{1}{2} \sum_{j=1}^{J-1} \left[S_j^x S_{j+1}^x + S_j^y S_{j+1}^y + ig \left(S_j^z - S_{j+1}^z \right) \right], \tag{3.6}$$

where for small values of the coupling constant g, it becomes spectral equivalence to the Hermitian Hamiltonian [54].

In Bargmann representation, 3.6 can be written analytically as

$$H_{s} = \frac{\hbar^{2}}{4} \sum_{j=1}^{J-1} \left(z_{j+1} w_{j} \frac{\partial^{2}}{\partial z_{j} \partial w_{j+1}} + z_{j} w_{j+1} \frac{\partial^{2}}{\partial w_{j} \partial z_{j+1}} \right)$$

$$+ \frac{i\hbar g}{4} \sum_{j=1}^{J-1} \left(z_{j} \frac{\partial}{\partial z_{j}} - z_{j+1} \frac{\partial}{\partial z_{j+1}} + w_{j+1} \frac{\partial}{\partial w_{j+1}} - w_{j} \frac{\partial}{\partial w_{j}} \right).$$

$$(3.7)$$

Since the Segal-Bargmann transform is unitary, the time evolution for spin chain Hamiltonians is the same both in operator and analytical form [53, 55].

4 The Swanson Hamiltonian

The Swanson Hamiltonian is [56, 57, 58, 59],

$$H(\omega, \alpha, \beta) = \hbar \omega \left(\hat{a}^{\dagger} \hat{a} + \frac{1}{2} \right) + \hbar \alpha \ \hat{a}^2 + \hbar \beta \ \hat{a}^{\dagger 2}, \tag{4.1}$$

where α and β are real unequal constants in general. When $\alpha = \beta$, 4.1 reduces to the squeezed harmonic oscillator, and the energy spectrum becomes real.

Eq.4.1 can be written in terms of position and momentum operators by plugging the exact form of creation and annihilation operators,

$$\hat{a} = \frac{1}{\sqrt{2}} \left(\frac{x}{\xi_0} + i \frac{\xi_0}{\hbar} p \right), \tag{4.2}$$

$$\hat{a}^{\dagger} = \frac{1}{\sqrt{2}} \left(\frac{x}{\xi_0} - i \frac{\xi_0}{\hbar} p \right), \qquad (4.3)$$

with ξ_0 being the characteristic length scale of the system. We obtain

$$H(\omega,\alpha,\beta) = \frac{1}{2}\hbar\left(\omega+\alpha+\beta\right)\left(\frac{x}{\xi_0}\right)^2 + \frac{1}{2}\hbar\left(\omega-\alpha-\beta\right)\left(\frac{\xi_0p}{\hbar}\right)^2 + \frac{1}{2}\hbar\left(\alpha-\beta\right)\left(\frac{2i}{\hbar}xp+1\right)(4.4)$$

It is convenient in our analysis to make the Swanson-harmonic Hamiltonian mapping. This can be done with proper definitions of the position and momentum operators. Following [58], we have

$$P = \left(p + i\hbar \frac{\alpha - \beta}{(\omega - \alpha - \beta)\xi_0^2} x \right), \tag{4.5}$$

$$X = x. \tag{4.6}$$

It is straightforward to verify the commutation relation $[X, P] = i\hbar$.

In terms of X and P, equation 4.4 becomes

$$H = \frac{1}{2m}P^2 + \frac{k}{2}X^2 \tag{4.7}$$

where

$$m = \frac{\hbar}{(\omega - \alpha - \beta)\xi_0^2} \in \mathbb{R}, \tag{4.8}$$

$$k = m\Omega^2 = m\sqrt{\omega^2 - 4\alpha\beta} \in \mathbb{C}.$$
(4.9)

As shown by Swanson in [56], 4.1 posses real and positive energy eigenvalues whenever the condition $\omega^2 - 4\alpha\beta \ge 0$ ($\Omega^2 \ge 0$) is satisfied. Indeed, the limiting case when $\Omega = 0$ and $\omega - \alpha - \beta \ne 0$ corresponds to a free particle[58]. The creation and annihilation operators are

$$\hat{A} = \sqrt{\frac{m\Omega}{2\hbar}} \left(X + \frac{i}{m\Omega} P \right), \qquad (4.10)$$

$$\hat{A}^{\dagger} = \sqrt{\frac{m\Omega}{2\hbar}} \left(X - \frac{i}{m\Omega} P \right) \tag{4.11}$$

Since Ω is complex in general, it can be written in the form $\Omega = |\Omega|e^{i\theta}$. Thus, 4.10 and 4.11 become

$$\hat{A} = \sqrt{\frac{m|\Omega|}{2\hbar}} \left(X e^{i\theta/2} + \frac{i}{m|\Omega|} P e^{-i\theta/2} \right), \qquad (4.12)$$

$$\hat{A}^{\dagger} = \sqrt{\frac{m|\Omega|}{2\hbar}} \left(X e^{-i\theta/2} - \frac{i}{m|\Omega|} P e^{i\theta/2} \right), \tag{4.13}$$

Obviously $[\hat{A}, \hat{A}^{\dagger}] = 1$. By virtue of Euler's formula, i.e., $e^{i\theta} = \cos \theta + i \sin \theta$, 4.12 and 4.13 becomes in matrix form

$$\begin{pmatrix} \hat{A} \\ \hat{A}^{\dagger} \end{pmatrix} = \begin{pmatrix} \cos\frac{\theta}{2} & i\sin\frac{\theta}{2} \\ -i\sin\frac{\theta}{2} & \cos\frac{\theta}{2} \end{pmatrix} \begin{pmatrix} \hat{b} \\ \hat{b}^{\dagger} \end{pmatrix} = \mathbf{M} \cdot \vec{\mathbf{b}}$$
(4.14)

where $\det(\mathbf{M}) = \cos \theta$. When $\theta = 2\pi N$ where N is integer, $\det(\mathbf{M}) = 1$ and this corresponds to unitary transformation, the energy spectrum in this case is real. The operators \hat{b} and \hat{b}^{\dagger} are the usual creation and annihilation

operators for a harmonic oscillator with frequency $|\Omega|$ and generalized momentum P. Next, consider $|\tilde{n}\rangle$ to be the energy eigenvectors for 4.7 and $|n\rangle$ to be the energy eigenvectors for a harmonic oscillator with frequency $|\Omega|$, then we have

$$\hat{A}|\tilde{n}\rangle = \sqrt{\tilde{n}}|\tilde{n}-1\rangle = \cos\frac{\theta}{2}\hat{b}|n\rangle + i\sin\frac{\theta}{2}\hat{b}^{\dagger}|n\rangle \qquad (4.15)$$
$$= \cos\frac{\theta}{2}\sqrt{n}|n-1\rangle + i\sin\frac{\theta}{2}\sqrt{n+1}|n+1\rangle$$
$$\hat{A}^{\dagger}|\tilde{n}\rangle = \sqrt{\tilde{n}+1}|\tilde{n}+1\rangle = \cos\frac{\theta}{2}\hat{b}^{\dagger}|n\rangle - i\sin\frac{\theta}{2}\hat{b}|n\rangle \qquad (4.16)$$
$$= \cos\frac{\theta}{2}\sqrt{n+1}|n+1\rangle - i\sin\frac{\theta}{2}\sqrt{n}|n-1\rangle$$

From these results we notice that when $\sin \frac{\theta}{2} = 0$, the Hamiltonian 4.7 assumes real energy eigenvalues. The potential term in 4.7 can acquire an imaginary value and thus break the \mathcal{PT} -symmetry. Its exponential Fourier transform has the form

$$\tilde{V}(P) = \frac{m|\Omega|^2}{2} e^{2i\theta} \int_{-\infty}^{\infty} X^2 e^{-iP \cdot X} dX = -\pi m |\Omega|^2 e^{2i\theta} \delta''(P)$$
(4.17)

where δ'' is the second-derivative of the Dirac delta function. The non-zero phase θ breaks the \mathcal{PT} -symmetry in the Swanson model. When $\theta = 0$, the energy eigenvalues of the Swanson Hamiltonian are real and the \mathcal{PT} -symmetry is exact.

The creation and annihilation operators 4.12, 4.13 can be written in terms of the operators \hat{a} and \hat{a}^{\dagger} as

$$\hat{A} = \frac{1}{2} \sqrt{\frac{m|\Omega|}{\hbar}} \left(\left(s\xi_0 + \frac{\hbar e^{-i\theta/2}}{m|\Omega|\xi_0} \right) \hat{a} + \left(s\xi_0 - \frac{\hbar e^{-i\theta/2}}{m|\Omega|\xi_0} \right) \hat{a}^{\dagger} \right), \quad (4.18)$$

$$\hat{A}^{\dagger} = \frac{1}{2} \sqrt{\frac{m|\Omega|}{\hbar}} \left(\left(s\xi_0 + \frac{\hbar e^{i\theta/2}}{m|\Omega|\xi_0} \right) \hat{a}^{\dagger} + \left(s\xi_0 - \frac{\hbar e^{i\theta/2}}{m|\Omega|\xi_0} \right) \hat{a} \right)$$
(4.19)

where $s = \left(e^{i\theta/2} - \frac{\hbar(\alpha-\beta)e^{-i\theta/2}}{m|\Omega|(\omega-\alpha-\beta)\xi_0^2}\right)$.

The operators \hat{A} and \hat{A}^{\dagger} belong to pseudo-boson annihilation and creation operators introduced in [60]. Using Bargmann representation, we may

write \hat{A} and \hat{A}^{\dagger} respectively, as

$$\frac{\partial}{\partial w} = \frac{1}{2} \sqrt{\frac{m|\Omega|}{\hbar}} \left(\left(s\xi_0 + \frac{\hbar e^{-i\theta/2}}{m|\Omega|\xi_0} \right) \frac{\partial}{\partial z} + \left(s\xi_0 - \frac{\hbar e^{-i\theta/2}}{m|\Omega|\xi_0} \right) z \right), \quad (4.20)$$
$$w = \frac{1}{2} \sqrt{\frac{m|\Omega|}{\hbar}} \left(\left(s\xi_0 + \frac{\hbar e^{i\theta/2}}{m|\Omega|\xi_0} \right) z + \left(s\xi_0 - \frac{\hbar e^{i\theta/2}}{m|\Omega|\xi_0} \right) \frac{\partial}{\partial z} \right), \quad (4.21)$$

with $\left[\frac{\partial}{\partial w}, w\right] = 1$. The eigenfunctions of the Swanson model are monomials of the phase-coordinate w i.e. $\frac{w^n}{\sqrt{n!}}$. While the eigenfunctions associated with operators \hat{a} and \hat{a}^{\dagger} are $\frac{z^n}{\sqrt{n!}}$ which by construction are the ordinary harmonic oscillator energy states.

5 Concluding Remarks

We have investigated the behavior of \mathcal{PT} -symmetric Hamiltonians under integral transforms. The fact that twice the Fourier transform serves as a parity operator is used in the inner products of pseudo-Hermitian Hamiltonians. We discussed the effect of applying the exponential Fourier transform to a non-Hermitian \mathcal{PT} -symmetric Hamiltonian and investigated the resulting eigenfunctions in the momentum space. More interestingly, we applied another type of integral transform called the Segal-Bargmann transform to write the Hamiltonian in terms of the phase-space coordinates starting from the Hamiltonians in the position space. This was applied to the case of non-Hermitian XX spin chains which is spectral equivalent to their Hermitian counterparts in the weak coupling regime. As a model example, we studied the Swanson model in detail from integral transforms point of view. One possible application of the Fourier transforms in the context of \mathcal{PT} -symmetric Hamiltonians can be found in the study of double well delta function potentials [61]

$$V(x) = (1 + i\Gamma) \,\delta\left(x + a\right) + (1 - i\Gamma) \,\delta\left(x - a\right) \tag{5.1}$$

which has applications in the study of \mathcal{PT} -symmetric Bose-Einstein condensates.

Acknowledgment. We are grateful to Professor H.F. Jones for useful email correspondence. We thank the anonymous reviewer(s) for their comments and suggestions. We acknowledge the support from USIM.

References

- C. M. Bender and S. Boettcher, Real Spectra in Non- Hermitian Hamiltonians Having PT Symmetry, Phys.Rev.Lett. 80(24) (1998), 5243-5246, https://doi.org/10.1103/PhysRevLett.80.5243
- [2] C. M. Bender, S. Boettcher, and P. N. Meisinger, PTsymmetric quantum mechanics, J. Math. Phys. 40, (1999), 2201, https://doi.org/10.1063/1.532860
- [3] P. Dorey, C. Dunning and R. Tateo, Spectral equivalences, Bethe Ansatz equations, and reality properties in PT-symmetric quantum mechanics, J. Phys. A 34 (2001), L391, https://doi.org/10.1088/0305-4470/34/28/305
- [4] Z. Ahmed, Real and complex discrete eigenvalues in an exactly solvable one-dimensional complex PT-invariant potential, Phys. Let. A 282(6), (2001), 343-348, https://doi.org/10.1016/S0375-9601(01)00218-3
- [5] C. M. Bender, D. C. Brody, H. F. Jones, Complex Extension of Quantum Mechanics, Phys. Rev. Lett. 89, 270401(2002);Erratum Phys. Rev. Lett. 119902 **92**, (2004),https://doi.org/10.1103/PhysRevLett.89.270401
- [6] C.F.M. Faria, A. Fring, Time evolution of non-Hermitian Hamiltonian systems, J. Phys. A **39**, (2006), 9269-9289, https://doi.org/10.1088/0305-4470/39/29/018
- [7] R. Ramirez, M. Reboiro, Dynamics of finite dimensional non-hermitian systems with indefinite metric, J. Math. Phys. 60, (2019), 012106, https://doi.org/10.1063/1.5075628
- [8] C. M. Bender, Making sense of non-Hermitian Hamiltonians, Rept. Prog. Phys. 70, (2007), 947, https://doi.org/10.1088/0034-4885/70/6/R03
- [9] A. Mostafazadeh, Pseudo-Hermitian representation of quantum mechanics, Int. J. Geom. Meth. Mod. Phys. 7, (2010), 1191-1306, https://doi.org/10.1142/S0219887810004816
- [10] C. M. Bender, PT-symmetry: In Quantum And Classical Physics, World Scientific Publishing (2018).

- [11] Z. Η. Musslimani, Κ. G. Makris, R. El-Ganainy, and D. N. Christodoulides. Optical Solitons in PT-Periodic Potentials, Phys. Rev. Lett. 100, (2008),030402 https://doi.org/10.1103/PhysRevLett.100.030402
- [12] S. Klaiman, U. Günther, and N. Moiseyev, Visualization of Branch Points in *PT*-Symmetric Waveguides, Phys. Rev. Lett. **101**, (2008), 080402, https://doi.org/10.1103/PhysRevLett.101.080402
- [13] S. Longhi, Bloch Oscillations in Complex Crystals with PT-Symmetry, Phys. Rev. Lett. 103, (2009),123601, https://doi.org/10.1103/PhysRevLett.103.123601
- [14] S. Longhi, PT-symmetric laser-absorber, Phys. Rev. A 82, (2010), 031801(R), https://doi.org/10.1103/PhysRevA.82.031801
- [15] Rüter, C., Makris, K., El-Ganainy, R. et al. Observation of parity-time symmetry in optics, Nature Phys 6, (2010), 192–195, https://doi.org/10.1038/nphys1515
- [16] B. Midya, R. Roychoudhury, Nonlinear localized modes in PTsymmetric Rosen-Morse potential well, Phys. Rev. A 87, (2013), 045803, https://doi.org/10.1103/PhysRevA.87.045803
- M. Gianfreda, N. Hatano, [17] S. Garmon, Bound states, scattering states and resonant states in**PT-symmetric** open quantum systems. Phys. Rev. Α **92**. (2015).022125,https://doi.org/10.1103/PhysRevA.92.022125
- [18] K. Shobe, K.Kuramoto, K-I. Imura, N.Hatano, Non-Hermitian Fabry-Perot Resonances in a PT-symmetric system, Phys. Rev. Research 3, (2021), 013223, https://doi.org/10.1103/PhysRevResearch.3.013223
- [19] M. Znojil, F. Cannata, B. Bagchi, and R. Roychoudhury, Supersymmetry without hermiticity, Physics Letters B 483, 284, (2000), https://doi.org/10.1016/S0370-2693(00)00569-4
- [20] M. W. AlMasri, Supersymmetry of PT-symmetric tridiagonal Hamiltonians, Mod. Phys. Lett. A 36 (35), (2021), 2150242, https://doi.org/10.1142/S0217732321502424
- [21] M. Ezawa, Non-Hermitian boundary and interface states in nonreciprocal higher-order topological metals and electrical circuits, Phys. Rev. B 99, (2019), 121411(R), https://doi.org/10.1103/PhysRevB.99.121411

- [22] M. Ezawa, Electric circuits for non-Hermitian Chern insulators, Phys. Rev. B 100, (2019), 081401, https://doi.org/10.1103/PhysRevB.100.081401
- [23] M. Ezawa, Electric-circuit simulation of the Schrödinger equation and non-Hermitian quantum walks, Phys. Rev. B 100, (2019), 165419, https://doi.org/10.1103/PhysRevB.100.165419
- [24] C. Hang, W. Li, and G. Huang, Nonlinear light diffraction by electromagnetically induced gratings with PT-symmetry in a Rydberg atomic gas, Phys. Rev. A 100, (2019), 043807, https://doi.org/10.1103/PhysRevA.100.043807
- [25] Xg. Wang, Gh. Guo, and J. Berakdar, Steering magnonic dynamics and permeability at exceptional points in a parity-time symmetric waveguide, Nature Communications 11, (2020), 5663, https://doi.org/10.1038/s41467-020-19431-3
- [26] J. Von Neumann, Mathematical Foundations of Quantum Mechanics, Princeton University Press (1955).
- [27] M. Lieber, Quantum mechanics in momentum space: An illustration, American Journal of Physics 43, (1975), 486, https://doi.org/10.1119/1.9774
- [28] J. P. Dahl and M. Springborg, The Morse oscillator in position space, momentum space, and phase space, J. Chem. Phys. 88, (1988), 4535, https://doi.org/10.1063/1.453761
- [29] M. I. Samar, V. M. Tkachuk, Exactly solvable problems in the momentum space with a minimum uncertainty in position, J. Math. Phys. 57,(2016), 042102, https://doi.org/10.1063/1.4945313
- [30] M. I. Samar, V. M. Tkachuk, One-dimensional Coulomb-like problem in general case of deformed space with minimal length, J. Math. Phys. 57, (2016), 082108, https://doi.org/10.1063/1.4961320
- [31] J. Gamboa, M. Loewe, and J.C. Rojas, Non-Commutative Quantum Mechanics, Phys.Rev. D 64 (2001), 067901, https://doi.org/10.1103/PhysRevD.64.067901
- [32] R. J. Szabo, Quantum Field Theory on Noncommutative Spaces, Phys. Rept. **378**, (2003), 207-299, https://doi.org/10.1016/S0370-1573(03)00059-0

- [33] M. W. AlMasri, Axial-Anomaly in Noncommutative QED and Pauli-Villars Regularization, int. J. Mod. Phys. A 34 (26), (2019), 1950150, https://doi.org/10.1142/S0217751X19501501
- [34] M. W. AlMasri, М. R. В. Wahiddin, Deformed Kac-Moody Algebra and Noncommutative Fermi Theory inTwo-Dimensions, Int. J. Mod. Phys. A 36 (26) (2021), 2150189, https://doi.org/10.1142/S0217751X2150189X
- [35] J. Gamboa, M. Loewe, F. Mendez, J. C. Rojas, The Landau problem and noncommutative quantum mechanics, Mod. Phys. Lett. A 16, (2001), 2075-2078, https://doi.org/10.1142/S0217732301005345
- [36] P. A.Horvathy, The non-commutative Landau problem, Ann.Phys. 299, (2002), 128-140, https://doi.org/10.1006/aphy.2002.6271
- [37] M. Demetrian, D. Kochan, Quantum mechanics on noncommutative plane, Acta Physica Slovaca 52 (1), (2002), pp. 1-9, https://doi.org/10.48550/arXiv.hep-th/0102050
- [38] R. J. Szabo, Magnetic Backgrounds and Noncommutative Field Theory". Int.J.Mod.Phys. A 19 (2004) 1837-1862, https://doi.org/10.1142/S0217751X04018099
- Κ. Li, [39] S. Dulat. Landau Problem in Noncommutative Quantum Mechanics, Chin. Phys. С **32**, (2008),92-95. https://doi.org/10.1088/1674-1137/32/2/003
- [40] H. Ishizuka, N. Nagaosa, Noncommutative quantum mechanics and skew scattering in ferromagnetic metals, Phys. Rev. B 96, (2017) 165202, https://doi.org/10.1103/PhysRevB.96.165202
- [41] A. N. Bogdanov, and C. Panagopoulos, Physical foundations and basic properties of magnetic skyrmions, Nature Reviews Physics 2, (2020) pages 492–498, https://doi.org/10.1038/s42254-020-0203-7.
- [42] M. W. AlMasri, SU(4) description of bilayer skyrmion-antiskyrmion pairs, EPL **129**, (2020), 10005, https://doi.org/10.1209/0295-5075/129/10005
- [43] F. R. Lux, F. Freimuth, S. Blugel, and Y. Mokrousov, The chiral Hall effect of magnetic skyrmions from a cyclic cohomology approach, Phys. Rev. Lett. **124** ,(2020), 096602 , https://doi.org/10.1103/PhysRevLett.124.096602

- [44] L. Debnath and D.Bhatta, Integral Transforms and Their Applications, Third edition, Taylor and Francis LLC (2015).
- [45] H. Bateman, and A. Erdelyi. Tables of Integral Transforms: Vols. 1 and 2. McGraw-Hill Book Co., New York, (1954).
- [46] T. W. Korner, Fourier Analysis, Cambridge University Press (1989).
- [47] N. Wiener, The Fourier Integral and Certain of Its Applications, Cambridge University Press (1933).
- [48] V. Bargmann, On a Hilbert space of analytic functions and an associated integral transform, Communications on Pure and Applied Mathematics, 14 (3): (1961), 187, https://doi.org/10.1002/cpa.3160140303
- [49] I. E. Segal, Mathematical problems of relativistic physics, in Kac, M. (ed.), Proceedings of the Summer Seminar, Boulder, Colorado, 1960, Vol. II, Lectures in Applied Mathematics, American Mathematical Society (1963).
- [50] A. M. Perelomov, Generalized Coherent States and Their Applications, Springer, Berlin (1986).
- [51] B. C. Hall, Holomorphic Methods in Mathematical Physics, Contemporary Mathematics, 260, (2000), pp. 1-59.
- [52] M. W. AlMasri, M. R. B. Wahiddin, Bargmann representation of quantum absorption refrigerators, Rep. Math. Phys. 89(2), (2022), Pages 185-198, https://doi.org/10.1016/S0034-4877(22)00022-2
- [53] M. W. AlMasri, M. R. B. Wahiddin. "Bargmann representation of spin chains", (2021), arXiv:2108.10626.
- [54] C. Korff, PT Symmetry of the non-Hermitian XX Spin-Chain: Nonlocal Bulk Interaction from Complex Boundary Fields, J. Phys. A 41 (2008) 295206, https://doi.org/10.1088/1751-8113/41/29/295206
- [55] M. W. AlMasri, M. R. B. Wahiddin, Quantum decomposition algorithm for master equations of stochastic processes: The damped spin case, Mod. Phys. Lett. A 37 (2022) 32, 2250216, https://doi.org/10.1142/S0217732322502169
- [56] M. S. Swanson. Transition elements for a non-Hermitian Phys. quadratic Hamiltonian, J. Math. $\mathbf{45}$ (2004),585, https://doi.org/10.1063/1.1640796

- [57] H. F. Jones, On Pseudo-Hermitian Hamiltonians and Their Hermitian Counterparts, J. Phys. A: Math. Gen. 38 (2005), 1741–1746, https://doi.org/10.1088/0305-4470/38/8/010
- [58] V. Fernández, R. Ramírez, M. Reboiro, Swanson Hamiltonian: non-PT-symmetry phase, J. Phys. A: Math. Theor. 55, 015303 , (2022), https://doi.org/10.1088/1751-8121/ac3a35
- [59] F. Bagarello, A Swanson-like Hamiltonian and the inverted harmonic oscillator, J. Phys. A: Math. Theor. 55 (2022) 225204, https://doi.org/10.1088/1751-8121/ac6a92
- [60] D. A. Trifonov, Pseudo-Boson Coherent and Fock States, in Differential Geometry, Complex Analysis and Mathematical Physics, eds. K. Sekigawa et al (W. Scientific 2009), pp. 241-250.
- [61] H. Cartarius, G. Wunner, Model of a PT symmetric Bose-Einstein condensate in a delta-functions double well, Phys. Rev. A 86, (2012), 013612, https://doi.org/10.1103/PhysRevA.86.013612