arXiv:2207.03319v3 [quant-ph] 6 Dec 2022

Topological Speed Limit

Tan Van Vu^{*} and Keiji Saito[†]

Department of Physics, Keio University, 3-14-1 Hiyoshi, Kohoku-ku, Yokohama 223-8522, Japan

(Dated: December 7, 2022)

Any physical system evolves at a finite speed that is constrained not only by the energetic cost but also by the topological structure of the underlying dynamics. In this Letter, by considering such structural information, we derive a unified topological speed limit for the evolution of physical states using an optimal transport approach. We prove that the minimum time required for changing states is lower bounded by the discrete Wasserstein distance, which encodes the topological information of the system, and the time-averaged velocity. The bound obtained is tight and applicable to a wide range of dynamics, from deterministic to stochastic, and classical to quantum systems. In addition, the bound provides insight into the design principles of the optimal process that attains the maximum speed. We demonstrate the application of our results to chemical reaction networks and interacting many-body quantum systems.

Introduction.—Investigating how fast a system can evolve is one of the central problems in classical and quantum mechanics. In a seminal work by Mandelstam and Tamm [1], a fundamental bound on the operational time required for the transformation between two orthogonal states for unitary dynamics was derived. Since then, generalizations of the bound for arbitrary states and nonunitary dynamics have been intensively studied [2–36], leading to the notion of speed limits (see Ref. [37] for a review). These speed limits establish the ultimate rate at which a system can evolve to a distinguishable state and have found diverse applications, for example, in quantum control [38–41], quantum metrology [42, 43], and thermodynamics of computation [33, 44–49].

Interacting systems generally form topological structures in their dynamics, such as chemical reaction networks that consist of several species (see the schematic in Fig. 1). In general, a state represented by a vector \boldsymbol{x}_t evolves over time and is significantly affected by the topology of the dynamics. For instance, a Markov jump process with dense connectivity may relax toward an equilibrium state faster than one with sparse connectivity. A many-body system with long-range interactions can change quantum states faster than one with shortrange interactions [50]. Although speed limits for state transformations have been intensively investigated, the topological nature arising from the network structure in the dynamics has not been fully accounted for. Note that conventional speed limits, which read $\tau \geq \mathcal{L}(\boldsymbol{x}_0, \boldsymbol{x}_{\tau})/\overline{\boldsymbol{v}}$, employed non-topological metrics \mathcal{L} , such as the Bures angle, trace norm, quantum Fisher information, etc., to quantify the distance between the initial and final states [37]. These metrics are always upper bounded by a constant that does not scale with the size of the system, whereas the dynamics strongly depends on the system size. Velocity \overline{v} is determined by the entire dynamics of the system [51], and hence it is generally of the order of system size. Consequently, conventional speed limits become trivial (i.e., $\tau \geq \mathcal{L}(\boldsymbol{x}_0, \boldsymbol{x}_{\tau})/\overline{v} \to 0$) as the system increases in terms of size [52]. This indicates that in or-




FIG. 1. (a) Generic time evolution of a physical state $\boldsymbol{x}_t = [x_1(t), \ldots, x_N(t)]^{\mathsf{T}}$ on a graph. $x_i(t)$ is evolved because of the flows $\{f_{ij}(t)\}$ exchanged between neighboring vertices and an external flow $f_i(t)$. (b) Examples include reactant concentrations in deterministic chemical reactions and boson numbers in interacting bosonic systems.

der to derive meaningful bounds, metrics that capture the topological nature and are scalable with system size should be considered.

In this Letter, we derive a speed limit for arbitrary states x_t using a topological metric defined through the network structure in the dynamics. The time evolution of such states is described by a graph in which each vertex exchanges flows with each other and may be pumped by an external flow. Examples include the probability distribution of discrete systems, mass concentrations in chemical reaction networks, and vectors of observables in quantum systems (see Fig. 1 again for illustration). We employ a generalized version of the discrete Wasserstein distance to quantify the distance between the states. This distance, widely used in optimal transport theory [53], encodes topological information and can grow proportional to the size of the system. We prove that the minimum operational time required to change the physical state is lower bounded by the Wasserstein distance divided by the average velocity [cf. Eqs. (5) and (6)]. The obtained speed limit is tight and can be saturated, even when the system size increases. Moreover, it is applicable to a broad range of dynamics ranging from deterministic and stochastic classical systems to isolated and open quantum systems. For example, we apply the theory to chemical reaction networks using the Wasserstein distance applicable to any reversible chemical reaction and provide a reaction speed formula that can discriminate between different chemical reactions [54]. Another important application is the interacting bosonic transport for arbitrary initial (mixed) states with and without a thermal environment, which is relevant to the Lieb–Robinson velocity [55]. Through the examples, we demonstrate that considering topological metrics does not only provide quantitatively tight bounds but also qualitatively reveals the physical mechanism of state transformations, which cannot be obtained with speed limits reported thus far.

General setup.—We consider a time-dependent vector state $\boldsymbol{x}_t \coloneqq [x_1(t), \ldots, x_N(t)]^{\mathsf{T}}$ and an undirected graph $G(\mathcal{V}, \mathcal{E})$ with the vertex set $\mathcal{V} = \{1, \ldots, N\}$ and edge set \mathcal{E} . Each element $x_i(t)$ corresponds to a vertex $i \in \mathcal{V}$. For example, \boldsymbol{x}_t can be a vector of the probability distribution of a discrete system, reactant concentrations of chemical reaction networks, or physical observables in classical and quantum systems (examples are provided later). For each vertex i, let $\mathcal{B}_i \coloneqq \{j \mid \langle i, j \rangle \in \mathcal{E}\}$ denote the set of neighboring vertices of i. We assume that the time evolution of \boldsymbol{x}_t is given by the following deterministic equation [see Fig. 1(a)]:

$$\dot{x}_i(t) = f_i(t) + \sum_{j \in \mathcal{B}_i} f_{ij}(t), \qquad (1)$$

where $f_{ij}(t) = -f_{ji}(t)$ denotes the flow exchange between vertices *i* and *j* for $i \neq j$ and $f_i(t)$ is an arbitrary external flow. In the absence of external flows [i.e., $f_i(t) = 0$ for all *i*], $\sum_{i=1}^{N} x_i(t)$ is invariant. Examples of Eq. (1) include the master equation of Markov jump processes, rate equation of chemical reaction networks, and time evolution of the observables in quantum systems. We define a time-dependent velocity [56], which is the sum of the absolute values of the external and exchanged flows, given by

$$\upsilon_{t,\lambda} \coloneqq \lambda \sum_{i} |f_i(t)| + \sum_{\langle i,j \rangle \in \mathcal{E}} |f_{ij}(t)|, \qquad (2)$$

where $\lambda \geq 0$ is a weighting factor, and the second summation is over all unordered pairs $\langle i, j \rangle \in \mathcal{E}$. For simplicity, we denote $v_{t,0}$ by v_t . We also define the Manhattan norm for an arbitrary vector \boldsymbol{x} as $\|\boldsymbol{x}\|_1 \coloneqq \sum_i |x_i|$ and the time average of an arbitrary time-dependent quantity w_t as $\langle w_t \rangle_{\mathcal{I}} \coloneqq \tau^{-1} \int_0^{\tau} w_t dt$.

Wasserstein distance.—Here we introduce the discrete L^1 -Wasserstein distance between two states \boldsymbol{x} and \boldsymbol{y} on the graph $G(\mathcal{V}, \mathcal{E})$. First, we consider the case in which \boldsymbol{x} and \boldsymbol{y} are balanced (that is, $\sum_i x_i = \sum_i y_i$), and then we generalize the distance to the unbalanced case (that is, $\sum_i x_i \neq \sum_i y_i$). Let d_{ij} denote the shortest path distance

between the vertices i and j in the graph. In other words, d_{ij} is the minimum length of paths connecting i and j. Graph G is assumed to be connected [57]; therefore, d_{ij} is always finite. Suppose that we have a transport plan that redistributes \boldsymbol{x} to \boldsymbol{y} by sending an amount of π_{ij} from x_j to y_i with a cost of d_{ij} per unit weight for all ordered pairs $\langle i, j \rangle$. The Wasserstein distance is then defined as the minimum transport cost for all feasible plans, given by

$$\mathcal{W}_1(\boldsymbol{x}, \boldsymbol{y}) \coloneqq \min_{\pi \in \Pi(\boldsymbol{x}, \boldsymbol{y})} \sum_{i, j} d_{ij} \pi_{ij}.$$
 (3)

Here, $\Pi(\boldsymbol{x}, \boldsymbol{y})$ denotes the set of all transport plans $\pi = [\pi_{ij}] \in \mathbb{R}_{\geq 0}^{N \times N}$ that satisfy $\sum_j \pi_{ij} = y_i$ and $\sum_j \pi_{ji} = x_i$. Previous studies have shown that the Wasserstein distance plays a crucial role in statistics and machine learning [58], computer vision [59], linguistics [60], molecular biology [61], and stochastic thermodynamics [33, 62–64].

Next, we describe the generalized Wasserstein distance for the unbalanced case. Transport between two unbalanced states can be enabled by allowing *add* and *remove* operations in addition to transportation between vertices. More precisely, an infinitesimal mass $\delta \boldsymbol{x}$ of \boldsymbol{x} can either be removed at cost $\lambda \|\delta \boldsymbol{x}\|_1$ or moved from \boldsymbol{x} to \boldsymbol{y} at cost $\mathcal{W}_1(\delta \boldsymbol{x}, \delta \boldsymbol{y})$. Mathematically, the generalized Wasserstein distance between unbalanced states can be defined as [65]

$$\mathcal{W}_{1,\lambda}(\boldsymbol{x},\boldsymbol{y}) \coloneqq \min \left\{ \lambda (\|\boldsymbol{x} - \tilde{\boldsymbol{x}}\|_1 + \|\boldsymbol{y} - \tilde{\boldsymbol{y}}\|_1) + \mathcal{W}_1(\tilde{\boldsymbol{x}}, \tilde{\boldsymbol{y}}) \right\},$$
(4)

where the minimum is over all the states $\tilde{\boldsymbol{x}}$ and $\tilde{\boldsymbol{y}}$ such that $\|\tilde{\boldsymbol{x}}\|_1 = \|\tilde{\boldsymbol{y}}\|_1$. By definition (4), distance $\mathcal{W}_{1,\lambda}$ always satisfies the triangle inequality [65]. If \boldsymbol{x} and \boldsymbol{y} are balanced states, then $\mathcal{W}_{1,\lambda}$ is reduced to \mathcal{W}_1 within the $\lambda \to +\infty$ limit. We also note that $\mathcal{W}_{1,\lambda}$ can be calculated numerically using the linear programming method [66].

Main results.—We now utilize the generalized Wasserstein distance (4) to derive a topological speed limit for any state \boldsymbol{x}_t obeying the general dynamics (1). Specifically, we prove that the minimum time required to transform \boldsymbol{x}_0 into \boldsymbol{x}_{τ} is lower bounded by the Wasserstein distance divided by the average velocity:

$$\tau \ge \frac{\mathcal{W}_{1,\lambda}(\boldsymbol{x}_0, \boldsymbol{x}_{\tau})}{\langle v_{t,\lambda} \rangle_{\tau}}, \ \forall \lambda \ge 0.$$
(5)

In the case that the external flows are absent [i.e., $f_i(t) = 0$], inequality (5) can be reduced to a simple bound by taking the $\lambda \to +\infty$ limit, which reads

$$\tau \ge \frac{\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_{\tau})}{\langle v_t \rangle_{\tau}}.$$
(6)

The inequalities (5) and (6) are our main results; the proof is postponed to the end of the Letter.

These results have several physically critical properties. (i) First, these bounds can be derived as long as the time evolution of x_t is described by Eq. (1), which is a general setting for both the classical and quantum cases. Notably, the bounds can be saturated if the time evolution (1) realizes an optimal transport plan. (ii) Second, our bounds utilize topological information about the system dynamics to provide a stringent constraint on the speed of changing states. Topological information is encoded into the Wasserstein distance, and this distance term can be as large as the order of the system's size. (iii) Third, by further upper bounding the time-averaged velocity $\langle v_{t,\lambda} \rangle_{\tau}$ by relevant quantities, such as the thermodynamic and kinetic costs, we can derive more interpretable bounds, which clarify the physical mechanism of the speed of state transformations. (iv) Finally, the speed limit for an arbitrary scalar observable defined in terms of state x_t can also be obtained as a consequence of Eq. (5) [66].

In the following, we illustrate the above remarks, especially (i)-(iii), through two applications to classical and quantum systems (see Ref. [66] for further applications in isolated and Markovian open quantum systems, measurement-induced quantum walk [67], and quantum communication [68, 69]).

Application 1: Chemical reaction networks.—We consider a chemical reaction system composed of several chemical species X_i $(i \in S)$ that interact through reversible elementary reaction channels $\rho \in \mathcal{R}$. Here, Sand \mathcal{R} denote the set of indices of the species and reaction channels, respectively. Each reaction channel is represented as

$$\sum_{i} \nu_{i}^{+\rho} X_{i} \stackrel{\kappa^{+\rho}}{\underset{\kappa^{-\rho}}{\rightleftharpoons}} \sum_{i} \nu_{i}^{-\rho} X_{i}, \tag{7}$$

where $+\rho$ and $-\rho$ correspond to the forward and backward reactions, respectively, $\{\kappa^{\pm\rho}\}$ are the macroscopic reaction rates, and $\{\nu_i^{\pm\rho}\}$ are the stoichiometric coefficients. Let \boldsymbol{x}_t denote the vector of the mass concentrations of species. The molar concentration \boldsymbol{c}_t can be related as $c_i(t) = x_i(t)/m_i$, where m_i denotes the molar mass of species X_i . The time evolution of \boldsymbol{x}_t can be described by the deterministic rate equation:

$$\dot{x}_i(t) = \sum_{\rho} m_i (\nu_i^{+\rho} - \nu_i^{-\rho}) J_t^{\rho}, \qquad (8)$$

where $J_t^{\rho} \coloneqq J_t^{-\rho} - J_t^{+\rho}$ is the net reaction current and $J_t^{\pm\rho} \coloneqq \kappa^{\pm\rho} \prod_i c_i(t)^{\nu_i^{\pm\rho}}$ are the reaction fluxes.

Next, we derive the speed limits for the system in terms of the Wasserstein distance defined on graph G. For simplicity, here we consider closed reaction networks, in which the total mass concentration is conserved [70]. The generalization for open reaction networks, wherein the total mass conservation may be violated, is presented in Ref. [66]. The total mass conservation law implies $\sum_{i} m_i (\nu_i^{+\rho} - \nu_i^{-\rho}) = 0$ for any ρ . Due to these conditions, there always exist matrices $Z^{\rho} = [z_{ij}^{\rho}]$ such that the rate

equation (8) can be expressed in the form of Eq. (1) with $f_{ij}(t) = \sum_{\rho} z_{ij}^{\rho} J_t^{\rho}$ and $f_i(t) = 0$ [66]. The graph *G* can be obtained by adding an undirected edge $\langle i, j \rangle$ to \mathcal{E} for any $z_{ij}^{\rho} \neq 0$. After some simple manipulations [66], we can prove that

$$\upsilon_t \le \sum_{\rho} \nu^{\rho} |J_t^{\rho}|,\tag{9}$$

where $\nu^{\rho} \coloneqq (1/2) \sum_{i} m_{i} |\nu_{i}^{+\rho} - \nu_{i}^{-\rho}|$. Combining Eqs. (6) and (9) yields the following speed limit:

$$\tau \ge \frac{\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_\tau)}{\left\langle \sum_{\rho} \nu^{\rho} | J_t^{\rho} | \right\rangle_{\tau}} \eqqcolon \tau_1.$$
(10)

Equation (10) implies that the operational time is lower bounded by the Wasserstein distance and the net reaction currents.

A thermodynamic speed limit can also be obtained using Eq. (10). The entropy production rate of a chemical reaction system can be defined as [71]

$$\sigma_t \coloneqq \sum_{\rho} J_t^{\rho} \ln \frac{J_t^{-\rho}}{J_t^{+\rho}},\tag{11}$$

where the gas constant is set to unity. We define the following kinetic quantity:

$$\ell_t \coloneqq \sum_{\rho} (\nu^{\rho})^2 \frac{J_t^{-\rho} - J_t^{+\rho}}{\ln(J_t^{-\rho}/J_t^{+\rho})},\tag{12}$$

which is the sum of the microscopic Onsager coefficients [33, 72]. Applying the Cauchy–Schwarz inequality, we prove that $\left\langle \sum_{\rho} \nu^{\rho} | J_t^{\rho} | \right\rangle_{\tau} \leq \left\langle \sqrt{\sigma_t \ell_t} \right\rangle_{\tau} \leq \sqrt{\langle \sigma_t \rangle_{\tau} \langle \ell_t \rangle_{\tau}}$. Consequently, we obtain the following thermodynamic speed limit:

$$\tau \ge \frac{\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_\tau)}{\sqrt{\langle \sigma_t \rangle_\tau \langle \ell_t \rangle_\tau}} \approx \tau_2.$$
(13)




FIG. 2. Numerical demonstration of the speed limits in the cascade reaction network with N = 10. The operational time τ , topological bounds τ_1 and τ_2 , and non-topological bound τ_3 are depicted by solid, dashed and dash-dotted, and dotted lines, respectively. The parameters are set to $k_f = 2$ and $k_b = 1$. The initial mass concentration is $\boldsymbol{x}_0 = [1, 0.9, \dots, 0.1]^{\mathsf{T}}$.

Inequality (13) implies that the minimum time required to transform \boldsymbol{x}_0 into \boldsymbol{x}_{τ} is determined by the product of the thermodynamic and kinetic costs.

We numerically demonstrate the derived bounds in a cascade reaction network with $|\mathcal{S}| = 10$ species and $|\mathcal{R}| = 9$ reaction channels (see Fig. 2). We also compare the results with a non-topological bound reported in Ref. [73], which reads $\tau \geq \tau_3 \coloneqq \mathcal{T}(\mathbf{c}_0, \mathbf{c}_\tau)/\sqrt{\langle \sigma_t \rangle_\tau} \langle d_t \rangle_\tau$. Here, \mathcal{T} denotes the total variation distance and $d_t \coloneqq$ $(|\mathcal{S}|/8) \sum_{\rho,i} (\nu_i^{+\rho} - \nu_i^{-\rho})^2 (J_t^{+\rho} + J_t^{-\rho})$ corresponds to the diffusion coefficient. We calculate and plot the lower bounds $\tau_i (1 \leq i \leq 3)$ in Fig. 2. As shown, the topological speed limits $\tau \geq \tau_1 \geq \tau_2$ are tight; especially, the bound $\tau \geq \tau_1$ is always saturated. On the contrary, the non-topological bound $\tau \geq \tau_3$ is loose and does not provide a meaningful bound for the speed of the system.

Application 2: Interacting bosonic systems.—Next, we describe an application for quantum many-body bosonic systems. We consider a model of bosons that hop on an arbitrary finite-dimensional lattice and interact with each other. Let Λ denote the set of all the sites in the lattice. The Hamiltonian can be expressed in the following generic form:

$$H_t \coloneqq -\gamma \sum_{\langle i,j \rangle} (b_i^{\dagger} b_j + b_j^{\dagger} b_i) + \sum_{Z \subseteq \Lambda} h_Z.$$
(14)

Here, the first summation is over neighboring lattice sites (which can be arbitrarily distant), $\gamma > 0$ describes the boson mobility, b_i and b_i^{\dagger} are the bosonic creation and annihilation operators for site *i*, respectively, $\hat{n}_i \coloneqq b_i^{\dagger} b_i$ is the number operator, and h_Z is an arbitrary function of $\{\hat{n}_i\}_{i\in Z}$. Examples include the Bose–Hubbard model, given by $\sum_{Z \subseteq \Lambda} h_Z = (U/2) \sum_i \hat{n}_i (\hat{n}_i - 1) - \mu \sum_i \hat{n}_i$, where U and μ are real constants. Note that the graph $G(\mathcal{V}, \mathcal{E})$ of the bosonic system is identical to the lattice topology (i.e., \mathcal{V} is the set of sites and \mathcal{E} is the set of edges that connect the two neighboring sites). The maximum vertex degree of the graph is denoted by d_G .

We assume that the bosonic system is weakly coupled to a Markovian thermal reservoir and can exchange particles with the reservoir, where the time evolution of the reduced density matrix is described by the Lindblad equation [74]:

$$\dot{\varrho}_t = -i[H_t, \varrho_t] + \sum_{i \in \Lambda} \left(\mathcal{D}[L_{i,+}] + \mathcal{D}[L_{i,-}] \right) \varrho_t, \qquad (15)$$

where $\mathcal{D}[L]\varrho \coloneqq L\varrho L^{\dagger} - (1/2)\{L^{\dagger}L, \varrho\}$ is the dissipator, $L_{i,+} = \sqrt{\gamma_{i,+}}b_i^{\dagger}$ and $L_{i,-} = \sqrt{\gamma_{i,-}}b_i$ are the jump operators that characterize the absorption and emission of bosons at site *i*, respectively. Hereafter, we set $\hbar = 1$ for simplicity.

We consider the vector of boson numbers occupied at each site, $x_i(t) = \operatorname{tr}\{\hat{n}_i \varrho_t\}$, and define the instantaneous total number of bosons as $\mathcal{N}_t \coloneqq \sum_{i \in \Lambda} x_i(t)$. Using the relation $[b_i, \hat{n}_i] = b_i$, we can show that the time evolution of $x_i(t)$ can be expressed in the form of Eq. (1) with $f_i(t) = \operatorname{tr} \{L_{i,+}\varrho_t L_{i,+}^{\dagger}\} - \operatorname{tr} \{L_{i,-}\varrho_t L_{i,-}^{\dagger}\}$ and $f_{ij}(t) = 2\gamma \operatorname{Im} [\operatorname{tr} \{b_j^{\dagger} b_i \varrho_t\}]$. By inserting these terms into $v_{t,\lambda}$, we can immediately obtain the speed limit (5) for bosonic transport.

Next, we derive a more physically interpretable speed limit by upper bounding the velocity $v_{t,\lambda}$. To this end, we introduce two relevant physical quantities. The first is the irreversible entropy production rate [75], which is the sum of the entropic changes in the system and environment, defined as $\sigma_t \coloneqq \sigma_t^{\text{sys}} + \sigma_t^{\text{env}}$. Here, $\sigma_t^{\text{sys}} \coloneqq$ $- \operatorname{tr}\{\dot{\varrho}_t \ln \varrho_t\}$ is the rate of von Neumann entropy of the bosonic system, and σ_t^{env} quantifies the heat dissipated to the environment as follows:

$$\sigma_t^{\text{env}} \coloneqq \sum_i \left(\operatorname{tr} \left\{ L_{i,+} \varrho_t L_{i,+}^{\dagger} \right\} - \operatorname{tr} \left\{ L_{i,-} \varrho_t L_{i,-}^{\dagger} \right\} \right) \ln \frac{\gamma_{i,+}}{\gamma_{i,-}}, \quad (16)$$

where we have assumed the local detailed balance condition [that is, $\ln(\gamma_{i,+}/\gamma_{i,-})$ is related to the heat dissipation of the boson exchange at site *i*]. The second is quantum dynamical activity [76, 77], which quantifies the boson exchange frequency between the system and reservoir, given by

$$a_t \coloneqq \sum_i \left(\operatorname{tr} \left\{ L_{i,+} \varrho_t L_{i,+}^{\dagger} \right\} + \operatorname{tr} \left\{ L_{i,-} \varrho_t L_{i,-}^{\dagger} \right\} \right).$$
(17)

Using these quantities, we can prove that the velocity $v_{t,\lambda}$ is upper bounded as [66]

$$\upsilon_{t,\lambda} \le \gamma d_G \mathcal{N}_t + \lambda \frac{\sigma_t}{2} \Phi \left(\frac{\sigma_t}{2a_t}\right)^{-1}, \tag{18}$$

where $\Phi(x)$ is the inverse function of $x \tanh(x)$. By combining Eqs. (5) and (18), we obtain the following speed limit:

$$\tau \ge \frac{\mathcal{W}_{1,\lambda}(\boldsymbol{x}_0, \boldsymbol{x}_{\tau})}{\langle \gamma d_G \mathcal{N}_t + \lambda \sigma_t \Phi(\sigma_t/2a_t)^{-1}/2 \rangle_{\tau}}.$$
 (19)

Equation (19) implies that the speed of bosonic transport is lower bounded by the lattice topology, boson mobility, and dissipation. The bound also indicates that dissipative controls can help accelerate the bosonic transport. The inequality (19) is valid for arbitrary initial states of the bosonic system.

It is worthwhile discussing the vanishing coupling limit (i.e., the case where the system becomes isolated). In this case, $\sigma_t = a_t = 0$ and $\mathcal{N}_t = \mathcal{N}$ for all times. Defining the boson concentration $\bar{x}_i(t) \coloneqq \mathcal{N}^{-1}x_i(t)$, we obtain $\sum_i \bar{x}_i(t) = 1$. By taking the $\lambda \to +\infty$ limit, Eq. (19) is reduced to a simple speed limit for an isolated bosonic system:

$$\tau \ge \frac{\mathcal{W}_1(\bar{\boldsymbol{x}}_0, \bar{\boldsymbol{x}}_\tau)}{\gamma d_G}.$$
(20)

Bound (20) has a remarkable implication for bosonic transport. Assume that all bosons are initially concentrated in a region R_1 , and we want to transport all of

them to a distinct region R_2 within a finite time τ . In this case, $W_1(\bar{x}_0, \bar{x}_\tau) \geq \text{dist}(R_1, R_2)$, where $\text{dist}(R_1, R_2)$ denotes the length of the shortest path connecting the regions R_1 and R_2 . Therefore, Eq. (20) implies that transporting bosons always takes at least a time proportional to the distance between the two regions: $\tau \geq \text{dist}(R_1, R_2)/(\gamma d_G)$, which cannot be obtained with conventional speed limits. This statement holds for *arbitrary* initial states, including the pure states considered in Ref. [78]. While the Lieb–Robinson bounds [79–83] imply a linear light cone for the operator spreading, Eq. (20) provides a useful bound for the operational time required for bosonic transport.

Proof of Eq. (5).—We consider the time discretization of Eq. (1) with time interval $\delta t = \tau/K$. For each $k \in [0, K-1]$ and $t = k\delta t$, we have

$$x_i(t+\delta t) = x_i(t) + \delta t \left[f_i(t) + \sum_{j \in \mathcal{B}_i} f_{ij}(t) \right].$$
(21)

Equation (21) indicates that we can transform \boldsymbol{x}_t to $\boldsymbol{x}_{t+\delta t}$ by adding $f_i(t)\delta t$ to $x_i(t)$ with cost $\lambda |f_i(t)|\delta t$ and exchanging $f_{ij}(t)\delta t$ between neighboring vertices i and jwith cost $|f_{ij}(t)|\delta t$. Such the transport plan takes the total cost of

$$\left(\lambda \sum_{i} |f_{i}(t)| + \sum_{\langle i,j \rangle \in \mathcal{E}} |f_{ij}(t)|\right) \delta t = \upsilon_{t,\lambda} \delta t, \qquad (22)$$

which should be larger than or equal to $\mathcal{W}_{1,\lambda}(\boldsymbol{x}_t, \boldsymbol{x}_{t+\delta t})$. Therefore, taking the sum of Eq. (22) from k = 0 to k = K - 1 and applying the triangle inequality for $\mathcal{W}_{1,\lambda}$ yield

$$\sum_{k=0}^{K-1} \upsilon_{t,\lambda} \delta t \ge \mathcal{W}_{1,\lambda}(\boldsymbol{x}_0, \boldsymbol{x}_{\tau}).$$
(23)

By taking the $\delta t \to 0$ limit in Eq. (23), we obtain $\tau \langle v_{t,\lambda} \rangle_{\tau} \geq \mathcal{W}_{1,\lambda}(\boldsymbol{x}_0, \boldsymbol{x}_{\tau})$, from which Eq. (5) is immediately derived.

Conclusion.—In this Letter, we derived the topological speed limit for vector states that accounts for the network structure in the underlying dynamics [84]. The speed limit provides a tight bound for the operational time and insight into the system speed from a topological perspective. We showed that the bound is applicable to various dynamics as long as the time evolution of the physical state can be described in terms of a graph. Because our speed limit is derived in a general setting, we expect that it can be applied to obtain fundamental bounds for several other dynamics.

We thank Tomotaka Kuwahara, Andreas Dechant, and Koudai Sugimoto for fruitful discussions, Marius Lemm and Ryusuke Hamazaki for helpful communications, and Mai Dan Nguyen for the help in preparing the figures. We also appreciate anonymous referees for valuable comments. This work was supported by Grants-in-Aid for Scientific Research (JP19H05603 and JP19H05791).

- [†] saitoh@rk.phys.keio.ac.jp
- L. Mandelstam and I. Tamm, The uncertainty relation between energy and time in non-relativistic quantum mechanics, J. Phys. USSR 9, 249 (1945).
- [2] A. Uhlmann, An energy dispersion estimate, Phys. Lett. A 161, 329 (1992).
- [3] N. Margolus and L. B. Levitin, The maximum speed of dynamical evolution, Physica D 120, 188 (1998).
- [4] A. del Campo, I. L. Egusquiza, M. B. Plenio, and S. F. Huelga, Quantum speed limits in open system dynamics, Phys. Rev. Lett. **110**, 050403 (2013).
- [5] S. Deffner and E. Lutz, Quantum speed limit for non-Markovian dynamics, Phys. Rev. Lett. **111**, 010402 (2013).
- [6] M. M. Taddei, B. M. Escher, L. Davidovich, and R. L. de Matos Filho, Quantum speed limit for physical processes, Phys. Rev. Lett. **110**, 050402 (2013).
- [7] D. P. Pires, M. Cianciaruso, L. C. Céleri, G. Adesso, and D. O. Soares-Pinto, Generalized geometric quantum speed limits, Phys. Rev. X 6, 021031 (2016).
- [8] D. Mondal, C. Datta, and S. Sazim, Quantum coherence sets the quantum speed limit for mixed states, Phys. Lett. A 380, 689 (2016).
- [9] S. Deffner, Geometric quantum speed limits: a case for Wigner phase space, New J. Phys. 19, 103018 (2017).
- [10] B. Shanahan, A. Chenu, N. Margolus, and A. del Campo, Quantum speed limits across the quantum-to-classical transition, Phys. Rev. Lett. **120**, 070401 (2018).
- [11] M. Okuyama and M. Ohzeki, Quantum speed limit is not quantum, Phys. Rev. Lett. **120**, 070402 (2018).
- [12] S. Ito, Stochastic thermodynamic interpretation of information geometry, Phys. Rev. Lett. 121, 030605 (2018).
- [13] N. Shiraishi, K. Funo, and K. Saito, Speed limit for classical stochastic processes, Phys. Rev. Lett. **121**, 070601 (2018).
- [14] F. Campaioli, F. A. Pollock, F. C. Binder, and K. Modi, Tightening quantum speed limits for almost all states, Phys. Rev. Lett. **120**, 060409 (2018).
- [15] K. Funo, N. Shiraishi, and K. Saito, Speed limit for open quantum systems, New J. Phys. 21, 013006 (2019).
- [16] L. P. García-Pintos and A. del Campo, Quantum speed limits under continuous quantum measurements, New J. Phys. 21, 033012 (2019).
- [17] T. D. Kieu, A class of time-energy uncertainty relations for time-dependent Hamiltonians, Proc. R. Soc. A 475, 20190148 (2019).
- [18] X. Hu, S. Sun, and Y. Zheng, Quantum speed limit via the trajectory ensemble, Phys. Rev. A 101, 042107 (2020).
- [19] S. Ito and A. Dechant, Stochastic time evolution, information geometry, and the Cramér-Rao bound, Phys. Rev. X 10, 021056 (2020).
- [20] S. B. Nicholson, L. P. García-Pintos, A. del Campo, and J. R. Green, Time-information uncertainty relations in thermodynamics, Nat. Phys. 16, 1211 (2020).
- [21] V. T. Vo, T. Van Vu, and Y. Hasegawa, Unified approach to classical speed limit and thermodynamic uncertainty relation, Phys. Rev. E 102, 062132 (2020).
- [22] T. Fogarty, S. Deffner, T. Busch, and S. Campbell, Orthogonality catastrophe as a consequence of the quantum

speed limit, Phys. Rev. Lett. 124, 110601 (2020).

- [23] N. Il'in and O. Lychkovskiy, Quantum speed limit for thermal states, Phys. Rev. A 103, 062204 (2021).
- [24] S. Sun, Y. Peng, X. Hu, and Y. Zheng, Quantum speed limit quantified by the changing rate of phase, Phys. Rev. Lett. 127, 100404 (2021).
- [25] T. Van Vu and Y. Hasegawa, Geometrical bounds of the irreversibility in Markovian systems, Phys. Rev. Lett. 126, 010601 (2021).
- [26] E. O'Connor, G. Guarnieri, and S. Campbell, Action quantum speed limits, Phys. Rev. A 103, 022210 (2021).
- [27] N. Shiraishi and K. Saito, Speed limit for open systems coupled to general environments, Phys. Rev. Research 3, 023074 (2021).
- [28] K. Bolonek-Lasoń, J. Gonera, and P. Kosiński, Classical and quantum speed limits, Quantum 5, 482 (2021).
- [29] A. del Campo, Probing quantum speed limits with ultracold gases, Phys. Rev. Lett. **126**, 180603 (2021).
- [30] J.-C. Delvenne and G. Falasco, Tight classical speed limits, arXiv preprint arXiv:2110.13050 (2021).
- [31] R. Hamazaki, Speed limits for macroscopic transitions, PRX Quantum 3, 020319 (2022).
- [32] Y. Hasegawa, Thermodynamic bounds via bulkboundary correspondence: speed limit, thermodynamic uncertainty relation, and Heisenberg principle, arXiv preprint arXiv:2203.12421 (2022).
- [33] T. Van Vu and K. Saito, Thermodynamic unification of optimal transport: Thermodynamic uncertainty relation, minimum dissipation, and thermodynamic speed limits, arXiv preprint arXiv:2206.02684 (2022).
- [34] S. Nakajima and Y. Utsumi, Speed limits of the trace distance in the interaction picture for open quantum system, arXiv preprint arXiv:2204.02884 (2022).
- [35] D. S. P. Salazar, Lower bound for entropy production rate in stochastic systems far from equilibrium, arXiv preprint arXiv:2204.00875 (2022).
- [36] A. Kolchinsky, A. Dechant, K. Yoshimura, and S. Ito, Information geometry of excess and housekeeping entropy production, arXiv preprint arXiv:2206.14599 (2022).
- [37] S. Deffner and S. Campbell, Quantum speed limits: from Heisenberg's uncertainty principle to optimal quantum control, J. Phys. A 50, 453001 (2017).
- [38] T. Caneva, M. Murphy, T. Calarco, R. Fazio, S. Montangero, V. Giovannetti, and G. E. Santoro, Optimal control at the quantum speed limit, Phys. Rev. Lett. 103, 240501 (2009).
- [39] S. Deffner, Optimal control of a qubit in an optical cavity, J. Phys. B 47, 145502 (2014).
- [40] S. Campbell and S. Deffner, Trade-off between speed and cost in shortcuts to adiabaticity, Phys. Rev. Lett. 118, 100601 (2017).
- [41] K. Funo, J.-N. Zhang, C. Chatou, K. Kim, M. Ueda, and A. del Campo, Universal work fluctuations during shortcuts to adiabaticity by counterdiabatic driving, Phys. Rev. Lett. 118, 100602 (2017).
- [42] V. Giovannetti, S. Lloyd, and L. Maccone, Quantum metrology, Phys. Rev. Lett. 96, 010401 (2006).
- [43] M. Beau and A. del Campo, Nonlinear quantum metrology of many-body open systems, Phys. Rev. Lett. 119, 010403 (2017).
- [44] S. Lloyd, Ultimate physical limits to computation, Nature 406, 1047 (2000).
- [45] K. Proesmans, J. Ehrich, and J. Bechhoefer, Finite-time Landauer principle, Phys. Rev. Lett. 125, 100602 (2020).

- [46] S. Deffner, Energetic cost of Hamiltonian quantum gates, Europhys. Lett. 134, 40002 (2021).
- [47] Y.-Z. Zhen, D. Egloff, K. Modi, and O. Dahlsten, Universal bound on energy cost of bit reset in finite time, Phys. Rev. Lett. **127**, 190602 (2021).
- [48] T. Van Vu and K. Saito, Finite-time quantum Landauer principle and quantum coherence, Phys. Rev. Lett. 128, 010602 (2022).
- [49] J. S. Lee, S. Lee, H. Kwon, and H. Park, Speed limit for a highly irreversible process and tight finite-time Landauer's bound, arXiv preprint arXiv:2204.07388 (2022).
- [50] J. Eisert, M. van den Worm, S. R. Manmana, and M. Kastner, Breakdown of quasilocality in long-range quantum lattice models, Phys. Rev. Lett. **111**, 260401 (2013).
- [51] F. Campaioli, F. A. Pollock, and K. Modi, Tight, robust, and feasible quantum speed limits for open dynamics, Quantum 3, 168 (2019).
- [52] M. Bukov, D. Sels, and A. Polkovnikov, Geometric speed limit of accessible many-body state preparation, Phys. Rev. X 9, 011034 (2019).
- [53] C. Villani, Optimal Transport: Old and New (Springer, Berlin, Heidelberg, 2008).
- [54] We note the related work [72] which discusses the application of Wasserstein distance to chemical reactions. However, this study uses the system-dependent L^2 -Wasserstein distance [25], which is defined for a given fixed chemical reaction system. Therefore, the reaction speeds cannot be compared between different chemical reactions.
- [55] E. H. Lieb and D. W. Robinson, The finite group velocity of quantum spin systems, Commun.Math. Phys. 28, 251 (1972).
- [56] Note that this velocity, which contains the absolute value of currents, is qualitatively and quantitatively different from dynamical activity in the literature, which involves the absolute value of unidirectional fluxes [85].
- [57] A graph is said to be connected if there always exists a path from any vertex to any other vertex in the graph. In the case that the graph is not connected, we can define the Wasserstein distance as the sum of the distances defined for each connected component.
- [58] S. Kolouri, S. R. Park, M. Thorpe, D. Slepcev, and G. K. Rohde, Optimal mass transport: Signal processing and machine-learning applications, IEEE Signal Process. Mag. 34, 43 (2017).
- [59] S. Haker, L. Zhu, A. Tannenbaum, and S. Angenent, Optimal mass transport for registration and warping, Int. J. Comput. Vision 60, 225 (2004).
- [60] G. Huang, C. Guo, M. J. Kusner, Y. Sun, F. Sha, and K. Q. Weinberger, Supervised word mover's distance, in Advances in Neural Information Processing Systems, Vol. 29 (2016).
- [61] G. Schiebinger, J. Shu, M. Tabaka, B. Cleary, V. Subramanian, A. Solomon, J. Gould, S. Liu, S. Lin, P. Berube, *et al.*, Optimal-transport analysis of single-cell gene expression identifies developmental trajectories in reprogramming, Cell **176**, 928 (2019).
- [62] E. Aurell, C. Mejía-Monasterio, and P. Muratore-Ginanneschi, Optimal protocols and optimal transport in stochastic thermodynamics, Phys. Rev. Lett. 106, 250601 (2011).
- [63] M. Nakazato and S. Ito, Geometrical aspects of entropy production in stochastic thermodynamics based

on wasserstein distance, Phys. Rev. Research **3**, 043093 (2021).

- [64] A. Dechant, Minimum entropy production, detailed balance and Wasserstein distance for continuous-time Markov processes, J. Phys. A 55, 094001 (2022).
- [65] B. Piccoli and F. Rossi, Generalized Wasserstein distance and its application to transport equations with source, Arch. Ration. Mech. Anal. 211, 335 (2013).
- [66] See the Supplemental Material for details of analytical calculations and further applications, which includes Refs. [86, 87].
- [67] A. Didi and E. Barkai, Measurement-induced quantum walks, Phys. Rev. E 105, 054108 (2022).
- [68] S. Bose, Quantum communication through an unmodulated spin chain, Phys. Rev. Lett. 91, 207901 (2003).
- [69] M. Murphy, S. Montangero, V. Giovannetti, and T. Calarco, Communication at the quantum speed limit along a spin chain, Phys. Rev. A 82, 022318 (2010).
- [70] Note that the mass conservation may not be fulfilled in open reaction networks. For example, in the presence of catalysts [88], the total mass concentration may vary over time.
- [71] R. Rao and M. Esposito, Nonequilibrium thermodynamics of chemical reaction networks: Wisdom from stochastic thermodynamics, Phys. Rev. X 6, 041064 (2016).
- [72] K. Yoshimura, A. Kolchinsky, A. Dechant, and S. Ito, Geometrical approach to excess/housekeeping entropy production in discrete systems, arXiv preprint arXiv:2205.15227 (2022).
- [73] K. Yoshimura and S. Ito, Thermodynamic uncertainty relation and thermodynamic speed limit in deterministic chemical reaction networks, Phys. Rev. Lett. **127**, 160601 (2021).
- [74] G. Lindblad, On the generators of quantum dynamical semigroups, Commun. Math. Phys. 48, 119 (1976).
- [75] G. T. Landi and M. Paternostro, Irreversible entropy production: From classical to quantum, Rev. Mod. Phys. 93, 035008 (2021).
- [76] Y. Hasegawa, Quantum thermodynamic uncertainty re-

lation for continuous measurement, Phys. Rev. Lett. **125**, 050601 (2020).

- [77] T. Van Vu and K. Saito, Thermodynamics of precision in Markovian open quantum dynamics, Phys. Rev. Lett. 128, 140602 (2022).
- [78] J. Faupin, M. Lemm, and I. M. Sigal, Maximal speed for macroscopic particle transport in the Bose-Hubbard model, Phys. Rev. Lett. **128**, 150602 (2022).
- [79] T. Kuwahara and K. Saito, Lieb-Robinson bound and almost-linear light cone in interacting boson systems, Phys. Rev. Lett. **127**, 070403 (2021).
- [80] M. C. Tran, A. Y. Guo, C. L. Baldwin, A. Ehrenberg, A. V. Gorshkov, and A. Lucas, Lieb-Robinson light cone for power-law interactions, Phys. Rev. Lett. **127**, 160401 (2021).
- [81] J. Faupin, M. Lemm, and I. M. Sigal, On Lieb-Robinson bounds for the Bose-Hubbard model, Communications in Mathematical Physics (2022).
- [82] C. Yin and A. Lucas, Finite speed of quantum information in models of interacting bosons at finite density, Phys. Rev. X 12, 021039 (2022).
- [83] T. Kuwahara, T. Van Vu, and K. Saito, Optimal light cone and digital quantum simulation of interacting bosons, arXiv preprint arXiv:2206.14736 (2022).
- [84] We notice the recent work [31] that has a similar motivation. Note that we consider the speed limits for vector states, which cover a wide range of state transformations, whereas Ref. [31] studies the speed limit only for a fixed scalar observable.
- [85] C. Maes, Frenesy: Time-symmetric dynamical activity in nonequilibria, Phys. Rep. 850, 1 (2020).
- [86] M. Fishman, S. R. White, and E. M. Stoudenmire, The ITensor Software Library for Tensor Network Calculations, SciPost Phys. Codebases, 4 (2022).
- [87] M. Yang and S. R. White, Time-dependent variational principle with ancillary Krylov subspace, Phys. Rev. B 102, 094315 (2020).
- [88] J. Unterberger and P. Nghe, Stoechiometric and dynamical autocatalysis for diluted chemical reaction networks, J. Math. Biol. 85 (2022).

Supplemental Material for "Topological Speed Limit"

Tan Van Vu and Keiji Saito

Department of Physics, Keio University, 3-14-1 Hiyoshi, Kohoku-ku, Yokohama 223-8522, Japan

This Supplemental Material describes the details of the analytical calculations presented in the main text and further applications of the topological speed limit for isolated and open quantum systems, measurement-induced quantum walk, and quantum communication. The equations and figure numbers are prefixed with S [e.g., Eq. (S1) or Fig. S1]. The numbers without this prefix [e.g., Eq. (1) or Fig. 1] refer to the items in the main text.

CONTENTS

S1.	Numerical calculation of the generalized	
	Wasserstein distance	1
S2.	Speed limit for scalar observables	1
S3.	Chemical reaction networks	2
	A. Graph construction for closed reaction	~
	networks	2
	B. Derivation of Eq. (9)	2
	C. Generalization to generic reaction networks	2
S4.	Bosonic transport	4
	A. Time evolution of boson number	4
	B. Derivation of Eq. (18)	4
	C. Numerical demonstration	5
S5.	Isolated quantum systems	6
S6.	Measurement-induced quantum walk	7
S7.	Quantum communication using spin systems	8
S8.	Markovian open quantum systems	9
	References	10

S1. NUMERICAL CALCULATION OF THE GENERALIZED WASSERSTEIN DISTANCE

The Wasserstein distance can be calculated using the linear programming method. Mathematically, it can be formulated as the following minimization problem:

minimize
$$\operatorname{tr}\{\mathbf{C}\pi\}$$

subject to $\pi\mathbf{1} = \mathbf{y}, \ \pi^{\mathsf{T}}\mathbf{1} = \mathbf{x}$ (S1)
 $\pi \ge 0.$

where $C = [d_{mn}]$ is the matrix of transport cost and 1 is the all-ones vector.

The generalized Wasserstein distance can be calculated in a similar way. Note that it has been shown that the generalized Wasserstein distance can be achieved using only *remove* and *transport* operations (i.e., we do not need *add* operation) [1]. Therefore, $\mathcal{W}_{1,\lambda}$ can be expressed as

$$\mathcal{W}_{1,\lambda}(\boldsymbol{x},\boldsymbol{y}) = \min \left\{ \lambda (\|\boldsymbol{x} - \tilde{\boldsymbol{x}}\|_1 + \|\boldsymbol{y} - \tilde{\boldsymbol{y}}\|_1) + \mathcal{W}_1(\tilde{\boldsymbol{x}}, \tilde{\boldsymbol{y}}) \right\},$$
(S2)

where the minimum is over all states $\tilde{\boldsymbol{x}}$ and $\tilde{\boldsymbol{y}}$ such that $\tilde{\boldsymbol{x}} \leq \boldsymbol{x}, \ \tilde{\boldsymbol{y}} \leq \boldsymbol{y}$, and $\|\tilde{\boldsymbol{x}}\|_1 = \|\tilde{\boldsymbol{y}}\|_1$. Here, $\boldsymbol{x} \leq \boldsymbol{y}$ means that $x_n \leq y_n$ for all n. Consequently, computing $\mathcal{W}_{1,\lambda}$ is equivalent to solving the following minimization problem:

minimize
$$\operatorname{tr} \{ \mathbf{C} \pi \} + \lambda [\sum_{n} (x_{n} - \tilde{x}_{n}) + \sum_{n} (y_{n} - \tilde{y}_{n})]$$

subject to $\pi \mathbf{1} = \tilde{y}, \ \pi^{\mathsf{T}} \mathbf{1} = \tilde{x}$
 $\tilde{x} \leq x, \ \tilde{y} \leq y$
 $\pi \geq 0.$ (S3)

The linear programming problems in Eqs. (S1) and (S3) can be efficiently solved using programming languages such as Python, Julia, or Mathematica. A Mathematica code that computes the (generalized) Wasserstein distance can be found on GitHub [2].

S2. SPEED LIMIT FOR SCALAR OBSERVABLES

Here we derive a speed limit for a scalar observable \mathcal{O}_t , defined in terms of state \boldsymbol{x}_t as

$$\mathcal{O}_t \coloneqq \sum_i o_i x_i(t) = \boldsymbol{o}^{\mathsf{T}} \boldsymbol{x}_t, \qquad (S4)$$

where $\boldsymbol{o} = [o_1, \ldots, o_N]^{\mathsf{T}}$ is a vector of real coefficients. For convenience, we define the spectral norm and Lipschitz constant of vector \boldsymbol{o} as follows:

$$\|\boldsymbol{o}\|_{\infty} \coloneqq \max|o_i|,\tag{S5}$$

$$\|\boldsymbol{o}\|_{\mathrm{Lip}} \coloneqq \max_{\langle i,j \rangle \in \mathcal{E}} |o_i - o_j|.$$
(S6)

We first consider the general case where external flows are present. According to Prop. 1, we obtain the following speed limit for observable \mathcal{O}_t from Eq. (5):

$$\tau \ge \frac{|\mathcal{O}_{\tau} - \mathcal{O}_{0}|}{\max\left\{\lambda^{-1} \|\boldsymbol{o}\|_{\infty}, \|\boldsymbol{o}\|_{\operatorname{Lip}}\right\} \langle v_{t,\lambda} \rangle_{\tau}}.$$
 (S7)

For the case that the external flows are absent, we can obtain a more compact speed limit by taking the $\lambda \rightarrow +\infty$

limit in Eq. (S7), which reads

$$\tau \ge \frac{|\mathcal{O}_{\tau} - \mathcal{O}_0|}{\|\boldsymbol{o}\|_{\operatorname{Lip}} \langle \boldsymbol{v}_t \rangle_{\tau}}.$$
(S8)

This bound recovers the result reported in Ref. [3] [see Eq. (19) therein].

Proposition 1. For arbitrary states x, y and a real vector o, the following inequality holds:

$$|\boldsymbol{o}^{\mathsf{T}}(\boldsymbol{x}-\boldsymbol{y})| \le \max\left\{\lambda^{-1} \|\boldsymbol{o}\|_{\infty}, \|\boldsymbol{o}\|_{\mathrm{Lip}}\right\} \mathcal{W}_{1,\lambda}(\boldsymbol{x},\boldsymbol{y}).$$
 (S9)

Proof. Let \tilde{x} and \tilde{y} be two states that realize $\mathcal{W}_{1,\lambda}(x, y)$. In other words, we have

$$\mathcal{W}_{1,\lambda}(\boldsymbol{x},\boldsymbol{y}) = \lambda(\|\boldsymbol{x} - \tilde{\boldsymbol{x}}\|_1 + \|\boldsymbol{y} - \tilde{\boldsymbol{y}}\|_1) + \mathcal{W}_1(\tilde{\boldsymbol{x}}, \tilde{\boldsymbol{y}}).$$
(S10)

Then, by applying the triangle inequality and the Kantorovich–Rubinstein duality formula [4]

$$\mathcal{W}_1(\tilde{\boldsymbol{x}}, \tilde{\boldsymbol{y}}) = \max\left\{ \boldsymbol{\phi}^{\mathsf{T}}(\tilde{\boldsymbol{x}} - \tilde{\boldsymbol{y}}) \mid \|\boldsymbol{\phi}\|_{\mathrm{Lip}} \leq 1 \right\},$$
 (S11)

we can prove Eq. (S9) as follows:

$$\begin{aligned} \left| \boldsymbol{o}^{\mathsf{T}}(\boldsymbol{x} - \boldsymbol{y}) \right| \\ \leq \left| \boldsymbol{o}^{\mathsf{T}}(\boldsymbol{x} - \tilde{\boldsymbol{x}}) \right| + \left| \boldsymbol{o}^{\mathsf{T}}(\boldsymbol{y} - \tilde{\boldsymbol{y}}) \right| + \left| \boldsymbol{o}^{\mathsf{T}}(\tilde{\boldsymbol{x}} - \tilde{\boldsymbol{y}}) \right| \\ \leq \left\| \boldsymbol{o} \right\|_{\infty} \left(\left\| \boldsymbol{x} - \tilde{\boldsymbol{x}} \right\|_{1} + \left\| \boldsymbol{y} - \tilde{\boldsymbol{y}} \right\|_{1} \right) + \left\| \boldsymbol{o} \right\|_{\operatorname{Lip}} \mathcal{W}_{1}(\tilde{\boldsymbol{x}}, \tilde{\boldsymbol{y}}) \\ \leq \max \left\{ \lambda^{-1} \left\| \boldsymbol{o} \right\|_{\infty}, \left\| \boldsymbol{o} \right\|_{\operatorname{Lip}} \right\} \mathcal{W}_{1,\lambda}(\boldsymbol{x}, \boldsymbol{y}). \end{aligned}$$
(S12)

S3. CHEMICAL REACTION NETWORKS

A. Graph construction for closed reaction networks

Here we describe in detail the construction of an undirected graph $G(\mathcal{V}, \mathcal{E})$ for the closed chemical system from which the Wasserstein distance can be defined immediately. Notice the total mass conservation $\sum_i m_i (\nu_i^{+\rho} - \nu_i^{-\rho}) = 0$ for any ρ . The set of vertices is defined as $\mathcal{V} = \mathcal{S}$, where the vertex *i* corresponds to the species X_i . For each reaction channel ρ , define $\mathcal{S}^{\rho}_+ \coloneqq \{i | \nu_i^{+\rho} > \nu_i^{-\rho}\}$ and $\mathcal{S}^{\rho}_- \coloneqq \{i | \nu_i^{+\rho} < \nu_i^{-\rho}\}$. Evidently, $\sum_{i \in \mathcal{S}^{\rho}_+} m_i (\nu_i^{+\rho} - \nu_i^{-\rho}) = \sum_{i \in \mathcal{S}^{\rho}_-} m_i (\nu_i^{-\rho} - \nu_i^{+\rho})$. According to Prop. 2, there exists a matrix $\mathbb{Z}^{\rho} = [z_{ij}^{\rho}] \in \mathbb{R}^{|\mathcal{S}| \times |\mathcal{S}|}$ such that

$$\sum_{i \in \mathcal{S}_{-}^{\rho}} z_{ij}^{\rho} = m_i (\nu_i^{+\rho} - \nu_i^{-\rho}), \quad \forall i \in \mathcal{S}_{+}^{\rho},$$
(S13)

$$\sum_{i\in\mathcal{S}_{+}^{\rho}} z_{ij}^{\rho} = m_j (\nu_j^{-\rho} - \nu_j^{+\rho}), \ \forall j \in \mathcal{S}_{-}^{\rho}, \tag{S14}$$

$$z_{ji}^{\rho} = -z_{ij}^{\rho} \le 0, \quad \forall i \in \mathcal{S}_{+}^{\rho}, \quad j \in \mathcal{S}_{-}^{\rho}, \tag{S15}$$

$$z_{ij}^{\rho} = 0$$
, otherwise. (S16)

For each $z_{ij}^{\rho} \neq 0$, we add an undirected edge $\langle i, j \rangle$ to \mathcal{E} (see Fig. S1 for illustration). Note that the existence of




FIG. S1. Illustration of the process of adding edges to the graph G for a reaction channel ρ . Here, for simplicity, the molar mass is set to unity for all species (i.e., $m_i = 1$ for any i). For each $z_{ij}^{\rho} \neq 0$, an edge between i and j is added to G.

 Z^{ρ} may not be unique; nevertheless, an instance of Z^{ρ} can always be found such that at most $|\mathcal{S}^{\rho}_{+}| + |\mathcal{S}^{\rho}_{-}| - 1$ edges are added to the graph for each reaction channel ρ . Repeating this process for all $\rho \in \mathcal{R}$, we can readily obtain the graph G.

By this construction, we can verify that

$$\dot{x}_{i}(t) = \sum_{\rho} m_{i} (\nu_{i}^{+\rho} - \nu_{i}^{-\rho}) J_{t}^{\rho}$$
$$= \sum_{j \in \mathcal{B}_{i}} \sum_{\rho} z_{ij}^{\rho} J_{t}^{\rho}$$
$$= \sum_{j \in \mathcal{B}_{i}} f_{ij}(t), \qquad (S17)$$

where $f_{ij}(t) = \sum_{\rho} z_{ij}^{\rho} J_t^{\rho}$.

B. Derivation of Eq. (9)

Furthermore, by applying the triangle inequality, we can prove that the velocity is upper bounded as follows:

$$\begin{aligned}
\upsilon_t &= \sum_{\langle i,j \rangle \in \mathcal{E}} |f_{ij}(t)| \\
&= \frac{1}{2} \sum_i \sum_{j \in \mathcal{B}_i} |f_{ij}(t)| \\
&\leq \frac{1}{2} \sum_i \sum_{j \in \mathcal{B}_i} \sum_{\rho} |z_{ij}^{\rho}| |J_t^{\rho}| \\
&= \frac{1}{2} \sum_{\rho} |J_t^{\rho}| \sum_i \sum_{j \in \mathcal{B}_i} |z_{ij}^{\rho}| \\
&= \frac{1}{2} \sum_{\rho} |J_t^{\rho}| \sum_i m_i |\nu_i^{+\rho} - \nu_i^{-\rho}| \\
&= \sum_{\rho} \nu^{\rho} |J_t^{\rho}|, \quad (S18)
\end{aligned}$$

where $\nu^{\rho} = (1/2) \sum_{i} m_{i} |\nu_{i}^{+\rho} - \nu_{i}^{-\rho}|$.

C. Generalization to generic reaction networks

Here we derive the topological speed limit for generic (open) chemical reaction networks, in which the total mass concentration may not be conversed. To this end, we construct the graph corresponding to the time evolution of \boldsymbol{x}_t . Once done, the topological speed limit Eq. (5) is immediately obtained. We need only consider reaction channels that do not conserve the total mass concentration because the conservative reactions have already been considered above.

Analogously, for each reaction channel ρ , we define $S^{\rho}_{+} \coloneqq \{i \mid \nu_{i}^{+\rho} > \nu_{i}^{-\rho}\}$ and $S^{\rho}_{-} \coloneqq \{i \mid \nu_{i}^{+\rho} < \nu_{i}^{-\rho}\}$. A non-conservative reaction channel implies $\sum_{i \in S^{\rho}_{+}} m_{i}(\nu_{i}^{+\rho} - \nu_{i}^{-\rho}) \neq \sum_{i \in S^{\rho}_{-}} m_{i}(\nu_{i}^{-\rho} - \nu_{i}^{+\rho})$. Then, according to Prop. 2, there exist a matrix $Z^{\rho} = \{z_{ij}^{\rho}\} \in \mathbb{R}^{|S| \times |S|}$ and a vector $\boldsymbol{\mu}^{\rho} \in \mathbb{R}^{|S|}$ such that

$$\sum_{j \in \mathcal{S}_{-}^{\rho}} z_{ij}^{\rho} = \mu_i^{\rho} \le m_i (\nu_i^{+\rho} - \nu_i^{-\rho}), \quad \forall i \in \mathcal{S}_{+}^{\rho}, \tag{S19}$$

$$\sum_{i \in \mathcal{S}^{\rho}_{+}} z_{ij}^{\rho} = -\mu_j^{\rho} \le m_i (\nu_j^{-\rho} - \nu_j^{+\rho}), \quad \forall j \in \mathcal{S}^{\rho}_{-},$$
(S20)

$$z_{ji}^{\rho} = -z_{ij}^{\rho} \le 0, \quad \forall i \in \mathcal{S}_{+}^{\rho}, \quad j \in \mathcal{S}_{-}^{\rho}, \tag{S21}$$

$$z_{ij}^{\rho} = 0$$
, otherwise, (S22)

$$\mu_i^{\rho} \ge 0, \ \forall i \in \mathcal{S}_+^{\rho}, \tag{S23}$$

$$\mu_j^{\rho} \le 0, \ \forall j \in \mathcal{S}_-^{\rho}, \tag{S24}$$

$$\mu_i^{\rho} = 0$$
, otherwise, (S25)

$$\sum_{i \in \mathcal{S}_{+}^{\rho}} \mu_{i}^{\rho} = \min\left\{\sum_{i \in \mathcal{S}_{+}^{\rho}} m_{i}(\nu_{i}^{+\rho} - \nu_{i}^{-\rho}), \sum_{i \in \mathcal{S}_{-}^{\rho}} m_{i}(\nu_{i}^{-\rho} - \nu_{i}^{+\rho})\right\}.$$
(S26)

For each $z_{ij}^{\rho} \neq 0$, we add an undirected edge $\langle i, j \rangle$ to \mathcal{E} . Again, the existence of Z^{ρ} and μ^{ρ} may not be unique; nevertheless, one can always find coefficients such that at most $|\mathcal{S}_{+}^{\rho}| + |\mathcal{S}_{-}^{\rho}| - 1$ edges are added to the graph for each reaction channel ρ . Repeating this process for all $\rho \in \mathcal{R}$, we readily obtain the graph G.

With the above construction, the time evolution of the reaction network can be expressed in the form of Eq. (1) with

$$f_i(t) = \sum_{\rho} [m_i(\nu_i^{+\rho} - \nu_i^{-\rho}) - \mu_i^{\rho}] J_t^{\rho}, \qquad (S27)$$

$$f_{ij}(t) = \sum_{\rho} z_{ij}^{\rho} J_t^{\rho}.$$
 (S28)

For convenience, we define

$$\chi^{\rho} \coloneqq \min\left\{ \sum_{i \in \mathcal{S}_{+}^{\rho}} m_{i} (\nu_{i}^{+\rho} - \nu_{i}^{-\rho}), \sum_{i \in \mathcal{S}_{-}^{\rho}} m_{i} (\nu_{i}^{-\rho} - \nu_{i}^{+\rho}) \right\},$$
(S29)

$$\eta^{\rho} \coloneqq \left| \sum_{i \in \mathcal{S}_{+}^{\rho}} m_i (\nu_i^{+\rho} - \nu_i^{-\rho}) - \sum_{i \in \mathcal{S}_{-}^{\rho}} m_i (\nu_i^{-\rho} - \nu_i^{+\rho}) \right|.$$
(S30)

Using these quantities, the velocity term $v_{t,\lambda}$ can be upper bounded as

$$\upsilon_{t,\lambda} = \lambda \sum_{i} \left| \sum_{\rho} [m_i (\nu_i^{+\rho} - \nu_i^{-\rho}) - \mu_i^{\rho}] J_t^{\rho} \right| + \sum_{\langle i,j \rangle \in \mathcal{E}} \left| \sum_{\rho} z_{ij}^{\rho} J_t^{\rho} \right|$$

$$\leq \sum_{\rho} |J_t^{\rho}| \left(\lambda \sum_i |m_i(\nu_i^{+\rho} - \nu_i^{-\rho}) - \mu_i^{\rho}| + \sum_{\langle i,j \rangle \in \mathcal{E}} |z_{ij}^{\rho}| \right)$$
$$= \sum_{\rho} (\lambda \eta^{\rho} + \chi^{\rho}) |J_t^{\rho}|. \tag{S31}$$

Consequently, we obtain the following speed limit for generic reaction networks:

$$\tau \ge \tau_1 \coloneqq \frac{\mathcal{W}_{1,\lambda}(\boldsymbol{x}_0, \boldsymbol{x}_{\tau})}{\left(\sum_{\rho} (\lambda \eta^{\rho} + \chi^{\rho}) |J_t^{\rho}|\right)_{\tau}}.$$
 (S32)

Define the generalized kinetic quantity $\ell_{t,\lambda}$ as

$$\ell_{t,\lambda} \coloneqq \sum_{\rho} (\lambda \eta^{\rho} + \chi^{\rho})^2 \frac{J_t^{-\rho} - J_t^{+\rho}}{\ln(J_t^{-\rho}/J_t^{+\rho})}, \qquad (S33)$$

we can prove that

$$\left\langle \sum_{\rho} (\lambda \eta^{\rho} + \chi^{\rho}) | J_t^{\rho} | \right\rangle_{\tau} \le \left\langle \sqrt{\sigma_t \ell_{t,\lambda}} \right\rangle_{\tau} \le \sqrt{\langle \sigma_t \rangle_{\tau} \langle \ell_{t,\lambda} \rangle_{\tau}}.$$
(S34)

Combining Eqs. (S32) and (S34) yields the following thermodynamic speed limit:

$$\tau \ge \tau_2 \coloneqq \frac{\mathcal{W}_{1,\lambda}(\boldsymbol{x}_0, \boldsymbol{x}_\tau)}{\sqrt{\langle \sigma_t \rangle_\tau \langle \ell_{t,\lambda} \rangle_\tau}}, \ \forall \lambda \ge 0.$$
(S35)

So far, we have derived the speed limits for the mass concentration \boldsymbol{x}_t . It is worth noting that speed limits for the molar concentration \boldsymbol{c}_t can be readily obtained by simply setting $m_i = 1$ for all species X_i . Note that by this setting, the graph G and coefficients χ^{ρ} and η^{ρ} may change accordingly. Eventually, the following inequalities hold:

$$\tau \ge \tau_1' \coloneqq \frac{\mathcal{W}_{1,\lambda}(\boldsymbol{c}_0, \boldsymbol{c}_\tau)}{\left\langle \sum_{\rho} (\lambda \eta^{\rho} + \chi^{\rho}) | J_t^{\rho} | \right\rangle_{\tau}} \ge \tau_2' \coloneqq \frac{\mathcal{W}_{1,\lambda}(\boldsymbol{c}_0, \boldsymbol{c}_\tau)}{\sqrt{\langle \sigma_t \rangle_{\tau} \langle \ell_{t,\lambda} \rangle_{\tau}}}.$$
(S36)

Next, we show that the $\lambda = 1/2$ speed limit is always tighter than a thermodynamic bound reported in Ref. [5], which reads

$$\tau \ge \tau_3 \coloneqq \frac{\mathcal{T}(\boldsymbol{c}_0, \boldsymbol{c}_\tau)}{\sqrt{\langle \sigma_t \rangle_\tau \langle d_t \rangle_\tau}},\tag{S37}$$

where $\mathcal{T}(\boldsymbol{c}_0, \boldsymbol{c}_{\tau}) \coloneqq \|\boldsymbol{c}_0 - \boldsymbol{c}_{\tau}\|_1/2$ is the total variation distance and $d_t \coloneqq (|\mathcal{S}|/8) \sum_{\rho,i} (\nu_i^{+\rho} - \nu_i^{-\rho})^2 (J_t^{+\rho} + J_t^{-\rho})$ corresponds to the diffusion coefficient. Note that $(1/2)\eta^{\rho} + \chi^{\rho} = (1/2) \sum_i |\nu_i^{+\rho} - \nu_i^{-\rho}|$. Applying the Cauchy–Schwarz inequality and the inequality $(x - y)/\ln(x/y) \leq (x + y)/2$, we have

$$\ell_{t,1/2} = \frac{1}{4} \sum_{\rho} \left(\sum_{i} |\nu_{i}^{+\rho} - \nu_{i}^{-\rho}| \right)^{2} \frac{J_{t}^{-\rho} - J_{t}^{+\rho}}{\ln(J_{t}^{-\rho}/J_{t}^{+\rho})}$$
$$\leq \frac{|\mathcal{S}|}{8} \sum_{\rho,i} (\nu_{i}^{+\rho} - \nu_{i}^{-\rho})^{2} (J_{t}^{-\rho} + J_{t}^{+\rho})$$
$$= d_{t}.$$
(S38)

By combining the above inequality and the relation $\mathcal{W}_{1,1/2}(\boldsymbol{c}_0, \boldsymbol{c}_{\tau}) = \mathcal{T}(\boldsymbol{c}_0, \boldsymbol{c}_{\tau})$, we readily obtain the following hierarchical relationship:

$$\tau_1' \ge \tau_2' \ge \tau_3. \tag{S39}$$

Proposition 2. Let $\boldsymbol{a} = [a_1, \ldots, a_n]^{\mathsf{T}}$ and $\boldsymbol{b} = [b_1, \ldots, b_m]^{\mathsf{T}}$ be two vectors of positive numbers. Then there exist matrix $\mathsf{Z} = [z_{ij}] \in \mathbb{R}_{\geq 0}^{n \times m}$ and nonnegative vectors $\tilde{\boldsymbol{a}} = [\tilde{a}_1, \ldots, \tilde{a}_n]^{\mathsf{T}}$ and $\tilde{\boldsymbol{b}} = [\tilde{b}_1, \ldots, \tilde{b}_m]^{\mathsf{T}}$ such that the following conditions are satisfied:

$$\sum_{i=1}^{m} z_{ij} = \tilde{a}_i \le a_i, \quad \forall 1 \le i \le n,$$
(S40)

$$\sum_{i=1}^{n} z_{ij} = \tilde{b}_j \le b_j, \quad \forall 1 \le j \le m,$$
(S41)

$$\sum_{i} \tilde{a}_{i} = \min\left\{\sum_{i} a_{i}, \sum_{j} b_{j}\right\}.$$
 (S42)

Proof. Without loss of generality, we can assume that $a_1 \leq \cdots \leq a_n$ and $b_1 \leq \cdots \leq b_m$. We prove by induction on $k = m + n \geq 2$. In the case of k = 2 (i.e., m = n = 1), we can set $z_{11} = \tilde{a}_1 = \tilde{b}_1 = \min\{a_1, b_1\}$. Suppose that the statement holds for all $k \leq \bar{k}$. We consider an arbitrary case with $k = \bar{k} + 1$. Assume that $a_1 \leq b_1$. If n = 1 then we can choose $z_{11} = a_1, z_{1j} = 0$ for all j > 1, $\tilde{a} = [a_1]$, and $\tilde{b} = [a_1, 0, \ldots, 0]^{\mathsf{T}}$. If $n \geq 2$, then consider two vectors $a' = [a_2, \ldots, a_n]^{\mathsf{T}}$ and $b' = [b_1 - a_1, b_2, \ldots, b_m]^{\mathsf{T}}$. There exist Z', \tilde{a}' , and \tilde{b}' such that

$$\sum_{j=1}^{m} z'_{ij} = \tilde{a}'_i \le a'_i, \ \forall 1 \le i \le n-1,$$
(S43)

$$\sum_{i=1}^{n-1} z'_{ij} = \tilde{b}'_j \le b'_j, \quad \forall 1 \le j \le m,$$
(S44)

$$\sum_{i} \tilde{a}'_{i} = \min\left\{\sum_{i} a'_{i}, \sum_{j} b'_{j}\right\}.$$
 (S45)

We construct $z_{11} = a_1$, $z_{1j} = 0$ for all j > 1, $z_{ij} = z'_{(i-1)j}$ for all $i \ge 2$, $\tilde{\boldsymbol{a}} = [a_1, \tilde{a}'_1, \dots, \tilde{a}'_{n-1}]^{\mathsf{T}}$, and $\tilde{\boldsymbol{b}} = [a_1 + \tilde{b}'_1, \tilde{b}'_2, \dots, \tilde{b}'_m]^{\mathsf{T}}$. It is easy to verify that this combination satisfies all conditions (S40), (S41), and (S42).

For example, for two vectors $\boldsymbol{a} = [4,5]^{\mathsf{T}}$ and $\boldsymbol{b} = [1,2,3]^{\mathsf{T}}$, we can construct an instance of matrix Z and vectors $\tilde{\boldsymbol{a}}, \tilde{\boldsymbol{b}}$ as follows:

$$\mathsf{Z} = \begin{bmatrix} 1 & 2 & 1 \\ 0 & 0 & 2 \end{bmatrix}, \quad \tilde{\boldsymbol{a}} = \begin{bmatrix} 4, 2 \end{bmatrix}^{\mathsf{T}}, \quad \tilde{\boldsymbol{b}} = \begin{bmatrix} 1, 2, 3 \end{bmatrix}^{\mathsf{T}}.$$
(S46)

S4. BOSONIC TRANSPORT

A. Time evolution of boson number

Using the relation $[b_i, \hat{n}_i] = b_i$ and $[\hat{n}_i, h_Z] = 0$, we can calculate as follows:

$$-i\operatorname{tr}\{\hat{n}_i[H_t,\varrho_t]\} = -i\operatorname{tr}\{[\hat{n}_i,H_t]\varrho_t\}$$

$$b_j^\dagger b_i] arrho_t \Big\}$$

4

$$= i\gamma \sum_{j \in \mathcal{B}_i} \operatorname{tr}\left\{ (b_i^{\dagger} b_j - b_j^{\dagger} b_i) \varrho_t \right\}$$
$$= 2\gamma \sum_{j \in \mathcal{B}_i} \operatorname{Im}\left[\operatorname{tr}\left\{ b_j^{\dagger} b_i \varrho_t \right\} \right].$$
(S47)

 $=i\gamma\sum_{i=n}$ tr{ $[\hat{n}_i, b_i^{\dagger}b_j +$

Similarly, by noting that $[b_i, b_i^{\dagger}] = 1$, we have

$$\operatorname{tr}\left\{\hat{n}_{i}\sum_{j\in\Lambda}\left(\mathcal{D}[L_{j,+}]+\mathcal{D}[L_{j,-}]\right)\varrho_{t}\right\}$$

$$=\operatorname{tr}\left\{\hat{n}_{i}\sum_{j\in\Lambda}\gamma_{j,+}\left(b_{j}^{\dagger}\varrho_{t}b_{j}-\{b_{j}b_{j}^{\dagger},\varrho_{t}\}/2\right)\right\}$$

$$+\operatorname{tr}\left\{\hat{n}_{i}\sum_{j\in\Lambda}\gamma_{j,-}\left(b_{j}\varrho_{t}b_{j}^{\dagger}-\{b_{j}^{\dagger}b_{j},\varrho_{t}\}/2\right)\right\}$$

$$=\operatorname{tr}\left\{\hat{n}_{i}\gamma_{i,+}\left(b_{i}^{\dagger}\varrho_{t}b_{i}-\{\hat{n}_{i}+1,\varrho_{t}\}/2\right)\right\}$$

$$+\operatorname{tr}\left\{\hat{n}_{i}\gamma_{i,-}\left(b_{i}\varrho_{t}b_{i}^{\dagger}-\{\hat{n}_{i},\varrho_{t}\}/2\right)\right\}$$

$$=\gamma_{i,+}\operatorname{tr}\left\{b_{i}^{\dagger}\varrho_{t}(b_{i}+\hat{n}_{i}b_{i})-\{\hat{n}_{i}^{2}+\hat{n}_{i},\varrho_{t}\}/2\right\}$$

$$=\gamma_{i,+}\operatorname{tr}\left\{b_{i}^{\dagger}\varrho_{t}b_{i}\right\}-\gamma_{i,-}\operatorname{tr}\left\{b_{i}\varrho_{t}b_{i}^{\dagger}\right\}$$

$$=\operatorname{tr}\left\{L_{i,+}\varrho_{t}L_{i,+}^{\dagger}\right\}-\operatorname{tr}\left\{L_{i,-}\varrho_{t}L_{i,-}^{\dagger}\right\}.$$
(S48)

Taking the time derivative of $x_i(t) = tr\{\hat{n}_i \varrho_t\}$, we can calculate the time evolution of $x_i(t)$ as follows:

$$\dot{x}_{i}(t) = \operatorname{tr}\{\hat{n}_{i}\dot{\varrho}_{t}\}$$

$$= \operatorname{tr}\left\{-i\hat{n}_{i}[H_{t},\varrho_{t}] + \hat{n}_{i}\sum_{i\in\Lambda}\left(\mathcal{D}[L_{i,+}] + \mathcal{D}[L_{i,-}]\right)\varrho_{t}\right\}$$

$$= 2\gamma\sum_{j\in\mathcal{B}_{i}}\operatorname{Im}\left[\operatorname{tr}\left\{b_{j}^{\dagger}b_{i}\varrho_{t}\right\}\right]$$

$$+ \operatorname{tr}\left\{L_{i,+}\varrho_{t}L_{i,+}^{\dagger}\right\} - \operatorname{tr}\left\{L_{i,-}\varrho_{t}L_{i,-}^{\dagger}\right\}$$

$$= f_{i}(t) + \sum_{j\in\mathcal{B}_{i}}f_{ij}(t), \qquad (S49)$$

where

$$f_i(t) = \operatorname{tr}\left\{L_{i,+} \varrho_t L_{i,+}^{\dagger}\right\} - \operatorname{tr}\left\{L_{i,-} \varrho_t L_{i,-}^{\dagger}\right\}, \qquad (S50)$$

$$f_{ij}(t) = 2\gamma \operatorname{Im}\left[\operatorname{tr}\left\{b_j^{\dagger} b_i \varrho_t\right\}\right].$$
(S51)

It can be verified that $f_{ij}(t) = -f_{ji}(t)$.

B. Derivation of Eq. (18)

Applying the Cauchy–Schwarz inequality, we can upper bound $|f_{ij}(t)|$ as follows:

$$|f_{ij}(t)| \leq 2\gamma \sqrt{\operatorname{tr}\{b_i^{\dagger}b_i\varrho_t\}\operatorname{tr}\{b_j^{\dagger}b_j\varrho_t\}}$$

$$\leq \gamma \left(\operatorname{tr}\{b_i^{\dagger}b_i\varrho_t\} + \operatorname{tr}\{b_j^{\dagger}b_j\varrho_t\}\right)$$

$$= \gamma [x_i(t) + x_j(t)].$$
(S52)

Taking the sum over all edges $\langle i, j \rangle \in \mathcal{E}$, we obtain

$$\sum_{\langle i,j \rangle} |f_{ij}(t)| \leq \sum_{\langle i,j \rangle} \gamma [x_i(t) + x_j(t)]$$
$$\leq \gamma d_G \sum_i x_i(t) = \gamma d_G \mathcal{N}_t.$$
(S53)

In addition, according to Prop. 3, we have

$$\sum_{i} |f_{i}(t)| = \sum_{i} \left| \operatorname{tr} \left\{ L_{i,+} \varrho_{t} L_{i,+}^{\dagger} \right\} - \operatorname{tr} \left\{ L_{i,-} \varrho_{t} L_{i,-}^{\dagger} \right\} \right|$$
$$\leq \frac{\sigma_{t}}{2} \Phi \left(\frac{\sigma_{t}}{2a_{t}} \right)^{-1}.$$
(S54)

By combining the inequalities (S53) and (S54), the velocity $v_{t,\lambda}$ can be upper bounded as

$$\upsilon_{t,\lambda} \le \gamma d_G \mathcal{N}_t + \lambda \frac{\sigma_t}{2} \Phi \left(\frac{\sigma_t}{2a_t}\right)^{-1}.$$
 (S55)

Proposition 3. The following inequality holds:

$$\sum_{i} \left| \operatorname{tr} \left\{ L_{i,+} \varrho_t L_{i,+}^{\dagger} \right\} - \operatorname{tr} \left\{ L_{i,-} \varrho_t L_{i,-}^{\dagger} \right\} \right| \le \frac{\sigma_t}{2} \Phi \left(\frac{\sigma_t}{2a_t} \right)^{-1},$$
(S56)

where $\Phi(x)$ is the inverse function of $x \tanh(x)$.

Proof. Let $\varrho_t = \sum_n p_n(t) |n_t\rangle \langle n_t|$ be the spectral decomposition of the density operator ϱ_t . Define $r_{mn}^{i,\pm}(t) \coloneqq |\langle m_t | L_{i,\pm} | n_t \rangle|^2$, then the rates of entropy production and dynamical activity can be expressed as

$$\sigma_{t} = \sum_{i} \sum_{m,n} [r_{mn}^{i,+}(t)p_{n}(t) - r_{nm}^{i,-}(t)p_{m}(t)] \ln \frac{r_{mn}^{i,+}(t)p_{n}(t)}{r_{nm}^{i,-}(t)p_{m}(t)}$$

$$= \sum_{i} \sum_{m,n} \sigma_{mn}^{i}(t), \qquad (S57)$$

$$a_{t} = \sum_{i} \sum_{m,n} [r_{mn}^{i,+}(t)p_{n}(t) + r_{nm}^{i,-}(t)p_{m}(t)]$$

$$= \sum_{i} \sum_{m,n} a_{mn}^{i}(t). \qquad (S58)$$

Note that $x\Phi(x/y)^{-1}$ is a is a concave function over $(0, +\infty) \times (0, +\infty)$. Applying Jensen's inequality yields

$$\sum_{i} \left| \operatorname{tr} \left\{ L_{i,+} \varrho_{t} L_{i,+}^{\dagger} \right\} - \operatorname{tr} \left\{ L_{i,-} \varrho_{t} L_{i,-}^{\dagger} \right\} \right|$$

$$= \sum_{i} \left| \sum_{m,n} \left[r_{mn}^{i,+}(t) p_{n}(t) - r_{nm}^{i,-}(t) p_{m}(t) \right] \right|$$

$$\leq \sum_{i} \sum_{m,n} \left| r_{mn}^{i,+}(t) p_{n}(t) - r_{nm}^{i,-}(t) p_{m}(t) \right|$$

$$= \sum_{i} \sum_{m,n} \frac{\sigma_{mn}^{i}(t)}{2} \Phi \left(\frac{\sigma_{mn}^{i}(t)}{2a_{mn}^{i}(t)} \right)^{-1}$$

$$\leq \frac{\sigma_{t}}{2} \Phi \left(\frac{\sigma_{t}}{2a_{t}} \right)^{-1}, \quad (S59)$$

which completes the proof.

C. Numerical demonstration

Here we numerically demonstrate the speed limits for bosonic transport in a one-dimensional Bose–Hubbard model with the size of N = 15 [see Fig. S2(a) for illustration]. Bosons can hop from site *i* to site *j* if and only if |i - j| = 1. The Hamiltonian is given by

$$H_t = -\gamma \sum_{i=1}^{N-1} (b_i^{\dagger} b_{i+1} + b_{i+1}^{\dagger} b_i) + \sum_{i=1}^N U_i(t) \frac{\hat{n}_i(\hat{n}_i - 1)}{2}.$$
 (S60)

For this isolated bosonic system, the flows are determined as follows:

$$f_i(t) = 0, \tag{S61}$$

$$f_{ij}(t) = \delta_{|i-j|,1} 2\gamma \operatorname{Im}\left[\operatorname{tr}\left\{b_j^{\dagger} b_i \varrho_t\right\}\right], \quad (S62)$$

and the maximal vertex degree is $d_G = 2$. Here $\delta_{x,y}$ denotes the Kronecker delta. The velocity v_t is thus given by

$$\upsilon_t = 2\gamma \sum_{i=1}^{N-1} \left| \operatorname{Im} \left[\operatorname{tr} \left\{ b_{i+1}^{\dagger} b_i \varrho_t \right\} \right] \right|.$$
 (S63)

Consequently, the derived speed limit reads

$$\tau \ge \frac{\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_\tau)}{\langle \upsilon_t \rangle_\tau} \eqqcolon \tau_1.$$
(S64)

For comparison, we also examine two other nontopological speed limits. The first one is the generalization of the Mandelstam–Tamm bound [6], which reads

$$\tau \ge \frac{\mathcal{L}(\varrho_0, \varrho_\tau)}{\langle \Delta_{\varrho_t} H_t \rangle_\tau} = \tau_{\rm MT},\tag{S65}$$

where $\mathcal{L}(\varrho, \sigma) \coloneqq \operatorname{arccostr}\{|\varrho^{1/2}\sigma^{1/2}|\}$ is the Bures angle. The second one is a speed limit using the total variation distance $\mathcal{T}(\boldsymbol{x}, \boldsymbol{y}) \coloneqq \|\boldsymbol{x} - \boldsymbol{y}\|_{1}/2$, which does not consider the topological nature and is given by

$$\tau \ge \frac{\mathcal{T}(\boldsymbol{x}_0, \boldsymbol{x}_\tau)}{\langle \boldsymbol{\upsilon}_t \rangle_\tau} \eqqcolon \tau_{\mathrm{TV}}.$$
 (S66)

Notice that $\tau_1 \geq \tau_{\text{TV}}$ since $\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_{\tau}) \geq \mathcal{T}(\boldsymbol{x}_0, \boldsymbol{x}_{\tau})$.

We set the initial state to the ground state of the Hamiltonian H_0 , which can be calculated using the algorithm of density matrix renormalization group [7]. Initially, bosons are almost uniformly distributed. The time-dependent Hamiltonian is modulated as $H_t = (1 - t/\tau)H_0 + (t/\tau)H_{\tau}$, where H_{τ} is the final Hamiltonian. More specifically, the interaction coefficients $U_i(t)$ $(1 \le i \le N)$ are given by

$$U_i(t) = \left(1 - \frac{t}{\tau}\right) U_0 + \frac{t}{\tau} U_{\tau,i}, \qquad (S67)$$

where $\{U_{\tau,i}\}$ are the interaction coefficients of the final Hamiltonian H_{τ} . By this protocol, bosons tend to gather




FIG. S2. (a) Schematic of the one-dimensional Bose–Hubbard chain with a finite length. Bosons can hop from site i to i + 1 and vice versa. (b) Time variation of the boson number at each site $i \in [1, 15]$. The bosonic system is initiated in the ground state of H_0 , in which bosons are almost uniformly distributed. After a period of time τ , bosons tend to gather at the central site. (c) Numerical verification of the speed limits. The operational time t, topological bound t_1 , and non-topological bounds $t_{\rm MT}$ and $t_{\rm TV}$ are depicted by solid, dashed, and dash-dotted and dotted lines, respectively. The parameters are fixed at $U_0 = 10$, $\gamma = 1$, and $U_{\tau} = [10, 9.5, 8.5, 7, 5, 2, 0.5, -5, 0.5, 2, 5, 7, 8.5, 9.5, 10]^{T}$.

at the central site after a period of time τ [see Fig. S2(b)]. We employ the time dependent variational principle algorithm [8] to simulate the time evolution of the bosonic system. For each time $t \leq \tau$ (= 10), we calculate all relevant quantities and plot the bounds in Fig. S2(c). As shown, the topological bound $t \geq t_1$ is tight and can be saturated. In contrast, the non-topological bound $t_{\rm TV}$ cannot accurately capture the operational time of bosonic transport. This is because the total variation distance does not consider topological information as the Wasserstein distance does. Although the Mandelstam–Tamm bound $t \geq t_{\rm MT}$ yields good prediction for $t \leq 6$, it subsequently becomes trivial as t increases.

S5. ISOLATED QUANTUM SYSTEMS

Here we derive a topological speed limit for isolated systems whose dynamics is described by the von Neumann equation:

$$\dot{\varrho}_t = -i[H_t, \varrho_t], \tag{S68}$$

where H_t is a time-dependent Hamiltonian. We consider a complete orthogonal set of projection operators $\{P_n\}_n$, i.e., $P_m P_n = \delta_{mn} P_n$ and $\sum_n P_n = \mathbb{1}$. Since the dynamics is invariant under transformation $H_t \to H_t + \alpha_t \mathbb{1}$, we can assume that $\operatorname{tr}\{H_t \tilde{\varrho}_t\} = 0$, where $\tilde{\varrho}_t \coloneqq \sum_n P_n \varrho_t P_n$ is a density matrix projected on the space of $\{P_n\}_n$. Let $G(\mathcal{V}, \mathcal{E})$ be the graph, in which $\mathcal{V} = \{n\}_n$ and $\mathcal{E} = \{\langle m, n \rangle \mid P_m H_t P_n \neq 0, m < n\}$, and d_G be the maximum vertex degree of the graph. Considering a vector of projective measurements, $x_n(t) = \operatorname{tr}\{P_n \varrho_t\}$, we obtain the time evolution of $x_n(t)$ in the following equation:

$$\dot{x}_n(t) = -i \sum_{m(\neq n)} \left(\operatorname{tr} \{ P_n H_t P_m \varrho_t \} - \operatorname{tr} \{ P_m H_t P_n \varrho_t \} \right)$$

$$=\sum_{m\in\mathcal{B}_n}f_{nm}(t),\tag{S69}$$

where $f_{mn}(t) = -f_{nm}(t)$ is given by

$$f_{nm}(t) = -i(\operatorname{tr}\{P_n H_t P_m \varrho_t\} - \operatorname{tr}\{P_m H_t P_n \varrho_t\}). \quad (S70)$$

Applying the speed limit in Eq. (6), we obtain the following bound on the operational time:

$$\tau \ge \frac{\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_\tau)}{\langle \boldsymbol{v}_t \rangle_\tau},\tag{S71}$$

(S74)

where v_t can be explicitly expressed as

$$\upsilon_t = \sum_{(m,n)\in\mathcal{E}} |\mathrm{tr}\{(P_n H_t P_m - P_m H_t P_n)\varrho_t\}|.$$
(S72)

By further upper bounding v_t , we can also obtain another speed limit that resembles the Mandelstam–Tamm bound. By applying the Cauchy–Schwarz inequality, we have

$$\begin{aligned} \left|\operatorname{tr}\{P_{n}H_{t}P_{m}\varrho_{t}\}\right| &\leq \operatorname{tr}\{P_{m}H_{t}P_{n}H_{t}P_{m}\varrho_{t}\}^{1/2}\operatorname{tr}\{P_{n}\varrho_{t}\}^{1/2}, \\ (S73) \end{aligned}$$
$$\left|\operatorname{tr}\{P_{m}H_{t}P_{n}\varrho_{t}\}\right| &\leq \operatorname{tr}\{P_{n}H_{t}P_{m}H_{t}P_{n}\varrho_{t}\}^{1/2}\operatorname{tr}\{P_{m}\varrho_{t}\}^{1/2}. \end{aligned}$$

Combining the above inequalities and applying the Cauchy–Schwarz inequality yield

$$|f_{nm}(t)| \leq \left[tr\{ (P_m H_t P_n H_t P_m + P_n H_t P_m H_t P_n) \varrho_t \} \right]^{1/2} \times \left[tr\{ (P_n + P_m) \varrho_t \} \right]^{1/2}.$$
 (S75)

Therefore, using the fact $\sum_n P_n = 1$, we have

$$\upsilon_t \leq \sum_{(m,n)\in\mathcal{E}} \left[\operatorname{tr} \left\{ \left(P_m H_t P_n H_t P_m + P_n H_t P_m H_t P_n \right) \varrho_t \right\} \right]^{1/2}$$

$$\times \left[\operatorname{tr} \{ (P_n + P_m) \varrho_t \} \right]^{1/2}$$

$$\leq \left[\sum_{\langle m,n \rangle \in \mathcal{E}} \operatorname{tr} \{ (P_m H_t P_n H_t P_m + P_n H_t P_m H_t P_n) \varrho_t \} \right]^{1/2}$$

$$\times \left[\sum_{\langle m,n \rangle \in \mathcal{E}} \operatorname{tr} \{ (P_n + P_m) \varrho_t \} \right]^{1/2}$$

$$\leq \left[\sum_n \operatorname{tr} \{ H_t^2 P_n \varrho_t P_n \} \right]^{1/2} \sqrt{d_G}$$

$$= \sqrt{d_G} \Delta_{\tilde{\varrho}_t} H_t, \qquad (S76)$$

where $\Delta_{\varrho} H_t \coloneqq \sqrt{\operatorname{tr} \{H_t^2 \varrho\} - (\operatorname{tr} \{H_t \varrho\})^2}$ is the energy fluctuation of the Hamiltonian H_t with respect to ϱ . Consequently, we obtain the following speed limit:

$$\tau \ge \frac{\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_\tau)}{\sqrt{d_G} \left\langle \Delta_{\tilde{\varrho}_t} H_t \right\rangle_\tau}.$$
 (S77)

We note that this speed limit is similar but different from the generalization of the Mandelstam–Tamm bound [6], which reads

$$\tau \ge \frac{\mathcal{L}(\varrho_0, \varrho_\tau)}{\langle \Delta_{\varrho_t} H_t \rangle_{\tau}},\tag{S78}$$

where $\mathcal{L}(\varrho, \sigma) \coloneqq \operatorname{arccostr}\{|\varrho^{1/2}\sigma^{1/2}|\}\$ is the Bures angle. Notably, the new speed limit can be applied to quantum systems under measurement, making it more applicable than the generalized Mandelstam–Tamm bound.

We now consider a specific case where $P_n = |n\rangle\langle n|$. In this case, $|f_{mn}(t)|$ can be upper bounded as

$$|f_{mn}(t)| = |\langle \langle n|H_t|m \rangle \langle m|\varrho_t|n \rangle - \langle m|H_t|n \rangle \langle n|\varrho_t|m \rangle)|$$

$$\leq 2|\langle m|H_t|n \rangle ||\langle n|\varrho_t|m \rangle|.$$
(S79)

According to the Cauchy-Schwarz inequality, we have

$$|\langle n|\varrho_t|m\rangle| = |\langle n|\varrho_t^{1/2}\varrho_t^{1/2}|m\rangle| \le \sqrt{\langle n|\varrho_t|n\rangle \langle m|\varrho_t|m\rangle}.$$
(S80)

Therefore,

$$|f_{mn}(t)| \le 2|\langle m|H_t|n\rangle|\sqrt{\langle n|\varrho_t|n\rangle\langle m|\varrho_t|m\rangle}.$$
 (S81)

Consequently, the velocity can be upper bounded in terms of the Hamiltonian and diagonal terms of density matrix as

$$\upsilon_t \le 2 \sum_{(m,n)\in\mathcal{E}} |h_{mn}(t)| \sqrt{p_n(t)p_m(t)} \rightleftharpoons \check{\upsilon}_t, \qquad (S82)$$

where we have defined $h_{mn}(t) \coloneqq \langle m|H_t|n \rangle$ and $p_n(t) \coloneqq \langle n|\varrho_t|n \rangle$ for simplicity. Then, Eq. (S71) yields the following speed limit:

$$\tau \ge \frac{\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_\tau)}{\langle \check{\boldsymbol{\upsilon}}_t \rangle_{\tau}},\tag{S83}$$

which is easier to compute than the original bound (S71).

S6. MEASUREMENT-INDUCED QUANTUM WALK

Here we demonstrate an application of the speed limits (S71), (S77), and (S83) for a quantum system under measurement. We consider a model of the continuous-time quantum walk [9], which is induced by measurements performed at discrete times. The system Hamiltonian is given by

$$H = \sum_{n=1}^{N-1} \gamma_n (|n| \langle n+1| + |n+1| \langle n|), \qquad (S84)$$

which describes hops between nearest neighbors on a finite line lattice. The projective measurements with operators $\{|n\rangle\langle n|\}_n$ are performed at times $t_k \coloneqq k\Delta t$ for $k = 0, \ldots, K (= \tau/\Delta t)$. Between each measurement event, the system unitarily evolves according to the von Neumman equation. The density matrix after the kth measurement is given by

$$\varrho_{t_k^+} = \sum_n \langle n | \varrho_{t_k} | n \rangle | n \rangle \langle n | .$$
 (S85)

For this system, we consider the set of projection operators $\{P_n = |n\rangle\langle n|\}$. The graph G thus has N vertices and N-1 edges that connect n and n+1 for all $1 \le n \le N-1$. The maximum degree of the graph is $d_G = 2$. Applying Eqs. (S71) and (S77) to the time evolution of the system between the kth and (k + 1)th measurements yields the following result:

$$\sqrt{2} \int_{t_k}^{t_{k+1}} \Delta_{\tilde{\varrho}_t} H \, dt \ge \int_{t_k}^{t_{k+1}} \upsilon_t \, dt \ge \mathcal{W}_1(\boldsymbol{x}_{t_k}, \boldsymbol{x}_{t_{k+1}}).$$
(S86)

Taking the sum of both sides of Eq. (S86) for $k = 0, \ldots, K-1$ and applying the triangle inequality for W_1 , we obtain

$$\sqrt{2} \int_0^\tau \Delta_{\tilde{\varrho}_t} H \, dt \ge \int_0^\tau \upsilon_t \, dt \ge \mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_\tau). \tag{S87}$$

Consequently, the following speed limits are derived:

$$\tau \ge \frac{\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_{\tau})}{\langle \boldsymbol{v}_t \rangle_{\tau}} \ge \frac{\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_{\tau})}{\sqrt{2} \langle \Delta_{\tilde{\varrho}_t} H \rangle_{\tau}}.$$
 (S88)

We note that the system is measured at discrete times; therefore, the generalized Mandelstam–Tamm bound cannot be applied directly.

By applying the speed limit (S83), we can also obtain another meaningful bound. Note that $\mathcal{E} = \{\langle n, n+1 \rangle | 1 \le n \le N-1\}$ and $h_{mn}(t) = \gamma_{\min\{m,n\}} \delta_{|m-n|,1}$ in this system. The quantity \check{v}_t can be upper bounded as follows:

$$\begin{split} \dot{v}_t &= 2\sum_{n=1}^{N-1} \gamma_n \sqrt{p_n(t)p_{n+1}(t)} \\ &\leq \sum_{n=1}^{N-1} \gamma_n [p_n(t) + p_{n+1}(t)] \end{split}$$

$$\leq 2 \max_{n} \gamma_n. \tag{S89}$$

Therefore, the operational time is lower bounded as

$$\tau \ge \frac{\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_\tau)}{2\max_n \gamma_n}.$$
 (S90)

S7. QUANTUM COMMUNICATION USING SPIN SYSTEMS

Here we derive a topological speed limit for quantum communication through an arbitrary graph $G(\mathcal{V}, \mathcal{E})$ of spins with ferromagnetic Heisenberg interactions [10]. The vertices $\{n \in \mathcal{V}\}_n$ represent spins and the edges $\{\langle n, m \rangle \in \mathcal{E}\}$ connect interacting spins. The Hamiltonian is given by

$$H_t = -\frac{\gamma}{2} \sum_{(n,m)\in\mathcal{E}} \vec{\sigma}_n \cdot \vec{\sigma}_m + \sum_{n\in\mathcal{V}} B_n(t)\sigma_n^z, \qquad (S91)$$

where $\gamma > 0$ denotes the coupling strength, $\vec{\sigma}_n = [\sigma_n^x, \sigma_n^y, \sigma_n^z]^{\mathsf{T}}$ is the vector of Pauli spin operators for the *n*th spin, and $B_n(t)$ is an external magnetic field in the *z* direction.

We assume that the sender, Alice, has a quantum state encoded in spin 1 and wants to relay it to the receiver, Bob, who can access and read out spin N. By manipulating the external magnetic field (which Alice can control), quantum information can be transmitted through the graph of spins. After a predetermined time when the state of spin 1 is transferred to spin N, Bob reads out the state of this site. The minimum time required for high-fidelity information transmission is the quantity of interest.

Consider state x_t with $x_n(t) = (\operatorname{tr}\{\sigma_n^z \varrho_t\} + 1)/2 \ge 0$. Notice that $x_n(t) = 0$ $(x_n(t) = 1)$ corresponds to spin down (up) with respect to the z direction. Taking the time derivative of $x_i(t)$, we can calculate as follows:

$$\begin{aligned} \dot{x}_n(t) &= \operatorname{tr}\{\sigma_n^z \dot{\varrho}_t\}/2 \\ &= (i/2) \operatorname{tr}\{[\sigma_n^z, H_t] \varrho_t\} \\ &= -(i\gamma/4) \sum_{m \in \mathcal{B}_n} \operatorname{tr}\{[\sigma_n^z, \vec{\sigma}_n \cdot \vec{\sigma}_m] \varrho_t\} \\ &= (\gamma/2) \sum_{m \in \mathcal{B}_n} \operatorname{tr}\{(\sigma_n^y \sigma_m^x - \sigma_n^x \sigma_m^y) \varrho_t\} \\ &= \sum_{m \in \mathcal{B}_n} f_{nm}(t), \end{aligned}$$
(S92)

where $f_{nm}(t) = -f_{mn}(t)$ is given by

$$f_{nm}(t) = (\gamma/2) \operatorname{tr} \{ (\sigma_n^y \sigma_m^x - \sigma_n^x \sigma_m^y) \varrho_t \}.$$
(S93)

Therefore, the velocity can be expressed as

$$\upsilon_t = \frac{\gamma}{2} \sum_{(m,n)\in\mathcal{E}} |\operatorname{tr}\{(\sigma_n^y \sigma_m^x - \sigma_n^x \sigma_m^y)\varrho_t\}|.$$
(S94)

Consequently, the derived topological speed limit reads

$$\tau \ge \frac{\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_\tau)}{\langle \boldsymbol{v}_t \rangle_\tau}.$$
 (S95)

It can be easily verified that

$$\sigma_n^y \sigma_m^x - \sigma_n^x \sigma_m^y \le (\sigma_n^z + \mathbb{1}_2) \otimes \mathbb{1}_2 + \mathbb{1}_2 \otimes (\sigma_m^z + \mathbb{1}_2), \quad (S96)$$

$$\sigma_n^x \sigma_m^y - \sigma_n^y \sigma_m^x \le (\sigma_n^z + \mathbb{1}_2) \otimes \mathbb{1}_2 + \mathbb{1}_2 \otimes (\sigma_m^z + \mathbb{1}_2), \quad (S97)$$

where $\mathbb{1}_2$ is the two-dimensional identity matrix. Here, $A \leq B$ means that B - A is positive semi-definite. Therefore, $|f_{nm}(t)|$ can be upper bounded as

$$|f_{nm}(t)| \le \gamma [x_n(t) + x_m(t)]. \tag{S98}$$

Using Eq. (S98), we can show that the velocity v_t is bounded from above as

$$\upsilon_t \le \sum_{(n,m)\in\mathcal{E}} \gamma[x_n(t) + x_m(t)] \le \gamma d_G \|\boldsymbol{x}_t\|_1.$$
 (S99)

It should be noted that the total spin $\|\boldsymbol{x}_t\|_1$ is invariant for all times. For convenience, we define $\mathcal{M} \coloneqq \|\boldsymbol{x}_t\|_1$. Using the inequality (S99) and the topological speed limit (S95), we obtain the following bound on the operational time required for transmitting information:

$$\tau \ge \frac{\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_\tau)}{\gamma d_G \mathcal{M}}.$$
 (S100)

This inequality implies that the speed of information transmission is constrained by the topology of the graph G and the coupling strength.

Now let us discuss a particular case where the graph G is a spin chain of length $N \ge 2$, and the spins n and n+1 interact with each other for all $1 \le n \le N-1$ [10, 11]. Alice prepares the spin chain in the initial state $\rho_0 = |\varphi_0\rangle\langle\varphi_0|$, with spin 1 in the excited state $|1\rangle$, and all other spins in the ground state $|0\rangle$. Specifically, $|\varphi_0\rangle$ is given by

$$|\varphi_0\rangle = |1\rangle \otimes |0\rangle \otimes \cdots \otimes |0\rangle.$$
 (S101)

In this setup, the maximum degree of the graph is $d_G = 2$, and the total spin is $\mathcal{M} = 1$. The initial and target vectors are given by $\boldsymbol{x}_0 = [1, 0, \dots, 0]^{\mathsf{T}}$ and $\boldsymbol{x}_{\tau} = [0, \dots, 0, 1]^{\mathsf{T}}$, respectively. The Wasserstein distance can also be analytically calculated as $\mathcal{W}_1(\boldsymbol{x}_0, \boldsymbol{x}_{\tau}) = N - 1$. Then, according to Eq. (S100), the minimum time required for transmitting the quantum state is lower bound as

$$\tau \ge \frac{N-1}{2\gamma}.\tag{S102}$$

Intriguingly, Eq. (S102) implies that it takes at least a time proportional to the distance between spins to reliably transfer a quantum state. The longer the distance, the more time is required. This is in agreement with the numerical result in Ref. [11], wherein an optimal control of the external magnetic field was used. It is noteworthy that this implication cannot be obtained from the conventional speed limits such as the Mandelstam–Tamm and Margolus–Levitin bounds.

S8. MARKOVIAN OPEN QUANTUM SYSTEMS

Here we derive a topological speed limit for Markovian open quantum systems. We consider a finite-dimensional quantum system that is weakly coupled to thermal reservoirs. The dynamics of the system's reduced density matrix is governed by the local Lindblad equation:

$$\dot{\varrho}_t = -i[H + V_t, \varrho_t] + \sum_k \mathcal{D}[L_k]\varrho_t, \qquad (S103)$$

where $H = \sum_{n} \epsilon_{n} |\epsilon_{n}\rangle\langle\epsilon_{n}|$ is the system Hamiltonian with $\epsilon_{n} \neq \epsilon_{m}$ for $n \neq m$, V_{t} is an external driving field, and $\{L_{k}\}$ are jump operators that characterize jumps between energy eigenstates with the same energy change ω_{k} (i.e., $[L_{k}, H] = \omega_{k}L_{k}$). We assume that the Hamiltonian has no energy degeneracy. Each jump operator L_{k} has a counterpart $L_{k'}$, which corresponds to the reversed jump and satisfies the local detailed balance condition $L_{k} = e^{s_{k}/2}L_{k'}^{\dagger}$. Here, $s_{k} = -s_{k'}$ denotes the change in environmental entropy due to the jump L_{k} . We consider the time evolution of the energetic population $x_{n}(t) = \langle \epsilon_{n} | \varrho_{t} | \epsilon_{n} \rangle$, which can be described by the following equation:

$$\dot{x}_n(t) = -i \langle \epsilon_n | [V_t, \varrho_t] | \epsilon_n \rangle + \sum_k \sum_{m(\neq n)} [r_{nm}^k x_m(t) - r_{mn}^{k'} x_n(t)]$$
(S104)

where $r_{mn}^k \coloneqq |\langle \epsilon_m | L_k | \epsilon_n \rangle|^2 \ge 0$ is the transition rate satisfying the local detailed balance $r_{mn}^k = e^{s_k} r_{nm}^{k'}$. Notice that Eq. (S104) can be expressed in the form of Eq. (1) with

$$f_n(t) = -i \langle \epsilon_n | [V_t, \varrho_t] | \epsilon_n \rangle, \qquad (S105)$$

$$f_{nm}(t) = \sum_{k} [r_{nm}^{k} x_{m}(t) - r_{mn}^{k'} x_{n}(t)].$$
(S106)

The graph $G(\mathcal{V}, \mathcal{E})$ is thus defined by $\mathcal{V} = \{1, \ldots, n, \ldots\}$ and $\mathcal{E} = \{\langle m, n \rangle | m < n, \exists k \text{ s.t. } r_{mn}^k \neq 0\}$. According to Eq. (5), we obtain the following speed limit:

$$\tau \ge \frac{\mathcal{W}_{1,\lambda}(\boldsymbol{x}_0, \boldsymbol{x}_{\tau})}{\langle \upsilon_{t,\lambda} \rangle_{\tau}}.$$
(S107)

Next, we derive an upper bound for $v_{t,\lambda}$. First, according to Prop. 4, we have

$$\sum_{n} |f_n(t)| \le \| [V_t, \varrho_t] \|_1 \le 2\Delta_{\varrho_t} V_t, \tag{S108}$$

where $||X||_p \coloneqq \operatorname{tr}\{|X|^p\}^{1/p}$ is the Schatten *p*-norm of operator X. Furthermore, according to Prop. 5, we can prove that

$$\sum_{(m,n)\in\mathcal{E}} |f_{mn}(t)| \le \frac{\sigma_t}{2} \Phi\left(\frac{\sigma_t}{2a_t}\right)^{-1}.$$
 (S109)

Here, $a_t \coloneqq \sum_k \operatorname{tr} \left\{ L_k \varrho_t L_k^{\dagger} \right\}$ is the dynamical activity rate, $\sigma_t \coloneqq \sigma_t^{\operatorname{pop}} + \sigma_t^{\operatorname{env}}$ is the total entropy production rate, $\sigma_t^{\operatorname{env}} \coloneqq \sum_k \operatorname{tr} \left\{ L_k \varrho_t L_k^{\dagger} \right\} s_k$ is the environmental entropy rate and $\sigma_t^{\operatorname{pop}} \coloneqq -\sum_n \dot{x}_n(t) \ln x_n(t) -$

 $i \sum_{n} \langle \epsilon_n | [V_t, \varrho_t] | \epsilon_n \rangle \ln x_n(t)$ is the sum of the Shannon entropy rate of the population distribution and the entropic change contributed by the external Hamiltonian V_t . Combining Eqs. (S108) and (S109) yields

$$\upsilon_{t,\lambda} \le 2\lambda \Delta_{\varrho_t} V_t + \frac{\sigma_t}{2} \Phi \left(\frac{\sigma_t}{2a_t}\right)^{-1}.$$
 (S110)

Consequently, we obtain the following thermodynamic speed limit:

$$\tau \ge \frac{\mathcal{W}_{1,\lambda}(\boldsymbol{x}_0, \boldsymbol{x}_{\tau})}{\langle 2\lambda \Delta_{\varrho_t} V_t + \sigma_t \Phi(\sigma_t/2a_t)^{-1}/2 \rangle_{\tau}}.$$
 (S111)

Proposition 4. The following inequality holds for arbitrary Hermitian matrix V and density matrix ϱ :

$$\sum_{n} |\langle \epsilon_{n} | [V, \varrho] | \epsilon_{n} \rangle| \le || [V, \varrho] ||_{1} \le 2\Delta_{\varrho} V.$$
 (S112)

Proof. Note that $\langle \epsilon_n | [V, \varrho] | \epsilon_n \rangle$ is pure imaginary. Let $U = \sum_n u_{nn} \epsilon_n$ be a diagonal matrix with elements $u_{nn} = \operatorname{sgn}(-i \langle \epsilon_n | [V, \varrho] | \epsilon_n \rangle)$. Here, $\operatorname{sgn}(x) = 1$ if $x \ge 0$ and $\operatorname{sgn}(x) = -1$ if x < 0. It is evident that all singular values of matrix U equal 1; thus, $||U||_{\infty} = 1$. According to von Neumann's trace inequality, the first inequality in Eq. (S112) can be proved as

$$\sum_{n} |\langle \epsilon_{n} | [V, \varrho] | \epsilon_{n} \rangle| = |\operatorname{tr} \{ U[V, \varrho] \}|$$

$$\leq ||U||_{\infty} ||[V, \varrho]||_{1}$$

$$= ||[V_{t}, \varrho_{t}]||_{1}. \quad (S113)$$

Next, we need only prove the second inequality in Eq. (S112). To this end, we follow the idea in Ref. [12]. Let $\rho = \sum_i p_i |i\rangle\langle i|$ be the spectral decomposition of the density matrix ρ acting on the *d*-dimensional Hilbert space \mathcal{H} . Let \mathcal{H}' be another copy of the Hilbert space \mathcal{H} with an orthonormal basis $\{|i'\rangle\}$. Then, $|\rho\rangle = \sum_i \sqrt{p_i} |i\rangle \otimes |i'\rangle \in \mathcal{H} \otimes \mathcal{H}'$ is the purification of the density matrix ρ_t . We also define $\tilde{V} \coloneqq V \otimes \mathbb{1} \in \mathcal{H} \otimes \mathcal{H}'$, where $\mathbb{1}$ is the identity matrix acting on the Hilbert space \mathcal{H}' . Let $\Phi(\cdot) = \operatorname{tr}_{\mathcal{H}'} \{\cdot\}$ denote the map of the partial trace with respect to \mathcal{H}' . By simple algebraic calculations, we can verify that

$$\Phi([\tilde{V}, |\varrho\rangle \langle \varrho|]) = [V, \varrho].$$
(S114)

Since the trace norm is contractive under a completely positive and trace-preserving map, we have

$$\begin{split} \| [V,\varrho] \|_{1} &= \| \Phi([V,|\varrho\rangle\langle\varrho|]) \|_{1} \\ &\leq \| [\tilde{V},|\varrho\rangle\langle\varrho|] \|_{1} \\ &= \Delta_{|\varrho\rangle\langle\varrho|} \tilde{V} \operatorname{tr} \left\{ \sqrt{|\varrho\rangle\langle\varrho| + |\varrho_{\perp}\rangle\langle\varrho_{\perp}|} \right\} \\ &= 2\Delta_{|\varrho\rangle\langle\varrho|} \tilde{V}. \end{split}$$
(S115)

Here, $|\varrho_{\perp}\rangle$ is a state orthogonal to $|\varrho\rangle$, given by

$$|\varrho_{\perp}\rangle \coloneqq \frac{\left(V - \langle \varrho | V | \varrho \rangle\right) |\varrho\rangle}{\Delta_{|\varrho\rangle\langle\varrho|} \tilde{V}}.$$
 (S116)

In addition, the energy fluctuation $\Delta_{|\varrho \chi_{\ell}|} \tilde{V}$ can be simplified as follows:

$$\Delta_{|\varrho\rangle\langle\varrho|}\tilde{V} = \left(\langle\varrho|\tilde{V}^{2}|\varrho\rangle - \langle\varrho|\tilde{V}|\varrho\rangle^{2}\right)^{1/2}$$
$$= \left(\operatorname{tr}\left\{V^{2}\varrho\right\} - \operatorname{tr}\left\{V\varrho\right\}^{2}\right)^{1/2}$$
$$= \Delta_{\varrho}V.$$
(S117)

Combining Eqs. (S115) and (S117) yields the desired second inequality in Eq. (S112). \Box

Proposition 5. The following inequality holds:

$$\sum_{(m,n)\in\mathcal{E}} \left| \sum_{k} \left[r_{nm}^{k} x_{m}(t) - r_{mn}^{k'} x_{n}(t) \right] \right| \le \frac{\sigma_{t}}{2} \Phi \left(\frac{\sigma_{t}}{2a_{t}} \right)^{-1},$$
(S118)

where $\Phi(x)$ is the inverse function of $x \tanh(x)$.

Proof. Note that $[L_k^{\dagger}L_k, H] = 0$ and $\langle \epsilon_n | L_k^{\dagger}L_k | \epsilon_m \rangle = 0$ for $m \neq n$. Therefore,

$$\operatorname{tr}\left\{L_{k}\varrho_{t}L_{k}^{\dagger}\right\} = \sum_{m} \left\langle\epsilon_{m}|\varrho_{t}|\epsilon_{m}\right\rangle \left\langle\epsilon_{m}|L_{k}^{\dagger}L_{k}|\epsilon_{m}\right\rangle$$
$$= \sum_{m} x_{m}(t)\sum_{n} \left\langle\epsilon_{m}|L_{k}^{\dagger}|\epsilon_{n}\right\rangle \left\langle\epsilon_{n}|L_{k}|\epsilon_{m}\right\rangle$$
$$= \sum_{m,n} r_{nm}^{k} x_{m}(t).$$
(S119)

We first calculate σ_t^{env} as follows:

$$\sigma_{t}^{\text{env}} = \sum_{k} \operatorname{tr} \{ L_{k} \varrho_{t} L_{k}^{\dagger} \} s_{k}$$

$$= \frac{1}{2} \sum_{k} (\operatorname{tr} \{ L_{k} \varrho_{t} L_{k}^{\dagger} \} - \operatorname{tr} \{ L_{k'} \varrho_{t} L_{k'}^{\dagger} \}) s_{k}$$

$$= \frac{1}{2} \sum_{k} \sum_{m,n} [r_{nm}^{k} x_{m}(t) - r_{mn}^{k'} x_{n}(t)] s_{k}$$

$$= \frac{1}{2} \sum_{k} \sum_{m,n} [r_{nm}^{k} x_{m}(t) - r_{mn}^{k'} x_{n}(t)] \ln \frac{r_{nm}^{k}}{r_{mn}^{k'}}.$$
 (S120)

Consequently, we can calculate

$$\sigma_{t} = \sigma_{t}^{\text{pop}} + \sigma_{t}^{\text{env}}$$

$$= -\sum_{k} \sum_{m\neq n} [r_{nm}^{k} x_{m}(t) - r_{mn}^{k'} x_{n}(t)] \ln x_{n}(t)$$

$$+ \frac{1}{2} \sum_{k} \sum_{m,n} [r_{nm}^{k} x_{m}(t) - r_{mn}^{k'} x_{n}(t)] \ln \frac{r_{nm}^{k}}{r_{mn}^{k'}}$$

$$= \frac{1}{2} \sum_{k} \sum_{m,n} [r_{nm}^{k} x_{m}(t) - r_{mn}^{k'} x_{n}(t)] \ln \frac{x_{m}(t)}{x_{n}(t)}$$

$$+ \frac{1}{2} \sum_{k} \sum_{m,n} [r_{nm}^{k} x_{m}(t) - r_{mn}^{k'} x_{n}(t)] \ln \frac{r_{nm}^{k}}{r_{mn}^{k'}}$$

$$= \frac{1}{2} \sum_{k} \sum_{m,n} [r_{nm}^{k} x_{m}(t) - r_{mn}^{k'} x_{n}(t)] \ln \frac{r_{nm}^{k} x_{m}(t)}{r_{mn}^{k'} x_{n}(t)}$$

$$= \sum_{k} \sum_{m,n} \sigma_{nm}^{k}(t). \qquad (S121)$$

Similarly, the dynamical activity rate can also be calculated as

$$a_{t} = \sum_{k} \operatorname{tr} \left\{ L_{k} \varrho_{t} L_{k}^{\dagger} \right\}$$
$$= \frac{1}{2} \sum_{k} \left[\operatorname{tr} \left\{ L_{k} \varrho_{t} L_{k}^{\dagger} \right\} + \operatorname{tr} \left\{ L_{k'} \varrho_{t} L_{k'}^{\dagger} \right\} \right]$$
$$= \frac{1}{2} \sum_{k} \sum_{m,n} \left[r_{nm}^{k} x_{m}(t) + r_{mn}^{k'} x_{n}(t) \right]$$
$$\Rightarrow \sum_{k} \sum_{m,n} a_{nm}^{k}(t).$$
(S122)

By applying the triangle inequality and Jensen's inequality, we obtain the desired inequality (S118) as

$$\sum_{\langle m,n\rangle\in\mathcal{E}} \left| \sum_{k} \left[r_{nm}^{k} x_{m}(t) - r_{mn}^{k'} x_{n}(t) \right] \right|$$

$$\leq \frac{1}{2} \sum_{k} \sum_{m,n} \left| r_{nm}^{k} x_{m}(t) - r_{mn}^{k'} x_{n}(t) \right|$$

$$= \sum_{k} \sum_{m,n} \frac{\sigma_{nm}^{k}(t)}{2} \Phi \left(\frac{\sigma_{nm}^{k}(t)}{2a_{nm}^{k}(t)} \right)^{-1}$$

$$\leq \frac{\sigma_{t}}{2} \Phi \left(\frac{\sigma_{t}}{2a_{t}} \right)^{-1}. \tag{S123}$$

- B. Piccoli and F. Rossi, Generalized Wasserstein distance and its application to transport equations with source, Arch. Ration. Mech. Anal. 211, 335 (2013).
- [2] T. Van Vu and K. Saito, Numerical calculation of Wasserstein distance, https://github.com/tvanvu/ WassersteinDistance (2022).
- [3] R. Hamazaki, Speed limits for macroscopic transitions, PRX Quantum 3, 020319 (2022).
- [4] C. Villani, Optimal Transport: Old and New (Springer,

Berlin, Heidelberg, 2008).

- [5] K. Yoshimura and S. Ito, Thermodynamic uncertainty relation and thermodynamic speed limit in deterministic chemical reaction networks, Phys. Rev. Lett. **127**, 160601 (2021).
- [6] A. Uhlmann, An energy dispersion estimate, Phys. Lett. A 161, 329 (1992).
- [7] M. Fishman, S. R. White, and E. M. Stoudenmire, The ITensor Software Library for Tensor Network Calcula-

tions, SciPost Phys. Codebases , 4 (2022).

- [8] M. Yang and S. R. White, Time-dependent variational principle with ancillary Krylov subspace, Phys. Rev. B 102, 094315 (2020).
- [9] A. Didi and E. Barkai, Measurement-induced quantum walks, Phys. Rev. E 105, 054108 (2022).
- [10] S. Bose, Quantum communication through an unmodulated spin chain, Phys. Rev. Lett. 91, 207901 (2003).
- [11] M. Murphy, S. Montangero, V. Giovannetti, and T. Calarco, Communication at the quantum speed limit along a spin chain, Phys. Rev. A 82, 022318 (2010).
- [12] K. Funo, N. Shiraishi, and K. Saito, Speed limit for open quantum systems, New J. Phys. 21, 013006 (2019).