

Universal description of dissipative Tomonaga-Luttinger liquids with $SU(N)$ spin symmetry: Exact spectrum and critical exponents

Kazuki Yamamoto* and Norio Kawakami

Department of Physics, Kyoto University, Kyoto 606-8502, Japan

(Dated: January 11, 2023)

Universal scaling relations for dissipative Tomonaga-Luttinger (TL) liquids with $SU(N)$ spin symmetry are obtained for both fermions and bosons, by using asymptotic Bethe-ansatz solutions and conformal field theory (CFT) in one-dimensional non-Hermitian quantum many-body systems with $SU(N)$ symmetry. We uncover that the spectrum of dissipative TL liquids with $SU(N)$ spin symmetry is described by the sum of one charge mode characterized by a complex generalization of $c = 1$ U(1) Gaussian CFT, and $N - 1$ spin modes characterized by level-1 $SU(N)$ Kac-Moody algebra with the conformal anomaly $c = N - 1$, and thereby dissipation only affects the charge mode as a result of spin-charge separation in one-dimensional non-Hermitian quantum systems. The derivation is based on a complex generalization of Haldane's ideal-gas description, which is implemented by the $SU(N)$ Calogero-Sutherland model with inverse-square long-range interactions.

I. INTRODUCTION

Recent advances in the study of ultracold atoms have opened a new arena to investigate open quantum many-body systems, where a variety of unique phenomena that have no counterpart in isolated quantum systems occur [1–10]. Experimentally, high controllability of ultracold atoms has facilitated investigations of nonequilibrium quantum dynamics in open quantum systems, where strong coupling to the environment plays a vital role [11–19]. In particular, non-Hermitian (NH) Hamiltonians have shed new light on investigations of dissipation-induced quantum many-body physics [20–22]; examples include loss-induced quantum phase transitions [23–25], measurement-induced entanglement dynamics [26–34], NH quantum phases as a result of postselections of measurement outcomes by means of quantum-gas microscopy [35–39]. Notably, one-dimensional (1D) NH quantum many-body systems show intriguing quantum critical phenomena induced by dissipation; e.g., correlation functions of dissipative Tomonaga-Luttinger (TL) liquids show unique critical behavior characterized by two TL parameters [36, 39], and exceptional points, where the effective Hamiltonian cannot be diagonalized, cause anomalous singularities that accompany the divergence of correlation length [40]. These studies have shown that dissipation drastically alters the universal properties of quantum many-body phenomena in isolated systems.

Another important aspect which has attracted broad interest in ultracold atoms is a multicomponent generalization of many-body phenomena, which show rich quantum phases as a result of strong correlation [41–50]. In particular, multicomponent fermions have been actively investigated in experiments over the last decade by controlling internal (nuclear) spin degrees of freedom, providing unique opportunities for quantum simulations of many-body systems. The milestone experiments have realized observations of antiferromagnetic correlations in $SU(2)$ Fermi-Hubbard models with degenerate ^{40}K and ^6Li [51–56], and exotic quantum phases stemming from $SU(N)$ ($N > 2$) spin symmetry with ultracold

alkaline-earth-like atoms such as ^{173}Yb and ^{87}Sr have recently been reported [57–63]. Multicomponent bosons are also successfully loaded into an optical lattice, where hyperfine states of bosonic atoms like ^{87}Rb can be used, offering the possibility to explore novel highly entangled many-body states [64, 65]. Moreover, it is worth noting that physics with $SU(N)$ symmetry in ultracold atoms has recently extended its research area to nonequilibrium quantum systems as represented by $SU(N)$ Hubbard models with two-body loss [10, 16, 23, 38, 40, 66, 67]. Thus, it is natural to consider how the universal properties of many-body physics with $SU(N)$ symmetry are affected by dissipation. However, it is a highly nontrivial problem to identify the universality class of dissipative quantum many-body systems with internal degrees of freedom.

To study universal properties of strongly-correlated systems, 1D critical systems have been a subject of intense research in condensed matter physics, because they realize TL liquids, which present a general description of low-energy quantum many-body phenomena [68–71]. Importantly, conformal invariance in 1D critical systems has brought about many valuable insights into many-body quantum systems [72–75], where TL liquids are characterized by U(1) Gaussian conformal field theory (CFT) with the central charge $c = 1$. To access the universality class of TL liquids, Haldane has proposed that quantum models with inverse-square long-range interactions, which were initially introduced by Calogero and Sutherland [76–80], give a unified understanding of 1D quantum critical phenomena [81–85]. He has demonstrated that the model gives an ideal-gas description in one dimension, providing a variant notion of fractional statistics, dubbed fractional exclusion statistics [86, 87], which has an intimate relation to the fractional quantum Hall effect [88, 89]. For example, semion, a particle with statistical $1/2$ interactions, appears in the $S = 1/2$ Heisenberg spin chain with inverse-square exchange [81–83]. Studies with inverse-square interactions have been successfully generalized to multicomponent systems, and they have demonstrated the universal properties of TL liquids with $SU(N)$ spin symmetry [90–95].

In this paper, we elucidate the universal properties of dissipative TL liquids with $SU(N)$ spin symmetry for both fermions and bosons in one dimension, based on a complex

* yamamoto.kazuki.72n@st.kyoto-u.ac.jp

generalization of Haldane's ideal-gas description with the NH SU(N) Calogero-Sutherland (CS) model. As a main result, we obtain critical exponents describing general NH quantum critical systems with SU(N) symmetry. By analyzing the NH CS model, we obtain the asymptotic Bethe-ansatz (ABA) solutions and the finite-size scaling formula in CFT. We demonstrate that the spectrum of dissipative TL liquids with SU(N) spin symmetry is described by one charge mode characterized by a complex generalization of $c = 1$ U(1) Gaussian CFT, and $N-1$ spin modes characterized by level-1 SU(N) Kac-Moody algebra with $c = N - 1$. Since the spin modes are protected by SU(N) symmetry, dissipation only affects the charge mode as a result of spin-charge separation in 1D NH quantum systems. Although our results are derived from specific integral models, universal scaling relations obtained can be applied to a wide range of NH quantum critical systems which are experimentally relevant, e.g., dissipative TL liquids with SU(2) spin symmetry describe the critical properties of the Fermi-Hubbard model with a complex-valued interaction.

The rest of this paper is organized as follows. First, in Sec. II, we summarize the main results of the universal scaling relations in NH quantum critical systems for SU(N) fermions and bosons, including SU(2) fermions as a special case. In the subsequent sections, we explain the detailed derivations of the universal properties of dissipative TL liquids with and without internal symmetry, based on a complex generalization of Haldane's ideal-gas description. In Sec. III, we analyze the NH CS model based on the ABA solutions and the finite-size scaling analysis in CFT. We generalize the results to multicomponent systems by analyzing NH SU(N) CS model in Sec. IV. We finally conclude with a summary and outlook in Sec. V.

II. MAIN RESULTS: UNIVERSAL SCALING RELATIONS

In this section, we present a summary of the main results of this paper, that is, universal scaling relations for dissipative TL liquids with SU(N) spin symmetry. The class of models that we want to consider throughout this paper is the critical system described by a NH Hamiltonian with SU(N) symmetry. We note that SU(N) symmetry is imposed not only on the original Hermitian Hamiltonian but also on dissipation, resulting in the whole SU(N) symmetry of the NH Hamiltonian. Although the scaling relations are obtained by an ideal-gas approach based on specific integral models as detailed in the following sections III and IV, they are universal and applicable to generic 1D dissipative many-body systems.

Before going to the main results of scaling relations, we introduce important concepts of two-types of correlation functions that appear in NH systems. In NH systems that are described by the effective Hamiltonian H_{eff} , a right eigenstate, which is defined by $H_{\text{eff}}|\Psi^R\rangle = E|\Psi^R\rangle$, and a left eigenstate, which is defined by $H_{\text{eff}}^\dagger|\Psi^L\rangle = E^*|\Psi^L\rangle$, are different from each other. Therefore, two types of correlation functions can emerge according to whether the right or left eigenstate is assigned to the bra vector in the expectation value. The first type is a mathematical generalization of correlation functions to NH systems defined by ${}_L\langle\cdots\rangle_R \equiv \langle\Psi_g^L|\cdots|\Psi_g^R\rangle/\langle\Psi_g^L|\Psi_g^R\rangle$,

where $|\Psi_g^L\rangle$ and $|\Psi_g^R\rangle$ are the left and right ground states (in the sense of the real part of the energy) of H_{eff} , respectively. This type of correlation functions is calculated through path integrals [23, 39], and has been shown that it is directly related to a complex extension of CFT [39]. The second type is defined by ${}_R\langle\cdots\rangle_R \equiv \langle\Psi_g^R|\cdots|\Psi_g^R\rangle/\langle\Psi_g^R|\Psi_g^R\rangle$, which is obtained in the postselected sector with no loss events as follows. First, the dynamics of the quantum state in such sectors is described by the Schrödinger equation $i\partial_t|\psi\rangle = H_{\text{eff}}|\psi\rangle$. Then, with the use of the postselected ground state $|\psi\rangle = |\Psi_g^R\rangle$, we obtain a standard quantum-mechanical expectation value ${}_R\langle\cdots\rangle_R$, which corresponds to an experimentally measurable physical quantity [36, 39]. We call the correlation functions ${}_L\langle\cdots\rangle_R$ and ${}_R\langle\cdots\rangle_R$ the biorthogonal correlation function and the right-state correlation function, respectively. We note that the subscripts L and R for the brackets are not related to the left and right branches of TL liquids. In the following two subsections, we give universal scaling relations for right-state correlation functions, which are physical observables and relevant to cold-atom experiments.

A. Critical exponents for fermions described by dissipative TL liquids with SU(2) spin symmetry

We first focus on fermions described by dissipative TL liquids with SU(2) spin symmetry separately, since such SU(2) models have been by far the best studied until now both in solid state systems and cold atom systems. In particular, SU(2) fermions are relevant to Fermi-Hubbard models, which has been a subject of intense study in condensed matter physics.

Now, we give universal scaling relations for dissipative TL liquids with SU(2) spin symmetry. In NH fermionic quantum critical systems with SU(2) symmetry (with zero magnetic field), we obtain the long-distance behavior of the right-state charge-density correlator as

$${}_R\langle\rho(x)\rho(0)\rangle_R \simeq A_1 \cos(4k_F x)x^{-\beta_1} + A_2 \cos(2k_F x)x^{-\beta_2} + A_0 \frac{1}{x^2}, \quad (1)$$

where $\rho(x) = \sum_\sigma c_\sigma^\dagger(x)c_\sigma(x)$, A_j is a correlation amplitude, and the critical exponents β_1 and β_2 are given by

$$\beta_1 = 4K_\rho^\phi, \quad (2)$$

$$\beta_2 = 1 + K_\rho^\phi. \quad (3)$$

As the ground state is given by the SU(2) singlet, we have defined the Fermi momentum $k_F = \pi n/2$ with the density $n = M/L$, where M is the total number of fermions and L is the circumference of the system imposed by periodic boundary conditions. Importantly, the exponent K_ρ^ϕ is obtained by the complex-valued TL parameter \tilde{K}_ρ as [39]

$$\frac{1}{K_\rho^\phi} = \text{Re} \left[\frac{1}{\tilde{K}_\rho} \right], \quad (4)$$

where \tilde{K}_ρ is defined through the $4k_F$ oscillating piece in the biorthogonal correlation function ${}_L\langle\rho(x)\rho(0)\rangle_R \simeq \cos(4k_F x)x^{-\tilde{\beta}_1}$ as

$$\tilde{K}_\rho = \frac{\tilde{\beta}_1}{4}. \quad (5)$$

We emphasize that the TL parameter is affected by dissipation, making it different from the one for the standard Hermitian TL liquid, and this fact leads to the emergence of two kinds of real TL exponents, which characterize experimentally measurable quantities as shown below. Here and henceforth, we use the symbol \tilde{A} to emphasize that the quantity \tilde{A} is complex-valued.

For the right-state fermion correlator, the long-distance behavior is written as

$${}_R\langle c_\sigma^\dagger(x)c_\sigma(0)\rangle_R \simeq C_1 \cos(k_F x)x^{-\eta_F}, \quad (6)$$

where C_1 is a correlation amplitude and the critical exponent η_F is given by

$$\eta_F = \frac{1}{2} + \frac{1}{4K_\rho^\theta} + \frac{K_\rho^\phi}{4}. \quad (7)$$

The exponent K_ρ^θ is obtained from the complex-valued TL parameter \tilde{K}_ρ as [39]

$$K_\rho^\theta = \text{Re}[\tilde{K}_\rho]. \quad (8)$$

We note that Eq. (6) does not depend on spin indices for the SU(2) singlet ground state.

We emphasize that the scaling relations (2), (3), and (7) are universal, and describe the general critical properties of fermions characterized by dissipative TL liquids with SU(2) spin symmetry. Eqs. (4) and (8) also highlight the universal properties of the U(1) charge part of dissipative TL liquids, which are characterized by the complex-valued TL parameter [39].

One of the prototypical dissipative models that have SU(2) symmetry is given by the NH Fermi-Hubbard model [10, 23, 38, 40, 67]

$$H_{\text{eff}}^{\text{Hubbard}} = -t \sum_{j\sigma} (c_{j\sigma}^\dagger c_{j+1\sigma} + \text{H.c.}) + \tilde{U} \sum_j n_{j\uparrow} n_{j\downarrow}, \quad (9)$$

where \tilde{U} is a complex-valued interaction as a result of dissipation, t is a hopping parameter, σ denotes up or down spin of fermions, $c_{j\sigma}$ is the annihilation operator of a spin- σ fermion at site j , and $n_{j\sigma} \equiv c_{j\sigma}^\dagger c_{j\sigma}$. Such model is realized by introducing two-body loss due to inelastic collisions between fermions as observed in cold-atom experiments [16, 19] e.g. with the use of photoassociation lasers. In the following, we consider dissipation that gives complex-valued interactions, and therefore other types of dissipation that cause e.g. asymmetric hopping of particles are not addressed in our study [96]. We note that, in the previous studies that deal with the NH Fermi-Hubbard model, such as superfluid states with attractive interactions [10, 23] and Mott insulators with repulsive

interactions [38, 40] have been considered, but the critical properties of metallic phases have not been investigated so far.

We demonstrate that critical properties of such NH SU(2) quantum models are described by dissipative TL liquids with SU(2) spin symmetry discussed in this subsection. In the NH Fermi-Hubbard model (9), critical properties of metallic phases are described by dissipative TL liquids with U(1)×SU(2) symmetry, and the universal scaling relations are given by Eqs. (2), (3), and (7) by taking the continuum limit as $c_\sigma(x) = c_{j\sigma}/\sqrt{a}$ with the lattice constant a .

B. Critical exponents for fermions and bosons described by dissipative TL liquids with SU(N) spin symmetry

In this subsection, motivated by the recent progress in SU(N) quantum phenomena in ultracold atoms, we summarize a generalization of universal scaling relations in NH quantum critical systems to SU(N) symmetric cases. We give universal scaling relations for both fermions and bosons described by dissipative TL liquids with SU(N) spin symmetry. The right-state charge-density correlator is the same for both fermions and bosons, and the long-distance behavior reads

$${}_R\langle\rho(x)\rho(0)\rangle_R \simeq \sum_{j=1}^N A_j \cos[2(N-j+1)k_F x]x^{-\beta_j} + \frac{A_0}{x^2}, \quad (10)$$

where A_j is a correlation amplitude, and the critical exponent β_j corresponding to the $2(N-j+1)k_F$ oscillation is given by

$$\beta_j = \frac{2(N-j+1)(j-1)}{N} + \frac{2(N-j+1)^2}{N} K_\rho^\phi. \quad (11)$$

Here, we have defined the Fermi momentum $k_F = \pi n/N$, which reflects the SU(N) singlet ground state. As in the SU(2) case, the critical exponent K_ρ^ϕ is obtained by the complex-valued TL parameter \tilde{K}_ρ through Eq. (4). In this case, \tilde{K}_ρ is generalized to the system with SU(N) symmetry, resulting in

$$\tilde{K}_\rho = \frac{\tilde{\beta}_1}{2N}, \quad (12)$$

which corresponds to the $2Nk_F$ oscillating piece of the biorthogonal correlation function ${}_L\langle\rho(x)\rho(0)\rangle_R \simeq \cos(2Nk_F x)x^{-\tilde{\beta}_1}$.

On the other hand, the single-particle correlator is different between fermions and bosons. For fermions described by dissipative TL liquids with SU(N) spin symmetry, we obtain the same correlator as Eq. (6), where the critical exponent η_F is generalized to SU(N) symmetric cases;

$$\eta_F = \frac{N-1}{N} + \frac{1}{2NK_\rho^\theta} + \frac{K_\rho^\phi}{2N}. \quad (13)$$

We note that the exponent K_ρ^θ is obtained from Eq. (8) with Eq. (12), and the fermion correlator (6) is independent of spin

indices for the $SU(N)$ singlet ground state as in the $SU(2)$ symmetric cases. For the right-state boson correlator, we obtain the long-distance behavior as

$${}_R\langle b_\sigma^\dagger(x)b_\sigma(0)\rangle_R \simeq B_1 x^{-\eta_B}, \quad (14)$$

where $b_\sigma(x)$ is the annihilation operator of bosons, B_1 is a correlation amplitude, and the critical exponent η_B is given by

$$\eta_B = \frac{N-1}{2N} + \frac{1}{2NK_\rho^\theta}. \quad (15)$$

We note that the bosonic correlator (14) does not depend on the spin σ as in the case of fermions. It is worth noting that the bosonic correlator (14) is not accompanied by the k_F oscillation that appears in the fermion case because of Bose statistics (see Sec. III for details).

We emphasize that the scaling relations (11), (13), and (15) are universal, and characterize the general critical properties of fermions and bosons described by dissipative TL liquids with $SU(N)$ spin symmetry, where the critical theory consists of one charge mode characterized by a complex generalization of $c = 1$ U(1) Gaussian CFT [39], and $N-1$ spin modes characterized by $c = N-1$ level-1 $SU(N)$ Kac-Moody algebra (see Sec. IV for details). Importantly, the algebraic structure of the spin sectors is fixed to the real one as a result of symmetry enhancement. In Sec. IV, we give the detailed derivation which includes the ABA solutions and the finite-size scaling analysis in CFT based on a complex generalization of Haldane's ideal-gas description.

III. DISSIPATIVE TOMONAGA-LUTTINGER LIQUIDS WITHOUT INTERNAL SYMMETRY

In this section, we analyze the NH CS model without internal symmetry, based on a complex generalization of Haldane's ideal-gas description, which is embodied by the ABA solution. We will see that the ideal-gas description captures universal properties of single-component dissipative TL liquids.

A. Model

We consider a 1D long-range interacting NH quantum system with the circumference L described by

$$H_{\text{eff}} = -\sum_{j=1}^M \frac{\partial^2}{\partial x_j^2} + \sum_{j>l} V(x_j - x_l), \quad (16)$$

$$V(x) = \tilde{g} \sum_{n=-\infty}^{\infty} (x + nL)^{-2} = \frac{\tilde{g}\pi^2}{L^2} \left[\sin\left(\frac{\pi x}{L}\right) \right]^{-2}, \quad (17)$$

where M is the number of particles, and \tilde{g} is the dimensionless complex parameter of the inverse-square long-range interaction. This model is a generalization of the CS model

[76–80, 84] to NH quantum many-body systems [97]. Models with inverse-square interactions have been investigated intensively to date, for example, in a quantum spin chain (Haldane-Shastry model [81–83]), in a supersymmetric $t - J$ model [91, 98, 99], and in a continuum model interacting by the inverse-square potential (CS model). These models are known to be integrable [100–102], for which the ground-state wave function is given by a Jastrow form, i.e., products of two-body wave functions [76–80].

An important aspect of this class of integrable models is that they can describe the essential properties of 1D quantum critical systems as an ideal gas, as demonstrated by Haldane [81, 83, 86]. To explain the essence of the idea, let us start by analyzing the ground state of the NH CS model (16). A little algebra leads to a complex generalization of the ground-state wave function of the Jastrow form (see Appendix A for detailed calculations)

$$\Psi_g = \prod_{j>l} \left| \sin \frac{\pi(x_j - x_l)}{L} \right|^{\tilde{\lambda}-s} \left(\sin \frac{\pi(x_j - x_l)}{L} \right)^s, \quad (18)$$

where $x_j > x_l$, $\tilde{\lambda} = [(1 + 2\tilde{g})^{\frac{1}{2}} + 1]/2$, $\text{Re}[\tilde{\lambda}] \geq 1/2$ is assumed, and we have introduced $s = 1$ for fermions and $s = 0$ for bosons, respectively. The wave function (18) reduces to the one for free fermions or hard-core bosons in the limit $\tilde{\lambda} \rightarrow 1$. The Jastrow-type wave function indicates that the two-body scattering is essential to the system. This observation motivated Sutherland to propose the ABA method [77–80], which states that the many-body scattering is decomposed into the product of two-body scatterings as a result of the integrability in spite of long-range nature of interactions [103]. The ABA solution provides a concrete way to implement the ideal-gas description, as described below.

B. Asymptotic Bethe ansatz: Ideal-gas description

Now, we derive the ABA solution of the NH CS model. The phase shift as a result of the two-body scattering is read off from the Jastrow wave function (18) as [84]

$$\theta(k) = \pi(\tilde{\lambda} - 1)\text{sgn}^*(k), \quad (19)$$

where we have defined the function $\text{sgn}^*(k)$ for the complex quasimomentum $k = \alpha e^{i\xi}$ ($\alpha \in \mathbb{R}$, $-\pi/2 < \xi < \pi/2$) as $\text{sgn}^*(k) = \text{sgn}(\alpha)$ (see Fig. 1). There is an important feature in Eq. (19): the phase shift is characterized by a single k -independent parameter $\tilde{\lambda}$ and a step-function in k space, which is contrasted to ordinary integrable models for which the corresponding phase shift is k -dependent. We therefore see that the above two-body phase shift incorporates the interaction effect via a level repulsion parameter $\tilde{\lambda}$ in a way similar to ideal gases such as free fermions or hard-core bosons [81, 83, 86].

The philosophy of the ABA (ideal-gas description) lies in the fact that the two-body phase shift (19) exactly describes the whole spectrum of the long-range interacting system (16)




FIG. 1. Distribution of the quasimomentum k_j in the ground state of long-range interacting systems (16). The quasimomentum k_j is equally spaced along the straight line with the interval $2\pi\tilde{\lambda}/L$, and $\pm Q$, which are called the quasi Fermi points, denote the edges of the quasimomentum distribution. The generalized signature between k_m and k_l gives the phase shift $\theta(k_m - k_l)$ (see text).

for arbitrary densities due to the integrability [103]. Based on this idea, we obtain the ABA equation as

$$k_j L = 2\pi I_j + \sum_{i=1}^M \theta(k_j - k_i), \quad (20)$$

where

$$I_j : \text{integer}, \quad (21)$$

for fermions, and

$$I_j = \frac{M+1}{2} \pmod{1}, \quad (22)$$

for bosons, respectively. It is worth noting that both fermions and bosons are described by the transcendental equation (20), and only the selection rule of the quantum number I_j distinguishes their statistics. By using the quasimomentum k_j , we obtain the energy as $E = \sum k_j^2$, and the momentum as $P = \sum k_j$. We note that, in NH systems, integrability has been shown to hold by explicitly constructing the Bethe equation for models having e.g. asymmetric hoppings and complex-valued interactions [20, 39, 40, 96, 104, 105].

We consider the ground state and obtain the quasimomentum distribution. The quantum numbers for the ground state $\{I_j\}$ are successive integers (half-odd integers), which are distributed symmetrically around the origin. Then, we obtain the quasimomentum as

$$k_j L = \pi\tilde{\lambda}(2j - M - 1) \quad (23)$$

for odd- M fermions and bosons by using $I_j = (2j - M - 1)/2$, and

$$k_j L = \pi\tilde{\lambda}(2j - M - 1) + \pi \quad (24)$$

for even- M fermions with the use of $I_j = (2j - M - 1)/2 + 1/2$. We see from Eqs. (23) and (24) that the quasimomentum

is equally spaced along the straight line in the complex plane as depicted in Fig. 1. This is nothing but the fact that each particle feels the complex-valued repulsion due to a step function in the complex phase shift given in Eq. (19). As shown in the following, the constant shift $1/2$ in the even- M fermion's quantum number I_j leads to the characteristic selection rule for fermions, which convey the current accompanied by the change of the particle number. It is worth noting that bosons do not know the Fermi point as a result of Bose statistics, and this fact causes the difference between the single-particle correlators of fermions and bosons.

C. Velocity of excitations

Next, we calculate the velocity of excitations that controls the bulk quantities, for which we first obtain the dressed energy and the distribution function in the thermodynamic limit. We note that the velocity of excitations becomes complex reflecting the effect of dissipation in open quantum systems. Here, we consider an elementary excitation from the ground state. By inserting a hole into the ground-state distribution of the quasimomentum, the dressed energy $\epsilon(k)$ is calculated by using the ABA equation (20) as

$$\epsilon(k) = k^2 - \tilde{\mu} - (\tilde{\lambda} - 1) \int_{-Q}^Q dk' \delta(k - k') \epsilon(k'), \quad (25)$$

which leads to

$$\epsilon(k) = \begin{cases} k^2 - \tilde{\mu} & (|k| > |Q|), \\ \frac{1}{\tilde{\lambda}}(k^2 - \tilde{\mu}) & (|k| < |Q|), \end{cases} \quad (26)$$

where $Q = \pi n \tilde{\lambda}$ with the density $n = M/L$. Here, we have introduced the delta function $\delta(k)$ that satisfies $\int_C dk \delta(k - k_0) f(k) = f(k_0)$ for an arbitrary function $f(k)$ on the contour C , which is the straight line shown in Fig. 1. As the dressed energy satisfies $\epsilon(\pm Q) = 0$ at the quasi Fermi points $\pm Q$, the complex-valued chemical potential is given by $\tilde{\mu} = \pi^2 n^2 \tilde{\lambda}^2$. Similarly, the distribution function in the thermodynamic limit $\sigma(k|\pm Q)$ is calculated as

$$\sigma(k|\pm Q) = \frac{1}{2\pi} - (\tilde{\lambda} - 1) \int_{-Q}^Q dk' \delta(k - k') \sigma(k'|\pm Q), \quad (27)$$

which leads to

$$\sigma(k|\pm Q) = \begin{cases} \frac{1}{2\pi} & (|k| > |Q|), \\ \frac{1}{2\pi\tilde{\lambda}} & (|k| < |Q|). \end{cases} \quad (28)$$

Finally, by using Eqs. (26) and (28), we obtain the velocity of excitations as

$$\tilde{v} = \frac{\epsilon'(Q)}{2\pi\sigma(Q|\pm Q)} = 2\pi\tilde{\lambda}n, \quad (29)$$

where $\epsilon'(k)$ denotes the derivative of $\epsilon(k)$ with respect to k . We emphasize that the velocity of excitations (29) becomes complex as a result of dissipation.

D. Finite-size scaling analysis in CFT

In this subsection, we perform the finite-size scaling analysis in CFT [106–111] by explicitly calculating the ground-state energy, the excitation energy, and the momentum transfer for both fermions and bosons [84]. First, we calculate the ground-state energy. By substituting Eqs. (23) and (24) into $E = \sum k_j^2$, we arrive at the ground state energy for fermions and bosons as

$$E_0 = \frac{\pi^2 \tilde{\lambda}^2}{L^2} \sum_{j=1}^M (2j - M - 1)^2 = L\epsilon_0 - \frac{\pi \tilde{v} \tilde{\lambda}}{6L}, \quad (30)$$

where we have introduced $\epsilon_0 = \pi^2 \tilde{\lambda}^2 n^3 / 3$, and the energy becomes complex as the system decays due to dissipation. We note that the calculation in Eq. (30) is exact, and we have shifted the ground-state energy for even- M fermions by $\pi^2 n / L$ because this term does not affect the universality. One notices that Eq. (30) is peculiar since the central charge could be given by $c = \tilde{\lambda}$, hence changes continuously in the complex plane. However, this artifact solely comes from the long-range nature of the interaction in a finite-size system, where we have applied the cylindrical geometry to the long-range interacting system [84], and the universality class is described by the central charge $c = 1$ as we see from the conformal dimensions below.

Next, we calculate the excitation energy from the ground state ΔE and the associated momentum transfer P . There are two types of excitations characterized by the quantum numbers ΔM and ΔD other than the particle-hole excitations. ΔM denotes the change of the particle number and ΔD describes the number of particles which move from the left Fermi point to the right one accompanied by the large momentum transfer P . Then, the excitation energy ΔE is calculated as

$$\begin{aligned} \Delta E &= \frac{\pi^2 \tilde{\lambda}^2}{L^2} \left[\sum_{j=1}^{M+\Delta M} \left(2j - M - \Delta M - 1 + \frac{2\Delta D}{\tilde{\lambda}} \right)^2 - \sum_{j=1}^M (2j - M - 1)^2 \right] \\ &= \tilde{\mu} \Delta M + \frac{2\pi \tilde{v}}{L} \left[\frac{\tilde{\lambda}}{4} (\Delta M)^2 + \frac{1}{\tilde{\lambda}} (\Delta D)^2 \right] + O\left(\frac{1}{L^2}\right). \end{aligned} \quad (31)$$

The associated momentum transfer P is obtained as

$$\begin{aligned} P &= \frac{\pi \tilde{\lambda}}{L} \sum_{j=1}^{M+\Delta M} \left(2j - M - \Delta M - 1 + \frac{2\Delta D}{\tilde{\lambda}} \right) \\ &= 2k_F \Delta D + \frac{2\pi}{L} \Delta M \Delta D, \end{aligned} \quad (32)$$

where $k_F = \pi n$ is the Fermi momentum. Importantly, the difference between fermions and bosons only appears in the

selection rule between ΔM and ΔD ;

$$\Delta M : \text{integer}, \quad (33)$$

$$\Delta D = \frac{\Delta M}{2} \pmod{1}, \quad (34)$$

for the Fermi case, and

$$\Delta M : \text{integer}, \quad (35)$$

$$\Delta D : \text{integer}, \quad (36)$$

for the Bose case, respectively. In the Fermi case, we cannot take the current ΔD to be independent of the change of the particle number ΔM . Finally, by adding particle-hole excitation terms characterized by the quantum numbers n^\pm , which are nonnegative integers, we arrive at the finite-size scaling formula for the excitation energy and the associated momentum transfer as

$$\Delta E = \tilde{\mu} \Delta M + \frac{2\pi \tilde{v}}{L} \left[\frac{\tilde{\lambda}}{4} (\Delta M)^2 + \frac{1}{\tilde{\lambda}} (\Delta D)^2 + n^+ + n^- \right], \quad (37)$$

$$P = 2k_F \Delta D + \frac{2\pi}{L} (\Delta M \Delta D + n^+ - n^-). \quad (38)$$

It is worth noting that the excitation energy becomes complex due to dissipation, but the momentum transfer remains real. From Eqs. (37) and (38), we can read off the conformal weights characterizing the holomorphic and antiholomorphic parts of the underlying Virasoro algebra as

$$\Delta^\pm(\Delta M; \Delta D; n^\pm) = \frac{1}{2} \left(\frac{\Delta M \sqrt{\tilde{\lambda}}}{2} \pm \frac{\Delta D}{\sqrt{\tilde{\lambda}}} \right)^2 + n^\pm. \quad (39)$$

The conformal dimensions (39) are typical for a complex generalization of $c = 1$ U(1) Gaussian CFT [39], and the universal behavior of the system is characterized by the complex-valued TL parameter $\tilde{K} = 1/\tilde{\lambda}$. The critical exponents of dissipative TL liquids have already been obtained in Ref. [39], where we have calculated both biorthogonal correlation functions and right-state correlation functions, by using path integrals and ground-state wave function approach, respectively. Summarizing this section, we have confirmed that an ideal-gas description based on the ABA solution of the NH CS model captures the essential properties of dissipative TL liquids in NH quantum critical systems.

IV. DISSIPATIVE TOMONAGA-LUTTINGER LIQUIDS WITH $SU(N)$ SPIN SYMMETRY

In this section, we generalize the NH CS model to multi-component systems with $SU(N)$ symmetry, and analyze the NH $SU(N)$ CS model based on a complex generalization of Haldane's ideal-gas description.

A. Model

We consider a complex generalization of the $SU(N)$ CS model to 1D NH quantum systems. The NH $SU(N)$ CS model with the circumference L is given by

$$H_{\text{eff}} = -\frac{1}{2} \sum_{i=1}^M \frac{\partial^2}{\partial x_i^2} + \sum_{i < j} D(x_i - x_j)^{-2} \tilde{\lambda}' (\tilde{\lambda}' + P_{ij}^\sigma), \quad (40)$$

where $D(x) = (L/\pi) |\sin(\pi x/L)|$ is the chord distance, P_{ij}^σ ($\sigma = 1, 2, \dots, N$) is an operator that exchanges particle spins between the sites i and j , $\tilde{\lambda}'$ is the dimensionless complex parameter of the long-range interaction, and we have assumed $\text{Re}[\tilde{\lambda}'] > 0$. We note that the $SU(N)$ CS model was originally solved by Ha and Haldane [90], and we use similar notations which are used in Ref. [90]; $\tilde{\lambda}'$ in Eq. (40) and $\tilde{\lambda}$ in Eq. (16) are related by $\tilde{\lambda}' = \tilde{\lambda} - 1$, and Eqs. (40) and (16) are different from each other by a factor 2 in the single-component limit. Such difference in notations does not affect the universality.

The ground-state wave function of the $SU(N)$ CS model is a Jastrow type, and it has been demonstrated that the Jastrow factor does not contain the internal spin degrees of freedom [90–93]. Then, we obtain the ground-state wave function by generalizing the results obtained in the Hermitian case [90] to the NH case as

$$\Psi_g = \prod_{l > m} |z_l - z_m|^{\tilde{\lambda}' - s'} \Psi_0, \quad (41)$$

where

$$\Psi_0 = \prod_j z_j^{J_{\sigma_j}} \prod_{l > m} \left\{ (z_l - z_m)^{s' + \delta_{\sigma_l, \sigma_m}} \times \exp \left[\frac{i}{2} \pi \text{sgn}(\sigma_l - \sigma_m) \right] \right\}. \quad (42)$$

Here, $z_m = \exp(2\pi i x_m/L)$, σ_l is the ordered spin index, J_σ is the global current of the spin- σ particle, δ is the Kronecker's delta, and we have introduced $s' = 0$ for fermions and $s' = 1$

for bosons, respectively. Thus, we see from Eq. (41) that the complex interaction parameter $\tilde{\lambda}'$ in the Jastrow factor only affects the charge degrees of freedom even in the NH case.

B. Nested asymptotic Bethe ansatz

Next, to exploit an ideal-gas description, we derive the ABA solution of the NH $SU(N)$ CS model. By generalizing the S -matrices in the Hermitian multicomponent systems [91, 93, 112–114], we obtain the S -matrices of the NH $SU(N)$ CS model as

$$S_{ij} = \exp(-i\tilde{\lambda}'\theta(k_i - k_j)) \lim_{\eta \rightarrow 0} \frac{k_i - k_j + i\eta P_{ij}^\sigma}{k_i - k_j + i\eta}, \quad (43)$$

for fermions, and

$$S_{ij} = \exp(-i(\tilde{\lambda}' - 1)\theta(k_i - k_j)) \lim_{\eta \rightarrow 0} \frac{k_i - k_j + i\eta P_{ij}^\sigma}{k_i - k_j - i\eta}, \quad (44)$$

for bosons, respectively. Here, we have introduced $\theta(k) = \pi \text{sgn}^*(k)$, where the function $\text{sgn}^*(k)$ for the complex quasimomentum $k = \alpha e^{i\xi}$ ($\alpha \in \mathbb{R}$, $-\pi/2 < \xi < \pi/2$) is defined by $\text{sgn}^*(k) = \text{sgn}(\alpha)$ (see Fig. 2) in the same way as in the case without internal symmetry. It is noted that, for $k_i - k_j = \alpha e^{i\xi}$ in the S -matrices, we take the limit $\eta \rightarrow 0$ as $\eta' \rightarrow +0$ with $\eta = \eta' e^{i\xi}$ ($\eta' \in \mathbb{R}$). The additional phase shift in the exponential in the S -matrices $\pi(\tilde{\lambda}' - s') \text{sgn}^*(k_i - k_j)$ comes from the Jastrow factor in Eq. (41). The S -matrices in Eqs. (43) and (44) satisfy the Yang-Baxter equation $S_{jk} S_{ik} S_{ij} = S_{ij} S_{ik} S_{jk}$, hence we can construct the nested ABA [91, 93, 112–114] in NH multicomponent systems with $SU(N)$ symmetry. We note that, in the limit $\tilde{\lambda}' \rightarrow 0$, the S -matrix for fermions (43) reduces to the one for free fermions with $SU(N)$ internal spin symmetry [112, 113]. As for the limit $\tilde{\lambda}' - 1 \rightarrow 0$, the S -matrix for bosons (44) corresponds to the one for the $SU(N)$ Haldane-Shastry model [91, 114].

By using the S -matrices (43) and (44), the nested ABA equations for fermions ($s' = 0$) and bosons ($s' = 1$) are calculated as [91, 93, 112–114]

$$\exp(ik_j^{(1)} L) = (-1)^{(M_1-1)s'} \prod_{m=1}^{M_2} f_1(k_m^{(2)} - k_j^{(1)}) \prod_{l \neq j}^{M_1} \exp \left[-i\tilde{\lambda}'\theta(k_l^{(1)} - k_j^{(1)}) \right], \quad (1 \leq j \leq M_1), \quad (45)$$

$$\prod_{l \neq m}^{M_\sigma} f_2(k_m^{(\sigma)} - k_l^{(\sigma)}) = \prod_j^{M_{\sigma-1}} f_1(k_m^{(\sigma)} - k_j^{(\sigma-1)}) \prod_q^{M_{\sigma+1}} f_1(k_m^{(\sigma)} - k_q^{(\sigma+1)}), \quad (1 \leq m \leq M_\sigma, 2 \leq \sigma \leq N-1), \quad (46)$$

$$\prod_{l \neq q}^{M_N} f_2(k_q^{(N)} - k_l^{(N)}) = \prod_j^{M_{N-1}} f_1(k_q^{(N)} - k_j^{(N-1)}), \quad (1 \leq q \leq M_N), \quad (47)$$

respectively. Here, we have introduced $f_m(x) = \lim_{\eta \rightarrow 0} (x - i m \eta / 2) / (x + i m \eta / 2)$, and the number of the quasimomentum $k_j^{(\sigma)}$ as $M_\sigma = \sum_{\tau=\sigma}^N N_\tau$, where N_σ is the number of particles with spin σ . The nested ABA equations (45)–(47) describe a

charge excitation ($\sigma = 1$) characterized by the complex-valued interaction parameter $\tilde{\lambda}'$, and $N - 1$ kinds of spin excitations ($2 \leq \sigma \leq N$), where $\tilde{\lambda}'$ does not appear. The particles that appear in the spin sectors are called spinons, which are frequently used to describe spin excitations in the Haldane-Shastry model [83]. We see that the only difference between fermions and bosons is the factor of $(-1)^{M_1-1}$ in the right-hand side in Eq. (45), hence is included into the quantum number in the charge sector $I_j^{(1)}$ as shown below. Then, by taking the logarithm, we obtain the transcendental ABA equations as

$$k_j^{(1)} L = 2\pi I_j^{(1)} + \sum_m^{M_2} \theta(k_m^{(2)} - k_j^{(1)}) + \tilde{\lambda}' \sum_l^{M_1} \theta(k_j^{(1)} - k_l^{(1)}), \quad (1 \leq j \leq M_1), \quad (48)$$

$$\sum_l^{M_\sigma} \theta(k_m^{(\sigma)} - k_l^{(\sigma)}) + 2\pi I_m^{(\sigma)} = \sum_j^{M_{\sigma-1}} \theta(k_m^{(\sigma)} - k_j^{(\sigma-1)}) + \sum_q^{M_{\sigma+1}} \theta(k_m^{(\sigma)} - k_q^{(\sigma+1)}), \quad (1 \leq m \leq M_\sigma, 2 \leq \sigma \leq N-1), \quad (49)$$

$$\sum_l^{M_N} \theta(k_q^{(N)} - k_l^{(N)}) + 2\pi I_q^{(N)} = \sum_j^{M_{N-1}} \theta(k_q^{(N)} - k_j^{(N-1)}), \quad (1 \leq q \leq M_N), \quad (50)$$

where the quantum number $I_j^{(1)}$ that characterizes a charge excitation is given by

$$I_j^{(1)} = \frac{M_2}{2}, \quad (\text{mod } 1), \quad (51)$$

for fermions, and

$$I_j^{(1)} = \frac{M_1 + M_2 + 1}{2}, \quad (\text{mod } 1), \quad (52)$$

for bosons, respectively. On the other hand, $N - 1$ kinds of spin excitations ($2 \leq \sigma \leq N$) are described by the same quantum numbers in the case of fermions and bosons;

$$I_j^{(\sigma)} = \frac{M_{\sigma-1} + M_\sigma + M_{\sigma+1} + 1}{2}, \quad (\text{mod } 1, 2 \leq \sigma \leq N), \quad (53)$$

where $M_{N+1} = 0$. The energy of the system is given in terms of the quasimomentum $k_j^{(1)}$ in the charge sector as $E = (1/2) \sum_j [k_j^{(1)}]^2$. Importantly, dissipation parameter $\tilde{\lambda}'$ only appears in the charge sector in Eq. (45) and Eq. (48) in the nested ABA equations. This fact is nothing but the protection of spin sectors by $SU(N)$ internal symmetry in NH quantum systems, and can be seen as a NH generalization of the spin-charge separation in 1D quantum many-body systems [38, 39].

C. Velocity of excitations

In this subsection, we obtain the velocity of excitations, for which we generalize the distribution function and the dressed energy to NH quantum systems with $SU(N)$ symmetry. We first consider the ground state for general cases, where the chemical potential imbalance between different spin- σ particles is included. In such ground states, quantum numbers $\{I_j^{(\sigma)}\}$ are the successive integers distributed symmetrically around the origin. As the phase shift in the NH $SU(N)$ CS model is a step function with a complex coefficient, the

quasimomentum is equally spaced in each region C_τ between $\pm Q_\tau$ and $\pm Q_{\tau+1}$, where $|Q_1| \geq |Q_2| \geq \dots \geq |Q_N|$ and $Q_{N+1} = 0$ (see Fig. 2). As we see from Fig. 2, the quasimomentum $k_j^{(\sigma)}$ is distributed in the region $\sum_{\tau=\sigma}^N C_\tau$, where $\pm Q_\sigma$ denote the endpoints of the quasimomentum distribution of $k_j^{(\sigma)}$. Accordingly, the distribution function in the thermodynamic limit $\rho_\sigma(k)$ introduced for the quasimomentum $k_j^{(\sigma)}$ takes a constant complex value in each region C_τ , and is given by

$$\rho_\sigma(k) = \frac{\tau - \sigma + 1}{2\pi(1 + \tau\tilde{\lambda}')} , \quad (\tau \geq \sigma, \sigma = 1, 2, \dots, N). \quad (54)$$




FIG. 2. Distribution of the quasimomentum $k_j^{(\sigma)}$ in the ground state of the NH $SU(N)$ CS model (40) (Chemical potential imbalance between different spin- σ particles is included). Each region C_τ is constructed from two straight lines which are symmetric with respect to the origin; C_τ^+ (the region between Q_τ and $Q_{\tau+1}$) and C_τ^- (the region between $-Q_\tau$ and $-Q_{\tau+1}$). Quasimomentum is equally spaced in each region C_τ , and $k_j^{(\sigma)}$ is distributed in the region $\sum_{\tau=\sigma}^N C_\tau$, where $\pm Q_\sigma$ denote the endpoints of the quasimomentum distribution of $k_j^{(\sigma)}$. The generalized signature between $k_m^{(\sigma)}$ and $k_l^{(\sigma')}$ gives the phase shift $\theta(k_m^{(\sigma)} - k_l^{(\sigma')})$ (see text).

By using the distribution function (54), the particle density with spin σ is obtained as

$$n_\sigma = \frac{N_\sigma}{L} = \int_{\sum_{\tau=\sigma}^N C_\tau} \rho_\sigma(k) dk - \int_{\sum_{\tau=\sigma+1}^N C_\tau} \rho_{\sigma+1}(k) dk, \quad (55)$$

where we have assumed $n_1 \geq n_2 \geq \dots \geq n_N$, and the endpoint of each quasiparticle distribution Q_σ is given by the following set of equations;

$$Q_N = \pi n_N (1 + N\tilde{\lambda}'), \quad (56)$$

$$Q_{N-1} - Q_N = \pi(n_{N-1} - n_N)(1 + (N-1)\tilde{\lambda}'), \quad (57)$$

$$\vdots$$

$$Q_1 - Q_2 = \pi(n_1 - n_2)(1 + \tilde{\lambda}'), \quad (58)$$

which become complex reflecting the quasimomentum distribution in the complex plane.

Now, we concentrate on the $SU(N)$ singlet ground state, where $Q_1 = \dots = Q_N = \pi n(1 + N\tilde{\lambda}')/N$ with the total density $n = M/L$. In this case, only the region C_N exists and the quasimomentum $k_j^{(\sigma)}$ is distributed in the region C_N depicted by the straight line (see Fig. 2). By inserting a hole into the ground-state distribution of the quasimomentum, the dressed energy is calculated with the help of the nested ABA equations (48)–(50) as

$$\epsilon_1(k) = \frac{1}{2}k^2 - \tilde{\mu} + \int_{-Q_N}^{Q_N} \delta(k - k') \epsilon_2(k') dk' - \tilde{\lambda}' \int_{-Q_N}^{Q_N} \delta(k - k') \epsilon_1(k') dk', \quad (59)$$

$$\epsilon_\sigma(k) = \sum_{q=-1,0,1} (-1)^{q+1} \int_{-Q_N}^{Q_N} \delta(k - k') \epsilon_{\sigma+q}(k') dk', \quad (\sigma = 2, 3, \dots, N-1), \quad (60)$$

$$\epsilon_N(k) = \sum_{q=-1,0} (-1)^{q+1} \int_{-Q_N}^{Q_N} \delta(k - k') \epsilon_{N+q}(k') dk', \quad (61)$$

where the complex chemical potential $\tilde{\mu}$ is obtained by using $\epsilon_\sigma(\pm Q_N) = 0$ as $\tilde{\mu} = \pi^2 n^2 (1 + N\tilde{\lambda}')^2 / 2N^2$. Finally, we arrive at the velocity of excitations for the $SU(N)$ singlet ground state as

$$\tilde{v}_\sigma = \frac{\epsilon'_\sigma(Q_N)}{2\pi\rho_\sigma(Q_N)} = \frac{\pi n}{N} (1 + N\tilde{\lambda}') \equiv \tilde{v}, \quad (62)$$

where $\epsilon'_\sigma(k)$ denotes the derivative of $\epsilon_\sigma(k)$ with respect to k . Here, \tilde{v}_1 characterizes the charge velocity and \tilde{v}_σ ($\sigma = 2, 3, \dots, N$) describes the $N-1$ kinds of spin velocity. We note that the velocity for the charge excitation \tilde{v}_1 and that for the spin excitations \tilde{v}_σ ($\sigma = 2, 3, \dots, N$) are the same in long-range interacting systems (40). It is worth noting that both the charge velocity and the spin velocity are affected by dissipation through $\tilde{\lambda}'$ as a result of the nested equations of ABA.

D. Finite-size spectrum

In this subsection, we obtain the finite-size spectrum that leads to the universal scaling relations for dissipative TL liquids with $SU(N)$ spin symmetry in NH quantum critical systems. In Hermitian systems, $SU(N)$ CS model has been shown to describe the universal properties of TL liquids with $SU(N)$ spin symmetry as an ideal gas, e.g., $SU(2)$ CS model

successfully describes the $SU(2)$ spin symmetric TL liquids, which characterize the critical properties of Hubbard models [92, 93, 115–117]. The excitation energy in a finite system for the NH $SU(N)$ CS model can be obtained by using the nested ABA equations (48)–(50) [91]. This method is useful to understand conformal properties stemming from $U(1) \times SU(N)$ symmetry underlying in the ABA equations as we see below. On the other hand, the finite-size spectrum for the Hermitian $SU(N)$ CS model was originally obtained by Ha and Haldane [90] with the use of the bosonization method, which also embodies an ideal-gas description. This method is beneficial to interpret the particle picture and the origin of associated selection rules for Fermi and Bose statistics. To understand the effect of dissipation on each particle more clearly, we start from the particle picture introduced in Ref. [90] and then transform it to the ABA picture underlying in Eqs. (48)–(50).

By constructing excited states over the $SU(N)$ singlet ground state in Eq. (41) systematically, we end up with the finite-size spectrum, which is given by analytically continuing the one obtained in Ref. [90] to the case of the NH $SU(N)$ CS model. The result is written down as

$$\Delta E = \frac{\pi \tilde{v}}{L} \sum_{\sigma\sigma'} \left[\frac{1}{2} A_{\sigma\sigma'} \Delta M_\sigma \Delta M_{\sigma'} + 2 A_{\sigma\sigma'}^{-1} \Delta J_\sigma \Delta J_{\sigma'} \right], \quad (63)$$

where we have to pay careful attention to the fact that not

only the interaction parameter $\tilde{\lambda}'$ but also the velocity of excitations \tilde{v} becomes complex-valued as a result of dissipation [39]. We also note that particle-hole excitation terms that form conformal towers inherent in the underlying Virasoro algebra are omitted in Eq. (63) because they do not affect the critical exponents of leading order that we consider in the next subsection. In Eq. (63), the coefficient $A_{\sigma\sigma'} = \tilde{\lambda}' + \delta_{\sigma\sigma'}$ is the matrix element of the $N \times N$ matrix \mathbf{A} given by

$$\mathbf{A} = \begin{pmatrix} \tilde{\lambda}' + 1 & \tilde{\lambda}' & \tilde{\lambda}' & \cdots \\ \tilde{\lambda}' & \tilde{\lambda}' + 1 & \tilde{\lambda}' & \\ \tilde{\lambda}' & \tilde{\lambda}' & \ddots & \\ \vdots & & & \end{pmatrix}, \quad (64)$$

and $A_{\sigma\sigma'}^{-1} = -\tilde{\lambda}'/(1+N\tilde{\lambda}') + \delta_{\sigma\sigma'}$ is that of the inverse matrix \mathbf{A}^{-1} , which is given by

$$\mathbf{A}^{-1} = \begin{pmatrix} \chi + 1 & \chi & \chi & \cdots \\ \chi & \chi + 1 & \chi & \\ \chi & \chi & \ddots & \\ \vdots & & & \end{pmatrix}, \quad (65)$$

where we have introduced $\chi = -\tilde{\lambda}'/(1+N\tilde{\lambda}')$. We note that ΔM_σ and ΔJ_σ are the vector elements of the $N \times 1$ column vectors that characterize two types of excitations given by $\Delta \mathbf{M}$ and $\Delta \mathbf{J}$, respectively. Here, ΔM_σ denotes the particle-number excitations, and ΔJ_σ describes the excitations accompanied by the large momentum transfer P_σ for a spin- σ particle, which is given by

$$P_\sigma = 2k_F \Delta J_\sigma, \quad (66)$$

where the Fermi momentum $k_F = \pi n/N$ is generalized to the $SU(N)$ singlet ground state.

As we see in Eq. (64), dissipation affects all the elements of the matrix \mathbf{A} equally via the complex-valued interaction parameter $\tilde{\lambda}'$. We emphasize that the difference of statistics between fermions and bosons are incorporated into the selection rules of quantum numbers, which are given by

$$\Delta M_\sigma : \text{integer}, \quad (67)$$

$$\Delta J_\sigma = \frac{\Delta M_\sigma}{2} \pmod{1}, \quad (68)$$

for fermions, and

$$\Delta M_\sigma : \text{integer}, \quad (69)$$

$$\Delta J_\sigma : \text{integer}, \quad (70)$$

for bosons, respectively. We see that the selection rules for each spin- σ particle (67)–(70) are the same as those in the single component cases (33)–(36).

The basis of the matrices used in Eqs. (64) and (65) are the one which was introduced by Ha and Haldane [90], and is different from the one used in the ABA equations (48)–(50). The basis transformation between them is conducted by using the following $N \times N$ matrix α [94]

$$\alpha = \begin{pmatrix} 1 & -1 & & & \\ & 1 & -1 & \mathbf{0} & \\ & & \ddots & \ddots & \\ \mathbf{0} & & & \ddots & -1 \\ & & & & 1 \end{pmatrix}, \quad (71)$$

which gives the inverse matrix as

$$\alpha^{-1} = \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 \\ & 1 & 1 & \cdots & 1 \\ & & 1 & \cdots & 1 \\ \mathbf{0} & & & \ddots & 1 \\ & & & & 1 \end{pmatrix}. \quad (72)$$

Then, with the help of Eq. (71), the excitation energy (63) is rewritten as $\Delta E = (\pi \tilde{v}/L)(\mathbf{m}^t \mathbf{D} \mathbf{m}/2 + 2\mathbf{d}^t \mathbf{D}^{-1} \mathbf{d})$, where the matrix \mathbf{A} is transformed into the matrix \mathbf{D} , which is calculated as

$$\mathbf{D} \equiv \alpha^t \mathbf{A} \alpha = \begin{pmatrix} \tilde{\lambda}' + 1 & -1 & & & \\ -1 & 2 & -1 & \mathbf{0} & \\ & -1 & 2 & \ddots & \\ \mathbf{0} & & \ddots & \ddots & -1 \\ & & & -1 & 2 \end{pmatrix}. \quad (73)$$

Similarly, with the use of Eq. (72), the inverse matrix \mathbf{D}^{-1} is written as follows;

$$\mathbf{D}^{-1} \equiv \alpha^{-1} \mathbf{A}^{-1} (\alpha^t)^{-1} = \frac{1}{1+N\tilde{\lambda}'} \begin{pmatrix} N & N-1 & N-2 & \cdots & 1 \\ N-1 & (N-1)(\tilde{\lambda}'+1) & (N-2)(\tilde{\lambda}'+1) & \cdots & \tilde{\lambda}'+1 \\ N-2 & (N-2)(\tilde{\lambda}'+1) & (N-2)(2\tilde{\lambda}'+1) & & \vdots \\ \vdots & \vdots & \vdots & (N-2)\tilde{\lambda}'+1 & \\ 1 & \tilde{\lambda}'+1 & 2\tilde{\lambda}'+1 & \cdots & (N-1)\tilde{\lambda}'+1 \end{pmatrix}. \quad (74)$$

Accordingly, the basis transformations for the quantum num-

ber vectors $\Delta \mathbf{M}$ and $\Delta \mathbf{J}$ are given by

$$\mathbf{m} = \alpha^{-1} \Delta \mathbf{M}, \quad (75)$$

$$\mathbf{d} = \alpha^t \Delta \mathbf{J}, \quad (76)$$

where the quantum numbers m_1 and d_1 characterize excitations in the charge sector with the Fermi point Nk_F , and m_σ and d_σ ($2 \leq \sigma \leq N$) describe excitations in the spin sectors with the Fermi point $(N - \sigma + 1)k_F$. In detail, m_σ denotes the change of the number of charge particles for $\sigma = 1$, and that of spinons for $2 \leq \sigma \leq N$. Accordingly, d_σ carries the large momentum P'_σ associated with charge ($\sigma = 1$) or spin ($2 \leq \sigma \leq N$) excitations as follows;

$$P'_\sigma = 2(N - \sigma + 1)k_F d_\sigma. \quad (77)$$

By using the quantum numbers m_σ and d_σ , the selection rules are rewritten as

$$m_\sigma : \text{integer}, \quad (78)$$

$$d_\sigma = \frac{m_{\sigma-1} + m_{\sigma+1}}{2} \pmod{1}, \quad (79)$$

with $m_0 = m_1$ and $m_{N+1} = 0$ for fermions, and

$$m_\sigma : \text{integer}, \quad (80)$$

$$d_\sigma : \text{integer}, \quad (81)$$

for bosons, respectively. The selection rules for m_σ and d_σ given in Eqs. (78)–(81) are rather complicated than those for ΔM_σ and ΔJ_σ given in Eqs. (67)–(70), but are easily obtained by using the basis transformation given by the matrix α in Eq. (71).

Notably, the matrix D in Eq. (73) reflects the $SU(N)$ internal spin symmetry underlying in the present dissipative TL liquids. $(N-1) \times (N-1)$ matrix obtained by deleting the first column and the first row in D is called the $SU(N)$ Cartan matrix, which characterizes $N-1$ kinds of spin excitations with $c = N-1$ level-1 $SU(N)$ Kac-Moody algebra [118–122]. On the other hand, we see in Eq. (73) that the effect of dissipation through the complex parameter $\tilde{\lambda}'$ only appears in the charge sector, which is characterized by a complex generalization of $c = 1$ U(1) Gaussian CFT [39] as in the case without internal spin symmetry discussed in Sec. III.

E. Universal scaling relations

Finally, we obtain the critical exponents which provide universal properties of dissipative TL liquids with $SU(N)$ spin symmetry. From the excitation energy spectrum (63) and the basis transformation given in Eq. (73), the general formula of the (complex) critical exponents that express dissipative TL liquids with $SU(N)$ spin symmetry are read off as [123]

$$\tilde{\eta} = \frac{1}{2} \Delta M^t A \Delta M + 2 \Delta J^t A^{-1} \Delta J \quad (82)$$

$$= \frac{1}{2} m^t D m + 2 d^t D^{-1} d, \quad (83)$$

where the first line (82) corresponds to the basis introduced by Ha and Haldane [90], and the second line (83) corresponds to the one used in the ABA equations (48)–(50). It is noted that $\tilde{\eta}$ is the critical exponent of biorthogonal correlation functions

in NH systems [39], hence is a complex value. In the following, we obtain the scaling formula for both biorthogonal correlation functions and right-state correlation functions in NH systems based on the formula of $\tilde{\eta}$ given in Eqs. (82) and (83).

1. Critical exponents for biorthogonal correlation functions

We first calculate the critical exponents for biorthogonal correlation functions. The exponents of biorthogonal correlators become complex, and are directly given by the formula $\tilde{\eta}$ given in Eqs. (82) and (83). First, the long-distance behavior of the biorthogonal charge-density correlator is given by

$${}_L \langle \rho(x) \rho(0) \rangle_R \simeq \sum_{j=1}^N \tilde{A}_j \cos[2(N-j+1)k_F x] x^{-\tilde{\beta}_j} + \frac{\tilde{A}_0}{x^2}, \quad (84)$$

which has the same form both in the case of fermions and bosons. Here, the exponent $\tilde{\beta}_j$ denotes the $2(N-j+1)k_F$ oscillating part ($1 \leq j \leq N$) of the biorthogonal charge-density correlator (84), and \tilde{A}_j is a complex-valued correlation amplitude. As Eq. (84) conserves the particle number, the selection rules for ΔM and m for this type of excitations reduce to

$$\Delta M = m = 0, \quad (85)$$

which brings about the same form of the charge density correlators for both fermions and bosons. For the excitations that convey the large momentum transfer given in Eqs. (66) and (77), the selection rules corresponding to the $2(N-j+1)k_F$ oscillations ($1 \leq j \leq N$) associated with the momentum transfer $2Nk_F, 2(N-1)k_F, \dots, 2k_F$ are given by

$2Nk_F$ oscillation,

$$\Delta J = (1, 1, \dots, 1)^t, \quad d = (1, 0, 0, \dots, 0)^t, \quad (86)$$

$2(N-1)k_F$ oscillation,

$$\Delta J = (0, 1, \dots, 1)^t, \quad d = (0, 1, 0, \dots, 0)^t, \quad (87)$$

\vdots

$2k_F$ oscillation,

$$\Delta J = (0, \dots, 0, 1)^t, \quad d = (0, 0, \dots, 0, 1)^t, \quad (88)$$

respectively. Equations (86)–(88) provide the critical exponent $\tilde{\beta}_j$ corresponding to the $2(N-j+1)k_F$ oscillating piece ($1 \leq j \leq N$) as

$$\tilde{\beta}_j = \frac{2(N-j+1)(j-1)}{N} + \frac{2(N-j+1)^2}{N} \tilde{K}_\rho, \quad (89)$$

where the complex-valued TL parameter \tilde{K}_ρ is given by $\tilde{K}_\rho = 1/(1 + N\tilde{\lambda}')$ with the use of Eq. (12). Notably, from Eq. (89), we see that the effect of dissipation is included only in the charge excitation described by \tilde{K}_ρ .

On the other hand, fermion correlators and boson correlators are different from each other. The long-distance behavior of the biorthogonal fermion correlator, which accompanies the k_F oscillation, is written as

$${}_L\langle c_\sigma^\dagger(x)c_\sigma(0)\rangle_R \simeq \tilde{C}_1 \cos(k_F x) x^{-\tilde{\eta}_F}, \quad (90)$$

where \tilde{C}_1 is a complex-valued correlation amplitude. It should be noted that the biorthogonal fermion correlator (90) is independent of spin indices for the $SU(N)$ singlet ground state. As the correlation function (90) changes the particle number by one, the selection rules for the biorthogonal fermion correlator (90) are given by

$$\sigma = 1$$

$$\Delta \mathbf{M} = (1, 0, \dots, 0)^t, \quad \mathbf{m} = (1, 0, \dots, 0)^t, \quad (91)$$

$$\Delta \mathbf{J} = (\frac{1}{2}, 0, \dots, 0)^t, \quad \mathbf{d} = (\frac{1}{2}, -\frac{1}{2}, 0, \dots, 0)^t, \quad (92)$$

$$\sigma = 2$$

$$\Delta \mathbf{M} = (0, 1, 0, \dots, 0)^t, \quad \mathbf{m} = (1, 1, 0, \dots, 0)^t, \quad (93)$$

$$\Delta \mathbf{J} = (0, \frac{1}{2}, 0, \dots, 0)^t, \quad \mathbf{d} = (0, \frac{1}{2}, -\frac{1}{2}, 0, \dots, 0)^t, \quad (94)$$

$$\vdots$$

$$\sigma = N$$

$$\Delta \mathbf{M} = (0, \dots, 0, 1)^t, \quad \mathbf{m} = (1, 1, \dots, 1)^t, \quad (95)$$

$$\Delta \mathbf{J} = (0, \dots, 0, \frac{1}{2})^t, \quad \mathbf{d} = (0, \dots, 0, \frac{1}{2})^t, \quad (96)$$

all of which lead to the same critical exponent $\tilde{\eta}_F$ as

$$\tilde{\eta}_F = \frac{N-1}{N} + \frac{1}{2N\tilde{K}_\rho} + \frac{\tilde{K}_\rho}{2N}. \quad (97)$$

Again, we see from Eq. (97) that dissipation only affects the exponent of the charge excitation characterized by \tilde{K}_ρ , and the exponent stemming from spin excitations, $(N-1)/N$, is fixed to be real as a result of the underlying $SU(N)$ symmetry.

The long-distance behavior of the biorthogonal boson correlator, which does not accompany any oscillations, are given by

$${}_L\langle b_\sigma^\dagger(x)b_\sigma(0)\rangle_R \simeq \tilde{B}_1 x^{-\tilde{\eta}_B}, \quad (98)$$

where \tilde{B}_1 is a complex-valued correlation amplitude. We note that the boson correlator in Eq. (98) is independent of spin indices for the $SU(N)$ singlet ground state as in the fermion case (90). As the leading part of the biorthogonal boson correlator (98) does not carry the large momentum from the left Fermi point to the right one, the selection rules for $\Delta \mathbf{J}$ and \mathbf{d} read

$$\Delta \mathbf{J} = \mathbf{d} = \mathbf{0}. \quad (99)$$

The boson correlator (98) changes the number of particles by one as in the fermion case (90), hence the selection rules for

$\Delta \mathbf{M}$ and \mathbf{m} are the same as Eqs. (91)–(96). Then, we obtain the critical exponent $\tilde{\eta}_B$ as

$$\tilde{\eta}_B = \frac{N-1}{2N} + \frac{1}{2N\tilde{K}_\rho}, \quad (100)$$

where the piece proportional to the TL parameter does not appear in contrast to the one in the fermion case (97). Importantly, the critical exponents of the biorthogonal correlation functions (89), (97), and (100) provide the universal scaling relations for dissipative TL liquids with $SU(N)$ spin symmetry.

2. Critical exponents for right-state correlation functions

Next, we analyze the right-state correlation functions, which are experimentally relevant in ultracold atoms. We emphasize that the exponents of right-state correlators are real because right-state correlation functions are obtained as a standard quantum-mechanical expectation value for the ground state [36, 39].

The critical exponents of the right-state correlation functions are obtained from Eqs. (89), (97), and (100) by replacing \tilde{K}_ρ with K_ρ^ϕ , and $1/\tilde{K}_\rho$ with $1/K_\rho^\theta$, respectively. This simple transformation comes from the fact that dissipation only affects the charge excitation characterized by the complex-valued TL parameter \tilde{K}_ρ . As \tilde{K}_ρ describes the charge degrees of freedom, the relations among the TL parameters \tilde{K}_ρ , K_ρ^ϕ , and K_ρ^θ given in Eqs. (4) and (8) hold for dissipative TL liquids both with and without internal symmetry [39]. As a result, two types of real TL parameters K_ρ^ϕ and K_ρ^θ describe the universal properties of right-state correlation functions through the general formula (4) and (8) in dissipative TL liquids with $SU(N)$ spin symmetry. Therefore, in terms of K_ρ^ϕ and K_ρ^θ , we obtain the universal scaling relations for β_j , η_F , and η_B in the right-state correlation functions as Eqs. (11), (13), and (15) as summarized in the main results in Sec. II. These exponents characterize the universal behavior of dissipative TL liquids with $SU(N)$ spin symmetry, and are relevant to experiments.

In Fig. 3, we plot the critical exponents K_ρ^θ given in Eq. (8) and η_F given in Eq. (13) in long-range interacting $SU(N)$ symmetric systems (40) for the case of $N=2$ and $N=4$. We note that K_ρ^ϕ is not affected by dissipation in the present case due to an artifact of long-range interactions. From Fig. 3(a), we see that the TL parameter K_ρ^θ is suppressed as dissipation $\text{Im}\tilde{\lambda}'$ is increased. This is a similar behavior obtained in the NH Lieb-Liniger model [36], where the TL parameter is suppressed due to dissipation as a result of the continuous quantum Zeno effect [7, 10–12, 14, 23]. As we deal with the continuum model, the increase in TL parameters as a result of the Umklapp scattering reported in lattice models is not seen [39]. Accordingly, the behavior of η_F shown in Fig. 3(b) reflects the suppression of K_ρ^θ via dissipation $\text{Im}\tilde{\lambda}'$, and shows the enhancement due to dissipation. The enhancement in η_F gives the suppression of the fermion correlator (6), the behavior of which is consistent with the one obtained in the NH




FIG. 3. Critical exponents (a) K_ρ^θ and (b) η_F as a function of the complex-valued interaction parameter $\tilde{\lambda}'$ for $N = 2$ (orange) and $N = 4$ (blue) in the NH $SU(N)$ CS model (40). $\text{Re}\tilde{\lambda}'$ and $\text{Im}\tilde{\lambda}'$ denote a (real-valued) interaction, and dissipation, respectively. K_ρ^θ and η_F do not depend on the sign of $\text{Im}\tilde{\lambda}'$.

Lieb-Liniger model [36]. In Fig. 3(b), we see that the enhancement of η_F due to dissipation $\text{Im}\tilde{\lambda}'$ becomes significant in the limit of free fermions, and the difference of η_F induced by dissipation between $N = 2$ and $N = 4$ also becomes significant in this limit. We note that almost the same behavior as that of η_F is obtained for η_B given in Eq. (15).

V. CONCLUSIONS

We have explored the universal properties of dissipative TL liquids with $SU(N)$ spin symmetry in one dimension based on a complex generalization of Haldane's ideal-gas description, which is realized by the $SU(N)$ Calogero-Sutherland model. As the main results, we have obtained the universal scaling relations for dissipative TL liquids with $SU(N)$ spin symmetry for both fermions and bosons. We have demonstrated that the spectrum of dissipative TL liquids with $SU(N)$ spin symmetry is described by the sum of one charge mode characterized by a complex generalization of $c = 1$ U(1) Gaussian CFT, and $N - 1$ spin modes characterized by level-1 $SU(N)$ Kac-Moody algebra with the conformal anomaly $c = N - 1$. We have elucidated that dissipation only affects the charge sector in the correlation exponents by deriving ABA solutions and

using CFT in NH quantum critical many-body systems. The scaling relations obtained in the present work are relevant to a wide variety of cold-atom experiments in NH quantum critical systems with $SU(N)$ symmetry, e.g., ultracold alkaline-earth-like atom ^{173}Yb is a promising candidate, where the Fermi-Hubbard model is successfully loaded into an optical lattice, and dissipation is introduced by using photoassociation techniques [19]. As a useful tool to postselect special measurement outcomes that realize NH quantum many-body systems, quantum-gas microscopy can be utilized [36–39] to observe dissipative TL liquids with $SU(N)$ spin symmetry.

There remains an interesting question how the present results for dissipative TL liquids are related to the critical behavior studied in open quantum systems [124, 125]. Critical exponents of correlation functions and correlation length are essential to investigate the microscopic origin of quantum phase transitions. As critical exponents in open quantum systems have been investigated by using the framework of Lindblad master equations [124, 125], it is important to clarify the universality class which lies in dissipative quantum critical systems from the perspective of CFT. As for the relations between NH quantum systems and Markovian open quantum systems, it is known that the spectrum of the Liouvillian is obtained from that of the NH Hamiltonian when the system follows loss-only dynamics [40, 105, 126]. This may benefit us for finding out universal relations of critical exponents between NH quantum systems and Markovian open quantum systems. Moreover, in view of recent advancement of CFT description in NH quantum systems [127–129] and measurement-induced entanglement dynamics [130–133], it is interesting to investigate how CFT in NH quantum many-body systems is related to the one in measurement-induced dynamics.

Dissipation makes the theory nonunitary and the entire spectrum of the NH quantum many-body systems can be complicated. However, low-energy physics shows universal properties in NH quantum systems with internal degrees of freedom, which leads to unconventional quantum critical phenomena characterized by a complex extension of CFT. Since systems with inverse-square long-range interactions are known to be closely related to the fractional quantum Hall effect [86, 87], characterization of NH fractional quantum Hall effects [134, 135] by using the present framework remains a future research subject. As ultracold mixtures with $SU(N) \times SU(N')$ symmetry have been realized [136], it is of interest to extend our theory to more general multicomponent systems beyond $SU(N)$ symmetry. Our results in this paper will certainly stimulate further study on multicomponent extension of dissipative TL liquids in open quantum systems [32, 36, 37, 125, 137–144].

ACKNOWLEDGMENTS

This work was supported by KAKENHI (Grants No. JP18H01140 and No. JP19H01838). K.Y. was supported by WISE Program, MEXT and JSPS KAKENHI Grant-in-Aid for JSPS fellows Grant No. JP20J21318.

Appendix A: Ground-state wave function

In this appendix, we give a proof that a complex generalization of the wave function of the Jastrow form (18) provides the eigenstate of the effective Hamiltonian (16) with the complex-valued ground-state energy E_0 given in Eq. (30). Here, we consider the case of bosons for simplicity, and that of fermions can be proved similarly by taking the anticommutation relations of fermions into account. We calculate the

we obtain

$$\begin{aligned} -\frac{\partial^2}{\partial x_j^2} \Psi_g &= -\frac{\pi^2 \tilde{\lambda}}{L^2} \left[\tilde{\lambda} \left(\sum_{i(\neq j)} \cot \frac{\pi(x_j - x_i)}{L} \right)^2 - \sum_{i(\neq j)} \frac{1}{\sin^2(\pi(x_j - x_i)/L)} \right] \Psi_g \\ &= -\frac{\pi^2 \tilde{\lambda}}{L^2} \left(\sum_{i(\neq j)} \frac{\tilde{\lambda} - 1}{\sin^2(\pi(x_j - x_i)/L)} - \tilde{\lambda}(N - 1) + 2\tilde{\lambda} \sum_{\langle i, k \rangle (\neq j)} \cot \frac{\pi(x_j - x_i)}{L} \cot \frac{\pi(x_j - x_k)}{L} \right) \Psi_g, \end{aligned} \quad (\text{A3})$$

where we have used the identity $\cot^2 x = 1/\sin^2 x - 1$ in the second equality, $\langle i, k \rangle$ denotes the combination over the pair of i and k with $i \neq k$, and $\langle i, k \rangle (\neq j)$ means that both i and k are different from j . By taking the sum of Eq. (A3), we get

$$\begin{aligned} -\sum_j \frac{\partial^2}{\partial x_j^2} \Psi_g &= \frac{\pi^2 \tilde{\lambda}}{L^2} \left[-2 \sum_{i > j} \frac{\tilde{\lambda} - 1}{\sin^2(\pi(x_i - x_j)/L)} + \tilde{\lambda}N(N - 1) - 2\tilde{\lambda} \sum_{\langle i, j, k \rangle} \left(\cot \frac{\pi(x_j - x_i)}{L} \cot \frac{\pi(x_j - x_k)}{L} + (i \rightarrow j \rightarrow k) \right) \right] \Psi_g, \end{aligned} \quad (\text{A4})$$

where $\langle i, j, k \rangle$ denotes the combination over the pair of $\{i, j, k\}$ which are different from each other, and $(i \rightarrow j \rightarrow k)$ stands for two terms of cyclic permutations of the first term in the same parentheses. Finally, by using the identity

$$\cot x \cot y + \cot y \cot z + \cot z \cot x = 1, \quad (x + y + z = 0), \quad (\text{A5})$$

we arrive at

$$-\sum_j \frac{\partial^2}{\partial x_j^2} \Psi_g = \left(\frac{\pi^2 N(N^2 - 1)\tilde{\lambda}^2}{3L^2} - \frac{2\pi^2 \tilde{\lambda}(\tilde{\lambda} - 1)}{L^2} \sum_{i > j} \frac{1}{\sin^2(\pi(x_i - x_j)/L)} \right) \Psi_g. \quad (\text{A6})$$

One notices that the second term in the right-hand side of Eq. (A6) has the same form as the interaction term (17). Therefore, by taking $2\tilde{\lambda}(\tilde{\lambda} - 1) = \tilde{g}$, the wave function (18) becomes the eigenstate with the ground state energy E_0 given in Eq. (30).

-
- [1] M. Müller, S. Diehl, G. Pupillo, and P. Zoller, “Engineered open systems and quantum simulations with atoms and ions,” *Adv. Atom. Mol. Opt. Phys.* **61**, 1 (2012).
 - [2] A. J. Daley, “Quantum trajectories and open many-body quantum systems,” *Adv. Phys.* **63**, 77 (2014).
 - [3] L. M. Sieberer, M. Buchhold, and S. Diehl, “Keldysh field theory for driven open quantum systems,” *Rep. Prog. Phys.* **79**, 096001 (2016).

- [4] Y. Ashida, Z. Gong, and M. Ueda, “Non-hermitian physics,” *Adv. Phys.* **69**, 249 (2020).
- [5] S. Diehl, A. Micheli, A. Kantian, B. Kraus, H. Büchler, and P. Zoller, “Quantum states and phases in driven open quantum systems with cold atoms,” *Nat. Phys.* **4**, 878 (2008).
- [6] B. Kraus, H. P. Büchler, S. Diehl, A. Kantian, A. Micheli, and P. Zoller, “Preparation of entangled states by quantum Markov processes,” *Phys. Rev. A* **78**, 042307 (2008).

- [7] J. J. García-Ripoll, S. Dürr, N. Syassen, D. M. Bauer, M. Lettner, G. Rempe, and J. I. Cirac, “Dissipation-induced hardcore boson gas in an optical lattice,” *New J. Phys.* **11**, 013053 (2009).
- [8] S. Dürr, J. J. García-Ripoll, N. Syassen, D. M. Bauer, M. Lettner, J. I. Cirac, and G. Rempe, “Lieb-Liniger model of a dissipation-induced Tonks-Girardeau gas,” *Phys. Rev. A* **79**, 023614 (2009).
- [9] K. Yamamoto, Y. Ashida, and N. Kawakami, “Rectification in nonequilibrium steady states of open many-body systems,” *Phys. Rev. Research* **2**, 043343 (2020).
- [10] K. Yamamoto, M. Nakagawa, N. Tsuji, M. Ueda, and N. Kawakami, “Collective Excitations and Nonequilibrium Phase Transition in Dissipative Fermionic Superfluids,” *Phys. Rev. Lett.* **127**, 055301 (2021).
- [11] N. Syassen, D. M. Bauer, M. Lettner, T. Volz, D. Dietze, J. J. García-Ripoll, J. I. Cirac, G. Rempe, and S. Dürr, “Strong dissipation inhibits losses and induces correlations in cold molecular gases,” *Science* **320**, 1329 (2008).
- [12] B. Yan, S. A. Moses, B. Gadway, J. P. Covey, K. R. Hazzard, A. M. Rey, D. S. Jin, and J. Ye, “Observation of dipolar spin-exchange interactions with lattice-confined polar molecules,” *Nature (London)* **501**, 521 (2013).
- [13] G. Barontini, R. Labouvie, F. Stubenrauch, A. Vogler, V. Guarnera, and H. Ott, “Controlling the Dynamics of an Open Many-Body Quantum System with Localized Dissipation,” *Phys. Rev. Lett.* **110**, 035302 (2013).
- [14] B. Zhu, B. Gadway, M. Foss-Feig, J. Schachenmayer, M. L. Wall, K. R. A. Hazzard, B. Yan, S. A. Moses, J. P. Covey, D. S. Jin, J. Ye, M. Holland, and A. M. Rey, “Suppressing the Loss of Ultracold Molecules Via the Continuous Quantum Zeno Effect,” *Phys. Rev. Lett.* **112**, 070404 (2014).
- [15] T. Tomita, S. Nakajima, I. Danshita, Y. Takasu, and Y. Takahashi, “Observation of the Mott insulator to superfluid crossover of a driven-dissipative Bose-Hubbard system,” *Sci. Adv.* **3**, e1701513 (2017).
- [16] K. Sponselee, L. Freystatzky, B. Abeln, M. Diem, B. Hundt, A. Kochanek, T. Ponath, B. Santra, L. Mathey, K. Sengstock, and C. Becker, “Dynamics of ultracold quantum gases in the dissipative Fermi-Hubbard model,” *Quantum Sci. Technol.* **4**, 014002 (2018).
- [17] T. Tomita, S. Nakajima, Y. Takasu, and Y. Takahashi, “Dissipative Bose-Hubbard system with intrinsic two-body loss,” *Phys. Rev. A* **99**, 031601(R) (2019).
- [18] R. Bouganne, M. B. Aguilera, A. Ghermaoui, J. Beugnon, and F. Gerbier, “Anomalous decay of coherence in a dissipative many-body system,” *Nat. Phys.* **16**, 21 (2020).
- [19] K. Honda, S. Taie, Y. Takasu, N. Nishizawa, M. Nakagawa, and Y. Takahashi, “Observation of the Sign Reversal of the Magnetic Correlation in a Driven-Dissipative Fermi-Hubbard System,” *arXiv:2205.13162*.
- [20] M. Nakagawa, N. Kawakami, and M. Ueda, “Non-Hermitian Kondo Effect in Ultracold Alkaline-Earth Atoms,” *Phys. Rev. Lett.* **121**, 203001 (2018).
- [21] J. A. S. Lourenço, R. L. Eneias, and R. G. Pereira, “Kondo effect in a \mathcal{PT} -symmetric non-Hermitian Hamiltonian,” *Phys. Rev. B* **98**, 085126 (2018).
- [22] R. Hanai and P. B. Littlewood, “Critical fluctuations at a many-body exceptional point,” *Phys. Rev. Research* **2**, 033018 (2020).
- [23] K. Yamamoto, M. Nakagawa, K. Adachi, K. Takasan, M. Ueda, and N. Kawakami, “Theory of Non-Hermitian Fermionic Superfluidity with a Complex-Valued Interaction,” *Phys. Rev. Lett.* **123**, 123601 (2019).
- [24] R. Hamazaki, K. Kawabata, and M. Ueda, “Non-Hermitian Many-Body Localization,” *Phys. Rev. Lett.* **123**, 090603 (2019).
- [25] R. Hanai, A. Edelman, Y. Ohashi, and P. B. Littlewood, “Non-Hermitian Phase Transition from a Polariton Bose-Einstein Condensate to a Photon Laser,” *Phys. Rev. Lett.* **122**, 185301 (2019).
- [26] Y. Li, X. Chen, and M. P. A. Fisher, “Quantum Zeno effect and the many-body entanglement transition,” *Phys. Rev. B* **98**, 205136 (2018).
- [27] A. Chan, R. M. Nandkishore, M. Pretko, and G. Smith, “Unitary-projective entanglement dynamics,” *Phys. Rev. B* **99**, 224307 (2019).
- [28] B. Skinner, J. Ruhman, and A. Nahum, “Measurement-Induced Phase Transitions in the Dynamics of Entanglement,” *Phys. Rev. X* **9**, 031009 (2019).
- [29] Y. Fuji and Y. Ashida, “Measurement-induced quantum criticality under continuous monitoring,” *Phys. Rev. B* **102**, 054302 (2020).
- [30] S. Goto and I. Danshita, “Measurement-induced transitions of the entanglement scaling law in ultracold gases with controllable dissipation,” *Phys. Rev. A* **102**, 033316 (2020).
- [31] Q. Tang and W. Zhu, “Measurement-induced phase transition: A case study in the nonintegrable model by density-matrix renormalization group calculations,” *Phys. Rev. Research* **2**, 013022 (2020).
- [32] M. Buchhold, Y. Minoguchi, A. Altland, and S. Diehl, “Effective Theory for the Measurement-Induced Phase Transition of Dirac Fermions,” *Phys. Rev. X* **11**, 041004 (2021).
- [33] M. Block, Y. Bao, S. Choi, E. Altman, and N. Y. Yao, “Measurement-Induced Transition in Long-Range Interacting Quantum Circuits,” *Phys. Rev. Lett.* **128**, 010604 (2022).
- [34] T. Minato, K. Sugimoto, T. Kuwahara, and K. Saito, “Fate of Measurement-Induced Phase Transition in Long-Range Interactions,” *Phys. Rev. Lett.* **128**, 010603 (2022).
- [35] H. Ott, “Single atom detection in ultracold quantum gases: a review of current progress,” *Rep. Prog. Phys.* **79**, 054401 (2016).
- [36] Y. Ashida, S. Furukawa, and M. Ueda, “Quantum critical behavior influenced by measurement backaction in ultracold gases,” *Phys. Rev. A* **94**, 053615 (2016).
- [37] Y. Ashida, S. Furukawa, and M. Ueda, “Parity-time-symmetric quantum critical phenomena,” *Nat. Commun.* **8**, 15791 (2017).
- [38] M. Nakagawa, N. Tsuji, N. Kawakami, and M. Ueda, “Dynamical Sign Reversal of Magnetic Correlations in Dissipative Hubbard Models,” *Phys. Rev. Lett.* **124**, 147203 (2020).
- [39] K. Yamamoto, M. Nakagawa, M. Tezuka, M. Ueda, and N. Kawakami, “Universal properties of dissipative Tomonaga-Luttinger liquids: Case study of a non-Hermitian XXZ spin chain,” *Phys. Rev. B* **105**, 205125 (2022).
- [40] M. Nakagawa, N. Kawakami, and M. Ueda, “Exact Liouvillian Spectrum of a One-Dimensional Dissipative Hubbard Model,” *Phys. Rev. Lett.* **126**, 110404 (2021).
- [41] C. Wu, J.-P. Hu, and S.-C. Zhang, “Exact $SO(5)$ Symmetry in the Spin-3/2 Fermionic System,” *Phys. Rev. Lett.* **91**, 186402 (2003).
- [42] C. Honerkamp and W. Hofstadter, “Ultracold Fermions and the $SU(N)$ Hubbard Model,” *Phys. Rev. Lett.* **92**, 170403 (2004).
- [43] R. W. Cherng, G. Refael, and E. Demler, “Superfluidity and Magnetism in Multicomponent Ultracold Fermions,” *Phys. Rev. Lett.* **99**, 130406 (2007).
- [44] M. A. Cazalilla, A. Ho, and M. Ueda, “Ultracold gases of ytterbium: ferromagnetism and Mott states in an $SU(6)$ Fermi

- system,” *New J. Phys.* **11**, 103033 (2009).
- [45] M. Hermele, V. Gurarie, and A. M. Rey, “Mott Insulators of Ultracold Fermionic Alkaline Earth Atoms: Underconstrained Magnetism and Chiral Spin Liquid,” *Phys. Rev. Lett.* **103**, 135301 (2009).
- [46] A. V. Gorshkov, M. Hermele, V. Gurarie, C. Xu, P. S. Julienne, J. Ye, P. Zoller, E. Demler, M. D. Lukin, and A. Rey, “Two-orbital SU (N) magnetism with ultracold alkaline-earth atoms,” *Nat. Phys.* **6**, 289 (2010).
- [47] M. A. Cazalilla and A. M. Rey, “Ultracold Fermi gases with emergent SU (N) symmetry,” *Rep. Prog. Phys.* **77**, 124401 (2014).
- [48] S. Capponi, P. Lecheminant, and K. Totsuka, “Phases of one-dimensional SU (N) cold atomic Fermi gases—From molecular Luttinger liquids to topological phases,” *Ann. Phys.* **367**, 50 (2016).
- [49] H. Yoshida and H. Katsura, “Rigorous Results on the Ground State of the Attractive SU(N) Hubbard Model,” *Phys. Rev. Lett.* **126**, 100201 (2021).
- [50] M. Nakagawa, H. Katsura, and M. Ueda, “Exact eigenstates of multicomponent Hubbard models: SU (N) magnetic η pairing, weak ergodicity breaking, and partial integrability,” *arXiv:2205.07235* (2022).
- [51] D. Greif, T. Uehlinger, G. Jotzu, L. Tarruell, and T. Esslinger, “Short-range quantum magnetism of ultracold fermions in an optical lattice,” *Science* **340**, 1307 (2013).
- [52] R. A. Hart, P. M. Duarte, T.-L. Yang, X. Liu, T. Paiva, E. Khatami, R. T. Scalettar, N. Trivedi, D. A. Huse, and R. G. Hulet, “Observation of antiferromagnetic correlations in the Hubbard model with ultracold atoms,” *Nature (London)* **519**, 211 (2015).
- [53] M. Boll, T. A. Hilker, G. Salomon, A. Omran, J. Nespolo, L. Pollet, I. Bloch, and C. Gross, “Spin-and density-resolved microscopy of antiferromagnetic correlations in Fermi-Hubbard chains,” *Science* **353**, 1257 (2016).
- [54] M. F. Parsons, A. Mazurenko, C. S. Chiu, G. Ji, D. Greif, and M. Greiner, “Site-resolved measurement of the spin-correlation function in the Fermi-Hubbard model,” *Science* **353**, 1253 (2016).
- [55] L. W. Cheuk, M. A. Nichols, K. R. Lawrence, M. Okan, H. Zhang, E. Khatami, N. Trivedi, T. Paiva, M. Rigol, and M. W. Zwierlein, “Observation of spatial charge and spin correlations in the 2D Fermi-Hubbard model,” *Science* **353**, 1260 (2016).
- [56] A. Mazurenko, C. S. Chiu, G. Ji, M. F. Parsons, M. Kanász-Nagy, R. Schmidt, F. Grusdt, E. Demler, D. Greif, and M. Greiner, “A cold-atom Fermi–Hubbard antiferromagnet,” *Nature* **545**, 462 (2017).
- [57] S. Taie, R. Yamazaki, S. Sugawa, and Y. Takahashi, “An SU (6) Mott insulator of an atomic Fermi gas realized by large-spin Pomeranchuk cooling,” *Nat. Phys.* **8**, 825 (2012).
- [58] F. Scazza, C. Hofrichter, M. Höfer, P. De Groot, I. Bloch, and S. Fölling, “Observation of two-orbital spin-exchange interactions with ultracold SU (N)-symmetric fermions,” *Nat. Phys.* **10**, 779 (2014).
- [59] X. Zhang, M. Bishof, S. L. Bromley, C. V. Kraus, M. S. Safronova, P. Zoller, A. M. Rey, and J. Ye, “Spectroscopic observation of SU (N)-symmetric interactions in Sr orbital magnetism,” *Science* **345**, 1467 (2014).
- [60] C. Hofrichter, L. Riegger, F. Scazza, M. Höfer, D. R. Fernandes, I. Bloch, and S. Fölling, “Direct probing of the mott crossover in the SU(n) fermi-hubbard model,” *Phys. Rev. X* **6**, 021030 (2016).
- [61] H. Ozawa, S. Taie, Y. Takasu, and Y. Takahashi, “Antiferromagnetic Spin Correlation of SU(N) Fermi Gas in an Optical Superlattice,” *Phys. Rev. Lett.* **121**, 225303 (2018).
- [62] D. Tusi, L. Franchi, L. Livi, K. Baumann, D. Benedicto Orenes, L. Del Re, R. Barfknecht, T.-W. Zhou, M. Inguscio, G. Cappellini, M. Capone, J. Catani, and L. Fallani, “Flavour-selective localization in interacting lattice fermions,” *Nat. Phys.* **18**, 1201 (2022).
- [63] S. Taie, E. Ibarra-García-Padilla, N. Nishizawa, Y. Takasu, Y. Kuno, H.-T. Wei, R. T. Scalettar, K. R. Hazzard, and Y. Takahashi, “Observation of antiferromagnetic correlations in an ultracold SU (N) Hubbard model,” *Nat. Phys.* (2022).
- [64] B. Yang, H. Sun, C.-J. Huang, H.-Y. Wang, Y. Deng, H.-N. Dai, Z.-S. Yuan, and J.-W. Pan, “Cooling and entangling ultracold atoms in optical lattices,” *Science* **369**, 550 (2020).
- [65] H. Sun, B. Yang, H.-Y. Wang, Z.-Y. Zhou, G.-X. Su, H.-N. Dai, Z.-S. Yuan, and J.-W. Pan, “Realization of a bosonic antiferromagnet,” *Nat. Phys.* **17**, 990 (2021).
- [66] L. Rosso, L. Mazza, and A. Biella, “Eightfold way to dark states in SU(3) cold gases with two-body losses,” *Phys. Rev. A* **105**, L051302 (2022).
- [67] L. Rosso, A. Biella, J. D. Nardis, and L. Mazza, “A dynamical theory for one-dimensional fermions with strong two-body losses: universal non-Hermitian Zeno physics and spin-charge separation,” *arXiv:2206.0683*.
- [68] F. D. M. Haldane, “General Relation of Correlation Exponents and Spectral Properties of One-Dimensional Fermi Systems: Application to the Anisotropic $S = \frac{1}{2}$ Heisenberg Chain,” *Phys. Rev. Lett.* **45**, 1358 (1980).
- [69] F. D. M. Haldane, “‘Luttinger liquid theory’ of one-dimensional quantum fluids. I. Properties of the Luttinger model and their extension to the general 1D interacting spinless Fermi gas,” *J. Phys. C* **14**, 2585 (1981).
- [70] F. D. M. Haldane, “Demonstration of the ‘Luttinger liquid’ character of Bethe-ansatz-soluble models of 1-D quantum fluids,” *Phys. Lett. A* **81**, 153 (1981).
- [71] F. D. M. Haldane, “Effective Harmonic-Fluid Approach to Low-Energy Properties of One-Dimensional Quantum Fluids,” *Phys. Rev. Lett.* **47**, 1840 (1981).
- [72] A. A. Belavin, A. M. Polyakov, and A. B. Zamolodchikov, “Infinite conformal symmetry in two-dimensional quantum field theory,” *Nucl. Phys. B* **241**, 333 (1984).
- [73] A. A. Belavin, A. M. Polyakov, and A. B. Zamolodchikov, “Infinite conformal symmetry of critical fluctuations in two dimensions,” *J. Stat. Phys.* **34**, 763 (1984).
- [74] D. Friedan, Z. Qiu, and S. Shenker, “Conformal Invariance, Unitarity, and Critical Exponents in Two Dimensions,” *Phys. Rev. Lett.* **52**, 1575 (1984).
- [75] E. Brézin and J. Zinn-Justin, eds., *Fields, strings and critical phenomena* (North-Holland, Amsterdam, 1988).
- [76] F. Calogero, “Ground State of a One-Dimensional N -Body System,” *J. Math. Phys.* **10**, 2197 (1969).
- [77] B. Sutherland, “Quantum Many-Body Problem in One Dimension: Ground State,” *J. Math. Phys.* **12**, 246 (1971).
- [78] B. Sutherland, “Quantum Many-Body Problem in One Dimension: Thermodynamics,” *J. Math. Phys.* **12**, 251 (1971).
- [79] B. Sutherland, “Exact Results for a Quantum Many-Body Problem in One Dimension,” *Phys. Rev. A* **4**, 2019 (1971).
- [80] B. Sutherland, “Exact Results for a Quantum Many-Body Problem in One Dimension. II,” *Phys. Rev. A* **5**, 1372 (1972).
- [81] F. D. M. Haldane, “Exact Jastrow-Gutzwiller Resonating-Valence-Bond Ground State of the Spin- $\frac{1}{2}$ Antiferromagnetic Heisenberg Chain with $1/r^2$ Exchange,” *Phys. Rev. Lett.* **60**, 635 (1988).

- [82] B. S. Shastry, “Exact Solution of an $S = \frac{1}{2}$ Heisenberg Antiferromagnetic Chain with Long-Range Interactions,” *Phys. Rev. Lett.* **60**, 639 (1988).
- [83] F. D. M. Haldane, ““Spinon Gas” Description of the $S = \frac{1}{2}$ Heisenberg Chain with Inverse-Square Exchange: Exact Spectrum and Thermodynamics,” *Phys. Rev. Lett.* **66**, 1529 (1991).
- [84] N. Kawakami and S.-K. Yang, “Finite-Size scaling in One-Dimensional Quantum Liquid with Long-Range Interaction,” *Phys. Rev. Lett.* **67**, 2493 (1991).
- [85] N. Kawakami, “Critical properties of quantum many-body systems with $1/r^2$ interaction,” *Prog. Theor. Phys.* **91**, 189 (1994).
- [86] F. D. M. Haldane, ““Fractional statistics” in Arbitrary Dimensions: A Generalization of the Pauli Principle,” *Phys. Rev. Lett.* **67**, 937 (1991).
- [87] N. Kawakami, “Novel Hierarchy of the SU(N) Electron Models and Edge States of Fractional Quantum Hall Effect,” *Phys. Rev. Lett.* **71**, 275 (1993).
- [88] R. B. Laughlin, “Anomalous Quantum Hall Effect: An Incompressible Quantum Fluid with Fractionally Charged Excitations,” *Phys. Rev. Lett.* **50**, 1395 (1983).
- [89] J. K. Jain, “Composite-Fermion Approach for the Fractional Quantum Hall Effect,” *Phys. Rev. Lett.* **63**, 199 (1989).
- [90] Z. N. C. Ha and F. D. M. Haldane, “Models with inverse-square exchange,” *Phys. Rev. B* **46**, 9359 (1992).
- [91] N. Kawakami, “Asymptotic Bethe-ansatz solution of multi-component quantum systems with $1/r^2$ long-range interaction,” *Phys. Rev. B* **46**, 1005 (1992).
- [92] N. Kawakami, “SU(N) generalization of the Gutzwiller-Jastrow wave function and its critical properties in one dimension,” *Phys. Rev. B* **46**, 3191(R) (1992).
- [93] N. Kawakami, “SU(N) Calogero-Sutherland Model and $t - J$ Model with $1/r^2$ Interaction,” *J. Phys. Soc. Jpn.* **62**, 2270 (1993).
- [94] N. Kawakami, “Correlation exponents for multicomponent Jastrow wave functions and effects of chiral constraint in one dimension,” *J. Phys. Soc. Jpn.* **62**, 2419 (1993).
- [95] D. Bernard, M. Gaudin, F. Haldane, and V. Pasquier, “Yang-Baxter equation in long-range interacting systems,” *J. Phys. A: Math. Gen.* **26**, 5219 (1993).
- [96] T. Fukui and N. Kawakami, “Breakdown of the Mott insulator: Exact solution of an asymmetric Hubbard model,” *Phys. Rev. B* **58**, 16051 (1998).
- [97] In the research field of high-energy physics, NH Calogero-Sutherland model with PT symmetry has been actively investigated (See reviews [145, 146] and references therein).
- [98] Y. Kuramoto and H. Yokoyama, “Exactly Soluble Supersymmetric $t - J$ -Type Model with Long-Range Exchange and Transfer,” *Phys. Rev. Lett.* **67**, 1338 (1991).
- [99] N. Kawakami, “Asymptotic Bethe ansatz: Application to the one-dimensional $t - J$ model with long-range exchange and transfer,” *Phys. Rev. B* **45**, 7525(R) (1992).
- [100] K. Hikami and M. Wadati, “Integrability of Calogero-Moser spin system,” *J. Phys. Soc. Jpn.* **62**, 469 (1993).
- [101] K. Hikami and M. Wadati, “Integrable spin- $\frac{1}{2}$ particle systems with long-range interactions,” *Phys. Lett. A* **173**, 263 (1993).
- [102] K. Hikami and M. Wadati, “Infinite symmetry of the spin systems with inverse square interactions,” *J. Phys. Soc. Jpn.* **62**, 4203 (1993).
- [103] B. Sutherland, *Beautiful models: 70 years of exactly solved quantum many-body problems* (World Scientific, 2004).
- [104] N. Shibata and H. Katsura, “Dissipative quantum Ising chain as a non-Hermitian Ashkin-Teller model,” *Phys. Rev. B* **99**, 224432 (2019).
- [105] B. Buča, C. Booker, M. Medenjak, and D. Jaksch, “Bethe ansatz approach for dissipation: exact solutions of quantum many-body dynamics under loss,” *New J. Phys.* **22**, 123040 (2020).
- [106] J. L. Cardy, “Conformal invariance and universality in finite-size scaling,” *J. Phys. A* **17**, L385 (1984).
- [107] J. L. Cardy, “Finite-size scaling in strips: antiperiodic boundary conditions,” *J. Phys. A* **17**, L961 (1984).
- [108] J. L. Cardy, “Operator content of two-dimensional conformally invariant theories,” *Nucl. Phys. B* **270**, 186 (1986).
- [109] J. L. Cardy, “Logarithmic corrections to finite-size scaling in strips,” *J. Phys. A* **19**, L1093 (1986).
- [110] H. W. J. Blöte, J. L. Cardy, and M. P. Nightingale, “Conformal Invariance, the Central Charge, and Universal Finite-Size Amplitudes at Criticality,” *Phys. Rev. Lett.* **56**, 742 (1986).
- [111] I. Affleck, “Universal Term in the Free Energy at a Critical Point and the Conformal Anomaly,” *Phys. Rev. Lett.* **56**, 746 (1986).
- [112] C. N. Yang, “Some Exact Results for the Many-Body Problem in One Dimension with Repulsive Delta-Function Interaction,” *Phys. Rev. Lett.* **19**, 1312 (1967).
- [113] B. Sutherland, “Further Results for the Many-Body Problem in One Dimension,” *Phys. Rev. Lett.* **20**, 98 (1968).
- [114] B. Sutherland, “Model for a multicomponent quantum system,” *Phys. Rev. B* **12**, 3795 (1975).
- [115] N. Kawakami and S.-K. Yang, “Luttinger anomaly exponent of momentum distribution in the Hubbard chain,” *Phys. Lett. A* **148**, 359 (1990).
- [116] H. J. Schulz, “Correlation Exponents and the Metal-Insulator Transition in the One-Dimensional Hubbard Model,” *Phys. Rev. Lett.* **64**, 2831 (1990).
- [117] H. Frahm and V. E. Korepin, “Critical exponents for the one-dimensional Hubbard model,” *Phys. Rev. B* **42**, 10553 (1990).
- [118] V. G. Kac and D. H. Peterson, “Infinite-dimensional Lie algebras, theta functions and modular forms,” *Adv. Math.* **53**, 125 (1984).
- [119] V. Knizhnik and A. Zamolodchikov, “Current algebra and Wess-Zumino model in two dimensions,” *Nucl. Phys. B* **247**, 83 (1984).
- [120] E. Witten, “Non-abelian bosonization in two dimensions,” *Commun. Math. Phys.* **92**, 455 (1984).
- [121] P. Goddard and D. Olive, “Kac-Moody and Virasoro algebras in relation to quantum physics,” *Int. J. Mod. Phys. A* **1**, 303 (1986).
- [122] A. Polyakov and P. B. Wiegmann, “Theory of nonabelian Goldstone bosons in two dimensions,” *Phys. Lett. B* **131**, 121 (1983).
- [123] A. Izergin, V. Korepin, and N. Y. Reshetikhin, “Conformal dimensions in Bethe ansatz solvable models,” *J. Phys. A: Math. Gen.* **22**, 2615 (1989).
- [124] M. Höning, M. Moos, and M. Fleischhauer, “Critical exponents of steady-state phase transitions in fermionic lattice models,” *Phys. Rev. A* **86**, 013606 (2012).
- [125] A. Bácsi, C. P. Moca, and B. Dóra, “Dissipation-Induced Luttinger Liquid Correlations in a One-Dimensional Fermi Gas,” *Phys. Rev. Lett.* **124**, 136401 (2020).
- [126] J. M. Torres, “Closed-form solution of Lindblad master equations without gain,” *Phys. Rev. A* **89**, 052133 (2014).
- [127] R. Couvreur, J. L. Jacobsen, and H. Saleur, “Entanglement in Nonunitary Quantum Critical Spin Chains,” *Phys. Rev. Lett.* **119**, 040601 (2017).
- [128] P.-Y. Chang, J.-S. You, X. Wen, and S. Ryu, “Entanglement spectrum and entropy in topological non-Hermitian systems and nonunitary conformal field theory,” *Phys. Rev. Research*

- 2, 033069 (2020).
- [129] K. Kawabata, T. Numasawa, and S. Ryu, “Entanglement Phase Transition Induced by the Non-Hermitian Skin Effect,” arXiv:2206.05384.
 - [130] Y. Li, X. Chen, and M. P. A. Fisher, “Measurement-driven entanglement transition in hybrid quantum circuits,” Phys. Rev. B **100**, 134306 (2019).
 - [131] C.-M. Jian, Y.-Z. You, R. Vasseur, and A. W. W. Ludwig, “Measurement-induced criticality in random quantum circuits,” Phys. Rev. B **101**, 104302 (2020).
 - [132] X. Chen, Y. Li, M. P. A. Fisher, and A. Lucas, “Emergent conformal symmetry in nonunitary random dynamics of free fermions,” Phys. Rev. Research **2**, 033017 (2020).
 - [133] Y. Li, X. Chen, A. W. W. Ludwig, and M. P. A. Fisher, “Conformal invariance and quantum nonlocality in critical hybrid circuits,” Phys. Rev. B **104**, 104305 (2021).
 - [134] T. Yoshida, K. Kudo, and Y. Hatsugai, “Non-Hermitian fractional quantum Hall states,” Sci. Rep. **9**, 16895 (2019).
 - [135] T. Yoshida, K. Kudo, H. Katsura, and Y. Hatsugai, “Fate of fractional quantum Hall states in open quantum systems: Characterization of correlated topological states for the full Liouvillian,” Phys. Rev. Research **2**, 033428 (2020).
 - [136] S. Taie, Y. Takasu, S. Sugawa, R. Yamazaki, T. Tsujimoto, R. Murakami, and Y. Takahashi, “Realization of a $SU(2) \times SU(6)$ system of fermions in a cold atomic gas,” Phys. Rev. Lett. **105**, 190401 (2010).
 - [137] M. Buchhold and S. Diehl, “Nonequilibrium universality in the heating dynamics of interacting Luttinger liquids,” Phys. Rev. A **92**, 013603 (2015).
 - [138] J.-S. Bernier, R. Tan, C. Guo, C. Kollath, and D. Poletti, “Melting of the critical behavior of a Tomonaga-Luttinger liquid under dephasing,” Phys. Rev. B **102**, 115156 (2020).
 - [139] B. Dóra and C. P. Moca, “Quantum Quench in \mathcal{PT} -Symmetric Luttinger Liquid,” Phys. Rev. Lett. **124**, 136802 (2020).
 - [140] Á. Bácsi, C. P. Moca, G. Zarand, and B. Dóra, “Dissipative dynamics in the free massive boson limit of the sine-Gordon model,” SciPost Phys. Core **5**, 004 (2022).
 - [141] A. Bácsi, C. P. Moca, G. Zaránd, and B. Dóra, “Vaporization Dynamics of a Dissipative Quantum Liquid,” Phys. Rev. Lett. **125**, 266803 (2020).
 - [142] C. P. Moca and B. Dóra, “Universal conductance of a \mathcal{PT} -symmetric luttinger liquid after a quantum quench,” Phys. Rev. B **104**, 125124 (2021).
 - [143] B. Dóra, D. Sticlet, and C. P. Moca, “Correlations at PT-Symmetric Quantum Critical Point,” Phys. Rev. Lett. **128**, 146804 (2022).
 - [144] B. Dóra and C. P. Moca, “Full counting statistics in the many-body hatano-nelson model,” arXiv:2208.12017.
 - [145] P. K. Ghosh, “Supersymmetric many-particle quantum systems with inverse-square interactions,” J. Phys. A: Math. Theor. **45**, 183001 (2012).
 - [146] A. Fring, “ \mathcal{PT} -symmetric deformations of integrable models,” Philos. Trans. Royal Soc. A **371**, 20120046 (2013).