

# G-dual teleparallel connections in Information Geometry

F. M. Ciaglia<sup>1,5</sup> , F. Di Cosmo<sup>1,2,6</sup> , A. Ibort<sup>1,2,7</sup> , G. Marmo<sup>3,4,8</sup> 

August 29, 2023

<sup>1</sup> Department of Mathematics, University Carlos III de Madrid, Leganés, Madrid, Spain

<sup>2</sup> ICMAT, Instituto de Ciencias Matemáticas (CSIC-UAM-UC3M-UCM)

<sup>3</sup> INFN-Sezione di Napoli, Naples, Italy

<sup>4</sup> Department of Physics “E. Pancini”, University of Naples Federico II, Naples, Italy

<sup>5</sup>fciaglia[at]math.uc3m.es, <sup>6</sup>fcosmo[at]math.uc3m.es, <sup>7</sup>albertoi[at]math.uc3m.es, <sup>8</sup>marmo[at]na.infn.it

## Abstract

Given a real, finite-dimensional, smooth parallelizable Riemannian manifold  $(\mathcal{N}, G)$  endowed with a teleparallel connection  $\nabla$  determined by a choice of a global basis of vector fields on  $\mathcal{N}$ , we show that the  $G$ -dual connection  $\nabla^*$  of  $\nabla$  in the sense of Information Geometry must be the teleparallel connection determined by the basis of  $G$ -gradient vector fields associated with a basis of differential one-forms which is (almost) dual to the basis of vector fields determining  $\nabla$ . We call any such pair  $(\nabla, \nabla^*)$  a  *$G$ -dual teleparallel pair*. Then, after defining a covariant  $(0, 3)$  tensor  $T$  uniquely determined by  $(\mathcal{N}, G, \nabla, \nabla^*)$ , we show that  $T$  being symmetric in the first two entries is equivalent to  $\nabla$  being torsion-free, that  $T$  being symmetric in the first and third entry is equivalent to  $\nabla^*$  being torsion free, and that  $T$  being symmetric in the second and third entries is equivalent to the basis vectors determining  $\nabla$  ( $\nabla^*$ ) being parallel-transported by  $\nabla^*$  ( $\nabla$ ). Therefore,  $G$ -dual teleparallel pairs provide a generalization of the notion of Statistical Manifolds usually employed in Information Geometry, and we present explicit examples of  $G$ -dual teleparallel pairs arising both in the context of both Classical and Quantum Information Geometry.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>G-dual teleparallel pairs</b>	<b>4</b>
<b>3</b>	<b>G-dual teleparallel pairs in Classical Information Geometry</b>	<b>9</b>
<b>4</b>	<b>G-dual teleparallel pairs in Quantum Information Geometry</b>	<b>11</b>
<b>5</b>	<b>Conclusions</b>	<b>15</b>
	<b>References</b>	<b>16</b>

# 1 Introduction

From a purely mathematical point of view, the structure theory of Classical Information Geometry is the theory of Riemannian smooth manifolds with an additional structure. This additional structure may be described, equivalently, using a pair of affine connections or a covariant  $(0, 3)$  tensor (cf. [2, 4, 40]). Specifically, in the first case we have a Riemannian smooth manifold  $(\mathcal{N}, G)$ , whose points parametrize probability distributions on a suitable outcome space, a torsion-free affine connection  $\nabla$ , and another torsion-free affine connection  $\nabla^*$  which is dual to  $\nabla$  with respect to  $G$  in the sense that

$$X(G(Y, Z)) = G(\nabla_X Y, Z) + G(Y, \nabla_X^* Z) \quad (1)$$

for all vector fields  $X, Y, Z$  on  $\mathcal{N}$ . In general,  $\nabla \neq \nabla^*$  unless  $\nabla$  is the Levi-Civita connection  $\nabla^G$  of  $G$ . However, both  $\nabla$  and  $\nabla^*$  can be expressed in terms of  $\nabla^G$  and a covariant tensor essentially because the space of affine connections is, quite appropriately, an affine space modelled on a vector space of tensor fields. Specifically, it holds

$$G(\nabla_X Y, Z) - G(\nabla_X^G Y, Z) = T(X, Y, Z) \quad (2)$$

with  $T$  a covariant  $(0, 3)$ -tensor field on  $\mathcal{N}$  which may also be written according to

$$G(Y, \nabla_X^G Z) - G(Y, \nabla_X^* Z) = T(X, Y, Z). \quad (3)$$

because of equation (1) and because  $\nabla^G$  is its own  $g$ -dual connection. If both  $\nabla$  and  $\nabla^*$  are torsion-free, it can be proved that  $T$  is completely symmetric.

If instead of the pair  $(\nabla, \nabla^*)$  of torsion-free,  $G$ -dual affine connections we endow the Riemannian smooth manifold  $(\mathcal{N}, G)$  with a covariant  $(0, 3)$ -tensor field  $T$  which is totally symmetric, we can again obtain a pair of torsion-free,  $G$ -dual affine connections simply reading equation (2) and (3) from right to left. Therefore, we conclude that the two approaches are equivalent. The triple  $(\mathcal{N}, G, T)$  is referred to as a **statistical manifold** (cf. [40]), and the tensor  $T$  is often called **Amari-Čencov** tensor in honour of the two pioneers who thoroughly studied this structure (cf. [2, 6, 56]).

It is worth noting how recently the theory of statistical manifolds has been approached from the point of view of Lie groupoids (cf. [30, 31, 32]) where it is shown how Information Geometry on statistical manifolds is but a particular example of Information Geometry on Lie groupoids. This point of view on Information Geometry may be particularly fruitful in connection with the recent reformulation of Quantum Theories in the framework of groupoids (cf. [13, 17, 18]), especially considering how this reformulation of Quantum theories heavily relies on a categorical background which shares many interesting similarities with the one exploited by Čencov in his pioneering work (cf. [56]). However, we only plan to investigate these matters in future works.

Coming back to statistical manifolds, of particular interest are those statistical manifolds whose Amari-Čencov tensor  $T$  gives rise to a pair  $(\nabla, \nabla^*)$  of  $G$ -dual connections which are both flat in the sense that, in addition to being torsion-free, their curvature vanishes. In this case, the statistical manifold is called **dually flat** and some interesting properties of both applied and purely mathematical flavour present themselves (cf. [3, 4, 50]). Of course, the request of flatness for the connections immediately imposes topological obstructions on the type of smooth manifold we can consider, and these kind of considerations also apply to the case of Quantum Information Geometry of finite-dimensional systems (cf. [7, 8, 27, 39]).

A typical example of a dually flat statistical manifold is given by the interior of the  $n$ -simplex endowed with the Fisher-Rao metric tensor  $G_{FR}$  and with the pair of  $G_{FR}$ -dual affine connections known as the *mixture* and *exponential connection*. The mixture connection on the interior of the  $n$ -simplex has a kind of privileged role because it is intimately connected with the convex structure of the set of probability distributions. We refer to section 3 for a thorough discussion of this example.

It is well-known that a statistical manifold  $(\mathcal{N}, G, T)$  can be built suitably expanding up to third order a smooth function  $S: \mathcal{N} \times \mathcal{N} \rightarrow \mathbb{R}$  (also known as a two-point function on  $\mathcal{N}$ ) satisfying some specific conditions (cf. [5, 14, 43]). It often happens that the function  $S$  is a relative entropy function, or a divergence function. The most relevant family of relative entropies in the classical setting is that investigated by Csizar (cf. [21]), and it turns out that every member in this family gives rise to the same Riemannian metric tensor which is precisely the Fisher-Rao metric tensor.

In the (finite-dimensional) quantum setting, it turns out that the situation is essentially different than in the classical setting. First of all, on the manifold of faithful quantum states  $\mathcal{S}(\mathcal{H})$  of a physical system described by a finite-dimensional complex Hilbert space  $\mathcal{H}$  (cf. section 4) there is an infinite number of Riemannian metric tensors whose theoretical (and often also applied) relevance has been thoroughly corroborated. These Riemannian metric tensors are known as *quantum monotone metric tensors* and have been completely classified by Petz (cf. [48]). Every quantum monotone metric tensor may be obtained suitably expanding a quantum relative entropy which, in some sense, generalizes the classical relative entropy of Csizar mentioned above (cf. [12, 41]). These instances are in stark contrast with what happens in the classical case where the Riemannian structure of essentially all the statistical manifolds considered are given by the Fisher-Rao metric tensor<sup>1</sup>, and where every relative entropy in the family investigated by Csizar leads to the Fisher-Rao metric tensor.

Of course, the dual pairs of affine connections generated by the quantum relative entropies associated with the quantum monotone metric tensors are necessarily torsion-free. In particular, it is possible to build a mixture connection on  $\mathcal{S}(\mathcal{H})$  because, analogously to the space of classical probability distributions, the space of quantum states is a convex set (cf. section 4), and it turns out that there is only one monotone metric tensor - the so-called Bogoliubov-Kubo-Mori metric tensor (cf. [45, 47, 49]) - for which the mixture connection admits a dual connection which is the quantum analogue of the exponential connection and which is torsion-free (cf. [34, 44]). In this case, the associated quantum relative entropy is the von Neumann-Umegaki relative entropy (cf. [24, 55, 57]). Incidentally, the resulting statistical manifold is dually flat.

However, it turns out that there are pairs of affine connections which are dual with respect to the quantum monotone metric tensors and are not both torsion-free (cf. [4, 26, 38]). In particular, examples of these dual pair with non-vanishing torsion are found when generalizing Amari's  $\alpha$ -representation (cf. [2]) to the quantum case (cf. [38]). Moreover, one of the results of this work is to show that the affine connection which is dual to the mixture connection on  $\mathcal{S}(\mathcal{H})$  with respect to a specific quantum monotone metric tensor always has non-vanishing torsion unless the gradient vector fields of the expectation value functions of quantum mechanical observables commute. Since the mixture connection is intimately connected with the convex structure of the space of quantum states and thus reflects a structural feature of  $\mathcal{S}(\mathcal{H})$  that goes

---

<sup>1</sup>We are here deliberately ignoring all the Wasserstein-type metric tensors not because we believe they are not relevant but only because their very definition depend on additional structures (e.g., a metric distance on the outcome space in the classical case or a metric distance on the space of pure states in the quantum case) that are not of interest for us here.

beyond the formalism of Information Geometry, the idea of discarding all the cases in which the affine connection dual to the mixture one presents torsion seems too restrictive. Indeed, already Amari and Nagaoka in their seminal work [4, p. 19] noted how “the incorporation of torsion into the framework of information geometry, which would relate it to such fields as quantum mechanics (noncommutative probability theory) and systems theory, is an interesting topic for the future.” Consequently, it seems that the notion of statistical manifold is not enough to completely understand the mathematical structures arising in Quantum Information Geometry.

Moreover, it is worth noting how the investigation of the appearance of torsion also in Classical Information Geometry is recently gaining interest (*cf.* [37, 59, 60, 61]). In particular, the present work may be thought of as complementing some theoretical aspects already discussed in [59] (especially in connection with the explicit construction of what is called the g-biorthogonal frame), and providing a natural extension to the case of Quantum Information Geometry.

The aim of this work is to introduce a *variation on the theme* of dually flat connections that allows the appearance of torsion and to show that this type of geometric structure naturally appears in Classical and Quantum Information Geometry. The starting point is the idea that, at least locally, a flat connection  $\nabla$  on a smooth manifold  $\mathcal{N}$  admits a basis of vector fields  $\{X_j\}_{j=1,\dots,n=\dim(\mathcal{N})}$  such that  $\nabla_{X_j}X_k = 0$  for all  $j, k$ . If the vector fields are globally defined so that  $\{X_j\}_{j=1,\dots,n=\dim(\mathcal{N})}$  is a basis for the module of vector fields on  $\mathcal{N}$  (note that this requires  $\mathcal{N}$  to be parallelizable), then the condition  $\nabla_{X_j}X_k = 0$  is what makes  $\nabla$  be the so-called teleparallel connection associated with  $\{X_j\}_{j=1,\dots,n=\dim(\mathcal{N})}$ . Moreover, a teleparallel connection associated with a basis  $\{X_j\}_{j=1,\dots,n=\dim(\mathcal{N})}$  of vector fields is uniquely determined by the condition  $\nabla_{X_j}X_k = 0$  for all  $j, k$ . In general, however, teleparallel connections need not be flat (i.e., curvature-free), nor even torsion-free (indeed, they are torsion-free if and only if the basis vector fields mutually commute as it can be easily seen from the very definition of the torsion tensor). In almost all the cases arising from Classical and Quantum Information Geometry, it turns out that there is one preferred teleparallel connection which is basically the mixture connection arising from the convex structure characteristic of both the space of probability distributions and the space of quantum states.

Motivated by the previous discussion, in section 2, given a fixed teleparallel connection on a smooth Riemannian manifold  $(\mathcal{N}, G)$ , we investigate the properties of the  $G$ -dual connection  $\nabla^*$  and show that it must necessarily be itself a teleparallel connection. We call any such pair  $(\nabla, \nabla^*)$  a ***G-dual teleparallel pair***. Moreover, we also investigate the symmetry properties of the tensor  $T$  generalizing the Amari-Čencov tensor to the case of  $G$ -dual teleparallel pairs. Then, in section 3 we review the case of the mixture and exponential connection on the interior  $\Delta_n$  of the  $n$ -simplex highlighting the teleparallel structure of both connections, and showing how the teleparallel structure of the exponential connection is connected with a particular homogeneous manifold structure for  $\Delta_n$ . In section 4 we investigate the quantum case and characterize the  $G_f$ -dual teleparallel pairs  $(\nabla, \nabla^*)$  on the manifold  $\mathcal{S}(\mathcal{H})$  of faithful quantum states in finite dimensions endowed with a monotone quantum metric tensor  $G_f$  in which  $\nabla$  is the mixture connection reflecting the convex structure of  $\mathcal{S}(\mathcal{H})$ . Finally, in section 5 we gather some concluding remarks and future perspectives.

## 2 G-dual teleparallel pairs

Let  $\mathcal{N}$  be a parallelizable smooth manifold and denote with  $\mathfrak{X}(\mathcal{N})$  the module of smooth vector fields on  $\mathcal{N}$ . Let  $\{X_j\}_{j=1,\dots,n}$ , with  $n = \dim(\mathcal{N})$ , be a global basis of vector fields on  $\mathcal{N}$ . Define

the *teleparallel connection*<sup>2</sup>  $\nabla$  associated with  $\{X_j\}_{j=1,\dots,n}$  by setting

$$\nabla_{X_j} X_k := 0 \quad \forall j, k = 1, \dots, n. \quad (4)$$

It is clear that every integral curve of  $X_j$  is a geodesic of  $\nabla$  for every  $j = 1, \dots, n$ . Moreover, since geodesics with fixed initial conditions are unique, every geodesic of  $\nabla$  starting at  $x \in \mathcal{N}$  with initial velocity  $v_x \in T_x \mathcal{N}$  can be realized as an integral curve starting at  $x$  of the vector field  $X = a^j X_j$  with  $a^j \in \mathbb{R}$  for all  $j = 1, \dots, n$  such that  $v_x = X(x)$ .

The torsion tensor  $T^\nabla$  of  $\nabla$ , in general, reads

$$T^\nabla(Z, W) := \nabla_Z W - \nabla_W Z - [Z, W] \quad (5)$$

for every  $Z, W \in \mathfrak{X}(\mathcal{N})$ . Therefore, equation (4) implies

$$T^\nabla(X_j, X_k) = [X_k, X_j] \quad (6)$$

for every  $j, k = 1, \dots, n$ , which means that, from the perspective of the basis  $\{X_j\}_{j=1,\dots,n}$ , the torsion tensor is connected with the commutator of the basis vector fields. Of course, from equation (6) it follows that  $\nabla$  is torsion-free if and only if the basis vector fields mutually commute.

The curvature tensor  $R^\nabla$  of  $\nabla$  reads

$$(R^\nabla(Z, W))(V) := \nabla_Z(\nabla_W V) - \nabla_W(\nabla_Z V) - \nabla_{[Z, W]} V \quad (7)$$

for every  $Z, W, V \in \mathfrak{X}(\mathcal{N})$ . Therefore, again equation (4) implies

$$(R^\nabla(X_j, X_k))(X_l) := -\nabla_{[X_j, X_k]} X_l = 0 \quad (8)$$

for every  $j, k, l = 1, \dots, n$ , because  $\{X_j\}_{j=1,\dots,n}$  is a basis of vector fields and thus  $[X_j, X_k] = F^l X_l$  with some smooth functions  $F^l$  globally defined on  $\mathcal{N}$ . We thus conclude that if  $\nabla$  is torsion-free then it is necessarily flat (in the sense of being free of curvature and torsion).

Let  $G$  be a smooth Riemannian metric tensor on  $\mathcal{N}$ . Motivated by the theory of Statistical Manifolds, we want to investigate the pairs  $(\nabla, \nabla^*)$  of  $G$ -dual connections on  $\mathcal{N}$  for which both  $\nabla$  and  $\nabla^*$  are teleparallel connections on  $\mathcal{N}$ . We call any such pair  $(\nabla, \nabla^*)$  a ***G-dual teleparallel pair***.

At this purpose, it is instrumental that we understand how  $\nabla$  behaves on differential one-forms. First of all, we consider a basis  $\{\theta^j\}_{j=1,\dots,n}$  of globally-defined differential one-forms on  $\mathcal{N}$  such that

$$\theta^j(X_k) = C_k^j. \quad (9)$$

where  $C_k^j$  is a number for every  $j, k = 1, \dots, n$ . We call any such basis ***almost dual*** to  $\{X_j\}_{j=1,\dots,n}$ . In particular, when  $C_k^j = \delta_k^j$  we obtain precisely the dual basis of  $\{X_j\}_{j=1,\dots,n}$ . We may think of  $C_{jk}$  as a scaled version of  $\delta_{jk}$  where the value at  $j = k$  actually depends on  $j$  and it is not necessarily always 1. The fact that all the constructions presented in the work are valid for  $C_{jk}$  instead of just  $\delta_{jk}$  simply reflects the fact that the choice of each single vector field in the basis is determined up to a constant.

---

<sup>2</sup>The term *teleparallel* may be translated with 'distantly parallel', and its use is mainly due to the fact that the affine connection we are interested in is used in the context of the so-called *Teleparallel Gravity* [1]. Note, however, that the teleparallel connection is also known as the **Weitzenböck connection** of  $\{X_j\}_{j=1,\dots,n}$ .

According to the general theory of affine connections (cf. [10]), the behaviour of  $\nabla$  on the almost dual basis  $\{\theta^j\}_{j=1,\dots,n}$  is encoded in the equation

$$(\nabla_Z \theta^j)(W) = \nabla_Z (\theta^j(W)) - \theta^j(\nabla_Z W) \quad (10)$$

where  $Z, W$  are arbitrary vector fields on  $\mathcal{N}$ . In particular, when  $Z = X_k$  and  $W = X_l$ , equation (10) becomes

$$(\nabla_{X_k} \theta^j)(X_l) = 0 \quad \forall j, k, l = 1, \dots, n \quad (11)$$

because of equation (4), and because the covariant derivatives of the  $C_{jk}$ 's in equation (9) vanish being the  $C_{jk}$ 's constant.

Equation (11) is equivalent to

$$\nabla_Z \theta^j = 0 \quad \forall j = 1, \dots, n \quad \forall Z \in \mathfrak{X}(\mathcal{N}) \quad (12)$$

because  $\{X_j\}_{j=1,\dots,n}$  is a basis for  $\mathfrak{X}(\mathcal{N})$  and, for every smooth function  $f$  on  $\mathcal{N}$ , it holds  $\nabla_f Z \alpha = f \nabla_Z \alpha$  for every differential one-form  $\alpha$  on  $\mathcal{N}$ .

Since the Riemannian metric tensor  $G$  is non-degenerate, for every  $\theta^j$  we can define its associated  $G$ -gradient (or simply 'gradient' if there is no risk of confusion) vector field  $Y_j$  uniquely determined by

$$\theta^j(Z) = G(Y_j, Z) \quad (13)$$

for every  $Z \in \mathfrak{X}(\mathcal{N})$ . Note that  $\{Y_j\}_{j=1,\dots,n}$  is a basis of globally-defined vector fields on  $\mathcal{N}$ . Setting  $Z = Y_k$  and  $W = X_l$  and exploiting equation (12), it follows that equation (10) becomes

$$0 = \theta^j(\nabla_{Y_k} X_l) \quad (14)$$

for all  $j, k, l = 1, \dots, n$ , which is equivalent to

$$\nabla_{Y_k} X_l = 0 \quad (15)$$

for all  $k, l = 1, \dots, n$ .

With these elements at our disposal, we now characterize the affine connection  $\nabla^*$  which is  $G$ -dual to the teleparallel connection  $\nabla$ . In particular, it is proved that  $\nabla^*$  is the teleparallel connection associated with  $\{Y_j\}_{j=1,\dots,n}$ . It is worth noting that the content of proposition 1 is essentially the same as that of Theorem 10 in [59].

**Proposition 1.** *Let  $\mathcal{N}$  be a parallelizable smooth manifold of dimension  $n = \dim(\mathcal{N})$ . Let  $\{X_j\}_{j=1,\dots,n}$  be the global basis of vector fields on  $\mathcal{N}$ , let  $\nabla$  be its associated teleparallel connection, and let  $\{\theta^j\}_{j=1,\dots,n}$  be an almost dual basis of differential one-forms for  $\{X_j\}_{j=1,\dots,n}$  (cf. equation (9)). For any smooth Riemannian metric tensor  $G$  on  $\mathcal{N}$ , let  $\{Y_j\}_{j=1,\dots,n}$  be the global basis of vector fields on  $\mathcal{N}$  made up of  $G$ -gradient vector fields of the  $\theta^j$ 's (cf. equation (13)). Then, the  $G$ -dual affine connection  $\nabla^*$  of  $\nabla$  (cf. equation (1)) is the teleparallel connection associated with  $\{Y_j\}_{j=1,\dots,n}$ .*

*Proof.* The  $G$ -dual connection  $\nabla^*$  is uniquely characterized by

$$Z(G(W, V)) = G(\nabla_Z W, V) + G(W, \nabla_Z^* V) \quad (16)$$

for every  $Z, W, V \in \mathfrak{X}(\mathcal{N})$ . Note that

$$Y_j(G(Y_k, X_l)) \stackrel{(13)}{=} Y_j(\theta^k(X_l)) \stackrel{(9)}{=} Y_j(C_l^k) = 0. \quad (17)$$



Therefore, setting  $Z = Y_j$ ,  $W = X_k$ , and  $V = Y_l$ , equation (16) becomes

$$0 \stackrel{(17)}{=} Y_j(G(X_k, Y_l)) = G(\nabla_{Y_j} X_k, Y_l) + G(X_k, \nabla_{Y_j}^* Y_l) \stackrel{(15)}{=} G(X_k, \nabla_{Y_j}^* Y_l) \quad (18)$$

for all  $j, k, l = 1, \dots, n$ , which is equivalent to

$$\nabla_{Y_j}^* Y_l = 0 \quad (19)$$

for all  $j, l = 1, \dots, n$  because  $\{X_j\}_{j=1, \dots, n}$  is a basis of vector fields on  $\mathcal{N}$ . We thus conclude that the  $G$ -dual connection  $\nabla^*$  is precisely the teleparallel connection determined by the basis  $\{Y_j\}_{j=1, \dots, n}$  of gradient vector fields on  $\mathcal{N}$  associated with the almost dual basis  $\{\theta^j\}_{j=1, \dots, n}$  of  $\{X_j\}_{j=1, \dots, n}$  through  $G$  as claimed.  $\square$   $\square$

**Remark 1.** We can introduce the basis  $\{\alpha^j\}_{j=1, \dots, n}$  of differential one-forms where each  $\alpha^j$  is the  $G$ -dual of  $X_j$  according to

$$\alpha^j(Z) = G(X_j, Z) \quad (20)$$

for every  $Z \in \mathfrak{X}(\mathcal{N})$ . Then, it follows that  $\{\alpha^j\}_{j=1, \dots, n}$  is an almost dual basis of  $\{Y_j\}_{j=1, \dots, n}$  because

$$\alpha^j(Y_k) \stackrel{(20)}{=} G(X_j, Y_k) \stackrel{(13)}{=} \theta^k(X_j) \stackrel{(9)}{=} C_j^k. \quad (21)$$

Consequently, proceeding in perfect analogy with what is done to obtain equation (15), we obtain

$$\nabla_{X_k}^* Y_l = 0 \quad (22)$$

for all  $k, l = 1, \dots, n$ .

**Remark 2.** Recalling the formula

$$d\beta(Z, W) = Z(\beta(W)) - W(\beta(Z)) - \beta([Z, W]) \quad (23)$$

valid for every differential one-form  $\beta$  and every pair of vector fields  $Z, W$  on  $\mathcal{N}$ , we immediately see that

$$d\theta^j(X_k, X_l) = X_k(\theta^j(X_l)) - X_l(\theta^j(X_k)) - \theta^j([X_k, X_l]) \stackrel{(9)}{=} -\theta^j([X_k, X_l]), \quad (24)$$

which means that  $\nabla$  is torsion-free (cf. equation (6)) if and only if  $\{\theta^j\}_{j=1, \dots, n}$  is a basis of closed differential one-forms. Similarly, we have

$$d\alpha^j(Y_k, Y_l) = Y_k(\alpha^j(Y_l)) - Y_l(\alpha^j(Y_k)) - \alpha^j([Y_k, Y_l]) \stackrel{(21)}{=} -\alpha^j([Y_k, Y_l]) \quad (25)$$

which means that  $\nabla^*$  is torsion-free (cf. equation (6)) if and only if  $\{\alpha^j\}_{j=1, \dots, n}$  is a basis of closed differential one-forms.

If  $(\nabla, \nabla^*)$  is a  $G$ -dual pair on  $(\mathcal{N}, G)$  and  $\nabla^G$  is the Levi-Civita connection of  $G$ , we may introduce the covariant  $(0, 3)$  tensor  $T$  setting

$$T(Z, W, V) := G(\nabla_Z W, V) - G(\nabla_Z^G W, V) \quad (26)$$

for every  $Z, W, V \in \mathfrak{X}(\mathcal{N})$ . Since  $\nabla^*$  is the  $G$ -dual connection of  $\nabla$ , equation (26) can be re-written in terms of  $\nabla^*$  as

$$T(Z, W, V) := G(W, \nabla_Z^G V) - G(W, \nabla_Z^* V). \quad (27)$$

If both  $\nabla$  and  $\nabla^*$  are torsion-free, then  $T$  is completely symmetric and it is precisely the Amari-Čencov tensor of the Statistical Manifold  $(\mathcal{N}, G, T)$ . In general, however,  $T$  is no longer completely symmetric and we now investigate its properties when  $(\nabla, \nabla^*)$  is a  $G$ -dual teleparallel pair.

**Proposition 2.** *Let  $(\nabla, \nabla^*)$  be a  $G$ -dual teleparallel pair on the real, finite-dimensional, smooth, parallelizable Riemannian manifold  $(\mathcal{N}, G)$  of dimension  $\dim(\mathcal{N}) = n$ , and let  $T$  be the covariant  $(0, 3)$  tensor defined in equation (26) (or equation (27)). Then:*

- *$T$  is symmetric in the first two entries if and only if  $\nabla$  is torsion-free or, equivalently, if and only if  $\{\theta^j\}_{j=1, \dots, n}$  is a basis of closed differential one-forms;*
- *$T$  is symmetric in the first and third entries if and only if  $\nabla^*$  is torsion-free or, equivalently, if and only if  $\{\alpha^j\}_{j=1, \dots, n}$  is a basis of closed differential one-forms;*
- *$T$  is symmetric in the second and third entries if and only if  $\nabla_{X_j}^* X_k = 0$  for every  $j, k = 1, \dots, n$ , or, equivalently, if and only if  $\nabla_{Y_j} Y_k = 0$   $j, k = 1, \dots, n$ .*

*Proof.* In this case it holds

$$T(X_j, X_k, Y_l) \stackrel{(26)(4)}{=} -G(\nabla_{X_j}^G X_k, Y_l). \quad (28)$$

Consequently, since  $\nabla^G$  is torsion-free, we have  $\nabla_{X_j}^G X_k - \nabla_{X_k}^G X_j = [X_j, X_k]$  and thus

$$T(X_j, X_k, Y_l) - T(X_k, X_j, Y_l) = G([X_k, X_j], Y_l) \stackrel{(13)}{=} -\theta^l([X_j, X_k]) \stackrel{(24)}{=} d\theta^j(X_k, X_l) \quad (29)$$

showing that  $T$  is symmetric in the first two entries if and only if  $\nabla$  is torsion-free or, equivalently, if and only if  $\{\theta^j\}_{j=1, \dots, n}$  is a basis of closed differential one-forms (cf. remark 2) as claimed. In particular, this is always true when we consider the mixture connection in both the classical and quantum case (cf. sections 3 and 4).

Analogously, we have

$$T(Y_j, X_k, Y_l) \stackrel{(27)(4)}{=} G(X_k, \nabla_{Y_j}^G Y_l), \quad (30)$$

so that, again since  $\nabla^G$  is torsion-free, we can use  $\nabla_{Y_j}^G Y_l - \nabla_{Y_l}^G Y_j = [Y_j, Y_l]$  and obtain

$$T(Y_j, X_k, Y_l) - T(Y_l, X_k, Y_j) = G(X_k, [Y_j, Y_l]) \stackrel{(20)}{=} -\alpha^k([Y_j, Y_l]) \stackrel{(25)}{=} d\alpha^k(Y_j, Y_l), \quad (31)$$

which means that  $T$  is symmetric in the first and third entries if and only if  $\nabla^*$  is torsion-free or, equivalently, if and only if  $\{\alpha^j\}_{j=1, \dots, n}$  is a basis of closed differential one-forms (cf. remark 2).

Finally, recalling that

$$Z(G(W, V)) = G(\nabla_Z^G W, V) + G(W, \nabla_Z^G V) \quad (32)$$

for all  $Z, W, V \in \mathfrak{X}(\mathcal{N})$  because  $\nabla^G$  is its own  $G$ -dual connection, we have

$$\begin{aligned} T(X_j, X_k, X_l) - T(X_j, X_l, X_k) &\stackrel{(27)}{=} G(X_k, \nabla_{X_j}^G X_l - \nabla_{X_j}^* X_l) - G(X_l, \nabla_{X_j}^G X_k - \nabla_{X_j}^* X_k) = \\ &\stackrel{(1)(4)(32)}{=} G(X_l, \nabla_{X_j}^* X_k), \end{aligned} \quad (33)$$



which means that  $T$  is symmetric in the second and third entries if and only if  $\nabla_{X_j}^* X_k = 0$  for every  $j, k = 1, \dots, n$ . Moreover, it also holds

$$\begin{aligned} T(Y_j, Y_k, Y_l) - T(Y_j, Y_l, Y_k) &\stackrel{(26)}{=} G(\nabla_{Y_j} Y_k - \nabla_{Y_j}^G Y_k, Y_l) - G(\nabla_{Y_j} Y_l - \nabla_{Y_j}^G Y_l, Y_k) = \\ &\stackrel{(1)(19)(32)}{=} -G(\nabla_{Y_j} Y_l, Y_k) \end{aligned} \quad (34)$$

which means that  $T$  is symmetric in the second and third entries if and only if  $\nabla_{Y_j} Y_k = 0$  for every  $j, k = 1, \dots, n$ . We thus conclude that

$$\nabla_{X_j}^* X_k = 0 \quad \forall j, k = 1, \dots, n \iff \nabla_{Y_j} Y_k = 0 \quad \forall j, k = 1, \dots, n. \quad (35)$$

Of course, if  $\nabla$  and  $\nabla^*$  are both torsion-free then it can easily be checked that  $T$  is completely symmetric.  $\square$

### 3 G-dual teleparallel pairs in Classical Information Geometry

Let  $\overline{\Delta_n}$  be the  $n$ -simplex in  $\mathbb{R}^n$ :

$$\overline{\Delta_n} := \left\{ \vec{p} = (p^1, \dots, p^n) \in \mathbb{R}^n \mid p^j \geq 0 \quad \forall j = 1, \dots, n, \text{ and } \sum_{j=1}^n p^j = 1 \right\}, \quad (36)$$

and  $\Delta_n$  its interior

$$\Delta_n := \left\{ \vec{p} = (p^1, \dots, p^n) \in \overline{\Delta_n} \mid p^j > 0 \quad \forall j = 1, \dots, n \right\}. \quad (37)$$

Note that both  $\overline{\Delta_n}$  and  $\Delta_n$  are convex subsets of  $\mathbb{R}^n$ . If  $\mathcal{X}_n$  denote a discrete set with  $n$  elements, then  $\overline{\Delta_n}$  can be interpreted as the set of probability distributions on  $\mathcal{X}_n$ , and  $\Delta_n$  as the subset of nowhere-vanishing probability distributions on  $\mathcal{X}_n$ . The space  $\Delta_n$  is a smooth manifold which is also parallelizable because it is essentially an open subset of the affine hyperplane determined by the normalization condition  $\sum_{j=1}^n p^j = 1$ .

A tangent vector at  $\vec{p} \in \Delta_n$  can be identified with an element  $\vec{a} \in \mathbb{R}^n$  such that  $\sum_{j=1}^n a^j = 0$ . If we choose  $(n-1)$  linearly independent vectors  $\vec{a}_1, \dots, \vec{a}_{n-1} \in \mathbb{R}^n$  such that  $\sum_{j=1}^n a_k^j = 0$  for all  $k = 1, \dots, (n-1)$ , we can define a basis  $\{L_k\}_{k=1, \dots, (n-1)}$  of global vector fields on  $\Delta_n$  setting

$$L_k(\vec{p}) = \vec{a}_k. \quad (38)$$

The notational change from  $X_k$  to  $L_k$  is made to remark the relation of these vector fields with the linear structure of  $\mathbb{R}^n$ , and thus with the convex structures of probability vectors. Indeed, every  $L_k$  generates a traslation on  $\mathbb{R}^n$ , and traslations are related with the linear structure of  $\mathbb{R}^n$ .

The teleparallel connection associated with  $\{L_k\}_{k=1, \dots, (n-1)}$  is known as the mixture connection on  $\Delta_n$  and it is often denoted by  $\nabla^m$ . This affine connection is intimately connected with the convex structure of  $\Delta_n$ . Indeed, as mentioned in section 2, every geodesic of  $\nabla^m$  starting at  $\vec{p}$  and with intial velocity  $\vec{v} = v^k a_k = v^k L_k(\vec{p})$ , where  $v^k \in \mathbb{R}$  for all  $k = 1, \dots, (n-1)$ , can be realized as an integral curve of the vector field  $V = v^k L_k$ , and has the simple expression

$$\vec{p}_{\vec{v}}(t) = \vec{p} + t \vec{v}. \quad (39)$$

Therefore, given two arbitrary points  $\vec{p}_1, \vec{p}_2 \in \Delta_n$ , we can build a geodesic starting at  $\vec{p}_1$  and ending at  $\vec{p}_2$  at time  $t = 1$  setting  $\vec{v} = \vec{p}_2 - \vec{p}_1$  thus obtaining the segment  $(1 - t)\vec{p}_1 + t\vec{p}_2$  describing all the possible convex combinations of  $\vec{p}_1$  and  $\vec{p}_2$ .

It is perhaps seldom observed that, from the geometrical point of view,  $\Delta_n$  is actually an homogeneous space of the multiplicative Lie group  $\mathbb{R}_+^n$ , i.e., the product of  $n$  copies of  $\mathbb{R}_+$ . Specifically, let  $(q^1, \dots, q^n) = \vec{q} \in \mathbb{R}_+^n$  and set

$$\gamma(\vec{q}, \vec{p}) := \frac{1}{N_{\vec{q}}^{\vec{p}}} (q^1 p^1, \dots, q^n p^n) \quad (40)$$

with  $N_{\vec{q}}^{\vec{p}} = \sum_{j=1}^n q^j p^j$ . It is a matter of direct computation to check that  $\gamma$  is indeed a smooth, transitive left action of  $\mathbb{R}_+^n$  on  $\Delta_n$ . Then, setting  $\vec{b}_0 = (1, \dots, 1)$  and selecting  $(n - 1)$  linearly independent vectors  $\vec{b}_1, \dots, \vec{b}_{n-1} \in \mathbb{R}^n$  such that  $\sum_{j=1}^n b_k^j = 0$  for all  $k = 1, \dots, (n - 1)$ , we obtain a basis  $\{\vec{b}_j\}_{j=0, \dots, (n-1)}$  of the Lie algebra  $\mathbb{R}^n$  of  $\mathbb{R}^n$  such that the fundamental vector field of  $\gamma$  associated with  $\vec{b}_0$  identically vanishes, while the fundamental vector field of  $\gamma$  associated with  $\vec{b}_j$  reads

$$Y_j(\vec{p}) = \left( \left( b_j^1 - N_{\vec{b}_j}^{\vec{p}} \right) p^1, \dots, \left( b_j^n - N_{\vec{b}_j}^{\vec{p}} \right) p^n \right). \quad (41)$$

It turns out that the teleparallel connection determined by the basis  $\{Y_j\}_{j=1, \dots, (n-1)}$  of vector fields on  $\Delta_n$  is precisely the exponential connection which is dual to  $\nabla^m$  with respect to the Fisher-Rao metric tensor.

According to Čencov's pioneering work (cf. [56]), the Riemannian manifold structure which is relevant in the context of Classical Information Geometry is the so-called Fisher-Rao metric tensor  $G_{FR}$  given by

$$(G_{FR}(X, Y))(\vec{p}) := \langle X(\vec{p}), Y(\vec{p}) \rangle_{\vec{p}} \quad (42)$$

where  $\langle \cdot, \cdot \rangle_{\vec{p}}$  is the weighted scalar product on  $\mathbb{R}^n$  given by

$$\langle \vec{a}, \vec{b} \rangle_{\vec{p}} := \sum_{j=1}^n \frac{a^j b^j}{p^j}. \quad (43)$$

It is well-known that the  $G_{FR}$ -dual connection of  $\nabla^m$  is the so-called exponential connection  $\nabla^e$  (cf. [4, 56]). Moreover, from proposition 1 we know that  $\nabla^e$  is the teleparallel connection associated with the gradient vector fields of an almost dual basis of  $\{L_k\}_{k=1, \dots, (n-1)}$ . At this purpose, an almost dual basis  $\{\theta^k\}_{k=1, \dots, (n-1)}$  of  $\{L_k\}_{k=1, \dots, (n-1)}$  is associated with a choice of  $(n - 1)$  linearly independent vectors  $\{\vec{b}_k\}_{k=1, \dots, (n-1)}$  such that  $\sum_{j=1}^n b_k^j = 0$  for all  $k = 1, \dots, (n - 1)$ . Indeed, setting  $\theta^k = \text{d}l^k$  with

$$l^k(\vec{p}) := \vec{p} \cdot \vec{b}_k, \quad (44)$$

it immediately follows that

$$\theta^k(L_j) = \frac{\text{d}}{\text{d}t} \left( (\vec{p} + t\vec{a}_j) \cdot \vec{b}_k \right)_{t=0} = \vec{a}_j \cdot \vec{b}_k, \quad (45)$$

and the right hand side clearly does not depend on the point  $\vec{p}$ , thus showing that  $\{\theta^k\}_{k=1, \dots, (n-1)}$  is indeed an almost dual basis of  $\{L_k\}_{k=1, \dots, (n-1)}$ . Then, a direct computation using the very definition of gradient vector field (cf. equation (13)), and equations (42) and (45) shows that the gradient vector field associated with  $\theta^k$  by the Fisher-Rao metric tensor  $G_{FR}$  is precisely

the vector field  $Y_k$  given in equation (41) as claimed. Since the exponential connection  $\nabla^e$  is the teleparallel connection determined by the basis  $\{Y_j\}_{j=1,\dots,(n-1)}$ , every geodesic of  $\nabla^e$  starting at  $\vec{p}$  and with initial velocity  $\vec{v} = v^j a_j = v^j Y_j(\vec{p})$ , where  $v^j \in \mathbb{R}$  for all  $j = 1, \dots, (n-1)$ , can be realized as an integral curve of the vector field  $V = v^j Y_j$ . Then, since the  $Y_j$ 's are the fundamental vector fields of the action of  $\mathbb{R}_+^n$  on  $\Delta_n$  described in equation (40), it immediately follows that every integral curve of  $V$  starting at  $\vec{p}$  reads

$$\vec{p}_{\vec{v}}(t) = \frac{1}{\sum_{j=1}^n e^{tv^j} p^j} \left( e^{tv^1} p^1, \dots, e^{tv^n} p^n \right), \quad (46)$$

from which it is evident that  $\nabla^e$  is complete.

## 4 G-dual teleparallel pairs in Quantum Information Geometry

The purpose of this section is to discuss the  $G$ -dual teleparallel pairs that naturally arise in the context of finite-dimensional Quantum Information Geometry when the reference teleparallel connection comes from the natural convex structure of the space of quantum states, and the metric tensors are the so-called monotone metric tensors replacing the Fisher-Rao metric tensor in the quantum context and classified by Petz [48].

Let  $\mathcal{H}$  be a complex Hilbert space of dimension  $n < \infty$ . Let  $\langle \cdot, \cdot \rangle$  denote the Hilbert space inner product on  $\mathcal{H}$ ,  $\mathcal{B}(\mathcal{H})$  denote the space of bounded linear operators on  $\mathcal{H}$ , and  $\mathcal{B}_{sa}(\mathcal{H})$  the space of self-adjoint elements in  $\mathcal{B}(\mathcal{H})$ . The vector space  $\mathcal{B}_{sa}(\mathcal{H})$  is a real Hilbert space with respect to the so-called Hilbert-Schmidt product

$$\langle \mathbf{a}, \mathbf{b} \rangle_{HS} := \text{Tr}(\mathbf{a}\mathbf{b}), \quad (47)$$

where  $\text{Tr}$  is the Hilbert space trace. We can always select a basis  $\{\sigma_j\}_{j=0,\dots,n^2-1}$  for  $\mathcal{B}_{sa}(\mathcal{H})$  in such a way that  $\sigma_0 = \mathbb{I}$ , the identity operator on  $\mathcal{H}$ , and  $\text{Tr}(\sigma_j) = 0$  for all  $j = 1, \dots, n^2 - 1$ . When  $\mathcal{H} \cong \mathbb{C}^2$ , the typical example of such a basis is the one in which  $\sigma_j$  is the  $j$ -th Pauli matrix for  $j = 1, 2, 3$ .

Every element  $\mathbf{a} \in \mathcal{B}_{sa}(\mathcal{H})$  can then be expressed as

$$\mathbf{a} = x^0 \sigma_0 + \vec{x} \cdot \vec{\sigma} \quad (48)$$

where  $\vec{x} = (x^1, \dots, x^{n^2-1}) \in \mathbb{R}^{n^2-1}$ ,  $\vec{\sigma} = (\sigma^1, \dots, \sigma^{n^2-1})$ , and  $\cdot$  is the Euclidean scalar product. The real numbers  $(x^0, \vec{x})$  determine a Cartesian coordinate system on  $\mathcal{B}_{sa}(\mathcal{H})$ .

An element in  $\mathcal{B}_{sa}(\mathcal{H})$  is called positive if its spectrum lies on the non-negative half-line, and strictly positive when its spectrum lies on the positive half-line. In particular, a strictly positive operator is invertible. It is customary to write  $\rho \geq 0$  to denote a positive operator, and  $\rho > 0$  to denote a strictly positive one. The space  $\mathcal{P}(\mathcal{H})$  of strictly positive operators on  $\mathcal{H}$  is an open convex cone in  $\mathcal{B}_{sa}(\mathcal{H})$ , which means it is an open submanifold of  $\mathcal{B}_{sa}(\mathcal{H})$ , and thus a parallelizable smooth manifold since  $\mathcal{B}_{sa}(\mathcal{H})$  is a vector space.

The space of faithful quantum states of a finite-level quantum systems associated with  $\mathcal{H}$  is identified with the space of invertible density operators on  $\mathcal{H}$  given by

$$\mathcal{S}(\mathcal{H}) := \{\rho \in \mathcal{P}(\mathcal{H}) \mid \text{Tr}(\rho) = 1\}. \quad (49)$$

Accordingly,  $\mathcal{S}(\mathcal{H})$  is a codimension-one submanifold of  $\mathcal{P}(\mathcal{H})$  which is also parallelizable because it is essentially an open subset of the affine hyperplane in  $\mathcal{B}_{sa}(\mathcal{H})$  determined by the trace-normalization condition. Moreover, it is also a convex set.

The space of all quantum states  $\overline{\mathcal{S}(\mathcal{H})}$  is the norm-closure of  $\mathcal{S}(\mathcal{H})$  inside  $\mathcal{B}(\mathcal{H})$  (or  $\mathcal{B}_{sa}(\mathcal{H})$ ), and it is a compact, convex subset of  $\mathcal{B}_{sa}(\mathcal{H})$  which is also a stratified manifold [22, 33].

Recalling equation (48) and the normalization condition in equation (49), it is clear that every  $\rho \in \overline{\mathcal{S}(\mathcal{H})}$  is such that  $x^0 = \frac{1}{n}$ . However, it is not in general possible to explicitly find all the constraints the vector  $\vec{x}$  in equation (48) must satisfy in order to ensure the positivity condition  $\rho \geq 0$ . The two-dimensional case is the only case in which a direct computation of the eigenvalues shows that  $\rho \geq 0$  is equivalent to  $\vec{x} \cdot \vec{x} \leq \frac{1}{4}$ .

Because of the normalization condition, and because of dimensional reasons related with the fact that  $\mathcal{S}(\mathcal{H})$  is a codimension-one submanifold of the open submanifold  $\mathcal{P}(\mathcal{H}) \subset \mathcal{B}_{sa}(\mathcal{H})$ , the tangent space at  $\rho$  can be expressed as the linear span

$$T_\rho \mathcal{S}(\mathcal{H}) \cong \text{span}(\sigma_1, \dots, \sigma_{n^2-1}), \quad (50)$$

and we can define the global basis  $\{L_k\}_{k=1, \dots, n^2-1}$  of vector fields on  $\mathcal{S}(\mathcal{H})$  setting

$$L_k(\rho) = \sigma_k. \quad (51)$$

Again, the notational change from  $X_k$  to  $L_k$  is made to emphasize the relation of these vector fields with the linear structure of  $\mathcal{B}_{sa}(\mathcal{H})$  (cf. the comment after equation (38)).

The teleparallel connection associated with  $\{L_k\}_{k=1, \dots, n^2-1}$  is known as the mixture connection on  $\mathcal{S}(\mathcal{H})$  and it is denoted as  $\nabla^m$ . Very much like it happens in the classical setting for the mixture connection on the interior of the n-simplex, the mixture connection on  $\mathcal{S}(\mathcal{H})$  turns out to have a natural geometric interpretation in terms of the convex structure of  $\mathcal{S}(\mathcal{H})$ . Indeed, every geodesic of  $\nabla^m$  starting at  $\rho$  with initial tangent vector  $\mathbf{v} = v^k \sigma_k = v^k L_k(\rho)$ , where  $v^k \in \mathbb{R}$  for every  $k = 1, \dots, (n^2 - 1)$ , can be realized as the integral curve of the vector field  $V = v^k L_k$  (cf. section 2) which reads

$$\rho_{\mathbf{v}}(t) = \rho + t\mathbf{v}. \quad (52)$$

Therefore, given two arbitrary points  $\rho_1, \rho_2 \in \mathcal{S}(\mathcal{H})$ , we can build a geodesic starting at  $\rho_1$  and ending at  $\rho_2$  at time  $t = 1$  by setting  $\mathbf{v} = \rho_2 - \rho_1$ , thus obtaining the segment  $(1 - t)\rho_1 + t\rho_2$  describing all the possible convex combinations of  $\rho_1$  and  $\rho_2$ . Moreover, it follows from equation (52) that  $\nabla^m$  is not complete on  $\mathcal{S}(\mathcal{H})$ .

In the context of Quantum Information Geometry, and, contrary to the situation in Classical Information Geometry, the Riemannian structure on  $\mathcal{S}(\mathcal{H})$  is not uniquely determined, and there is an infinite family of so-called **quantum monotone metric tensors** (cf. [48]). In particular, given an operator monotone function  $f: (0, \infty) \rightarrow (0, \infty)$  such that

$$f(x) = xf(x^{-1}), \quad f(1) = 1, \quad (53)$$

Petz's classification of quantum monotone metric tensor assures us that every such metric tensor  $G_f$  is associated with  $f$  by means of

$$(G_f(V, W))(\rho) = \text{Tr}(V_\rho T_\rho^f(W_\rho)), \quad (54)$$

where  $V, W$  are vector fields on  $\mathcal{S}(\mathcal{H})$ , and where  $T_\rho^f = (K_\rho^f)^{-1}$  with

$$K_\rho^f = f(L_\rho R_{\rho^{-1}}) R_\rho, \quad (55)$$

where  $L_\rho(\mathbf{a}) = \rho\mathbf{a}$  and  $R_\rho(\mathbf{a}) = \mathbf{a}\rho$ . Note that both  $L_\rho$  and  $R_\rho$  are (invertible) superoperators on  $\mathcal{B}(\mathcal{H})$ .

Being self-adjoint, every  $\rho$  can be diagonalized using an orthonormal basis  $\{|j\rangle\}_{j=1,\dots,\dim(\mathcal{H})}$  of  $\mathcal{H}$ . Writing  $\mathbf{e}_{lm}^\rho = |l\rangle\langle m|$ , we have

$$\rho = \sum_{j=1}^n p_\rho^j \mathbf{e}_{jj}^\rho. \quad (56)$$

Note that the collection of all  $\mathbf{e}_{lm}^\rho$ 's forms a basis of  $\mathcal{B}(\mathcal{H})$ .

We can also introduce the super-operators  $E_{kj}^\rho$  acting on  $\mathcal{B}(\mathcal{H})$  according to

$$E_{kj}^\rho(\mathbf{e}_{lm}^\rho) = \delta_{jl} \delta_{km} \mathbf{e}_{jk}^\rho, \quad (57)$$

and it is then a matter of straightforward computation to check that

$$K_\rho^f = \sum_{j,k=1}^n p_\rho^k f\left(\frac{p_\rho^j}{p_\rho^k}\right) E_{kj}^\rho, \quad (58)$$

and

$$T_\rho^f = \sum_{j,k=1}^n \left( p_\rho^k f\left(\frac{p_\rho^j}{p_\rho^k}\right) \right)^{-1} E_{kj}^\rho. \quad (59)$$

As a side remark, note that, for every vector field  $W$  such that its associated tangent vector  $W_\rho$  at  $\rho$  commutes with  $\rho$ , and for every vector field  $V$ , using equations (59) and (42) it follows that

$$(G_f(V, W))(\rho) = (G_{FR}(\vec{V}_d, \vec{W}_d))(\vec{p}_\rho) \quad (60)$$

with  $\vec{p}_\rho = (p_\rho^1, \dots, p_\rho^n)$ , and  $\vec{V}_d(\vec{p}_\rho), \vec{W}_d(\vec{p}_\rho)$  the vectors built out of the diagonal elements of  $V_\rho$  and  $W_\rho$  with respect to the basis of eigenvectors of  $\rho$ . Note that equation (60) holds for every choice of the operator monotone function  $f$ .

From proposition 1 we know that the  $G_f$ -dual connection  $\nabla^f$  of the mixture connection  $\nabla^m$  is the teleparallel connection associated with the gradient vector fields of an almost dual basis of  $\{L_k\}_{k=1,\dots,n}$ . An almost dual basis  $\{\theta^k\}_{k=1,\dots,n^2-1}$  of  $\{L_k\}_{k=1,\dots,n^2-1}$  is then easily seen to be associated with a choice of basis  $\{\omega_k\}_{k=0,\dots,n^2-1}$  for  $\mathcal{B}_{sa}(\mathcal{H})$  in such a way that  $\omega_0 = \mathbb{I}$  and  $Tr(\omega_k) = 0$ . Indeed, setting  $\theta^k := dl^k$  with

$$l^k(\rho) := Tr(\rho\omega_k) \quad (61)$$

the expectation-value function associated with  $\sigma_k$ , a direct computation shows that

$$\theta^k(L_j) = \frac{d}{dt} \left( l^k(\rho_j(t)) \right)_{t=0} \stackrel{(52)}{=} Tr(\sigma_j\omega_k), \quad (62)$$

and the right-hand-side does not depend on the point  $\rho$ . Then, we have

$$Tr(\sigma_j\omega_k) \stackrel{(62)}{=} (\theta^k(L_j))(\rho) \stackrel{(13)}{=} (G_f(L_j, Y_k^f))(\rho) \stackrel{(54)(51)}{=} Tr(\sigma_j T_\rho^f(Y_k^f(\rho))) \quad (63)$$

for all  $k = 1, \dots, (n^2 - 1)$ , from which it follows that the gradient vector field  $Y_k^f$  associated with  $\theta^k$  by means of  $G_f$  must be such that

$$\omega_j - T_\rho^f(Y_k^f(\rho)) = c_\rho \mathbb{I} \equiv c_\rho \omega_0. \quad (64)$$

Therefore, since  $T_\rho^f = (K_\rho^f)^{-1}$ , we have

$$Y_k^f(\rho) = K_\rho^f(\omega_k - c_\rho \mathbb{I}) \stackrel{(58)}{=} K_\rho^f(\omega_k) - c_\rho \rho, \quad (65)$$

with  $c_\rho = \text{Tr}(K_\rho^f(\omega_k))$  because it must be  $\text{Tr}(Y_k^f(\rho)) = 0$  since  $Y_k^f(\rho) \in T_\rho \mathcal{S}(\mathcal{H})$ .

From equation (6) we know that the torsion tensor of a teleparallel connection is connected with the commutator of the basis vector fields defining the teleparallel connection. It is then relevant to review the results presented in [11, 15, 16] in relation with the teleparallel connection  $\nabla^f$  determined by  $\{Y_k^f\}_{k=1,\dots,n}$ . In the above-mentioned works, even if their role with respect to  $G_f$ -dual pairs was not known, the gradient vector fields associated with the expectation value functions by means of monotone quantum metric tensors have been studied in relation with the actions of Lie groups suitably extending the unitary group  $\mathcal{U}(\mathcal{H})$  and its canonical action

$$\rho \mapsto \alpha(\mathbf{U}, \rho) = \mathbf{U} \rho \mathbf{U}^\dagger \quad (66)$$

on  $\mathcal{S}(\mathcal{H})$ , and a great deal of attention has been given to the commutators between these vector fields.

When the operator monotone function is  $f(x) = \kappa(x-1)/\ln(x)$  with  $\kappa > 0$ , the Riemannian metric tensor  $G_f$  reduces to the Bogoliubov-Kubo-Mori metric tensor (cf. [48]) and the gradient vector fields mutually commute. Consequently, the teleparallel connection  $\nabla^f$  is torsion-free and flat. This is not surprising because it is well-known that the connection which is dual to  $\nabla^m$  with respect to the Bogoliubov-Kubo-Mori metric tensor is the quantum version of the exponential connection which is torsion-free, flat, and also complete (cf. [24, 34, 44]). What is surprising is the following fact (we refer to [11, 15, 16] for more details on the construction we are about to resume). Together with the fundamental vector fields of the standard action of  $\mathcal{U}(\mathcal{H})$  on  $\mathcal{S}(\mathcal{H})$  (cf. equation (66)), the gradient vector fields giving rise to connection  $\nabla^f$  can be used to build an anti-representation of the Lie algebra of the cotangent group  $T^*\mathcal{U}(\mathcal{H})$ . This anti-representation integrates to a transitive and smooth left action on  $\mathcal{S}(\mathcal{H})$  which may be thought of as the quantum counterpart of the action  $\gamma$  (cf. equation (40)) of  $\mathbb{R}_+^n$  on  $\Delta_n$ . In particular, every geodesic of  $\nabla^f$  starting at  $\rho$  with initial velocity  $\mathbf{v}$  is of the form

$$\rho_{\mathbf{v}}(t) = \frac{e^{\ln \rho + t\mathbf{v}}}{\text{Tr}(e^{\ln \rho + t\mathbf{v}})} \quad (67)$$

from which it clearly follows that  $\nabla^f$  is complete on  $\mathcal{S}(\mathcal{H})$ . Moreover, when  $[\mathbf{v}, \rho] = \mathbf{0}$ , the geodesic in equation (67), if diagonalized, has basically the form of the geodesic of the classical exponential connection given in equation (46). Since both  $\nabla^m$  and  $\nabla^f$  are torsion-free in this case, the covariant tensor  $T$  is completely symmetric and  $(\mathcal{S}(\mathcal{H}), G_f, T)$  is a Statistical Manifold in the sense of [40].

It is also worth noting that, when  $f(x) = \frac{\kappa(x-1)(x^\kappa+1)}{2(x^\kappa-1)}$  with  $0 < \kappa < 1$ , a similar instance manifests itself. Indeed, again together with the fundamental vector fields of the standard action of  $\mathcal{U}(\mathcal{H})$  on  $\mathcal{S}(\mathcal{H})$  (cf. equation (66)), the gradient vector fields giving rise to connection  $\nabla^f$  provide a Lie algebra anti-representation. However, this time the anti-representation is of the Lie algebra  $\mathfrak{gl}(\mathcal{H})$  of the general linear group  $\mathcal{GL}(\mathcal{H})$ . This representation integrates to a transitive and smooth left action of  $\mathcal{GL}(\mathcal{H})$  on  $\mathcal{S}(\mathcal{H})$ . In particular, every geodesic of  $\nabla^f$  starting at  $\rho$  and with initial tangent vector  $\mathbf{v} = v^j \sigma_j$ , where  $v^j \in \mathbb{R}$  for every  $j = 1, \dots, (n^2 - 1)$ , is of the form

$$\rho_{\mathbf{v}}(t) = \frac{\left(e^{t\mathbf{v}} \rho^{\sqrt{\kappa}} e^{t\mathbf{v}}\right)^{\frac{1}{\sqrt{\kappa}}}}{\text{Tr}\left(\left(e^{t\mathbf{v}} \rho^{\sqrt{\kappa}} e^{t\mathbf{v}}\right)^{\frac{1}{\sqrt{\kappa}}}\right)}, \quad (68)$$



from which it is clear that  $\nabla^f$  is always complete.

When  $\kappa = 1$ , the monotone quantum metric tensor  $G_f$  coincides with the Bures-Helstrom metric tensor (cf. [9, 20, 23, 36, 54, 58]) ubiquitous in quantum estimation theory (cf. [19, 25, 42, 46, 51, 52, 53]), while when  $\kappa = \frac{1}{4}$ , the monotone quantum metric tensor  $G_f$  coincides with the Wigner-Yanase metric tensor [28, 29, 35, 39]. In all these cases, the commutator between gradient vector fields is a fundamental vector field of  $\alpha$  and, in general, it does not vanish so that the connection  $\nabla^f$  presents torsion. Therefore, the tensor  $T$  determined by  $\nabla^m, \nabla^f$ , and  $G_f$  will not be symmetric in the first and second entries so that  $(\mathcal{S}(\mathcal{H}), G_f, T)$  is not a Statistical Manifold in the sense of [40].

**Remark 3.** *Incidentally, for a two-level quantum system, the cases  $f(x) = \kappa(x-1)/\ln(x)$  with  $\kappa > 0$  and  $f(x) = \frac{\kappa(x-1)(x^\kappa+1)}{2(x^\kappa-1)}$  with  $0 < \kappa < 1$  are the only cases of monotone quantum metric tensors for which the gradient vector fields and the fundamental vector fields of the standard action of  $\mathcal{U}(\mathcal{H})$  on  $\mathcal{S}(\mathcal{H})$  give rise to a transitive group action on  $\mathcal{S}(\mathcal{H})$  (cf. [15]).*

## 5 Conclusions

In this work we investigated the notion of  $G$ -dual teleparallel pairs  $(\nabla, \nabla^*)$  of affine connections on a real, finite-dimensional, smooth, parallelizable Riemannian manifold  $(\mathcal{N}, G)$ . We showed how every such pair gives rise to a covariant  $(0, 3)$  tensor  $T$  generalizing the Amari-Čencov tensor of Statistical Manifolds to the case where connections present torsion. Then, we reviewed some examples of  $G$ -dual teleparallel pairs in the context of Classical and Quantum Information Geometry, where the existence of a suitable (torsion-free and flat) mixture connection  $\nabla^m$  (associated with a suitable convex structure) gifts us with a preferred teleparallel connection.

In the classical case, it turns out that the dual connection is also torsion-free and flat, and it is known as the exponential connection, a fact which is well-known. Following the theory developed in section 2, we find the explicit form of a basis of vector fields determining the exponential connection as the associated teleparallel connection, and show how these vector fields provide a homogeneous manifold structure for the interior of the  $n$ -simplex  $\Delta_n$  such that the geodesics of the exponential connection coincides with the images of one-parameter subgroups. As far as we know, these geometric considerations regarding the exponential connection are new.

In the quantum case, we investigate  $G_f$ -dual pairs  $(\nabla, \nabla^*)$  where  $G_f$  is a monotone quantum metric tensor, and  $\nabla$  is the mixture connection  $\nabla^m$  on the manifold  $\mathcal{S}(\mathcal{H})$  of faithful quantum states of a quantum system with finite-dimensional Hilbert space  $\mathcal{H}$ . The theory developed in section 2 allows us to conclude that  $\nabla^* \equiv \nabla^f$  is the teleparallel connection associated with the gradient vector fields of the expectation value functions. These gradient vector fields already appeared in the literature in connection with transitive actions on  $\mathcal{S}(\mathcal{H})$  of suitable extensions of the unitary group  $\mathcal{U}(\mathcal{H})$ , and we are here able to make a point of contact between these group actions and the properties of  $\nabla^f$ . In particular, whenever these groups appear, we are able to explicitly write down the geodesics of  $\nabla^f$  again as the images of one-parameter subgroups, from which it follows that  $\nabla^f$  is a complete connection. However, it is in general true that  $\nabla^f$  presents torsion unless  $G_f$  is the so-called Bogoliubov-Kubo-Mori metric tensor, in which case  $\nabla^f$  is the quantum counterpart of the exponential connection.

A natural evolution of the ideas presented here would be the investigation of how  $G$ -dual teleparallel pairs “projects down” on homogeneous manifolds. Indeed, it is clear that  $G$ -dual teleparallel pairs only exist on parallelizable manifolds, but not all manifolds of interest in

Classical and Quantum Information Geometry are parallelizable. A very relevant example is given by the manifold of pure states of a finite-level quantum system, which is diffeomorphic with the complex projective space  $\mathbb{P}(\mathcal{H})$ . In this and other cases, however, the manifold of interest  $\mathcal{N}$  is actually a homogeneous manifold for some Lie group  $\mathcal{G}$  (e.g., the unitary group  $\mathcal{U}(\mathcal{H})$  for the manifold of pure quantum states), and Lie groups are obviously parallelizable manifolds. Then, we may pull-back the Riemannian metric tensor on  $\mathcal{N}$  “coming from” Information Geometry to the Lie group  $\mathcal{G}$  by means of the canonical projection  $\pi: \mathcal{G} \rightarrow \mathcal{N}$ , and complement it to a Riemannian metric tensor  $G$  on  $\mathcal{G}$  for which  $G$ -dual teleparallel pairs exist. At this point,  $\pi$  becomes a Riemannian submersion and the idea would be to understand if and how it is possible to “project down” a  $G$ -dual teleparallel pair on the homogeneous manifold in a suitable way, perhaps exploiting a suitable projector to describe the module of vector fields on  $\mathcal{N}$  in terms of the module of vector fields on  $\mathcal{G}$ . We are currently investigating this problem, and we plan to discuss it in a future publication.

## Funding

This work has been supported by the Madrid Government (Comunidad de Madrid-Spain) under the Multiannual Agreement with UC3M in the line of “Research Funds for Beatriz Galindo Fellowships” (C&QIG-BG-CM-UC3M), and in the context of the V PRICIT (Regional Programme of Research and Technological Innovation). The authors acknowledge financial support from the Spanish Ministry of Economy and Competitiveness, through the Severo Ochoa Programme for Centres of Excellence in RD (SEV-2015/0554), the MINECO research project PID2020-117477GB-I00, and Comunidad de Madrid project QUITEMAD++, S2018/TCS-A4342. G. M. is also a member of the Gruppo Nazionale di Fisica Matematica (INDAM), Italy. F. D. C. acknowledges support from the CONEX-Plus programme funded by Universidad Carlos III de Madrid and the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 801538.

## References

- [1] R. Aldrovandi and J. G. Pereira. *Teleparallel Gravity*. Springer Netherlands, 2013. DOI: [10.1007/978-94-007-5143-9](https://doi.org/10.1007/978-94-007-5143-9). ↓ 5
- [2] S. I. Amari. *Differential-Geometrical Methods in Statistics*, volume 28 of *Lecture Notes in Statistics*. Springer-Verlag, Berlin Heidelberg, 1985. DOI: [10.1007/978-1-4612-5056-2](https://doi.org/10.1007/978-1-4612-5056-2). ↓ 2, 3
- [3] S. I. Amari. *Information Geometry and its Application*. Springer, Japan, 2016. DOI: [10.1007/978-4-431-55978-8](https://doi.org/10.1007/978-4-431-55978-8). ↓ 2
- [4] S. I. Amari and H. Nagaoka. *Methods of Information Geometry*. American Mathematical Society, Providence, RI, 2000. DOI: [10.1090/mmono/191](https://doi.org/10.1090/mmono/191). ↓ 2, 3, 4, 10
- [5] N. Ay and S. I. Amari. A novel approach to canonical divergences within information geometry. *Entropy*, 17(12):8111 – 8129, 2015. DOI: [10.3390/e17127866](https://doi.org/10.3390/e17127866), MPI-MIS Preprint. ↓ 3

- [6] N. Ay, J. Jost, H. V. Le, and L. Schwachhöfer. *Information Geometry*. Springer International Publishing, 2017. DOI: [10.1007/978-3-319-56478-4](https://doi.org/10.1007/978-3-319-56478-4). ↓ 2
- [7] N. Ay and W. Tuschmann. Dually flat manifolds and global information geometry. *Open Systems & Information Dynamics*, 9(2):195–200, 2002. DOI: [10.1023/a:1015604927654](https://doi.org/10.1023/a:1015604927654), MPI-MIS Preprint. ↓ 2
- [8] N. Ay and W. Tuschmann. Duality versus dual flatness in quantum information geometry. *Journal of Mathematical Physics*, 44(4):1512 – 1518, 2003. DOI: [10.1063/1.1556192](https://doi.org/10.1063/1.1556192), MPI-MIS Preprint. ↓ 2
- [9] I. Bengtsson and K. Życzkowski. *Geometry of Quantum States: An Introduction to Quantum Entanglement*. Cambridge University Press, New York, 2006. DOI: [10.1017/cbo9780511535048](https://doi.org/10.1017/cbo9780511535048). ↓ 15
- [10] A. L. Besse. *Einstein Manifolds*. Springer, Berlin, Heidelberg, 1987. DOI: [10.1007/978-3-540-74311-8](https://doi.org/10.1007/978-3-540-74311-8). ↓ 6
- [11] F. M. Ciaglia. Quantum states, groups and monotone metric tensors. *European Physical Journal Plus*, 135:530–16pp, 2020. DOI: [10.1140/epjp/s13360-020-00537-y](https://doi.org/10.1140/epjp/s13360-020-00537-y), arXiv:2006.10595 [math-ph]. ↓ 14
- [12] F. M. Ciaglia, F. Di Cosmo, F. Di Nocera, and P. Vitale. Monotone metric tensors in Quantum Information Geometry. *arXiv*, 2022. arXiv:2203.10857 [quant-ph]. ↓ 3
- [13] F. M. Ciaglia, F. Di Cosmo, A. Ibort, and G. Marmo. Schwinger’s Picture of Quantum Mechanics. *International Journal of Geometric Methods in Modern Physics*, 17(04):2050054–14, 2020. DOI: [10.1142/S0219887820500541](https://doi.org/10.1142/S0219887820500541), arXiv:2002.09326 [math-ph]. ↓ 2
- [14] F. M. Ciaglia, F. Di Cosmo, M. Laudato, G. Marmo, G. Mele, F. Ventriglia, and P. Vitale. A Pedagogical Intrinsic Approach to Relative Entropies as Potential Functions of Quantum Metrics: the q-z family. *Annals of Physics*, 395:238–274, 2018. DOI: [10.1016/j.aop.2018.05.015](https://doi.org/10.1016/j.aop.2018.05.015), arXiv:1711.09769 [quant-ph]. ↓ 3
- [15] F. M. Ciaglia and F. Di Nocera. Group Actions and Monotone Metric Tensors: The Qubit Case. In Frank Nielsen and Frédéric Barbaresco, editors, *Geometric Science of Information 2021*, volume 12829 of *Lecture Notes in Computer Science*, pages 145–153. Springer International Publishing, 2021. DOI: [10.1007/978-3-030-80209-7\\_17](https://doi.org/10.1007/978-3-030-80209-7_17), arXiv: 2105.15067 [quant-ph]. ↓ 14, 15
- [16] F. M. Ciaglia and F. Di Nocera. Group Actions and Monotone Quantum Metric Tensors. *Mathematics*, 10(15):2613, 2022. DOI: [10.3390/math10152613](https://doi.org/10.3390/math10152613), arXiv:2206.10394 [math-ph]. ↓ 14
- [17] F. M. Ciaglia, A. Ibort, and G. Marmo. A gentle introduction to Schwinger’s formulation of quantum mechanics: the groupoid picture. *Modern Physics Letters A*, 33(20):1850122–8, 2018. DOI: [10.1142/s0217732318501225](https://doi.org/10.1142/s0217732318501225), arXiv:1807.00519 [quant-ph]. ↓ 2

- [18] F. M. Ciaglia, A. Ibort, and G. Marmo. Schwinger’s Picture of Quantum Mechanics I: Groupoids. *International Journal of Geometric Methods in Modern Physics*, 16(08):1950119 (31), 2019. DOI: [10.1142/S0219887819501196](https://doi.org/10.1142/S0219887819501196), arXiv:1905.12274 [math-ph]. ↓2
- [19] F. M. Ciaglia, J. Jost, and L. Schwachhöfer. Differential geometric aspects of parametric estimation theory for states on finite-dimensional C\*-algebras. *Entropy*, 22(11):1332, 2020. DOI: [10.3390/e22111332](https://doi.org/10.3390/e22111332), arXiv: 2010.14394 [math-ph]. ↓15
- [20] F. M. Ciaglia, J. Jost, and L. Schwachhöfer. From the Jordan product to Riemannian geometries on classical and quantum states. *Entropy*, 22(06):637–27, 2020. DOI: [10.3390/e22060637](https://doi.org/10.3390/e22060637), arXiv:2005.02023 [math-ph]. ↓15
- [21] I. Csizár. Eine informationstheoretische Ungleichung und ihre Anwendung auf den Beweis der Ergodizität von Markoffschen Ketten. *A Magyar Tudományos Akadémia Matematikai és Fizikai Tudományok Osztályának Közleményei*, 8:85–108, 1963. ↓3
- [22] F. D’Andrea and D. Franco. On the pseudo-manifold of quantum states. *Differential Geometry and its Applications*, 78:101800, 2021. DOI: [10.1016/j.difgeo.2021.101800](https://doi.org/10.1016/j.difgeo.2021.101800), arXiv:2011.04488 [math.OA]. ↓12
- [23] J. Dittmann. On the Riemannian metric on the space of density matrices. *Reports on Mathematical Physics*, 36(3):309–315, 1995. DOI: [10.1016/0034-4877\(96\)83627-5](https://doi.org/10.1016/0034-4877(96)83627-5). ↓15
- [24] D. Felice, S. Mancini, and N. Ay. Canonical Divergence for Measuring Classical and Quantum Complexity. *Entropy*, 21(4):435, 2019. DOI: [10.3390/e21040435](https://doi.org/10.3390/e21040435), arXiv: 1903.09797 [math-ph]. ↓3, 14
- [25] C. A. Fuchs. *Distinguishability and Accessible Information in Quantum Theory*. PhD thesis, Departement IRO, Université de Montréal, 1996. arXiv:9601020 [quant-ph]. ↓15
- [26] A. Fujiwara. Geometry of Quantum Information Systems. In O. E. Barndorff-Nielsen and E. B. V. Jensen, editors, *Geometry in Present Day Science*, pages 35–48, 1999. DOI: [10.1142/3958](https://doi.org/10.1142/3958). ↓3
- [27] Fujiwara, A. Dually flat structures induced from monotone metrics on a two-level quantum state space. *European Physical Journal Plus*, 135(10), 2020. DOI: [10.1140/epjp/s13360-020-00877-9](https://doi.org/10.1140/epjp/s13360-020-00877-9), arXiv: [ ]. ↓2
- [28] P. Gibilisco and T. Isola. A characterization of Wigner-Yanase skew information among statistically monotone metrics. *Infinite Dimensional Analysis, Quantum Probability and Related Topics*, 4(4):553–557, 2001. DOI: [10.1142/s0219025701000644](https://doi.org/10.1142/s0219025701000644), UniRoma Preprint. ↓15
- [29] P. Gibilisco and T. Isola. Wigner-Yanase information on quantum state space: the geometric approach. *Journal of Mathematical Physics*, 44(9):3752–3762, 2003. DOI: [10.1063/1.1598279](https://doi.org/10.1063/1.1598279), arXiv:0304170 [math.PR]. ↓15
- [30] K. Grabowska, J. Grabowski, M. Kuś, and G. Marmo. Lie groupoids in information geometry. *Journal of Physics A: Mathematical and General*, 52(50):505202, 2019. DOI: [10.1088/1751-8121/ab542e](https://doi.org/10.1088/1751-8121/ab542e). ↓2

- [31] K. Grabowska, J. Grabowski, M. Kuś, and G. Marmo. Information Geometry on Groupoids: The Case of Singular Metrics. *Open Systems & Information Dynamics*, 27(03):2050015, 2020. DOI: [10.1142/S1230161220500158](#). ↓ 2
- [32] K. Grabowska, J. Grabowski, M. Kuś, and G. Marmo. Lifting statistical structures. *Reviews in Mathematical Physics*, 35(02), 2023. DOI: [10.1142/S0129055X22500428](#), [arXiv:2203.02938 \[math.DG\]](#). ↓ 2
- [33] J. Grabowski, M. Kuś, and G. Marmo. Geometry of quantum systems: density states and entanglement. *Journal of Physics A: Mathematical and General*, 38(47):10217–10244, 2005. DOI: [10.1088/0305-4470/38/47/011](#). ↓ 12
- [34] M. R. Grasselli and R. F. Streater. On the Uniqueness of the Chentsov Metric in Quantum Information Geometry. *Infinite Dimensional Analysis, Quantum Probability and Related Topics*, 4(2):173–182, 2001. DOI: [10.1142/S0219025701000462](#). ↓ 3, 14
- [35] H. Hasegawa. Dual geometry of the Wigner-Yanase-Dyson information content. *Infinite Dimensional Analysis, Quantum Probability and Related Topics*, 6(3):413–430, 2003. DOI: [10.1142/S021902570300133X](#). ↓ 15
- [36] C. W. Helstrom. Minimum mean-squared error of estimates in quantum statistics. *Physics Letters A*, 25(2):101–102, 1967. DOI: [10.1016/0375-9601\(67\)90366-0](#). ↓ 15
- [37] M. Henmi and H. Matsuzoe. *Statistical Manifolds Admitting Torsion and Partially Flat Spaces*, pages 37–50. Springer International Publishing, 2019. DOI: [10.1007/978-3-030-02520-5\\_3](#). ↓ 4
- [38] A. Jenčová. Geometry of Quantum States: Dual Connections and Divergence Functions. *Reports in Mathematical Physics*, 47(1):121 – 138, 2001. ↓ 3
- [39] A. Jenčová. Flat connections and Wigner-Yanase-Dyson metrics. *Reports on Mathematical Physics*, 52(3):331–351, 2003. DOI: [10.1016/S0034-4877\(03\)80033-2](#). ↓ 2, 15
- [40] S. Lauritzen. Statistical Manifolds. In S. I. Amari, O. E. Barndorff-Nielsen, R. E. Kass, S. L. Lauritzen, and C. R. Rao, editors, *Differential geometry in statistical inference*, volume 10, pages 163–216. Institute of Mathematical Statistics, 1987. DOI: [10.1214/lnms/1215467061](#), Project Euclid. ↓ 2, 14, 15
- [41] A. Lesniewski and M. B. Ruskai. Monotone riemannian metrics and relative entropy on noncommutative probability spaces. *Journal of Mathematical Physics*, 40(11):5702–5724, 1999. DOI: [10.1063/1.533053](#). ↓ 3
- [42] J. Liu, H. Yuan, X.-M. Lu, and X. Wang. Quantum Fisher information matrix and multiparameter estimation. *Journal of Physics A: Mathematical and Theoretical*, 53(2):023001–69, 2020. DOI: [10.1088/1751-8121/ab5d4d](#), [arXiv:1907.08037 \[quant-ph\]](#). ↓ 15
- [43] T. Matumoto. Any statistical manifold has a contrast function: on the  $C^3$ -functions taking the minimum at the diagonal of the product manifold. *Hiroshima mathematical journal*, 23(2):327 – 332, 1993. ↓ 3



- [44] H. Nagaoka. Differential Geometrical Aspects of Quantum State Estimation and Relative Entropy. In V. P. Belavkin, O. Hirota, and R. L. Hudson, editors, *Quantum Communications and Measurement*, Springer US, pages 449–452, 1994. DOI: [10.1007/978-1-4899-1391-3\\_44](#). ↓ [3](#), [14](#)
- [45] J. Naudts, A. Verbeure, and R. Weder. Linear Response Theory and the KMS Condition. *Communications in Mathematical Physics*, 44:87–99, 1975. DOI: [10.1007/BF01609060](#). ↓ [3](#)
- [46] M. G. A. Paris. Quantum Estimation for Quantum Technology. *International Journal of Quantum Information*, 7(1):125–137, 2009. DOI: [10.1142/S0219749909004839](#), arXiv:0804.2981 [quant-ph]. ↓ [15](#)
- [47] D. Petz. Geometry of canonical correlation on the state space of a quantum system. *Journal of Mathematical Physics*, 35(2):780–795, 1994. DOI: [10.1063/1.530611](#). ↓ [3](#)
- [48] D. Petz. Monotone metrics on matrix spaces. *Linear Algebra and its Applications*, 244:81–96, 1996. DOI: [10.1016/0024-3795\(94\)00211-8](#). ↓ [3](#), [11](#), [12](#), [14](#)
- [49] D. Petz and G. Toth. The Bogoliubov Inner Product in Quantum Statistics. *Letters in Mathematical Physics*, 27:205–216, 1993. DOI: [10.1007/BF00739578](#). ↓ [3](#)
- [50] H. Shima. *The geometry of Hessian structures*. World Scientific Publishing Company Pte. Ltd., 2007. DOI: [10.1142/9789812707536](#). ↓ [2](#)
- [51] J. Suzuki. Information Geometrical Characterization of Quantum Statistical Models in Quantum Estimation Theory. *Entropy*, 21(7):703, 2019. DOI: [10.3390/e21070703](#). ↓ [15](#)
- [52] J. Suzuki, Y. Yang, and M. Hayashi. Quantum state estimation with nuisance parameters. *Journal of Physics A: Mathematical and Theoretical*, 53(45):453001, 2020. DOI: [10.1088/1751-8121/ab8b78](#), arXiv:1911.02790 [quant-ph]. ↓ [15](#)
- [53] M. Tsang, F. Albarelli, and A. Datta. Quantum Semiparametric Estimation. *Physical Review X*, 10(3):031023, 2020. DOI: [10.1103/PhysRevX.10.031023](#). ↓ [15](#)
- [54] A. Uhlmann. The Metric of Bures and the Geometric Phase. In R. Gielerak, J. Lukierski, and Z. Popowicz, editors, *Groups and Related Topics*, pages 267–274. Springer, Dordrecht, 1992. DOI: [10.1007/978-94-011-2801-823](#). ↓ [15](#)
- [55] H. Umegaki. Conditional expectation in an operator algebra IV: Entropy and information. *Kodai Mathematical Seminar Reports*, 14(2):59–85, 1962. DOI: [10.2996/kmj/1138844604](#). ↓ [3](#)
- [56] N. N. Čencov. *Statistical Decision Rules and Optimal Inference*. American Mathematical Society, Providence, RI, 1982. DOI: [10.1090/mmono/053](#). ↓ [2](#), [10](#)
- [57] J. von Neumann. *Mathematical Foundations of Quantum Mechanics*. Princeton University Press, Princeton, NJ, 1955. ↓ [3](#)
- [58] D. Šafránek. Simple expression for the quantum Fisher information matrix. *Physical Review A*, 97:042322–06, 2018. DOI: [10.1103/PhysRevA.97.042322](#), arXiv:1801.00945 [quant-ph]. ↓ [15](#)



- [59] J. Zhang and G. Khan. From Hessian to Weitzenböck: manifolds with torsion-carrying connections. *Information Geometry*, 8, 2019. DOI: [10.1007/s41884-019-00018-x](https://doi.org/10.1007/s41884-019-00018-x). ↓ [4](#), [6](#)
- [60] J. Zhang and G. Khan. New Geometry of Parametric Statistical Models. In F. Nielsen and F. Barbaresco, editors, *Geometric Science of Information*, pages 288–296. Springer International Publishing, 2019. DOI: [10.1007/978-3-030-26980-7\\_30](https://doi.org/10.1007/978-3-030-26980-7_30). ↓ [4](#)
- [61] J. Zhang and G. Khan. Statistical Mirror Symmetry. *Differential Geometry and its Applications*, 73:101678, 2020. DOI: [10.1016/j.difgeo.2020.101678](https://doi.org/10.1016/j.difgeo.2020.101678). ↓ [4](#)